Regional oceanographic features and hydrothermal activity influence protist diversity and biogeography in the Okinawa Trough

Margaret Mars Brisbin[a*], Asa E. Conover[a], Satoshi Mitarai[a]

[a] Marine Biophysics Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan
* Corresponding Author, margaret.marsbrisbin@oist.jp

10 ## Abstract

Microbial eukaryotes (protists) contribute substantially to ecological functioning in marine ecosystems, but factors shaping protist diversity, such as dispersal barriers and environmental selection, remain difficult to parse. Deep-sea water masses, which form geographic barriers, and hydrothermal vents, which represent isolated productivity

15 hotspots, are ideal opportunities for studying the effects of dispersal barriers and environmental selection on protist communities. The Okinawa Trough, a deep, back-arc spreading basin, contains distinct water masses in the bottom waters of northern and southern regions and at least twenty-five active hydrothermal vents. In this study, we used metabarcoding to characterize protist communities from fourteen stations

20 spanning the length of the Okinawa Trough, including three hydrothermal vent sites. Significant differences in community structure reflecting regional oceanography and water mass composition were present, indicating the importance of geographic factors in shaping protist communities. Protist communities in bottom waters affected by hydrothermal activity were significantly different from communities in other bottom

25 waters, suggesting that environmental factors can be especially important in shaping community composition under specific conditions. Amplicon sequence variants that were enriched in hydrothermally influenced bottom waters largely derived from cosmopolitan protists that were present, but rare, in other near-bottom samples, thus highlighting the importance of the rare biosphere.

Introduction

Microbial unicellular eukaryotes (protists) are important contributors to all marine ecosystems, from the sunlit surface waters [1] to the deep, dark bathypelagic [2]. Extensive phylogenetic and functional diversity among protists—especially in extreme

35   environments like the deep sea and hydrothermal vents [3]—has been revealed by high-throughput sequencing with environmental samples, but factors influencing protist community structure remain difficult to parse [4, 5]. Physicochemical factors, such as temperature, light and nutrient availability, have historically been regarded as having the strongest influence on microbial community structure, but geographic restrictions—

40   between ocean basins [4] and water masses [5, 6]—and biological interactions [7] can also contribute substantially to shaping microbial community structure. Furthermore, physicochemical, biological, and geographical factors can vary in their relative influences on community structure at different depths and in different oceanic regions [8].

45   The Okinawa Trough (OT), a back-arc spreading basin (> 2000 m) within the East China Sea, is an ideal environment to investigate factors shaping protist community structure. The Kuroshio Current, the western boundary current of the North Pacific subtropical gyre, enters the OT to the east of Taiwan, and transports warm, high-salinity water northward [9]. The bottom water in the southern OT is formed from

50   Kuroshio Intermediate water (KIW), which is 55% high-salinity South China Sea Intermediate Water (SCSIW) and 45% low-salinity North Pacific Intermediate Water (NPIW), flowing over the sill (775 m) east of Taiwan and into the southern OT [10]. Additional surface water and intermediate water are entrained into the OT through the Kerama Gap, which, with a sill depth of 1100 m, is the deepest channel connecting the

55   OT and the Philippine Sea. NPIW consistently enters the deep regions of the OT through the Kerama Gap [11], increasing the contribution of NPIW to bottom waters in the northern OT to 65% [10]. However, surface flow in the Kerama Gap is periodically reversed, likely due to the arrival of mesoscale eddies from the interior of the North Pacific [11]. The topography and current structure in the OT lead to varying

60   environmental conditions in the northern and southern trough, as well as in the Kerama Gap. Most notably, water mass compositions are distinct in the bottom waters of the

southern (45% NPIW, 55% SCSIW) and northern OT (65% NPIW, 35% SCSIW) and the Kerama Gap (75% NPIW, 25% SCSIW) [10], thus facilitating study of how water mass composition impacts on protist community structure.

65   As a back-arc spreading basin, there is extensive hydrothermal activity throughout the Okinawa Trough [12]. Twenty-five active OT hydrothermal vent sites are listed in The InterRidge Database v3.4 [13], but this figure likely remains an underestimate as new vents continue to be discovered [14, 15]. The hydrothermal vents in the OT are distinct from those found at mid-ocean ridges, primarily due to thick layers

70 of terrigenous sediments overlaying vent sites. Compared to sediment-starved vent systems, OT vent fluids are typically low pH with high concentrations of CO_2, NH_4^+, boron, iodine, potassium, lithium [16] and methane [17]. Unique physicochemical conditions associated with different vent systems—such as those associated with OT vents—give rise to heterogeneous biological communities [18], warranting careful study

75 of diverse hydrothermal systems. In addition, organic matter and inorganic nutrients resuspended by vents in the OT may locally increase microbial production in bottom waters [19]. However, vent fields in the Okinawa Trough have not been studied nearly as extensively as hydrothermal vents in other oceanic regions [12].

       In this study, we present the first comprehensive survey of protist community

80 structure in the Okinawa Trough and Kuroshio Current. We analyzed protist community composition in replicate samples collected from the sea surface, subsurface chlorophyll maximum, middle water column, and bottom waters at sites spanning the entire length of the Okinawa Trough and within the Kerama Gap. In addition, bottom waters at three of our sampling sites were influenced by nearby hydrothermal vents, as evidenced by

85 increased turbidity, CDOM (colored dissolved organic matter), NH_4, and total carbon concentrations—conditions typical of hydrothermal vent plumes in the OT [16, 20]. The main objectives of the study were to: (i) assess broad patterns in protist diversity in the Okinawa Trough and Kuroshio Current, (ii) determine the extent to which OT protist diversity is affected by regional ocean circulation and water mass composition, and (iii)

90 investigate the influence of hydrothermal activity on deep-sea protist communities in the OT.

Materials and Methods

Sampling locations

95   Water samples were collected from 14 sites in the Okinawa Trough and Kerama Gap during the Japan Agency for Marine-Earth Science and Technology (JAMSTEC) MR17-03C cruise from May 29 to June 13, 2017 (Figure 1). Stations 8, 9, 10, and 11 are located in the Southern Okinawa Trough (SOT); stations 3, 4, 2, 5 make up a transect of the Kerama Gap (KG) from east to west; stations 12, 13, 14, 15, 17, and 18 are in the

100   Northern Okinawa Trough (NOT) stations. Five stations are located near known active hydrothermal vent sites; station 10 is at the Hatoma Knoll [16], station 2 is at the ANA vent site of the Daisan-Kume Knoll [14], station 11 is near the Dai-Yon Yonaguni Knoll [21], station 5 is near the Higa vent site [14], and station 12 is near the Iheya North vent field [22]. Physicochemical data collected by in-situ sensors and chemical analysis of

105   water samples were used to discern whether or not samples were influenced by hydrothermal vent plumes. Vertical profiles from stations 2, 10, and 11 showed near-bottom turbidity and CDOM maxima (Supplemental Figure 1A), which is typical for OT vent plumes [16, 20]. In addition, nutrient analysis showed elevated NH_4 concentrations in bottom water samples from stations 2 and 10, providing further evidence of

110   hydrothermal influence [16, 20] (Supplemental Figure 1B). Total carbon was only measured for samples from stations 2, 10, 14, and 15; consistent with hydrothermal influence [16, 20], stations 2 and 10 exhibit near bottom peaks in carbon concentrations (Supplemental Figure 1C). Stations 2, 10 and 11 are, therefore, included in analyses aimed at determining how hydrothermal activity influences protist community structure

115   in the OT, while the other sampling stations near vent sites are not.

Figure 1. Sampling locations in the Okinawa Trough. Numbered red circles denote sampling stations where water was collected during the Japan Agency for Marine-Earth Science and Technology (JAMSTEC) MR17-03C cruise in May and June 2017. Blue triangles indicate active hydrothermal sites according to the InterRidge Vents Database v3.4 (https://vents-data.interridge.org/ventfields-osm-map, accessed 05/08/2019). Bathymetry data was accessed and plotted through the function getNOAA.bathy in R package marmap. The 200, 1000, 2000, and 3000 m isobaths are plotted and labeled. Station 2 is at the ANA site of the Daisan-Kume Knoll, station 10 is at the Hatoma Knoll, station 11 is near the Dai-Yon Yonaguni Knoll, station 5 is near the Higa vent site, and station 12 is near the Iheya North vent field.

Sample collection

A Niskin rosette with 30 10-liter bottles and fitted with a conductivity-temperature-depth (CTD) probe (SBE 911plus, Sea-Bird Scientific, Bellevue, WA) was deployed at each cruise station to collect water from the subsurface chlorophyll maximum (SCM, 50–100 m), the middle water column (mid, 700–1500 m), and approximately 10 m above the seafloor (bottom, 772–2957 m) (Supplemental Table 1). Surface seawater was collected by bucket alongside the research vessel. Two replicates of 4.5 liters (surface) from separate bucket casts or 5 liters from separate Niskin bottles (SCM, mid, bottom) were sequentially filtered under a gentle vacuum through 10.0-μm and 0.2-μm pore-size

polytetrafluoroethylene (PTFE) filters (Millipore, Burlington, MA). Filters were flash-
frozen in liquid nitrogen and stored at -80°C. Temperature, salinity, dissolved oxygen,

140   fluorescence, and turbidity profiles were recorded by CTD probe at each station. CDOM
and chemical analyses for nitrate, nitrite, NH_4, phosphate, silicate and total carbon were
performed by Marine Works Japan Ocean Chemistry Analysis Section onboard with
water collected by Niskin bottle.

145   ***DNA extraction and sequencing library preparation***
DNA was extracted from PTFE filters (n = 224, two replicates of two filter pore-sizes at
four depths from 14 stations) following manufacturer's protocols for the DNeasy
PowerWater Kit (Qiagen, Hilden, Germany) including the optional heating step for 10
min at 65°C to fully lyse cells. Sequencing libraries were prepared following the Illumina

150   16S Metagenomic Sequencing Library Preparation manual, but with universal
eukaryotic primers for the V4 region of the eukaryotic 18S rRNA gene (F:
CCAGCASCYGCGGTAATTCC [23], R: ACTTTCGTTCTTGATYR [24]) and 58°C
annealing temperature in the initial PCR. Amplicon libraries were sequenced by the
Okinawa Institute of Science and Technology (OIST) DNA Sequencing Section on the

155   Illumina MiSeq platform with 2x300-bp v3 chemistry. Amplification and sequencing were
successful for 211 samples and at least one replicate succeeded for each sample type.

Amplicon sequence analysis
Sequence data from each of four flow-cells were denoised separately using the Divisive

160   Amplicon Denoising Algorithm [25] through the DADA2 plug-in for QIIME 2 [26]. We
analyzed amplicon sequence variants (ASVs) to maximize the amount of diversity
included in the study [27] and detect changes in community composition that may not
be apparent if analyzed at higher taxonomic levels. Denoised ASV tables were merged
before taxonomy was assigned to ASVs with a naive Bayes classifier trained on the

165   Protist Ribosomal Reference (PR2) database [28] using the QIIME 2 feature-classifier
plug-in [29]. We imported results into the R statistical environment [30] for further
processing with the R package phyloseq [31]. Sequences were initially filtered to
remove those that were not assigned taxonomy at the Kingdom level—likely data

artifacts—and all metazoan sequences. We did not apply additional prevalence or

170   minimum abundance filtering since samples were from varied locations and depths, but the DADA2 algorithm discards singletons.

Statistical analyses

To test whether community composition varied significantly by depth, filter-size, region

175   or presence or absence of hydrothermal influence in bottom water samples, we computed the Aitchison distance between samples, which minimizes compositional bias inherent in metabarcoding data [32], and performed Permutational Multivariate Analyses of Variance (PERMANOVA, 999 permutations) with the adonis2 function from the R package vegan [33]. We further used pairwise PERMANOVA (999 permutations) to test

180   which regions differed in community composition from each other. Because stations 3 and 18 were much deeper than other stations (Supplemental Table 1), we used mid-water samples from these stations in the bottom-water analyses and excluded these stations from the mid-water analyses. This is further justified because the mid-water samples at these stations were from the salinity layer contributing to bottom waters at

185   the other stations (Supplemental Figure 2). To determine if environmental variables contributed to changes in community composition, we performed Redundancy Analysis (RDA) and applied ANOVA to RDA results to test whether RDA models were statistically significant and which environmental variables significantly contributed to variation in community composition [34]. Before performing RDA, we tested for

190   collinearity of environmental variables by computing Pearson correlation coefficients between environmental variables in each depth layer (Supplemental Figure 3). If the absolute value of correlation coefficients was greater than 0.8, we only included one of the correlated variables in the RDA for that depth [35]. We further applied variance partitioning on variables that significantly contributed to variation in community

195   composition to determine the percent of variation attributable to each variable [36]. Lastly, we used the DESeq2 Bioconductor package [37] to determine which ASVs were differentially abundant in bottom water from sites influenced by hydrothermal activity compared to bottom water at other sites. The data and code necessary to reproduce the statistical analyses are available on github

200 (https://github.com/maggimars/OkinawaTroughProtists), including an interactive online document: https://maggimars.github.io/OkinawaTroughProtists/OTprotists.html.

Results

Protist diversity in the Okinawa Trough

205 Overall, 31.5 million sequencing reads were generated for this study, with 34,631–421,992 sequencing reads per sample (mean = 144,604). All sequences are available from the NCBI Sequencing Read Archive with accession PRJNA546472. Following denoising, 16.8 million sequences remained, with 1,724–215,842 sequences per sample (mean = 77,342). A total of 22,656 unique ASVs were identified in our dataset,

210 with 49–1,906 observed ASVs per sample (mean = 730). We used pairwise Wilcox tests on observed richness and Shannon indices to determine whether alpha diversity differed between depths or was influenced by hydrothermal activity. The SCM had significantly higher richness than surface, mid, and bottom waters ($p < 0.001$, Supplemental Figure 4A). Shannon indices for SCM and surface samples were not

215 significantly different but were both significantly higher than for mid and bottom waters ($p < 0.001$, Supplemental Figure 4B). Hydrothermally influenced bottom waters at stations 11, 10, and 2 did not have significantly different observed richness or Shannon indices from other bottom water samples (Supplemental Figure 4C–D).

Samples clustered by depth in a Principal Coordinate Analysis (PCoA) of

220 Aitchison distance between samples, with SCM and surface samples clustering separately from mid and bottom water samples on the primary axis (Figure 2). SCM and surface samples further separated on the secondary axis, but mid and bottom water samples clustered together. PERMANOVA results were significant at $p < 0.05$ when performed by depth layer on all samples. When samples from each depth were

225 analyzed separately, samples clustered by filter pore-size on the primary axis for surface and SCM samples, and on the secondary axis for mid and bottom water samples (Supplemental Figure 5A–D), but PERMANOVA results by filter pore-size were not statistically significant.

Surface samples were dominated by Dinoflagellata, Haptophyta, and Radiolaria

230 in the larger size-fraction and Dinoflagellata, Haptophyta, Ochrophyta and MAST groups

in the smaller size-fraction (Figure 3). In the SCM, Dinoflagellata, Haptophyta, Ochrophyta and Radiolaria were most abundant in the larger fraction and Chlorophyta, Dinoflagellata, Haptophyta, Ochrophyta, Radiolaria, and MAST groups were most abundant in the smaller size-fraction. The mid and bottom water samples were

235   dominated by Dinoflagellata and Radiolaria in both size-fractions. Interestingly, bottom water samples from hydrothermally influenced stations (2, 10, 11) included noticeably higher proportions of Ciliophora, Ochrophyta, and MAST groups.

240   **Figure 2. Principal coordinates analysis of Aitchison distances between size-fractionated protist communities from four depths in the Okinawa Trough.** Point color indicates the depth layer from which samples were collected and point shape reflects the filter pore-size used to collect samples in μm. Samples cluster in three main groups by depth: surface, subsurface chlorophyll maximum (SCM), and mid/bottom waters. Community composition was significantly
245   different by depth layer (PERMANOVA, 999 permutations, $p < 0.05$).

Figure 3. Relative abundance of major protist groups in size-fractionated samples from four depths in the Okinawa Trough (OT). Sampling stations are ordered on the x-axis from south to north: Southern Okinawa Trough (SOT) stations include 11, 10, 9, and 8; Kerama Gap (KG) stations include 3, 4, 2, and 5; Northern Okinawa Trough (NOT) stations include 12, 13, 14, 15, 17, and 18. The plot is faceted by sampling depth (columns) and filter pore-size in μm (rows). When replicates were available for a particular station/depth/filter combination, replicates were collapsed and represented with a single stacked bar. Regional community differences are not visible at the high taxonomic level represented in the plot, but there are visible differences by depth and filter pore-size. Hydrothermally influenced sites include station 10 at the Hatoma Knoll vent field, station 11 near the Dai-Yon Yonaguni Knoll, and station 2 at the ANA vent at the Daisan-Kume Knoll (highlighted in red).

Protist biogeography in the Okinawa Trough

To investigate biogeography of protist communities in the OT, we merged smaller and larger size-fractions for replicates and subset samples by depth. Permutational Multivariate Analyses of Variance (PERMANOVA, 999 permutations) by region (SOT, KG, NOT) on the Aitchison distances between samples were significant in surface, SCM, mid-water, and bottom-water samples ($p < 0.05$). Among surface samples, NOT and SOT samples were significantly different from KG samples, but NOT and SOT

samples were not significantly different from each other. Among SCM samples, NOT and SOT samples were significantly different, but neither were significantly different

270   from KG samples. For mid- and bottom-water samples, all pairwise comparisons were significant.

        Redundancy analysis was applied to evaluate the contribution of environmental factors to regional differences in protist community composition (Figure 4). ANOVA results for the surface water RDA model were significant ($p < 0.05$) and ANOVA by

275   environmental variable showed that temperature significantly affected community composition. Variance partitioning indicated that temperature explained 2.6% of variance, whereas region explained 5.5% (multiple testing adjusted R^2). Dissolved Oxygen (DO) was correlated with temperature (Supplemental Figure 3) in surface samples, so DO was excluded from the model. The RDA model for SCM samples

280   additionally included the depth of the SCM at each site and no variables covaried. The SCM model results were not significant ($p = 0.182$); variance partitioning indicated region explained 3.4% of variance. In the mid-water samples, temperature covaried with DO, CDOM, nitrate, silicate, and phosphate and, of these, only temperature was included in the RDA model. The mid water model results were statistically significantly

285   and salinity and NH_4 both significantly contributed to model results. Variance partitioning showed that region explained 7.3% of variance, whereas NH_4 and salinity explained 2.4 and 1.2%, respectively. In bottom water samples, temperature covaried with CDOM, nitrate, phosphate, silicate and, of these, only temperature was included in the RDA model. The RDA model was statistically significant and turbidity, NH_4, temperature and

290   DO significantly contributed to model results. Variance partitioning indicated that region explained 6.9% of variance, whereas turbidity explained 3.2%, NH_4 explained 1.4%, temperature explained 1.5% and DO explained 1.7 %. It should be noted that DO and temperature had very high variance inflation factors in this test (> 150), even though their correlation coefficient was < 0.8.

Figure 4. Distance based redundancy analysis (RDA) of protist communities and explanatory environmental variables for four depth-layers in the Okinawa Trough. If environmental variables covaried with a Pearson correlation coefficient greater than 0.8 (absolute value), only one covarying variable was included in the analysis; temperature covaried with dissolved oxygen (DO) in surface waters, no variables covaried at the SCM, temperature covaried with DO, colored dissolved organic matter (CDOM), nitrate (NITRAT), silicate (SILICAT), and phosphate (PHSPHT) in the mid-water column, and temperature covaried with CDOM, nitrate, phosphate, silicate in the bottom waters. The RDA model results were significant in surface (A), mid (C), and bottom waters (D) and variables that significantly contributed to variation in community composition are in bold and marked with an asterisk (ANOVA, *p* < 0.05). KG is the Kerama Gap (light blue), NOT is the Northern Okinawa Trough (dark blue), and SOT is the Southern Okinawa Trough (gold). For bottom water samples (D), hydrothermally influenced sites are demarcated as triangles.

295
300
305
310

Protist communities at hydrothermal vent sites

To evaluate the effect of hydrothermal activity on protist diversity, we first checked whether alpha diversity was significantly different at hydrothermally influenced sites;

315 hydrothermal influence did not significantly increase or decrease observed richness or Shannon indices in bottom water samples (Supplemental Figure 4C–D). To determine if hydrothermal influence altered community composition, we performed PERMANOVA by presence or absence of hydrothermal influence on the Aitchison distances between bottom water samples; community compositions at hydrothermally influenced sites were

320 significantly different from other bottom water samples (p = 0.001).

Lastly, we investigated if there were protist ASVs specifically associated with hydrothermally influenced sites by testing for differentially abundant ASVs in the hydrothermally influenced bottom samples compared to the rest of the bottom water samples using DESeq2 [37]. An ASV was considered significantly differentially

325 abundant when the False Discovery Rate adjusted p-value was less than or equal to 0.01. Overall, 45 ASVs were significantly differentially abundant between samples from hydrothermally influenced sites and the rest of the bottom water samples. Of the differentially abundant ASVs, 30 were more abundant at sites with hydrothermal influence than at the other sites, and included Dinophyceae, Syndiniales,

330 Chrysophyceae, MAST, RAD-A and RAD-B Radiolarians, Oligohymenophorea and Spirotrichea ciliates, and Picozoa ASVs (Figure 5).

Figure 5. Log2 fold-change of significantly differentially abundant ASVs in bottom water influenced by hydrothermal activity. The DESeq function in the DESeq2 package was used to test whether ASVs were significantly more (positive log2 fold-change) or less (negative log2 fold-change) abundant in hydrothermally influenced bottom water samples from Stations 10, 11, and 2 compared to bottom water samples from the other sites. Each point represents a single significantly differentially abundant ASV and taxonomic groupings are indicated on the x-axis and by color. ASVs were considered significantly differentially abundant if False Discovery Rate adjusted p-values were < 0.01.

Discussion

Due to their small size, protists, like bacteria, face few dispersal barriers in the ocean [38]. In addition, rapid asexual reproduction in many protists, as well as the ability to form resting cysts, increases their dispersal potential. As a result, many protists are cosmopolitan, suggesting that global protist diversity is low while local diversity in any given place is relatively high [39, 40]. Whether or not this is the case depends largely on how protist species and diversity are defined. For example, if a higher level of genetic diversity is included in a species definition, global diversity will be lower. Conversely, if less genetic differentiation is included in a species definition, global diversity will be higher [41]. This has been demonstrated in bacteria through studies of biogeography at deep sea hydrothermal vents: only when microdiversity—diversity at the single

355 nucleotide polymorphism level—is considered does geographic population structure emerge [17, 42]. By analyzing ASVs, which capture the highest amount of diversity available in metabarcoding datasets, we investigated whether biogeographic patterns exist among protists in the Okinawa Trough in regard to regional ocean circulation patterns, water mass composition, and hydrothermal influence. Our results demonstrate relatively weak biogeography in surface waters through to the mid-water column, but

360 show strong regional differentiation of protist communities in deeper water. In addition, protist communities in bottom waters affected by hydrothermal vent activity were significantly different from communities in other bottom water samples.

Protist biogeography in the Okinawa Trough

365 Consistent with previous studies on protist biogeography, the strongest overall influence on community composition was the depth sampled [43]. Depth-dependent diversity patterns were similar to metabarcoding results from other regions, with higher diversity in the surface and SCM than the meso- and bathypelagic [8, 43]. In addition, overall community structure by depth was comparable to global patterns: Dinoflagellata,

370 Haptophyta, Ochrophyta, Picozoa, Radiolaria and Marine Stramenopiles (MASTs) were abundant in the surface and SCM while Dinoflagellata and Radiolaria dominated in the meso- and bathypelagic [5].

While depth was the strongest determinant of community composition overall, there were significant regional differences in community composition in all depth layers

375 and region contributed more to variance than environmental variables. The Okinawa Trough is divided into Southern (SOT) and Northern (NOT) regions based on the location of the Kerama Gap (KG) and bottom water characteristics [10]. The fast-flowing, northward Kuroshio current is the major oceanographic feature in the upper water column [9]. Intermediate water consistently enters the trough through the KG,

380 whereas surface water regularly enters and occasionally exits [11, 44]. Regional differences in protist community composition are mostly consistent with these prevailing current patterns. In surface and mid-water samples, NOT and SOT communities were both significantly different from KG samples, but NOT and SOT communities were not different from each other in the surface waters, where the Kuroshio current is strongest.

385   At the SCM, NOT and SOT samples differed significantly but KG samples did not differ from NOT or SOT samples. These results may be due, in part, to Changjiang (a.k.a. Yangtze) Diluted Water (CDW) and Kuroshio water were mixing at the continental shelf near station 13. The Changjiang/Yangtze is the longest river in China and transports terrestrial nutrients and anthropogenic pollutants into the East China Sea, causing

390   eutrophication that regularly triggers causes diatom blooms [45]. CTD and nutrient measurements at station 13 showed decreased subsurface salinity and increased nutrient concentrations and chlorophyll fluorescence compared to other sampling sites (Supplemental Figure 6). In addition, there as was an increased contribution of Bacillariophyceae diatoms (Ochrophyta, Figure 4) to the community composition at

395   station 13.

        The regional pattern in protist community was most robust in the bottom waters, where protist communities were significantly distinct in each region and where each region has distinct water mass composition; bottom water in the SOT is comprised of 45% low-salinity North Pacific Intermediate Water (NPIW) and 55% high-salinity South

400   China Sea Intermediate Water (SCSIW), while KG bottom water is 75% NPIW and 25% SCSIW and NOT bottom water is 65% NPIW, 35% SCSIW [10]. Similarly, Agogue et al. (2011) found water mass composition was the strongest predictor of prokaryotic community composition in the Atlantic Ocean [6], and Pernice et al. (2016) found water mass composition was the strongest predictor of protist community composition in the

405   deep waters of the Atlantic, South and Equatorial Pacific, and Indian oceans [5]. Water masses in the ocean may either act as geographic boundaries to dispersal or select for microbes adapted to specific environmental conditions [5, 6]. If differences in community composition were due to distinct chemical properties of water masses, it would support the latter [6]. In the OT, salinity is the major marker for water mass composition [10],

410   and the salinities measured during our sampling were consistent with those reported for the different water mass compositions (Supplemental Figure 2A–B, [10]). However, salinity did not significantly contribute to variation in community composition in the bottom waters of the OT (Figure 4D). Therefore, our results support the idea that particular water mass histories influence community composition and suggest that water

415   masses may act as barriers to microbial dispersal [6].

Protist communities at hydrothermal vent sites

The protist communities in samples from hydrothermally influenced sites were significantly different from the communities in bottom waters at sites without

420   hydrothermal influence. Furthermore, environmental factors that significantly contributed to variation in community composition in bottom waters—turbidity, NH_4—are related to hydrothermal activity in the OT, where vents resuspend overlying terrigenous material creating turbid vent plumes [16]. Syndiniales, Ciliophora, and MAST (Marine Stramenopiles) ASVs were significantly enriched in hydrothermally influenced OT

425   bottom waters, which is consistent with previous work comparing protists in bottom waters with varying proximity to hydrothermal vents [3]; Ochrophyta, environmental clades of Radiolaria, and Picozoa ASVs were additionally enriched near hydrothermally influenced sites in this study (Figures 2, 5). Quantitative models of deep-water circulation suggest that vents in back-arc basins, such as the Okinawa Trough, are well-

430   connected, but basin to basin dispersal may be restricted [12]. A remaining question, then, is whether protists enriched near vents in the Okinawa Trough are similar to protists in other hydrothermal systems and other marine ecosystems.

       Dinoflagellate ASVs made up the majority of differentially abundant ASVs at hydrothermally influenced sites (22 out of 45, Figure 5) and half of these were more

435   abundant near vents. Among the enriched ASVs, eight are in the parasitic order Syndiniales, including group I (clades 2 and 7) and group II (clades 1, 6, 12, 13, 14, and 16). The remaining enriched dinoflagellate ASVs belong to the Order Dinophyceae but could not be further classified. Group II Syndiniales are common in sunlit, surface waters, whereas group I is common in suboxic and anoxic ecosystems. However, both

440   groups I and II are found in surface and deep water and have clades that have only been recovered from suboxic or anoxic ecosystems [46]. The majority of the enriched Syndiniales ASVs shared 100% identity with GenBank sequences (nr/nt, accessed 5/16/2019, [47, 48]) from a wide variety of ecosystems throughout the global ocean, including surface waters near the poles, mesopelagic water near the California coast,

445   100 and 200 m oxygenated and micro-oxic North Atlantic waters, and at the sediment interface near the Juan de Fuca Ridge vent system [49, 50]; the Syndinales ASVs

enriched in hydrothermally influenced samples clearly derive from cosmopolitan organisms. Although they are not restricted to vent systems, parasitic Syndiniales do seem to thrive in vent systems and may take advantage of increased host availability in
450   such regions [51].

        Ochrophyta accounted for the second-most enriched ASVs in hydrothermally influenced samples (Figure 5). The enriched Ochrophyta ASVs belong to several clades within the order Chrysophyceae. Two belong to clades with unknown morphology made up of environmental sequences (clades EC2H and EC2I) recovered from a diverse
455   array of marine and freshwater environments [52]. The remaining ASVs belong to the genus *Paraphysomonas*, which are colorless phagotrophs that are globally distributed in freshwater, marine, and soil ecosystems and have previously been recovered from vent sites [53]. Although several of the enriched ASVs share > 99%–100% identity with previously published vent-associated sequences [53], they are equally similar to
460   sequences recovered from many other marine environments, indicating that these enriched ASVs also derive from cosmopolitan organisms.

        Five of the ASVs enriched in hydrothermally influenced samples belong to marine stramenopile lineages—MAST-1, -7, -8 (Figure 5). Most of what is known about MAST lineages has been learned through environmental molecular surveys. Aggregate
465   analyses of sequences collected through such studies have found that MAST-1, -7, and -8 are highly diverse, globally-distributed, and abundant in surface waters [54, 55]. Culture-independent techniques have further demonstrated that MAST-1 [56], MAST-7 [57], and MAST-8 [54] are bactivorous flagellates. Therefore, it is likely that the MAST ASVs enriched in hydrothermally influenced samples represent cosmopolitan
470   bacterivores. Indeed, MAST ASVs from this study each share 99% (MAST-7) to 100% (MAST-1, -8) identity with multiple sequences recovered from surface waters, including from (anoxic) Saanich Inlet, Vancouver, Canada [58], the Scotian Shelf in the North Atlantic [59], and the Southern Ocean [60].

        The RAD-A and RAD-B radiolarian groups represent environmental clades that
475   have no morphological description—other than being picoplankton—or known ecological roles [61]. The RAD-A ASV enriched in hydrothermally influenced samples (Figure 5) shares > 99% identity with many GenBank sequences recovered from oxic

and anoxic waters globally, including the Cariaco basin in the Caribbean Sea [50], the Gulf Stream [49], and the South East Pacific [62]. The enriched RAD-B ASV shares
480   100% identity with multiple sequences recovered from the East Pacific Rise (1500 m and 2500 m), the Arctic Ocean (500 m), oxygenated water in the Cariaco Basin [50], the Juan de Fuca Ridge [63], and the Southern Ocean [64]. While the radiolarian ASVs enriched in hydrothermally influenced samples appear to be cosmopolitan, it is notable that identical and similar sequences have repeatedly been isolated from vent fields and
485   anoxic regions.

        Previous studies indicated Ciliophora are abundant in sediments and bacterial mats near hydrothermal vents [65–67]. It is unsurprising, then, that two Ciliophora ASVs were significantly more abundant in hydrothermally influenced samples (Figure 5). The enriched Spirotrichea ASV was classified to genus level by our classifier and belongs to
490   Leegaardiella, a recently described genus collected in the North Atlantic [68]. The Leegardiella ASV shared 100% coverage and identity with one sequence in GenBank— an uncultured eukaryote clone recovered from 2500 m at the East Pacific Rise—and had greater than 99% shared identity with another sequence recovered from the East Pacific Rise, also from 2500 m [49]. However, the Leegardiella ASV also had 100%
495   coverage and > 99% identity matches with several sequences recovered from Arctic surface water [69]. This Leegardiella ASV may, therefore, derive from a widely distributed genus that opportunistically becomes more abundant at vent sites. The Oligohymenophorea ASV belongs to the environmental OLIGO 5 clade, but did not have > 96% identity matches to any sequences in GenBank; the closest match was to
500   an uncultured eukaryote clone recovered from 500 m depth off the coast of California [49]. The EukRef-Ciliophora curated Oligohymenophorea phylogeny groups the OLIGO 5 clade with the subclass Scuticiliatia [70]. Interestingly, Zhao and Zu (2016) also found Spirotrichea and Scuticiliatia enriched among ciliates found near a vent site in the Northern Okinawa Trough [71]. While spirotrich ciliates are generally bacterivores [65],
505   Scuticiliatia ciliates are known to be symbionts of aquatic organisms [72], including giant hydrothermal bivalves [3]. If the Scuticiliata ASV is a symbiont, it could provide some explanation as to why more similar sequences have not been recovered elsewhere; coevolution with a dispersal-limited host could also limit symbiont dispersal [73].

However, it is equally likely that sequences with high similarity to the Scuticiliata ASV

510   have not yet been recovered elsewhere simply due to insufficient sampling.

A single ASV enriched near vent sites belonged to Picozoa (Figure 5). The first Picozoa species was described in 2013 [74] and Picozoa ecology remains poorly resolved. Information regarding their distribution derives from environmental molecular surveys, which have established five clades within Picozoa. While all Picozoa are

515   marine, biogeographical patterns vary between clades: clade BP2 has only been found in surface waters, group 2 is mostly found in deep waters, and the other three groups are globally distributed throughout the water column [75]. The enriched Picozoa ASV belongs to one of the cosmopolitan clades (group 1). The sequence shares 100% identity with three others in Genbank, from a variety of environments—surface water in

520   the Arctic [76] and Southern [77] oceans, as well as mesopelagic water near southern California [49].

Conclusions and future directions

While the Okinawa Trough provides an ideal setting to investigate protist diversity and

525   biogeography, this study represents the first comprehensive survey of protists in the region. Our results show that protist communities largely reflect regional oceanography and are influenced by water mass composition in near-bottom waters. Disentangling environmental and geographic effects on communities is challenging since many variables, such as temperature, covary with latitude. However, region consistently

530   contributed more to variance in community composition than any environmental factors in this study. We additionally found distinct protist communities associated with hydrothermally influenced sites, with bacterivore, parasitic, and putatively symbiotic protists enriched in hydrothermally influenced bottom waters. The enriched protists were mostly globally distributed, and many were previously found at vent sites in other parts

535   of the world. These results emphasize the importance of the rare biosphere; most protists seem to be widely distributed, even if rare, and can become opportunistically more abundant under certain conditions. A major limitation of metabarcoding studies with DNA is that it is impossible to know whether sequences derive from actively metabolizing cells, dead or inactive cells, or even environmental DNA; future studies will

540   benefit from incorporating methods that better delineate metabolic state. Additionally,
       metabarcoding studies suffer from focusing on a short sequence to assess diversity;
       metagenomic and metatranscriptomic approaches are becoming increasingly feasible
       for large-scale protist surveys, which will allow for finer resolution of diversity. In the
       future, comparisons between the Okinawa Trough and other hydrothermal regions will
545   continue to advance our understanding of important protists in hydrothermal
       ecosystems and how hydrothermal activity influences protist diversity and biogeography
       in the global ocean.

Acknowledgements

550   We thank the captain and crew of the JAMSTEC R/V Mirai for their assistance and
       support in sample collection. Hiroyuki Yamamoto, Hiromi Watanabe, Dhugal Lindsay,
       Mary M. Grossmann, and Yuko Hasagawa were instrumental in organizing and
       facilitating cruise sampling and Otis Brunner substantially contributed to seawater
       filtration. We are further grateful to Hiromi Watanabe for helping to access and interpret
555   environmental data and for comments on the manuscript. We thank the OIST DNA
       sequencing section (Onna, Okinawa) for carrying out the sequencing. This work was
       funded by the Marine Biophysics Unit of the Okinawa Institute of Science and
       Technology Graduate University. MMB was supported by a Japan Society for the
       Promotion of Science DC1 graduate student fellowship.

References

560
       1. Massana R. Eukaryotic picoplankton in surface oceans. *Annu Rev Microbiol* 2011; **65**: 91–
           110.
       2. Edgcomb VP. Marine protist associations and environmental impacts across trophic levels
           in the twilight zone and below. *Curr Opin Microbiol* 2016; **31**: 169–175.
565   3. Sauvadet A-L, Gobet A, Guillou L. Comparative analysis between protist communities from
           the deep-sea pelagic ecosystem and specific deep hydrothermal habitats. *Environ Microbiol*
           2010; **12**: 2946–2964.
       4. de Vargas C, Audic S, Henry N, Decelle J, Mahé F, Logares R, et al. Ocean plankton.
           Eukaryotic plankton diversity in the sunlit ocean. *Science* 2015; **348**: 1261605.

570   5.   Pernice MC, Giner CR, Logares R, Perera-Bel J, Acinas SG, Duarte CM, et al. Large
              variability of bathypelagic microbial eukaryotic communities across the world's oceans.
              ISME J 2016; **10**: 945–958.

        6.   Agogué H, Lamy D, Neal PR, Sogin ML, Herndl GJ. Water mass-specificity of bacterial
              communities in the North Atlantic revealed by massively parallel sequencing. *Mol Ecol*
575   2011; **20**: 258–274.

        7.   Lima-Mendez G, Faust K, Henry N, Decelle J, Colin S, Carcillo F, et al. Ocean plankton.
              Determinants of community structure in the global plankton interactome. *Science* 2015;
              348: 1262073.

        8.   Giner CR, Balagué V, Pernice MC, Duarte CM, Gasol JM, Logares R, et al. Marked
580   changes in diversity and relative activity of picoeukaryotes with depth in the global ocean.

        9.   Barkley RA. The Kuroshio current. *Science Journal* 1970; **6**: 54–60.

        10.   Nakamura H, Nishina A, Liu Z, Tanaka F. Intermediate and deep water formation in the
              Okinawa Trough. *Journal of Geophysical Research Oceans* 2013.

        11.   Na H, Wimbush M, Park J-H, Nakamura H, Nishina A. Observations of flow variability
585   through the Kerama Gap between the East China Sea and the Northwestern Pacific. *J
              Geophys Res C: Oceans* 2014; **119**: 689–703.

        12.   Mitarai S, Watanabe H, Nakajima Y, Shchepetkin AF, McWilliams JC. Quantifying dispersal
              from hydrothermal vent fields in the western Pacific Ocean. *Proc Natl Acad Sci U S A* 2016;
              113: 2976–2981.

590   13.   Beaulieu SE, Szafranski K. InterRidge Global Database of Active Submarine Hydrothermal
              Vent Fields, Version 3.4. http://vents-data.interridge.org. Accessed 5 May 2019.

        14.   Makabe A., Tsutsumi S., Chen C., Torimoto J., Matsui Y., Shibuya T., Miyazaki J., Kitada
              K., Kawagucci S. Discovery of New Hydrothermal Vent Fields in the Mid- and southern-
              Okinawa Trough. *Goldschmidt Abstracts*. 2016. p 1945.

595   15.   Miyazaki J, Kawagucci S, Makabe A, Takahashi A, Kitada K, Torimoto J, et al. Deepest and
              hottest hydrothermal activity in the Okinawa Trough: the Yokosuka site at Yaeyama Knoll.
              R Soc Open Sci 2017; **4**: 171570.

        16.   Toki T, Itoh M, Iwata D, Ohshima S, Shinjo R, Ishibashi J-I, et al. Geochemical
              characteristics of hydrothermal fluids at Hatoma Knoll in the southern Okinawa Trough.
600   *Geochem J* 2016; **50**: 493–525.

        17.   Mino S, Makita H, Toki T, Miyazaki J, Kato S, Watanabe H, et al. Biogeography of
              Persephonella in deep-sea hydrothermal vents of the Western Pacific. *Front Microbiol*
              2013; **4**: 107.

18. Dick GJ. The microbiomes of deep-sea hydrothermal vents: distributed globally, shaped locally. *Nat Rev Microbiol* 2019; **17**: 271–283.

19. Nagata T, Tamburini C, Arístegui J, Baltar F, Bochdansky AB, Fonda-Umani S, et al. Emerging concepts on microbial processes in the bathypelagic ocean – ecology, biogeochemistry, and genomics. *Deep Sea Research Part II: Topical Studies in Oceanography* . 2010. , **57**: 1519–1536

20. Zhang X, Sun Z, Fan D, Xu C, Wang L, Zhang X, et al. Compositional characteristics and sources of DIC and DOC in seawater of the Okinawa Trough, East China Sea. *Cont Shelf Res* 2019; **174**: 108–117.

21. Konno U, Tsunogai U, Nakagawa F, Nakaseama M, Ishibashi J-I, Nunoura T, et al. Liquid CO 2 venting on the seafloor: Yonaguni Knoll IV hydrothermal system, Okinawa Trough. *Geophys Res Lett* 2006; **33**: 725.

22. Nakajima R, Yamamoto H, Kawagucci S, Takaya Y, Nozaki T, Chen C, et al. Post-drilling changes in seabed landscape and megabenthos in a deep-sea hydrothermal system, the Iheya North field, Okinawa Trough. *PLoS One* 2015; **10**: e0123095.

23. Stoeck T, Bass D, Nebel M, Christen R, Jones MDM, Breiner H-W, et al. Multiple marker parallel tag environmental DNA sequencing reveals a highly complex eukaryotic community in marine anoxic water. *Mol Ecol* 2010; **19 Suppl 1**: 21–31.

24. Mars Brisbin M, Mesrop LY, Grossmann MM, Mitarai S. Intra-host Symbiont Diversity and Extended Symbiont Maintenance in Photosymbiotic Acantharea (Clade F). *Front Microbiol* 2018; **9**: 1998.

25. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods* 2016; **13**: 581–583.

26. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol* 2019.

27. Callahan BJ, McMurdie PJ, Holmes SP. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME J* 2017; **11**: 2639–2643.

28. Guillou L, Bachar D, Audic S, Bass D, Berney C, Bittner L, et al. The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote small sub-unit rRNA sequences with curated taxonomy. *Nucleic Acids Res* 2013; **41**: D597–604.

29. Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, et al. Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* 2018; **6**: 90.

30. R Core Team. R: A language and environment for statistical computing. 2018.

31. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* 2013; **8**: e61217.

32. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. Microbiome Datasets Are Compositional: And This Is Not Optional. *Front Microbiol* 2017; **8**.

33. Oksanen J, Guillaume Blanchet F, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan: Community Ecology Package. R package version 2.5-4. 2019.

34. Legendre P, Oksanen J. Testing the significance of canonical axes in redundancy analysis. *Methods Ecol Evol* 2011.

35. Ramette A. Multivariate analyses in microbial ecology. *FEMS Microbiol Ecol* 2007; **62**: 142–160.

36. Legendre P. Studying beta diversity: ecological variation partitioning by multiple regression and canonical analysis. *J Plant Ecol* 2008; **1**: 3–8.

37. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014; **15**: 550.

38. Whittaker KA, Rynearson TA. Evidence for environmental and ecological selection in a microbe with no geographic limits to gene flow. *Proc Natl Acad Sci U S A* 2017; **114**: 2651–2656.

39. Finlay BJ. Global dispersal of free-living microbial eukaryote species. *Science* 2002; **296**: 1061–1063.

40. Fenchel T, Finlay BJ. The Ubiquity of Small Species: Patterns of Local and Global Diversity. *Bioscience* 2004; **54**: 777–784.

41. Caron DA. Past President's address: protistan biogeography: why all the fuss? *J Eukaryot Microbiol* 2009; **56**: 105–112.

42. Mino S, Nakagawa S, Makita H, Toki T, Miyazaki J, Sievert SM, et al. Endemicity of the cosmopolitan mesophilic chemolithoautotroph Sulfurimonas at deep-sea hydrothermal vents. *ISME J* 2017; **11**: 909–919.

43. Countway PD, Gast RJ, Dennett MR, Savai P, Rose JM, Caron DA. Distinct protistan assemblages characterize the euphotic zone and deep sea (2500 m) of the western North Atlantic (Sargasso Sea and Gulf Stream). *Environ Microbiol* 2007; **9**: 1219–1232.

44. Jin B, Wang G, Liu Y, Zhang R. Interaction between the East China Sea Kuroshio and the Ryukyu Current as revealed by the self-organizing map. *J Geophys Res* 2010; **115**: 937.

45. Lin YC, Chung CC, Gong GC, Chiang KP. Diversity and abundance of haptophytes in the East China Sea. *Aquat Microb Ecol* 2014; **72**: 227–240.

46. Guillou L, Viprey M, Chambouvet A, Welsh RM, Kirkham AR, Massana R, et al. Widespread occurrence and genetic diversity of marine parasitoids belonging to Syndiniales (Alveolata). *Environ Microbiol* 2008; **10**: 3349–3365.

675   47. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+: architecture and applications. *BMC Bioinformatics* 2009; **10**: 421.

48. Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. GenBank. *Nucleic Acids Res* 2016; **44**: D67–72.

49. Lie AAY, Liu Z, Hu SK, Jones AC, Kim DY, Countway PD, et al. Investigating microbial

680   eukaryotic diversity from a global census: insights from a comparison of pyrotag and full-length sequences of 18S rRNA genes. *Appl Environ Microbiol* 2014; **80**: 4363–4373.

50. Edgcomb V, Orsi W, Bunge J, Jeon S, Christen R, Leslin C, et al. Protistan microbial observatory in the Cariaco Basin, Caribbean. I. Pyrosequencing vs Sanger insights into species richness. *ISME J* 2011; **5**: 1344–1356.

685   51. Moreira D, López-García P. Are hydrothermal vents oases for parasitic protists? *Trends Parasitol* 2003; **19**: 556–558.

52. Scoble JM, Cavalier-Smith T. Scale evolution in Paraphysomonadida (Chrysophyceae): Sequence phylogeny and revised taxonomy of Paraphysomonas, new genus Clathromonas, and 25 new species. *Eur J Protistol* 2014; **50**: 551–592.

690   53. Atkins MS, Teske AP, Anderson OR. A Survey of Flagellate Diversity at Four Deep-Sea Hydrothermal Vents in the Eastern Pacific Ocean Using Structural and Molecular Approaches. *J Eukaryot Microbiol* 2000; **47**: 400–411.

54. Massana R, del Campo J, Sieracki ME, Audic S, Logares R. Exploring the uncultured microeukaryote majority in the oceans: reevaluation of ribogroups within stramenopiles.

695   *ISME J* 2014; **8**: 854–866.

55. Massana R, Castresana J, Balagué V, Guillou L, Romari K, Groisillier A, et al. Phylogenetic and ecological analysis of novel marine stramenopiles. *Appl Environ Microbiol* 2004; **70**: 3528–3534.

56. Massana R, Terrado R, Forn I, Lovejoy C, Pedrós-Alió C. Distribution and abundance of

700   uncultured heterotrophic flagellates in the world oceans. *Environ Microbiol* 2006; **8**: 1515–1522.

57. Frias-Lopez J, Thompson A, Waldbauer J, Chisholm SW. Use of stable isotope-labelled cells to identify active grazers of picocyanobacteria in ocean surface waters. *Environ Microbiol* 2009; **11**: 512–525.

705   58. Orsi W, Song YC, Hallam S, Edgcomb V. Effect of oxygen minimum zone formation on

communities of marine protists. *ISME J* 2012; **6**: 1586–1601.

59. Dasilva CR, Li WKW, Lovejoy C. Phylogenetic diversity of eukaryotic marine microbial plankton on the Scotian Shelf Northwestern Atlantic Ocean. *J Plankton Res* 2014; **36**: 344–363.

710 60. Massana R, Guillou L, Díez B, Pedrós-Alió C. Unveiling the organisms behind novel eukaryotic ribosomal DNA sequences from the ocean. *Appl Environ Microbiol* 2002; **68**: 4554–4558.

61. Not F, Gausling R, Azam F, Heidelberg JF, Worden AZ. Vertical distribution of picoeukaryotic diversity in the Sargasso Sea. *Environ Microbiol* 2007; **9**: 1233–1252.

715 62. Shi XL, Marie D, Jardillier L, Scanlan DJ, Vaulot D. Groups without cultured representatives dominate eukaryotic picophytoplankton in the oligotrophic South East Pacific Ocean. *PLoS One* 2009; **4**: e7657.

63. Jungbluth SP, Grote J, Lin H-T, Cowen JP, Rappé MS. Microbial diversity within basement fluids of the sediment-buried Juan de Fuca Ridge flank. *ISME J* 2013; **7**: 161–172.

720 64. Clarke LJ, Bestley S, Bissett A, Deagle BE. A globally distributed Syndiniales parasite dominates the Southern Ocean micro-eukaryote community near the sea-ice edge. *ISME J* 2019; **13**: 734–737.

65. Coyne KJ, Countway PD, Pilditch CA, Lee CK, Caron DA, Cary SC. Diversity and distributional patterns of ciliates in Guaymas Basin hydrothermal vent sediments. *J*

725 *Eukaryot Microbiol* 2013; **60**: 433–447.

66. Pasulka A, Hu SK, Countway PD, Coyne KJ, Cary SC, Heidelberg KB, et al. SSU-rRNA Gene Sequencing Survey of Benthic Microbial Eukaryotes from Guaymas Basin Hydrothermal Vent. *J Eukaryot Microbiol* 2019; **66**: 637–653.

67. López-García P, Vereshchaka A, Moreira D. Eukaryotic diversity associated with

730 carbonates and fluid-seawater interface in Lost City hydrothermal field. *Environ Microbiol* 2007; **9**: 546–554.

68. Santoferrara LF, Alder VV, McManus GB. Phylogeny, classification and diversity of Choreotrichia and Oligotrichia (Ciliophora, Spirotrichea). *Mol Phylogenet Evol* 2017; **112**: 12–22.

735 69. Terrado R, Scarcella K, Thaler M, Vincent WF, Lovejoy C. Small phytoplankton in Arctic seas: vulnerability to climate change. *Biodiversity* 2013; **14**: 2–18.

70. Boscaro V, Santoferrara LF, Zhang Q, Gentekaki E, Syberg-Olsen MJ, Del Campo J, et al. EukRef-Ciliophora: a manually curated, phylogeny-based database of small subunit rRNA gene sequences of ciliates. *Environ Microbiol* 2018; **20**: 2218–2230.

740 71. Zhao F, Xu K. Molecular diversity and distribution pattern of ciliates in sediments from deep-sea hydrothermal vents in the Okinawa Trough and adjacent sea areas. *Deep Sea Res Part I* 2016; **116**: 22–32.

72. Fan X, Hu X, Al-Farraj SA, Clamp JC, Song W. Morphological description of three marine ciliates (Ciliophora, Scuticociliatia), with establishment of a new genus and two new
745   species. *Eur J Protistol* 2011; **47**: 186–196.

73. Peek AS, Feldman RA, Lutz RA, Vrijenhoek RC. Cospeciation of chemoautotrophic bacteria and deep sea clams. *Proc Natl Acad Sci U S A* 1998; **95**: 9962–9966.

74. Seenivasan R, Sausen N, Medlin LK, Melkonian M. Picomonas judraskeda gen. et sp. nov.: the first identified member of the Picozoa phylum nov., a widespread group of
750   picoeukaryotes, formerly known as 'picobiliphytes'. *PLoS One* 2013; **8**: e59565.

75. Moreira D, López-García P. The rise and fall of Picobiliphytes: how assumed autotrophs turned out to be heterotrophs. *Bioessays* 2014; **36**: 468–474.

76. Terrado R, Medrinal E, Dasilva C, Thaler M, Vincent WF, Lovejoy C. Protist community composition during spring in an Arctic flaw lead polynya. *Polar Biol* 2011; **34**: 1901–1914.

755 77. Díez B, Pedrós-Alió C, Massana R. Study of genetic diversity of eukaryotic picoplankton in different oceanic regions by small-subunit rRNA gene cloning and sequencing. *Appl Environ Microbiol* 2001; **67**: 2932–2941.