# Sensory-modality independent activation of the brain network for language

Sophie Arana[1,2,*], André Marquand[1], Annika Hultén[1,2], Peter Hagoort[1,2], Jan-Mathijs Schoffelen[1]

**1 Donders Institute, Radboud University, Nijmegen, The Netherlands**
**2 Max Planck Institute for Psycholinguistics**

**\* s.arana@donders.ru.nl**

## Abstract

The meaning of a sentence can be understood, whether presented in written or spoken form. Therefore it is highly probable that brain processes supporting language comprehension are at least partly independent of sensory modality. To identify where and when in the brain language processing is independent of sensory modality, we directly compared neuromagnetic brain signals of 200 human subjects (102 males) either reading or listening to sentences. We used multiset canonical correlation analysis to align individual subject data in a way that boosts those aspects of the signal that are common to all, allowing us to capture word-by-word signal variations, consistent across subjects and at a fine temporal scale. Quantifying this consistency in activation across both reading and listening tasks revealed a mostly left hemispheric cortical network. Areas showing consistent activity patterns include not only areas previously implicated in higher-level language processing, such as left prefrontal, superior & middle temporal areas and anterior temporal lobe, but also parts of the control-network as well as subcentral and more posterior temporal-parietal areas. Activity in this supramodal sentence processing network starts in temporal areas and rapidly spreads to the other regions involved. The findings do not only indicate the involvement of a large network of brain areas in supramodal language processing, but also indicate that the linguistic information contained in the unfolding sentences modulates brain activity in a word-specific manner across subjects.

## Introduction

Language can be realized in different modalities: amongst others through writing or in speech. Depending on whether the sensory input modality is visual or auditory, different dedicated networks of brain areas are activated in order to derive meaning from the physical stimulus. Besides the obvious differences in the recruitment of brain circuits that process low-level sensory information, differences in linguistic features across sensory modalities prompt a differential activation of brain areas involved in higher-order

processing as well. For instance, speech is enriched with meaningful prosodic cues, but also requires a coarticulated signal to be parsed into individual words. Written text contains cues in the form of punctuation and spaces, and has the advantage of instantaneous availability of full information compared to the temporally unfolding nature of speech. If text is presented word-by-word, as was done in the present as well as many previous studies, the temporal structure of auditory stimulation can be approximated. Nonetheless, the auditory information necessary for word recognition is still temporally distributed, whereas in the visual modality, the entire wordform is instantly available on screen.These, and other, differences are paralleled in the brain's response, and thus the sensory modality in which language stimuli are presented determines the dominant spatiotemporal patterns that will be elicited [19].

Regardless of these differences, the same core message can be conveyed in either modality. Therefore, language processing models of the past and present ([16], [18],[21]) usually include not only early sensory (up to 200ms) processing steps, but also contain late (200 - 500 ms), more abstract, and supposedly supramodal processing steps. While early processing is largely unimodal, and supported by brain regions in the respective primary and associative sensory areas, later processes (for instance lexical retrieval and integration) that activate several areas within the temporo-frontal language network are assumed to do so independent of modality.

In order to gain insight into the location and timing of brain processes representing this latter, higher order processing of the linguistic content, researchers so far had to rely on carefully manipulated experimental conditions. As a result, our current understanding of how the brain processes language across different modalities reflects a large variety in tasks (semantic decision task [10], error detection task [8], [11], passive hearing/listening [25], orthogonal tasks like size judgment [31]) and stimulus material (ranging from words [10] to sentences [3] and stories [5, 14, 38]). Despite this wealth of experimental findings and resulting insights, an important interpretational limitation stems from the fact that the majority of studies employ modality specific low-level baseline conditions (tone pairs and lines, spectrally rotated speech and false fonts, non-words, white noise [29]) to remove the sensory component of the processing. It can be difficult to assess whether the resulting effects are comparable across auditory and visual experiments. Recent fMRI work has demonstrated sensory-modality independent brain activity by directly

comparing the BOLD response across visual and auditory presentations [14; and 38]. Yet, fMRI signals lack the temporal resolution to allow for a temporally sufficiently fine grained investigation of the response to individual words.

There are only few studies using magnetoencephalography (MEG) to study the supramodal brain response and they all rely on aggregate activation measures (i.e. based on event-related averaging) to study the supramodal brain response [31, 3, 36, 47]. Aggregate measures capture only generic components in the neural response. While these generic components make a large contribution to the neural activity measured during language processing, we also know that there exist meaningful variability in the neural response that is stimulus-specific and robust[4]. This stimulus-specific activity is likely to play a big role in the higher-level language processes we are interested in. A complete analysis of the supramodal language network, therefore, needs to tap into these subtle variations as well.

Here, we overcome previous limitation by achieving a direct comparison that does not require modality-specific baseline conditions and that leverages word-by-word variation in the brain response. Using magnetoencephalography (MEG) signals from 200 subjects, we performed a quantitative assessment of the sensory modality independent brain activity following word onset during sentence processing. The MEG data forms part of a large publicly available dataset [40], and has been used in other publications as well [28], [39], [24], [27]. We identified widespread left hemispheric involvement, starting from 325 ms after word onset in the temporal lobe and rapidly spreading to anterior areas. These findings not only provide a quantitative confirmation of earlier findings in a large study sample, they also indicate that supramodal linguistic information conveyed by the individual words in their sentence context leads to subtle fluctuations in brain activation patterns that are correlated across different subjects.

## A Temporal alignment procedure

## B Cross-validated multiset canonical correlation analysis

**Figure 1.** **Analysis pipeline.** Caption on next page

**Figure 1.** (Previous page.) (A) Temporal alignment procedure. MEG signals of auditory and visual subjects differed in length due to different presentation rates. To achieve alignment between signals of auditory and visual subjects, auditory signals were epoched into overlapping segments. Each segment's first sample corresponds to the auditory word onset but each segment's length depends on the duration of the equivalent visual stimulus. Segments were then concatenated in original order to recover signal for the full sentence length. This way, the neural response to each word is fully taken into account in further comparisons, including in the case of short words for which stimulus late processing coincided with the next word presentation. (B) Starting point for the multi-set canonical correlation analysis were parcel-based neural signals for all subjects, consisting of five spatial components each. 1. Signals for all sentence trials were split into five subsets and for cross-validation one subset of sentences was left out as test data, while the remaining four subsets served as training data. 2. Based on the training dataset only an unmixing matrix was found, per parcel, defining the linear combination of the five spatial components so that the correlation across sets (subjects) and time samples were maximized. The cross-covariance was computed between all subjects' spatial components and across time collapsing over sentence trials. 3. The projection was applied to the test data to compute canonical variables for the left out sentence trials (purple outline) for all subjects. Steps 2 and 3 were repeated for all folds until each sentence subset had been left out once and the resulting canonical variables were concatenated until the entire signal was transformed. 4. Canonical variables were epoched according to word onsets and for each time point a subject-by-subject correlation matrix was computed across words. Correlation between cross-modal subjects (pink outline) were interpreted as quantifying supramodal activation.

## Materials and Methods

### Subjects

A total of 204 native Dutch speakers (102 males), with an age range of 18–33 years (mean of 22 years), participated in the experiment. In the current analysis, data from 200 subjects were included. Exclusion of four subjects was due to technical issues during acquisition, which made their datasets not suitable for our analysis pipeline. All subjects were right-handed, had normal or corrected-to-normal vision, and reported no history of neurological, developmental, or language deficits. The study was approved by the local ethics committee (CMO, the local "Committee on Research Involving Human Participants" in the Arnhem–Nijmegen region) and followed the guidelines of the Helsinki declaration. All subjects gave written informed consent before participation and received monetary compensation for their participation.

## Experimental Design 

The subjects were seated comfortably in a magnetically shielded room and presented with
Dutch sentences. From the total stimulus set of 360 sentences six subsets of 120 sentences
were created. This resulted in six different groups of subjects who were presented with
the same subset of stimuli although in a different (randomized) order with some overlap
of items between groups. Within each group of subjects half of them performed the task
in only the visual, the other half in only the auditory modality. In the visual modality,
words were presented sequentially on a back-projection screen, placed in front of them
(vertical refresh rate of 60 Hz) at the center of the screen within a visual angle of 4
degrees, in a black mono-spaced font, on a grey background. Each word was separated
by an empty screen for 300 ms and the inter-sentence interval was jittered between 3200
and 4200 ms. Mean duration of words was 351 ms (minimum 300 ms and maximum
1400 ms), depending on word length. The median duration of whole sentences was 8.3 s
(range 6.2 - 12 s). Auditory sentences had a median duration of 4.2 s (range 2.8 - 6.0
s) spoken in a natural pace. The duration of each visual word was determined by a
function of the relative duration of each letter in the auditory task plus a constant of
150 ms. In the auditory task the stimuli were presented via plastic tubes and ear pieces
to both ears. Before the experiment, the hearing threshold was determined individually
and the stimuli were then presented at an intensity of 50 dB above the hearing threshold.
A female native Dutch speaker recorded the auditory versions of the stimuli. The audio
files were recorded in stereo at 44100 Hz. During the post processing the audio files were
low-pass filtered at 8500 Hz and normalized so that all audio files had the same peak
amplitude, and same peak intensity. All stimuli were presented using the Presentation
software (Version 16.0, Neurobehavioral Systems, Inc). Sentences were presented in small
blocks, of five sentences each, along with blocks containing scrambled sentences, which
were not used here. See Lam et al. [28] for more details about the stimulus material
used. In order to check for compliance, 20% of the trials were randomly followed by
a yes/no question about the content of the previous sentence/word list. Half of the
questions addressed the content of the sentence (e.g. Did grandma give a cookie to the
girl?) whereas the other half, addressed one of the main content words (e.g. Was the
word 'grandma' mentioned?). Subjects answered the question by pressing a button for

'Yes'/ 'No' with their left index and middle finger, respectively. 

## MEG Data Acquisition & Structural imaging 

MEG data were collected with a 275 axial gradiometer system (CTF). The signals were 
analog low-pass-filtered at 300 Hz and digitized at a sampling frequency of 1,200 Hz. 
The subject's head was registered to the MEG-sensor array using three coils attached to 
the subject's head (nasion, and left and right ear canals). Throughout the measurement, 
the head position was continuously monitored using custom software [44]. During breaks 
the subject was allowed to reposition to the original position if needed. Participants 
were able to maintain a head position within 5 mm of their original position. Three 
bipolar Ag/AgCl electrode pairs were used to measure the horizontal and vertical 
electrooculogram and the electrocardiogram. 

A T1-weighted magnetization-prepared rapid gradient-echo (MP-RAGE) pulse se- 
quence was used for the structural images, with the following parameters: volume TR = 
2300 ms, TE = 3.03 ms, 8 degree flip-angle, 1 slab, slice-matrix size = $256 \times 256$, slice 
thickness = 1 mm, field of view = 256 mm, isotropic voxel-size = $1.0 \times 1.0 \times 1.0$ mm. 
A vitamin-E capsule was placed as fiducial behind the right ear to allow a visual identifi- 
cation of left-right consistency. 

## Preprocessing 

Data were bandpass filtered between 0.5 and 20 Hz, and epoched according to sentence 
onset, each epoch varying in length, depending on the number of words within each 
sentence. Samples contaminated by artifacts due to eye movements, muscular activity, 
and superconducting quantum interference device jumps were replaced by NaN before 
further analysis. Since all sentences had been presented in random order, we reordered 
sentences for each subject to yield the same order across subjects. Subsequently, the sig- 
nals of the auditory subjects were temporally aligned to the signals of the visual subjects, 
ensuring coincidence of the onset of the individual words across modalities (Figure 1A). 
This alignment was needed to accommodate for differences in word presentation rate. 
The alignment was achieved by first epoching the auditory subject's signals into smaller 
overlapping segments. Each segment's first sample corresponded to one of the word 

onsets as annotated manually according to the audio file while each segment's length 137
depended on the duration of the visual presentation of the corresponding word. Finally, 138
all segments were concatenated again in the original order. By defining segments that 139
were longer than the corresponding auditory word duration, the neural response to each 140
word is fully taken into account and matched to the visual signal, even in the case of 141
short words where the response partly coincided with the next word presentation. MEG 142
data were then downsampled to 120 Hz. 143

## Source Reconstruction 144

We used linearly constrained minimum variance beamforming (LCMV) [46] to reconstruct 145
activity onto a parcellated cortically constrained source model. For this, we computed 146
the covariance matrix between all MEG-sensor pairs, as the average covariance matrix 147
across the cleaned single trial covariance estimates. This covariance matrix was used 148
in combination with the forward model, defined on a set of 8,196 locations on the 149
subject-specific reconstruction of the cortical sheet to generate a set of spatial filters, one 150
filter per dipole location. Individual cortical sheets were generated with the Freesurfer 151
package [12, version 5.1] (surfer.nmr.mgh.harvard.edu), coregistered to a template with 152
a surface-based coregistration approach, using Caret software [45] (download here 153
and here), and subsequently downsampled to 8,196 nodes, using the MNE software 154
[17] (martinos.org/mne/stable/index.html)). The forward model was computed using 155
FieldTrip's singleshell method [34], where the required brain/skull boundary was obtained 156
from the subject-specific T1-weighted anatomical images. We further reduced the 157
dimensionality of the data to 191 parcels per hemisphere [39]. For each parcel, we 158
obtained a parcel-specific spatial filter as follows: We concatenated the spatial filters 159
of the dipoles comprising the parcel, and used the concatenated spatial filter to obtain 160
a set of time courses of the reconstructed signal at each parcel. Next, we performed a 161
principal component analysis and selected for each parcel the first five spatial components 162
explaining most of the variance in the signal. 163

## Multi-set Canonical Correlation Analysis

Multi-set canonical correlation analysis (MCCA) [37], [13] was applied to find projections of those five spatial components that would transform the subject-specific signals so as to boost similarities between them. Canonical correlation analysis (CCA) is a standard multivariate statistical method often used to investigate underlying relationships between two sets of variables. Classically, canonical variates are estimated by transforming the two sets in a way that optimizes their correlation. We applied a generalized version of the classical approach (MCCA) [26], which extends the method to multiple sets, here multiple subjects. In our case, we find linear combinations of the five spatial components for each of two subjects, so that the correlation across time between those subjects is maximized. Since we have more than two subjects, we find for each subject its own linear combination, which maximizes the correlation across time between all subjects from both modality groups (auditory and visual stimulation). Following Parra we obtained the optimal projection as the eigenvector with the largest eigenvalue of a square matrix $D^{-1}R$, where R and D are square matrices:

$$R = \begin{pmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,N} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,N} \\ \vdots & \vdots & \ddots & \vdots \\ a_{N,1} & a_{N,2} & \cdots & a_{N,N} \end{pmatrix}, D = \begin{pmatrix} a_{1,1} & 0 & \cdots & 0 \\ 0 & a_{2,2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{N,N} \end{pmatrix}$$

Where $a^{lk}$ are cross-covariance matrices between subject pairs and D contains only the diagonal blocks of within subject covariances [37]. In our case the cross-covariance matrices are of size 5-by-5 containing the cross-covariance between all five spatial components for a given subject (pair). The cross-correlation is computed across time points for each sentence and subsequently averaged across sentences. It is important to note here, that the canonical variates resulting from the optimal projection do not reflect sentence averages anymore but have the same temporal resolution as the original source signals. CCA is prone to overfitting and known to be unstable [15]. For reliable CCA estimates the number of samples should be much larger than the number of features, i.a. a sample-to-feature ratio of 20/1 is recommended [43]. We estimated the canonical variables over concatenated data, which included between 756 and 1453 samples per sentence compared to only five features (spatial components) which provides a decent

sample-to-feature ratio. Further, we estimated our canonical variables out-of sample 192
using 5-fold cross-validation to limit overfitting. We randomly split all sentences into 193
five subsets, estimating projections on 96 sentences and applying them to the 24 left out 194
sentences (Figure 1B). 195

## Statistical Analysis 196

As per the study design, the subjects were assigned to one of six stimulus sets. Different 197
groups of subjects were presented with different sets of sentences. Since MCCA relies 198
on commonalities across datasets, we could only combine data from subjects who 199
received the exact same stimulation. We therefore applied MCCA for each subgroup of 200
subjects who listened to or saw the same stimuli separately. Initially, we constrained our 201
analysis to the first set of 33 subjects (henceforth exploratory dataset). After applying 202
the projection to the data we computed a time-resolved Pearson correlation between 203
all possible subject pairings. To this end, we first epoched the resulting canonical 204
components according to individual word onsets and selected only content words (nouns, 205
adjectives & verbs) for subsequent steps. We then subtracted the mean across samples 206
and subsequently computed the correlation across words for each point in time and 207
subject pairing. Correlation coefficients of cross-modality pairings, that is correlations 208
between subjects reading and subjects listening to the sentences are interpreted as 209
capturing supramodal processing. We used a permutation test with clustering over time 210
and space (parcels) for family-wise error rate correction for statistical inference [32], 211
using 1000 randomizations of the epoched words. To this end, we randomized word 212
order for the source-reconstructed parcel time series of the auditory subjects to test 213
for exchangeability of the exact word pairing across sensory modalities. For a more 214
conservative test, we additionally tested the observed correlation patterns against a null 215
distribution obtained by permuting sentence order and doing this prior to MCCA. The 216
results and a description of this additional test can be found in supporting Figure 3-1. 217

We then analyzed the remaining sets of subjects (confirmatory dataset) using the 218
same analysis pipeline. We evaluated the overlap in the results across all six subgroups 219
using information prevalence inference [1]. Prevalence inference allows to formulate a 220
complex null hypothesis, i.e. that the prevalence of the effect is smaller than or equal 221

to a threshold prevalence, where the threshold can be realized by different values. For 222
each of the six sets of data, we obtained spatial maps of time-resolved supramodal 223
correlations, as well as 1000 permutation estimates after word order shuffling (see above). 224
We used the smallest observed average correlation across subgroups as the second-level 225
test statistic. We then tested the majority null hypothesis of the prevalence of the effect 226
being smaller or equal to a threshold prevalence. For this, we computed the largest 227
threshold such that the corresponding null hypothesis could still be rejected at the given 228
significance level $\alpha$. This was done after concatenating the minimum statistic from all 229
parcels and time points, using the maximum statistic to correct for multiple comparisons 230
in time and space (parcels). For each parcel we evaluated the highest threshold at which 231
the prevalence null hypothesis could be rejected at a level of $\alpha = 0.05$ (see supporting 232
Figures 4-1 and 4-2 for cortical maps showing thresholds averaged and per time point). 233

## Code Accessibility                                                         234

All analyses were done with custom-written MATLAB scripts and FieldTrip [35] and 235
the corresponding code is available upon request. 236

# Results                                                                      237

## Modality-specific activation                                               238

We first quantified the similarity between different subjects' brain response within 239
only the exploratory dataset (33 subjects) by correlating word-by-word fluctuations in 240
brain activity between all possible pairs of subjects. Averaging the correlations across 241
those subject pairings for which subjects were stimulated either in the same sensory 242
modality, or each in a different modality, allowed us to evaluate the modality-specific 243
brain response and the supramodal response, respectively. As displayed in Figure 2, early 244
sensory cortical areas only show correlated activity for the group of subjects receiving 245
the stimuli in the corresponding sensory modality, for the visual (red), and auditory 246
(blue) modalities. We found that MCCA is a crucial analysis step in order to reveal 247
meaningful inter-subject correlations. Only after MCCA does cortical activity in visual 248
and auditory areas become significantly correlated across those subjects performing the 249

task in the visual or auditory domain respectively (Figure 2A).

## Supramodal activation patterns

We averaged between-subject correlations over all cross-modal subject pairings as a metric for supramodal activity. We observed significant supramodal correlated activation patterns in mostly left-lateralized cortical areas (Figure 3). The effect has a large spatial and temporal extent, becoming apparent as early as 250 ms and lasting until 700 ms after word onset. Parcels in middle superior temporal gyrus (STG) contribute to the effect at the earliest time points, followed by the posterior and anterior part of the STG and about 50 ms later the anterior temporal pole. Supramodal correlated activation in ventral temporal cortex follows a similar temporal and spatial pattern, with supramodal correlations starting out more posterior around 292 ms and evolution towards the middle anterior temporal lobe at 308 ms. Other areas that express supramodal activity at relatively early time points are medial prefrontal cortex and primary auditory cortex (250 ms), followed by subcentral parietal regions and supramarginal gyrus at around 300 ms, and finally dorsolateral frontal cortex (DLFC, 325 ms). By the time 375 ms have passed, the entire orbito-frontal cortex, anterior and DLFC as well as inferior frontal gyrus (IFG) show strong supramodal subject correlation. Supramodal activation in the frontal lobe further extends towards posterior regions including pre- and postcentral gyrus. At around 400 ms supramodal subject correlation in the anterior temporal pole reaches its peak. In addition to the lateral cortical areas, correlated activity also extends to left dorsal and ventral anterior cingulate cortex as well as left fusiform gyrus. The spatio-temporal patterns of supramodal activation described so far are robust, also when ordinal word position and MCCA overfitting is controlled for in the statistical evaluation (see supporting Figure 3-1).

## Prevalence Inference

Our confirmatory analysis combined over all six datasets and tested whether the spatiotemporal patterns observed in the exploratory dataset would generalize to the population. For those parcels at which the global null hypothesis could be rejected, we infer that at least in one of the datasets an effect of supramodal processing was present

(Figure 4). In addition, we evaluated the majority null hypothesis of whether, in the majority of subgroups in the population, the data contains an effect (threshold > 0.5, outlined in black in figure 4B).

The global null hypothesis (no information in any set of subjects in the population) could be rejected at a level of $\alpha = 0.05$ in on average 40 parcels per time point (between 325 and 617 ms after onset, std. = 31.48). For those parcels, the average largest lower bound $\gamma 0$ at which the prevalence null hypothesis can be rejected is shown in supporting Figures 4-1 and 4-2. For those parcels for which the largest bound $\gamma 0$ is larger than or equal to 0.5 we can infer that in the majority of the datasets the activity patterns were similar across subjects independent of modality.

This majority null hypothesis could be rejected (at a level of $\alpha = 0.05$) in 90% of parcels that also showed a global effect (Figure 4, see supporting Figure 4-3 for results in the right medial hemisphere). Compared to the temporal pattern of the largest nominal suprathreshold cluster from the cluster-based permutation test conducted on the exploratory dataset, the effect became significant in the majority of datasets later in time and was less long lasting (325 - 608 ms). Given this time span, the majority null hypothesis was rejected in on average 42% of those parcels contributing most to the largest cluster. The orbitofrontal cortex and IFG showed an involvement in supramodal processing in both analyses, but the effect there was much more temporally sustained in the exploratory dataset. In addition, according to the exploratory dataset, supramodal activation of STG occurred almost 100 ms earlier as compared to IFG. Based on the confirmatory dataset, however, supramodal correlated activation in IFG and STG appeared almost simultaneously. Finally, the exploratory analysis revealed supramodal activation in primary and premotor areas extending over the entire left dorsolateral surface, of which only the most ventral parcels close to the Sylvian fissure were significantly supramodal in the majority of datasets. Thus, the spatial extent of the effect was partly reduced for prevalence inference, compared to the cluster-based permutation approach on the exploratory data. Nevertheless, widely overlapping anatomical regions were indicated by both analyses, encompassing dorsolateral frontal gyrus and the middle & superior parts of the temporal lobe at first and later also inferior frontal and orbito-frontal cortex, as well as anterior temporal lobe.

**Figure 2. Specificity of the within-modality correlated activity patterns.** (A) Time-resolved correlation values averaged across all visual subject pairings for a parcel in left primary visual cortex (upper panel) and all auditory subject pairings for a parcel in left primary auditory cortex (lower panel), before (dark grey line), and after MCCA (blue and red lines). Light grey lines show recomputed correlation values for 1000 random permutations of word order across subjects. Notably, signals of auditory subjects highly correlate even before word onset. This is likely due to a more varied distribution of information in the auditory signal caused by the continuous nature of auditory stimulation and as a result differing time points at which individual words become uniquely recognizable. The MCCA is blind to the stimulus timing and will thus find canonical variables that yield maximal correlations at any timepoint if possible. (B) Cortical map of the spatial distribution of correlations, comparing within modality visual subject pairs (red) with auditory subject pairs (blue). Correlation strength is expressed as the Pearson correlation coefficient averaged over a time window from 150 to 200 ms post word-onset and normalized by the maximum value of that window.

**Figure 3. Supramodal correlated activity patterns.** Time-resolved spatial maps of supramodal correlated activity patterns (averaged over all possible cross-modal subject pairings) in the left hemisphere. Medial views of the brain surface are depicted in the first and third row, lateral views in the second and fourth row. Color codes for strength of correlation. Colored parcels were most strongly correlated between cross-modal subject pairs (nonparametric permutation test, corrected for multiple comparisons).

**Figure 4.** **Supramodal correlated activity patterns consistent across the majority of datasets.** Caption on next page

**Figure 4.** (Previous page.) Supramodal correlated activity patterns of word-specific activity consistent across the majority of datasets. (A) Averaged correlation time courses (mean over all possible cross-modal subject pairings) are shown for selected parcels in inferior frontal Gyrus (IFG, green), supramarginal Gyrus (red), Subcentral Cortex (orange), Anterior cingulate cortex (ACC, pink), anterior middle temporal Gyrus (aMTG, purple), and middle superior temporal gyrus (mSTG, blue). Time courses are shown for each dataset individually (light-colored lines) as well as averaged (dark lines). Grey shaded areas mark statistically significant time points. (B) Time-resolved spatial maps of cross-modal correlations in the left hemisphere. Medial views of the brain surface are depicted in the first, third and fifth row, lateral views in the second, fourth and sixth row. For those parcels that were part of the largest nominal suprathreshold cluster tested on only the exploratory dataset, the mean correlation over all six datasets is shown. Color codes for strength of correlation. In addition, the parcels at which the majority null hypothesis according to prevalence inference could be rejected are outlined in black.

# Discussion 310

The aim of this research was to quantify similarities of the brain response across reading 311
and listening at a fine temporal scale. To this end we correlated word-by-word fluctuations 312
in the neural activity across subjects receiving either auditory or visual stimulation. We 313
identified a widespread left-lateralized brain network, which was activated independent 314
of modality starting from 325 ms after word onset. Importantly, dividing our large 315
study sample into six independent subsets, we could directly quantify the consistency 316
and generalizability of these observed activity patterns. The spatial distribution of the 317
supramodal activation is in line with the known involvement of left hemispheric areas, 318
including parts of left temporal cortex, left inferior parietal lobe, as well as portions of 319
prefrontal cortex [48], [10], [11], [7], [30], [29], [42], [23]. The involvement of both STG 320
and IFG also fits with predictions from the Memory, Unification and Control model 321
(MUC), in which activity reverberating within a posterior-frontal network ([2], [18]) is 322
thought to be crucial for adequate language processing. According to the MUC model 323
temporal and parietal areas support the retrieval of lexical information, while unification 324
processes are supported by inferior frontal cortex. Bidirectional communication [39] 325
between these areas is facilitated by white matter connections. Concerning the temporal 326
dynamics, we observe that temporal areas are supramodally activated at earliest time 327
points and sustain this activation for the longest as compared to other regions. Over 328
time, supramodal activation spreads from middle and posterior left STG to the anterior 329
temporal pole. This rapid progression of activity from posterior to anterior regions 330
mirrors previous observations ([31], [47]), adding to those findings a direct quantitative 331

comparison of the supramodal brain activity. <sub>332</sub>

## Beyond the core language network and the single word level <sub>333</sub>

Whereas prior studies on supramodal processes have reported involvement of mainly <sub>334</sub>
inferior parts of the frontal cortex [25], [29], [11], [31], [23], [33], we also observed <sub>335</sub>
modality-independent activity in more dorsal regions of the frontal cortex. This dorsal <sub>336</sub>
frontal cortical involvement could well be caused by our use of linguistically rich sentence <sub>337</sub>
material, of varying syntactic complexity, as opposed to single words or short phrases <sub>338</sub>
as stimuli. Indeed, discrepancies with respect to frontal lobe involvement in modality- <sub>339</sub>
independent processing have been argued to mainly arise from differences in stimulus <sub>340</sub>
material and task demands [7], and a recent meta-analysis has identified that more <sub>341</sub>
complex syntax robustly activates more dorsal parts of the left inferior frontal gyrus [20]. <sub>342</sub>
Further, a previously published analysis on the visual input modality of this MEG data <sub>343</sub>
showed DLFC to be sensitive to sentence progression effects, when compared against a <sub>344</sub>
word list condition [24]. <sub>345</sub>

Two previous fMRI studies have reported supramodal effects in response to narratives, <sub>346</sub>
which similarly can be considered complex stimuli [14, 38]. Regev et al. directly correlated <sub>347</sub>
the BOLD response evoked by different modalities. They report supramodal activation <sub>348</sub>
in the left frontal lobe that extends beyond inferior frontal regions but not necessarily <sub>349</sub>
in DLFC. Deniz et al. study modality-independent brain areas by directly modelling <sub>350</sub>
semantic features of the stimulus in one modality and predicting BOLD responses based <sub>351</sub>
on the learned weights in the other modality. Deniz et al. report an extensive set of <sub>352</sub>
voxels across the prefrontal cortex to be well predicted across modalities. They further <sub>353</sub>
show that semantic representations do not differ across stimulus-modality within DLFC. <sub>354</sub>
In summary, while it seems like complex stimuli like sentences and narratives consistently <sub>355</sub>
activate prefrontal areas beyond only inferior frontal cortex, the reasons for those areas <sub>356</sub>
to be supramodally activated are not yet clear. <sub>357</sub>

In contrast to ours and previous studies on supramodal processing, both fMRI studies <sub>358</sub>
report supramodal activation not to be restricted to the left hemisphere only. However, <sub>359</sub>
only Regev et al. additionally report extra-linguistic areas like the precuneus to be <sub>360</sub>
supramodally active. According to the authors the involvement of the precuneus might <sub>361</sub>

be due to their narratives being more strongly contextually modulated as compared to sentences. Our results also point towards extra-linguistic areas being involved in supramodal processing. Specifically, we find bilateral supramodal activation within anterior cingulate cortex (ACC). The ACC is a midline structure, forming part of a large domain-general executive control network supporting language processing [18], [9]. It has been shown to be sensitive to statistical contingencies in the language input supporting a potential role in mediating learning and adaptation in response to predictive regularities in both local experimental as well as global environment [49]. It should be noted that deep sources are normally poorly detectable in MEG [22] and we thus consider any interpretations with respect to the midline structures as tentative. Nonetheless, we generally observe that supramodal processing involves more than the traditional language areas.

## Supramodal orthography-phonology mapping

We observed supramodal activation in post-central and subcentral gyrus, as well as supramarginal gyrus. These areas show significant supramodal activation in the majority of datasets, which coincides temporally with supramodal activation of primary auditory cortex. Increases of neural activity in supramarginal gyrus have been repeatedly elicited by cross-modal tasks [41], such as rhyming judgments to visually presented words [6], for which conversion between orthographic and phonological representations is likely needed. At the same time post- and subcentral areas partly span articulatory motor and somatosensory areas for the mouth and tongue. Together, the modality-independent activation of these areas suggests that retrieval of phonetic and articulatory mappings is not limited to speech perception only but also occurs during passive reading.

## Leveraging word-by-word variability of the neural response

Brain activity exhibits a high degree of moment-to-moment variability. While some of this variability is related to the experimental stimulation and therefore associated with cognitive activity, some of it is simply unrelated ongoing neural activity. By applying MCCA across subjects we reduced the noise and made subtle word-by-word fluctuations in the MEG signal interpretable. Comparing neural activity across subjects

is challenging due to differing position or orientation of neuronal sources relative to <sub>391</sub> the MEG sensors. The use of parcellated MEG source reconstruction in combination <sub>392</sub> with exact temporal alignment of individual sentences across subjects, allowed for the <sub>393</sub> extraction of signal components that are shared across subjects, and provided a way <sub>394</sub> to reduce the spatial variability across subjects, which is commonly observed in more <sub>395</sub> traditional (for instance, dipole fitting) procedures [47]. MCCA thus allowed us to more <sub>396</sub> directly investigate time-resolved inter-subject correlations and move beyond expressing <sub>397</sub> activation as event-related averages [31]. Importantly, our analysis approach allows us to <sub>398</sub> conclude that the identified cortical areas are implicated in supramodal processing in a <sub>399</sub> word-specific way. Our findings therefore go beyond showing a general activation of those <sub>400</sub> areas as compared to baseline but rather reveal consistent word-by-word fluctuations of <sub>401</sub> activation within the recruited areas. <sub>402</sub>

## Latency of supramodal processing <sub>403</sub>

The temporal alignment procedure of the subject data, as a necessary preparation <sub>404</sub> step for the MCCA procedure, followed by the estimation of time-resolved intersubject <sub>405</sub> correlations, focused on common signal aspects that are exactly synchronized across <sub>406</sub> subjects. The differences in sensory modality specific characteristics of the input signal <sub>407</sub> require dedicated processing with likely different processing latencies, which may also <sub>408</sub> lead to latency differences in the activation of supramodal areas. For example, in <sub>409</sub> previous work investigating supramodal language processing [31], shorter reaction times <sub>410</sub> during the visual task were observed, yet earlier activity peaks were found for the <sub>411</sub> auditory task in corresponding early sensory cortex and the left anterior temporal lobe. <sub>412</sub> In contrast, other work observed earlier anterior temporal lobe activation for visual, <sub>413</sub> compared to auditory stimulation [3]. Our results indicate that there is a certain degree <sub>414</sub> of overlap across modalities in the temporal window within which supramodal cortical <sub>415</sub> areas are activated. It is possible, that we were able to observe more temporally extensive <sub>416</sub> activation, for instance, activity related to unification processes, given that we used longer <sub>417</sub> sentences as compared to using words [10],[31],[6],[30],[47] or short phrases [3],[8],[7]. In <sub>418</sub> addition, any overlap may have been amplified as a necessary consequence of the MCCA <sub>419</sub> procedure. Evidently, time-resolved correlations between signals from auditory subjects <sub>420</sub>

were boosted with less temporal specificity compared to visual subjects (Figure 2B). <sup>421</sup>
This observation was unexpected and may be due to a more continuous stimulation in <sup>422</sup>
the auditory experiment. For example, as the sound of a spoken word unfolds, the timing <sup>423</sup>
at which it becomes uniquely recognizable will vary across word. Thus, the distribution <sup>424</sup>
of information in the auditory signal is much more varied as compared to the visual. <sup>425</sup>
MCCA will pick up on any common relationship across subjects regardless of timing. In <sup>426</sup>
our specific application, projections were estimated on concatenated data, effectively <sup>427</sup>
making the method blind to word onset boundaries. <sup>428</sup>

In conclusion, this study provides direct neurophysiological evidence for sensory <sup>429</sup>
modality independent processes supporting language comprehension in multiple left <sup>430</sup>
hemispheric brain areas. We identified a network of areas including domain general <sup>431</sup>
control areas as well as phonological mapping circuits over and above traditional higher- <sup>432</sup>
level language areas in frontal and temporal-parietal regions, by quantifying between- <sup>433</sup>
subject consistency of their respective word-specific activation patterns. These consistent <sup>434</sup>
activation patterns were word-specific, and thus likely reflect more than just generic <sup>435</sup>
activation during language processing. Based on a prevalence statistical analysis, we <sup>436</sup>
conclude that these findings are generalizable to the wider population. Finally, this <sup>437</sup>
work indicates that alignment of individual subject data through multi-set canonical <sup>438</sup>
correlation analysis might be a promising tool for investigating subtle word-in-context <sup>439</sup>
specific modulations of brain activity in the language processing system. <sup>440</sup>

were boosted with less temporal specificity compared to visual subjects (Figure 2B). [421]
This observation was unexpected and may be due to a more continuous stimulation in [422]
the auditory experiment. For example, as the sound of a spoken word unfolds, the timing [423]
at which it becomes uniquely recognizable will vary across word. Thus, the distribution [424]
of information in the auditory signal is much more varied as compared to the visual. [425]
MCCA will pick up on any common relationship across subjects regardless of timing. In [426]
our specific application, projections were estimated on concatenated data, effectively [427]
making the method blind to word onset boundaries. [428]

In conclusion, this study provides direct neurophysiological evidence for sensory [429]
modality independent processes supporting language comprehension in multiple left [430]
hemispheric brain areas. We identified a network of areas including domain general [431]
control areas as well as phonological mapping circuits over and above traditional higher- [432]
level language areas in frontal and temporal-parietal regions, by quantifying between- [433]
subject consistency of their respective word-specific activation patterns. These consistent [434]
activation patterns were word-specific, and thus likely reflect more than just generic [435]
activation during language processing. Based on a prevalence statistical analysis, we [436]
conclude that these findings are generalizable to the wider population. Finally, this [437]
work indicates that alignment of individual subject data through multi-set canonical [438]
correlation analysis might be a promising tool for investigating subtle word-in-context [439]
specific modulations of brain activity in the language processing system. [440]

# Acknowledgments 

We thank Phillip Alday for providing helpful comments. This work was supported by The Netherlands Organisation for Scientific Research (NWO Vidi: 864.14.011, awarded to J.M.S.). 

# References

1. C. Allefeld, K. Görgen, and J. D. Haynes. Valid population inference for information-based imaging: From the second-level t-test to prevalence inference. *NeuroImage*, 141:378–392, 2016.

2. G. Baggio and P. Hagoort. The balance between memory and unification in semantics: A dynamic account of the N400. *Language and Cognitive Processes*, 26(9):1338–1367, 2011.

3. D. K. Bemis and L. Pylkkänen. Basic linguistic composition recruits the left anterior temporal lobe and left angular gyrus during both listening and reading. *Cerebral Cortex*, 23(8):1859–1873, 2013.

4. A. Ben-Yakov, C. J. Honey, Y. Lerner, and U. Hasson. Loss of reliable temporal structure in event-related averaging of naturalistic stimuli. *NeuroImage*, 63(1):501 – 506, 2012.

5. M. M. Berl, E. S. Duke, J. Mayo, L. R. Rosenberger, E. N. Moore, J. VanMeter, N. B. Ratner, C. J. Vaidya, and W. D. Gaillard. Functional anatomy of listening and reading comprehension during development. *Brain and Language*, 114(2):115–125, 2010.

6. J. R. Booth, D. D. Burman, J. R. Meyer, D. R. Gitelman, T. B. Parrish, and M. M. Mesulam. Modality independence of word comprehension. *Human Brain Mapping*, 16(4):251–261, 2002.

7. D. Braze, W. E. Mencl, W. Tabor, K. R. Pugh, R. Todd Constable, R. K. Fulbright, J. S. Magnuson, J. A. Van Dyke, and D. P. Shankweiler. Unification of sentence processing via ear and eye: An fMRI study. *Cortex*, 47(4):416–431, 2011.

8. A. Carpentier, K. R. Pugh, M. Westerveld, C. Studholme, O. Skrinjar, J. L. Thompson, D. D. Spencer, and R. T. Constable. Functional MRI of language processing: Dependence on input modality and temporal lobe epilepsy. *Epilepsia*, 42(10):1241–1254, 2001.

9. I. Cattinelli, N. A. Borghese, M. Gallucci, and E. Paulesu. Reading the reading brain: A new meta-analysis of functional imaging data on reading. *Journal of Neurolinguistics*, 26(1):214–238, 2013.

10. M. W. L. Chee, K. M. O'Craven, R. Bergida, B. R. Rosen, and R. L. Savoy. Auditory and visual word processing studied with fMRI. *Human Brain Mapping*, 7(1):15–28, 1999.

11. R. T. Constable, K. R. Pugh, E. Berroya, W. E. Mencl, M. Westerveld, W. Ni, and D. Shankweiler. Sentence complexity and input modality effects in sentence comprehension: An fMRI study. *NeuroImage*, 22(1):11–21, 2004.

12. A. M. Dale, B. Fischl, and M. I. Sereno. Cortical surface-based analysis: I. segmentation and surface reconstruction. *NeuroImage*, 9(2):179 – 194, 1999.

13. A. de Cheveigné, G. M. Di Liberto, D. Arzounian, D. D. Wong, J. Hjortkjær, S. Fuglsang, and L. C. Parra. Multiway canonical correlation analysis of brain data. *NeuroImage*, 186:728–740, 2018.

14. F. Deniz, A. O. Nunez-Elizalde, A. G. Huth, and J. L. Gallant. The Representation of Semantic Information Across Human Cerebral Cortex During Listening Versus Reading Is Invariant to Stimulus Modality. *The Journal of Neuroscience*, 39(39):7722–7736, 2019.

15. R. Dinga, L. Schmaal, B. W. Penninx, M. J. van Tol, D. J. Veltman, L. van Velzen, M. Mennes, N. J. van der Wee, and A. F. Marquand. Evaluating the evidence for biotypes of depression: Methodological replication and extension of Drysdale et al. (2017). *NeuroImage: Clinical*, 22:101796, 2019.

16. N. Geschwind. Specializations of the human brain. *Scientific American*, 2(241):180–201, 1979.

17. A. Gramfort, M. Luessi, E. Larson, D. A. Engemann, D. Strohmeier, C. Brodbeck, L. Parkkonen, and M. S. Hämäläinen. Mne software for processing meg and eeg data. *NeuroImage*, 86:446 – 460, 2014.

18. P. Hagoort. The core and beyond in the language-ready brain. *Neuroscience and Biobehavioral Reviews*, 81:194–204, 2017.

19. P. Hagoort and C. M. Brown. ERP effects of listening to speech compared to reading: the P600 to syntactic visual presentation. *Neuropsychologia*, 38:1531–1549, 2000.

20. P. Hagoort and P. Indefrey. The Neurobiology of Language Beyond Single Words. *Annual Review of Neuroscience*, 37(1):347–362, 2014.

21. G. Hickok and D. Poeppel. The Cortical Organisation fo Speech Processing. *Nature*, 8(5):393–402, 2007.

22. A. Hillebrand and G. R. Barnes. A quantitative assessment of the sensitivity of whole-head MEG to activity in the adult human cortex. *NeuroImage*, 16(3A):638–650, 2002.

23. F. Homae, R. Hashimoto, K. Nakajima, Y. Miyashita, and K. L. Sakai. From perception to sentence comprehension: The convergence of auditory and visual information of language in the left inferior frontal cortex. *NeuroImage*, 16(4):883–900, 2002.

24. A. Hultén, J.-M. Schoffelen, J. Uddén, N. H. Lam, and P. Hagoort. How the brain makes sense beyond the processing of single words – An MEG study. *NeuroImage*, 186:586–594, 2019.

25. G. Jobard, M. Vigneau, B. Mazoyer, and N. Tzourio-Mazoyer. Impact of modality and linguistic complexity during reading and listening tasks. *NeuroImage*, 34(2):784–800, 2007.

26. J. R. Kettenring. Canonical analysis of several sets of variables. *Biometrika*, 58(3):433–451, 1971.

27. N. H. Lam, A. Hultén, P. Hagoort, and J. M. Schoffelen. Robust neuronal oscillatory entrainment to speech displays individual variation in lateralisation. *Language, Cognition and Neuroscience*, 33(8):943–954, 2018.

28. N. H. Lam, J.-M. Schoffelen, J. Uddén, A. Hultén, and P. Hagoort. Neural activity during sentence processing as reflected in theta, alpha, beta, and gamma oscillations. *NeuroImage*, 142:43–54, 2016.

29. R. Lindenberg and L. Scheef. Supramodal language comprehension: Role of the left temporal lobe for listening and reading. *Neuropsychologia*, 45(10):2407–2415, 2007.

30. A. G. Liuzzi, R. Bruffaerts, R. Peeters, K. Adamczuk, E. Keuleers, S. De Deyne, G. Storms, P. Dupont, and R. Vandenberghe. Cross-modal representation of spoken and written word meaning in left pars triangularis. *NeuroImage*, 150:292–307, 2017.

31. K. Marinkovic, R. P. Dhond, A. M. Dale, M. Glessner, V. Carr, and E. Halgren. Spatiotemporal dynamics of modality-specific and supramodal word processing. *Neuron*, 38(3):487–497, 2003.

32. E. Maris and R. Oostenveld. Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, 164(1):177–190, 2007.

33. E. Michael, T. A. Keller, P. Carpenter, and M. A. Just. fMRI investigation of sentence comprehension by eye and by ear: Modality fingerprints on cognitive processes. *Human Brain Mapping*, 13(4):239–252, 2001.

34. G. Nolte. The magnetic lead field theorem in the quasi-static approximation and its use for magnetoenchephalography forward calculation in realistic volume conductors. *Physics in Medicine and Biology*, 48(22):3637–3652, 2003.

35. R. Oostenveld, P. Fries, E. Maris, and J. M. Schoffelen. FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, 2011:1–9, 2011.

36. A. C. Papanicolaou, M. Kilintari, R. Rezaie, S. Narayana, and A. Babajani-Feremi. The Role of the Primary Sensory Cortices in Early Language Processing. *Journal of Cognitive Neuroscience*, 29(10), 2017.

37. L. C. Parra. Multi-set Canonical Correlation Analysis simply explained. *arXiv:1802.03759*, (11 February 2018), 2018.

38. M. Regev, C. J. Honey, E. Simony, and U. Hasson. Selective and invariant neural responses to spoken and written narratives. *Journal of Neuroscience*, 33(40):15978–15988, 2013.

39. J.-M. Schoffelen, A. Hultén, N. Lam, A. F. Marquand, J. Uddén, and P. Hagoort. Frequency-specific directed interactions in the human brain network for language. *Proceedings of the National Academy of Sciences*, 114(30):8083–8088, 2017.

40. J.-M. Schoffelen, R. Oostenveld, N. H. L. Lam, J. Uddén, A. Hultén, and P. Hagoort. A 204-subject multimodal neuroimaging dataset to study language processing. *Scientific Data*, 6(1):1–17, 2019.

41. M. W. Sliwinska, M. Khadilkar, J. Campbell-Ratcliffe, F. Quevenco, and J. T. Devlin. Early and sustained supramarginal gyrus contributions to phonological processing. *Frontiers in Psychology*, 3:1–10, 2012.

42. G. Spitsyna, J. E. Warren, S. K. Scott, F. E. Turkheimer, and R. J. S. Wise. Converging Language Streams in the Human Temporal Lobe. *Journal of Neuroscience*, 26(28):7328–7336, 2006.

43. J. P. Stevens. Canonical Correlation. In *Applied multivariate statistics for the social sciences*, pages 395–412. Routledge, 2012.

44. A. Stolk, A. Todorovic, J. M. Schoffelen, and R. Oostenveld. Online and offline tools for head movement compensation in MEG. *NeuroImage*, 68:39–48, 2013.

45. D. C. Van Essen, H. A. Drury, J. Dickson, J. Harwell, D. Hanlon, and C. H. Anderson. An Integrated Software Suite for Surface-based Analyses of Cerebral Cortex. *Journal of the American Medical Informatics Association*, 8(5):443–459, 09 2001.

46. B. D. Van Veen, W. van Drongelen, M. Yuchtman, and A. Suzuki. Localization of brain electrical activity via linearly constrained minimum variance spatial filtering. *IEEE Transactions on Biomedical Engineering*, 44(9):867–880, 1997.

47. J. Vartiainen, T. Parviainen, and R. Salmelin. Spatiotemporal Convergence of Semantic Processing in Reading and Speech Perception. *Journal of Neuroscience*, 29(29):9271–9280, 2009.

48. M. Vigneau, V. Beaucousin, B. Mazoyer, M. Vigneau, N. Tzourio-Mazoyer, L. Petit, G. Jobard, P.-Y. Hervé, F. Crivello, L. Zago, and E. Mellet. What is right-hemisphere contribution to phonological, lexico-semantic, and sentence processing? *NeuroImage*, 54(1):577–593, 2010.

49. K. Weber, E. F. Lau, B. Stillerman, and G. R. Kuperberg. The Yin and the Yang of prediction: An fMRI study of semantic predictive processing. *PLoS ONE*, 11(3):1–25, 2016.

**Figure 3-1. Figure 3-1. Significant supramodal correlated activity patterns as assessed by additional permutation test.**

In an additional significance test, we shuffled the sentence order 500 times prior to MCCA, controlling for the possibility that MCCA as a preprocessing step may artificially increase correlations between subjects (through overfitting). Importantly, this permutation was not fully unconstrained, since we aimed at aligning sentences across modalities with the same number of words, to avoid loss of data and to preserve ordinal word position. Thus, we did a random pairing between sentences with the same number of words, after binning the sentences according to their word count. Sentences consisting of 9, 14 or 15 words were infrequent, with fewer than 5 occurrences each. After each permutation, we performed the temporal alignment between sensory modalities (aligning the word onsets), followed by cross-validated MCCA and computation of the time-resolved correlations of crossmodal subject pairs. Due to the long computation time of the canonical variates, we created this null distribution for the exploratory data only. In the figure, color codes for strength of correlation. Colored parcels were most strongly correlated between cross-modal subject pairs (corrected for multiple comparisons).

**Figure 4-1. Figure 4-1. Cortical map of maximum threshold $\gamma 0$.**
For those parcels at which the global null hypothesis could be rejected, the mean (over time) maximum threshold is plotted, for which the null hypothesis can be rejected ($\alpha = 0.05$). Given the sample size of six datasets, the number of second-level permutations and a significance level of $\alpha = 0.05$ the maximally possible threshold that can be reached is 0.5633.

**Figure 4-2. Figure 4-2. Cortical map of prevalence threshold $\gamma0$.**
For those parcels at which the global null hypothesis could be rejected, the maximum threshold is plotted, for which the null hypothesis can be rejected at a level of $\alpha = 0.05$.

**Figure 4-3. Figure 4-3. Time-resolved spatial maps of cross-modal correlations for the right hemisphere.**

The average correlation over all six datasets is shown. Color codes for strength of correlation. In addition, the parcels at which the majority null hypothesis according to prevalence inference could be rejected are outlined in black.