Sensory-modality independent activation of the brain network for language

Sophie Arana[1,2,*], André Marquand[1], Annika Hultén[1,2], Peter Hagoort[1,2], Jan-Mathijs Schoffelen[1]

1 Donders Institute, Radboud University, Nijmegen, The Netherlands
2 Max Planck Institute for Psycholinguistics

* s.arana@donders.ru.nl

Abstract

The meaning of a sentence can be understood, whether presented in written or spoken form. Therefore it is highly probable that brain processes supporting language comprehension are at least partly independent of sensory modality. To identify where and when in the brain language processing is independent of sensory modality, we directly compared neuromagnetic brain signals of 200 human subjects (102 males) either reading or listening to sentences. We used multiset canonical correlation analysis to align individual subject data in a way that boosts those aspects of the signal that are common to all, allowing us to capture word-by-word signal variations, consistent across subjects and at a fine temporal scale. Quantifying this consistency in activation across both reading and listening tasks revealed a mostly left hemispheric cortical network. Areas showing consistent activity patterns include not only areas previously implicated in higher-level language processing, such as left prefrontal, superior & middle temporal areas and anterior temporal lobe, but also parts of the control-network as well as subcentral and more posterior temporal-parietal areas. Activity in this supramodal sentence processing network starts in temporal areas and rapidly spreads to the other regions involved. The findings do not only indicate the involvement of a large network of brain areas in supramodal language processing, but also indicate that the linguistic information contained in the unfolding sentences modulates brain activity in a word-specific manner across subjects.

Introduction

Language can be realized in different modalities: amongst others through writing or speech. Depending on whether the sensory input modality is visual or auditory, different networks of brain areas are activated to derive meaning from the stimulus. Besides different brain circuits being recruited to process low-level sensory information, differences in linguistic features across sensory modalities prompt a differential activation of brain areas involved in higher-order processing as well. For instance, speech is enriched

with meaningful prosodic cues, but also requires coarticulated signals to be parsed into individual words. Written text has the advantage of instantaneous availability of full information compared to the temporally unfolding nature of speech. These differences are paralleled in the brain's response, and thus the sensory modality in which language stimuli are presented determines the dominant spatiotemporal patterns that will be elicited [19].

Regardless of low-level differences, the same core message can be conveyed in either modality. Therefore, language processing models of the past and present ([16], [18],[21]) include not only early sensory (up to 200ms) processing steps, but also contain late (200 - 500 ms), more abstract, and supposedly supramodal processing steps. While early processing is largely unimodal, and supported by brain regions in the respective primary and associative sensory areas, later processes (for instance lexical retrieval and integration) that activate several areas within the temporo-frontal language network are assumed to do so independent of modality.

In order to gain insight into the location and timing of brain processes representing this latter, higher order processing of the linguistic content, researchers so far relied on carefully manipulated experimental conditions. As a result, our current understanding of how the brain processes language across different modalities reflects a large variety in tasks (semantic decision task [10], error detection task [8], [11], passive hearing/listening [25], size judgment [31]) and stimulus material (words [10],sentences [3], and stories [5, 14, 38]). Despite this wealth of experimental findings and resulting insights, an important interpretational limitation stems from the fact that the majority of studies employ modality specific low-level baseline conditions (tone pairs and lines, spectrally rotated speech and false fonts, non-words, white noise [29]) to remove the sensory component of the processing. It is difficult to assess in how far such baselines are comparable across auditory and visual experiments. Recent fMRI work has demonstrated sensory-modality independent brain activity by directly comparing the BOLD response across visual and auditory presentations [14, 38]. Yet, fMRI signals lack the temporal resolution to allow for a temporally sufficiently fine-grained investigation of the response to individual words.

Few studies used magnetoencephalography (MEG) to study supramodal brain activity and all are based on event-related averaging [31, 3, 36, 47]. Averaged measures capture

only generic components in the neural response. While generic components make a large 40 contribution to the neural activity measured during language processing, there also exist 41 meaningful variability in the neural response that is stimulus-specific and robust [4]. A 42 complete analysis of the supramodal language network needs to tap into these subtle 43 variations as well. 44

Here, we overcome previous limitation by achieving a direct comparison without 45 relying on modality-specific baseline conditions, leveraging word-by-word variation in 46 the brain response. Using MEG signals from 200 subjects, we performed a quantitative 47 assessment of the sensory modality independent brain activity following word onset 48 during sentence processing. The MEG data forms part of a large publicly available 49 dataset [40], and has been used in other publications [28], [39], [24], [27]. We identified 50 widespread left hemispheric involvement, starting from 325 ms after word onset in 51 the temporal lobe and rapidly spreading to anterior areas. These findings provide a 52 quantitative confirmation of earlier findings in a large study sample. Importantly, they 53 also indicate that supramodal linguistic information conveyed by the individual words 54 in sentence context leads to subtle fluctuations in brain activation patterns that are 55 correlated across different subjects. 56

A Temporal alignment procedure

B Cross-validated multiset canonical correlation analysis

Figure 1. Analysis pipeline. Caption on next page

Figure 1. (Previous page.) (A) Temporal alignment procedure. MEG signals of auditory and visual subjects differed in length due to different presentation rates. To achieve alignment between signals of auditory and visual subjects, auditory signals were epoched into overlapping segments. Each segment's first sample corresponds to the auditory word onset but each segment's length depends on the duration of the equivalent visual stimulus. Segments were then concatenated in original order to recover signal for the full sentence length. This way, the neural response to each word is fully taken into account in further comparisons, including in the case of short words for which stimulus late processing coincided with the next word presentation. (B) Starting point for the multi-set canonical correlation analysis were parcel-based neural signals for all subjects, consisting of five spatial components each. 1. Signals for all sentence trials were split into five subsets and for cross-validation one subset of sentences was left out as test data, while the remaining four subsets served as training data. 2. Based on the training dataset only an unmixing matrix was found, per parcel, defining the linear combination of the five spatial components so that the correlation across sets (subjects) and time samples were maximized. The cross-covariance was computed between all subjects' spatial components and across time collapsing over sentence trials. 3. The projection was applied to the test data to compute canonical variables for the left out sentence trials (purple outline) for all subjects. Steps 2 and 3 were repeated for all folds until each sentence subset had been left out once and the resulting canonical variables were concatenated until the entire signal was transformed. 4. Canonical variables were epoched according to word onsets and for each time point a subject-by-subject correlation matrix was computed across words. Correlation between cross-modal subjects (pink outline) were interpreted as quantifying supramodal activation.

Materials and Methods 57

Subjects 58

A total of 204 native Dutch speakers (102 males), with an age range of 18–33 years 59

(mean of 22 years), participated in the experiment. In the current analysis, data from 60

200 subjects were included. Exclusion of four subjects was due to technical issues 61

during acquisition, which made their datasets not suitable for our analysis pipeline. All 62

subjects were right-handed, had normal or corrected-to-normal vision, and reported no 63

history of neurological, developmental, or language deficits. The study was approved by 64

the local ethics committee (CMO, the local "Committee on Research Involving Human 65

Participants" in the Arnhem–Nijmegen region) and followed the guidelines of the Helsinki 66

declaration. All subjects gave written informed consent before participation and received 67

monetary compensation for their participation. 68

Experimental Design

The subjects were seated comfortably in a magnetically shielded room and presented with Dutch sentences. From the total stimulus set of 360 sentences six subsets of 120 sentences were created. This resulted in six different groups of subjects who were presented with the same subset of stimuli although in a different (randomized) order with some overlap of items between groups. Within each group of subjects half of them performed the task in only the visual, the other half in only the auditory modality. In the visual modality, words were presented sequentially on a back-projection screen, placed in front of them (vertical refresh rate of 60 Hz) at the center of the screen within a visual angle of 4 degrees, in a black mono-spaced font, on a grey background. Each word was separated by an empty screen for 300 ms and the inter-sentence interval was jittered between 3200 and 4200 ms. Mean duration of words was 351 ms (minimum 300 ms and maximum 1400 ms), depending on word length. The median duration of whole sentences was 8.3 s (range 6.2 - 12 s). Auditory sentences had a median duration of 4.2 s (range 2.8 - 6.0 s) spoken in a natural pace. The duration of each visual word was determined by the following quantities: (i) the total duration of the audio-version of the sentence/word list (audiodur), (ii) the number of words in the sentence (nwords), (iii) the number of letters per word (nletters), and (iv) the total number of letters in the sentence (sumnletters). Specifically, the duration (in ms) of a single word was defined as: (nletters/ sumnletters) * (audiodur + 2000–150 * nwords). In the auditory task the stimuli were presented via plastic tubes and ear pieces to both ears. Before the experiment, the hearing threshold was determined individually and the stimuli were then presented at an intensity of 50 dB above the hearing threshold. A female native Dutch speaker recorded the auditory versions of the stimuli. The audio files were recorded in stereo at 44100 Hz. During the post processing the audio files were low-pass filtered at 8500 Hz and normalized so that all audio files had the same peak amplitude, and same peak intensity. All stimuli were presented using the Presentation software (Version 16.0, Neurobehavioral Systems, Inc). Sentences were presented in small blocks, of five sentences each, along with blocks containing scrambled sentences, which were not used here. See Lam et al. [28] for more details about the stimulus material used. In order to check for compliance, 20% of the trials were randomly followed by a yes/no question

about the content of the previous sentence/word list. Half of the questions addressed the 100
content of the sentence (e.g. Did grandma give a cookie to the girl?) whereas the other 101
half, addressed one of the main content words (e.g. Was the word 'grandma' mentioned?). 102
Subjects answered the question by pressing a button for 'Yes'/ 'No' with their left index 103
and middle finger, respectively. 104

MEG Data Acquisition & Structural imaging 105

MEG data were collected with a 275 axial gradiometer system (CTF). The signals were 106
analog low-pass-filtered at 300 Hz and digitized at a sampling frequency of 1,200 Hz. 107
The subject's head was registered to the MEG-sensor array using three coils attached to 108
the subject's head (nasion, and left and right ear canals). Throughout the measurement, 109
the head position was continuously monitored using custom software [44]. During breaks 110
the subject was allowed to reposition to the original position if needed. Participants 111
were able to maintain a head position within 5 mm of their original position. Three 112
bipolar Ag/AgCl electrode pairs were used to measure the horizontal and vertical 113
electrooculogram and the electrocardiogram. 114

A T1-weighted magnetization-prepared rapid gradient-echo (MP-RAGE) pulse se- 115
quence was used for the structural images, with the following parameters: volume TR = 116
2300 ms, TE = 3.03 ms, 8 degree flip-angle, 1 slab, slice-matrix size = 256 × 256, slice 117
thickness = 1 mm, field of view = 256 mm, isotropic voxel-size = $1.0 \times 1.0 \times 1.0$ mm. 118
A vitamin-E capsule was placed as fiducial behind the right ear to allow a visual identifi- 119
cation of left-right consistency. 120

Preprocessing 121

Data were bandpass filtered between 0.5 and 20 Hz, and epoched according to sentence 122
onset, each epoch varying in length, depending on the number of words within each 123
sentence. Samples contaminated by artifacts due to eye movements, muscular activity, 124
and superconducting quantum interference device jumps were replaced by NaN before 125
further analysis. Since all sentences had been presented in random order, we reordered 126
sentences for each subject to yield the same order across subjects. Subsequently, the sig- 127
nals of the auditory subjects were temporally aligned to the signals of the visual subjects, 128

ensuring coincidence of the onset of the individual words across modalities (Figure 1A). 129
This alignment was needed to accommodate for differences in word presentation rate. 130
The alignment was achieved by first epoching the auditory subject's signals into smaller 131
overlapping segments. Each segment's first sample corresponded to one of the word 132
onsets as annotated manually according to the audio file while each segment's length 133
depended on the duration of the visual presentation of the corresponding word. Finally, 134
all segments were concatenated again in the original order. By defining segments that 135
were longer than the corresponding auditory word duration, the neural response to each 136
word is fully taken into account and matched to the visual signal, even in the case of 137
short words where the response partly coincided with the next word presentation. MEG 138
data were then downsampled to 120 Hz. 139

Source Reconstruction 140

We used linearly constrained minimum variance beamforming (LCMV) [46] to reconstruct 141
activity onto a parcellated cortically constrained source model. For this, we computed 142
the covariance matrix between all MEG-sensor pairs, as the average covariance matrix 143
across the cleaned single trial covariance estimates. This covariance matrix was used 144
in combination with the forward model, defined on a set of 8,196 locations on the 145
subject-specific reconstruction of the cortical sheet to generate a set of spatial filters, one 146
filter per dipole location. Individual cortical sheets were generated with the Freesurfer 147
package [12, version 5.1] (surfer.nmr.mgh.harvard.edu), coregistered to a template with 148
a surface-based coregistration approach, using Caret software [45] (download here 149
and here), and subsequently downsampled to 8,196 nodes, using the MNE software 150
[17] (martinos.org/mne/stable/index.html)). The forward model was computed using 151
FieldTrip's singleshell method [34], where the required brain/skull boundary was obtained 152
from the subject-specific T1-weighted anatomical images. We further reduced the 153
dimensionality of the data to 191 parcels per hemisphere [39]. For each parcel, we 154
obtained a parcel-specific spatial filter as follows: We concatenated the spatial filters 155
of the dipoles comprising the parcel, and used the concatenated spatial filter to obtain 156
a set of time courses of the reconstructed signal at each parcel. Next, we performed a 157
principal component analysis and selected for each parcel the first five spatial components 158

explaining most of the variance in the signal.

Multi-set Canonical Correlation Analysis

Multi-set canonical correlation analysis (MCCA) [37], [13] was applied to find projections
of those five spatial components that would transform the subject-specific signals so as
to boost similarities between them. Canonical correlation analysis (CCA) is a standard
multivariate statistical method often used to investigate underlying relationships between
two sets of variables. Classically, canonical variates are estimated by transforming the
two sets in a way that optimizes their correlation. We applied a generalized version of
the classical approach (MCCA) [26], which extends the method to multiple sets, here
multiple subjects. In our case, we find linear combinations of the five spatial components
for each of two subjects, so that the correlation across time between those subjects is
maximized. Since we have more than two subjects, we find for each subject its own linear
combination, which maximizes the correlation across time between all subjects from
both modality groups (auditory and visual stimulation). Following Parra we obtained
the optimal projection as the eigenvector with the largest eigenvalue of a square matrix
$D^{-1}R$, where R and D are square matrices:

$$
R = \begin{pmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,N} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,N} \\ \vdots & \vdots & \ddots & \vdots \\ a_{N,1} & a_{N,2} & \cdots & a_{N,N} \end{pmatrix}, D = \begin{pmatrix} a_{1,1} & 0 & \cdots & 0 \\ 0 & a_{2,2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{N,N} \end{pmatrix}
$$

Where a^{lk} are cross-covariance matrices between subject pairs and D contains only
the diagonal blocks of within subject covariances [37]. In our case the cross-covariance
matrices are of size 5-by-5 containing the cross-covariance between all five spatial
components for a given subject (pair). The cross-correlation is computed across time
points for each sentence and subsequently averaged across sentences. It is important to
note here, that the canonical variates resulting from the optimal projection do not reflect
sentence averages anymore but have the same temporal resolution as the original source
signals. CCA is prone to overfitting and known to be unstable [15]. For reliable CCA
estimates the number of samples should be much larger than the number of features,
i.a. a sample-to-feature ratio of 20/1 is recommended [43]. We estimated the canonical

variables over concatenated data, which included between 756 and 1453 samples per ₁₈₆ sentence compared to only five features (spatial components) which provides a decent ₁₈₇ sample-to-feature ratio. Further, we estimated our canonical variables out-of sample ₁₈₈ using 5-fold cross-validation to limit overfitting. We randomly split all sentences into ₁₈₉ five subsets, estimating projections on 96 sentences and applying them to the 24 left out ₁₉₀ sentences (Figure 1B). ₁₉₁

Statistical Analysis

As per the study design, the subjects were assigned to one of six stimulus sets. Different ₁₉₃ groups of subjects were presented with different sets of sentences. Since MCCA relies on ₁₉₄ commonalities across datasets, we could only combine data from subjects who received the ₁₉₅ exact same stimulation. We therefore applied MCCA for each subgroup of subjects who ₁₉₆ listened to or saw the same stimuli separately. Initially, we constrained our analysis to the ₁₉₇ first set of 33 subjects (henceforth exploratory dataset). After applying the projection to ₁₉₈ the data we computed a time-resolved Pearson correlation between all possible subject ₁₉₉ pairings. To this end, we first epoched the resulting canonical components according ₂₀₀ to individual word onsets and selected only content words (nouns, adjectives & verbs) ₂₀₁ for subsequent steps. Before computing the correlation we subtracted the mean across ₂₀₂ samples. For each pair of subjects, we computed the correlation between two sets of ₂₀₃ observations, i.e. a pair of vectors with each data point reflecting the subject-specific ₂₀₄ neural signal for each of the individual words (lexical items), at a given time point relative ₂₀₅ to word onset, and at a given cortical location. Correlation coefficients of cross-modality ₂₀₆ pairings, that is correlations between subjects reading and subjects listening to the ₂₀₇ sentences are interpreted as capturing supramodal processing. We used a permutation ₂₀₈ test with clustering over time and space (parcels) for family-wise error rate correction ₂₀₉ for statistical inference (Maris and Oostenveld, 2007), using 1000 randomizations of the ₂₁₀ epoched words. To this end, we randomized word order for the source-reconstructed ₂₁₁ parcel time series of the auditory subjects to test for exchangeability of the exact word ₂₁₂ pairing across sensory modalities. By destroying the one-to-one mapping of individual ₂₁₃ lexical items, the null distribution allowed for a distinction between individual item ₂₁₄ specific shared variance, and shared variance due to a more generic response. For a more ₂₁₅

conservative test, we additionally tested the observed correlation patterns against a null ²¹⁶
distribution obtained by permuting sentence order and doing this prior to MCCA. The ²¹⁷
results and a description of this additional test can be found in supporting Figure 3-1. ²¹⁸

We then analyzed the remaining sets of subjects (confirmatory dataset) using the ²¹⁹
same analysis pipeline. We evaluated the overlap in the results across all six subgroups ²²⁰
using information prevalence inference [1]. Prevalence inference allows to formulate a ²²¹
complex null hypothesis, i.e. that the prevalence of the effect is smaller than or equal ²²²
to a threshold prevalence, where the threshold can be realized by different values. For ²²³
each of the six sets of data, we obtained spatial maps of time-resolved supramodal ²²⁴
correlations, as well as 1000 permutation estimates after word order shuffling (see above). ²²⁵
We used the smallest observed average correlation across subgroups as the second-level ²²⁶
test statistic. We then tested the majority null hypothesis of the prevalence of the effect ²²⁷
being smaller or equal to a threshold prevalence. For this, we computed the largest ²²⁸
threshold such that the corresponding null hypothesis could still be rejected at the given ²²⁹
significance level α. This was done after concatenating the minimum statistic from all ²³⁰
parcels and time points, using the maximum statistic to correct for multiple comparisons ²³¹
in time and space (parcels). For each parcel we evaluated the highest threshold at which ²³²
the prevalence null hypothesis could be rejected at a level of $\alpha = 0.05$ (see supporting ²³³
Figures 4-1 and 4-2 for cortical maps showing thresholds averaged and per time point). ²³⁴

Code Accessibility ²³⁵

All analyses were done with custom-written MATLAB scripts and FieldTrip [35] and ²³⁶
the corresponding code is available upon request. ²³⁷

Results ²³⁸

Modality-specific activation ²³⁹

We first quantified the similarity between different subjects' brain response within ²⁴⁰
only the exploratory dataset (33 subjects) by correlating word-by-word fluctuations in ²⁴¹
brain activity between all possible pairs of subjects. Averaging the correlations across ²⁴²
those subject pairings for which subjects were stimulated either in the same sensory ²⁴³

modality, or each in a different modality, allowed us to evaluate the modality-specific ₂₄₄ brain response and the supramodal response, respectively. As displayed in Figure 2, early ₂₄₅ sensory cortical areas only show correlated activity for the group of subjects receiving ₂₄₆ the stimuli in the corresponding sensory modality, for the visual (red), and auditory ₂₄₇ (blue) modalities. We found that MCCA is a crucial analysis step in order to reveal ₂₄₈ meaningful inter-subject correlations. Only after MCCA does cortical activity in visual ₂₄₉ and auditory areas become significantly correlated across those subjects performing the ₂₅₀ task in the visual or auditory domain respectively (Figure 2A). ₂₅₁

Supramodal activation patterns

We averaged between-subject correlations over all cross-modal subject pairings as a ₂₅₃ metric for supramodal activity. We observed significant supramodal correlated activation ₂₅₄ patterns in mostly left-lateralized cortical areas (Figure 3). The effect has a large spatial ₂₅₅ and temporal extent, becoming apparent as early as 250 ms and lasting until 700 ms ₂₅₆ after word onset. Parcels in middle superior temporal gyrus (STG) contribute to the ₂₅₇ effect at the earliest time points, followed by the posterior and anterior part of the STG ₂₅₈ and about 50 ms later the anterior temporal pole. Supramodal correlated activation in ₂₅₉ ventral temporal cortex follows a similar temporal and spatial pattern, with supramodal ₂₆₀ correlations starting out more posterior around 292 ms and evolution towards the middle ₂₆₁ anterior temporal lobe at 308 ms. Other areas that express supramodal activity at ₂₆₂ relatively early time points are medial prefrontal cortex and primary auditory cortex ₂₆₃ (250 ms), followed by subcentral parietal regions and supramarginal gyrus at around 300 ₂₆₄ ms, and finally dorsolateral frontal cortex (DLFC, 325 ms). By the time 375 ms have ₂₆₅ passed, the entire orbito-frontal cortex, anterior and DLFC as well as inferior frontal ₂₆₆ gyrus (IFG) show strong supramodal subject correlation. Supramodal activation in the ₂₆₇ frontal lobe further extends towards posterior regions including pre- and postcentral ₂₆₈ gyrus. At around 400 ms supramodal subject correlation in the anterior temporal pole ₂₆₉ reaches its peak. In addition to the lateral cortical areas, correlated activity also extends ₂₇₀ to left dorsal and ventral anterior cingulate cortex (ACC) as well as left fusiform gyrus. ₂₇₁ The spatio-temporal patterns of supramodal activation described so far are robust, also ₂₇₂ when ordinal word position and MCCA overfitting is controlled for in the statistical ₂₇₃

Prevalence Inference   275

Our confirmatory analysis combined over all six datasets and tested whether the spa-   276
tiotemporal patterns observed in the exploratory dataset would generalize to the pop-   277
ulation. For those parcels at which the global null hypothesis could be rejected, we   278
infer that at least in one of the datasets an effect of supramodal processing was present   279
(Figure 4). In addition, we evaluated the majority null hypothesis of whether, in the   280
majority of subgroups in the population, the data contains an effect (threshold > 0.5,   281
significant parcels under the majority null hypothesis outlined in black in figure 4B).   282

The global null hypothesis (no information in any set of subjects in the population)   283
could be rejected at a level of $\alpha = 0.05$ in on average 40 parcels per time point (between   284
325 and 617 ms after onset, std. $= 31.48$). For those parcels, the average largest lower   285
bound $\gamma 0$ at which the prevalence null hypothesis can be rejected is shown in supporting   286
Figures 4-1 and 4-2. For those parcels for which the largest bound $\gamma 0$ is larger than or   287
equal to 0.5 we can infer that in the majority of the datasets the activity patterns were   288
similar across subjects independent of modality.   289

This majority null hypothesis could be rejected (at a level of $\alpha = 0.05$) in 90% of   290
parcels that also showed a global effect (Figure 4, see supporting Figure 4-3 for results   291
in the right medial hemisphere). Compared to the temporal pattern of the largest   292
nominal suprathreshold cluster from the cluster-based permutation test conducted on   293
the exploratory dataset, the effect became significant in the majority of datasets later   294
in time and was less long lasting (325 - 608 ms). Given this time span, the majority   295
null hypothesis was rejected in on average 42% of those parcels contributing most   296
to the largest cluster. The orbitofrontal cortex and IFG showed an involvement in   297
supramodal processing in both analyses, but the effect there was much more temporally   298
sustained in the exploratory dataset. In addition, according to the exploratory dataset,   299
supramodal activation of STG occurred almost 100 ms earlier as compared to IFG.   300
Based on the confirmatory dataset, however, supramodal correlated activation in IFG   301
and STG appeared almost simultaneously. Finally, the exploratory analysis revealed   302
supramodal activation in primary and premotor areas extending over the entire left   303

dorsolateral surface, of which only the most ventral parcels close to the Sylvian fissure ³⁰⁴ were significantly supramodal in the majority of datasets. Thus, the spatial extent ³⁰⁵ of the effect was partly reduced for prevalence inference, compared to the cluster- ³⁰⁶ based permutation approach on the exploratory data. Nevertheless, widely overlapping ³⁰⁷ anatomical regions were indicated by both analyses, encompassing dorsolateral frontal ³⁰⁸ gyrus and the middle & superior parts of the temporal lobe at first and later also inferior ³⁰⁹ frontal and orbito-frontal cortex, as well as anterior temporal lobe. ³¹⁰

Figure 2. Specificity of the within-modality correlated activity patterns. (A) Time-resolved correlation values averaged across all visual subject pairings for a parcel in left primary visual cortex (upper panel) and all auditory subject pairings for a parcel in left primary auditory cortex (lower panel), before (dark grey line), and after MCCA (blue and red lines). Light grey lines show recomputed correlation values for 1000 random permutations of word order across subjects. Notably, signals of auditory subjects highly correlate even before word onset. This is likely due to a more varied distribution of information in the auditory signal caused by the continuous nature of auditory stimulation and as a result differing time points at which individual words become uniquely recognizable. The MCCA is blind to the stimulus timing and will thus find canonical variables that yield maximal correlations at any timepoint if possible. (B) Cortical map of the spatial distribution of correlations, comparing within modality visual subject pairs (red) with auditory subject pairs (blue). Correlation strength is expressed as the Pearson correlation coefficient averaged over a time window from 150 to 200 ms post word-onset and normalized by the maximum value of that window.

Figure 3. Supramodal correlated activity patterns. Time-resolved spatial maps of supramodal correlated activity patterns (averaged over all possible cross-modal subject pairings) in the left hemisphere. Medial views of the brain surface are depicted in the first and third row, lateral views in the second and fourth row. Color codes for strength of correlation. Colored parcels were most strongly correlated between cross-modal subject pairs (nonparametric permutation test, corrected for multiple comparisons).

Figure 4. Supramodal correlated activity patterns consistent across the majority of datasets. Caption on next page

Figure 4. (Previous page.) Supramodal correlated activity patterns of word-specific activity consistent across the majority of datasets. (A) Averaged correlation time courses (mean over all possible cross-modal subject pairings) are shown for selected parcels in inferior frontal Gyrus (IFG, green), supramarginal Gyrus (red), Subcentral Cortex (orange), Anterior cingulate cortex (ACC, pink), anterior middle temporal Gyrus (aMTG, purple), and middle superior temporal gyrus (mSTG, blue). Time courses are shown for each dataset individually (light-colored lines) as well as averaged (dark lines). Grey shaded areas mark statistically significant time points. (B) Time-resolved spatial maps of cross-modal correlations in the left hemisphere. Medial views of the brain surface are depicted in the first, third and fifth row, lateral views in the second, fourth and sixth row. For those parcels that were part of the largest nominal suprathreshold cluster tested on only the exploratory dataset, the mean correlation over all six datasets is shown. Color codes for strength of correlation. In addition, the parcels at which the majority null hypothesis according to prevalence inference could be rejected are outlined in black.

Discussion

Our aim was to quantify similarities of the brain response across reading and listening at a fine temporal scale. To this end we correlated word-by-word fluctuations in the neural activity across subjects receiving either auditory or visual stimulation. We identified a widespread left-lateralized brain network, activated independently of modality starting 325 ms after word onset. Importantly, dividing our large study sample into six subsets, we could directly quantify the consistency and generalizability of these activity patterns. The spatial distribution of the supramodal activation is in line with the known involvement of left hemispheric areas, including parts of left temporal cortex, left inferior parietal lobe, as well as prefrontal cortex [48], [10], [11], [7], [30], [29], [42], [23]. The involvement of both STG and IFG fits predictions from the Memory, Unification and Control model (MUC), in which activity reverberating within a posterior-frontal network ([2], [18]) is thought to be crucial for language processing. According to the MUC model temporal and parietal areas support the retrieval of lexical information, while unification processes are supported by inferior frontal cortex. Bidirectional communication [39] between these areas is facilitated by white matter connections. We observe that temporal areas are supramodally activated at earliest time points and sustain activation for the longest compared to other regions. Over time, supramodal activation spreads from middle and posterior left STG to the anterior temporal pole. This rapid progression of activity from posterior to anterior regions mirrors previous observations ([31], [47]), adding to those findings a direct quantitative comparison of the supramodal brain activity.

Beyond the core language network and the single word level

We observed modality-independent activity in dorsal frontal cortex, in addition to more widely reported inferior parts of the frontal cortex [25], [29], [11], [31], [23], [33]. This could be due to us using linguistically rich sentence material, of varying syntactic complexity, as opposed to single words [10],[31],[6],[30],[47] or short phrases [3],[8],[7]. Indeed, discrepancies with respect to frontal lobe involvement in modality-independent processing seem to mainly arise from differences in stimulus material and task demands [7]. A recent meta-analysis has identified that more complex syntax robustly activates dorsal parts of the left IFG [20]. Further, a previously published analysis of these MEG data showed DLFC to be sensitive to sentence progression effects [24]. Two previous fMRI studies using narratives [14, 38] add to the debate. Regev et al. correlated BOLD responses evoked by different modalities. They report supramodal activation in the left frontal lobe, extending beyond inferior frontal regions. Deniz et al. study modality-independent brain areas by modelling semantic features of the stimulus in one modality and used the model to predict the BOLD signal in the other modality. They report BOLD signals in prefrontal cortex to be well predicted across modalities. In summary, while complex stimuli consistently activate prefrontal areas beyond inferior frontal cortex, the exact stimulus features which cause this supramodal activation are still debated.

Some previous studies, using narratives and fMRI, report supramodal activation not to be restricted to the left hemisphere [14, 38, 25]. It could be that the previously observed bilateral involvement is due to differences in context-based semantic processing during narratives, as compared to the processing of isolated sentences in our experiment. Menteni and colleagues have specifically contrasted BOLD activity in response to sentences presented within a neutral or a local context. The authors indeed reported right frontal cortex to be more sensitive to local discourse context as compared to its left-hemispheric homolog [32]. Further research is needed to determine whether this effect of presence and absence of narrative thread similarly affects lateralisation of brain activity in MEG.

Even though mostly restricted to the left hemisphere, our results also implicate extra-linguistic areas in supramodal processing. Specifically, we find bilateral supramodal activation within ACC. The ACC is a midline structure, forming part of a domain-general

executive control network supporting language processing [18], [9]. It is sensitive to statistical contingencies in the language input and thus might play a role in mediating learning and adaptation in response to predictive regularities in both local experimental as well as global environment [49]. It should be noted that deep sources are normally poorly detectable in MEG [22] and we thus consider any interpretations with respect to the midline structures as tentative.

Supramodal orthography-phonology mapping

We observed supramodal activation in post-central and subcentral gyrus, as well as supramarginal gyrus, which coincides temporally with supramodal activation of primary auditory cortex. Activity in supramarginal gyrus has been repeatedly elicited by cross-modal tasks [41], such as rhyming judgments to visually presented words [6], for which conversion between orthographic and phonological representations is likely needed. At the same time, post- and subcentral areas partly span articulatory motor and somatosensory areas for the mouth and tongue. Together, the supramodal activation of these areas suggests that retrieval of phonetic and articulatory mappings is not limited to speech perception only but also occurs during passive reading.

Leveraging word-by-word variability of the neural response

Neuroelectric brain signals exhibit strong moment-to-moment variability. While some of this variability is related to the experimental stimulation, and therefore associated with specific cognitive activity, some of it is unrelated, ongoing neural activity. By applying MCCA across subjects we reduced this type of noise and made subtle word-by-word fluctuations in the MEG signal interpretable. Comparing neural activity across subjects is challenging due to differing position or orientation of neuronal sources relative to the MEG sensors. We used parcellated MEG source reconstruction in combination with exact temporal alignment of individual sentences across subjects. This allowed for the extraction of signal components that are shared across subjects, thus reducing the intersubject spatial variability, which is commonly observed in more traditional (for instance, dipole fitting) procedures [47]. MCCA thus allowed us to more directly investigate time-resolved inter-subject correlations and move beyond event-

related averages [31]. Importantly, our analysis approach allows us to conclude that ₃₉₂ the identified supramodal activity is word-specific. Our findings therefore go beyond ₃₉₃ showing a general activation of those areas as compared to baseline but rather reveal ₃₉₄ consistent word-by-word fluctuations of activation within the recruited areas. ₃₉₅

Latency of supramodal processing ₃₉₆

The temporal alignment procedure, as a necessary preparation step for the MCCA ₃₉₇ procedure, followed by the estimation of time-resolved intersubject correlations, focused ₃₉₈ on common signal aspects that are exactly synchronized across subjects. The differences ₃₉₉ in sensory modality specific characteristics of the input signal require dedicated processing ₄₀₀ with likely different processing latencies, which may also lead to latency differences in the ₄₀₁ activation of supramodal areas. For example, Marinkovic and colleagues report shorter ₄₀₂ reaction times during the visual task, yet found earlier activity peaks for the auditory task ₄₀₃ in corresponding early sensory cortex and left anterior temporal lobe [31]. In contrast, ₄₀₄ other work observed earlier anterior temporal lobe activation for visual, compared to ₄₀₅ auditory stimulation [3]. Our results indicate a certain degree of overlap across modalities ₄₀₆ in the temporal window within which supramodal cortical areas are activated. It is ₄₀₇ possible, that we observed more temporally extensive activation, for instance, related to ₄₀₈ unification processes, because we used longer sentences. In addition, any overlap may ₄₀₉ have been amplified as a necessary consequence of the MCCA procedure. Evidently, ₄₁₀ correlations between signals from auditory subjects were boosted with less temporal ₄₁₁ specificity compared to visual subjects (Figure 2B). This observation was unexpected ₄₁₂ and may be due to more continuous stimulation in the auditory experiment. As the ₄₁₃ sound of a spoken word unfolds, the timing at which it becomes uniquely recognizable ₄₁₄ will vary across word. Thus, the distribution of information in the auditory signal is ₄₁₅ much more varied as compared to the visual. MCCA will pick up on any common ₄₁₆ relationship across subjects regardless of timing. In our specific application, projections ₄₁₇ were estimated on concatenated data, effectively making the method blind to word onset ₄₁₈ boundaries. ₄₁₉

In conclusion, this study provides direct neurophysiological evidence for sensory ₄₂₀ modality independent processes supporting language comprehension in multiple left ₄₂₁

hemispheric brain areas. We identified a network of areas including domain general 422
control areas as well as phonological mapping circuits over and above traditional higher- 423
level language areas in frontal and temporal-parietal regions, by quantifying between- 424
subject consistency of their respective word-specific activation patterns. These consistent 425
activation patterns were word-specific, and thus likely reflect more than just generic 426
activation during language processing. Finally, we show that alignment of individual 427
subject data through MCCA is a promising tool for investigating subtle word-in-context 428
specific modulations of brain activity in the language system. 429

Acknowledgments

We thank Phillip Alday for providing helpful comments. This work was supported by The Netherlands Organisation for Scientific Research (NWO Vidi: 864.14.011, awarded to J.M.S.).

References

1. C. Allefeld, K. Görgen, and J. D. Haynes. Valid population inference for information-based imaging: From the second-level t-test to prevalence inference. *NeuroImage*, 141:378–392, 2016.

2. G. Baggio and P. Hagoort. The balance between memory and unification in semantics: A dynamic account of the N400. *Language and Cognitive Processes*, 26(9):1338–1367, 2011.

3. D. K. Bemis and L. Pylkkänen. Basic linguistic composition recruits the left anterior temporal lobe and left angular gyrus during both listening and reading. *Cerebral Cortex*, 23(8):1859–1873, 2013.

4. A. Ben-Yakov, C. J. Honey, Y. Lerner, and U. Hasson. Loss of reliable temporal structure in event-related averaging of naturalistic stimuli. *NeuroImage*, 63(1):501 – 506, 2012.

5. M. M. Berl, E. S. Duke, J. Mayo, L. R. Rosenberger, E. N. Moore, J. VanMeter, N. B. Ratner, C. J. Vaidya, and W. D. Gaillard. Functional anatomy of listening and reading comprehension during development. *Brain and Language*, 114(2):115–125, 2010.

6. J. R. Booth, D. D. Burman, J. R. Meyer, D. R. Gitelman, T. B. Parrish, and M. M. Mesulam. Modality independence of word comprehension. *Human Brain Mapping*, 16(4):251–261, 2002.

7. D. Braze, W. E. Mencl, W. Tabor, K. R. Pugh, R. Todd Constable, R. K. Fulbright, J. S. Magnuson, J. A. Van Dyke, and D. P. Shankweiler. Unification of sentence processing via ear and eye: An fMRI study. *Cortex*, 47(4):416–431, 2011.

8. A. Carpentier, K. R. Pugh, M. Westerveld, C. Studholme, O. Skrinjar, J. L. Thompson, D. D. Spencer, and R. T. Constable. Functional MRI of language processing: Dependence on input modality and temporal lobe epilepsy. *Epilepsia*, 42(10):1241–1254, 2001.

9. I. Cattinelli, N. A. Borghese, M. Gallucci, and E. Paulesu. Reading the reading brain: A new meta-analysis of functional imaging data on reading. *Journal of Neurolinguistics*, 26(1):214–238, 2013.

10. M. W. L. Chee, K. M. O'Craven, R. Bergida, B. R. Rosen, and R. L. Savoy. Auditory and visual word processing studied with fMRI. *Human Brain Mapping*, 7(1):15–28, 1999.

11. R. T. Constable, K. R. Pugh, E. Berroya, W. E. Mencl, M. Westerveld, W. Ni, and D. Shankweiler. Sentence complexity and input modality effects in sentence comprehension: An fMRI study. *NeuroImage*, 22(1):11–21, 2004.

12. A. M. Dale, B. Fischl, and M. I. Sereno. Cortical surface-based analysis: I. segmentation and surface reconstruction. *NeuroImage*, 9(2):179 – 194, 1999.

13. A. de Cheveigné, G. M. Di Liberto, D. Arzounian, D. D. Wong, J. Hjortkjær, S. Fuglsang, and L. C. Parra. Multiway canonical correlation analysis of brain data. *NeuroImage*, 186:728–740, 2018.

14. F. Deniz, A. O. Nunez-Elizalde, A. G. Huth, and J. L. Gallant. The Representation of Semantic Information Across Human Cerebral Cortex During Listening Versus Reading Is Invariant to Stimulus Modality. *The Journal of Neuroscience*, 39(39):7722–7736, 2019.

15. R. Dinga, L. Schmaal, B. W. Penninx, M. J. van Tol, D. J. Veltman, L. van Velzen, M. Mennes, N. J. van der Wee, and A. F. Marquand. Evaluating the evidence for biotypes of depression: Methodological replication and extension of Drysdale et al. (2017). *NeuroImage: Clinical*, 22:101796, 2019.

16. N. Geschwind. Specializations of the human brain. *Scientific American*, 2(241):180–201, 1979.

17. A. Gramfort, M. Luessi, E. Larson, D. A. Engemann, D. Strohmeier, C. Brodbeck, L. Parkkonen, and M. S. Hämäläinen. Mne software for processing meg and eeg data. *NeuroImage*, 86:446 – 460, 2014.

18. P. Hagoort. The core and beyond in the language-ready brain. *Neuroscience and Biobehavioral Reviews*, 81:194–204, 2017.

19. P. Hagoort and C. M. Brown. ERP effects of listening to speech compared to reading: the P600 to syntactic visual presentation. *Neuropsychologia*, 38:1531–1549, 2000.

20. P. Hagoort and P. Indefrey. The Neurobiology of Language Beyond Single Words. *Annual Review of Neuroscience*, 37(1):347–362, 2014.

21. G. Hickok and D. Poeppel. The Cortical Organisation fo Speech Processing. *Nature*, 8(5):393–402, 2007.

22. A. Hillebrand and G. R. Barnes. A quantitative assessment of the sensitivity of whole-head MEG to activity in the adult human cortex. *NeuroImage*, 16(3A):638–650, 2002.

23. F. Homae, R. Hashimoto, K. Nakajima, Y. Miyashita, and K. L. Sakai. From perception to sentence comprehension: The convergence of auditory and visual information of language in the left inferior frontal cortex. *NeuroImage*, 16(4):883–900, 2002.

24. A. Hultén, J.-M. Schoffelen, J. Uddén, N. H. Lam, and P. Hagoort. How the brain makes sense beyond the processing of single words – An MEG study. *NeuroImage*, 186:586–594, 2019.

25. G. Jobard, M. Vigneau, B. Mazoyer, and N. Tzourio-Mazoyer. Impact of modality and linguistic complexity during reading and listening tasks. *NeuroImage*, 34(2):784–800, 2007.

26. J. R. Kettenring. Canonical analysis of several sets of variables. *Biometrika*, 58(3):433–451, 1971.

27. N. H. Lam, A. Hultén, P. Hagoort, and J. M. Schoffelen. Robust neuronal oscillatory entrainment to speech displays individual variation in lateralisation. *Language, Cognition and Neuroscience*, 33(8):943–954, 2018.

28. N. H. Lam, J.-M. Schoffelen, J. Uddén, A. Hultén, and P. Hagoort. Neural activity during sentence processing as reflected in theta, alpha, beta, and gamma oscillations. *NeuroImage*, 142:43–54, 2016.

29. R. Lindenberg and L. Scheef. Supramodal language comprehension: Role of the left temporal lobe for listening and reading. *Neuropsychologia*, 45(10):2407–2415, 2007.

30. A. G. Liuzzi, R. Bruffaerts, R. Peeters, K. Adamczuk, E. Keuleers, S. De Deyne, G. Storms, P. Dupont, and R. Vandenberghe. Cross-modal representation of spoken and written word meaning in left pars triangularis. *NeuroImage*, 150:292–307, 2017.

31. K. Marinkovic, R. P. Dhond, A. M. Dale, M. Glessner, V. Carr, and E. Halgren. Spatiotemporal dynamics of modality-specific and supramodal word processing. *Neuron*, 38(3):487–497, 2003.

32. L. Menenti, K. M. Petersson, R. Scheeringa, and P. Hagoort. When elephants fly: Differential sensitivity of right and left inferior frontal gyri to discourse and world knowledge. *Journal of Cognitive Neuroscience*, 21(12):2358–2368, 2009. PMID: 19016600.

33. E. Michael, T. A. Keller, P. Carpenter, and M. A. Just. fMRI investigation of sentence comprehension by eye and by ear: Modality fingerprints on cognitive processes. *Human Brain Mapping*, 13(4):239–252, 2001.

34. G. Nolte. The magnetic lead field theorem in the quasi-static approximation and its use for magnetoenchephalography forward calculation in realistic volume conductors. *Physics in Medicine and Biology*, 48(22):3637–3652, 2003.

35. R. Oostenveld, P. Fries, E. Maris, and J. M. Schoffelen. FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, 2011:1–9, 2011.

36. A. C. Papanicolaou, M. Kilintari, R. Rezaie, S. Narayana, and A. Babajani-Feremi. The Role of the Primary Sensory Cortices in Early Language Processing. *Journal of Cognitive Neuroscience*, 29(10), 2017.

37. L. C. Parra. Multi-set Canonical Correlation Analysis simply explained. *arXiv:1802.03759*, (11 February 2018), 2018.

38. M. Regev, C. J. Honey, E. Simony, and U. Hasson. Selective and invariant neural responses to spoken and written narratives. *Journal of Neuroscience*, 33(40):15978–15988, 2013.

39. J.-M. Schoffelen, A. Hultén, N. Lam, A. F. Marquand, J. Uddén, and P. Hagoort. Frequency-specific directed interactions in the human brain network for language. *Proceedings of the National Academy of Sciences*, 114(30):8083–8088, 2017.

40. J.-M. Schoffelen, R. Oostenveld, N. H. L. Lam, J. Uddén, A. Hultén, and P. Hagoort. A 204-subject multimodal neuroimaging dataset to study language processing. *Scientific Data*, 6(1):1–17, 2019.

41. M. W. Sliwinska, M. Khadilkar, J. Campbell-Ratcliffe, F. Quevenco, and J. T. Devlin. Early and sustained supramarginal gyrus contributions to phonological processing. *Frontiers in Psychology*, 3:1–10, 2012.

42. G. Spitsyna, J. E. Warren, S. K. Scott, F. E. Turkheimer, and R. J. S. Wise. Converging Language Streams in the Human Temporal Lobe. *Journal of Neuroscience*, 26(28):7328–7336, 2006.

43. J. P. Stevens. Canonical Correlation. In *Applied multivariate statistics for the social sciences*, pages 395–412. Routledge, 2012.

44. A. Stolk, A. Todorovic, J. M. Schoffelen, and R. Oostenveld. Online and offline tools for head movement compensation in MEG. *NeuroImage*, 68:39–48, 2013.

45. D. C. Van Essen, H. A. Drury, J. Dickson, J. Harwell, D. Hanlon, and C. H. Anderson. An Integrated Software Suite for Surface-based Analyses of Cerebral Cortex. *Journal of the American Medical Informatics Association*, 8(5):443–459, 09 2001.

46. B. D. Van Veen, W. van Drongelen, M. Yuchtman, and A. Suzuki. Localization of brain electrical activity via linearly constrained minimum variance spatial filtering. *IEEE Transactions on Biomedical Engineering*, 44(9):867–880, 1997.

47. J. Vartiainen, T. Parviainen, and R. Salmelin. Spatiotemporal Convergence of Semantic Processing in Reading and Speech Perception. *Journal of Neuroscience*, 29(29):9271–9280, 2009.

48. M. Vigneau, V. Beaucousin, B. Mazoyer, M. Vigneau, N. Tzourio-Mazoyer, L. Petit, G. Jobard, P.-Y. Hervé, F. Crivello, L. Zago, and E. Mellet. What is right-hemisphere contribution to phonological, lexico-semantic, and sentence processing? *NeuroImage*, 54(1):577–593, 2010.

49. K. Weber, E. F. Lau, B. Stillerman, and G. R. Kuperberg. The Yin and the Yang of prediction: An fMRI study of semantic predictive processing. *PLoS ONE*, 11(3):1–25, 2016.

Figure 3-1. Figure 3-1. Significant supramodal correlated activity patterns as assessed by additional permutation test.

In an additional significance test, we shuffled the sentence order 500 times prior to MCCA, controlling for the possibility that MCCA as a preprocessing step may artificially increase correlations between subjects (through overfitting). Importantly, this permutation was not fully unconstrained, since we aimed at aligning sentences across modalities with the same number of words, to avoid loss of data and to preserve ordinal word position. Thus, we did a random pairing between sentences with the same number of words, after binning the sentences according to their word count. Sentences consisting of 9, 14 or 15 words were infrequent, with fewer than 5 occurrences each. After each permutation, we performed the temporal alignment between sensory modalities (aligning the word onsets), followed by cross-validated MCCA and computation of the time-resolved correlations of crossmodal subject pairs. Due to the long computation time of the canonical variates, we created this null distribution for the exploratory data only. In the figure, color codes for strength of correlation. Colored parcels were most strongly correlated between cross-modal subject pairs (corrected for multiple comparisons).

Figure 4-1. Figure 4-1. Cortical map of maximum threshold $\gamma 0$.
For those parcels at which the global null hypothesis could be rejected, the mean (over time) maximum threshold is plotted, for which the null hypothesis can be rejected ($\alpha = 0.05$). Given the sample size of six datasets, the number of second-level permutations and a significance level of $\alpha = 0.05$ the maximally possible threshold that can be reached is 0.5633.

Figure 4-2. Figure 4-2. Cortical map of prevalence threshold $\gamma 0$.
For those parcels at which the global null hypothesis could be rejected, the maximum threshold is plotted, for which the null hypothesis can be rejected at a level of $\alpha = 0.05$.

Figure 4-3. Figure 4-3. Time-resolved spatial maps of cross-modal correlations for the right hemisphere.
The average correlation over all six datasets is shown. Color codes for strength of correlation. In addition, the parcels at which the majority null hypothesis according to prevalence inference could be rejected are outlined in black.