High-throughput identification of MHC class I binding

peptides using an ultradense peptide array*

Amelia K. Haj[1], Meghan E. Breitbach[1], David A. Baker[1], Mariel S. Mohns[1], Gage K. Moreno[1],

Nancy A. Wilson[2], Victor Lyamichev[3], Jigar Patel[3], Kim L. Weisgrau[4], Dawn M. Dudley[1], David H.

O'Connor[1]

[1]Department of Pathology and Laboratory Medicine, University of Wisconsin-Madison

[2]Department of Medicine, University of Wisconsin-Madison

[3]Nimble Therapeutics, Madison, WI

[4]Wisconsin National Primate Research Center, Madison, WI

* This work was supported by R24OD017850 and National Library of Medicine Grant

5T15LM007359.

Abstract

Rational vaccine development and evaluation requires identifying and measuring the magnitude of epitope-specific CD8 T cell responses. However, conventional CD8 T cell epitope discovery methods are labor-intensive and do not scale well. Here, we accelerate this process by using an ultradense peptide array as a high-throughput tool for screening peptides to identify putative novel epitopes. In a single experiment, we directly assess the binding of four common Indian rhesus macaque MHC class I molecules – Mamu-A1*001, -A1*002, -B*008, and -B*017 – to approximately 61,000 8-mer, 9-mer, and 10-mer peptides derived from the full proteomes of 82 simian immunodeficiency virus (SIV) and simian-human immunodeficiency virus (SHIV) isolates. Many epitope-specific CD8 T cell responses restricted by these four MHC molecules have already been identified in SIVmac239, providing an ideal dataset for validating the array; up to 64% of these known epitopes are found in the top 192 SIVmac239 peptides with the most intense MHC binding signals in our experiment. To assess whether the peptide array identified putative novel CD8 T cell epitopes, we validated the method by IFN-γ ELISPOT assay and found three novel peptides that induced CD8 T cell responses in at least two Mamu-A1*001-positive animals; two of these were validated by *ex vivo* tetramer staining. This high-throughput identification of peptides that bind class I MHC will enable more efficient CD8 T cell response profiling for vaccine development, particularly for pathogens with complex proteomes where few epitope-specific responses have been defined.

Introduction

CD8 T cell receptors typically recognize epitopes – linear peptides of 8-11 amino acids – presented by major histocompatibility complex (MHC) class I molecules. These peptides are usually derived from self proteins that under normal circumstances do not induce an immune response. Cancerous cells, transplanted cells, and infected cells, however, present non-self peptides that are recognized as foreign by CD8 T cells (1). CD8 T cell recognition of foreign epitopes, including those derived from viral proteins, along with concurrent recognition of costimulatory molecules on the antigen-presenting cell, leads to T cell activation and production of cytotoxic compounds that trigger apoptosis in the affected cell (2). Polymorphisms in the peptide binding cleft of the MHC glycoprotein determine the variety of peptides each molecule is able to present, and specificity is determined in part by anchor residues in the peptide binding to specific pockets in the binding cleft (3). Identifying which peptides are presented by particular MHC molecules and are likely to elicit epitope-specific CD8 T cell responses is a critical task in designing prophylactic and therapeutic vaccines against infectious diseases and cancers (4, 5).

Characterizing MHC-peptide binding can be a time- and resource-intensive task. Competitive binding assays, which calculate the concentration (IC_{50}) at which a peptide of interest outcompetes 50% of a radiolabeled probe peptide for binding to MHC class I molecules, are used to estimate MHC-peptide binding affinity (6). These methods are low-throughput and difficult to scale. Computational tools have been developed that have the potential to improve throughput, enabling prediction of the likelihood a given peptide will bind to a specific MHC molecule. However, these methods exploit our understanding of anchor residues and rely on previously generated peptide binding datasets for pattern recognition, and so are limited to MHC molecules whose binding repertoires have already been well studied, creating a self-fulfilling prophecy of sorts (6, 7). A different computational technique, molecular docking, uses the three-dimensional

crystal structure of MHC molecules to predict peptides likely to bind (8, 9). While this method has

the benefit of working for both MHC class I and II molecules, it still cannot serve as a substitute

for *in vitro* binding assays.

CD8 T cell epitope identification is a similarly involved task, often requiring large quantities of cells

exposed *in vivo* to the pathogen of interest. In our previous work identifying CD8 T cell epitopes,

we performed IFN-γ ELISPOTs using pools of 15-mer peptides to stimulate peripheral blood

mononuclear cells (PBMCs), followed by successive rounds of deconvolution of positive wells to

determine stimulatory peptides and minimal optimal epitopes within positive 15-mers (10). This is

followed by generation of MHC-peptide tetramers, which stably bind T cell receptors and can be

used for cell staining and sorting (11). While successful, these methods are not ideal – using 15-

mer peptides, which are too large to bind class I MHC *in vivo*, can lead to false negatives in

ELISPOT assays (12). Currently, there is no high-throughput *in vitro* tool for assessing whether

shorter peptides are likely to bind to MHC and therefore are more likely to be CD8 T cell epitopes.

Rhesus macaques infected with simian immunodeficiency virus (SIV) and simian-human

immunodeficiency virus (SHIV) have long been used as models for HIV infection studies. Rhesus

macaque class I MHC class I A (Mamu-A) and class I B (Mamu-B) genes bind and present

peptides to CD8 T cells like their HLA-A and HLA-B counterparts in humans. Previous studies

have characterized the peptide binding specificity of common Mamu-A and Mamu-B molecules

to SIV-derived epitopes, and have identified epitopes that elicit CD8 T cell responses (13–16).

Knowledge of these epitopes enables tracking of SIV-specific responses after infection and

vaccination (17), and identification of the peptide binding specificities of MHC alleles commonly

found in elite controllers has guided the development of epitope-based vaccines against SIV (18).

Yet, even in this well-characterized system, T cell epitopes have only been defined for a small

minority of the hundreds of known MHC class I alleles, while nearly no T cell epitopes have been

defined for other macaque viruses or bacterial pathogens. While considerably more effort has

been dedicated to defining T cell epitopes in humans, these efforts are similarly concentrated on

a small number of pathogens (e.g., HIV, influenza). Methods that facilitate the high-throughput

identification of T cell epitopes, particularly for pathogens with large proteomes where T cells play

an essential role in protective immunity (e.g., *Mycobacterium tuberculosis*), are urgently needed.

Ultradense peptide arrays have previously been used as a high-throughput tool for assessing

antibody responses to pathogens; here, we repurpose them to identify peptides that bind specific

MHC class I molecules. Briefly, up to six million linear peptides are synthesized on a single array

chip, molecules of interest – antibodies in serum, or other molecules such as MHC proteins – are

allowed to bind to the peptides, and binding affinity is reported in fluorescence intensity units

based on the detection of fluorophore-conjugated secondary antibodies bound to the molecules.

Our group and others have designed arrays to simultaneously assess IgG and IgM binding to

linear pathogen-derived peptides (19, 20). Here, we generated a peptide array with 61,066 unique

8-mer, 9-mer, and 10-mer peptides derived from 82 SIV and SHIV isolates, and measured binding

to four rhesus macaque MHC class I molecules, Mamu-A1*001, -A1*002, -B*008, and -B*017, all

of which have peptide binding repertoires that have been extensively studied by conventional

methods. While we focus our attention here on a particular well-studied strain, SIVmac239, as a

way of validating the array, the large capacity of the array gave us the opportunity to generate the

same data for several dozen isolates, creating a sizable dataset that allows for comparisons

among strains. We tested the top 86 SIVmac239-derived peptides identified by the array as

having the highest binding affinity for Mamu-A1*001 for their ability to induce IFN-γ production in

peripheral blood mononuclear cells isolated from SIVmac239-infected, Mamu-A1*001-positive

rhesus macaques. We demonstrate that peptides that exhibit strong MHC binding on our array

frequently correspond to CD8 T cell epitopes that have been identified by conventional assays,

and that our high-binding peptides are enriched for putative novel epitopes that elicit CD8 T cell

responses. This suggests that the MHC-peptide array can be used to predict immunologically

relevant MHC-peptide interactions, offering a high-throughput alternative to traditional methods.

Materials and Methods

SIV/SHIV peptidome array design

Peptide sequences were generated from SIV and SHIV amino acid sequences (Supplemental table I). We downloaded 82 SIV and SHIV GenBank files from NCBI and extracted amino acid sequences for all open reading frames into FASTA files. The GenBank SIVmac239 nef protein sequence is truncated and therefore the correct full-length sequence was translated from the full GenBank sequence and added manually. The FASTA files were sent to Roche NimbleGen (Madison, WI; now Nimble Therapeutics) to generate the array. Non-redundant 8-mer, 9-mer, and 10-mer peptide sequences were generated with 1-amino acid offsets, and 5 replicates for each peptide were included on the array in a 12-plex array configuration.

Amino acid substitution array design

Substitution libraries were designed by introducing all possible single amino-acid substitutions and single-amino-acid deletions for the Gag CM9 epitope (CTPYDINQM) and the Tat SL8 epitope (STPESANL) using all 20 L-amino acids. The library was synthesized in five replicates. For this array, 1785 peptides were synthesized ($9 \times 21 \times 5 + 8 \times 21 \times 5 = 1785$, where 9 and 8 are the peptide lengths, 21 is the number of possible amino acids/deletion at each position, and 5 is the number of replicates).

Production of MHC molecules

Inclusion bodies were prepared essentially according to the method of Hutchinson, et al (21). In brief, bacterial expression plasmids containing the sequence for human beta-2 microglobulin (β2M) or rhesus MHC class I Mamu-A1*001, Mamu-A1*002, Mamu-B*008, or Mamu-B*017 were transformed into competent BL21DE3 *E. coli* bacteria. Single colonies were grown in 1 L TYP media (16 g Yeast extract, 16 g Tryptone, 5 g NaCl, 1 g K_2HPO_4) with appropriate antibiotic

markers, induced with IPTG during logarithmic growth phase, and allowed to express protein

overnight. Bacterial pellets were incubated with lysis buffer (10 mM Tris pH 8.1, 150 mM NaCl,

10 mM EDTA with DNAse and $MgCl_2$). Following sonication, the inclusion bodies were purified in

50 mM Tris pH 8.1 10 mM EDTA 100 mM NaCl buffer with 0.5% Triton X-100, washed in

resuspension buffer without Triton-X and the pellet dissolved in denaturing buffer. β2M inclusion

bodies were dissolved in 8M Urea with 50 mM Tris pH 8.1, 100 mM NaCl, 10 mM EDTA and 10

mM DTT. MHC heavy chains were dissolved in 6 M guanidine, 50 mM Tris pH 8.1, 100 mM NaCl,

10 mM EDTA with 10 mM DTT. After a Bradford assay to determine the concentration of protein,

inclusions were snap frozen in LN_2 and frozen at -80°C in aliquots of 30 mg for MHC heavy chain

and 10 mg for β2M.

MHC complex preparation

To prepare MHC I complexes for peptide array binding, 45 µl di-sodium tetraborate/sodium

hydroxide pH 10 buffer (Sigma-Aldrich), 10 µl 2 mg/ml β2M, and 60 µg of MHC class I heavy

chain stock were mixed in 400 µl water. This mixture was concentrated with AmiconUltra 0.5 ml,

10K centrifugal filter (Millipore) at 12,000 g for 5 min without pre-incubation. Concentrated material

was mixed with 400 µl 0.05X pH10 buffer to further dilute urea and 2-ME and concentrated in the

same Amicon filter at 12,000 g for 7 min. The concentrated sample was transferred to a 1.5 ml

Eppendorf tube, centrifuged at 20,000 g for 5 min to remove precipitated material, and stored at

4ºC before binding to the peptide array.

MHC-peptide binding assay

Peptide synthesis was accomplished through light-directed array synthesis in a Maskless Array

Synthesizer (MAS) using an amino-functionalized substrate as previously reported (22). Each

peptide replicate were randomly positioned on the array. After synthesis and de-protection, the

array was incubated in 0.67 µg/ml Cy5-labeled streptavidin (SA-Cy5) (Amersham), 1% alkali-

soluble casein (Novagen) at room temperature for 1 hour to stain technical SA-binding features

used for array QC and gridding. After a quick rinse in water, the array was dried by spinning in a

microcentrifuge equipped with an array holder and a 12-plex incubation chamber was applied to

the array surface. Before loading onto the array, MHC I complexes were 1:6 diluted in 20 mM tris-

HCl, pH7.4, 1% BSA and 6 µl aliquots were loaded into 5 chambers of a 12-plex chamber well.

An additional chamber was loaded without the MHC I complexes as a control. The loading ports

were sealed with port seals (Grace Bio-Labs, Bend, OR) and the array was incubated at 4ºC

overnight. After incubation the chamber was removed, and slides were rinsed in water and stained

by dipping with Alexa Fluor 647 MEM-123 1:1,000 (Novus Biologicals, Centennial, CO). After

rinsing in water and drying as described above, the array was scanned to collect fluorescence

data. Cy5 fluorescence intensity of the array was measured with an MS200 scanner at resolution

2 µm, wavelength 635 nm. Cy5 fluorescence intensities were extracted using Image Extraction

Software.

Peptide array data analysis

The array data contains the fluorescence intensity, reported in arbitrary signal intensity units, for

each peptide sequence replicate for each MHC molecule. We took the base 2 logarithm of the

fluorescence intensity and used the median value for each of the unique peptide

sequences. Using the median reduces noise induced by the peptide position on the array and

reduces the overall variance. We assessed the top 24, 48, 96, and 192 peptides with the highest

binding intensities in each virus-MHC class I molecule combination for the presence of known

epitopes.

IFN-γ ELISPOT assay

ELISPOTs were performed in duplicate using frozen rhesus macaque PBMCs isolated from

EDTA-treated whole blood. Frozen cells were thawed slowly at room temperature and washed

twice in R10 media containing RPMI, 10% FBS, 1% antibiotic/antimycotic, and 1% L-glutamine

(Hyclone, Logan, UT). Approximately 100,000 cells in 100 µl of R10 media were layered on pre-

coated monkey IFN-γ ELISpot-Plus plates (Mabtech Inc., Mariemont, Ohio, USA) with a single

peptide per well at a final concentration of 1 µM. Peptides were synthesized by GenScript

(Piscataway, NJ, USA). Concanavalin A (ConA, 10 µM) was used as a positive control and cells

with no peptide were used as a negative control.

ELISPOT analysis

Developed ELISPOT plates were read with an AID Elispot reader using the IFN-γ setting and a

threshold of 100 spots per million cells. ConA wells were counted using custom parameters

defined for ConA (Min intensity: 10, Min size: 20, Min gradient: 1, Max intensity: 57, Max size:

295, Max gradient: 50, Basic algorithm settings emphasis: tiny, Algorithm: C); the rest were read

using custom IFN-γ parameters (Min intensity: 20, Min size: 40, Min gradient: 1, Max intensity:

254, Max size: 5000, Max gradient: 90, Basic algorithm settings emphasis: small, Algorithm: C).

Positive ELISPOT responses were determined as previously described (23).

Tetramers and tetramer staining

Tetramers conjugated to APC or PE were generously produced by the NIH Tetramer Core Facility

using peptides generated by GenScript. Thawed PBMC resuspended to 1 x 10^6 cells/100 µl of

R10 medium were stained with tetramers at a concentration of 5 µg/ml in the dark for one hour at

37ºC, and were subsequently stained with CD3 Alexa Fluor 700 (BD Biosciences, San Jose, CA)

and CD8 BV421 (BD Biosciences) for 15 minutes at room temperature, washed, and fixed with

2% paraformaldehyde. Samples were run on a BD LSRII flow cytometer within 24 hours and were

analyzed with FlowJo software, version 10.5.3.

Results

Identification of SIVmac239 peptides that bind MHC class I molecules

We tested binding of purified rhesus MHC class I Mamu-A1*001, -A1*002, -B*008, and -B*017 molecules to the full proteomes of 82 SIV and SHIV isolates (Supplemental table I). Previous studies have identified SIVmac239 epitopes that bind each of the four MHC class I molecules tested here, which enables us to validate the peptide array as a method for screening for putative novel CD8 T cell epitopes (14–16, 24). To determine whether the array identified previously-established SIVmac239 epitopes, we determined the percentage of known CD8 T cell epitopes that were identified in the top 24, 48, 96, and 192 (convenient fractions/multiples of a 96-well ELISPOT plate) highest-binding peptides for each MHC molecule. The top 192 peptides for each of the four MHC molecules encompassed 27-64% of the known T cell epitopes restricted by each molecule (Table I). Figure 1 shows the complete data for SIVmac239 peptides binding to Mamu-A1*001, -A1*002, -B*008, and -B*017, with previously-identified epitopes present in the top 192 peptides (out of 9912 SIVmac239 peptides, or 1.9%) for each molecule indicated.

Critical residues for Mamu-A1*001 peptide binding identified through substitution analysis

Peptide anchor residues define the binding motif for a given MHC molecule and enable prediction of other peptides likely to bind. The binding motif for Mamu-A1*001 was previously defined by sequence analysis of peptides bound to the molecule in vitro and by performing competitive binding assays using peptides with amino acid substitutions (24, 25). These studies identified positions two and three (P2 and P3), and the C-terminus, as important for binding. We performed a full substitution analysis at each amino acid for two known immunodominant CD8 T cell epitopes that bind to Mamu-A1*001, Gag CM9 (Gag 181-189) and Tat SL8 (Tat 28-35). We tested the influence of all 20 amino acids, plus an amino acid deletion, at each position on the binding of the peptides to Mamu-A1*001 molecules. As would be expected based on prior studies, substitutions

at P2 (threonine) and P3 (proline) for both peptides were generally poorly tolerated and resulted

in dramatically decreased signal intensity, indicating reduced binding affinity (Figure 2). At P2, a

substitution with serine was tolerated; P3 did not tolerate any substitutions. These results

corroborated those found by competitive binding assays using Gag CM9 peptides with single

amino acid substitutions (24). A previous study found that L, I, V, M, F, W, Y, and T are tolerated

at the C-terminus; however, this study did not test every possible amino acid at that position, and

our results suggest that R, K, and H may also be tolerated (24).

Measuring IFN-γ ELISPOT responses to SIVmac239 peptides that bind to Mamu-A1*001

We assessed whether the MHC peptide array can be used to efficiently screen for putative novel

CD8 T cell epitopes by performing IFN-γ ELISPOTs for top-binding peptides. We tested six

previously-established Mamu-A1*001 CD8 T cell epitopes plus the top 86 SIVmac239 peptides

that bound Mamu-A1*001 with the highest binding scores, using PBMCs from four SIVmac239-

infected Mamu-A1*001-positive rhesus macaques. PBMC from each animal ranged from 12 to 47

weeks post infection. Of the established epitopes, Gag CM9 was positive in all four animals; Tat

SL8 was positive in two (TVPWPNASL was positive in one, and LGPHYTPKIV, CAPPGYAL, and

CAPPGYALL were positive in none). Of the 86 peptides selected from the array, 10 were positive

in at least one animal and three were positive in three animals (Supplemental table II). All three

of the peptides positive in multiple animals (STPESANLG, STPESANLGE, TPESANLGE)

overlapped Tat SL8, and in one animal were positive even though SL8 itself was not. Given the

extensive research on Mamu-A1*001-restricted T cell responses to SIVmac239 over the past two

decades, it is perhaps not surprising that we did not identify additional shared responses to

epitopes in this virus restricted by this allele.

To quantify and confirm these responses, we produced Mamu-A1*001 tetramers folded with Gag

CM9, Tat SL8, and the three peptides identified by ELISPOT. We tested these for CD8 T cell

binding using PBMCs from two Mamu-A1*001-positive animals whose cells we used in our

ELISPOT experiments (one animal that did and one that did not show responses to the three

ELISPOT-positive peptides) and one animal that did not have Mamu-A1*001. For Gag CM9, Tat

SL8, STPESANLG, and STPESANLGE (but not TPESANLGE), we identified populations of

tetramer-positive CD8 T cells within the PBMCs from the Mamu-A1*001-positive, ELISPOT-

positive animal, and much smaller populations in those from the Mamu-A1*001-positive,

ELISPOT-negative animal (Figure 3). As expected, we did not identify populations of tetramer-

positive CD8 T cells in the Mamu-A1*001-negative animal.

Comparison of Mamu-A1*001 binding peptides across SIV strains

Due to its large capacity, the peptide array offers the ability to compare binding of MHC molecules

to peptides across several related pathogen strains. As a proof of concept, we compared Mamu-

A1*001 binding to SIVmac239 and SIVmac251 peptides. Most of the known Mamu-A1*001-

restricted SIVmac239 epitopes have the same sequence in SIVmac251, but a few have amino

acid variations. Of the six SIVmac239 epitopes that had amino acid variations in SIVmac251, we

identified one (QSPGGLDKGL in SIVmac239, QSLGGLGKGL in SIVmac251) that lost its proline

in position 3; as we would expect based on the importance of position 3 in binding, this epitope

was in the top 192 highest-binding peptides for SIVmac239 but not for SIVmac251 (Supplemental

table III).

We additionally compared Mamu-A1*001 binding at the region corresponding to the SIVmac239

Tat SL8 epitope across all viruses on the array. CD8 T cell responses to the SL8 epitope in

animals carrying the Mamu-A1*001 allele have been associated with suppression of viral

replication, and mutations in this region are commonly implicated in viral escape (26, 27).

Identifying whether Mamu-A1*001 is likely to be able to bind to this region despite amino acid

sequence differences will be important in determining whether this epitope region is likely to be

immunodominant in viral strains other than SIVmac239. SIVmac251 strains, also commonly used

in research, are split between having the SL8 sequence and TTPESANL (TL8). TL8 has

previously been found to be immunodominant; however, this was first established in experiments

studying macaques infected with SIVmac239 – suggesting that there is some latitude in Mamu-

A1*001's ability to bind to that region (13). The peptide array results affirm TL8 as a strong binder

to Mamu-A1*001, with a higher binding score (8.72) than SL8 (8.33). We found that all SIVsm

(isolated from sooty mangabeys) strains had the sequence PTPESANL (PL8), an escape variant

seen in SIVmac239 infection that exhibits reduced binding to Mamu-A1*001 (27). Similarly, on

our array we found that PL8 had a lower binding score than SL8 and TL8 (6.28), but one that was

still above baseline.

Correlation of binding scores to IC_{50} values

IC_{50} values obtained in competitive binding assays are customarily used to represent MHC-

peptide binding affinity; however, the equilibrium dissociation constant (k_d) is more commonly

used to quantify protein-ligand binding affinity and is generally preferred as it does not depend on

experimental conditions. The peptide array analysis outputs binding scores for each MHC-peptide

combination measured in arbitrary fluorescence intensity units, which reflect binding affinity. While

we expect these values to correlate with those obtained from competitive binding assays, recent

studies have found that IC_{50} values do not accurately estimate k_d values, making it challenging to

make a direct comparison between IC_{50} and fluorescence intensity units (28). However, using the

IC_{50} data available in the literature for peptides tested for binding to Mamu-A1*001, -A1*002, -

B*008, and -B*017, we do find that peptide array binding scores for peptides that were previously

found to bind an MHC molecule ($IC_{50} \leq 500$ nM) were significantly higher (Median = 0.51; 95%

CI = 0.34, 0.70; $p < 0.001$, using a Wilcoxon rank-sum test) than for those that did not bind MHC

($IC_{50} > 500$ nM) (Figure 4). This dataset had a small selection of binding peptides for all MHC

molecules with non-binding peptides reported only for Mamu-B*008; using a dataset of IC_{50} values

for all peptide/MHC combinations generated using Stabilized Matrix Method (SMM) MHC-peptide

binding prediction software where experimental values were unavailable yielded the same

significant difference (data not shown).

Discussion

Identification of CD8 T cell epitopes has historically relied on low-throughput experiments with cumbersome peptide screening methods. In an effort to develop a high-throughput method for screening peptides for MHC binding prior to determining CD8 T cell responses, we used ultradense peptide array technology to test the binding of class I MHC molecules to viral peptides. Compared to conventional methods, ultradense peptide arrays provide massive amounts of information regarding binding of class I MHC molecules to thousands of individual peptides in a single experiment. In this study, of the 86 peptides that bound Mamu-A1*001 with the highest fluorescence intensity scores on the array, we identified 10 peptides that were positive in at least one animal and three that were positive in three animals using IFN-γ ELISPOTs and PBMCs from four SIVmac239-infected Mamu-A1*001-positive rhesus macaques. These peptides were not reported as epitopes previously, thus demonstrating the power of the peptide array method as an alternative approach to screen for putative novel immunogenic epitopes. We additionally demonstrate that many of the same epitopes previously found to bind each molecule and induce CD8 T cell responses were identified as high-binding peptides in our array; 27-64% of the previously-identified epitopes for each MHC molecule were found in the top 192 high-binding peptides on our array.

Despite these initial promising results, many known epitopes were not detected by our method. This may be due in part to the choice of 192 peptides as a convenient cutoff for "high binding." Because the peptide array binding scores do not directly correspond to IC_{50} values from competitive binding assays, we refrain from choosing a hard binding score cutoff to distinguish binding from non-binding peptides. While there was a significant difference ($p < 0.001$) in fluorescence intensity values between peptides that were previously found to bind or not bind to MHC molecules based on competitive binding assays, there were some instances where

fluorescence intensity values did not reflect the results of the competitive binding assays. For

example,the Tat SL8 epitope's high fluorescence intensity (7.15 sd above the median for Mamu-

A1*001) and low IC_{50} value both indicate strong peptide binding. However, the Gag CM9 epitope

showed a substantially lower fluorescence intensity (2.36 sd above the median for Mamu-

A1*001), but also had a low IC_{50} value. These discrepancies may be attributable to differences in

how certain peptides or MHC molecules perform on the array; certainly, it would be reasonable

to expect that some differences may arise because peptides are anchored on the array rather

than free-floating. Unfortunately, neither competitive binding assays nor peptide arrays truly mimic

in vivo MHC-peptide binding, so without further optimization of the method, it would be premature

to label one method as "more true" than the other.

We tested a portion of the high-binding peptides identified here by IFN-γ ELISPOT as a proof of

concept. We identified three SIVmac239 peptides that strongly bound Mamu-A1*001 that had

positive responses in at least two animals and were not previously identified epitopes; two of the

tetramers generated to test these responses were functional. All three peptides overlapped the

Tat SL8 epitope, and the two functional tetramers had weaker responses than the SL8 tetramer,

which may suggest that these are not separate epitopes but simply extensions of the SL8 minimal

optimal epitope. Still, this indicates that a researcher using the array to look for MHC binding to

an unknown pathogen's proteome could reasonably expect to find CD8 T cell epitopes in a subset

of high-binding peptides identified by the array; moreover, this demonstrates that, at least in some

cases, peptide binding to clusters of related sequences (ie., 8-mer, 9-mer, and 10-mer peptides

containing the same component 8-mer) could be used as a filter to identify interesting peptides

for follow-up. Using the array as a screening tool could thus drastically reduce the amount of work

and resources required to identify novel epitopes.

MHC-peptide array technology has not yet been optimized to work with class II MHC molecules.

CD8 T cell responses are not the sole arbiter of protective immunity in HIV vaccination; a better

understanding of CD4 responses is necessary for developing a more comprehensive picture of

the determinants of a successful vaccine. Assessing CD4 T cell responses in PBMCs and

developing class II tetramers is currently more difficult than for CD8 T cells and class I MHC.

While this will present challenges in validating a class II array, it underscores the need for a

method for rapidly screening for potential CD4 T cell responses, and optimizing a class II array

will be an important future direction.

Ultradense peptide arrays offer a method for rapidly mapping class I MHC binding to thousands

of individual peptides. While we used peptides derived from SIV and SHIV here, the scalability of

ultradense peptide arrays makes them an attractive option for characterizing MHC class I binding

to peptides from larger, more complex pathogens. *Mycobacterium tuberculosis* (Mtb) Erdman

strain, for instance, has 4,246 annotated protein sequences, in contrast to nine for SIV. The size

of the Mtb proteome makes thorough analyses of MHC binding across the full proteome nearly

impossible with conventional methods, but just three peptide array chips would be sufficient to

accommodate a full 8-, 9-, and 10-mer peptide analysis for a single MHC allele. The ability to

screen peptides for MHC binding en masse prior to performing *in vitro* tests will streamline epitope

identification studies that might otherwise be prohibitively large and expensive to conduct.

Acknowledgments

The authors would like to thank Amy Ellis for her help in analyzing the tetramer flow data.

References

1. Hewitt, E. W. 2003. The MHC class I antigen presentation pathway: strategies for viral immune evasion. *Immunology* 110: 163-169.

2. Halle, S., O. Halle, and R. Förster. 2017. Mechanisms and Dynamics of T Cell-Mediated Cytotoxicity In Vivo. *Trends Immunol* 38: 432-443.

3. Wieczorek, M., E. T. Abualrous, J. Sticht, M. Álvaro-Benito, S. Stolzenberg, F. Noé, and C. Freund. 2017. Major Histocompatibility Complex (MHC) Class I and MHC Class II Proteins: Conformational Plasticity in Antigen Presentation. *Front Immunol* 8: 292.

4. McMichael, A. J., and W. C. Koff. 2014. Vaccines that stimulate T cell immunity to HIV-1: the next step. *Nat Immunol* 15: 319-322.

5. Zhao, L., M. Zhang, and H. Cong. 2013. Advances in the study of HLA-restricted epitope vaccines. *Hum Vaccin Immunother* 9: 2566-2577.

6. Peters, B., H. H. Bui, J. Sidney, Z. Weng, J. T. Loffredo, D. I. Watkins, B. R. Mothé, and A. Sette. 2005. A computational resource for the prediction of peptide binding to Indian rhesus macaque MHC class I molecules. *Vaccine* 23: 5212-5224.

7. Di Carluccio, A. R., C. F. Triffon, and W. Chen. 2018. Perpetual complexity: predicting human CD8. *Immunol Cell Biol* 96: 358-369.

8. E Silva, R. F., L. F. Ferreira, M. Z. Hernandes, M. E. de Brito, B. C. de Oliveira, A. A. da Silva, O. P. de-Melo-Neto, A. M. Rezende, and V. R. Pereira. 2016. Combination of In Silico Methods in the Search for Potential CD4(+) and CD8(+) T Cell Epitopes in the Proteome of Leishmania braziliensis. *Front Immunol* 7: 327.

9. Mahdavi, M., and V. Moreau. 2016. In silico designing breast cancer peptide vaccine for binding to MHC class I and II: A molecular docking study. *Comput Biol Chem* 65: 110-116.

10. Budde, M. L., J. J. Lhost, B. J. Burwitz, E. A. Becker, C. M. Burns, S. L. O'Connor, J. A. Karl, R. W. Wiseman, B. N. Bimber, G. L. Zhang, W. Hildebrand, V. Brusic, and D. H.

O'Connor. 2011. Transcriptionally abundant major histocompatibility complex class I

alleles are fundamental to nonhuman primate simian immunodeficiency virus-specific

CD8+ T cell responses. *J Virol* 85: 3250-3261.

11.   Sims, S., C. Willberg, and P. Klenerman. 2010. MHC-peptide tetramers for the analysis of

antigen-specific T cells. *Expert Rev Vaccines* 9: 765-774.

12.   Fiore-Gartland, A., B. A. Manso, D. P. Friedrich, E. E. Gabriel, G. Finak, Z. Moodie, T.

Hertz, S. C. De Rosa, N. Frahm, P. B. Gilbert, and M. J. McElrath. 2016. Pooled-Peptide

Epitope Mapping Strategies Are Efficient and Highly Sensitive: An Evaluation of Methods

for Identifying Human T Cell Epitope Specificities in Large-Scale HIV Vaccine Efficacy

Trials. *PLoS One* 11: e0147812.

13.   Allen, T. M., B. R. Mothé, J. Sidney, P. Jing, J. L. Dzuris, M. E. Liebl, T. U. Vogel, D. H.

O'Connor, X. Wang, M. C. Wussow, J. A. Thomson, J. D. Altman, D. I. Watkins, and A.

Sette. 2001. CD8(+) lymphocytes from simian immunodeficiency virus-infected rhesus

macaques recognize 14 different epitopes bound by the major histocompatibility complex

class I molecule mamu-A*01: implications for vaccine design and testing. *J Virol* 75: 738-

749.

14.   Mothé, B. R., J. Sidney, J. L. Dzuris, M. E. Liebl, S. Fuenger, D. I. Watkins, and A. Sette.

2002. Characterization of the peptide-binding specificity of Mamu-B*17 and identification

of Mamu-B*17-restricted epitopes derived from simian immunodeficiency virus proteins. *J

Immunol* 169: 210-219.

15.   Loffredo, J. T., J. Sidney, C. Wojewoda, E. Dodds, M. R. Reynolds, G. Napoé, B. R.

Mothé, D. H. O'Connor, N. A. Wilson, D. I. Watkins, and A. Sette. 2004. Identification of

seventeen new simian immunodeficiency virus-derived CD8+ T cell epitopes restricted by

the high frequency molecule, Mamu-A*02, and potential escape from CTL recognition. *J

Immunol* 173: 5064-5076.

16.   Loffredo, J. T., J. Sidney, A. T. Bean, D. R. Beal, W. Bardet, A. Wahl, O. E. Hawkins, S.

Piaskowski, N. A. Wilson, W. H. Hildebrand, D. I. Watkins, and A. Sette. 2009. Two MHC

class I molecules associated with elite control of immunodeficiency virus replication,

Mamu-B*08 and HLA-B*2705, bind peptides with sequence similarity. *J Immunol* 182:

7763-7775.

17.   John T. Loffredo, L. E. V., David I. Watkins. 2006. Beyond Mamu-A*01+ Indian Rhesus

Macaques: Continued Discovery of New MHC Class I Molecules that Bind Epitopes from

he Simian AIDS Viruses. *HIV Molecular Immunology* 29-51.

18.   Martins, M. A., D. C. Tully, M. A. Cruz, K. A. Power, M. G. Veloso de Santana, D. J. Bean,

C. B. Ogilvie, R. Gadgil, N. S. Lima, D. M. Magnani, K. Ejima, D. B. Allison, M. Piatak, J.

D. Altman, C. L. Parks, E. G. Rakasz, S. Capuano, R. Galler, M. C. Bonaldo, J. D. Lifson,

T. M. Allen, and D. I. Watkins. 2015. Vaccine-Induced Simian Immunodeficiency Virus-

Specific CD8+ T-Cell Responses Focused on a Single Nef Epitope Select for Escape

Variants Shortly after Infection. *J Virol* 89: 10802-10820.

19.   Heffron, A. S., E. L. Mohr, D. Baker, A. K. Haj, C. R. Buechler, A. Bailey, D. M. Dudley, C.

M. Newman, M. S. Mohns, M. Koenig, M. E. Breitbach, M. Rasheed, L. M. Stewart, J.

Eickhoff, R. S. Pinapati, E. Beckman, H. Li, J. Patel, J. C. Tan, and D. H. O'Connor. 2018.

Antibody responses to Zika virus proteins in pregnant and non-pregnant macaques. *PLoS*

Negl Trop Dis 12: e0006903.

20.   Mishra, N., A. Caciula, A. Price, R. Thakkar, J. Ng, L. V. Chauhan, K. Jain, X. Che, D. A.

Espinosa, M. Montoya Cruz, A. Balmaseda, E. H. Sullivan, J. J. Patel, R. G. Jarman, J. L.

Rakeman, C. T. Egan, C. B. E. M. Reusken, M. P. G. Koopmans, E. Harris, R. Tokarz, T.

Briese, and W. I. Lipkin. 2018. Diagnosis of Zika Virus Infection by Peptide Array and

Enzyme-Linked Immunosorbent Assay. *MBio* 9:

21.   Hutchinson, S. L., L. Wooldridge, S. Tafuro, B. Laugel, M. Glick, J. M. Boulter, B. K.

Jakobsen, D. A. Price, and A. K. Sewell. 2003. The CD8 T cell coreceptor exhibits

disproportionate biological activity at extremely low binding affinities. *J Biol Chem* 278:

24285-24293.

22.   Forsström, B., B. B. Axnäs, K. P. Stengele, J. Bühler, T. J. Albert, T. A. Richmond, F. J.

Hu, P. Nilsson, E. P. Hudson, J. Rockberg, and M. Uhlen. 2014. Proteome-wide epitope

mapping of antibodies using ultra-dense peptide arrays. *Mol Cell Proteomics* 13: 1585-

1597.

23.   Reynolds, M. R., A. M. Weiler, S. M. Piaskowski, M. Piatak, H. T. Robertson, D. B. Allison,

A. J. Bett, D. R. Casimiro, J. W. Shiver, N. A. Wilson, J. D. Lifson, W. C. Koff, and D. I.

Watkins. 2012. A trivalent recombinant Ad5 gag/pol/nef vaccine fails to protect rhesus

macaques from infection or control virus replication after a limiting-dose heterologous SIV

challenge. *Vaccine* 30: 4465-4475.

24.   Sidney, J., J. L. Dzuris, M. J. Newman, R. P. Johnson, A. Kaur, K. Amitinder, C. M.

Walker, E. Appella, B. Mothe, D. I. Watkins, and A. Sette. 2000. Definition of the Mamu

A*01 peptide binding specificity: application to the identification of wild-type and optimized

ligands from simian immunodeficiency virus regulatory proteins. *J Immunol* 165: 6387-

6399.

25.   Allen, T. M., J. Sidney, M. F. del Guercio, R. L. Glickman, G. L. Lensmeyer, D. A. Wiebe,

R. DeMars, C. D. Pauza, R. P. Johnson, A. Sette, and D. I. Watkins. 1998.

Characterization of the peptide binding motif of a rhesus MHC class I molecule (Mamu-

A*01) that binds an immunodominant CTL epitope from simian immunodeficiency virus. *J*

Immunol 160: 6062-6071.

26.   Loffredo, J. T., B. J. Burwitz, E. G. Rakasz, S. P. Spencer, J. J. Stephany, J. P. Vela, S. R.

Martin, J. Reed, S. M. Piaskowski, J. Furlott, K. L. Weisgrau, D. S. Rodrigues, T. Soma, G.

Napoé, T. C. Friedrich, N. A. Wilson, E. G. Kallas, and D. I. Watkins. 2007. The antiviral

efficacy of simian immunodeficiency virus-specific CD8+ T cells is unrelated to epitope

specificity and is abrogated by viral escape. *J Virol* 81: 2624-2634.

27.   Allen, T. M., D. H. O'Connor, P. Jing, J. L. Dzuris, B. R. Mothé, T. U. Vogel, E. Dunphy, M.

E. Liebl, C. Emerson, N. Wilson, K. J. Kunstman, X. Wang, D. B. Allison, A. L. Hughes, R.

C. Desrosiers, J. D. Altman, S. M. Wolinsky, A. Sette, and D. I. Watkins. 2000. Tat-specific

cytotoxic T lymphocytes select for SIV escape variants during resolution of primary

viraemia. *Nature* 407: 386-390.

28.   Barbet, J., and S. Huclier-Markai. 2019. Equilibrium, affinity, dissociation constants, IC5O:

Facts and fantasies. *Pharm Stat*

Table I: The percentage of known epitopes for each MHC molecule that were found in the top N

peptides with high signal intensity.

Figure 1: MHC-peptide binding plots. Previously-identified SIVmac239 8-mer, 9-mer, and 10-

mer epitopes found in the top 192 peptides for each MHC class I molecule. Parentheticals

indicate overlapping epitopes.

Figure 2: Identification of anchor residues by substitution analysis. Single amino acid

substitutions were made at each position of the Gag CM9 (A) and SL8 (B) epitopes. Each box

represents a peptide with the indicated substitution, and the fold change in signal intensities for

each substitution (compared to the wild-type sequence) is reported. Position numbers are

reported along the left axis of the heatmap; the fold change color scale is indicated at right.

Figure 3: Tetramer analysis. Mamu-A1*001 MHC-peptide tetramers were generated and

tested for binding to PBMC from SIV-infected animals. (A) A representative sequential gating

strategy for the flow cytometry data is shown. (B) Tetramer binding to PBMC from three animals

was tested; shown are the ELISPOT responses to each peptide (STPESANLG, STPESANLGE,

and TPESANLG are abbreviated as SG9, SE10, and TE9 respectively), and the percentage of

CD8 T cells that were tetramer-positive.

Figure 4: Comparison of IC_{50} and fluorescence intensity scores. Peptides with

experimentally obtained IC_{50} values for binding to Mamu-A1*001, -A1*002, -B*008, or -B*017

were grouped by IC_{50} value into binding ($IC_{50} \leq 500$ nM) and non-binding ($IC_{50} > 500$ nM)

peptides. Binding peptides had significantly higher fluorescence intensity scores than non-

binding peptides ($p < 0.001$, using a Wilcoxon rank-sum test).

	N = 24	N = 48	N = 96	N = 192
Mamu-A1*001	22%	44%	47%	64%
Mamu-A1*002	0%	0%	16%	37%
Mamu-B*008	1%	8%	12%	27%
Mamu-B*017	12%	18%	29%	41%
Average coverage	9%	18%	26%	42%

Table I

A.

Substituted amino acid

B.

Substituted amino acid

Figure 3

B.

Animal:		A		B		C	
A1*001:		**+**		**+**		**-**	
WPI:		**12**		**30**		**11**	
ELISPOT/ Tetramer:	E	T	E	T	E	T
CM9	+	4.09	+	1.43	n/a	0
SL8	+	1.05	-	0.10	n/a	0.002
SG9	+	0.77	-	0.035	n/a	0.005
SE10	+	0.36	-	0.043	n/a	0.004
TE9	+	0.004	-	0	n/a	0