Testing the ability of species distribution models to infer variable importance

Running head: Inferring variable importance

Authors

Adam B. Smith[1*] & Maria J. Santos[2]

[1] Center for Conservation and Sustainable Development, Missouri Botanical Garden, Saint Louis
MO 63116 USA. Email: adam.smith@mobot.org

[2] University Research Priority Program in Global Change and Biodiversity and Department of
Geography, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland

* Corresponding author

Abstract: 290 words

Main text: 5992 words

References cited: 64

Number of tables: 1

Number of boxes: 1

Number of figures: 4

Number of supplemental appendices: 10

Abstract

Models of species' distributions and niches are frequently used to infer the importance of range- and niche-defining variables. However, the degree to which these models can reliably identify important variables and quantify their influence remains unknown. Here we use a series of simulations to explore how well models can 1) discriminate between variables with different influence and 2) calibrate the magnitude of influence relative to an "omniscient" model. To quantify variable importance, we trained generalized additive models (GAMs), Maxent, and boosted regression trees (BRTs) on simulated data and tested their sensitivity to permutations in each predictor. Importance was inferred by calculating the correlation between permuted and unpermuted predictions, and by comparing predictive accuracy of permuted and unpermuted predictions using AUC and the Continuous Boyce Index. In scenarios with one influential and one uninfluential variable, models were unable to discriminate reliably between variables in conditions that are normally challenging for generating accurate predictions: training occurrences <8-64; prevalence >0.5; small spatial extent; environmental data with coarse resolution when spatial autocorrelation is low; and correlation between environmental variables where $|r| > 0.7$. When two variables influenced the distribution equally, importance was underestimated when species had narrow or intermediate niche breadth. Interactions between variables in how they shaped the niche did not affect inferences about their importance. When variables acted unequally, the effect of the stronger variable was overestimated. GAMs and Maxent discriminated between variables more reliably than BRTs, but no algorithm was consistently well-calibrated vis-à-vis the omniscient model. Algorithm-specific measures of importance like Maxent's change-in-gain metric were less robust than the permutation test. Overall, high predictive accuracy did not connote robust inferential capacity. As a result, requirements for

42   reliably measuring variable importance are likely more stringent than for creating models with

43   high predictive accuracy.

44   **Key words**

45   species distribution model, ecological niche model, variable importance, statistical inference,

46   sample size, collinearity, spatial autocorrelation, spatial scale

Introduction

What environmental factors determine species' ranges and environmental tolerances? The answer to this question remains elusive for most species on Earth despite being crucial for addressing long-standing issues in theoretical and applied ecology. Although manipulative field- and laboratory-based studies can identify factors that shape niches and ranges (Hargreaves et al. 2016; Lee-Yaw et al. 2016), for most species logistical difficulties preclude examining large numbers of variables and studying range limits across wide geographic scales. Alternatively, environmental limits can be inferred using models of species' geographic ranges and niches. These ecological niche models and species distribution models (SDMs) are constructed by correlating observations of occurrence with data on environmental conditions at occupied sites. Indeed, one of the most common uses for SDMs is to identify important variables (e.g., 227 inferential studies analyzed by Bradie & Leung 2017 using the Maxent algorithm alone). However, we are aware of no studies that systematically evaluate how well SDMs measure variable importance. Compared to the attention devoted to understanding predictive accuracy of models of niches and distributions (e.g., Elith et al. 2006; Smith et al. 2013; Buklin et al. 2015; Guevara et al. 2018; Norberg et al. 2019), the lack of research on the efficacy of these same models for identifying important variables is a striking oversight.

The ability of an SDM to infer variable importance will likely be affected by factors that are extrinsic to the species and thus at least nominally under control of the modeler (e.g., sample size, study region extent, etc.) and by factors intrinsic to the species (e.g., niche breadth). Here we utilize a reductionist approach based on virtual species (Meynard et al. 2019) to systematically evaluate a set of extrinsic and intrinsic factors expected to influence variable inference. Our goal was to identify the minimal conditions under which each factor allows robust

inference of variable importance, assuming all other conditions are optimal. We conducted nine

simulation experiments to explore circumstances that affect inference. We start with the simplest

case in which a species' range is determined by a single "TRUE" variable, but the SDMs are

presented with data on this variable plus an uncorrelated "FALSE" variable with no effect on

distribution. We then explore the effects of sample size, spatial scale, and collinearity between

variables. Finally, we examine cases where the species' distribution is shaped by two collinear

TRUE variables that can be correlated and can interact to define the niche.

Methods

General approach

We assume a variable is "important" if a model has high predictive accuracy and if the

predictions are highly sensitive to changes in values of that variable (Meinshausen & Bühlmann

2010). To measure sensitivity of the model to changes in the variable, we use the permute-after-

calibration test (Breiman 2001) which compares unpermuted and permuted predictions. We

compared unpermuted and permuted predictions by calculating for each the Continuous Boyce

Index (CBI), which indicates how well a model's predictions serve as an index of the probability

of presence (Boyce et al. 2002; Hirzel et al. 2006), and the area under the receiver-operator curve

(AUC), which indicates a model's ability to differentiate between presence and non-presence

sites. We also calculated the correlation between unpermuted and permuted predictions (COR;

Breiman 2001), which reflects differences between the two sets of predictions. Increasing the

effect of a variable in a model should decrease CBI, AUC, or COR relative to an unimportant

variable.

To simulate species' distributions, we defined a generative function of one or two variables on

the landscape, then used it to produce a raster of the probability of occurrence. For each cell, true

occupancy was determined using a Bernoulli draw with the probability of success equal to the

simulated probability of presence (Meynard & Kaplan 2013). We then calibrated and evaluated

three SDM algorithms: generalized additive models (GAMs; Wood 2006), Maxent (Phillips et al.

2006), and boosted regression trees (BRTs; Elith et al. 2008). Full details of model calibration

are presented in Appendix 1. Briefly, (unless otherwise stated) SDMs were calibrated using 200

occurrences and 10000 (Maxent and GAMs) or 200 (BRTs) background sites (Baret-Massin et

al. 2012). Predictive accuracy and inferential capacity were evaluated using 200 distinct test

occurrences plus either 10,000 background sites (CBI, AUC_{bg}, and COR_{bg}) or 200 absences

(AUC_{pa} and COR_{pa}; Meynard et al. 2019). In addition to the permute-after-calibration test, for

each of the experiments we also evaluated algorithm-specific measures of variable importance:

AIC-based variable weighting for GAMs (Burnham & Anderson 2002); contribution,

permutation, and change-in-gain tests for Maxent (Phillips & Dudík 2008); and deviance

reduction for BRTs (Elith et al. 2008). To streamline discussion, we present results only for

Maxent and CBI in the main text (Appendices 3-9 present the complete set of results).

In our simulations there is no process-based spatial autocorrelation in species occurrences arising

from dispersal, disturbance, or similar processes. As a result, occurrences at sites are statistically

independent of one another regardless of their proximity, which obviates the need to use

geographically distinct training and test sites. This is a convenience that is unlikely to be met in

real-world situations where robust inference requires test data that is geographically and/or

temporally as independent as possible from training data (Roberts et al. 2017; Fourcade et al.

2018).

For each level of a factor we manipulated in an experiment (e.g., landscape size), we generated

100 landscapes with training and test data sets, then calibrated and evaluated GAM, Maxent, or

BRT models on each set. As a benchmark, we used an "omniscient" (OMNI) model which had

the same variables and parameter values as the generative model used to create the species'

probability of occurrence (Meynard et al. 2019). We evaluated the OMNI model with the same

set of test data used to evaluate the SDMs. Since OMNI does not use training data variability in

its results is due solely to stochastic differences in test sites between iterations. To generate

predictions from permuted variables, for a given level of a factor, data iteration, and SDM

algorithm, we created 30 permutations of each variable, calculated test statistics (CBI, AUC, and

COR) for each, then took the average test statistic value across these 30 sets.

We know a priori that levels of a factor are different, so our interest is in the effect size of each

factor level (White et al. 2014), which can be discerned by eye. For a given test statistic (CBI,

AUC, or COR), we assessed the capacity of the permute-after-calibration test to assess

discrimination (qualitative differences between variables with different influence) and

calibration (how well the distribution of the SDM's test statistic matches that of the test statistic

generated using the unbiased OMNI model). We designate a test as having "reliable

discrimination" if there is complete lack of overlap between the inner 95% of the distribution of

the test statistic between the permuted and unpermuted predictions across the 100 iterations. We

designate a test as having "reliable calibration" by comparing the distribution of the test statistic

between the SDM and OMNI: the range of the SDM's inner 95% of values across data iterations

are within ±10% of OMNI's range, and the SDM's median value is within the 40th and 60th

percentile of OMNI's median value. Under this definition, neither CBI, COR_{pa}, nor COR_{bg} were

ever well-calibrated, although in a few cases AUC_{pa} and AUC_{bg} yielded well-calibrated

outcomes (Appendices 5 and 6). Hence, hereafter we focus on discrimination accuracy.

Experiment 1: Simple scenario

In the simplest scenario we assumed the species' probability of occurrence is determined by a

logistic generative function of a single TRUE variable:

Eq. 1 $$\Pr(occ) = \frac{exp(\beta \times TRUE)}{1 + exp(\beta \times TRUE)}$$

where β represents the strength of the response. We set $\beta = 2$ which produced in a moderate

gradient in the probability of presence across the landscape. The TRUE variable has a linear

gradient in geographic space ranging from -1 to 1 across a square landscape 1024 cells on a side

(Appendix 2 Fig. S1). As a result, the species' probability of occurrence is symmetrically

distributed with an inflection point midway across the landscape. For each of the 100 data

iterations, a Maxent model was trained using values of TRUE plus values of a spatially random

FALSE variable with the range (-1, 1). The FALSE variable represents a variable "mistakenly"

assumed to influence the species' distribution.

Experiment 2: Training sample size

Next, we examined how the number of occurrences used in the training sample affects estimates

of variable importance. We used the same landscape and probability of occurrence as in

Experiment 1. The number of training occurrences was varied across the doubling series 8, 16,

32, … , 1024, but the number of test sites was kept the same (200 occurrences and 10,000

background sites).

Experiment 3: Prevalence

We then explored the effects of prevalence (mean probability of occurrence). The species

responded to TRUE as per a logistic function,

Eq. 2 $$\Pr(occ) = \frac{exp(\beta_1 \times (TRUE - \beta_2))}{1 + exp(\beta_1 \times (TRUE - \beta_2))},$$

where non-zero values of the offset parameter β_2 shift the range across the landscape, thereby

altering prevalence (Appendix 2 Fig. S2). This allows us to manipulate prevalence while not

changing study region extent, which is almost impossible in real-world situations. We set β_1

equal to 2 and chose values of β_2 that varied prevalence in 9 steps from 0.05 to 0.95.

Experiment 4: Study region extent

In real-world situations enlarging a study region typically decreases prevalence (Anderson &

Raza 2010) while also increasing the range of variability in environmental variables (VanDerWal

et al. 2009). In this experiment we isolate the effect of increasing the extent of the study region

on the range of environmental variation in the TRUE variable. Landscape size was varied along

the doubling series 128, 256, 512, …, and 8192 cells on a side, each matched with an increasing

range of the TRUE variable from (-0.125, 0.125) for the smallest landscape to (-8, 8) for the

largest (Appendix 2 Fig. S3). FALSE was spatially randomly distributed and had the range (-1,

1). The species responded to TRUE as per a logistic function (Eq. 1). As a result, increasing

extent did not affect prevalence.

Experiment 5: Spatial resolution and autocorrelation of environmental data

In this experiment we explored the effects of spatial autocorrelation and spatial resolution (grain

size) of environmental data on estimates of variable importance. We began with a linear TRUE

and random FALSE variable distributed across a landscape with 1024 cells on a side. When

unperturbed, the linear gradient of the TRUE variable has a high degree of spatial autocorrelation

because cells close to one another have similar values. To manipulate spatial autocorrelation, we

randomly swapped values of a set proportion of cell values (no cells, or one third, two thirds, or

all of the cells). Swapping values reduces spatial autocorrelation in the TRUE variable because

cells with dissimilar values were more likely to be close to one another (Appendix 2 Fig. S4).

FALSE had a level of spatial autocorrelation no different from random, regardless of swapping.

We assumed the species responded to the environment at the "native" 1/1024th scale of the

landscape. Probability of occupancy was modeled with Eq. 1, and training and test sites were

sampled at this "native" resolution. In some of the simulations we changed the spatial resolution

of the environmental data presented to the SDMs by resampling the environmental rasters to a

finer resolution with 16,384 cells or to a coarser resolution with 64 cells on a side using bilinear

interpolation (sample size was kept the same regardless of resolution). In summary, we created a

landscape, (possibly) swapped cells, modeled the species' distribution and located training and

test sites at the "native" scale, then assigned environmental values to sites based on the landscape

at the (possibly) resampled resolution. This recreates a realistic situation where occurrences

represent the scale of the true response but environmental data used to predict the response are

available at a (potentially) different resolution. We explored all combinations of grain size (cell

size of 1/16,384th, 1/1024th, and 1/64th of the landscape's linear dimension) and spatial

autocorrelation (swapping no cells, or one third, two thirds, or all of the cells; Appendix 2 Fig.

S4).

Experiment 6: Collinearity between environmental variables

Next, we explored the effects collinearity (correlation) between FALSE and TRUE predictors.

As before, the species had a logistic response (Eq. 1) to TRUE, which has a linear gradient

across the landscape. In contrast to previous experiments, FALSE also has a linear trend which is

rotated relative to the gradient in TRUE to alter the correlation between the variables. We used a

circular landscape to ensure no change in the univariate frequencies of the variables with

rotation. The two variables are uncorrelated when their relative rotation is 90°, and positive or

negative if less than or more than 90°, respectively (Appendix 2 Fig. S5). We used rotations of

FALSE relative to TRUE from 22.5 to 157.5° in steps of 22.5°, which produced correlations

between the two variables ranging from -0.91 to 0.91. Both variables had values in the range (-1,

1).

Experiments 7, 8, and 9: Two influential variables

In the final experiments the species' niche was shaped by two influential variables, T1 and T2,

which both have linear gradients on a circular landscape and values in the range (-1, 1). The

species responds to T1 and T2 as per a Gaussian bivariate function:

Eq. 3 $\Pr(occ) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} exp\left(-\frac{(T1-\overline{T1})^2}{2\sigma_1^2(1-\rho^2)} + \rho\frac{(T1-\overline{T1})(T2-\overline{T2})}{\sigma_1\sigma_2(1-\rho^2)} - \frac{(T2-\overline{T2})^2}{2\sigma_2^2(1-\rho^2)}\right).$

Niche breadth in T1 and T2 is determined by σ_1 and σ_2. Importantly, decreasing σ_i increases the

degree to which a variable restricts distribution, meaning that σ_i and variable importance are

inversely related. Changing σ_1 and σ_2 also alters the distribution of the species (Appendix 2 Fig.

S6). Parameter ρ determines the degree of "niche covariance," or interaction between variables in

shaping the niche.

In *Experiment 7* we examined the effects of niche breadth by varying σ_1 and σ_2 across all

combinations of 0.1, 0.3, and 0.5. We set niche covariance $\rho = 0$ and kept T1 and T2

uncorrelated on the landscape. In *Experiment 8* we investigated the effects of niche covariance

by varying ρ from -0.75 to 0 to 0.75. We used intermediate niche breadth (σ_1 and $\sigma_2 = 0.3$) and

kept T1 and T2 uncorrelated. Finally, in *Experiment 9* we explored all combinations of niche

breadth (varying σ_1 and σ_2 from 0.1, 0.3, to 0.5), niche covariance (varying ρ from -0.5, 0, to

0.5), and collinearity between T1 and T2 on the landscape (varying r from -0.71, 0, to 0.71).

Reproducibility

We created the R (R Core Team 2018) package [URL redacted for double-blind peer review] for

generating virtual species and assessing variable importance (available at [URL redacted]). Code

for the experiments and figures in this article is available at [URL redacted]. The package and

code for the experiments depend primarily on the *dismo* (Hijmans et al. 2017), *raster* (Hijmans

2018), and [redacted] packages for R.

Results

In each experiment we assessed six metrics (Box 1). Results from OMNI serve as a benchmark

for the SDM because they represent the best an SDM could be expected to do given only

variation in test data. Thus, when unpermuted predictions from OMNI have poor predictive

accuracy, or when OMNI with permuted predictions fails to discriminate between variables, the

SDM should also fail. Importantly, results for a given level of a manipulation represent outcomes

across multiple data iterations, each of which typically spanned a much smaller range (during

permutation) than the full set of models. Modelers typically have just one set of data for a

species, so variation for a single data instance will underestimate the uncertainty inherent in the

data sampling process.

Experiment 1: Simple scenario

OMNI with unpermuted variables had high predictive accuracy (Fig. 1a). OMNI TRUE and

FALSE permuted did not overlap, meaning that the variables could be successfully

differentiated. Maxent performed similarly, although with more variation around TRUE

permuted compared to OMNI.

Experiment 2: Training sample size

OMNI does not use training data, so always correctly discriminated TRUE from FALSE

regardless of training sample size (Fig. 1b). Maxent unpermuted performed as well as OMNI

unpermuted when sample size was ≥ 64, but below this Maxent had much greater variability than

OMNI and was unable to reliably discriminate between TRUE and FALSE at $n < 32$. At the

smallest sample size ($n = 8$), Maxent often yielded intercept-only models that could not be used

to calculate CBI, which requires variation in predictions for calculation.

Experiment 3: Prevalence

Increasing prevalence (mean probability of occurrence) reduced unpermuted OMNI's predictive

accuracy and increased variability. As a result, OMNI often performed no better than random

when prevalence was ≥ 0.85 (Fig. 2a) and could not discriminate between TRUE and FALSE.

Maxent was qualitatively the same, although variation in permuted and unpermuted CBI was

greater at high prevalence than in OMNI (Fig. 2a).

Experiment 4: Study region extent

Increasing landscape extent (the range of the TRUE variable) caused predictive accuracy of

OMNI unpermuted to peak at intermediate extents where the range of TRUE was from 2 to 4,

which occurs when the landscape was from 1024 to 2048 cells on a side (Fig. 2b). OMNI failed

to reliably discriminate between TRUE and FALSE on the smallest landscapes (range of TRUE

≤0.5). Maxent was less reliable, failing when the range of TRUE was ≤1 (Fig. 2b). The

worsening performance of OMNI unpermuted at large extents might be due to the sensitivity of

CBI to test presences that are located in areas with a very low probability of presence and to

potential underfitting of Maxent (Appendix 10).

Experiment 5: Spatial resolution and autocorrelation of environmental data

The "true" importance of TRUE and FALSE is indicated by OMNI at the "native" resolution of

1/1024 (middle column of subpanels in Fig. 2c). Results for OMNI at the other resolutions

represent the outcome that would be obtained if a modeler had perfect knowledge of the species'

response to the environment but only had data available at finer or coarser resolutions. Both

OMNI and Maxent reliably discriminated between TRUE and FALSE at the "native" 1/1024th

resolution and at the finer 1/6384th resolution, plus at the coarser 1/64th resolution when spatial

autocorrelation was high. However, when the environmental data was coarse (1/64th scale) and

spatial autocorrelation low (proportion swapped two-thirds or 1), both OMNI and Maxent

unpermuted overlapped or nearly overlapped 0, indicating some models performed no better than

random. Neither model could reliably discriminate between TRUE and FALSE in these cases

(cf. Meynard et al. 2019).

Experiment 6: Collinearity between environmental variables: OMNI completely ignores the

FALSE variable and thus had high performance regardless of the magnitude of correlation

between TRUE and FALSE (Fig. 3). Although Maxent unpermuted was slightly more variable

than OMNI, Maxent had fairly high predictive accuracy across the entire range of correlation

between TRUE and FALSE. Maxent also reliably discriminated between TRUE and FALSE

except when collinearity was high ($|r| > 0.71$), although even small amounts of correlation caused

TRUE permuted to vary more than OMNI TRUE permuted. The increasing range of Maxent

FALSE permuted at high magnitudes of correlation indicate that Maxent sometimes used

information in the FALSE variable (Fig. 3).

Experiments 7 through 9: Two influential variables

In *Experiment 7* we manipulated niche breadth of two influential variables while keeping all

other factors "off" (no niche covariance, no correlation). In cases where both variables had equal

influence on the niche ($\sigma_1 = \sigma_2$), OMNI unpermuted had high predictive accuracy across all

combinations of niche breadth in T1 and T2 (median CBI ranging from 0.89 to 0.93; Fig. 4a).

Maxent unpermuted also had high predictive accuracy. Permuting a variable for which niche

breadth is narrow should reduce CBI more than when it is broad, which is what we observed

with OMNI. For example, changing niche breadth from broad ($\sigma_1 = \sigma_2 = 0.5$) to medium ($\sigma_1 = \sigma_2$

$= 0.3$) to narrow ($\sigma_1 = \sigma_2 = 0.1$) reduced median CBI from 0.68 to 0.63 to 0.46, respectively

(similar values were achieved for T2). However, Maxent did not show a monotonic decline;

respective values were 0.68, 0.76, and 0.64 (Fig. 4a). (Permuting T2 yielded similar anomalies).

Thus, Maxent always underestimated the importance of T1 and T2 when niche breadth was

moderate or narrow and variables acted equally.

In cases where variables had asymmetrical influence ($\sigma_1 \neq \sigma_2$), OMNI unpermuted had high

predictive accuracy, and permuting T1 or T2 caused CBI to decrease monotonically with

decreasing niche breadth in the respective variable (Fig. 4a). Maxent always reliably

discriminated between T1 and T2 (i.e., no overlap between distributions of permuted CBI for T1

and T2). However, Maxent tended to overestimate the importance of the more influential

variable. Estimates of importance for the less influential variable tended to be more uncertain

compared to the more influential variable.

In *Experiment 8* we altered niche covariance (interaction between variables) but kept niche

breadth and correlation between variables fixed. Increasing the magnitude of covariance

increased the actual and estimated importance of the variables even though niche breadth was

held constant ($\sigma_1 = \sigma_2 = 0.3$; Fig. 4b). On average, Maxent underestimated the importance of the

variables at all levels of covariance.

In *Experiment 9* we examined all possible combinations of niche breadth and covariance and

correlation between variables. Results were qualitatively similar to the preceding two

experiments (Appendix 4 Fig. S9).

Results using AUC and COR, algorithm-specific tests, and different algorithms

Results using the permute-after-calibration test paired with AUC_{pa}, AUC_{bg}, COR_{pa}, and COR_{bg}

and algorithm-specific tests are summarized in Appendix 3 and presented in detail in Appendices

4 through 9, so are only recapitulated here. We found notable interactions between model

algorithm and the metric used by the permute-after-calibration test. For example, GAMs were

capable of discriminating between TRUE and FALSE at training sample sizes as low as 8

(though few models converged) when using CBI or COR (either variant), but required 128 or

more occurrences when using AUC (either variant). However, results using AUC were well-

calibrated at large sample sizes. Across all experiments, there was no best test statistic, although

AUC was much less variable than CBI, and COR was much more variable. Maximum values of

AUC for unpermuted OMNI models were always substantially <1.

Although Maxent and GAMs performed differently in most experiments, neither consistently

outperformed the other across experiments. GAMs tended to show less variation in simple cases

(TRUE versus FALSE) but more in complex cases (TRUE versus TRUE). In all experiments

BRTs had much greater variation and thus less reliable discrimination than the other two

algorithms.

In general, algorithm-specific tests were less robust to challenging situations than the permute-

after calibration-test. For example, Maxent's change-in-gain test was only able to discriminate

between TRUE and FALSE when sample size was ≥128 (Appendix 9 Fig. S2), but minimum

necessary sample size was 64 using COR or Maxent's permutation or contribution tests, and just

32 when using CBI or AUC.

Discussion

Our objective was to delineate the minimal necessary conditions under which species distribution

models and ecological niche models can be used to infer variable importance. We found that the

permute-after-calibration test was capable of discriminating between variables under many

situations, but results typically differed from expectations established by an "omniscient" model.

Notably, high predictive accuracy did not necessarily connote high inferential capacity.

However, situations that are challenging for generating SDMs with high predictive accuracy

were also challenging for estimating variable importance (results summarized in Table 1): small

sample size, high prevalence, low spatial extent (low environmental variability), high

collinearity, and using environmental data that is coarser than the perceptual scale of the species

when spatial autocorrelation is low. When more than one variable shaped a species' distribution,

SDMs were able to discriminate between two variables with different influence, but mis-

calibrated importance when variables acted equally. Interactions between variables in how they

shaped the niche had little effect on discriminatory power. In general, factors that shape

inference can be classified into those that are extrinsic to the species (e.g., choice of modeling

algorithm, sample size) and those that are intrinsic to the species (e.g., niche breadth). Extrinsic

factors are at least nominally under the control of the modeler and thus offer the potential for

amelioration, whereas confounding intrinsic factors likely require development of new

techniques and robust data to control for their influence. We structure the discussion around what

modelers typically can control and what they typically cannot.

Model algorithm and inferential method

The reliability of tests of variable importance depended on the algorithm, type of test, and metric

used to evaluate the test. We found GAMs and Maxent had much less variability and thus greater

discriminatory capacity than BRTs, despite extensive efforts to tune BRTs (Appendix 1). Choice

of modeling algorithm is one of the largest contributors to variation in predictive capacity

(Dormann et al. 2008; Barbet-Massin et al. 2012; Rapacciuolo et al. 2012), with no one

algorithm necessarily best for all species (Qiao et al. 2015). Our results show that model choice

also affects inferential power and thus underscores the importance of evaluating variable

importance using multiple algorithms.

We also found inferential capacity varied by the nature of the test (permute-after-calibration test

versus algorithm-specific tests) and associated test metric (e.g., CBI, AUC, change in Maxent's

gain). Depending on the situation and algorithm, the choice of test metric (CBI, AUC, COR)

affected the reliability of the permute-after-calibration test, but no one metric consistently out-

performed the others in discrimination capacity. AUC was occasionally better-calibrated than

CBI and COR, especially when paired with GAMs, but also had less reliable discrimination in

these same circumstances. In contrast, algorithm-specific tests were less robust to challenging

circumstances than the permute-after-calibration test (Appendix 3).

Despite the better performance of the permute-after-calibration test, care needs to be used to

ensure the metric with which it is paired provides unconfounded interpretation. Namely, tests of

variable importance can only reliably indicate differences between variables if the original model

(i.e., with unpermuted predictions) has high predictive accuracy (Meinshausen & Bühlmann

2010). Neither AUC nor COR are capable metrics in this respect. In particular, maximum AUC

(either variant) is typically depressed well below 1 (Jiménez-Valverde et al. 2013; Smith 2013a),

which is evident even in our simplest scenario (*Experiment 1*) where median unpermuted AUC_{bg}

and AUC_{pa} for the OMNI model was only ~0.78 and ~0.64, respectively. Likewise, COR does

not indicate the predictive capacity of a model. Given these considerations, we recommend 1)

employing multiple modeling algorithms that can be compared using 2) algorithm-independent

tests like the permute-after-calibration test; and 3) employing metrics that can be objectively

interpreted as measures of predictive accuracy and that are not known to be influenced by study-

specific aspects like prevalence or sample size (Jiménez-Valverde et al. 2013; Jiménez-Valverde

2020; Smith 2013a).

Sample size

We found that inferential power was compromised when training sample size was between 8-

128, depending on the algorithm and test. However, owing to the fact that simulations ensured

other conditions were optimal (e.g., no dispersal, perfect detection), we expect the actual

minimum sample size for reliable application real-world situations to be larger, especially given

biases in data collection (Feeley & Silman 2011). Although new techniques amenable to

modeling rare species (Lomba et al. 2010; Breiner et al. 2015) might be able to lower the

threshold sample size necessary for predictive accuracy, small samples can still induce spurious

correlations between predictors (Ashcroft et al. 2011) and may not adequately capture the full

extent of species' environmental tolerances (Feeley & Silman 2011). On these bases, we expect

that minimal sample size for reliable inference will be several times larger than sample size

necessary for generating models with high predictive accuracy.

Spatial scale: Extent, prevalence, resolution, and autocorrelation

We found inferential power declined rapidly when prevalence was >0.5 (Fig. 2a) and when the

study region extent was too small to encompass sufficient environmental variation to distinguish

occurrences from non-occurrences (Fig. 2b). In real-world situations, decisions regarding spatial

extent of the study region typically affect prevalence, the range of environmental variability in

training data, and degree of spatial autocorrelation and collinearity between predictors (Seo et al.

2008; VanDerWal et al. 2009; Lauzeral et al. 2013). Thus, there are likely interactions and

cascading effects of decisions about scale that are not apparent in our results. For example, study

extent can interact with spatial autocorrelation to affect the variables that appear important in a

model (VanDerWal et al. 2009; Connor et al. 2017).

We also found that inference was compromised when spatial resolution was coarser than the

species' scale of perception and spatial autocorrelation was low (Fig. 2c). Best practices

recommend using environmental data at a resolution matching the scale of species' response to

the environment (Mertes & Jetz 2018; Araújo et al. 2019), although scale mismatch can be

ameliorated when spatial autocorrelation is high (Fig. 2c; Moudrý & Šímová 2012; Mertes &

Jetz 2018). To date, the finest resolution climate data with global-scale coverage has a resolution

on the order of ~1 km (Fick & Hijmans 2017; Karger et al. 2017), which is much larger than the

scale of perception of the environment of most sessile and many mobile organisms. Hence, scale

mismatch will likely remain a problem for many studies.

Based on our results, we recommend at the minimum ensuring the region from which

background sites are drawn is large enough to encompass sufficient environmental variation and

that the species occupies less than about half the landscape. Likewise, when fine-scale

environmental data is not available, we recommend at least measuring spatial autocorrelation to

assess the degree to which scale mismatch could confound inference (Naimi et al. 2014).

Modelers must be aware that the results of an inferential study will be dependent on all aspects

of scale and that these aspects can interact to affect inference in ways not explored here (e.g.,

VanDerWal et al. 2009; Hanberry 2013; Connor et al. 2017). As a result, comparisons between

inferential studies that vary in aspects of scale need to be made with these complications in mind.

Collinearity

Our results indicate that inferential power is low when the magnitude of pairwise correlation is

>0.7 (Fig. 3). Alarmingly, unpermuted predictions often had high predictive accuracy even when

high collinearity caused them to mistakenly use information in the FALSE variable. This is

surprising but supported by other work that finds using predictors with no actual relationship to a

species' occurrence can yield models as accurate when using "real" variables (Buklin et al. 2015;

Fourcade et al. 2018). Thus, the predictive accuracy of a model is not a reliable indicator of its

inferential capacity.

Of all of our findings, the inability of models to differentiate between influential and

uninfluential correlated variables, yet produce seemingly accurate predictions is the most

troubling (Warren et al. 2020). Environmental variables are often collinear (Jiménez-Valverde et

al. 2009), so this is likely a very frequent challenge to successful inference. However, modelers

have some means to modulate collinearity. The simplest solution is to simply select variables

that have low pairwise-correlations (Dorman et al. 2013). Unfortunately, discarding correlated

variables inherently assumes dropped variables have zero influence with absolute uncertainty.

Another solution is to employ modeling algorithms with regularization or regularization-like-

behavior, but the methods used here already do that (e.g., Maxent LASSO; Tibshirani 1996;

Phillips et al. 2006) and were not entirely robust to collinearity (Dormann et al. 2013). A third

potential solution may be to construct multiple models with different sets of relatively

uncorrelated variables (Barbet-Massin et al. 2014; Petitpierre et al. 2017) then average variables'

importance across them.

Qualities of the niche

Niche breadth and interactions between variables in shaping the niche are inherent to species and

thus not under control by the modeler. Niche breadth has the most obvious relationship to

variable importance since narrower environmental tolerance should translate into increased

sensitivity of a model to changes in that variable. Surprisingly, we found that when two variables

act to shape the niche equally, reducing niche breadth does not lead to a monotonic increase in

estimated importance of the variables (Fig. 4a). Likewise, when two variables acted unequally to

influence the niche, SDMs overestimated the importance of the more important variable. As a

result, the relative difference between the permuted and unpermuted values of a test statistic

should not be interpreted as a measure of the absolute importance of a variable. Rather, we

recommend interpreting only qualitative (rank) importance (Barbet-Massin & Jetz 2014).

Niche covariance occurs when, for example, negative effects of high temperature on a species'

fitness can be offset by high values of precipitation (Smith 2013b). Interaction between niche

dimensions rotates the orientation of the niche in niche space, thereby changing the range of

environments occupied (Appendix 2 Fig. S7). As a result, the importance of niche covariance are

not always obvious from examination of univariate niche breadth (Smith 2013b). We found that

increasing the magnitude of niche covariance ($\rho \neq 0$) increased actual and estimated importance

compared to cases where variables acted independently ($\rho = 0$), but importance was still

miscalibrated vis-à-vis an omniscient model (Fig. 4b). We did not find strong interactions

between niche breadth, niche covariance, and collinearity for the range of each investigated here,

although the simplicity of our simulations does not preclude different outcomes in real-world

situations.

Variable and model selection

Our work calls into question the common practice of using automated methods for variable and

model selection (Barbet-Massin & Jetz 2014; Gobeyn et al. 2017; Guisande et al. 2017; Cobos et

al. 2019). We found SDMs using uninfluential variables could still yield measures of predictive

accuracy that qualified them as "good" models (Fig. 3; Buklin et al. 2015; Fourcade et al. 2018).

As a result, we echo others' recommendations to use expert-based selection of variables before

conducting algorithmic-based screening (Mod et al. 2016; Gardener et al. 2019).

Future directions

The simplified nature of our scenarios likely means that conditions we identify for reliable

inference (Table 1) represent the minimum circumstances under which these tests perform

robustly. Real-world applications will surely require larger sample sizes, less collinearity,

smaller disparities in scale, et cetera, to be reliable. Given the many ecological questions

informed by measures of variable importance, understanding the domain in which inferential

tests can be trusted is a pressing priority. To this end, many questions must be addressed: How

do tests of variable importance fare against real-world factors like biotic interactions, local

adaptation, disturbance, dispersal, bias in sampling, realistic environmental variation, and so on?

How does high-dimensional niche space affect inference? How do other tests of importance

compare to the ones evaluated here? How does data type (presence/background versus

presence/absence versus abundance) affect inference (Gábor et al. 2020)? Answering these

questions will require expanding beyond the reductionist approach used in this work. One

alternative is to simulate niches or distributions as realistically as possible, including

realistically-structured landscapes, biotic interactions, dispersal limitation, and other ecological

processes (e.g., Zurell et al. 2016; Warren et al. 2020), then apply a battery of procedures to

identify situations that are conducive to measuring variable importance accurately (Groves &

Lempert 2007 describe an analogous approach in policy analysis). Alternatively, the small subset

of Earth's species for which there is extensive field-based knowledge of range-shaping factors

could be used to validate model-based inferences of variable importance (e.g., Angert et al.

2018).

Conclusions

Our work represents the first systematic assessment of conditions under which SDMs can

reliably estimate variable importance. The good news is that SDMs were able to discriminate

between variables under conditions conducive to generating models with high predictive

accuracy. The bad news is that high predictive accuracy did not necessarily connote reliable

inference (cf. Warren et al. 2020). Factors extrinsic to species that can be influenced by modelers

and factors intrinsic to species affect the ability to measure variable importance. Given the

ubiquity with which these models are used to measure the importance of environmental factors in

shaping species' distributions and niches (Bradie & Leung 2017), we see a great opportunity and

a great need for further research in this area.

Acknowledgements

We wish to thank three anonymous reviewers and the subject editor who dedicated the

unreimbursed time and attention to improve the manuscript. This work was supported by the

[redacted] to [redacted].

Literature cited

Angert, A.L., Bayly, M., Sheth, S.N., and Paul, J.R. 2018. Testing range-limit hypotheses using

range-wide habitat suitability and occupancy for the Scarlet Monkeyflower (*Erythranthe*

cardinanis). American Naturalist 191:E76-E89.

Anderson, R.P. and A. Raza. 2010. The effect of the extent of the study region on GIS models

of species geographic distributions and estimates of niche evolution: Preliminary tests with

montane rodents (genus *Nephelomys*) in Venezuela. Journal of Biogeography 37:1378-1393.

Araújo, M.B, Anderson, R.P., Barbosa, A.M., Beale, C.M., Dormann, C.F., Early, R., Garcia,

R.A., Guisan, A., Maiorano, L., Naimi, B., O'Hara, R.B., Zimmermann, N.E., and Rahbek, C.

2019. Standards for distribution models in biodiversity assessments. Science Advances

5:eaat4858.

Ashcroft, M.B., French, K.O., and Chisholm, L.A. 2011. An evaluation of environmental

factors affecting species distributions. Ecological Modeling 222:524-531.

Barbet-Massin, M. and Jetz, W. 2014. A 40-year, continent-wide, multispecies assessment of

relevant climate predictors for species distribution modeling. Diversity and Distributions

20:1285-1295.

Barbet-Massin, M., F. Jiguet, C.H. Albert, and W. Thuiller. 2012. Selecting pseudo-absences

for species distribution models: How, where and how many? Methods in Ecology and

Evolution 3:327-338.

Boyce, M.S., Vernier, P.R., Nielsen, S.E., and Schmiegelow, F.K.A. 2002. Evaluating resource

selection functions. Ecological Modeling 157:281-300.

Bradie, J. and Leung, B. 2017. A quantitative synthesis of the importance of variables used in

MaxEnt species distribution models. Journal of Biogeography 44:1344-1361.

Breiman, L. 2001. Random forests. Machine Learning 45:5-32.

Breiner, F.T., Guisan, A., Bergamini, A., and Nobis, M.P. 2015. Overcoming limitations of

modeling rare species by using ensembles of small models. Methods in Ecology and

Evolution 6:1210-1218.

Buklin, D.N., Basille, M., Benscoter, A.M., Brandt, L.A., Mazotti, F.J., Romañach, S.S.,

Speroterra, C., and Watling, J.I. 2015. Comparing species distribution models constructed

with different subsets of environmental predictors. Diversity and Distributions 21:23-35.

Cobos, M.E., Peterson, A.T., Osorio-Olvera, L., and Jiménez-García, D. 2019. An exhaustive

analysis of heuristic methods for variable selection in ecological niche modeling and species

distribution modeling. Ecological Informatics 53:100983.

Connor, T., Hull, V., Viña, A., Shortridge, A., Tang, Y., Zhang, J., Wang, F., and Liu, J. 2017.

Effects of grain size and niche breadth on species distribution modeling. Ecography

41:1270-1282.

Dormann, C.F., Purchke, O., Márquez, J.R.G., Lautenbach, S., and Schroder, B. 2008.

Components of uncertainty in the species distribution analysis: A case study of the Great

Gray Shrike. Ecology 89:3371-3386.

Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl G., Carré, G., Marquéz, J.R.G., Gruber,

B,. Lafourcade, B., Leitão, P.J., Münkemüller, T., McClean, C., Osborne, P.E., Reineking,

B., Schröder, B., Skidmore, A.K., Zurell, D., and Lautenbach, S. 2013. Collinearity: A

review of methods to deal with it and a simulation study evaluating their performance.

Ecography 36:27-46.

Elith, J., C.H. Graham, R.P. Anderson, M. Dudík, S. Ferrier, A. Guisan, R.J. Hijmans, F.

Huettmann, J.R. Leathwick, A. Lehmann, J. Li, L.G. Lohmann, B.A. Loiselle, G. Manion, C.

Moritz, M. Nakamura, Y. Nakazawa, J.McC. Overton, A.T. Peterson, S.J. Phillips, K.

Richardson, R. Scachetti-Pereira, R.E. Schapire, J. Soberón, S. Williams, M.S. Wisz, and

N.E. Zimmermann. 2006. Novel methods improve prediction of species' distributions from occurrence data. Ecography 29:129-151.

Elith, J., J.R. Leathwick, and T. Hastie. 2008. A working guide to boosted regression trees. Journal of Animal Ecology 77:802-813.

Feeley, K.J. and Silman, M.R. 2011. Keep collecting: Accurate species distribution modeling requires more collections than previously thought. Diversity and Distributions 17:1132-1140.

Fick, S.E. and R.J. Hijmans, 2017. Worldclim 2: New 1-km spatial resolution climate surfaces for global land areas. International Journal of Climatology 37:4302-4315.

Fourcade, Y., Besnard, A.G., and Secondi, J. 2018. Paintings predict the distribution of species, or the challenge of selecting environmental predictors and evaluation statistics. Global Ecology and Biogeography 27:245-256. Francis, A.P and D.J. Currie. 2003. A globally consistent richness-climate relationship for angiosperms. American Naturalist 161:523-536.

Gábor, L., Moudrý, V., Lecours, V., Malavasi, M., Barták, V., Fogl, M., Šímová, P., Rocchini, D., and Václavík, T. 2020. The effect of positional error on fine scale species distribution models increases for specialist species. Ecography 43:256-269.

Gardener, A.S., Mclean, I.M.D., and Gaston, K.J. 2019. Climatic predictors of species distributions neglect biophysiologically meaningful variables. Diversity and Distributions 25:1273-1288.

Groves, D.G., and Lempert, R.J. 2007. A new analytic method for finding policy-relevant scenarios. Global Environmental Change 17:73-85.

Guevara, L., Gerstner, B.E., Kass, J.M., and Anderson, R.P. 2018. Toward ecologically realistic

predictions of species distributions: A cross-time example from tropical montane cloud

forests. Global Change Biology 24:1511-1522.

Guisande, C., García-Roselló, Heine, J., González-Dacosta, González Vilas, L., García Pérez,

B.J., and Lobo, J.M. 2017. SPEDInstandR: An algorithm based on a fluctuation index for

selecting predictors in species distribution modeling. Ecological Informatics 37:18-23.

Hanberry, B.B. 2013. Finer grain size increases effects of error and changes influence of

environmental predictors on species distribution models. Ecological Informatics 15:8-13.

Hargreaves, A.L., Samis, K.E., and Eckert, C.G. 2014. Are species' range limits simply niche

limits writ large? A review of transplant experiments beyond the range. American Naturalist

183:157-173.

Hijmans, R.J. 2019. raster: Geographic Data Analysis and Modeling. R package version 2.9-23.

https://CRAN.R-project.org/package=raster.

Hijmans, R.J., Phillips, S.J., Leathwick, J., and Elith, J. 2017. dismo: Species Distribution

Modeling. R package version 1.1-4. https://CRAN.R-project.org/package=dismo.

Hirzel, A.H., Le Lay, G., Helfer, V., Randin, C., and Guisan, A. 2006. Evaluating the ability of

habitat suitability models to predict species presences. Ecological Modeling 199:142-152.

Jiménez-Valverde, A. 2020. Sample size for the evaluation of presence-absence models.

Ecological Indicators 114:106289.

Jiménez-Valverde, A., Nakazawa, Y., Lira-Noriega, A., and Peterson, A.T. 2009.

Environmental correlation structure and ecological niche projections. Biodiversity

Informatics 6:28-35.

Jiménez-Valverde, A., Acevedo, P., Barbosa, M., Lobo, J.M., and Real, R. 2013.

Discrimination capacity in species distribution models depends on the representativeness of

the environmental domain. Global Ecology and Biogeography 22:508-516.

Júnior, P.D.M. and Nóbrega, C.C. 2018. Evaluating collinearity effects on species distribution

models: An approach based on virtual species simulation. Public Library of Science ONE

13:e0202403.

Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., Zimmermann,

N.E., Linder, H.P. & Kessler, M. 2017 Climatologies at high resolution for the earth's land

surface areas. Scientific Data 4:170122.

Lauzeral, C., Grenouillet, G., and Brosse, S. 2013. Spatial range shape drives grain size effects

in species distribution models. Ecography 36:778-787.

Lee-Yaw, J.A., Kharouba, H.M., Bontrager, M., Mahony, C., Csergö, A.M., Noreen, A.M.E., Li,

Q., Schuster, R., and Angert, A.L. 2016. A synthesis of transplant experiments and

ecological niche models suggest that range limits are often niche limits. Ecology Letters

19:710-722.

Lomba, A., L. Pellissier, C. Randin, J. Vicente, J. Horondo, and A. Guisan. 2010. Overcoming

the rare species modeling complex: A novel hierarchical framework applied to an Iberian

endemic plant. Biological Conservation 143:2647-2657.

Meinshausen, N. & Bühlmann, P. 2010. Stability selection. Journal of the Royal Statistical

Society: Statistical Methodology Series B 72:417-473.

Mertes, K. and Jetz, W. 2018. Disentangling scale dependencies in species environmental

niches and distributions. Ecography 41:1604-1615.

Meynard, C.N. and D.N. Kaplan. 2013. Using virtual species to study species distributions and

model performance. Journal of Biogeography 40:1-8.

Meynard, C.N., Leroy, B., and Kaplan, D.M. 2019. Testing methods in species distribution

modelling using virtual species: what have we learnt and what are we missing? Ecography

42:2021-2036.

Mod, H.K., Scherrer, D., and Guisan, A. 2016. What we use is not what we know:

Environmental predictors in plant distribution models. Journal of Vegetation Science

27:1308-1322.

Moudrý, V. and Šímová, P. 2012. Influence of positional accuracy, sample size and scale on

modeling species distributions: A review. International Journal of Geographic Information

Science 26:2083-2095.

Naimi, B., Hamm, N.A.S., Groen, T.A., Skidmore, A.K., and Toxopeus, A.G. 2014. Where is

positional uncertainty a problem for species distribution modeling? Ecography 57:191-203.

Norberg, A., Abrego, N., Blanchet, F.G. Adler, F.R., Anderson, B.J., Anttila, J., Araújo, M.B.,

Dallas, T., Dunson, D., Elith, J., et al. 2019. A comprehensive evaluation of predictive

performance of 33 species distribution models at species and community levels. Ecological

Monographs 89:e01370.

Petitpierre, B., Broennimann, O., Kueffer, C., Daehler, C., and Guisan, A. 2017. Selecting

predictors to maximize the transferability of species distribution models: Lessons from cross-

continental plant invasions. Global Ecology and Biogeography 26:275-287.

Phillips, S.J. and Dudík, M. 2008. Modeling species distributions with Maxent: New extensions

and a comprehensive evaluation. Ecography 31:161-175.

Phillips, S.J., Anderson, R.P., and Schapire, R.E. 2006. Maximum entropy modeling of species

geographic distributions. Ecological Modelling 190:231-259.

Qiao, H., Soberón, J., and Peterson, A.T. 2015. No silver bullets in correlative ecological niche

modeling: Insights from testing among many potential algorithms for niche estimation.

Methods in Ecology and Evolution 6:1126-1136.

R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org.

Rapacciuolo, G., Roy, D.B., Gillings, S., Fox, R., Walker, K., and Purvis, A. 2012. Climatic

associations of British species distributions show good transferability in time but low

predictive accuracy for range change. Public Library of Science ONE 7:e40212.

Roberts, D.R., Bahn, V., Ciuti, S., Boyce, M.S., Elith, J., Guillera-Arroita, G., Hausentein, S.,

Lahoz-Monfort, J.J., Schröder, B., Thuiller, W., Warton, D.I., Wintle, B.A., Hartig, F., and

Dormann, C.F. 2017. Cross-validation strategies for data with temporal, spatial,

hierarchical, or phylogenetic structure. Ecography 40:913-929.

Seo, C., Thorne, J.H., Hannah, L., and Thuiller, W. 2008. Scale effects in species distribution

models: Implications for conservation planning under climate change. Biology Letters 5:39-

43.

Smith, A.B. 2013a. On evaluating species distribution models with random background sites in

place of absences when test presences disproportionately sample suitable habitat. Diversity

and Distributions 19:867-872.

Smith, A.B. 2013b. The relative influence of temperature, moisture and their interaction on

range limits of mammals over the past century. Global Ecology and Biogeography 22:334-

343.

Smith, A.B. 2019. enmSdm: Tools for modeling species niches and distributions. R package

version 0.3.8.2. URL https://github.com/adamlilith/enmSdm.

Smith, A.B., M.J. Santos, M. S. Koo, K.C. Rowe, K.M.C. Rowe, J.L. Patton, S. Beissinger, and

C. Moritz. 2013. Evaluation of species distribution models by resampling of sites surveyed

a century ago by Joseph Grinnell. Ecography 36:1017-1031.

Tibshirani, R. 1996. Regression shrinkage and selection via the Lasso. Journal of the Royal

Statistical Society, Series B 58:267-288.

VanDerWal, J., Shoo, L.P., Graham, C., and Williams, S.E. 2009. Selecting pseudo-absence

data for presence-only distribution modeling: How far should you stray from what you

know? Ecological Modeling 220:589-594.

Warren, D.L., Matzke, N.J., and Iglesias, T.L. 2020. Evaluating presence‐only species

distribution models with discrimination accuracy is uninformative for many applications.

Journal of Biogeography 47:167-180.

White, J.W., Rassweiler, A., Samhouri, J.F., Stier, A.C., and White, C. 2014. Ecologists should

not use statistical significance tests to interpret simulation model results. Oikos 123:385-388.

Wood, S.N. 2006. Generalized Additive Models: An Introduction with R. Chapman and

Hall/CRC, Boca Raton. 392 pp.

Zurell, D., Thuiller, W., Pagel, J., Cabral, J.S., Münkenmüller, T., Gravel, D., Dullinger, S.,

Normand, S., Schiffers, K.H., Moore, K.A., and Zimmermann, N.E. 2016. Benchmarking

novel approaches for modeling species range dynamics. Global Change Biology 22:2651-

2664.

Table 1. Summary of results across the nine simulation experiments for the permute-after-calibration test. Discrimination refers to the ability to differentiate between two variables with different influence. Few tests met our quantitative standard for reliable calibration, so here "good calibration" refers to a subjective comparison between the tests result and the results using an "omniscient" model (i.e., OMNI). Asterisks indicate the algorithm sometimes had convergence issues or yielded intercept-only models which disallowed calculation of test statistics. Results in this table are particular to using the permute-after-calibration test paired with CBI. Thresholds using other metrics or algorithm-specific tests were sometimes different, but recommendations remain unchanged.

Experiment	OMNI	GAMs	Maxent	BRTs	Recommendations
1 Simple	No issues.	Reliable discrimination, mediocre calibration.	Reliable discrimination, mediocre calibration.	Reliable discrimination, worst calibration.	Assess robustness of estimates of variable importance using multiple algorithms.
2 Sample size	No issues.	Reliable discrimination at all n,	Unreliable discrimination at $n < 32$,	Unreliable discrimination at $n < 64$, poor	For measuring importance, adequate sample size may be larger than that needed for generating accurate models.

		unreliable calibration at n<32.*	unreliable calibration at n <128.*	calibration at all n.*	
3 Prevalence	Increasingly poor predictive accuracy as prevalence increases, especially >0.5, poor discrimination at prevalence >0.75.	Unreliable discrimination and poor calibration at prevalence > 0.75.	Unreliable discrimination and poor calibration at prevalence > 0.75.	Unreliable discrimination at prevalence >0.62, poor calibration when prevalence >0.15.*	Ensure species occupy <60-70% (preferably <50%) of the space from which training background sites can be drawn.
4 Extent	Highest predictive accuracy at	Unreliable discrimination at small	Unreliable discrimination and calibration	Unreliable discrimination at small extents.	Use study regions that encompass sufficient amount of variation in predictors; if unsure, err in direction of

	medium extents, poor discrimination at small extents.	extents, unimodal relationship between calibration and extent.*	at small extents. More sensitive to small extent than GAMs. *	Calibration always poor. More sensitive to small extent than GAMs. *	using larger regions/variables with sufficient variation.
5 Spatial resolution and autocorrelation	Poor predictive accuracy and discrimination when resolution is coarse and autocorrelation low.	Unreliable discrimination and calibration when resolution is coarse relative to niche breadth and environmental autocorrelation low.			Use environmental data at spatial resolutions that match or are finer than the scale of habitat selection of the species. If only coarse data is available, ensure spatial autocorrelation is high to reduce effects of scale disparity.
6 Collinearity	No issues.	Unreliable discrimination and	Unreliable discrimination when $|r| > 0.71$,	Unreliable discrimination when $r < -0.71$	Use variables with demonstrated effects on fitness, physiology, and/or population growth. Ensure variable

	calibration poor when $|r|$ >0.71.	poor calibration for all r.	or $r > 0.38$, poorest calibration for all r.	pairs have correlations such that $|r|<0.7$. High model performance does not connote reliable inference.
7 Niche breadth	No issues.	When variables act equally, inverse relationship between estimated importance and niche breadth. When variables act unequally, stronger variable overestimated. Reliable discrimination in all cases. Maxent has better calibration than GAMs and BRTs.		Interpret only qualitative (rank) differences, not quantitative differences, between variables.
8 Niche covariance	Importance declines with increasing magnitude of covariance.	As OMNI but underestimates importance relative to OMNI.	As Maxent but more variable.	As Maxent but greatest variability.	Compare models with/out interaction terms to estimate effect of covariance.

9 Collinearity,   Qualitatively the same as *Experiments 7* and *8.*

niche breadth,

covariance

Figure 1. (a) *Experiment 1: Simple scenario*. The species' range is influenced solely by a TRUE

variable, but data on an uninfluential FALSE variable is also "mistakenly" presented to models.

Maxent successfully discriminates between the two variables (no overlap between green and red

bars). See Box 1 for further interpretation. (b) *Experiment 2: Sample size*. Maxent cannot

differentiate between TRUE and FALSE at *n* <32. Asterisks indicate cases where OMNI and

Maxent reliably discriminate between TRUE and FALSE. Bar width is proportional to the

number of Maxent models that are more than intercept-only models (typically *n* = 100; CBI

cannot be calculated if there is no variation the response). See Box 1 for further guidance on

interpretation.

Figure 2. (a) *Experiment 3: Prevalence*—Effect of prevalence on inferential power. Neither

Maxent nor OMNI reliably discriminate when prevalence is >0.75. (b) *Experiment 4: Study*

region extent—Maxent measures variable importance most reliably when the study region is

large enough to encompass sufficient environmental variation. Study region extent is indicated

by the range of the TRUE variable which is proportional to the size of the landscape. (c)

Experiment 5: Spatial resolution and autocorrelation—The species perceives the environment at

a "native" resolution such that cells are 1/1024th of the linear dimension of the landscape.

Environmental data were downscaled to cells 1/16,384th on a side or upscaled to 1/64th on a

side. Spatial autocorrelation was decreased by randomly swapping cell values of 1/3, 2/3 or all of

the cells. Maxent fails when resolution is coarse and autocorrelation low. In all panels asterisks

indicate both OMNI and Maxent reliably discriminate between TRUE and FALSE. See Box 1

for further guidance on interpretation.

Figure 3. *Experiment 6: Collinearity*. The species' range is determined by a single TRUE

variable, but Maxent is presented data on this variable plus a correlated FALSE variable. FALSE

is increasingly used by Maxent as the magnitude of correlation increases. Asterisks indicate both

OMNI and Maxent reliably discriminate between TRUE and FALSE. See Box 1 for further

guidance on interpretation.

Figure 4. (a) *Experiment 7: Niche breadth*. Each subpanel represents results from modeling a

species on a landscape with two influential variables, T1 and T2, with niche breadth set by σ_1

and σ_2. Narrower niche breadth increases limitation by that variable so should yield lower CBI

when the variable is permuted. The y-axis on each subpanel represents CBI. The case shown

here is for a landscape with no correlation between variables ($r = 0$) and no niche covariance (ρ

$= 0$). Asterisks indicate both OMNI and Maxent reliably discriminate between T1 and T2. (b)

Experiment 8: *Niche covariance*. Variables interact to define the niche. The case shown here is

for a landscape with no correlation ($r = 0$) with moderate and equal niche breadth in both

variables ($\sigma_1 = \sigma_2 = 0.3$). See Box 1 for further interpretation.

Online appendices

Appendix 1: Model calibration

Appendix 2: Maps of landscapes and species

Appendix 3: Summary of results for CBI, AUC, COR, and algorithm-specific tests

Appendix 4: Using AUC calculated with presences and absences (AUC_{pa}) as a test statistic for

the permute-after-calibration test of variable importance

Appendix 5: Using AUC calculated with presences and absences (AUC_{bg}) as a test statistic for

the permute-after-calibration test of variable importance

Appendix 6: Using the correlation test statistic calculated with presences and absences (COR_{pa})

as a test statistic for the permute-after-calibration test of variable importance

Appendix 7: Using the correlation test statistic calculated with presences and absences (COR_{bg})

as a test statistic for the permute-after-calibration test of variable importance

Appendix 8: Algorithm-specific metrics of variable importance

Appendix 9: Sensitivity of the Continuous Boyce Index to improbable test presences