Functional innovation in the evolution of the calcium-dependent system of the eukaryotic endoplasmic reticulum

Daniel E. Schäffer[1,2†], Lakshminarayan M. Iyer [1], A. Maxwell Burroughs[1], L. Aravind[1*]

[1]National Center for Biotechnology Information, National Library of Medicine, National Institutes of
Health, Bethesda, MD, USA

[2]Science, Mathematics, and Computer Science Magnet Program, Montgomery Blair High School,
Silver Spring, MD, USA.

* Correspondence:
L. Aravind
aravind@mail.nih.gov

†School of Computer Science, Carnegie Mellon University, Pittsburgh, PA, USA

Keywords: calcium binding, calcium stores, calmodulin, channels, eukaryote origins,
endomembranes, SOCE pathway, wolframin.

Abstract

The origin of eukaryotes was marked by the emergence of several novel subcellular systems. One
such is the calcium (Ca^{2+})-stores system of the endoplasmic reticulum, which profoundly influences
diverse aspects of cellular function including signal transduction, motility, division, and
biomineralization. We use comparative genomics and sensitive sequence and structure analyses to
investigate the evolution of this system. Our findings reconstruct the core form of the Ca^{2+}- stores
system in the last eukaryotic common ancestor as having at least 15 proteins that constituted a basic
system for facilitating both Ca^{2+} flux across endomembranes and Ca^{2+}-dependent signaling. We
present evidence that the key EF-hand Ca^{2+}-binding components had their origins in a likely bacterial
symbiont other than the mitochondrial progenitor, whereas the protein phosphatase subunit of the
ancestral calcineurin complex was likely inherited from the asgardarchaeal progenitor of the stem
eukaryote. This further points to the potential origin of the eukaryotes in a Ca^{2+}-rich biomineralized
environment such as stromatolites. We further show that throughout eukaryotic evolution there were
several acquisitions from bacteria of key components of the Ca^{2+}-stores system, even though no
prokaryotic lineage possesses a comparable system. Further, using quantitative measures derived
from comparative genomics we show that there were several rounds of lineage-specific gene
expansions, innovations of novel gene families, and gene losses correlated with biological innovation
such as the biomineralized molluscan shells, coccolithophores, and animal motility. The burst of
innovation of new genes in animals included the wolframin protein associated with Wolfram
syndrome in humans. We show for the first time that it contains previously unidentified Sel1, EF-
hand, and OB-fold domains, which might have key roles in its biochemistry.

1   Introduction

The emergence of a conserved endomembrane system marks the seminal transition in cell structure
that differentiates eukaryotes from their prokaryotic progenitors (Jekely, 2007). This event saw the
emergence of a diversity of eukaryotic systems and organelles such as the nucleus, the endoplasmic
reticulum (ER), vesicular trafficking, and several, novel signaling systems that are uniquely
associated with this sub-cellular environment. A major eukaryotic innovation in this regard is the
intracellular ER-dependent calcium (Ca^{2+})-stores system that regulates the cytosolic concentration of
Ca^{2+} (Ashby and Tepikin, 2001). Although Ca^{2+} ions are maintained at a 4000 to 10,000-fold higher
concentration in the endoplasmic reticulum (ER) lumen as compared to the cytoplasm (Woo et al.,
2018), upon appropriate stimulus they are released into the cytoplasm by Ca^{2+}-release channels such
as the inositol trisphosphate receptors (IP3Rs) and the ryanodine receptors (RyRs). The process is
then reversed and Ca^{2+} is pumped back into the ER by the ATP-dependent SERCA
(sarcoplasmic/endoplasmic reticulum calcium ATPase) pumps, members of the P-type ATPase
superfamily (Ashby and Tepikin, 2001; Altshuler et al., 2012). In addition to the above core
components that mediate the flux of Ca^{2+} from and into the ER-dependent stores, several other
proteins have been linked to the regulation of this process and transmission of Ca^{2+}-dependent
signals, including: 1) chaperones such as calreticulin, calnexin, and calsequestrin in the ER lumen
(Kozlov et al., 2010); 2) diverse EF-hand proteins such as calmodulin (and its relatives), calcineurin
B, and sorcin that bind Ca^{2+} and regulate the response to the Ca^{2+} flux (Denessiouk et al., 2014); 3)
channel proteins such as the voltage-gated calcium channels (VGCC) and trimeric intracellular cation
channels (TRIC) that influence the flow of Ca^{2+} (Lanner et al., 2010; Zhou et al., 2014); and 4)
protein kinases (calcium/calmodulin-dependent kinases (CaMKs)) and protein phosphatases
(calcineurin A) that mediate the Ca^{2+}-dependent signaling response (Berridge, 2012). Together, the
Ca^{2+}-stores system and intracellular Ca^{2+}-dependent signaling apparatus regulate a variety of cellular
functions required for eukaryotic life, such as transcription, cellular motility, cell growth, stress
response, and cell division (Clapham, 2007; Berridge, 2012; Krebs et al.).

Comparative evolutionary analyses of the proteins in the ER Ca^{2+}-stores and -signaling system have
revealed that some components were either present in the last eukaryotic common ancestor (LECA)
(e.g. Calmodulin and SERCA) or derived early in the evolution of the eukaryotes (e.g. IP3R) (Nolan
et al., 1994; Moreno and Docampo, 2003; Reiner et al., 2003; Prole and Taylor, 2011; Plattner and
Verkhratsky, 2013; Verkhratsky and Parpura, 2014; Perez-Gordones et al., 2015). Other proteins
show a patchier distribution in lineages outside of metazoans (e.g. calreticulin and calnexin) (Moreno
and Docampo, 2003; Banerjee et al., 2007), or were reconstructed to have been derived in lineages
closely related to the metazoans (e.g. RyR, which diverged from the ancestor of IP3R at the base of
filozoans) (Alzayady et al., 2015). A substantial number of the components that have been studied in
this system are primarily found in the metazoans, with no identifiable homologs outside of metazoa
(Cai et al., 2015). Most studies have focused on animal proteins of these systems, highlighting the
general lack of knowledge regarding the regulation of ER Ca^{2+}-stores and the potential diversity in
the regulatory systems present in other eukaryotes. To our knowledge, a systematic assessment of the
evolutionary origins of the entire Ca^{2+}-stores system and its regulatory components, as currently
understood, has yet to be attempted.

Given our long-term interest in the origin and evolution of the eukaryotic subcellular systems, we
conducted a comprehensive analysis of the core and regulatory components of the Ca^{2+}-stores
system, analyzing their known and predicted interactions and inferring the evolutionary depth of
various components. We show that an ancient core of at least 15 protein families was already in place
at the stem of the eukaryotic lineage. Of these, a subset of proteins is of recognizable bacterial

ancestry, although there is no evidence of a bacterial Ca^{2+}-stores system resembling those in
eukaryotes. We also show that gene loss and lineage-specific expansions of these components shaped
the system in different eukaryotic lineages, and sometimes corresponds to recognized adaptive
features unique to particular organisms or lineages. Further, we conducted a systematic domain
analysis of the proteins in the system, uncovering three novel unreported domains in the enigmatic
wolframin protein. These provide further testable hypotheses on the functions of wolframin in the
context of the Ca^{2+}-stores system and in protecting cells against the response stresses that impinge on
the ER.

88 2   Methods

89 2.1   Sequence analysis

Iterative sequence profile searches were performed using the PSI-BLAST program (RRID:
SCR_001010) (Altschul et al., 1997) against a curated database of 236 eukaryotic proteomes
retrieved from the National Center for Biotechnology Information (NCBI), with search parameters
varying based on the query sequence (see Supplementary Figure S1) and composition. For building a
curated dataset of eukaryotic proteomes, completely sequenced eukaryotic genomes were culled from
Refseq and Genbank, and representative genomes (with a preference for reference genomes) were
chosen from different phyletic groups using the eukaryotic phylogenetic tree as guide. If more than
one genome was available for a species, we typically chose the one listed as a reference genome, or
one that had the best assembly, or one with the most complete proteome set. The list of genomes is
given in the supplementary data of the supplementary material. The program HHpred
(RRID:SCR_010276) (Soding, 2005; Alva et al., 2016) was used for profile-profile comparisons. The
BLASTCLUST program[1] (RRID: SCR_016641) was used to cluster protein sequences based on
BLAST similarity scores. Support for inclusion of a protein in an orthologous cluster involved
reciprocal BLAST searches, conservation of domain architectures, and, when required, construction
of phylogenetic trees with FastTree 2.1.3 (RRID: SCR_015501) (Price et al., 2010) with default
parameters. The trees were visualized using FigTree[2] (RRID: SCR_008515). Selected taxonomic
absences were further investigated with targeted BLASTP (RRID: SCR_001010) (Altschul et al.,
1990) and TBLASTN (RRID: SCR_011822) (Gertz et al., 2006) searches against NCBI's non-
redundant (nr) and nucleotide (nt) databases (Benson et al., 2013), respectively. Multiple sequence
alignments were constructed using the MUSCLE (Edgar, 2004) and GISMO (Neuwald and Altschul,
2016) programs with default parameters. Alignments were manually adjusted using BLAST high-
score pair (hsp) results as guides. Secondary structure predictions were performed with the Jpred 4
program (RRID: SCR_016504) with default settings (Drozdetskiy et al., 2015). EMBOSS (RRID:
SCR_008493) pepwheel[3] was used to generate renderings of amino acid positions on the
circumference of an α-helix.

115 2.2   Protein network construction

116 Protein-protein interactions were extracted from published data sources, updating any outdated
117 gene/protein names and making substantial efforts to disambiguate between paralogs (Supplementary
118 Data). High-throughput/predicted protein-protein interactions were extracted from the FunCoup

---

[1] ftp://ftp.ncbi.nih.gov/blast/documents/blastclust.html

[2] http://tree.bio.ed.ac.uk/software/figtree

[3] http://www.bioinformatics.nl/cgi-bin/emboss/pepwheel

database (Ogris et al., 2018). Networks were visualized using the R-language implementations of the
iGraph and qGraph packages. For network rendering, the Fruchterman-Reingold force-directed
algorithm was used (Fruchterman and Reingold, 1991).

2.3   Comparative genome analyses

Quantitative analysis of phyletic patterns and paralog counts for different proteins/families were first
obtained using a combination of sequence similarity searches as outlined above. We filtered the
counts to exclude multiple identical sequences annotated with the same gene name, using the latest
genome assemblies for each of these organisms as available in NCBI GenBank (RRID:
SCR_002760) as anchors. Proteomes with identifiable quality issues were removed from downstream
analyses (e.g. genomes containing sequences with ambiguous strain assignment), leaving counts for
216 organisms. These counts and their phyletic patterns defined two sets of vectors, namely the
distribution by organism for a given protein and the complement of proteins for a given organism.
These vectors for the protein families and organisms were used to compute the inter-protein or inter-
organism Canberra distances (Lance and Williams, 1966), which is best suited for such integer data
of the form of presences and absences. The Canberra distance between two vectors $\vec{p}$ and $\vec{q}$ is defined
as:

$$d(\vec{p}, \vec{q}) = \sum_{i=1}^{n} \frac{|p_i - q_i|}{|p_i| + |q_i|}$$

These distances were used to cluster the protein families and organisms through agglomerative
hierarchical clustering using Ward's method (Kaufman and Rousseeuw, 1990). The results were
rendered as dendrograms. Ward's method takes the distance between two clusters, A and B to be the
amount by which the sum of squares from the center of the cluster will increase when they are
merged. Ward's method then tries to keep this growth as small as possible. It tends to merge smaller
clusters that are at the same distance from each other as larger ones, a behavior useful in lumping
"stragglers" in terms of both organisms and proteins with correlated phyletic patterns.
142
The protein complement vectors for organisms were also used to perform principal component
analysis to detect spatial clustering upon reducing dimensionality. The variables were scaled to have
unit variance for this analysis. Similarly, a linear discriminant analysis was performed on these
vectors using representatives of the major eukaryotic evolutionary lineages (see Supplementary Table
S1 for list) as the prior groups for classification. This was then used on our complete phyletic pattern
data for classification of the organisms based on their protein complements.
149
150   Organism polydomain scores were calculated as follows: if $c(o, p)$ counts the number of paralogs of
151   some protein domain family p in some organism o, P is the set of all proteins studied, and O is the
152   set of all organisms studied, then the polydomain score for an organism $o \in O$ is defined as:

$$PD(o) = \sum_{p \in P} c(o, p) \cdot (f(p) - \bar{f})$$

153   where $\bar{f}$ is the mean of $f(p)$ for all proteins $p \in P$, and $f(p)$ is defined as:

$$f(p) = \log_2 \left(\frac{\sum_{o \in O} c(o, p)}{\sum_{q \in P} \sum_{o \in O} c(o, q)} \right)$$

Computations and visualizations were performed using the R language.

3   Results

3.1   Protein-protein interaction network for the ER-dependent calcium stores
regulatory/signaling apparatus

Metazoan Ca^{2+} stores have been extensively studied, leading to the identification of several proteins
directly or indirectly involved in calcium transport or signaling and regulation of these processes. In
order to apprehend the global structure of this system, we used published literature and the FunCoup
database to derive a network of protein-protein interactions (PPIs) containing human genes that have
roles in Ca^{2+} stores, centering the network on the three families of ER Ca^{2+} channels (SERCA, IP3R,
RyR) (see Methods). The resulting network totaled 173 protein nodes and 761 interaction edges
(Figure 1A, Supplementary Data). The distribution of the number of connections per node (degree
distribution) in this network displays an inverse (rectangular hyperbolic) relationship (R^2= 0.88;
Figure 1C). This is a notable departure from typical PPI networks which tend to show power-law
degree distributions (Bader and Hogue, 2002; Rodrigues et al., 2011). To better understand this
pattern, we studied its most densely connected subnetworks by searching for cliques, where every
node is connected to every other node in the subnetwork. The largest cliques in this network have ten
nodes. As the degree distribution graph shows an inflection around degree 6, we merged all cliques
of size 6 or greater resulting in a subnetwork of 46 nodes comprising close to 50% of the edges of the
overall network (Figure 1B). This suggests that the inverse relationship of the degree distribution is a
consequence of the presence of a core of several highly connected nodes (6 or more edges), which is
in contrast to other system-specific PPI networks showing a power-law degree distribution, as in the
case of the ubiquitin network (Venancio et al., 2009).

Analysis of the proteins in this highly connected sub-network suggests that the 46 proteins can be
broadly classified into 5 groups: 1) the channels and ATP-dependent pumps which constitute the core
Ca^{2+} transport system for ER stores; 2) EF-hand domain proteins such as calmodulin and sorcin that
bind Ca^{2+} and consequently interact with and regulate the biochemistry of numerous other proteins;
3) proteins involved in folding and stability of other proteins, such as chaperones, redox proteins, and
protein disulfide isomerases; 4) components of the protein phosphorylation response that is
downstream of Ca^{2+} flux into the cytoplasm; and 5) proteins linking the network to other major
functional systems, such as Bcl2, which is involved in the apoptosis response in animals, and beclin-
1, which is involved in autophagy.

3.2   Phyletic distribution of key proteins and implications for the evolution of the ER Ca^{2+}-
stores and -signaling pathway

This densely connected sub-network invites questions about its evolutionary origin, particularly
given that the wet-lab results that inform the connections are predominantly drawn from mammalian
studies (see Methods). To understand better the emergence of this sub-network and the conservation
of its nodes across eukaryotes, we systematically analyzed their phyletic patterns (Figure 2A,
Supplementary Figure S1 and Data). A list of the 34 proteins and protein families studied, as well as
their domain architectures, is in Supplementary Table S2. Further, Ward clustering analysis of the
core components based on their phyletic patterns (see Methods) revealed the presence of 5 distinct
clusters (Figure 2C). These clusters appear to have an evolutionary basis with distinct clusters
accommodating proteins that could be inferred as having been in the LECA (e.g. cluster 1) and those
that emerged in a metazoan-specific expansion of the ER Ca^{2+}-stores system (e.g. clusters 4 and 5).

3.2.1 The core LECA complement of the Ca^{2+}-stores system

At least 15 proteins of the ER Ca^{2+}-stores and signaling system are found across all or most
eukaryotic lineages, suggesting that they were present in the LECA. These proteins include key
components of the dense sub-network, such as 1) the SERCA Ca^{2+} pumps; 2) the Ca^{2+}-binding EF-

hand proteins like calmodulin; 3) chaperones involved in protein folding that are mostly found in the
ER and that sometimes act as either Ca^{2+} binding proteins (e.g. calreticulin, calnexin) or as regulators
of other components of the Ca^{2+}-stores and -signaling system (e.g. ERp57/PDIA3, calstabin, TMX1,
ERdj5/DNAJC10); and 4) core enzymes of the Ca^{2+}-dependent phosphorylation-based signaling
system including the CaMKs and the calcineurin A phosphatase. This set of components is likely to
have comprised the minimal ER-Ca^{2+}-stores and -signaling system in the LECA and suggests that
there were already sub-systems in place to mediate: 1) the dynamic transport of Ca^{2+} ions across the
emerging eukaryotic intracellular membrane system and 2) the transmission of signals affecting a
wide-range of subcellular processes based on the sensing of Ca^{2+} ions. 3) A potential stress response
system that channels Ca-dependent signals to regulation of protein folding (Krebs et al.).
212
Notably, the IP3R channels are absent in lineages that are often considered the basal-most
eukaryotes, namely the parabasalids and diplomonads; however, they are present in some other early-
branching eukaryotes such as kinetoplastids (Cavalier-Smith et al.). Their absence in certain extant
eukaryotes (Figure 2A) suggests that they can be dispensable, or that their role can be taken up by
other channels in eukaryotes that lack them. A comparable phyletic pattern is also seen for certain
other key components of the densely connected sub-network, namely the ERdj5 and calnexin
chaperones.
220

3.2.2 Components with clearly identifiable prokaryotic origins

Deeper sequence-based homology searches revealed that at least three protein families of the ER
Ca^{2+}-stores system have a clearly-identifiable bacterial provenance, namely the P-type ATPase pump
SERCA, sarcalumenin, and calmodulin and related EF-hand proteins. Phylogenetic analyses suggest
that the P-type ATPases SERCA and plasma membrane calcium-transporting ATPase (PMCA) were
both present in the LECA. They are most closely-related to bacterial P-type ATPases (Plattner and
Verkhratsky, 2015), which commonly associate with transporters (e.g. Na$^+$-Ca^{2+} antiporters), ion
exchangers (Na$^+$-H$^+$ exchangers), permeases, and other distinct P-type ATPases in conserved gene-
neighborhoods (Supplementary Figure S2) suggesting a role in maintenance of ionic homeostasis
even in bacteria.
231

The GTPases EHD and sarcalumenin, whose GTPase domains belong to the dynamin family (Leipe
et al., 2002), also show clear bacterial origins based on their phyletic patterns and phylogenetic
affinities. The closest bacterial homologs possess a pair of transmembrane helices C-terminal to the
GTPase domain, suggesting a possible role in membrane dynamics (Figure 3B, Supplementary Data).
Phylogenetic trees show that although they are related to the dynamins, the progenitor of the
eukaryotic sarcalumenin and EHD was acquired independently of the dynamins via a separate
transfer from a proteobacterial lineage early in eukaryotic evolution (Figure 3A) (Leipe et al., 2002).
This gave rise to the EH-domain-containing EHD clade of GTPases, which are involved in regulating
vesicular trafficking and membrane/Golgi reorganization. A further secondary transfer, likely from
the kinetoplastid lineage to the metazoans, gave rise to sarcalumenin proper, which has characterized
roles in the Ca^{2+}-stores system (Figure 3A). This raises the possibility that in other eukaryotes, EHD
performs additional roles in the Ca^{2+}-stores system overlapping with metazoan sarcalumenin.
244

Our analysis revealed that the bacterial calmodulin-like EF-hands show a great diversity of domain-
architectural associations. Versions closest to the eukaryotic ones are found in actinobacteria,
cyanobacteria, proteobacteria, and verrucomicrobiae. These prokaryotic calmodulin-like EF-hand
domains are found fused to a variety of other domains (Figure 3E, Supplementary Data), such as a 7-
transmembrane domain (7TM) (cyanobacteria and to a lesser extent in actinobacteria), the

prokaryotic Tic110-like α-helical domain (cyanobacteria), heme-oxygenases (actinobacteria), the
nitric oxide synthase and NADPH oxidase with ferredoxin and nucleotide-binding domains
(cyanobacteria and δ-proteobacteria), as well as cNMP-binding, thioredoxin, cytochrome, and
sulfatase domains (verrucomicrobiae). Comparable architectural associations with several of these
domains, such as the fusions to the sulfatases and the redox-regulator domains thioredoxin,
cytochromes, glyoxylases, and NADPH oxidases, are also observed in eukaryotes. However, our
analysis showed that these fusions appear to be independently derived in the two superkingdoms.
These diverse associations suggest that even in bacteria the calmodulin-like EF-hands function in the
context of membrane-associated signaling and redox reactions, possibly regulated in a Ca^{2+}-flux-
dependent manner (Zhou et al., 2006; Dominguez et al., 2015). We infer that a version of these was
transferred to the stem eukaryote, probably from the cyanobacteria or the actinobacteria, and had
already triplicated by the LECA, giving rise to the ancestral versions of calmodulin and calcineurin
B, which function as part of the Ca^{2+}-stores system, and the centrins, which were recruited for a
eukaryote-specific role in cell division is association with the centrosome (Dantas et al., 2012).

Among components with indirect regulatory roles in the ER Ca^{2+}-dependent system is the peptidyl-
prolyl cis-trans isomerase (PPIase) calstabin (FKBP1A/B in humans), which is a member of the
FKBP family of PPIase chaperones. Phyletic pattern analysis indicates that calstabin was present in
the LECA (Figure 2A). Eukaryotic PPIases are more similar to bacterial PPIases (Supplementary
Data); hence, the ancestral eukaryotic PPIase was likely acquired from the alphaproteobacterial
mitochondrial progenitor. This is also supported by the evidence from extant pathogenic/symbiotic
bacteria wherein the bacterial FKBP-like PPIases play a role in establishing associations with
eukaryotic hosts (Unal and Steinert, 2014). In the stem eukaryotes, the ancestral PPIase acquired
from the bacterial source underwent a large radiation resulting in diverse PPIases that probably went
hand-in-hand with the eukaryotic expansion of low-complexity proteins with potential substrate
prolines. Thus, eukaryotes acquired a wide range of substrate proteins in several eukaryote-specific
pathways and function in several cellular compartments (Trandinh et al., 1992). Calstabin is one of
the paralogs that arose as part of this radiation and appears to have been dedicated to the ER Ca^{2+}-
stores system.

Beyond the above-mentioned, several other components inferred to be part of the LECA complement
of the ER Ca^{2+}-stores system are likely of bacterial origin. However, they do not have obvious
bacterial orthologs and might have diverged considerably from their bacterial precursors in the stem
eukaryote itself. These include the TRIC-like channels (Silverio and Saier, 2011), the CaMKs, and
the thioredoxin domains found in ERp57, ERdj5, and TMX1.

In contrast to the several components of bacterial provenance, the large eukaryotic assemblage of
calcineurin-like protein phosphatases, which includes the Ca^{2+}-stores regulator calcineurin A, are
specifically related to an archaeal clade to the exclusion of all other members of the superfamily
(Figure 3A, Supplementary Data). Notably, these close relatives are present in several
Asgardarchaea, suggesting the eukaryotes may have directly inherited the ancestral version of these
phosphoesterases from their archaeal progenitor (Zaremba-Niedzwiedzka et al., 2017). Strikingly, the
archaeal calcineurin-like phosphatases occur in a conserved operon (Figure 3A) with genes for two
others proteins, one combining a zinc ribbon fused to a phosphopeptide-recognition FHA domain and
the second with a vWA domain fused to a β-barrel-like domain. This supports a similar role for these
archaeal calcineurin-like phosphatases to their eukaryotic counterparts in transducing a signal
through dephosphorylation of a protein-substrate.

Thus, the core, ancestrally-conserved components of the ER-dependent stores system predominantly
descend from the bacteria, although at least one component was inherited from the archaea. While the
roles for some of these domains in possible bacterial Ca^{2+}-dependent systems are apparent, there is
no evidence that these versions function in a coordinated fashion in any single bacterial species or
clade. Further, it is also clear that there was likely more than one bacterial source for the proteins:
components such as SERCA and calstabin, whose closest relatives are proteobacterial, are likely to
have been acquired from the mitochondrial ancestor, whereas calmodulin is likely to have been
derived from a cyanobacterium or actinobacterium. Thus, the ER-dependent Ca^{2+}-stores network was
assembled in the stem eukaryote from diverse components drawn from different prokaryotic lineages.
This assembly of the system in eukaryotes is strikingly illustrated by the case of the calcineurin
complex. Here, the protein phosphatase component has a clear-cut origin from the archaeal precursor
of the eukaryotes, whereas the Ca^{2+}-binding calcineurin B component descends from a bacterial
source. It was the combination of these proteins with very distinct ancestries that allowed the
emergence of a Ca^{2+}-signaling system.
312

3.2.3 Lineage-specific expansions and gene loss shape the Ca2+ response across eukaryotes

In order to obtain some insights regarding the major developments in the evolution of the eukaryotic
Ca^{2+}-stores system, we systematically assembled protein complement vectors for each of the
organisms in the curated proteome dataset (see Methods). After computing the pairwise Canberra
distance between these vectors, we performed clustering using the Ward algorithm (see Methods).
The resulting clusters recapitulate aspects of eukaryote phylogeny, with animals, fungi, plants,
kinetoplastids and apicomplexans forming distinct monotypic clusters (Figure 2B). These
observations together indicate that most major eukaryotic lineages have likely evolved specific
component around the core Ca^{2+}-stores system inherited from the LECA that distinguish them from
related lineages. We then used principal component and linear discriminant analysis (see Methods)
on these vectors to obtain a global quantitative view of the diversification of the Ca^{2+}-stores system.
Plotting the first two principal components/discriminants reveals discernable spatial separation of the
metazoans from their nearest sister group, the fungi (Figure 2D-E). Further, diverse photosynthetic
eukaryotes tended to be spatially colocalized. These observations suggest that certain accretions to
the core Ca^{2+}-stores system probably occurred alongside unique functional developments such as
motility and multicellularity (metazoa) and autotrophy (photosynthetic lineages). Hence,
investigations on individual lineages are likely to unearth novel, lineage-specific regulatory devices,
which potentially reflect adaptations to their respective environments and provide hints about the
biological contexts in which they were found.
332

We further investigated these potential lineage-specific developments by defining a measure, the
polydomain score (see Methods), which captures the overall amplification of the Ca^{2+}-stores system
of an organism in terms of the contribution of the different constituent protein families of the system
(Figure 2F). While PCA and LDA help identify the overall tendencies among organisms in terms of
the system under consideration, the polydomain score is better-suited for identifying specific unusual
features in terms of individual protein families/domains in particular organisms or clades. This score
allowed us to capture some of the key developments in the reconstructed Ca^{2+}-stores system of
particular organisms and lineages. Notably, this led to the identification some of the distinctions
between Ca^{2+}-regulatory systems that may reflect differential strategies for the incorporation of
calcium into exo/endo-skeletal structures in eukaryotes.
343

Comparison of polydomain scores within metazoa (Figure 2G) showed a striking reduction of the
Ca^{2+}-stores system in arthropods as well as in some molluscs and other marine lineages, consistent

with their distinct grouping in the PCA/LDA plots. This appears to be generally related to the lower
dependence on biomineralized structures in these organisms – arthropods utilize chitin as the primary
exoskeletal material as opposed to calcareous structures, and the molluscan lineages showing this
pattern have lost their calcified shells (e.g. octopi). Similarly, fungi, which use chitin as a central cell
wall component, show lower polydomain scores relative to sister eukaryotic lineages. Conversely, we
noted elevated polydomain scores in molluscs with calcareous shells. Likewise, ciliates, which have
evolved a distinct extension to the ancestral eukaryotic Ca^{2+}-stores system in the form of the alveoli
(Plattner, 2017), also have elevated polydomain scores. Ca^{2+}-based signaling has been shown to be
important for defensive trichocyst exocytosis, ciliary action, and other pathways in ciliates (Plattner,
2015). These observations suggest a linkage between strategies for structural and signaling usage of
calcium and the regulation of Ca^{2+} distribution between intracellular compartments.
357
In qualitative terms, the distinct spatial positioning of different eukaryotic lineages in the PCA/LDA
plots and the difference in their polydomain scores could be explained on the basis of lineage-specific
expansions (LSEs), gene losses, and domain architectural diversity between lineages (Lespinet et al.,
2002). We identified specific gains and losses based on inferred ancestral protein complements
(Supplementary Figure S1). One of the most frequent proteins displaying LSEs is the CaMK:
expansions have occurred independently in animals, plants, oomycetes and alveolates (Figure 3C,
Supplementary Data). Ciliates in particular show a dramatic expansion of the family with over one
hundred copies in *Paramecium* and *Stentor*. Other proteins that show LSEs in specific lineages
(Supplementary Data) include calmodulin (in metazoans, certain fungal lineages, and plants),
calstabin (independently in different stramenopile lineages and haptophytes), PP2R3-like proteins (in
kinetoplastids and *Trichomonas*), calcineurin A (in ciliates and *Entamoeba*), calnexins (in
Trichomonas and diatoms), ORAI (in *Emiliania*, Figure 3D), and SERCA (in certain fungi). The
metazoans also display LSEs for several protein families that appear to have specifically emerged in
metazoa (see below).
372
Some of these LSEs have clear biological correlates that were also suggested by the polydomain
scores. For example, LSEs of the calmodulin family in the shelled molluscs (20-46 copies) might
correspond to or be involved in the regulation of Ca^{2+} concentration during the biomineralization
processes involved in the formation of calcareous shells. Concomitant with this expansion, in shelled
molluscs several transporters, including P-type ATPases, voltage-gated channels, and ion exchangers,
have been adapted for the transportation of Ca^{2+} ions for shell formation (Sillanpaa et al., 2018).
Similarly, the development of the distinct set of Ca^{2+} stores in the form of the alveoli, which play
several key ciliate-specific roles (Plattner, 2015; 2017), might explain the expansions in these
organisms. Thus, the very large LSEs of the CaMK and calcineurin A-like phosphoesterase likely
reflect the many diverse pathways into which Ca^{2+}-based signaling is incorporated in these
organisms. Additionally, experimental evidence shows that some pan-eukaryotic members of the ER
Ca^{2+} stores system have acquired additional roles in ciliates; for example, SERCA-like Ca^{2+}-pumps
are also in the membranes of *Paramecium* alveolar calcium stores (Plattner et al., 1999).
386
ORAI family LSEs in *Emiliania* might correspond to the need to regulate Ca^{2+} transport for
mineralizing the calcareous coccoliths (Yin et al., 2018). Further, calcium for coccolith formation
may also supplied in part by an ER-membrane-localized Ca^{2+}/H$^+$ exchanger, as well as by increased
activity of SERCA and a plasma membrane Ca^{2+}/H$^+$ exchanger (Mackinder et al., 2011). Here, as in
ciliates, these expansions might have accompanied the emergence of a distinct Ca^{2+}-stores system
associated with the nuclear envelope (an extension of the ER), which is close in proximity to the
envelope of the coccolith and the site of biomineralization (Brownlee et al., 2015).
394

In contrast, we observed extensive gene losses of proteins including ERdj5, sarcalumenin, IP3R, and
ORAI across several or all fungi and losses of at least 10 conserved proteins in *Entamoeba* (Figure
2A). However, experimental studies have shown that both fungi and *Entamoeba* have Ca^{2+} transients
and associated signaling systems (Makioka et al., 2002; Kim et al., 2012); hence, despite these losses,
the evidence favors these organisms retaining at least a limited Ca^{2+}-store-dependent signaling
network. The CREC (calumenin, reticulocalbin, and Cab45) family displays an unusual phyletic
pattern of particular note, being found only in metazoans and land plants (Figure 2A). Barring the
unusual possibility of lateral transfer between these two lineages, this would imply extensive loss of
these proteins across most other major eukaryotic lineages. Within metazoa, several proteins display
unusual loss patterns, including sorcin, calsequestrin, and SelN (Figure 2A).

Notably, the land plants and their sister group the green algae have entirely lost ancient core Ca^{2+}-
dependent signaling proteins such as calcineurin A and HOMER. The IP3R channels are retained by
the chlorophytes but lost entirely by the land plants, while the ORAI channels are present in the basal
land plants *Physcomitrella* and *Selaginella* but are not observed in crown land plants (Edel et al.,
2017). Land plants have also lost the TRIC channels, but the basal streptophyte *Chara braunii* retains
a copy. The differential retention and expansion of various ancestral Ca^{2+} components (Figure 2A) in
the land plants might be seen as adaptations to a sessile lifestyle no longer requiring Ca^{2+}-signaling
components associated with active cell or organismal motility. However, the paradoxical LSEs of the
calmodulin-like and EF-hand-fused CaMK-like proteins (CDPK family) (Edel et al., 2017) in land
plants relative to their chlorophyte sister-group (Figure 2A, 3C). These might have emerged as part
of Ca^{2+} sequestration and signaling mechanisms relevant in the Ca^{2+}-poor freshwater ecosystems,
wherein the land plants originated from algal progenitors (Delwiche and Cooper, 2015).

Little in the way of domain architectural diversity is observed in the core conserved proteins of the
animal Ca^{2+}-stores system. A notable exception is seen in the CaMKs: metazoan CaMKs show a
higher architectural complexity via their fusion to several distinct globular domains. These domains
act as adaptors in interactions with Ca^{2+}-binding regulatory domains, effectively broadening the total
range of signaling pathways in which the CaMKs participate in metazoa (Wang et al., 2015). In
contrast, land plants display lower architectural complexity with CaMK orthologs directly fused to
Ca^{2+}-binding domains (Klimecka and Muszynska, 2007).

3.2.4 Accretion of novel Ca2+ signaling pathway components in the Metazoa

Case by case examination revealed that the distinct position of the metazoans in the PCA/LDA plots
relative to other eukaryotes (Figure 2D-E) is due to a major accretion of novel signaling and flux-
related Ca^{2+}-stores components at the base of the metazoan lineage (Figure 2A). Our analyses suggest
three distinct origins of these proteins: 1) several emerged as paralogs of domains already present in
the LECA and functioning in the context of Ca^{2+}-signaling, including the EF-hand domains found in
the STIM1/2, calumenin, SelN, S100, sorcin, and NCS1 proteins, the thioredoxin domain found in
the ERp44 and calsequestrin proteins, and the RyR channel proteins. 2) Proteins or component
domains which are involved in a broader range of functions but have been recruited to roles in Ca^{2+}-
stores regulation specifically in the animal lineages, such as the SAM domain in the STIM1/2
proteins, the PDZ domain in the neurabin protein, and the TRPC channels. 3) Proteins containing
domains which appear to be either novel metazoan innovations, such as the KRAP domain of the
Tespa1 protein, or whose origins have yet to be traced, such as VGCC and wolframin. Further, the
origin of certain metazoan-specific proteins such as phospholamban, which inhibits the SERCA
ATPase, remain difficult to trace because of their small size and highly-biased composition as
membrane proteins. Inspection of the neighbors of these proteins in the constructed network suggests

that in almost all instances, these proteins were added to the Ca^{2+} signaling network via interactions
with one or more of the ancient components of the system (Figure 1A).
445
This sudden accretion of Ca^{2+}-stores components likely coincided with emergence of well-studied
aspects of differentiated metazoan tissues, such as muscle contraction and neurotransmitter release
(Zucchi and Ronca-Testoni, 1997; Clapham, 2007). Other additions to the network likely arose via
interface with other pathways like apoptosis and autophagy in the context of ER stress response
(Smaili et al., 2013). Notable in this regard is the metazoan-specific Bcl2 family of membrane-
associated proteins associated with regulation of apoptosis. They appear to have been derived via
rapid divergence in metazoans from pore-forming toxin domains of ultimately bacterial provenance
(Peng et al., 2009), which are found in pathogenic bacteria and fungi (Aravind et al., 2012).
454
Still other proteins were recruited to the system as part of newly-emergent regulatory subnetworks,
such as the Store-Operated Calcium Entry (SOCE) pathway for re-filling the ER from extracellular
Ca^{2+} stores (Prakriya and Lewis, 2015; Ong et al., 2016). Of the proteins identified in this pathway,
the ORAI Ca^{2+} channels are present in the early-branching kinetoplastid lineage but were lost in
several later-branching lineages (Figure 2A), while other components like the ORAI-regulating
STIM1/2 proteins and the TRPC channels emerged around the metazoan accretion event, suggesting
the core SOCE pathway came together at or near the base of the animal lineage. In course of this
analysis, we identified a common origin for the ORAI and Jiraiya/TMEM221 ER channels, the latter
of which are characterized in BMP signaling (Aramaki et al., 2010). The Jiraiya channels are
observed in animals but are absent in earlier-branching eukaryotic lineages, suggesting they emerged
from a duplication of an ORAI channel early in metazoan evolution (Figure 3D, Supplementary
Data). Jiraiya channel domains lack the Ca^{2+} binding residues seen in ORAI channels, suggesting
they are unlikely to directly bind Ca^{2+}. However, it is possible that Jiraiya/TMEM221 physically
associates with components of the Ca^{2+}-stores system to regulate them.

3.3   Domain architectural anatomy and functional analysis of the wolframin protein

As noted above, one of the uniquely metazoan proteins in the Ca^{2+}-stores system is wolframin, a
transmembrane protein localized to the ER membrane (Hildebrand et al., 2008; Rigoli et al., 2011;
Qian et al., 2015) (Figure 2A, Supplementary Data). Wolframin, along with the structurally-unrelated
and more widely phyletically distributed (Figure 2A) Wolfram syndrome 2 (WFS2) protein, is
implicated in Wolfram syndrome (Inoue et al., 1998; Strom et al., 1998; Amr et al., 2007; Urano,
2016). Experimental studies have attributed biological roles to unannotated regions upstream and
downstream of the TM region in wolframin, and our analyses revealed four uncharacterized globular
regions therein (Figure 4A). The cytosolic N-terminal region was identified through iterative
database searches (see Methods) as containing Sel1-like repeats (SLRs; query: NP_005996, hit:
OYV16035, iteration 2, e-value: 5×10^{-16}), which are α-helical superstructure forming repeats
structurally comparable to the tetratricopeptide repeats (TPRs) (Ponting et al., 1999; Karpenahalli et
al., 2007). Profile-profile searches (see Methods) unified the remaining two globular regions
respectively with various EF-hand domains (e.g., query: XP_011608878, hit: 1SNL_A, p-value
2.3×10^{-4}) and OB fold-containing domains (e.g. query: XP_017330582, hit: 2FXQ_A, p-value:
1.3×10^{-4}).
485
The wolframin SLR possess several conserved residues seen in the classical SLRs (Figure 4C,
Supplementary Table S4)(Mittl and Schneider-Brachert, 2007), and additionally display a truncated
first loop when compared to known SLRs (Figure 4C). The SLRs of wolframin appear most closely-
related to bacterial versions, suggesting a possible horizontal transfer at the base of animals (Ponting

et al., 1999); however, the short length and rapid divergence of such repeats complicates definitive
ascertainment of such evolutionary relationships. SLRs and related α/α repeats are often involved in
coordinating interactions within protein complexes (Karpenahalli et al., 2007; Mittl and Schneider-
Brachert, 2007) and have been characterized specifically in ER stress and misfolded protein
degradation responses (Jeong et al., 2016) and in mediating interactions of membrane-associated
protein complexes (Mittl and Schneider-Brachert, 2007). As the wolframin SLRs roughly correspond
to an experimentally-determined calmodulin-binding region (Yurimoto et al., 2009), they might
specifically mediate that protein interaction.

The EF-hand region of wolframin contains the two characteristic copies of the bihelical repeat that
form the basic EF-hand unit. However, it lacks the well-characterized Ca^{2+}-binding DxDxDG motif
or any comparable residue conservation (Figure 4D) (Gifford et al., 2007; Denessiouk et al., 2014). It
also features striking loop length variability in between the two helix-loop-helix motifs (Figure 4D).
Such "inactive" EF-hand units typically dimerize with other EF-hand proteins (Kawasaki et al., 1998;
Gifford et al., 2007), suggesting that wolframin could be self-dimerizing or that its EF-hand could
interact at the ER membrane with calmodulin and/or a distinct EF-hand protein.

The C-terminal OB fold domain (Figure 4B,E) lacks the conserved polar residues typical of nucleic
acid-binding OB-fold domains (Watson et al., 2007; Guardino et al., 2009). Further, the localization
of this region to the ER lumen suggests that it is unlikely to be involved in nucleic acid binding
(Arcus, 2002; Flynn and Zou, 2010; Krishna et al., 2010). Alternatively, this OB-fold domain could
mediate protein-protein interactions as has been observed for other members of the fold (Flynn and
Zou, 2010) and is also consistent with previous experimental studies implicating this region of
wolframin in binding the pre-folded form of ATP1B1 (Zatyka et al., 2008). Strikingly, in the region
N-terminal to the OB fold domain we observe a further globular region containing a set of six
absolutely-conserved cysteines (Figure 4E). This region is not unifiable with any known domains and
could conceivably represent an extension to the core OB fold domain. These conserved cysteine
residues could contribute to disulfide-bond-mediated cross-linking, a well-studied regulatory
mechanism of Ca^{2+}-stores regulation (Ushioda et al., 2016). Additionally, C-terminal to the OB-fold,
wolframin has a hydrophobic helix that might be involved in intra- or inter-molecular interactions
(Figure 4E).

The wolframin TM region, located in the central region of the protein (Figure 4A), consists of nine
transmembrane helices (Hofmann et al., 2003; Rigoli et al., 2011; Qian et al., 2015). Despite
extensive studies on the TM region, its precise role in affecting Ca^{2+} flow across the ER membrane
remains the subject of some debate (Osman et al., 2003; Aloi et al., 2012; Zatyka et al., 2015;
Cagalinec et al., 2016). Inspection of a multiple sequence alignment of the wolframin transmembrane
region (Supplementary Figure S3) revealed a concentration of polar residues which are spatially
alignable in helices 4 and 5 (Supplementary Figure S4). This is reminiscent of membrane associated
polar residue configurations seen in proteins that allow transmembrane flux of ions. Hence, it would
be of interest to investigate if these residues might play a role in ion transport by wolframin.

4   Discussion

4.1   Evolutionary and functional considerations

4.1.1 Early and later landmarks in the evolution of the ER Ca^{2+}-stores system

The ER Ca^{2+}-stores system displays several parallels in its evolutionary history to other
endomembrane-dependent systems such as the nuclear membrane and vesicular trafficking systems

(Mans et al., 2004; Jekely, 2008). Like in the case of these systems, dedicated ER Ca^{2+}-stores
systems are absent in the prokaryotes, despite the presence of Ca^{2+} transients and Ca^{2+}-dependent
signaling pathways in them (Dominguez et al., 2015). However, several of the more ancient
individual components of eukaryotic Ca^{2+}-stores systems are of clear-cut prokaryotic origin. Notably,
we show here that not all of these proteins originated from the proto-mitochondrion; notably,
calmodulin is of likely cyanobacterial or actinobacterial provenance and the calcineurin-like
phosphoesterases originally descended from the archaea. The former observation adds to
accumulating evidence of LECA acquiring bacterial contributions from non-α-proteobacterial
lineages. It is therefore possible that LECA had a more extensive set of associated symbionts than
what was fixed as the mitochondrion in eukaryotic evolution (Huang and Gogarten; Burroughs et al.,
2017; Verma et al., 2018).

The cyanobacterial/actinobacterial origin of calmodulin and the role for the closely-related and early-
branching centrin family of EF-hand proteins in microtubule dynamics during cell division suggest
that the LECA already had a strong Ca^{2+} dependency. This raises the possibility that the prokaryotic
ancestors of the eukaryotes might have existed in a calcium-rich environment such as the
biomineralized structures (e.g. stromatolites) formed by cyanobacteria (Bosak et al., 2013). This is
consistent with the diversity of domain architectures for calmodulin-like proteins with ramifications
into various functional systems in the cyanobacteria that we reported here. This diversified pool of
Ca^{2+}-binding domains could have contributed raw materials needed during the initial emergence of
Ca^{2+} flux-based signaling and regulation across endomembranes in eukaryotes. The newly emerged
intracellular Ca^{2+} gradients were likely fixed by the myriad advantages it bestowed in the stem
eukaryotes, including increased signaling capacity and a regulatory mechanism for processes like
growth/proliferation, secretion, and motility.

Figure 5 shows a summary of key acquisitions and losses of Ca^{2+}-stores proteins during eukaryotic
evolution, placed onto a simplified model of eukaryotic evolution. In inferring ancestral states, and
thereby gains and losses, we followed the (relatively) certain contours of the eukaryotic tree (see also
Supplementary Figure S1). For example, we assumed that the root of eukaryotes lies in the excavates
and that the SAR (stramenopile/alveolate/rhizarian) group is monophyletic. In general, we strove to
make the least number of assumptions possible; therefore, many of our estimates – for example of the
number of Ca^{2+}-stores proteins in the LECA – are lower bounds and conservative. By these criteria,
we infer that the ancestral eukaryotic ER-dependent Ca^{2+}-stores system likely consisted of a
combination of the SERCA pump, a cation channel, EF-hand-containing proteins, and
phosphorylation enzymes (Figure 5). Early in eukaryotic evolution, chaperone domains associated
with protein folding and thioredoxin fold domains were added to the system, likely recruited from
roles as general regulatory domains, some of which could also bind Ca^{2+} (Figure 5). Association with
thioredoxin fold domains, involved in disulfide-bond isomerization, is of note as this points to early
emergence of a link between redox-dependent folding of cysteine-rich proteins and Ca^{2+}
concentrations. The striking presence of cysteine-rich domains associated with cyanobacterial
calmodulin homologs (Figure 3E; DES, LMI, and LA unpublished observations) suggests that such a
connection might have emerged even before the origin of the eukaryotic Ca^{2+}-stores system. These
functional links might have persisted until later in eukaryotic evolution as hinted by the cysteine-rich
domain present in wolframin. This led to the basic system as reconstructed in the LECA (Figure 2A).

Waves of additional accretion events added components to the Ca^{2+}-stores system at distinct points in
eukaryotic evolution, often appearing to correlate with adaptations to distinct lifestyles, such as the
evolution of motile multicellular forms, loss of motility (the crown plant lineage), or the evolution of

calcium-rich biomineralized skeletons and shells (Figures 2A, 5, Supplementary Figure S1).
Strikingly, we appear to observe some correlation between the loss-and-gain patterns of the
regulatory components of Ca^{2+}-stores systems and the degree of structural utilization of calcium. For
example, a relative dearth of these components is observed in the arthropod and fungal lineage,
which use chitin-based structural components as opposed to the calcium-based biomineralized
skeletons and shells of vertebrates and certain molluscs (Figure 2F-G). "Because the data in this
study necessarily relies on experimental findings primarily from animals, it is best suited to
characterizing the system from the viewpoint of metazoa. However, variations in presence/absence
across lineages and LSEs provide insight into the dynamic evolution of Ca^{2+} stores regulation and
suggests there are further complexities to explore in more poorly-characterized eukaryotes/

4.1.2 Wolframin domain architecture and interactors

Even within animal Ca^{2+}-stores systems, several proteins with important regulatory roles remain
poorly-understood in terms of their functional mechanisms. Wolframin is such a protein whose
domain composition has eluded researchers for over two decades (Inoue et al., 1998; Strom et al.,
1998; Osman et al., 2003). Assignment of domains at the N- and C-termini of the central TM region
(Figure 4A), as well as the positioning of wolframin in the assembled interaction network, (Figure
1A) (Zatyka et al., 2015) supports a role in coordinating interactions on both sides of the ER
membrane. These are likely to take the form of PPIs with Ca^{2+}-binding proteins or through disulfide
bond interactions (see above).
603
However, outside of a possible bacterial origin for the SLR region (see above, Figure 4C), the precise
evolutionary origins of the remaining domains comprising wolframin remain mostly unclear. It
appears likely these domains were derived from paralogs of existing EF-hand and OB fold domains,
both of which had already undergone extensive domain radiations in the eukaryotes prior to the
emergence of the animals, and then assembled and recruited to a Ca^{2+}-stores regulatory role at the ER
(Lespinet et al., 2002). Their rapid divergence, evident by the lack of recognizable relationships to
known families with their respective folds, could have resulted from the extraordinary selective
pressures occurring with the major burst of evolutionary changes during the emergence of the
metazoan lineage.
613
Despite observations that genic disruptions contribute to similar though not identical phenotypes
(Urano, 2016), WFS1 and WFS2 are structurally unrelated. Our interaction network analysis further
failed to uncover any shared interactors (Figure 1A), although both have been linked in the past to
calpain activity (Lu et al., 2014), mitochondrial dysfunction (Chang et al., 2012; Cagalinec et al.,
2016), and apoptosis (Yamada et al., 2006; Chang et al., 2010). Recent research on WFS2 has
particularly focused on its role in calcium stores regulation at the intersection of the ER and
mitochondrial membranes (Rouzier et al., 2017). We believe that our identification of the constituent
domains of wolframin reported herein might help clarify its function better through target deletion
and mutagenesis of these domains.

4.2   Conclusions

Reconstruction of the evolution of the eukaryotic Ca^{2+}-stores regulatory system points to a core of
domains inherited from distinct prokaryotic sources conserved across most eukaryotes. Lineage-
specific differentiation of the system across eukaryotes is driven by complexities stemming from
both the loss and/or expansion of the core complement of domains by the addition of components via
LSEs or recruitment of domains of diverse provenance. We analyze in depth one such striking
example of the latter, namely wolframin, which has been previously implicated in human disease.

The evolution of wolframin provides a model for how regulatory components of the Ca^{2+}-stores
system emerged: through the combining of existing mediators of Ca^{2+} signaling, like the EF-hand
domains, with other domains originally not found in the system. Such transitions often happened at
the base of lineages that subsequently underwent substantial diversification. We hope the findings
presented here open novel avenues for the ongoing research on the regulation of calcium stores
across eukaryotes, including providing new handles for understanding the functional mechanisms of
wolframin and its dysregulation in Wolfram syndrome.

5   Conflict of Interest

638   The authors declare that the research was conducted in the absence of any commercial or financial
639   relationships that could be construed as a potential conflict of interest.

6   Author Contributions

641   Conceptualization, DS, AMB, LA; formal analysis, DS, AMB, LMI; analytical tools, DS, LA;
642   project administration, AMB, LMI, LA; visualization, DS; writing—original draft DS; AMB.;
643   writing—review and editing, AMB, LMI, and LA

7   Funding

645   DES, LMI., AMB., and LA are supported by the Intramural Research Program of the NIH, National
646   Library of Medicine.

8   Acknowledgments

648   This is a short text to acknowledge the contributions of specific colleagues, institutions, or agencies
649   that aided the efforts of the authors.

9   Data Availability Statement

651   All datasets analyzed for this study are included in the manuscript and the supplementary files.

10   Figure legends

Figure 1. Protein-protein interaction network of the ER-dependent Ca^{2+}-stores and -signaling system.
(**A**) The network. Edges are classified as follows: interactions reported in the literature in which each
partner gene either has no other paralog or has been specifically identified (solid black), reported
interactions where the specific paralog for at least one of the two genes is unclear (dashed gray), and
interactions found in FunCoup 4.0 with P ≥ 0.9 (light pink). Medium- and large-sized nodes represent
the proteins whose phyletic distribution was selected for detailed analysis and are colored based on
phylogeny as follows: pan-eukaryotic, green; metazoans and close relatives, yellow; metazoan-
specific, orange; chordate-specific, red. Nodes labeled by HUGO nomenclature to capture paralog-
specific interactions. The borders of nodes involved in selected intracellular systems are colored
based on the key given in the lower left. (**B**) The highly-connected subnetwork. Edge coloring has no
significance; nodes are colored and sized as in (A). (**C**) A histogram of the degrees of the nodes of
the network shown in (A). A curve of best fit is shown in gray.

Figure 2. Comparative genome analyses. (**A**) A visualization of the number of paralogs of the 34 key
proteins/protein families (vertical axis) across 216 eukaryotes (horizontal axis). Each circle

represents the number of paralogs of one protein/family in one organism; the radius of each circle is
scaled by the hyperbolic arcsine of the represented number of paralogs. The proteins and protein
families shown can be broadly divided into two categories: those that are largely pan-eukaryotic
(calmodulin to ORAI) and those that originate in the metazoans or their close relatives (STIM to
S100). Eukaryotic clades are color-coded and labeled below the plot. Where needed, protein/protein
family names are supplemented in parenthesis with human gene names from Figure 1, where node
labels distinguish between paralogs. (**B**) A dendrogram of the 216 eukaryotes based on the counts of
their paralogs of the 34 proteins/families. Tip labels are colored by clade using the same system as in
panel A. (**C**) A dendrogram of the 34 proteins/families based on the counts of their paralogs. Five
major clusters are numbered to their left. Branch coloring corresponds to major clusters. (**D-E**) Plots
of the first two (**D**) principal components and (**E**) discriminants of each of the 216 eukaryotes
resulting from principal component and linear discriminant analyses on the paralog count dataset.
Color keys linking colors to high-level clades are given in the upper right and lower right,
respectively, of the plots. (**F-G**) Histograms of the polydomain scores of (**F**) all 216 eukaryotes,
shown on a base-10 log scale, and (**G**) 50 metazoans, shown on a linear scale. Contributions of some
clades to each bar of the histograms are shown through coloring given by the keys in the upper right
of the histograms.

Figure 3. (**A-D**) Stylized phylogenetic trees showing (**A**) sarcalumenin, EHD, and related bacterial
dynamins; (**B**) a partial tree of the calcineurin-like superfamily, containing the classical calcineurin A
phosphatases with their immediate eukaryotic relatives and newly-recognized archaeal orthologs,
along with the related MRE11/rad32/sbcD-like phosphoesterases as an outgroup; (**C**) the expansions
of the calcium/calmodulin-dependent kinase in eukaryotes; and (**D**) ORAI and Jiraiya. Collapsed
groups are colored as follows: universal distribution, yellow; bacterial, blue; archaeal, red; pan-
eukaryotic, dark green; metazoan, light green; restricted to non-metazoan eukaryotes, blue-green.
Branches with bootstrap support of greater than 85% are marked with a black circle. Genome
contextual associations pertinent to a given clade are provided within context of trees. Conserved
gene neighborhoods are depicted as boxed arrows and protein domain architectures as boxes linked
in the same polypeptide. The trees are also provided in Newick format in the Supplementary Data.
(**E**) Domain architectures of proteins containing calmodulin-like EF-hands. Each EF-hand is a dyad
of EF repeats. Long (>200 residue) regions without an annotated domain are collapsed using "//".

Figure 4. Structural and sequence overview of wolframin. (A) Domain architecture of wolframin. (B)
Topology diagram of the wolframin OB-fold. The labeled secondary structure elements correspond
with the labeled secondary structure elements and coloring in the OB-fold alignment in part E. The β-
strand shaded in gray is a possible sixth strand stacking with the core OB-fold domain barrel. (C-E)
Multiple sequence alignments of the (C) Sel1-like repeats, (**D**) EF-hand, (**E**) and cysteine-rich and
OB-fold domains of wolframin. Sequences are labeled to the left of the alignments by organism
abbreviation (see Supplementary Table S3) and NCBI GenBank accession number. Secondary
structure is provided above the alignments; green arrows represent strands and red cylinders represent
helices. The two EF motifs are marked with blue arrows above the secondary structure line. The
boundaries of the core OB-fold and the cysteine-rich region (E) are marked with pentagonal arrows
above the secondary structure line and pointing towards the center of the domain. The six secondary
structure elements that comprise the OB-fold are labeled with blue text in the secondary structure
line. Conserved cysteines are marked with an asterisk. A 90% consensus line is provided below the
alignments; the coloring and abbreviations used are: h (hydrophobic), l (aliphatic), and a (aromatic)
are shown on a yellow background; o (alcohol) is shown in salmon font; p (polar) is shown in blue
font; + (positively charged), - (negatively charged), and c (charged) are shown in pink font; s (small)
is shown in green font; u (tiny) is shown on a green background; b (big) is shaded gray.

Figure 5. Summary of the evolution of the Ca^{2+} stores system. The temporal timing of the emergence
of the core components of the system are overlaid on a consensus phylogenetic tree depicting
eukaryotic evolution. Colors indicate component provenance, as labeled in the key provided in upper-
left corner. Components with identified losses or expansions within a particular lineage are listed to
the right, with '-' and '*' labels denoting loss and expansion of a particular component, respectively.
A detailed breakdown of losses/expansions within the labeled lineages are provided in Figure 2 and
Supplementary Figure S1.

11   References

Aloi, C., Salina, A., Pasquali, L., Lugani, F., Perri, K., Russo, C., et al. (2012). Wolfram syndrome:
723   new mutations, different phenotype. *PLoS One* 7(1)**,** e29150. doi:
724   10.1371/journal.pone.0029150.
725   Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990). Basic local alignment
726   search tool. *J Mol Biol* 215(3)**,** 403-410. doi: 10.1016/s0022-2836(05)80360-2.
727   Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W., et al. (1997). Gapped
728   BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic
729   Acids Res* 25(17)**,** 3389-3402.
730   Altshuler, I., Vaillant, J.J., Xu, S., and Cristescu, M.E. (2012). The evolutionary history of
731   sarco(endo)plasmic calcium ATPase (SERCA). *PLoS One* 7(12)**,** e52617. doi:
732   10.1371/journal.pone.0052617.
733   Alva, V., Nam, S.Z., Soding, J., and Lupas, A.N. (2016). The MPI bioinformatics Toolkit as an
734   integrative platform for advanced protein sequence and structure analysis. *Nucleic Acids Res*
735   44(W1)**,** W410-415. doi: 10.1093/nar/gkw348.
736   Alzayady, K.J., Sebe-Pedros, A., Chandrasekhar, R., Wang, L., Ruiz-Trillo, I., and Yule, D.I. (2015).
737   Tracing the Evolutionary History of Inositol, 1, 4, 5-Trisphosphate Receptor: Insights from
738   Analyses of Capsaspora owczarzaki Ca2+ Release Channel Orthologs. *Mol Biol Evol* 32(9)**,**
739   2236-2253. doi: 10.1093/molbev/msv098.
740   Amr, S., Heisey, C., Zhang, M., Xia, X.J., Shows, K.H., Ajlouni, K., et al. (2007). A homozygous
741   mutation in a novel zinc-finger protein, ERIS, is responsible for Wolfram syndrome 2. *Am J
742   Hum Genet* 81(4)**,** 673-683. doi: 10.1086/520961.
743   Aramaki, T., Sasai, N., Yakura, R., and Sasai, Y. (2010). Jiraiya attenuates BMP signaling by
744   interfering with type II BMP receptors in neuroectodermal patterning. *Dev Cell* 19(4)**,** 547-
745   561. doi: 10.1016/j.devcel.2010.09.001.
746   Aravind, L., Anantharaman, V., Zhang, D., de Souza, R.F., and Iyer, L.M. (2012). Gene flow and
747   biological conflict systems in the origin and evolution of eukaryotes. *Front Cell Infect
748   Microbiol* 2**,** 89. doi: 10.3389/fcimb.2012.00089.
749   Arcus, V. (2002). OB-fold domains: a snapshot of the evolution of sequence, structure and function.
750   *Curr Opin Struct Biol* 12(6)**,** 794-801.
751   Ashby, M.C., and Tepikin, A.V. (2001). ER calcium and the functions of intracellular organelles.
752   *Semin Cell Dev Biol* 12(1)**,** 11-17. doi: 10.1006/scdb.2000.0212.
753   Bader, G.D., and Hogue, C.W. (2002). Analyzing yeast protein-protein interaction data obtained
754   from different sources. *Nat Biotechnol* 20(10)**,** 991-997. doi: 10.1038/nbt1002-991.
755   Banerjee, S., Vishwanath, P., Cui, J., Kelleher, D.J., Gilmore, R., Robbins, P.W., et al. (2007). The
756   evolution of N-glycan-dependent endoplasmic reticulum quality control factors for
757   glycoprotein folding and degradation. *Proc Natl Acad Sci U S A* 104(28)**,** 11676-11681. doi:
758   10.1073/pnas.0704862104.
759   Benson, D.A., Cavanaugh, M., Clark, K., Karsch-Mizrachi, I., Lipman, D.J., Ostell, J., et al. (2013).
760   GenBank. *Nucleic Acids Res* 41(Database issue)**,** D36-42. doi: 10.1093/nar/gks1195.

Berridge, M.J. (2012). Calcium signalling remodelling and disease. *Biochem Soc Trans* 40(2)**,** 297-
309. doi: 10.1042/BST20110766.
Bosak, T., Knoll, A.H., and Petroff, A.P. (2013). The Meaning of Stromatolites. *Annual Review of*
Earth and Planetary Sciences 41(1)**,** 21-44. doi: 10.1146/annurev-earth-042711-105327.
Brownlee, C., Wheeler, G.L., and Taylor, A.R. (2015). Coccolithophore biomineralization: New
questions, new answers. *Semin Cell Dev Biol* 46**,** 11-16. doi: 10.1016/j.semcdb.2015.10.027.
Burroughs, A.M., Kaur, G., Zhang, D., and Aravind, L. (2017). Novel clades of the HU/IHF
superfamily point to unexpected roles in the eukaryotic centrosome, chromosome
partitioning, and biologic conflicts. *Cell Cycle* 16(11)**,** 1093-1103. doi:
10.1080/15384101.2017.1315494.
Cagalinec, M., Liiv, M., Hodurova, Z., Hickey, M.A., Vaarmann, A., Mandel, M., et al. (2016). Role
of Mitochondrial Dynamics in Neuronal Development: Mechanism for Wolfram Syndrome.
PLoS Biol 14(7)**,** e1002511. doi: 10.1371/journal.pbio.1002511.
Cai, X., Wang, X., Patel, S., and Clapham, D.E. (2015). Insights into the early evolution of animal
calcium signaling machinery: a unicellular point of view. *Cell Calcium* 57(3)**,** 166-173. doi:
10.1016/j.ceca.2014.11.007.
Cavalier-Smith, T., Chao, E.E., and Lewis, R. (2015). Multiple origins of Heliozoa from flagellate
ancestors: New cryptist subphylum Corbihelia, superclass Corbistoma, and monophyly of
Haptista, Cryptista, Hacrobia and Chromista. *Mol Phylogenet Evol* 93**,** 331-362. doi:
10.1016/j.ympev.2015.07.004.
Chang, N.C., Nguyen, M., Germain, M., and Shore, G.C. (2010). Antagonism of Beclin 1-dependent
autophagy by BCL-2 at the endoplasmic reticulum requires NAF-1. *Embo j* 29(3)**,** 606-618.
doi: 10.1038/emboj.2009.369.
Chang, N.C., Nguyen, M., and Shore, G.C. (2012). BCL2-CISD2: An ER complex at the nexus of
autophagy and calcium homeostasis? *Autophagy* 8(5)**,** 856-857. doi: 10.4161/auto.20054.
Clapham, D.E. (2007). Calcium signaling. *Cell* 131(6)**,** 1047-1058. doi: 10.1016/j.cell.2007.11.028.
Dantas, T.J., Daly, O.M., and Morrison, C.G. (2012). Such small hands: the roles of
centrins/caltractins in the centriole and in genome maintenance. *Cell Mol Life Sci* 69(18)**,**
2979-2997. doi: 10.1007/s00018-012-0961-1.
Delwiche, C.F., and Cooper, E.D. (2015). The Evolutionary Origin of a Terrestrial Flora. *Curr Biol*
25(19)**,** R899-910. doi: 10.1016/j.cub.2015.08.029.
Denessiouk, K., Permyakov, S., Denesyuk, A., Permyakov, E., and Johnson, M.S. (2014). Two
structural motifs within canonical EF-hand calcium-binding domains identify five different
classes of calcium buffers and sensors. *PLoS One* 9(10)**,** e109287. doi:
10.1371/journal.pone.0109287.
Dominguez, D.C., Guragain, M., and Patrauchan, M. (2015). Calcium binding proteins and calcium
signaling in prokaryotes. *Cell Calcium* 57(3)**,** 151-165. doi: 10.1016/j.ceca.2014.12.006.
Drozdetskiy, A., Cole, C., Procter, J., and Barton, G.J. (2015). JPred4: a protein secondary structure
prediction server. *Nucleic Acids Res* 43(W1)**,** W389-394. doi: 10.1093/nar/gkv332.
Edel, K.H., Marchadier, E., Brownlee, C., Kudla, J., and Hetherington, A.M. (2017). The Evolution
of Calcium-Based Signalling in Plants. *Curr Biol* 27(13)**,** R667-r679. doi:
10.1016/j.cub.2017.05.020.
Edgar, R.C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res 32(5)**,** 1792-1797. doi: 10.1093/nar/gkh340.
Flynn, R.L., and Zou, L. (2010). Oligonucleotide/oligosaccharide-binding fold proteins: a growing
family of genome guardians. *Crit Rev Biochem Mol Biol* 45(4)**,** 266-275. doi:
10.3109/10409238.2010.488216.
Fruchterman, T.M.J., and Reingold, E.M. (1991). Graph drawing by force-directed placement.
Software: Practice and Experience 21(11)**,** 1129-1164. doi: 10.1002/spe.4380211102.

Gertz, E.M., Yu, Y.K., Agarwala, R., Schaffer, A.A., and Altschul, S.F. (2006). Composition-based
statistics and translated nucleotide searches: improving the TBLASTN module of BLAST.
BMC Biol 4**,** 41. doi: 10.1186/1741-7007-4-41.
Gifford, J.L., Walsh, M.P., and Vogel, H.J. (2007). Structures and metal-ion-binding properties of the
Ca2+-binding helix-loop-helix EF-hand motifs. *Biochem J* 405(2)**,** 199-221. doi:
10.1042/BJ20070255.
Guardino, K.M., Sheftic, S.R., Slattery, R.E., and Alexandrescu, A.T. (2009). Relative stabilities of
conserved and non-conserved structures in the OB-fold superfamily. *Int J Mol Sci* 10(5)**,**
2412-2430. doi: 10.3390/ijms10052412.
Hildebrand, M.S., Sorensen, J.L., Jensen, M., Kimberling, W.J., and Smith, R.J. (2008). Autoimmune
disease in a DFNA6/14/38 family carrying a novel missense mutation in WFS1. *Am J Med*
Genet A 146A(17)**,** 2258-2265. doi: 10.1002/ajmg.a.32449.
Hofmann, S., Philbrook, C., Gerbitz, K.D., and Bauer, M.F. (2003). Wolfram syndrome: structural
and functional analyses of mutant and wild-type wolframin, the WFS1 gene product. *Hum*
Mol Genet 12(16)**,** 2003-2012.
Huang, J., and Gogarten, J.P. (2007). Did an ancient chlamydial endosymbiosis facilitate the
establishment of primary plastids? *Genome Biol* 8(6)**,** R99. doi: 10.1186/gb-2007-8-6-r99.
Inoue, H., Tanizawa, Y., Wasson, J., Behn, P., Kalidas, K., Bernal-Mizrachi, E., et al. (1998). A gene
encoding a transmembrane protein is mutated in patients with diabetes mellitus and optic
atrophy (Wolfram syndrome). *Nat Genet* 20(2)**,** 143-148. doi: 10.1038/2441.
Jekely, G. (2007). Origin of eukaryotic endomembranes: a critical evaluation of different model
scenarios. *Adv Exp Med Biol* 607**,** 38-51. doi: 10.1007/978-0-387-74021-8_3.
Jekely, G. (2008). Origin of the nucleus and Ran-dependent transport to safeguard ribosome
biogenesis in a chimeric cell. *Biol Direct* 3**,** 31. doi: 10.1186/1745-6150-3-31.
Jeong, H., Sim, H.J., Song, E.K., Lee, H., Ha, S.C., Jun, Y., et al. (2016). Crystal structure of SEL1L:
Insight into the roles of SLR motifs in ERAD pathway. *Sci Rep* 6**,** 20261. doi:
10.1038/srep20261.
Karpenahalli, M.R., Lupas, A.N., and Soding, J. (2007). TPRpred: a tool for prediction of TPR-,
PPR- and SEL1-like repeats from protein sequences. *BMC Bioinformatics* 8**,** 2. doi:
10.1186/1471-2105-8-2.
Kaufman, L., and Rousseeuw, P.J. (1990). *Finding Groups in Data: An Introduction to Cluster*
Analysis. New York: John Wiley & Sons.
Kawasaki, H., Nakayama, S., and Kretsinger, R.H. (1998). Classification and evolution of EF-hand
proteins. *Biometals* 11(4)**,** 277-295.
Kim, H.S., Czymmek, K.J., Patel, A., Modla, S., Nohe, A., Duncan, R., et al. (2012). Expression of
the Cameleon calcium biosensor in fungi reveals distinct Ca(2+) signatures associated with
polarized growth, development, and pathogenesis. *Fungal Genet Biol* 49(8)**,** 589-601. doi:
10.1016/j.fgb.2012.05.011.
Klimecka, M., and Muszynska, G. (2007). Structure and functions of plant calcium-dependent
protein kinases. *Acta Biochim Pol* 54(2)**,** 219-233.
Kozlov, G., Bastos-Aristizabal, S., Maattanen, P., Rosenauer, A., Zheng, F., Killikelly, A., et al.
(2010). Structural basis of cyclophilin B binding by the calnexin/calreticulin P-domain. *J Biol*
Chem 285(46)**,** 35551-35557. doi: 10.1074/jbc.M110.160101.
Krebs, J., Agellon, L.B., and Michalak, M. (2015). Ca(2+) homeostasis and endoplasmic reticulum
(ER) stress: An integrated view of calcium signaling. *Biochem Biophys Res Commun* 460(1)**,**
114-121. doi: 10.1016/j.bbrc.2015.02.004.
Krishna, S.S., Aravind, L., Bakolitsa, C., Caruthers, J., Carlton, D., Miller, M.D., et al. (2010). The
structure of SSO2064, the first representative of Pfam family PF01796, reveals a novel two-
domain zinc-ribbon OB-fold architecture with a potential acyl-CoA-binding role. *Acta*

Crystallogr Sect F Struct Biol Cryst Commun 66(Pt 10)**,** 1160-1166. doi:
10.1107/s1744309110002514.
Lance, G.N., and Williams, W.T. (1966). Computer Programs for Hierarchical Polythetic
Classification ("Similarity Analyses"). *The Computer Journal* 9(1)**,** 60-64. doi:
10.1093/comjnl/9.1.60.
Lanner, J.T., Georgiou, D.K., Joshi, A.D., and Hamilton, S.L. (2010). Ryanodine receptors: structure,
expression, molecular details, and function in calcium release. *Cold Spring Harb Perspect*
Biol 2(11)**,** a003996. doi: 10.1101/cshperspect.a003996.
Leipe, D.D., Wolf, Y.I., Koonin, E.V., and Aravind, L. (2002). Classification and evolution of P-loop
GTPases and related ATPases. *J Mol Biol* 317(1)**,** 41-72. doi: 10.1006/jmbi.2001.5378.
Lespinet, O., Wolf, Y.I., Koonin, E.V., and Aravind, L. (2002). The role of lineage-specific gene
family expansion in the evolution of eukaryotes. *Genome Res* 12(7)**,** 1048-1059. doi:
10.1101/gr.174302.
Lu, S., Kanekura, K., Hara, T., Mahadevan, J., Spears, L.D., Oslowski, C.M., et al. (2014). A
calcium-dependent protease as a potential therapeutic target for Wolfram syndrome. *Proc*
Natl Acad Sci U S A 111(49)**,** E5292-5301. doi: 10.1073/pnas.1421055111.
Mackinder, L., Wheeler, G., Schroeder, D., von Dassow, P., Riebesell, U., and Brownlee, C. (2011).
Expression of biomineralization-related ion transport genes in Emiliania huxleyi. *Environ*
Microbiol 13(12)**,** 3250-3265. doi: 10.1111/j.1462-2920.2011.02561.x.
Makioka, A., Kumagai, M., Kobayashi, S., and Takeuchi, T. (2002). Possible role of calcium ions,
calcium channels and calmodulin in excystation and metacystic development of Entamoeba
invadens. *Parasitol Res* 88(9)**,** 837-843. doi: 10.1007/s00436-002-0676-6.
Mans, B.J., Anantharaman, V., Aravind, L., and Koonin, E.V. (2004). Comparative genomics,
evolution and origins of the nuclear envelope and nuclear pore complex. *Cell Cycle* 3(12)**,**
1612-1637. doi: 10.4161/cc.3.12.1345.
Mittl, P.R., and Schneider-Brachert, W. (2007). Sel1-like repeat proteins in signal transduction. *Cell*
Signal 19(1)**,** 20-31. doi: 10.1016/j.cellsig.2006.05.034.
Moreno, S.N., and Docampo, R. (2003). Calcium regulation in protozoan parasites. *Curr Opin*
Microbiol 6(4)**,** 359-364.
Neuwald, A.F., and Altschul, S.F. (2016). Bayesian Top-Down Protein Sequence Alignment with
Inferred Position-Specific Gap Penalties. *PLoS Comput Biol* 12(5)**,** e1004936. doi:
10.1371/journal.pcbi.1004936.
Nolan, D.P., Reverlard, P., and Pays, E. (1994). Overexpression and characterization of a gene for a
Ca(2+)-ATPase of the endoplasmic reticulum in Trypanosoma brucei. *J Biol Chem* 269(42)**,**
26045-26051.
Ogris, C., Guala, D., and Sonnhammer, E.L.L. (2018). FunCoup 4: new species, data, and
visualization. *Nucleic Acids Res* 46(D1)**,** D601-D607. doi: 10.1093/nar/gkx1138.
Ong, H.L., de Souza, L.B., and Ambudkar, I.S. (2016). Role of TRPC Channels in Store-Operated
Calcium Entry. *Adv Exp Med Biol* 898**,** 87-109. doi: 10.1007/978-3-319-26974-0_5.
Osman, A.A., Saito, M., Makepeace, C., Permutt, M.A., Schlesinger, P., and Mueckler, M. (2003).
Wolframin expression induces novel ion channel activity in endoplasmic reticulum
membranes and increases intracellular calcium. *J Biol Chem* 278(52)**,** 52755-52762. doi:
10.1074/jbc.M310331200.
Peng, J., Ding, J., Tan, C., Baggenstoss, B., Zhang, Z., Lapolla, S.M., et al. (2009). Oligomerization
of membrane-bound Bcl-2 is involved in its pore formation induced by tBid. *Apoptosis*
14(10)**,** 1145-1153. doi: 10.1007/s10495-009-0389-8.
Perez-Gordones, M.C., Serrano, M.L., Rojas, H., Martinez, J.C., Uzcanga, G., and Mendoza, M.
(2015). Presence of a thapsigargin-sensitive calcium pump in Trypanosoma evansi:

Immunological, physiological, molecular and structural evidences. *Exp Parasitol* 159**,** 107-
117. doi: 10.1016/j.exppara.2015.08.017.
Plattner, H. (2015). Molecular aspects of calcium signalling at the crossroads of unikont and bikont
eukaryote evolution--the ciliated protozoan Paramecium in focus. *Cell Calcium* 57(3)**,** 174-
185. doi: 10.1016/j.ceca.2014.12.002.
Plattner, H. (2017). Signalling in ciliates: long- and short-range signals and molecular determinants
for cellular dynamics. *Biol Rev Camb Philos Soc* 92(1)**,** 60-107. doi: 10.1111/brv.12218.
Plattner, H., Flotenmeyer, M., Kissmehl, R., Pavlovic, N., Hauser, K., Momayezi, M., et al. (1999).
Microdomain arrangement of the SERCA-type Ca2+ pump (Ca2+-ATPase) in
subplasmalemmal calcium stores of paramecium cells. *J Histochem Cytochem* 47(7)**,** 841-
854. doi: 10.1177/002215549904700701.
Plattner, H., and Verkhratsky, A. (2013). Ca2+ signalling early in evolution--all but primitive. *J Cell*
Sci 126(Pt 10)**,** 2141-2150. doi: 10.1242/jcs.127449.
Plattner, H., and Verkhratsky, A. (2015). The ancient roots of calcium signalling evolutionary tree.
Cell Calcium 57(3)**,** 123-132. doi: 10.1016/j.ceca.2014.12.004.
Ponting, C.P., Aravind, L., Schultz, J., Bork, P., and Koonin, E.V. (1999). Eukaryotic signalling
domain homologues in archaea and bacteria. Ancient ancestry and horizontal gene transfer. *J*
Mol Biol 289(4)**,** 729-745. doi: 10.1006/jmbi.1999.2827.
Prakriya, M., and Lewis, R.S. (2015). Store-Operated Calcium Channels. *Physiol Rev* 95(4)**,** 1383-
1436. doi: 10.1152/physrev.00020.2014.
Price, M.N., Dehal, P.S., and Arkin, A.P. (2010). FastTree 2--approximately maximum-likelihood
trees for large alignments. *PLoS One* 5(3)**,** e9490. doi: 10.1371/journal.pone.0009490.
Prole, D.L., and Taylor, C.W. (2011). Identification of intracellular and plasma membrane calcium
channel homologues in pathogenic parasites. *PLoS One* 6(10)**,** e26218. doi:
10.1371/journal.pone.0026218.
Qian, X., Qin, L., Xing, G., and Cao, X. (2015). Phenotype Prediction of Pathogenic
Nonsynonymous Single Nucleotide Polymorphisms in WFS1. *Sci Rep* 5**,** 14731. doi:
10.1038/srep14731.
Reiner, D.S., Hetsko, M.L., Meszaros, J.G., Sun, C.H., Morrison, H.G., Brunton, L.L., et al. (2003).
Calcium signaling in excystation of the early diverging eukaryote, Giardia lamblia. *J Biol*
Chem 278(4)**,** 2533-2540. doi: 10.1074/jbc.M208033200.
Rigoli, L., Lombardo, F., and Di Bella, C. (2011). Wolfram syndrome and WFS1 gene. *Clin Genet*
79(2)**,** 103-117. doi: 10.1111/j.1399-0004.2010.01522.x.
Rodrigues, F.A., Costa Lda, F., and Barbieri, A.L. (2011). Resilience of protein-protein interaction
networks as determined by their large-scale topological features. *Mol Biosyst* 7(4)**,** 1263-
1269. doi: 10.1039/c0mb00256a.
Rouzier, C., Moore, D., Delorme, C., Lacas-Gervais, S., Ait-El-Mkadem, S., Fragaki, K., et al.
(2017). A novel CISD2 mutation associated with a classical Wolfram syndrome phenotype
alters Ca2+ homeostasis and ER-mitochondria interactions. *Hum Mol Genet* 26(9)**,** 1599-
1611. doi: 10.1093/hmg/ddx060.
Sillanpaa, J.K., Sundh, H., and Sundell, K.S. (2018). Calcium transfer across the outer mantle
epithelium in the Pacific oyster, Crassostrea gigas. *Proc Biol Sci* 285(1891). doi:
10.1098/rspb.2018.1676.
Silverio, A.L., and Saier, M.H., Jr. (2011). Bioinformatic characterization of the trimeric intracellular
cation-specific channel protein family. *J Membr Biol* 241(2)**,** 77-101. doi: 10.1007/s00232-
011-9364-8.
Smaili, S.S., Pereira, G.J., Costa, M.M., Rocha, K.K., Rodrigues, L., do Carmo, L.G., et al. (2013).
The role of calcium stores in apoptosis and autophagy. *Curr Mol Med* 13(2)**,** 252-265.

Soding, J. (2005). Protein homology detection by HMM-HMM comparison. *Bioinformatics* 21(7)**,**
951-960. doi: 10.1093/bioinformatics/bti125.
Strom, T.M., Hortnagel, K., Hofmann, S., Gekeler, F., Scharfe, C., Rabl, W., et al. (1998). Diabetes
insipidus, diabetes mellitus, optic atrophy and deafness (DIDMOAD) caused by mutations in
a novel gene (wolframin) coding for a predicted transmembrane protein. *Hum Mol Genet*
7(13)**,** 2021-2028.
Trandinh, C.C., Pao, G.M., and Saier, M.H., Jr. (1992). Structural and evolutionary relationships
among the immunophilins: two ubiquitous families of peptidyl-prolyl cis-trans isomerases.
Faseb j 6(15)**,** 3410-3420. doi: 10.1096/fasebj.6.15.1464374.
Unal, C.M., and Steinert, M. (2014). Microbial peptidyl-prolyl cis/trans isomerases (PPIases):
virulence factors and potential alternative drug targets. *Microbiol Mol Biol Rev* 78(3)**,** 544-
571. doi: 10.1128/mmbr.00015-14.
Urano, F. (2016). Wolfram Syndrome: Diagnosis, Management, and Treatment. *Curr Diab Rep*
16(1)**,** 6. doi: 10.1007/s11892-015-0702-6.
Ushioda, R., Miyamoto, A., Inoue, M., Watanabe, S., Okumura, M., Maegawa, K.I., et al. (2016).
Redox-assisted regulation of Ca2+ homeostasis in the endoplasmic reticulum by disulfide
reductase ERdj5. *Proc Natl Acad Sci U S A* 113(41)**,** E6055-E6063. doi:
10.1073/pnas.1605818113.
Venancio, T.M., Balaji, S., Iyer, L.M., and Aravind, L. (2009). Reconstructing the ubiquitin network:
cross-talk with other systems and identification of novel functions. *Genome Biol* 10(3)**,** R33.
doi: 10.1186/gb-2009-10-3-r33.
Verkhratsky, A., and Parpura, V. (2014). Calcium signalling and calcium channels: evolution and
general principles. *Eur J Pharmacol* 739**,** 1-3. doi: 10.1016/j.ejphar.2013.11.013.
Verma, R., Reichermeier, K.M., Burroughs, A.M., Oania, R.S., Reitsma, J.M., Aravind, L., et al.
(2018). Vms1 and ANKZF1 peptidyl-tRNA hydrolases release nascent chains from stalled
ribosomes. *Nature* 557(7705)**,** 446-451. doi: 10.1038/s41586-018-0022-5.
Wang, Y.Y., Zhao, R., and Zhe, H. (2015). The emerging role of CaMKII in cancer. *Oncotarget*
6(14)**,** 11725-11734. doi: 10.18632/oncotarget.3955.
Watson, E., Matousek, W.M., Irimies, E.L., and Alexandrescu, A.T. (2007). Partially folded states of
staphylococcal nuclease highlight the conserved structural hierarchy of OB-fold proteins.
Biochemistry 46(33)**,** 9484-9494. doi: 10.1021/bi700532j.
Woo, J.S., Srikanth, S., and Gwack, Y. (2018). "Modulation of Orai1 and STIM1 by Cellular
Factors," in *Calcium Entry Channels in Non-Excitable Cells,* eds. J.A. Kozak & J.W. Putney,
Jr. (Boca Raton (FL): CRC Press/Taylor & Francis
Yamada, T., Ishihara, H., Tamura, A., Takahashi, R., Yamaguchi, S., Takei, D., et al. (2006). WFS1-
deficiency increases endoplasmic reticulum stress, impairs cell cycle progression and triggers
the apoptotic pathway specifically in pancreatic beta-cells. *Hum Mol Genet* 15(10)**,** 1600-
1609. doi: 10.1093/hmg/ddl081.
Yin, X., Ziegler, A., Kelm, K., Hoffmann, R., Watermeyer, P., Alexa, P., et al. (2018). Formation and
mosaicity of coccolith segment calcite of the marine algae Emiliania huxleyi. *J Phycol* 54(1)**,**
85-104. doi: 10.1111/jpy.12604.
Yurimoto, S., Hatano, N., Tsuchiya, M., Kato, K., Fujimoto, T., Masaki, T., et al. (2009).
Identification and characterization of wolframin, the product of the wolfram syndrome gene
(WFS1), as a novel calmodulin-binding protein. *Biochemistry* 48(18)**,** 3946-3955. doi:
10.1021/bi900260y.
Zaremba-Niedzwiedzka, K., Caceres, E.F., Saw, J.H., Backstrom, D., Juzokaite, L., Vancaester, E., et
al. (2017). Asgard archaea illuminate the origin of eukaryotic cellular complexity. *Nature*
541(7637)**,** 353-358. doi: 10.1038/nature21031.

Zatyka, M., Da Silva Xavier, G., Bellomo, E.A., Leadbeater, W., Astuti, D., Smith, J., et al. (2015).
Sarco(endo)plasmic reticulum ATPase is a molecular partner of Wolfram syndrome 1 protein,
which negatively regulates its expression. *Hum Mol Genet* 24(3)**,** 814-827. doi:
10.1093/hmg/ddu499.
Zatyka, M., Ricketts, C., da Silva Xavier, G., Minton, J., Fenton, S., Hofmann-Thiel, S., et al. (2008).
Sodium-potassium ATPase 1 subunit is a molecular partner of Wolframin, an endoplasmic
reticulum protein involved in ER stress. *Hum Mol Genet* 17(2)**,** 190-200. doi:
10.1093/hmg/ddm296.
Zhou, X., Lin, P., Yamazaki, D., Park, K.H., Komazaki, S., Chen, S.R., et al. (2014). Trimeric
intracellular cation channels and sarcoplasmic/endoplasmic reticulum calcium homeostasis.
Circ Res 114(4)**,** 706-716. doi: 10.1161/circresaha.114.301816.
Zhou, Y., Yang, W., Kirberger, M., Lee, H.W., Ayalasomayajula, G., and Yang, J.J. (2006).
Prediction of EF-hand calcium-binding proteins and analysis of bacterial EF-hand proteins.
Proteins 65(3)**,** 643-655. doi: 10.1002/prot.21139.
Zucchi, R., and Ronca-Testoni, S. (1997). The sarcoplasmic reticulum Ca2+ channel/ryanodine
receptor: modulation by endogenous effectors, drugs and disease states. *Pharmacol Rev*
49(1)**,** 1-51.

1020

Identified gene loss/expansion events within lineages

loss ⊖   expansion ✱

Legend:
- Clearly-discernable archaeal origin
- Clearly-discernable bacterial origin
- Emergence in eukaryotes, derived from a bacterial origin
- Eukaryotic provenance or innovation of novel family in eukaryotes

Calstabin, CaMKs, Calcineurin B, TRIC, Homer, ERp57, ERdj5, TMX1, PPP2R3

Calcineurin A

Calreticulin, Calnexin

archaea

bacteria

LECA

Calmodulin family, SERCA

EHD/Sarcalumenin

IP3R

NCS1, Calcipressin

L-type VGCC, TRPC, WFS2

STIM, ERp44, RyR, Sorcin

ORAI

CREC family, SelN, Wolframin, Neurabin, Tespa1, Calsequestrin, S100

Bcl2

Diplomonadida

Parabasalia — PPP2R3 ✱, calnexin ✱

Heterolobosea

Kinetoplastids — PPP2R3 ✱

Haptophyta

Rhizaria

SAR

Stramenopiles — calstabin ✱, ORAI ✱

Apicomplexa — CaMK ✱, calstabin ✱, calnexin ✱

Ciliates — CaMK ✱, calcineurin A ✱

Archaeplastida

Animals — sorcin ⊖, calsequestrin ⊖, SelN ⊖, CaMK ✱, calmodulin ✱

Choanoflagellates

Filasterea

Fungi — ERdj5 ⊖, sarcalumenin ⊖, IP3R ⊖, ORAI ⊖, calmodulin ✱, SERCA ✱

Apusomonada

Amoebozoa — calcineurin A ✱

Plants — calcineurin A ⊖, HOMER ⊖, IP3R ⊖, TRIC ⊖, CaMK ✱, calmodulin ✱

Rhodophyta