Quantitative, super-resolution localization of small RNAs with sRNA-PAINT

Kun Huang [1,2], Feray Demirci [3], Blake C. Meyers [4,5]*, Jeffrey L. Caplan [1,2]*

[1] Department of Plant and Soil Sciences, University of Delaware, Newark, DE 19716, USA

[2] Bio-Imaging Center, Delaware Biotechnology Institute, University of Delaware, Newark, DE 19716, USA

[3] FiDoSoft Software Consulting, Redmond, WA, 98053, USA

[4] Donald Danforth Plant Science Center, 975 North Warson Road, St. Louis, Missouri 63132, USA

[5] University of Missouri – Columbia, Division of Plant Sciences, 52 Agriculture Lab, Columbia, MO 65211

* To whom correspondence should be addressed: Jeffrey L. Caplan; Tel: (302) 831-3403; Fax: (302) 831-4841; Email: jcaplan@udel.edu

Correspondence may also be addressed to: Blake C. Meyers; Tel: (314) 587-1422; Email: bmeyers@danforthcenter.org

Keywords

Small RNA, RNA detection, DNA-PAINT, *in situ* hybridization, super-resolution, single molecule, microscopy, maize anther, quantification

Abstract

Small RNAs are non-coding RNAs that play important roles in the life of both animals and plants. They

are 21- to 24-nt in length and around 10 nm in size. Their small size and high diversity has made it

challenging to develop methods that have sufficient resolution and specificity to multiplex and quantify.

We created a method for the detection of small RNA with nanometer resolution by combining the

super-resolution method, DNA-based points accumulation in nanoscale topography (DNA-PAINT), and

the specificity of locked nucleic acid (LNA) probes for the *in situ* detection of multiple small RNAs. The

method relies on designing probes to target small RNAs that combine the DNA oligos for PAINT with the

LNA-containing oligos for hybridization; therefore, we developed an online tool called "Vetting &

Analysis of RNA for *in situ* Hybridization probes" (VARNISH) for probe design. Our method utilizes

advances in DNA-PAINT methodologies, including qPAINT for quantification and Exchange-PAINT for

multiplexing. We demonstrated these capabilities or sRNA-PAINT by detecting and quantifying small

RNAs in different cell layers of early developmental stage maize anther that are important for male

sexual reproduction.

Introduction

In plants, 21- to 24-nucleotide (nt), non-coding small RNAs (sRNAs) regulate many important biological

processes [1]. Plant and animal genomes encode various sRNAs that can be divided in two major

categories: microRNAs (miRNAs) and small interfering RNAs (siRNAs) [2]. miRNAs derived from long,

hairpin precursor RNAs processed by precise cleavage, yielding mature miRNAs which influence gene

transcript levels and translation [3]. Plant siRNAs also derive from long precursors that are made double-

stranded by RNA-dependent RNA polymerases (RDRPs), are processed to siRNAs, and also regulate gene

transcript levels, through post-transcriptional silencing and epigenetically via RNA-directed DNA

methylation (RdDM) [4]. Some plant miRNAs trigger phased, secondary siRNAs (phasiRNAs), which are

either 21 or 24 nt in length[5-7]. Anthers, the plant male reproductive organ, are a particularly rich source

of miRNAs and phasiRNAs, particularly in grasses like maize [7]; during the meiotic stage, they are

composed of cell layers that include the epidermis, endothecium, middle, and tapetum, that all

surround the pollen mother cells. In maize anthers, the abundance and distribution of miRNAs and

phasiRNAs varies in different cell layers [7], and our interest in the spatial organization and concentration

of these sRNAs, particularly at the subcellular level, provided us a strong motivation to develop new

methods for fluorescent detection.

sRNAs are always on the move [8], moving within and between organelles, from cell-to-cell, long distances

within an organism, and there is increasing interest in their functions across organismal boundaries, in

plant-pathogen interactions. For example, within a cell, sRNAs move between the nucleus and

cytoplasm, bind Argonaute (AGO) proteins, Dicing bodies (D-bodies), Cajal bodies (CBs), and may be

polysome-associated or found with organelles such as the endoplasmic reticulum [9-12]. Intercellular

movement of sRNAs, presumably via plasmodesmata in plants, regulates tissue and organ development

by silencing and regulating developmentally important genes [13]. Systemic translocation of sRNAs has

been demonstrated by epigenetic silencing across graft junctions and sRNA profiling of vasculature [8, 14].

Remarkably, sRNAs also move between organisms at host-pathogen interfaces [15]. Pathogens, including

parasitic plants, regulate host defense responses via miRNAs and *trans*-species sRNAs, and *vice versa* [16].

In animals, extracellular vesicles (EVs) transport miRNAs from cell-to-cell and over long-distances [17].

Plant EVs are populated with miRNAs and fragments as small as 10 to 17 nt, the so-called tiny RNAs [2]. All

of these sRNAs vary in abundance from low to extremely high in abundance, as measured from

sequencing data, which typically utilizes gram quantities of plant tissue, substantially limiting spatial

analyses. And at the subcellular level, the localizations of these RNAs are not well-characterized. Thus,

future approaches to sRNA characterization will require nanometer resolution, high specificity, and both

multiplexing and quantitative capabilities.

For *in situ* hybridization of sRNAs, the use of locked nucleic acid (LNA) oligonucleotide probes has shown

specificity and high fidelity [18]. However, colorimetric and even fluorescent localization of sRNAs using

LNA-based methods are able to resolve only tissue-level localization patterns. Because of the limits of

light diffraction, lens-based microscopy cannot resolve spots below ~200 nm in diameter and 400 to 700

nm in axial length [19]. New techniques known as Photoactivated Localization Microscopy (PALM),

Structured Illumination microscopy (SIM), direct Stochastic Optical Reconstruction Microscopy

(dSTORM), and Points Accumulation for Imaging in Nanoscale Topography (PAINT) have been used to

drive microscopic detection beyond the diffraction limit, yielding so-called "super-resolution"

microspopy [19]. We have previously demonstrated an sRNA-FISH method combined with SIM and

dSTORM [20]. However, PALM and dSTORM utilize the blinking property of fluorescent dyes and rely on

specific excitation and buffer conditions to achieve proper imaging [21]. Photoswitching of the dye

molecule is hard to predict, which made absolute quantification difficult using these methods. The

number of dyes with the appropriate photoswitching properties within the visible spectrum is also a

limiting factor for multiplexed detection [21]. DNA-PAINT decouples the blinking events from dye

photophysics, using the binding and dwelling kinetics of a short dye-labeled oligonucleotide "imager

strand" to localize target-specific "docking strands" [22, 23]. The docking strand in DNA-PAINT is linked to an

antibody for single-molecule protein detection, with imager strands introduced by perfusion, one

imager strand at a time, each specific to a given docking strand and protein. After the image is acquired,

the imager strand is washed off and then the next imager strand is introduced [21]. As a result, a single dye

can be linked to different imager strands and applied for detection of numerous protein targets via use

of different imager strand-docking strand combinations. This multiplexing technique, known as

"Exchange-PAINT" has been used to *in situ* imaging of protein targets, such as microtubules and

mitochondria [23], and *in vitro* profiling of miRNAs to a synthetic DNA origami [24].

To achieve subcellular, nanometer resolution imaging for the localization and analysis of sRNAs, we

created a detection method called sRNA-PAINT. This method combines the high resolution and precise

quantification of DNA-PAINT with the efficiency and specificity of LNA-based *in situ* hybridization. We

created a probe design tool for our method and show that these sRNA-PAINT probes can used for robust

sRNA detection. And, we demonstrate that it can be combined with qPAINT to localize and quantify

distinct sRNAs in fixed biological sample and Exchange-PAINT for multiplexed target detection.

Results

sRNA-PAINT probe design

Most sRNA backbones are between 21 to 24 nt in length. To perform sRNA-PAINT, we designed a probe

that is composed of three sequences: the probe backbone sequence with LNA bases, the DNA-PAINT

docking strand sequence, and a linker sequence that connects those two (shown in Fig. 1). The probe

design is the most critical step of the sRNA-PAINT method, and therefore, we created an online tool

called VARNISH (Vetting & Analysis of RNA for *in situ* Hybridization probes) for automated design of

sRNA-PAINT probes (https://wasabi.ddpsc.org/~apps/varnish/). The tool requires the input of a target

sRNA sequence and the hybridization parameters (defaults provided), including sodium (50 mM) and

temperature (25°C). VARNISH first will reverse complement the sRNA sequence and will then chose

between 19-22 nt of the sequence such that the melting point temperature (T_m) is lower than 60°C, with

preference for lower T_m values for the longest sequence length. The melting temperature is calculated

by calling the "Analyze" function from the IDT web application program interface (API)

(https://www.idtdna.com/AnalyzerService/AnalyzerService.asmx). The final sequence is the probe

backbone (Fig. 1, cyan). Next, LNA bases and a linker sequence are introduced to the selected probe

backbone by VARNISH. The default linker sequence is "tattcgt", which has no match to the sRNA

sequences in miRbase, but this can be changed to any custom linker sequence. VARNISH introduces

between five and nine LNA bases (default is eight) and it avoids stretches of more than four LNA bases

and three or more Gs or Cs. The algorithm exhausts all the possible combinations of the positions for

LNA bases, and it calculates the approximate T_m using previously estimated T_m values for different LNAs

[25]. Next, VARNISH takes the top 200 probe backbone candidates with the highest TM T_m and it calculates

the exact T_m using the IDT web API. The last step is adding the DNA-PAINT docking strand. In the

VARNISH tool, the docking strand can be either chosen from a drop down menu containing the 13

previously used docking strands [23] or the sequence can be entered manually in the VARNISH tool. The

software will conduct a homo-dimer and secondary structure analysis from all or a subset of the 13

provided docking strands in combination with the linker and the top 200 probe backbone candidates.

These two steps are performed by calling the "SelfDimer" and "UNAFoldRun" functions from the IDT

web API. Finally, the algorithm will choose the top 10 candidates for each docking stand with the highest

T_m and lowest ΔG for homo-dimer and secondary structure predictions. Once the computations are

complete, VARNISH will send an email with a link to an output page containing the 10 top probe

candidates and the ordering information for both the probes and their corresponding imager strands.

sRNA-PAINT detects cellular sRNAs at single molecule resolution

Figure 2a shows the general pipeline for sRNA-PAINT. First, the samples are fixed, embedded in paraffin,

and sectioned. Then, thin sections were layered onto coverslips with gold fiducials sealed with silicon

dioxide (SiO_2); for this step, we highly recommend that no additional coatings are added, such as poly-l-

lysine, since they resulted in non-specific binding. Next, during the hybridization process, probes are

applied to fixed and sectioned samples (Fig. 2a). Excess probes are washed off in the washing process.

Afterward, imager strands are perfused in. When the un-annealed imager strands drift within the

buffer, the speed of the movement is too fast to be caught on by the microscope. However, when an

imager strand finds its docking partner, and dwells on it, the short time period before the imager strand

dissociates from the docking creates a blinking event that can be detected using a super-resolution

microscope. As a result, when thousands of these blinks events are collected, detection of all the

docking sites within the sample is eventually completed, defining the locations of the sRNA targets.

For sRNA-PAINT probe labeling, we evaluated Alexa Fluor® 647 (AF 647), ATTO 665, and ATTO 488. The

AF647 dye resulted in the lowest levels of background binding in the negative control samples, whereas

ATTO 665 bound to paraffin-embedded plant tissue non-specifically, and ATTO 488 dye emission

wavelength overlaps with the strong autofluorescence that is typical of paraffin embedded plant tissue –

thus we selected AF674 as the best label (data not shown). The sRNA probes designed by VARNISH were

hybridized and the coverslip was mounted in perfusion chamber. The corresponding, 3'-dye-labeled

imager strands were continuously perfused and imaging was conducted using total internal reflection

fluorescence (TIRF) microscopy. As a demonstration, we performed sRNA-PAINT on an abundant 24-nt

phasiRNA – a class of plant secondary siRNAs abundant in the early developmental stages of maize

anthers [7] (Fig. 1b); the 24-nt phasiRNAs are triggered by a miRNA, miR2275, providing us with two

different classes of small RNAs in one tissue. After the 1st round of imaging, buffer was perfused into the

imaging chamber, and the signal was diminished rapidly within 1 min, suggesting that stripping off the

imager strand is very efficient. And then, sRNA-PAINT signal was re-achieved by re-applying the imager

strands (Fig. 1a, b). Different imager strands had no or very little non-specific binding, which is similar to

the original DNA-PAINT method [23]. The scrambled, control probe detected very little to no background

(Supplementary Figure 1), which showed that the imager strand does not bind non-specifically to the

plant tissue we used. Our sRNA-PAINT experiment was consistent with our previous findings that the 24-

nt phasiRNA was present in all of the anther cell layers, and most abundant in the tapetum layer and

pollen mother cells [7]; however, our, sRNA-PAINT approach achieved a much higher resolution of below

10 nm (Fig. 1 c).

Calculation of sRNA copy number with qPAINT

qPAINT can count molecules by analyzing the predictable and programmable binding kinetics of the

imager strand to the docking strand [22]. This quantitative method is based solely on binding kinetics, and

therefore, individual molecules do not need to be resolved. Picasso software is publicly available and

can perform this calculation for the binding sites within a selected region [21]. We selected a 3.18 μm (20

pixel) diameter circle in the cytoplasm to represent the copy number in the cytoplasmic region of each

cell in each layer, and we performed qPAINT using the same image in Figure 2b. After the count, we

calculated the number of binding sites in 63.25 μm^2, which is roughly the area of the cytoplasm of cells

in our sections. As shown in Figure 3, we detected 10.41 copies of 24-nt phasiRNA per selected region

per cell in epidermal cells, 11.04 copies in endothecial cells, 22.64 copies in middle layer cells, 72.68

copies in pollen mother cells, and most abundantly, 61.05 copies in tapetal cells. Pollen mother cells and

tapetum have a greater abundance of the 24-nt phasiRNA compared with the other cell layers. qPAINT,

in combination with sRNA-PAINT, allowed the detection of the exact copy number of sRNAs in cell

sections and the relative sRNA targets in the different cell layers. These data will be useful to ground-

truth sequencing data, currently the predominant method of measuring sRNA.

Detection of miRNA and phasiRNA with sRNA-Exchange-PAINT

Sequential detection of sRNAs can be done by multiplex VARNISH probes with different docking strand

combinations. We name this technique sRNA-Exchange-PAINT, as a variation of the protein-focused,

multiplexed Exchange-PAINT method. For this analysis, we measured both miR2275 and the most

abundant 24-nt phasiRNA it triggers, again in maize anthers. As shown in the Figure 2b, the imager

strand used for miR2275 (P0*) and 24-nt phasiRNA (P1*) do not non-specifically bind to each other's

docking strand. Fixed sample were first perfused with P0* imager strand. After the image of miR2275

was acquired, buffer wash was performed to completely remove the P0* imager strand. And then

imager strand P1* was applied, and generated the image for 24-nt phasiRNA. Using traditional *in situ*

hybridization methods, miR2275 and 24-nt phasiRNA were thought to co-localize to each other in the

tapetum layer and pollen mother cells. Using sRNA-PAINT, which has the resolution limit down to 10

nm, we confirmed that they both localize to the tapetum layer and pollen mother cells (Fig. 4 B), but our

results show that they do not co-localize at the single molecule level (Supplementary Fig. 2).

Discussion

sRNA-PAINT provides a robust method that quantitatively detects small RNAs with single-molecule

resolution. It greatly improves the resolution and precision of sRNA localization compared to other FISH

methods, namely those that are limited by the diffraction limit of light. Quantification via dSTORM is

hindered by photobleaching that results in an underestimation of the targets, particularly, sRNAs [20]. The

small size of sRNAs are generally only detected by one fluorophore per sRNA, as a result, each dye play

an important role in terms of quantification. In contrast to other super-resolution methods, such as SR-

SIM and dSTORM, absolute quantification of binding sites can be deduced from the known binding

kinetics of the imager strands to the docking strands with sRNA-PAINT. Another benefit of using sRNA-

PAINT is that it takes less time than regular *in situ* hybridization methods, since it by-passes the antibody

incubation and amplification steps that typify those methods. Samples are directly imaged after

hybridization and washing, bringing the sample preparation time down from 4 days to overnight. The

signal amplification of sRNA-PAINT occurs at the same time as image acquisition, although the time of

image acquisition for sRNA-PAINT is longer than traditional sRNA-FISH methods due to the need to take

thousands of images for one field of view for detection. Another benefit of bypassing the antibody

amplification steps of traditional *in situ* techniques is that the primary and secondary antibodies add

significant amount of bulk, decreasing the precision of localization by 24 - 30 nm [26]. Similarly, our simple,

single-oligo VARNISH probe design does not require biotin and streptavidin like the original DNA-PAINT

method. The primary drawback to sRNA-PAINT is simply the cost of the LNA oligos, which makes it

expensive for new probes, although this cost is amortized if the probe is used for many samples, as a

typical yield for a purchased LNA oligo is enough for innumerable sRNA-PAINT experiments.

With the ability to count the exact copy number, this method could be used to investigate many

biological questions that require precise subcellular localization or quantitative data, from small RNA

biogenesis to the function of sRNAs. Images acquired with sRNA-PAINT may even change the way we

consider sRNA localization. The size of a small RNA is about 10 nm [27] and the imaging resolution of our

sRNA-PAINT is sub-10 nm. Colocalization of sRNA-PAINT should be evaluated in the context of spatial

information at the tissue, subcellular and molecular level. In case of miR2275 and the 24-nt phasiRNA,

in situ hybridization of these two sRNAs showed that these two co-localize at tissue level in tapetal layer

and pollen mother cells, and, at the subcellular level mainly in the cytoplasm. However, at the molecular

level, we observed minimal colocalization because sRNA-PAINT has single molecule resolution. Co-

localization of miR2275 and its phasiRNA at the molecular level would suggest that they physically

interact with each other or that their density is too high for the method to resolve individual target sRNA

molecules. Future plan includes generating an analysis pipeline for sRNA-PAINT co-localization that will

be most useful for sRNA-mRNA and sRNA-protein interactions, to examine outstanding questions in

sRNA biogenesis and processing.

A distinct advantage of sRNA-PAINT is that it is compatible with Exchange-PAINT, for multiplexing using

multiple docking strands that are linked to of the same fluorophore [23]. Our previously describe sRNA-

FISH [20] is limited to only 2-3 targets, mainly due to the limited choices of antibody-based amplification

and fluorophore choice. sRNA-PAINT only requires the use of single fluorophore and therefore it is not

prone to differences in photophysical properties of different flourophores for different targets. Our

antibody-free method potentially can be massively multiplexed using numerous probes designed via

VARNISH to query many targets in the same sample. The Exchange-PAINT for protein targets predicts

that there should be hundreds of non-interacting docking sequences for multiplexing [23], and our sRNA-

Exchange-PAINT should have similar multiplexing potential.

Finally, our sRNA-PAINT method provides an alternative to conventional sRNA-FISH, as it is capable of

higher resolution, quantification, and multiplexing. All FISH-based methods are dependent on the actual

hybridization efficiency of the probes prior to imaging. The development of LNA technology increased

the specificity and efficiency of sRNA-FISH methods; future improvements in probes may provide

thorough yet specific labeling of target sRNAs that will benefit all FISH-based RNA detection methods,

including sRNA-PAINT. One strategy for increasing specificity is to use a split probe for detection of one

sRNA, such that each probe backbone will be 9-10 nt in length. By combining signals from both probe,

specific signal could be obtained for each target. Another drawback of the sRNA-PAINT method is that it

can only applied to fixed tissues or cells. Live-cell imaging, especially aptamer-based sensor imaging for

small RNAs, was made possible by fluorescent RNA aptamers such as Spinach and Mango [28-30]. Creating a

live-cell, single-molecule level of sRNA-PAINT would shed light on the dynamics and ever changing

contents of cellular RNAs.

Methods

Plant materials and oligonucleotides.

Maize anthers from W23 were provided by the Walbot lab at Stanford University (Palo Alto, CA). Plants

were grown in Palo Alto, CA under greenhouse conditions. Anther dissection and measurements were

performed as previously described [31]. LNA-modified oligonucleotide probes were designed using our

VARNISH software and synthesized by Exiqon (QIAGEN; Germantown, MD). Imager strands coupled with

AF647 were ordered from IDT (Coralville, Iowa).

Sample preparation.

Sample preparation for *in situ* hybridization was performed as previously described [20]. Briefly, anthers

were dissected and fixed in a 50 ml tube using 4% paraformaldehyde in 1xPHEM buffer (5 mM HEPES, 60

mM PIPES, 10 mM EGTA, 2 mM $MgSO_4$ at pH 7). Samples were then processed in a vacuum chamber

(0.08 MPa) three times, 15 min each. After fixation, samples were embedded in paraffin at the

Histochemistry and Tissue Processing Core Lab at Nemours/Alfred I. duPont Hospital for Children

(Wilmington, DE). Paraffin samples were sectioned at 6 µm thickness using a paraffin microtome and

dried on a Wide Spectral Band 600+/- 100 nm Gold Fiducials coverglass (600-100AuF; Hestzig LLC,

Leesburg, VA) at 37° C degree on a slider warmer.

In situ *hybridization.*

The *in situ* hybridization step was performed following our previously-published protocol [20]. Briefly,

samples were de-paraffinized using Histo-Clear (item 50-899-90147; Fisher Scientific, Pittsburgh, PA) and

re-hydrated by going through an ethanol series of 95, 80, 70, 50, 30, 10% (vol/vol) and water (1 min) at

room temperature. Then samples were treated with protease (item P5147; Sigma-Aldrich, St. Louis, MO)

for 20 min at 37°C. Excess formaldehyde background was removed by treating samples with 0.2%

glycine (item G8898; Sigma-Aldrich) for 20 min. After two washes in 1xPBS buffer (phosphate-buffered

saline), samples were de-hydrated by going through an ethanol series of 10, 30, 50, 70, 80, 95%, and

100% (vol/vol). Hybridization was done with 10 uM probe at 53.3°C in a hybridization oven. After

hybridization, samples were washed twice with 0.2x SSC buffer (saline-sodium citrate). Sample were

then washed in 1x BPS buffer for 10 min. Hybridized samples can be kept in 1xPBS at 4°C until imaging.

Super-resolution optical.

Super-resolution imaging was carried out on an inverted Zeiss Elyra PS.1 super-resolution microscope

(Carl Zeiss, Lberkochen, Germany). TIRF illumination was done using a 100% 642 nm laser and α-Plan-

Apochromat 100x/1.46 oil objective. For each imager strand, as well as control imager strand, images

were taken with an exposure time of 100 ms, an EMCCD Gain 30, and 20,000 frames in total.

PAINT.

A DH40iL culture dish incubate system (model 640388; Warner Instruments LLC, Hamden, CT) and a

quick release magnetic chamber for 25 mm low profile, round coverslips (model 641943; Warner

Instruments LLC) were assembled to be used as the perfusion chamber. A ValveLink8.2 Perfusion System

(AutoMate Scientific, Berkeley, CA) was used for perfusing buffer and imager strand solutions into the

chamber. All imager strands were diluted to 0.5-2 nM in buffer C (1× PBS, 500 mM NaCl, pH 8). Imagers

were taken with constant flow of imager stand. A Masterflex C/L peristaltic pump (60 RPM, model

77120-62; Cole-Parmer Instrument Company LLC, Vernon Hills, IL) was used to collect the buffer flow.

Super-resolution image reconstruction.

Each image was analyzed in Zen software (Carl Zeiss). The images were processed using the same

parameters, which are listed below: Ignore overlapping molecules; Peak Mask Size (6); Peak Intensity to

Noise (6.0); Fit model (x,y 2D Gauss Fit). Drift correction was done by using fiducial-based algorithm. For

image rendering, pixel resolution was set to 16 nm/pixel and 2 nm/pixel for zoomed in images with 1x

PSF (point spread function) expansion factor.

qPAINT data analysis.

qPAINT data analysis was done following the protocol by Schnitzbauer et al. [21] using the Picasso

software package. In brief, 8000 fames of the raw movie file of PAINT data were processed with

"Picasso: Localize". We adjusted the threshold until only the PAINT spots were detected and selected.

After "Localize (Identify & Fit)", a .hdf5 file was generated and this was used as the input for the next

module. The .hdf5 file was opened in "Picasso: Render", and marker-based drift correction or redundant

cross-correlative drift correction was performed. A 20 pixel diameter region of interest in the cytoplasm

of each anther cell layer was picked by using "Tools: Pick". The dark time under "View: Show info" was

used as the input for calculating the influx rate via $\xi = k_{on} \times c$, and the influx rate was used to calculate

the copy number.

sRNA sequences.

All small RNA probe sequences and imager strands used are listed below, designing an LNA nucleotide

after the "+", the lowercase nucleotides indicate the linker, and the docking strand is underscored:

Scrambled control probe with P0 docking strand:

GTG+T+AA+CACG+T+C+T+A+TACGCCCAtattcgat<u>TTTCAATGTAT</u>

miR2275: TTCAGTTTCCTCTAATATCTCA

Probe for miR2275 with linker and P0 docking strand:

T+GA+GA+TAT+TA+GAG+GAAA+C+TGAAtattcgat<u>TTTCAATGTAT</u>

P0* imager strand: 5'-CATACATTGA-AF647

24 nt-phasiRNA: AGTTGTTGGACCGGACCTTGGCCC

Probe for 24-nt phasiRNA with linker and P1 docking strand:

GGC+C+AAG+G+TC+C+GGTCCA+A+CAACTtattcgat<u>TTATACATCTA</u>

P1* imager strand: 5'-CTAGATGTAT-AF647

Acknowledgements

This project was supported by the US NSF Plant Genome Research Program, awards 1649424, 1611853

and 1754097. We thank members of the Meyers and Caplan labs for help and support. We thank

Virginia Walbot (Stanford University) and members of her lab for supplying maize materials and for

useful discussions on anther small RNAs. We thank Prof. Dr. Ralf Jungmann for his support in DNA-

PAINT, qPAINT and analysis with Picasso. Microscopy equipment was acquired with a shared

321   instrumentation grant (S10 OD016361) and access was supported by the NIH-NIGMS (P20 GM103446),

322   the NSF (IIA-1301765) and the State of Delaware.

References

1. Lewsey, M.G. et al. Mobile small RNAs regulate genome-wide DNA methylation. *Proc Natl Acad Sci U S A* **113**, E801-810 (2016).

2. Baldrich, P. et al. Plant Extracellular Vesicles Contain Diverse Small RNA Species and Are Enriched in 10-to 17-Nucleotide "Tiny" RNAs. *Plant Cell* **31**, 315-324 (2019).

3. Wight, C., Ross, D.A. & Weinberger, D.R. Small RNAs May Answer Big Questions in Mental Illness. *Biological Psychiatry* **83**, E1-E3 (2018).

4. Borges, F. & Martienssen, R.A. The expanding world of small RNAs in plants. *Nature Reviews Molecular Cell Biology* **16**, 727-741 (2015).

5. Fei, Q., Xia, R. & Meyers, B.C. Phased, secondary, small interfering RNAs in posttranscriptional regulatory networks. *Plant Cell* **25**, 2400-2415 (2013).

6. Xia, R. et al. 24-nt reproductive phasiRNAs are broadly present in angiosperms. *Nature Communications* **10** (2019).

7. Zhai, J.X. et al. Spatiotemporally dynamic, cell-type-dependent premeiotic and meiotic phasiRNAs in maize anthers. *P Natl Acad Sci USA* **112**, 3146-3151 (2015).

8. Chitwood, D.H. & Timmermans, M.C.P. Small RNAs are on the move. *Nature* **467**, 415-419 (2010).

9. Fang, Y.D. & Spector, D.L. Identification of nuclear dicing bodies containing proteins for microRNA biogenesis in living Arabidopsis plants. *Current Biology* **17**, 818-823 (2007).

10. Li, C.F. et al. An ARGONAUTE4-containing nuclear processing center colocalized with Cajal bodies in Arabidopsis thaliana. *Cell* **126**, 93-106 (2006).

11. Li, S.B. et al. Biogenesis of phased siRNAs on membrane-bound polysomes in Arabidopsis. *Elife* **5** (2016).

12. Song, L., Han, M.H., Lesicka, J. & Fedoroff, N. Arabidopsis primary microRNA processing proteins HYL1 and DCL1 define a nuclear body distinct from the Cajal body. *P Natl Acad Sci USA* **104**, 5437-5442 (2007).

13. Dunoyer, P., Melnyk, C., Molnar, A. & Slotkin, R.K. Plant Mobile Small RNAs. *Cold Spring Harbor Perspectives in Biology* **5** (2013).

14. Yoo, B.C. et al. A systemic small RNA signaling system in plants. *Plant Cell* **16**, 1979-2000 (2004).

15. Johnson, N.R. & Axtell, M.J. Small RNA warfare: exploring origins and function of trans-species microRNAs from the parasitic plant Cuscuta. *Curr Opin Plant Biol* **50**, 76-81 (2019).

16. Cai, Q. et al. Plants send small RNAs in extracellular vesicles to fungal pathogen to silence virulence genes. *Science* **360**, 1126-1129 (2018).

17. Liu, T. et al. EVmiRNA: a database of miRNA profiling in extracellular vesicles. *Nucleic Acids Research* **47**, D89-D93 (2019).

18. Javelle, M. & Timmermans, M.C.P. In situ localization of small RNAs in plants by using LNA probes. *Nature Protocols* **7**, 533-541 (2012).

19. Hell, S.W. Microscopy and its focal switch. *Nature Methods* **6**, 24-32 (2009).

20. Huang, K., Baldrich, P., Meyers, B.C. & Caplan, J.L. sRNA-FISH: versatile fluorescent in situ detection of small RNAs in plants. *Plant Journal* **98**, 359-369 (2019).

21. Schnitzbauer, J., Strauss, M.T., Schlichthaerle, T., Schueder, F. & Jungmann, R. Super-resolution microscopy with DNA-PAINT. *Nature Protocols* **12**, 1198-1228 (2017).

22. Jungmann, R. et al. Quantitative super-resolution imaging with qPAINT. *Nature Methods* **13**, 439-442 (2016).

23. Jungmann, R. et al. Multiplexed 3D cellular super-resolution imaging with DNA-PAINT and Exchange-PAINT. *Nature Methods* **11**, 313-318 (2014).

24. Xu, W.D., Yin, P. & Dai, M.J. Super-resolution Geometric Barcoding for Multiplexed miRNA
Profiling. *Angewandte Chemie-International Edition* **57**, 14075-14079 (2018).
25. McTigue, P.M., Peterson, R.J. & Kahn, J.D. Sequence-dependent thermodynamic parameters; for
locked nucleic acid (LNA)-DNA duplex formation. *Biochemistry* **43**, 5388-5405 (2004).
26. Pleiner, T. et al. Nanobodies: site-specific labeling for super-resolution imaging, rapid epitope-
mapping and native protein complex isolation. *Elife* **4** (2015).
27. Yoffe, A.M. et al. Predicting the sizes of large RNA molecules. *P Natl Acad Sci USA* **105**, 16153-
16158 (2008).
28. Autour, A. et al. Fluorogenic RNA Mango aptamers for imaging small non-coding RNAs in
mammalian cells. *Nature Communications* **9** (2018).
29. Huang, K. et al. FASTmiR: an RNA-based sensor for in vitro quantification and live-cell
localization of small RNAs. *Nucleic Acids Research* **45** (2017).
30. Paige, J.S., Wu, K.Y. & Jaffrey, S.R. RNA Mimics of Green Fluorescent Protein. *Science* **333**, 642-
646 (2011).
31. Kelliher, T. & Walbot, V. Emergence and patterning of the five cell types of the *Zea mays* anther
locule. *Dev Biol* **350**, 32-49 (2011).
386
387

 Figure legends

Figure 1. Illustration of an sRNA-PAINT probe.

An sRNA-PAINT probe is composed of three parts: a probe backbone (cyan) that is complementary to

the target sRNA (orange), a docking strand (green) and a linker sequence (dark blue) connecting them.

The imager strand is conjugated to the fluorophore, and it will find its corresponding docking strand

during the imaging process.

Figure 2. sRNA-PAINT on a 24 nt-phasiRNA in an early-stage maize anther.

a. A work flow of sRNA-PAINT. Sample were fixed and embedded, and a thin section is placed on the

coverglass. During the hybridization process, the VARNISH probes will bind to sRNAs in the sample. **b.** an

example of the sRNA-PAINT imaging process. In the first imaging step, imager strands are added to the

probe-hybridized sample on the coverglass. After the image is acquired, the imager strands are washed

off with a short buffer wash step. In the second imaging step, the signal can be re-detected when re-

applying the imager strand to the sample. A non-specific imager stand was used as a negative control. **c.**

Detection of a 24-nt phasiRNA in a cross-section of maize anther. The sRNA-PAINT signal was detected in

all cell layers but it is most abundant in the tapetal layer and pollen mother cells. **c.** Zoomed images of

boxes 1 and 2 in panel b. There are more 24-nt phasiRNA localization events in the pollen mother cells

(box 1) compared to the epidermal and endothecium layers (box 2). The dotted line (light blue) marks

the division between the epidermal (above the line) and endothecium layers (below the line). Expanded

boxed areas (white) show representative areas at higher magnification. A red-to-white lookup table was

used to display low-to-high intensity values.

Figure 3. Quantification of binding site per cell in each layer.

Cartoon diagram representing each cell layer of a cross section of a maize anther lobe: epidermis

(yellow), endothecium (green), middle layer (red), tapetum (blue), and pollen mother cells (purple). The

Picasso software was used to calculate the binding sites by choosing a region of interest in the

cytoplasmic area of each cell layer. 10.41, 22.64, 72.68 and 61.05 copies of 24-nt phasiRNA per selected

region per cell in epidermal cells, endothecial cells, middle layer cells, and pollen mother cells

respectively.

Figure 4. sRNA-Exchange-PAINT

a. A diagram showing sRNA-Exchange-PAINT, which differs from sRNA-PAINT via the use of multiple

imager strands. Imager strand 1 is first introduced to the sample, and after imaging, imager strand 1 is

removed with a buffer wash. Next, imager strand 2 is introduced. This process can be repeated until the

needed number of targets is reached. **b.** sRNA-Exchange-PAINT on miR2275 (yellow) and a 24-nt

phasiRNA (blue) in maize anther. After the buffer wash, imaging showed little or no signal following each

round of detection using the imager strands. Scale bar = 10 µm for all images.

Supplemental figure legends

Supplementary Figure 1. sRNA-PAINT showed limited to no background with a scrambled control probe.

a. sRNA-PAINT with probes for a 24-nt phasiRNA and a scrambled control. Both docking strands were

detected with corresponding imager strands. Arrows indicate 2 out of ~66 fiducials used in experiments.

Scrambled control detected little or no signal. Scale bar = 10 µm for all images. **b.** qPAINT analysis of

image in panel a. Number showed the binding events for each cell (80 pixel size diameter circle in

Picasso software) in different cell layers.

Supplementary Figure 2. Subcellular localization analysis of miR2275 and 24-nt phasiRNA in maize

anther.

Right panel: zoomed image of the marked region in the left panel. These two sRNAs do not co-localize at

a nanometer resolution.

Fig 1.

Figure 1. Illustration of an sRNA-PAINT probe.
An sRNA-PAINT probe is composed of three parts: a probe backbone (cyan) that is complementary to the target sRNA (orange), a docking strand (green) and a linker sequence (dark blue) connecting them. The imager strand is conjugated to the fluorophore, and it will find its corresponding docking strand during the imaging process.

Fig. 2

Figure 2. sRNA-PAINT on a 24 nt-phasiRNA in an early-stage maize anther.
a. A work flow of sRNA-PAINT. Sample were fixed and embedded, and a thin section is placed on the coverglass. During the hybridization process, the VARNISH probes will bind to sRNAs in the sample. b. an example of the sRNA-PAINT imaging process. In the first imaging step, imager strands are added to the probe-hybridized sample on the coverglass. After the image is acquired, the imager strands are washed off with a short buffer wash step. In the second imaging step, the signal can be re-detected when re-applying the imager strand to the sample. A non-specific imager stand was used as a negative control. c. Detection of a 24-nt phasiRNA in a cross-section of maize anther. The sRNA-PAINT signal was detected in all cell layers but it is most abundant in the tapetal layer and pollen mother cells. c. Zoomed images of boxes 1 and 2 in panel b. There are more 24-nt phasiRNA localization events in the pollen mother cells (box 1) compared to the epidermal and endothecium layers (box 2). The dotted line (light blue) marks the division between the epidermal (above the line) and endothecium layers (below the line). Expanded boxed areas (white) show representative areas at higher magnification. A red-to-white lookup table was used to display low-to-high intensity values.

Fig.3

Figure 3. Quantification of binding site per cell in each layer.
Cartoon diagram representing each cell layer of a cross section of a maize anther lobe: epidermis (yellow), endothecium (green), middle layer (red), tapetum (blue), and pollen mother cells (purple). The Picasso software was used to calculate the binding sites by choosing a region of interest in the cytoplasmic area of each cell layer. 10.41, 22.64, 72.68 and 61.05 copies of 24-nt phasiRNA per selected region per cell in epidermal cells, endothecial cells, middle layer cells, and pollen mother cells respectively.

Fig.4

Figure 4. sRNA-Exchange-PAINT
a. A diagram showing sRNA-Exchange-PAINT, which differs from sRNA-PAINT via the use of multiple imager strands. Imager strand 1 is first introduced to the sample, and after imaging, imager strand 1 is removed with a buffer wash. Next, imager strand 2 is introduced. This process can be repeated until the needed number of targets is reached. b. sRNA-Exchange-PAINT on miR2275 (yellow) and a 24-nt phasiRNA (blue) in maize anther. After the buffer wash, imaging showed little or no signal following each round of detection using the imager strands. Scale bar = 10 μm for all images.

Supplementary Fig. 1

Supplementary Figure 1. sRNA-PAINT showed limited to no background with a scrambled control probe.
a. sRNA-PAINT with probes for a 24-nt phasiRNA and a scrambled control. Both docking strands were detected with corresponding imager strands. Arrows indicate 2 out of ~66 fiducials used in experiments. Scrambled control detected little or no signal. Scale bar = 10 μm for all images. b. qPAINT analysis of image in panel a. Number showed the binding events for each cell (80 pixel size diameter circle in Picasso software) in different cell layers.

Supplementary Fig. 2

Supplementary Figure 2. Subcellular localization analysis of miR2275 and 24-nt phasiRNA in maize anther. Right panel: zoomed image of the marked region in the left panel. These two sRNAs do not co-localize at a nanometer resolution.