Title page

Title

State and trait characteristics of anterior insula time-varying functional connectivity

Authors

Lorenzo Pasquini[1]*, Gianina Toller[1], Adam Staffaroni[1], Jesse A. Brown[1], Jersey Deng[1], Alex Lee[1], Katarzyna

Kurcyus[2,3], Suzanne M. Shdo[5], Isabel Allen[1], Virginia E. Sturm[1], Yann Cobigo[1], Valentina Borghesani[1],

Giovanni Battistella[1], Maria Luisa Gorno-Tempini[1], Katherine P. Rankin[1], Joel Kramer[1], Howard H. Rosen[1],

Bruce L. Miller[1], William W. Seeley[1,4]

Affiliations

Memory and Aging Center, Departments of Neurology[1] and Pathology[4], University of California, San

Francisco, California USA; Departments of Neuroradiology[2] and Nuclear Medicine[3], Klinikum rechts der

Isar, Technische Universität München, Munich, Germany, [5]Psychology, University of California Berkeley

*Corresponding author: William W. Seeley, MD. 675 Nelson Rising Lane 94158, San Francisco, California

USA. Email: bill.seeley@ucsf.edu. Phone: +1-415-476-2793

Running title: Time-varying functional connectivity of the anterior insula

Key words: Anterior insula, socio-emotional functioning, time-varying functional connectivity, trait, state

Abbreviations

Anterior insula = aINS; BOLD = blood oxygen level dependent; inferior frontal gyrus = IFG; task-free

functional MRI = tf-fMRI

Abstract

The human anterior insula (aINS) is a topographically organized brain region, in which ventral portions
contribute to socio-emotional function through limbic and autonomic connections, whereas the dorsal
aINS contributes to cognitive processes through frontal and parietal connections. Open questions remain,
however, regarding how aINS connectivity varies over time. We implemented a novel approach combining
seed-to-whole-brain sliding-window functional connectivity MRI and k-means clustering to assess time-
varying functional connectivity of aINS subregions. We studied three independent large samples of
healthy participants and longitudinal datasets to assess inter- and intra-subject stability, and related aINS
time-varying functional connectivity profiles to dispositional empathy. We identified four robust aINS
time-varying functional connectivity modes that displayed both "state" and "trait" characteristics: while
modes featuring connectivity to sensory regions were modulated by eye closure, modes featuring
connectivity to higher cognitive and emotional processing regions were stable over time and related to
empathy measures.

Introduction

The human anterior insula (aINS) is a functionally heterogeneous region implicated in functions ranging from interoceptive awareness and emotion processing to time perception and cognitive control (1). In humans, neuroimaging studies have begun to parcellate the aINS, based on its patterns of functional and structural connectivity. For example, task-free fMRI (tf-fMRI) studies, which measure the brain-wide correlation structure in slow (< 0.1 Hz), spontaneous blood oxygen level dependent (BOLD) signal fluctuations (2, 3), have shown that the ventral, agranular aINS is functionally connected to limbic and autonomic processing regions that include the pregenual anterior cingulate cortex, the amygdala, and subcortical structures such as the thalamus and periaqueductal gray (4–8). These regions make up the "salience network", a large-scale distributed system that represents the homeostatic significance of prevailing stimuli and conditions (9–13). In contrast, the dorsal, dysgranular aINS is connected to a cingulo-opercular "task-control network" whose nodes include dorsolateral and opercular prefrontal, anterior midcingulate, and anterior parietal areas involved in cognitive control processes such as task-set initiation and maintenance (4, 7, 8, 14). Under task-free conditions, both aINS subregions, but perhaps especially the dorsal aINS (8, 15), show activity that is anticorrelated with the "default mode network", a system including the posterior cingulate cortex, inferior parietal lobules, and precuneus (16). In addition to the dorsal-ventral axis, hemispheric lateralization of the aINS is proposed to help maintain bodily homeostasis by adjusting and balancing autonomic outflow based on bioenergetics demands (1, 10, 11). Recent evidence suggests that while the left-sided (dominant hemisphere) aINS controls parasympathetic tone, the homotopic right (non-dominant) aINS is more closely linked to sympathetic tone and responses (1, 10, 11). Structural and functional changes in aINS subregions have been reported in a variety of neuropsychiatric conditions, ranging from mood and anxiety disorders to schizophrenia, autism, and frontotemporal dementia (13, 17–20).

Standard tf-fMRI has helped to reveal the functional organization of the human aINS and other brain areas by providing a snapshot of functional connectivity as averaged across the duration of a scanning session. Although long-range structural connections are assumed to be relatively stable in the adult brain (21), coordinated functional activity is dynamic, with the brain continuously reshaping network configurations in response to prevailing conditions or task demands (22–24). Approaches that capture time-varying connectivity bring the potential to clarify how brain dynamics are organized and relate to function. We reasoned that this approach could help clarify how the aINS may change its network partners in response to salient internal and external stimuli and possibly contribute to the development of more granular and personalized fingerprints of brain function in health and disease. Such time-varying functional connectivity has been shown to relate to clinical outcomes (25–27); task performance (28); and behaviorally relevant measures of cognitive (29–31), emotional (32, 33), and attentional processing (34, 35). One of the few studies applying time-varying functional connectivity analyses to the insula parcellated it into posterior, middle, and dorsal and ventral anterior components, revealing partially overlapping time-varying connectivity profiles for ventral and dorsal aINS subregions. Ventral and dorsal profiles diverged based on distinct contributions from limbic/emotional processing and cingulo-opercular/cognitive regions mirroring findings from static tf-fMRI studies (15).

Despite this body of research, it remains unclear how distinct aINS subregions dynamically engage the salience and task-control networks over time. Moreover, it is largely unknown whether time-varying functional connectivity of the aINS — or any region for that matter — is modulated by externally-driven

states or instead displays trait characteristics such as within-subject temporal stability and relationships
to behavioral or dispositional measures. To gain a deeper understanding of these issues, we implemented
a novel approach combining seed-to-whole-brain sliding-window functional connectivity and k-means
clustering to derive time-varying functional connectivity profiles of aINS subregions across three large,
independent samples of healthy participants, including eyes open vs. eyes closed conditions and
longitudinal datasets. Across methods and samples, we found that aINS subregions display shared and
distinct time-varying connectivity "modes" that bind together cognitive and/or emotional processing
areas versus upstream sensory and motor cortices. Overall, the findings suggest that short-term temporal
variability in aINS connectivity reflects both state and trait characteristics, revealing a path toward use of
such data for assessing psychopharmacological treatment efficacy as well as long-term therapeutic
disease modification in neuropsychiatric conditions.

Materials and Methods

Participants. All participants were healthy, cognitively normal adults, recruited from two different centers
(Table 1, Supplementary Figure S1). Elderly participants were retrospectively selected from the Hillblom
Aging Network, an extensively characterized longitudinal cohort assessed at the University of California,
San Francisco (UCSF) Memory and Aging Center. Subjects were required to have a Clinical Dementia Rating
Scale score (36) of 0 (range 0-3) and a Mini-Mental State Examination score (37) of 28 (range 0-30) or
higher. Secondary inclusion criteria were based on availability of the Interpersonal Reactivity Index, a
widely used questionnaire measuring emotional and cognitive empathy. Out of 224 elderly subjects with
a tf-fMRI scan and at least one Interpersonal Reactivity Index available, a cross-sectional cohort of 121
elderly adults was selected based on availability of the Interpersonal Reactivity Index within three months
of the tf-fMRI scan. A second sample was built based on availability of longitudinal tf-fMRI scans. Out of
total 184 older adults with longitudinally assessed tf-fMRI, 68 were excluded because they overlapped
with the cross-sectional sample (N = 121, described above) and 72 were excluded since they did not meet
our selection criteria of being scanned twice within a 5-13 month interval. This procedure resulted in a
final selection of 44 participants having longitudinal data and that did not overlap with the cross-sectional
sample. All visits included neuropsychological testing and a neurologic exam in addition to a structural
MRI and tf-fMRI scan. Exclusion criteria included a history of drug abuse, psychiatric or neurological
conditions, and current use of psychoactive medications.

Twenty additional younger participants were recruited from a simultaneous FDG-PET/tf-fMRI study at the
Klinikum rechts der Isar, Technische Universität München (38). Neuroimaging data was assessed twice
within an interval of one month, during task-free conditions with either eyes closed or eyes open.
Participants were randomly assigned to one of the two conditions, resulting in eight participants assessed
with eyes closed at the first and with eyes open at the second scan, and 12 participants assessed with
eyes open at the first and eyes closed at the second scan. Exclusion criteria included a history of psychiatric
or neurological conditions, use of psychoactive medications, pregnancy, and renal failure.

Empathy Assessment. The Interpersonal Reactivity Index (39) was completed by the participant's
informant within three months of tf-fMRI scan. This questionnaire is composed by four subscales, each
one consisting of 7 questions that can comprehensively reach a maximum score of 28. While empathic
concern and personal distress measure emotional aspects of empathy, perspective taking and the fantasy
score are designed to assess cognitive aspects of empathy. The Interpersonal Reactivity Index was chosen
since it has been widely used to assess different aspects of empathy in healthy and neuropsychiatric

conditions (39, 40). For example, studies in frontotemporal dementia, a neurodegenerative diseases of
socioemotional dysfunction, have shown that deficits in subscales of the Interpersonal Reactivity Index
are linked to structural deterioration of the aINS and regions connected to the aINS such as the anterior
cingulate and orbitofrontal cortices (40, 41).

Table 1. Demographics and sample characteristics

Dataset	Cross-sectional	Longitudinal	Eyes closed/open
Number of subjects	121	44	20
Age in years, mean (s.d.)	69.3 (3.6)	72.9 (7.6)	42.8 (6.4)
Gender (Female/Male)	73/48	25/19	9/11
Handedness (L/A/R)	0/0/121	6/0/38	0/0/20
Education in years, mean (s.d.)	17.6 (2.0)	17.5 (1.9)	14.4 (2.6)
IRI-EC, mean (s.d.)	27.4 (4.8)	NA	NA
IRI-PT, mean (s.d.)	24.3 (5.7)	NA	NA
IRI-PD, mean (s.d.)	13.1 (5.0)	NA	NA
IRI-FS, mean (s.d.)	18.3 (5.1)	NA	NA
Interval between scanning dates in months, mean (s.d.)	NA	9.2 (2.7)	1.0 (0.2)

A = ambidextrous; EC = empathic concern; FS = fantasy score; IRI = Interpersonal Reactivity Index; L = left;
NA = not applicable; PD = personal distress; PT = perspective taking; R = right; s.d. = standard deviation

Neuroimaging data acquisition. The cross-sectional and longitudinal cohorts from the Hillblom Aging
Network were scanned at the UCSF Neuroscience Imaging Center on a Siemens Trio 3T scanner. A T1-
weighted MP-RAGE structural scan was acquired with acquisition time=8 min 53 sec, sagittal orientation,
a field of view of 160 x 240 x 256 mm with an isotropic voxel resolution of 1 mm^3, TR=2300 ms, TE=2.98
ms, TI=900 ms, flip angle=9˚. Task-free T2*-weighted echoplanar fMRI scans were acquired with an
acquisition time=8 min 6 sec, axial orientation with interleaved ordering, field of view=230 x 230 x 129
mm, matrix size=92 x 92, effective voxel resolution=2.5 x 2.5 x 3.0 mm, TR=2000 ms, TE=27 ms, for a total
of 240 volumes. During the 8-minute tf-fMRI acquisition protocol, participants were asked to close their
eyes and concentrate on their breathing.

Data from the Klinikum rechts der Isar was acquired on an integrated Siemens Biograph scanner capable
of simultaneously acquiring PET and MRI data (3T). FDG-PET activity and tf-fMRI was simultaneously
measured during the initial 10 min immediately after bolus injection of the FDG tracer. Scanning was
performed in a dimmed environment obtained by switching off all lights, including those in the scanner
bore. Subjects were instructed to keep their eyes closed or open, to relax, to not think of anything in

particular, and to not fall asleep. MRI data were acquired using the following scanning parameters: Task-
free echoplanar fMRI scans: TR, 2.000 ms; TE, 30 ms/angle, 90°; 35 slices (gap, 0.6 mm) aligned to AC/PC
covering the whole brain; sFOV, 192 mm; matrix size, 64 x 64; voxel size, 3.0 x 3.0 x 3.0 mm^3 (each
measurement consists of 300 acquisitions in interleaved mode with a total scan time of 10 min and 8 s);
MP-RAGE: TR, 2.300 ms; TE, 2.98 ms; angle, 9°; 160 slices (gap, 0.5 mm) covering the whole brain; FOV,
256 mm; matrix size, 256 x 256; voxel size, 1.0 x 1.0 x 1.0 mm^3; total length of 5 min and 3 s.

Figure 1. Time-varying functional connectivity pipeline. (**A**) For each individual Si, average BOLD activity
time courses were extracted from subregions of the aINS, here the left ventral. (**B**) For each individual,
218 time-varying seed-to-whole-brain connectivity maps (S$_i$W$_1$-S$_i$W$_{218}$) were generated using a sliding-
windows of 18 TRs (36 s) in steps of 1 TR. (**C**) The derived connectivity windows were vectorized and
concatenated across subjects resulting in windows x voxels matrices for each aINS seed. K-means
clustering was then applied to the concatenated window matrices using a value of k = 4 to produce four
clusters representing four time-varying aINS modes present across the course of the functional scan. (**D**)
Group averaged maps of the four modes were generated using the cluster-specific centroid maps
generated through k-means (threshold at -0.5 > z > 0.5). The assignment of each window to a specific
mode, was used to derive fractional occupancy profiles for each mode (i.e. the time spent by an aINS
subregion in each mode), defined as the number of windows assigned to one mode divided through the
total number of generated windows. aINS = anterior insula; BOLD = blood oxygen level dependent signal

Neuroimaging data preprocessing. Before preprocessing, all images were visually inspected for quality
control. Images with excessive motion or image artifact were excluded. T1-weighted images underwent
segmentation using SPM12 (http://www.fil.ion.ucl.ac.uk/spm/software/spm12/). For each tf-fMRI scan,

the first five volumes were discarded. SPM12 and FSL (http://fsl.fmrib.ox.ac.uk/fsl) software were used
for subsequent tf-fMRI preprocessing. The remaining volumes were slice-time corrected, realigned to the
mean functional image and assessed for rotational and translational head motion. Volumes were next co-
registered to the MP-RAGE image, then normalized to the standard MNI-152 healthy adult brain template
using SPM segment, producing MNI-registered volumes with 2 mm^3 isotropic resolution. These volumes
were spatially smoothed with a 6 mm radius Gaussian kernel and temporally bandpass filtered in the
0.008-0.15 Hz frequency range using fslmaths. Nuisance parameters in the preprocessed data were
estimated for the CSF using a mask in the central portion of the lateral ventricles and for the white matter
using a highest probability cortical white matter mask as labeled in the FSL tissue prior mask. Additional
nuisance parameters included the 3 translational and 3 rotational motion parameters, the temporal
derivatives of the previous 8 terms (white matter/CSF/6 motion), and the squares of the previous 16 terms
(42). Subjects were included only if they met all of the following criteria: no inter-frame head translations
greater than 3 mm, no inter-frame head rotations greater than 3 degrees, and less than 24 motion spikes
(defined as inter-frame head displacements > 1 mm), less than 10% of the total number of frames.
Nuisance parameters were regressed out from the filtered data using fslmaths, and masked with a
binarized, skullstripped MNI-152 brain mask. The WM/CSF/head movement denoised data was used for
the subsequent time-varying functional connectivity analysis. Findings with global signal regression are
not presented in the main body of the manuscript but were generated for the longitudinal dataset and
are reported as sensitivity analysis in the Supplement.

Time-varying functional connectivity analysis. For each individual, average blood oxygen level dependent
signal time courses were extracted from the right and left ventral and dorsal aINS using four regions-of-
interest from the Brainnetome Atlas (http://atlas.brainnetome.org/). In order to assess the impact of seed
region selection, in the cross-sectional dataset activity time courses were also extracted from a language-
relevant region centered on the left inferior frontal gyrus (IFG), using a region-of-interest defined in a
previous study (43). Using in-house custom scripts based on Python (https://www.python.org/) and FSL,
a sliding-window approach was implemented to generate time-varying seed-to-whole-brain connectivity
maps (Figure 1A). The derived time series from the aINS and the entire tf-fMRI scan of each subject were
divided into sliding-windows of 18 TRs (36 s) in steps of 1 TR creating 218 (275 for eyes closed/open)
untapered, rectangular windows. At each window, linear regression was used to derive seed-to-whole-
brain time-varying functional connectivity maps for each aINS seed (Figure 1B). A window size of 36
seconds was chosen based on previous research showing that window sizes between 30 and 60 seconds
capture additional variations in functional connectivity not found in larger window sizes (24, 25, 44). Ideal
sliding-window size has been explored by additional methodological work assessing the relationship
between window length and cut-off frequencies, supporting the use of sliding-windows between lengths
of 30–60 seconds for tf-fMRI data preprocessed using a low-pass filter set at 0.15 Hz (45). These studies
are further supported by empirical findings showing that cognitive states can be discerned within such
window lengths (46, 47). Nevertheless, in order to assess the impact of sliding-window length, control
analyses were performed with window lengths of 72 TR (144 s). The resulting findings did not significantly
differ from the reported findings with windows of 36 s (Supplementary Results and Figure S2); therefore
36 s windows were used throughout. The derived time-varying functional connectivity maps were finally
standardized to z-scores (see Video 1 for left ventral aINS time-varying functional connectivity maps of a
typical study participant), masked with a binarized gray matter mask, vectorized, and concatenated across
subjects resulting in windows x voxels matrices for each aINS seed. K-means clustering was then applied
to the concatenated window matrices using a k = 4 to produce four clusters representing four time-varying

functional connectivity modes of aINS subregions present across the course of the functional scan (Figure
1C). The k-means algorithm used Euclidean distance, and an optimal solution was selected after 100
iterations and 10 replications. The optimal number of clusters, referred to henceforward as modes, was
determined using elbow and silhouette plots and by performing additional clustering solutions with k = 3,
5, and 6 (Supplementary Results and Figure S3). With lower k solutions, the identified modes merged,
resulting in information loss. Using higher k solutions resulted in the generation of redundant sub-modes,
exemplified by Mode 4 of the left ventral aINS that would split in anterior and posterior centered
components (Supplementary Figure S3B). A total of three clustering analyses per aINS subregion were
independently performed on time-varying functional connectivity windows: one on the cross-sectional
dataset; one on the longitudinal dataset; and one on the eyes closed/open dataset (see Supplementary
Figure S1 for a schematized summary of k-means clustering analyses performed).

Group-averaged maps of the four identified modes were generated, for each seed region, using the mode-
specific centroid maps rendered using k-means clustering. To assess the distinct spatial contribution of
major large-scale brain networks to time-varying connectivity modes, we used publicly available
templates of major brain networks from a study investigating the functional network organization of the
human brain (48)(https://www.jonathanpower.net/2011-neuron-bigbrain.html). Briefly, in this study
subgraphs corresponding to major brain systems were derived from tf-fMRI data of > 300 healthy adults
by retaining 2% of the strongest correlations. This procedure resulted in 12 binary templates spanning
cognitive, primary sensory and subcortical systems: the salience, default, cingulo-opercular task-control,
executive-control, ventral attention, dorsal attention, auditory, visual, ventral sensorimotor, dorsal
sensorimotor, medial temporal lobe ("memory retrieval") and subcortical networks (4, 48, 49)
(Supplementary Figure S4). Subsequently, for each aINS subregion the averaged z-score value enclosed
within these brain network templates was extracted from the mode-specific centroid maps generated in
the cross-sectional dataset.

Further, the assignment of each window to a specific mode was used to derive: (i) the number of
transitions from one mode to another; (ii) a fractional occupancy metric for each mode, defined as the
number of windows assigned to that mode divided by the total number of windows; and (iii) how often
aINS subregions simultaneously occupy distinct mode configurations in time. To identify meta-profiles of
aINS time-varying functional connectivity, individual subject fractional occupancy profiles, defined as a
vector of 16 elements summarizing the time fraction separately spent by individual aINS subregions on
the four identified modes, were derived from the cross-sectional dataset. Fractional occupancy profiles
were subsequently clustered using k-means with a clustering solution of k = 4 based on a silhouette
analysis (using Euclidean distance, 100 iterations, and 10 replication). On the cross-sectional sample, static
functional connectivity maps were also generated through voxel-wise regression analyses of each aINS
seed's time-course for the duration of the entire scan, following previous established methods (50, 51).

Statistical analysis. Statistical analyses were carried out using R (https://www.r-project.org/) and Matlab-
R2018 (https://www.mathworks.com/products/matlab.html). Pearson's correlation was used to assess
the spatial similarity of modes derived from different clustering analyses, using maps thresholded for z-
values higher than 0.3 and lower than -0.3.

In the cross-sectional dataset, the Shannon diversity index was used to analyze whether aINS subregions
of an individual subject transitioned repeatedly between distinct modes or showed instead a stereotypic
behavior spending most of the time only in a specific aINS mode:

$$\text{Shannon diversity index} = -\sum_{i=0}^{n-1} p_i log_2 p_i$$

wherein *p* is the fractional occupancy of an aINS subregion in a specific time-varying functional
connectivity mode. A Shannon diversity index of 1.4 indicates that all modes are equally occupied by the
aINS, while the lower the Shannon diversity index, the higher the fractional occupancy of the aINS on one
specific mode.

In the cross-sectional dataset, one-way ANOVAs were used to investigate whether aINS subregions
differed, within subjects, in terms of mode-specific fractional occupancy, number of transitions, and
Shannon diversity index. For descriptive purposes, averages and standard deviations derived over all aINS
subregions and modes are reported. Additionally, for each meta-profile cluster identified in the cross-
sectional dataset, one-way ANOVAs were used to assess differences in mode-specific fractional occupancy
averaged across aINS subregions. The assignment of participant to four meta-profiles of time-varying aINS
fractional occupancy was additionally used to test differences in aINS subregions' static connectivity using
ANOVA models and post-hoc t-tests implemented in SPM12 (height threshold p<0.005; extent threshold
p<0.05 FWE corrected for multiple comparisons).

Finally, in the cross-sectional dataset, four multiple linear regression models were used to test the
association of mode-specific fractional occupancies with subscales of the Interpersonal Reactivity Index
(39). Each model contained the mode-specific fractional occupancies averaged across aINS subregions,
and was corrected for age, gender, and sum frame-wise head displacement (p < 0.05 uncorrected for
multiple comparisons). To assess specificity of the aINS results, four additional models were performed,
testing the association between subscales of the Interpersonal Reactivity Index and fractional occupancy
of time-varying functional connectivity modes derived from the left IFG.

For the longitudinal datasets, cosine similarity was used to assess the similarity of fractional occupancy
profiles across distinct scanning dates.

$$\text{Cosine similarity} = 1 - \left(\frac{\sum_{i=1}^{n} A_i B_i}{\sqrt{\sum_{i=1}^{n} A_i^2} \sqrt{\sum_{i=1}^{n} B_i^2}} \right)$$

Where *A* and *B* are the vectorized fractional occupancy profiles of all aINS subregions at two different
scanning dates. Cosine similarity varies between 1 and 0, with 1 indicating identical fractional occupancy
profiles. Paired t-tests were performed to assess differences in fractional occupancy in the longitudinal
and in the eyes closed/open datasets. All statistical findings are reported at p < 0.05, two-tailed,
uncorrected for multiple comparisons except where specified otherwise.

Results

Bilateral ventral and dorsal aINS occupy four overlapping but distinct time-varying functional connectivity
modes

By combining seed-to-whole-brain functional connectivity, sliding-window analysis, and k-means
clustering, we found that aINS subregions adopt four overlapping but distinct large-scale configurations
or "modes" of time-varying functional connectivity (Figure 2A). All modes were characterized by bilateral
connectivity between the aINS and anterior cingulate cortices. Modes could be distinguished from each
other, however, by specific patterns of connectivity to other brain regions (Figure 2A and Supplementary

Table S1). In Mode 1, all four aINS subregions showed connectivity to the anterior midcingulate/pre-
supplementary motor area, right frontal operculum, and dorsal parietal and dorsolateral prefrontal areas
that together make up the task-control network. Mode 2 was characterized by negative connectivity to
primary visual and sensorimotor areas and prominent connectivity to the ventral striatum and thalamus.
Mode 3 showed inverted connectivity patterns to the same regions. Finally, Mode 4 showed connectivity
patterns aligned with more ventral salience network regions, including pre- and sub-genual anterior
cingulate and orbitofrontal cortices, with additional connectivity to the temporal poles for the right
ventral aINS. All four modes were identified in each aINS subregion, as shown by the correlation matrix
highlighting the spatial correspondence of equivalent modes (Figure 2B). Important distinctions were
found, however, when comparing dorsal and ventral aINS connectivity patterns within modes. In Mode 1,
the dorsal aINS showed prominent anti-correlations to the precuneus, angular gyrus, and medial
prefrontal cortex regions that make up the default mode network (16). In Mode 4, the ventral aINS
showed a more ventral connectivity pattern that encompassed subgenual anterior cingulate cortex and
temporal poles, when compared to the dorsal aINS (Figure 2C, Supplementary Figure S5A). Modes derived
from homologous left and right aINS regions differed from each other mainly with regard to the extent of
ipsilateral connectivity to neighboring regions (Supplementary Figure S5B-C). Importantly, similar aINS
time-varying functional connectivity modes were identified using longer sliding-window lengths
(Supplementary Figure S2), lower and higher clustering solutions (Supplementary Figure S3), and data
preprocessed using global signal regression (Supplementary Figure S6).

aINS subregions coherently transition between time-varying functional connectivity modes

Average fractional occupancy, i.e. the proportion of time spent by aINS subregions in the four time-varying
functional connectivity modes ranged from 0.18 (left ventral aINS in Mode 2) to 0.30 (right ventral aINS
in Mode 1), with an overall average of 0.25 +/- 0.20 (Figure 3A). Importantly, no significant differences in
mode-specific fractional occupancy were found across the four aINS subregions (Mode 1: F = 0.2, p = 0.82;
Mode 2: F = 0.7, p = 0.56; Mode 3: F = 0.4, p = 0.77; Mode 4: F = 0.5, p = 0.71). In only a few participants
did aINS subregions spend disproportionate time in only one mode. The number of transitions was also
comparable across aINS subregions (F = 0.4, p = 0.759; Figure 3B), with an overall average of 18.0 +/- 6.6
transitions over the task-free acquisition. In particular, our data shows that there were no direct
transitions between the primary sensory anticorrelated Mode 2 and the primary sensory correlated Mode
3 (Supplementary Figure S7), suggesting that aINS subregions need to transition to modes rooted in
cognitive networks before occupying primary sensory-centered modes characterized by opposing
connectivity patterns. The Shannon diversity index was used to quantify the diversity of fractional
occupancy and to compare this diversity across subregions. On average, aINS subregions showed high
Shannon diversity, suggesting that each subregion moved between modes in most subjects, rather than
stacking on a single mode. The overall average Shannon diversity index of fractional occupancy across
aINS subregions was 1.0 +/- 0.2 and did not significantly differ between subregions (F = 0.8; p = 0.503;
Figure 3C). Subject-level fractional occupancy profiles are illustrated in Figure 3D, which shows two
subjects with contrasting signatures. We finally sought to assess whether aINS subregions coherently
occupy the same modes at the same time, or for instance, how often the right ventral and dorsal aINS are
simultaneously connected to Mode 1. For each subject and each aINS subregion, we assessed how often
possible mode configurations were jointly occupied in time. This analysis revealed a tendency across aINS
subregions to occupy the same modes in time (Figure 3E). On average, the right ventral and dorsal aINS
occupied the same modes 50% of the time, the left ventral and dorsal aINS 53% of the time, the right and

left ventral aINS in 51% of the time, and the right and left dorsal aINS 61% of the time. These findings
suggest an overall tendency for aINS subregions to cohesively engage the same modes over time under
task-free scanning conditions.

Figure 2. Time-varying functional connectivity modes of the aINS. (A) Spatial maps of the four time-
varying functional connectivity modes identified across the four aINS subregions in the cross-sectional
dataset. Threshold at -0.5 > z > 0.5, negative connectivity is depicted in blue, positive in red. The right side
of the brain is shown on the right side of the image. **(B)** Correlation matrix reflecting the spatial similarity
of modes identified in distinct aINS subregions. **(C)** Differences in time-varying connectivity between
correspondent modes of the right dorsal and ventral aINS (threshold at -0.3 > z > 0.3, higher connectivity
of the dorsal aINS is depicted in red, higher connectivity of the ventral aINS in blue).

Individual aINS connectivity mode occupancy profiles are reproducible across samples and within subjects
over time

To assess the generalizability and reproducibility of our findings, we next turned to a longitudinal healthy
aging dataset, which consisted of 44 participants who were scanned twice over an interval between 5-13
months. Clustering of time-varying connectivity windows identified modes that strongly resembled the
four modes derived from the larger (and non-overlapping) cross-sectional dataset (Figure 4A-D). Average
fractional occupancies of aINS subregions in the four modes were comparable with the cross-sectional
sample (comparing Figure 3A to 4A-D). Across aINS subregions and modes, paired t-tests revealed no
significant differences in fractional occupancy when comparing the first scanning date with the second (p
< 0.05 uncorrected for multiple comparisons, for details on paired t-test statistics see Supplementary
Table S2). Within subjects, cosine similarity analysis showed that most participants had relatively stable
fractional occupancy profiles over time (mean cosine similarity = 0.70 +/- 0.19 s.d.), as further supported
by the inverted bell-shaped distribution of cosine similarity (Figure 5A). Stable fractional occupancies are
exemplified in Subjects 10 (first scanning date in red; second scanning date in blue; cosine similarity =
0.98) and 41 (cosine similarity = 0.92) (Figure 5B-C). A few participants, however, showed very different
fractional occupancy profiles when comparing the two scanning dates, as exemplified by Subjects 21
(cosine similarity = 0.46) and 26 (cosine similarity = 0.24) (Figure 5D-E). Overall, however, the temporal
stability of the individual profiles suggests a major contribution from trait-level factors.

Eye opening decreases the time aINS spends anticorrelated with visual cortices

Independently performed clustering of time-varying connectivity data from the eyes closed/open dataset
revealed similar aINS modes to those identified in the cross-sectional and longitudinal aging datasets
(Figure 6A-B). For all subregions except the right ventral aINS, participants showed higher fractional
occupancies in the anti-correlated primary sensory/motor Mode 2 with eyes closed and higher fractional
occupancies in the task-control Mode 1 with eyes open (p < 0.05 uncorrected, for details on paired t-test
statistics see Supplementary Table S2).

Time-varying functional connectivity meta-profiles show that aINS subregions cohesively occupy the same
modes

Individual subjects were clustered into meta-profiles (see Methods "*Time-varying functional connectivity*
analysis"), based on the tendency of their aINS subregions to occupy specific time-varying functional
connectivity modes. Individual fractional occupancy profiles of the cross-sectional sample (n = 121) were
clustered into four clusters using k-means. Silhouette plots were used to choose the ideal number of
clusters, k = 4 (Supplementary Figure S8). This analysis revealed four meta-profiles, characterized by
preferred fractional occupancy in one of the four time-varying functional connectivity modes (Figure 7A).
Meta-profile 1 (in green, 26 subjects) was characterized by a tendency to spend time in the task-control
Mode 1 (see Subject 22, Figure 7A); Meta-profile 2 (in blue, 31 subjects) was characterized by a tendency
to spend time in the sensory anticorrelated Mode 2 (see Subject 79, Figure 7A); Meta-profile 3 (in violet,
44 subjects) was characterized by a tendency to spend time in the sensory connected Mode 3 (see Subject
80, Figure 7A); while Meta-profile 4 (in red, 20 subjects) was characterized by a tendency to spend time
in the salience network Mode 4 (see Subject 88, Figure 7A). To evaluate statistical differences in mode
occupancy across meta-profiles, we averaged the time spent in each mode across aINS subregions and
used one-way ANOVAs to test for significant group differences in mode-specific fractional occupancy
across the distinct aINS meta-profiles (Meta-profiles 1, F = 50.1, p < 0.0001; Meta-profile 2, F = 39.8, p <
0.0001; Meta-profile 3, F = 79.6, p < 0.0001; Meta-profile 4, F = 46.1, p < 0.0001, see also Figure 7B). These
differences were further statistically assessed via post-hoc t-tests. In summary, these analyses revealed a

Figure 3. Temporal characteristics of the identified aINS time varying functional connectivity modes. (A) Violin plots reflecting group fractional occupancy in the four identified time-varying functional connectivity modes across the four aINS subregions. **(B)** Violin plots representing the number of transitions between time-varying functional connectivity modes across the four aINS subregions. **(C)** Violin plots representing the diversity of fractional occupancy profiles – measured using the Shannon diversity index – across aINS subregions. This measure reflects whether an aINS subregion tends to spend similar amounts of time in different modes or spends most of its time in only one mode. **(D)** Polar plots schematizing the fractional occupancies of aINS subregions into the four time-varying functional connectivity modes. On the left side, the fractional occupancy profile of Subject 47 transitions between several modes, and is characterized by a high number of transitions and by high Shannon diversity index averaged across aINS subregions. On the right side, we can appreciate the highly stereotyped fractional occupancy profile of Subject 22, with aINS subregions spending most of their time in a single mode. Regardless of the subregion, aINS subregions in this participant tend to spend time only in Mode 1, showing hence a low number of transitions and low Shannon diversity index averaged across aINS subregions. **(E)** Heat maps reflecting how specific modes are jointly occupied at the same time by distinct aINS subregions. Scale bar reflects the average group probability that distinct aINS subregions occupy certain mode configurations.

preference for aINS subregions to collectively spend similar fractions of time in a given mode, whereas distinct modes were primarily occupied by subjects in different meta-profiles.

Next, we asked whether the constituents of a given meta-profile would show group-level differences in terms of their static functional connectivity patterns. We therefore compared groups based on static functional connectivity of aINS subregions across the four meta-profiles described above. This analysis revealed that participants in Meta-profile 1 showed prominent static anticorrelation of the dorsal aINS to the default mode network when compared to participants clustered on other meta-profiles. Participants clustered on Meta-profile 2 and 3 showed marked visual and sensorimotor cortex static hypoconnectivity and hyperconnectivity, across all aINS subregions, when compared to participants of other meta-profiles; while participants in Meta-profile 4 showed prominent static connectivity of all aINS subregions to more ventral frontal regions (Supplementary Figure S9). In summary, key components of the time-varying connectivity modes were reflected in static connectivity differences between meta-profile-based groups.

Comparison of aINS and left IFG time-varying functional connectivity

To control for the location of the seed, we derived time-varying functional connectivity modes by seeding the left IFG. Time-varying functional connectivity modes of the left IFG shared some common themes with modes identified using aINS subregions (Supplementary Figure S10A). The four identified modes showed similar connectivity to regions of the language network, such as the left IFG, pre-supplementary motor area, and superior parietal lobule (43, 52). This contrasts with the modes identified in aINS subregions that showed overlapping time-varying functional connectivity to typical task-control and salience network regions (Supplementary Figure S10B). Modes derived from both the aINS and left IFG, however, showed similar patterns of default mode network anticorrelation (Mode 1), primary sensory anticorrelation (Mode 2), and primary sensory hypercorrelation (Mode 3). A fourth mode was identified using the left IFG, showing time-varying functional connectivity to regions of the dorsal attention network (49) (Mode 4).

Figure 4. Time-varying functional connectivity modes across time. In the longitudinal data sample, spatial patterns of time-varying functional connectivity modes where identified across the four aINS subregions bearing high similarity to the modes identified in the cross-sectional sample (maps averaged across both time points, sagittal plane only panels **A-D**). Fractional occupancy in the identified modes did not significantly differ between the first (red violin plots) and second scanning dates (blue violin plots) ~9 months apart (paired t-test; p < 0.05 uncorrected for multiple comparisons).

Time spent by the aINS in the task-control and salience networks correlates with metrics related to dispositional empathy

A key question of this work concerns whether individual differences in aINS time-varying connectivity relate to differences in aINS-associated traits and functions. Among the many tasks in which the anterior insula participates, empathy is one of the best documented (1, 18). The Interpersonal Reactivity Index is an informant-based questionnaire widely used to assess distinct aspects of empathy (39). Emotional aspects of empathy are covered by the empathic concern and persona distress measures, with the first assessing an other-centered emotional response, while the second reflects general anxiety and self-oriented emotional reactivity. Perspective taking and fantasy scores measure cognitive aspects of empathy, with the first assessing the tendency to spontaneously imagine the cognitive perspective of another person, while the second measures the tendency to project oneself into the experiences of fictional characters. Mode-specific fractional occupancies were averaged across aINS subregions and associated with the four subscales of the Interpersonal Reactivity Index. Multiple linear regression models corrected for age, gender, and sum frame-wise head displacement identified two significant positive associations. First, fractional occupancy in the task-control Mode 1 predicted higher levels of personal

distress, reflecting self-oriented feelings of personal anxiety and unease in tense interpersonal settings (β
= 7.1; p < 0.05 uncorrected for multiple comparisons; Figure 8A). Second, fractional occupancy in the
salience Mode 4 correlated with higher scores on the fantasy subscale, reflecting greater ability to
transpose one's self imaginatively into the feelings and actions of fictitious characters in books, movies,
and plays (β = 9.2; p < 0.05 uncorrected for multiple comparisons; Figure 8B and Supplementary Table S3).
These results remained significant when removing outliers (Mode 1 and personal distress β = 6.9, p < 0.05
uncorrected for multiple comparisons; Mode 4 and fantasy score β = 9.2, p < 0.05 uncorrected for multiple
comparisons). No significant associations were found between fractional occupancy in any mode and the
empathic concern or perspective taking subscales. As hypothesized, fractional occupancy of left IFG time-
varying connectivity modes was not significantly associated with any measure of dispositional empathy (p
< 0.05 uncorrected for multiple comparisons, see Supplementary Table S3).

Figure 5. Individual fractional occupancy of aINS subregions across time. (A) Violin plot representing the
similarity of aINS fractional occupancy profiles across scanning dates assessed using cosine similarity.
Subjects 10, 41, 21 and 26 are highlighted with red circles in the individual data plot. The individual
fractional occupancy profiles of these subjects and correspondent cosine similarity values are further
schematized in panels **B-E**, where the fractional occupancies are shown at the first (red) and second (blue)
scanning dates. aINS fractional occupancy in Subjects 10 and 41 did not substantially differ between
scanning dates (note the overlap in the polar plots and the high cosine similarity index), while Subjects 21

and 26 showed very different aINS fractional occupancy profiles when assessed at different time points
(note the little overlap in the polar plot and the low cosine similarity index).

Figure 6. Time-varying functional connectivity modes during scanning with eyes closed versus eyes
open. In the eyes closed/open dataset, spatial patterns of time-varying functional connectivity modes
where identified across the left dorsal, right dorsal, left ventral, and right ventral aINS bearing high
similarity to the modes identified in the cross-sectional sample (maps averaged across eyes closed/open
conditions, sagittal plane only panels **A-D**). With exception of the right ventral aINS, fractional occupancy
in the visually anticorrelated Mode 2 was significantly higher during eyes closed (red violin plots) than
during eyes open scanning (blue violin plots). On the other hand, in the eyes open condition participants
spent significantly more time in the task-control Mode 1 (paired t-test; $p < 0.05$ uncorrected for multiple
comparisons). *$p < 0.05$; **$p < 0.005$

Discussion

To date, most efforts to relate large-scale networks to individual differences have relied on metrics that
capture the topology or strength of node-to-node static functional (or "intrinsic") connections. Here, we
developed a novel approach combining seed-to-whole-brain functional connectivity, sliding-window
analysis, and k-means clustering on tf-fMRI data to identify distinct time-varying functional connectivity
modes of the aINS, a brain region critical for many human socio-emotional functions (1, 18). These robust
modes were identified across methods and samples, showing both "state" and "trait" characteristics. In
particular, while aINS modes related to sensory processing were modulated by visual input (eyes closed

vs. open conditions during scanning), modes featuring connectivity to cognitive and emotion processing
regions were stable over time and related to measures of dispositional empathy.

Partially overlapping but distinct time-varying aINS functional connectivity modes

Our pipeline reliably identified four time-varying functional connectivity modes of the aINS across
different subregions, methods, and samples. These modes were characterized by common connectivity
of aINS subregions to the contralateral insula and to the anterior cingulate cortex, but could be
differentiated from each other based on recruitment of other brain regions. Mode 1 was characterized
by a more dorsal connectivity pattern that resembles previous characterizations of a cingulo-opercular
task-control network (4, 14, 53) involving the bilateral aINS and anterior cingulate cortex, with additional
connectivity to dorsomedial parietal and dorsolateral frontal areas. Mode 2 was characterized by negative
connectivity to primary sensory areas and increased connectivity to the thalamus while Mode 3 showed
an inverted connectivity pattern to these regions. Finally, Mode 4 was characterized by fronto-insular
connectivity to more rostral and ventral portions of the anterior cingulate cortex, resembling the salience
network (4, 8), with additional connectivity to the temporal pole and medial orbitofrontal cortex,
components of the "semantic-appraisal network" (18, 54). These findings are consistent with those of a
previous region-of-interest-based time-varying functional connectivity study focusing on the insula, which
reported time-varying visual and sensorimotor hyperconnectivity of the ventral aINS. Our findings differ,
however, in that we did not detect positive default mode network connectivity to the dorsal aINS and that
patterns of hypoconnectivity to primary visual and sensorimotor areas were less apparent in the
aforementioned study (15). While left and right aINS subregions primarily differed with respect to
connectivity to regions in the ipsilateral hemisphere, consistent differences in time-varying functional
connectivity at the system-level were identified when comparing ventral to dorsal aINS subregions.
Mirroring static functional connectivity studies and the aforementioned time-varying connectivity study
on the insula (7, 8, 15), the task-control Mode 1 of the dorsal aINS showed stronger anti-correlations to
the default mode network than did its ventral counterpart. This observation is in line with salience
network models proposing that the dorsal aINS receives ventral aINS input streams regarding the
moment-to-moment condition of the body and then, based on these inputs, recruits task control (14) and
executive control (4) network resources to maintain cognitive set and guide behavior while inhibiting the
default mode network (13). The salience network Mode 4 showed stronger connectivity between ventral
aINS and ventromedial prefrontal, orbitofrontal, and temporopolar cortices when compared to its dorsal
aINS counterpart. This finding is in line with functional-anatomical models suggesting a close alliance
between the salience and the semantic appraisal networks (13, 18, 54), whose main components (the
temporal pole, ventral striatum, medial orbitofrontal cortex, and amygdala) have been proposed to
interact with autonomic representations in the aINS to construct the meaning and significance of social
and non-social stimuli under prevailing conditions (13, 18, 54). Importantly, we did not find any
differences across aINS subregions in the number of transitions or in the diversity of fractional occupancy
profiles. Moreover, our analyses exploring overlapping mode occupancy across aINS subregions revealed
that distinct aINS subregions tend to occupy the same mode at the same time. Further support for this
conclusion comes from the clustering analysis of individual fractional occupancy profiles, which led to the
identification of four meta-profiles characterized by occupancy of a dominant mode.

Figure 7. Meta-profiles of time-varying dynamic connectivity modes. (A) Individual fractional occupancy profiles of aINS subregions were clustered into four clusters, reflecting four meta-profiles of time-varying aINS functional connectivity. 26 subjects were clustered into Meta-profile 1 (triangles, green), 31 into Meta-profile 2 (+ sign, blue), 44 into Meta-profile 3 (circles, violet), and 20 into Meta-profile 4 (x's, red). Panels show individual fractional occupancy profiles representative of the identified meta-profiles (see black circles in the scatterplots). Across aINS subregions, Subject 22 tended to spend time in Mode 1, Subject 79 in Mode 2, Subject 80 in Mode 3, and Subject 88 in Mode 4. **(B)** When averaged across aINS subregions, one-way ANOVAs and related post-hoc t-tests reveal that the aINS of participants clustered in Meta-profile 1 showed significantly higher fractional occupancy in Mode 1. Similarly, the aINS of participants clustered on Meta-profiles 2, 3, or 4, spent more time on Modes 2, 3, or 4, respectively.

Although our findings are not conclusive on whether aINS subregions always engage the same mode in
time, they comprehensively suggest a tendency for coherent mode occupancy that could potentially
underpin specific trait characteristics or behavioral states. Future studies could leverage methods such as
directed and effective connectivity to assess the temporal relationships between intrinsic time-varying
activity of distinct aINS subregions, which could be lagged, phase-shifted, or anticorrelated over time. In
order to assess the specificity of aINS time-varying functional connectivity modes, we assessed and
compared time-varying functional connectivity of the left IFG, a region involved in language processing
and syntax production (43, 52, 55). The left IFG showed region-specific contributions from language
processing regions but also displayed pronounced anticorrelation to default mode network regions in
Mode 1 and similar time-varying connectivity to the same sensory regions found in Modes 2 and 3 derived
from the aINS. These findings suggest that distinct brain regions, although characterized by seed specific
connectivity patterns, may cohesively transition between broader modes characterized by shared regional
time-varying functional connectivity patterns, possibly revealing a fundamental property of functional
brain organization.

"State" characteristics of aINS time-varying functional connectivity modes

Having determined that two of the four major modes of aINS connectivity involved varying connectivity
to sensory, and, most notably, visual cortices, we chose to evaluate how sensory input modifies aINS
connectivity dynamics. To this end, we studied the same participants under eyes closed and eyes open
scanning conditions. Compared to the eyes open condition, participants with eyes closed spent
significantly less time in the task-control network Mode 1 but more time in the primary sensory anti-
correlated Mode 2. This finding is consistent with recent static connectivity studies revealing occipital and
sensorimotor functional changes in eyes closed versus open scanning conditions (56, 57) and suggests
that environmental conditions impact time-varying aINS connectivity by shifting the major modes
occupied. Therefore, a measurable component of time-varying connectivity may reflect external cues such
as conditions of eyes closure (38, 58), mental states (59, 60), or internally driven physiological conditions
of the body (12). Finally, it is tempting to speculate that aINS occupancy of the visually anticorrelated
Mode 2 might reflect levels of drowsiness and fluctuations in alertness. Intriguingly, a recent study
investigated wakefulness fluctuations as a source of time-varying functional connectivity by combining
simultaneously acquired tf-fMRI and EEG data (61). This study revealed progressive whole-brain
hypoconnectivity during deeper sleep stages (N2 and N3) when compared to wakefulness, raising the
possibility that the increased aINS fractional occupancy in Mode 2 during eyes closed is related to reduced
levels of alertness and wakefulness.

"Trait" characteristics of aINS time-varying functional connectivity modes

To the extent that modes and profiles of time-varying aINS functional connectivity represent traits,
individuals should show stability of these features over time. To assess this stability, we studied a
longitudinal dataset of cognitively healthy older adults. The four aINS modes were stable over an average
of 9 months in this sample, whether assessed by comparing fractional occupancy at the group level via
parametric tests or at the individual level via cosine similarity. These findings converge with several recent
studies showing that individual topologies in static functional brain organization are highly stable across
time and show unique features with promise for use in precision functional mapping of individual human
brains (62–65). Adding to previous work (15), we found that the time aINS subregions cohesively spent in
the task-control Mode 1 was positively associated with personal distress, while the time aINS subregions

spent in the salience Mode 4 was positively associated with the fantasy score from the Interpersonal
Reactivity Index.

Figure 8 Time-varying fractional occupancy of the aINS correlates with dispositional empathy. (A)
Fractional occupancy in the task-control Mode 1 averaged across aINS subregions showed a significant
positive association with the personal distress subscale from the Interpersonal Reactivity Index. This result
was stable also after removing one outlier circled in black in the scatterplot (β = 6.9; p < 0.05). **(B)**
Fractional occupancy in the salience Mode 4 averaged across aINS subregions showed a significant
positive association with the fantasy score subscale from the Interpersonal reactivity Index. This result
was stable also after removing one outlier outlined in black in the scatterplot (β = 9.2; p < 0.05). Multiple
linear regression models were corrected for age, gender and sum frame-wise head displacement. *p <
0.05, uncorrected for multiple comparisons.

Although the statistical association was moderate and did not survive multiple comparisons correction,
the identified associations add tentative support for the trait characteristics of aINS time-varying
functional connectivity. The personal distress subscale measures "self-oriented" feelings of personal
anxiety and unease in tense interpersonal settings. The level of personal distress was associated with
greater time spent in a mode characterized by dorsal cingulo-opercular regions overlapping with the task-
control network and anti-correlations to the default mode network, in line with findings associating
connectivity of the anterior cingulate cortex and insula with pre-scan anxiety levels, dispositional anxiety,
and affective symptoms in mood disorders (4, 20, 66). The fantasy score reflects the capacity of
participants to transpose themselves imaginatively into the feelings and actions of fictitious characters in
books, movies, and plays. This capacity requires a high level of social contextualization and involves
semantic processes typically associated with regions of the semantic appraisal network, such as the
temporal lobes and ventro-medial prefrontal areas (13, 18, 54), the same regions contributing to the
salience Mode 4. Importantly, fractional occupancies of time-varying functional connectivity modes
derived from the left IFG were not significantly associated with measures of dispositional empathy,
suggesting a degree of specificity of these findings to the aINS.

Limitations and future directions

Although time-varying functional connectivity analyses have been of increasing interest to the human
brain mapping community, consensus is still lacking about whether time-varying fluctuations in BOLD
signal coherence reflect brain physiology or are mainly due to noise stochastically occurring in the scanner
between the acquisition of distinct brain volumes (67). The sliding-window approach has been scrutinized
because of its moderate reliability (68) and since its output is prone to influence by head movement,
sliding-window length, and other methodical choices (69–71), such as global signal regression. In our
study, however, we did not find any significant association between time spent by distinct aINS subregions
in a specific mode and summary measures of head movement, as exemplified by the regression analyses
associating fractional occupancy and dispositional empathy, which were corrected for mean frame-wise
head displacement (42, 72). Importantly, our control analyses using the longitudinal dataset preprocessed
with global signal regression resulted in the identification of time-varying functional connectivity modes
that highly resembled the time-varying modes derived using data without global signal regression
(Supplementary results and Figure S6). The only exception was for the primary sensory hyperconnected
Mode 3, which was not identified in three out of four aINS subregions, suggesting that this mode is highly
modulated by the global signal. A recent report suggested that noise sources (for instance motion and
respiration) can have a temporally lagged effect on the BOLD signal, which is greatly reduced by global
signal regression (70). However, a recent study investigating time-varying functional connectivity in mice
reliably mapped distinct time-varying patterns of tf-fMRI co-activation that occurred at specific phases of
global signal fluctuations (73). These findings suggest that the global signal may partially relate to
oscillatory cycling of slowly propagating neural activity, as observed in lag threads of propagated tf-fMRI
signal in the human brain (74) and transient variation in calcium co-activation patterns in mice (75). Slow
oscillation in the global signal have been proposed to coordinate fluctuating periods of brain network
topologies characterized by heightened global integration and shifts in arousal (30, 76).

Finally, the limitations in temporal resolution affecting the sliding-window approach may also explain the
inability to detect mode-specific fractional occupancy differences between the left and right aINS,
although both regions play a distinct role in autonomic outflow and behavior (11, 12). More sophisticated
methods that preserve the temporal richness of tf-fMRI data, such as hidden Markov models and
Topological Data Analysis (28, 31), may help elucidate lateralized differences in time-varying connectivity
of the left and right aINS. However, the detection of static functional connectivity heterogeneity informed
by time-varying fractional occupancy profiles and the reproducibility of our findings across distinct
preprocessing pipelines, sliding-window lengths, clustering choices, samples, and time points (within
individuals)(77) increase confidence in the biological relevance of findings. In the future, our approach
could be extended to analyze time-varying functional connectivity of other brain regions, or to assess
time-varying fractional occupancy profiles in psychiatric and neurological disorders characterized by aINS
dysfunction.

Acknowledgements

We thank the participants for their contributions to neuroscience research.

Conflicts of interest

The authors declare no competing conflicts of interest.

Funding

This work was supported by the Larry L. Hillblom Foundation grants 2014-A-004-NET and 2018-A-025-FEL,

The Bluefield Project to Cure FTD, the Tau Consortium, and NIH grants R01AG032289, R01AG048234, and

UCSF ADRC P50 AG023501.

References

1.   Craig AD (2009) How do you feel - now? The anterior insula and human awareness. *Nat Rev*

Neurosci 10(1):59–70.

2.   Smith SM, et al. (2009) Correspondence of the brain's functional architecture during activation

and rest. *Proc Natl Acad Sci U S A* 106(31):13040–5.

3.   Fox MD, et al. (2005) The human brain is intrinsically organized into dynamic , anticorrelated

functional networks.

4.   Seeley WW, et al. (2007) Dissociable intrinsic connectivity networks for salience processing and

executive control. *J Neurosci* 27(9):2349–2356.

5.   Deen B, Pitskel NB, Pelphrey KA (2011) Three systems of insular functional connectivity identified

with cluster analysis. *Cereb Cortex* 21(7):1498–1506.

6.   Kurth F, Zilles K, Fox PT, Laird AR, Eickhoff SB (2010) A link between the systems: functional

differentiation and integration within the human insula revealed by meta-analysis. *Brain Struct*

Funct 214:1–16.

7.   Touroutoglou A, Hollenbeck M, Dickerson BC, Feldman Barrett L (2012) Dissociable large-scale

networks anchored in the right anterior insula subserve affective experience and attention.

Neuroimage 60(4):1947–1958.

8.   Uddin LQ (2014) Salience processing and insular cortical function and dysfunction. *Nat Rev*

Neurosci 16(1):55–61.

9.   Critchley HD (2005) Neural mechanisms of autonomic, affective, and cognitive integration. *J*

Comp Neurol 493(1):154–166.

10.   Critchley HD, Harrison NA (2013) Visceral Influences on Brain and Behavior. *Neuron* 77(4):624–

638.

11.   Sturm VE, et al. (2018) Network architecture underlying basal autonomic outflow: Evidence from

frontotemporal dementia. *J Neurosci* (415):0347–18.

12.   Guo CC, et al. (2016) Dominant hemisphere lateralization of cortical parasympathetic control as

revealed by frontotemporal dementia. *Proc Natl Acad Sci*:201509184.

13.   Zhou J, Seeley WW (2014) Network Dysfunction in Alzheimer ' s Disease and Frontotemporal

Dementia : Implications for Psychiatry. *Biol Psychiatry* 75(7):565–573.

14.   Dosenbach NUF, et al. (2006) A Core System for the Implementation of Task Sets. *Neuron*

50(5):799–812.

15.   Nomi JS, et al. (2016) Dynamic functional network connectivity reveals unique and overlapping

profiles of insula subdivisions. *Hum Brain Mapp* 37(5):1770–1787.

16.   Buckner RL, Andrews-Hanna JR, Schacter DL (2008) The brain's default network: Anatomy,

function, and relevance to disease. *Ann N Y Acad Sci* 1124:1–38.

17.   Menon V, Uddin LQ (2010) Saliency, switching, attention and control: a network model of insula

function. *Brain Struct Funct*:1–13.

18.   Seeley WW, Zhou J, Kim E-J (2012) Frontotemporal Dementia: What Can the Behavioral Variant

Teach Us about Human Brain Organization? *Neurosci* 18(4):373–385.

19.   Brandl F, et al. (2019) Specific Substantial Dysconnectivity in Schizophrenia: A Transdiagnostic

Multimodal Meta-analysis of Resting-State Functional and Structural Magnetic Resonance

Imaging Studies. *Biol Psychiatry* (10):1–11.

20.   Williams LM (2016) Precision psychiatry: A neural circuit taxonomy for depression and anxiety.

The Lancet Psychiatry 3(5):472–480.

21.   Sporns O, Tononi G, Kötter R (2005) The Human Connectome: A Structural Description of the

Human Brain. *PLoS Comput Biol* 1(4):e42.

22.   Lurie DJ, et al. (2018) On the nature of resting fMRI and time-varying connectivity. *Preprint*:1–33.

23.   Preti MG, Bolton TA, Van De Ville D (2017) The dynamic functional connectome: State-of-the-art

and perspectives. *Neuroimage* 160:41–54.

24.   Allen E, Damaraju E, Plis S, Erhardt E (2014) Tracking whole-brain connectivity dynamics in the

resting state. *Cerebral. Cortex* 24(3):663-76.

25.   Damaraju E, et al. (2014) Dynamic functional connectivity analysis reveals transient states of

dysconnectivity in schizophrenia. *NeuroImage Clin* 5:298–308.

26.   Kaiser RH, et al. (2016) Dynamic Resting-State Functional Connectivity in Major Depression.

Neuropsychopharmacology 41(7):1822–1830.

27.   Sourty M, Thoraval L, Roquet D (2016) Identifying dynamic functional connectivity changes in

dementia with lewy bodies based on product hidden markov models. *Front Comput Neurosc*

10:60*.*

28.   Saggar M, Sporns O, Gonzalez-castillo J, Bandettini PA, Carlsson G (2018) Towards a new approach to reveal dynamical organization of the brain using topological data analysis. *Nature Communication* 9:1–33.

29.   Madhyastha TM, Askren MK, Boord P, Grabowski TJ (2015) Tara M. Madhyastha, Mary K. Askren, Peter Boord, and Thomas J. Grabowski 1,2. 5(1):45–59.

30.   Shine JM, et al. (2016) The Dynamics of Functional Brain Networks: Integrated Network States during Cognitive Task Performance. *Neuron* 92(2):544–554.

31.   Vidaurre D, Smith SM, Woolrich MW (2017) Brain network dynamics are hierarchically organized in time. *Proc Natl Acad Sci* (48):12827-12832.

32.   Résibois M, et al. (2017) The neural basis of emotions varies over time: different regions go with onset- and offset-bound processes underlying emotion intensity. *Soc Cogn Affect Neurosci* 12(8):1261–1271.

33.   Tobia MJ, Hayashi K, Ballard G, Gotlib IH, Waugh CE (2017) Dynamic functional connectivity and individual differences in emotions during social stress. *Hum Brain Mapp* 38(12):6185–6205.

34.   Sadaghiani S, Poline J, Kleinschmidt A, Esposito MD (2015) Ongoing dynamics in large-scale functional connectivity predict perception. *Proc Natl Acad Sci* 112(27):8463–8468.

35.   Ekman M, Derrfuss J, Tittgemeyer M, Fiebach CJ (2012) Predicting errors from reconfiguration patterns in human brain networks. *Proc Natl Acad Sci* 109(41):16714–16719.

36.   Morris JC (1997) Clinical Dementia Rating: A reliable and valid diagnostic and staging measure for dementia of the Alzheimer type. *Int Psychogeriatrics* 9 :173–176.

37. Tombaugh TN, McIntyre NJ (1992) The mini-mental state examination: a comprehensive review. *J*

Am Geriatr Soc 40(9):922–35.

38. Riedl V, et al. (2016) Metabolic connectivity mapping reveals effective connectivity in the resting

human brain. *Proc Natl Acad Sci* 113(8):6260–6266.

39. Davis MHA (1980) A Multidimensional Approach to Individual Differences in Empathy. *JSAS*

Catalog of Selected Documents in Psychology 10:85.

40. Rankin KP, et al. (2006) Structural anatomy of empathy in neurodegenerative disease. *Brain*

129(11):2945–2956.

41. Nana AL, et al. (2018) Neurons selectively targeted in frontotemporal dementia reveal early stage

TDP-43 pathobiology. *Acta Neuropathol* 137(1):27–46.

42. Satterthwaite TD, et al. (2013) An improved framework for confound regression and filtering for

control of motion artifact in the preprocessing of resting-state functional connectivity data.

Neuroimage 64(1):240–256.

43. Gorno-Tempini ML (2004) Cognition and Anatomy in Three Variants of Primary Progressive

Aphasia. *Ann Neurol* 119(11):2658–2666.

44. Hutchison RM, et al. (2013) Dynamic functional connectivity: Promise, issues, and

interpretations. *Neuroimage* 80:360–378.

45. Leonardi N, Van De Ville D (2015) On spurious and real fluctuations of dynamic functional

connectivity during rest. *Neuroimage* 104:430–436.

46. Shirer WR, Ryali S, Rykhlevskaia E, Menon V, Greicius MD (2012) Decoding subject-driven

cognitive states with whole-brain connectivity patterns. *Cereb Cortex* 22(1):158–165.

47.   Wilson RS, et al. (2015) Influence of epoch length on measurement of dynamic functional

752   connectivity in wakefulness and behavioural validation in sleep. *Neuroimage* 112:169–179.

753   48.   Power JD, et al. (2011) Article Functional Network Organization of the Human Brain. *Neuron*

754   72(4):665–678.

755   49.   Dosenbach NUF, et al. (2007) Distinct brain networks for adaptive and stable task control in

756   humans. *Proc Natl Acad Sci* 104(26):11073–11078.

757   50.   Seeley WW, Crawford RK, Zhou J, Miller BL, Greicius MD (2009) Neurodegenerative Diseases

758   Target Large-Scale Human Brain Networks. *Neuron* 62(1):42–52.

759   51.   Toller G, et al. (2018) Individual differences in socioemotional sensitivity are an index of salience

760   network function. *Cortex* 1–13.

761   52.   Friederici AD, Gierhan SM (2013) The language network. *Curr Opin Neurobiol* 23(2):250–254.

762   53.   Sadaghiani S, D'Esposito M (2015) Functional Characterization of the Cingulo-Opercular Network

763   in the Maintenance of Tonic Alertness. *Cereb Cortex* 25(9):2763–2773.

764   54.   Guo CC, et al. (2013) Anterior temporal lobe degeneration produces widespread network-driven

765   dysfunction. *Brain* 136(10):2979–2991.

766   55.   Hickok G, Poeppel D (2007) The cortical organization of speech processing. *Nat Rev Neurosci*

767   8(5):393–402.

768   56.   Wei J, et al. (2018) Eyes-Open and Eyes-Closed Resting States With Opposite Brain Activity in

769   Sensorimotor and Occipital Regions: Multidimensional Evidences From Machine Learning

770   Perspective. *Front Hum Neurosci* 12:422.

57.   Agcaoglu O, Wilson T, Wang Y, Stephen J, Calhoun VD (2019) Resting state connectivity

differences in eyes open versus eyes closed conditions. *Hum Brain Mapp*:hbm.24539.

58.   Hahn A, et al. (2018) Task-relevant brain networks identified with simultaneous PET/MR imaging

of metabolism and connectivity. *Brain Struct Funct* 223(3):1369–1378.

59.   Shine JM, Koyejo O, Poldrack RA (2016) Temporal metastates are associated with differential

patterns of time-resolved connectivity, network topology, and attention. *Proc Natl Acad Sci U S A*

113(35):9888–91.

60.   Kucyi A (2018) Just a thought: How mind-wandering is represented in dynamic brain connectivity.

Neuroimage 180(Pt B):505–514.

61.   Haimovici A, Tagliazucchi E, Balenzuela P, Laufs H (2017) On wakefulness fluctuations as a source

of BOLD functional connectivity dynamics. *Sci Rep* 7(1):5908.

62.   Gordon EM, et al. (2017) Precision Functional Mapping of Individual Human NeuroResource

Precision Functional Mapping of Individual Human Brains. *Neuron* 95(4):791-807.

63.   Laumann TO, et al. (2015) Functional System and Areal Organization of a Highly Sampled

Individual Human Brain. *Neuron* 87(3):657–670.

64.   Poldrack RA, et al. (2015) Long-term neural and physiological phenotyping of a single human. *Nat*

Commun 6:8885.

65.   Guo CC, et al. (2012) One-year test–retest reliability of intrinsic connectivity network fMRI in

older adults. *Neuroimage* 61(4):1471–1483.

66.   Menon V (2011) Large-scale brain networks and psychopathology: A unifying triple network

model. *Trends Cogn Sci* 15(10):483–506.

67.   Laumann TO, et al. (2017) On the Stability of BOLD fMRI Correlations. *Cereb Cortex* 27(10):4719-

4732.

68.   Choe AS, et al. (2017) NeuroImage Comparing test-retest reliability of dynamic functional

795   connectivity methods. *Neuroimage* 158:155–175.

796   69.   Hindriks R, et al. (2016) Can sliding-window correlations reveal dynamic functional connectivity in

797   resting-state fMRI? *Neuroimage* 127:242–256.

798   70.   Byrge L, Kennedy DP (2017) Identifying and characterizing systematic temporally-lagged BOLD

799   artifacts. *Neuroimage* 171:376–392.

800   71.   Nalci A, Rao BD, Liu TT (2018) Nuisance Effects and the Limitations of Nuisance Regression in

801   Dynamic Functional Connectivity fMRI. *bioRxiv*:285239.

802   72.   Power JD, et al. (2014) NeuroImage Methods to detect , characterize , and remove motion

803   artifact in resting state fMRI. *Neuroimage* 84:320–341.

804   73.   Gutierrez-Barragan D, Basson MA, Panzeri S, Gozzi A (2018) Oscillatory brain states govern

805   spontaneous fMRI network dynamics. *bioRxiv*:393389.

806   74.   Mitra A, Snyder AZ, Constantino JN, Raichle ME (2015) The Lag Structure of Intrinsic Activity is

807   Focally Altered in High Functioning Adults with Autism. *Cereb Cortex* 27(2):1083-1093.

808   75.   Matsui T, Murakami T, Ohki K (2016) Transient neuronal coactivations embedded in globally

809   propagating waves underlie resting-state functional connectivity. *Proc Natl Acad Sci*

810   113(23):6556–6561.

811   76.   Liu X, et al. Subcortical evidence for a contribution of arousal to fMRI studies of brain activity. *Nat*

812   *Commun* (2018):1–10.

813   77.   Abrol A, et al. (2017) Replicability of time-varying connectivity patterns in large resting state fMRI

814   samples. *Neuroimage* 163(September):160–176.