Title page

Title

State and trait characteristics of anterior insula time-varying functional connectivity

Authors

Lorenzo Pasquini[1], Gianina Toller[1], Adam Staffaroni[1], Jesse A. Brown[1], Jersey Deng[1], Alex Lee[1], Katarzyna
Kurcyus[2,3], Suzanne M. Shdo[5], Isabel Allen[1], Virginia E. Sturm[1], Yann Cobigo[1], Valentina Borghesani[1],
Giovanni Battistella[1], Maria Luisa Gorno-Tempini[1], Katherine P. Rankin[1], Joel Kramer[1], Howard H. Rosen[1],
Bruce L. Miller[1], William W. Seeley[1,4]*

Affiliations

Memory and Aging Center, Departments of Neurology[1] and Pathology[4], University of California, San
Francisco, California USA; Departments of Neuroradiology[2] and Nuclear Medicine[3], Klinikum rechts der
Isar, Technische Universität München, Munich, Germany, [5]Psychology, University of California Berkeley

*Corresponding author: William W. Seeley, MD. 675 Nelson Rising Lane 94158, San Francisco, California
USA. Email: bill.seeley@ucsf.edu. Phone: +1-415-476-2793

Running title: Time-varying functional connectivity of the anterior insula

Key words: Anterior insula, socio-emotional functioning, time-varying functional connectivity, trait, state

Abbreviations

Anterior insula = aINS; BOLD = blood oxygen level dependent; inferior frontal gyrus = IFG; task-free
functional MRI = tf-fMRI

Abstract

The human anterior insula (aINS) is a topographically organized brain region, in which ventral portions
contribute to socio-emotional function through limbic and autonomic connections, whereas the dorsal
aINS contributes to cognitive processes through frontal and parietal connections. Open questions
remain, however, regarding how aINS connectivity varies over time. We implemented a novel approach
combining seed-to-whole-brain sliding-window functional connectivity MRI and k-means clustering to
assess time-varying functional connectivity of aINS subregions. We studied three independent large
samples of healthy participants and longitudinal datasets to assess inter- and intra-subject stability, and
related aINS time-varying functional connectivity profiles to dispositional empathy. We identified four
robust aINS time-varying functional connectivity modes that displayed both "state" and "trait"
characteristics: while modes featuring connectivity to sensory regions were modulated by eye closure,
modes featuring connectivity to higher cognitive and emotional processing regions were stable over
time and related to empathy measures.

Introduction

The human anterior insula (aINS) is a functionally heterogeneous region implicated in functions ranging
from interoceptive awareness and emotion processing to time perception and cognitive control (1). In
humans, neuroimaging studies have begun to parcellate the aINS, based on its patterns of functional
and structural connectivity. For example, task-free fMRI (tf-fMRI) studies, which measure the brain-wide
correlation structure in slow (< 0.1 Hz), spontaneous blood oxygen level dependent (BOLD) signal
fluctuations (2, 3), have shown that the ventral, agranular aINS is functionally connected to limbic and
autonomic processing regions that include the pregenual anterior cingulate cortex, the amygdala, and
subcortical structures such as the thalamus and periaqueductal gray (4–8). These regions make up the
"salience network", a large-scale distributed system that represents the homeostatic significance of
prevailing stimuli and conditions (9–13). In contrast, the dorsal, dysgranular aINS is connected to a
cingulo-opercular "task-control network" whose nodes include dorsolateral and opercular prefrontal,
anterior midcingulate, and anterior parietal areas involved in cognitive control processes such as task-
set initiation and maintenance (4, 7, 8, 14). Under task-free conditions, both aINS subregions, but
perhaps especially the dorsal aINS (8, 15), show activity that is anticorrelated with the "default mode
network", a system including the posterior cingulate cortex, inferior parietal lobules, and precuneus
(16). In addition to the dorsal-ventral axis, hemispheric lateralization of the aINS is proposed to help
maintain bodily homeostasis by adjusting and balancing autonomic outflow based on bioenergetics
demands (1, 10, 11). Recent evidence suggests that while the left-sided (dominant hemisphere) aINS
controls parasympathetic tone, the homotopic right (non-dominant) aINS is more closely linked to
sympathetic tone and responses (1, 10, 11). Structural and functional changes in aINS subregions have
been reported in a variety of neuropsychiatric conditions, ranging from mood and anxiety disorders to
schizophrenia, autism, and frontotemporal dementia (13, 17–20).

Standard tf-fMRI has helped to reveal the functional organization of the human aINS and other brain
areas by providing a snapshot of functional connectivity as averaged across the duration of a scanning
session. Although long-range structural connections are assumed to be relatively stable in the adult
brain (21), coordinated functional activity is dynamic, with the brain continuously reshaping network
configurations in response to prevailing conditions or task demands (22–24). Approaches that capture
time-varying connectivity bring the potential to clarify how brain dynamics are organized and relate to
function. We reasoned that this approach could help clarify how the aINS may change its network
partners in response to salient internal and external stimuli and possibly contribute to the development
of more granular and personalized fingerprints of brain function in health and disease. Such time-varying
functional connectivity has been shown to relate to clinical outcomes (25–27); task performance (28);
and behaviorally relevant measures of cognitive (29–31), emotional (32, 33), and attentional processing
(34, 35). One of the few studies applying time-varying functional connectivity analyses to the insula
parcellated it into posterior, middle, and dorsal and ventral anterior components, revealing partially
overlapping time-varying connectivity profiles for ventral and dorsal aINS subregions. Ventral and dorsal
profiles diverged based on distinct contributions from limbic/emotional processing and cingulo-
opercular/cognitive regions mirroring findings from static tf-fMRI studies (15).

Despite this body of research, it remains unclear how distinct aINS subregions dynamically engage the
salience and task-control networks over time. Moreover, it is largely unknown whether time-varying
functional connectivity of the aINS — or any region for that matter — is modulated by externally-driven

states or instead displays trait characteristics such as within-subject temporal stability and relationships
to behavioral or dispositional measures. To gain a deeper understanding of these issues, we
implemented a novel approach combining seed-to-whole-brain sliding-window functional connectivity
and k-means clustering to derive time-varying functional connectivity profiles of aINS subregions across
three large, independent samples of healthy participants, including eyes open vs. eyes closed conditions
and longitudinal datasets. Across methods and samples, we found that aINS subregions display shared
and distinct time-varying connectivity "modes" that bind together cognitive and/or emotional
processing areas versus upstream sensory and motor cortices. Overall, the findings suggest that short-
term temporal variability in aINS connectivity reflects both state and trait characteristics, revealing a
path toward use of such data for assessing psychopharmacological treatment efficacy as well as long-
term therapeutic disease modification in neuropsychiatric conditions.

Materials and Methods

Participants. All participants were healthy, cognitively normal adults, recruited from two different
centers (Table 1, Supplementary Figure S1). Elderly participants were retrospectively selected from the
Hillblom Aging Network, an extensively characterized longitudinal cohort assessed at the University of
California, San Francisco (UCSF) Memory and Aging Center. Subjects were required to have a Clinical
Dementia Rating Scale score (36) of 0 (range 0-3) and a Mini-Mental State Examination score (37) of 28
(range 0-30) or higher. Secondary inclusion criteria were based on availability of the Interpersonal
Reactivity Index, a widely used questionnaire measuring emotional and cognitive empathy. Out of 224
elderly subjects with a tf-fMRI scan and at least one Interpersonal Reactivity Index available, a cross-
sectional cohort of 121 elderly adults was selected based on availability of the Interpersonal Reactivity
Index within three months of the tf-fMRI scan. A second sample was built based on availability of
longitudinal tf-fMRI scans. Out of total 184 older adults with longitudinally assessed tf-fMRI, 68 were
excluded because they overlapped with the cross-sectional sample (N = 121, described above) and 72
were excluded since they did not meet our selection criteria of being scanned twice within a 5-13 month
interval. This procedure resulted in a final selection of 44 participants having longitudinal data and that
did not overlap with the cross-sectional sample. All visits included neuropsychological testing and a
neurologic exam in addition to a structural MRI and tf-fMRI scan. Exclusion criteria included a history of
drug abuse, psychiatric or neurological conditions, and current use of psychoactive medications.

Twenty additional younger participants were recruited from a simultaneous FDG-PET/tf-fMRI study at
the Klinikum rechts der Isar, Technische Universität München (38). Neuroimaging data was assessed
twice within an interval of one month, during task-free conditions with either eyes closed or eyes open.
Participants were randomly assigned to one of the two conditions, resulting in eight participants
assessed with eyes closed at the first and with eyes open at the second scan, and 12 participants
assessed with eyes open at the first and eyes closed at the second scan. Exclusion criteria included a
history of psychiatric or neurological conditions, use of psychoactive medications, pregnancy, and renal
failure.

For all samples, written informed consent was obtained from all involved participants and the study was
approved by the institutional review board where the data was acquired (UCSF/Klinikum rechts der Isar).

Empathy Assessment. The Interpersonal Reactivity Index (39) was completed by the participant's
informant within three months of tf-fMRI scan. This questionnaire is composed by four subscales, each

one consisting of 7 questions that can comprehensively reach a maximum score of 35. While empathic
concern and personal distress measure emotional aspects of empathy, perspective taking and the
fantasy score are designed to assess cognitive aspects of empathy. The Interpersonal Reactivity Index
was chosen since it has been widely used to assess different aspects of empathy in healthy and
neuropsychiatric conditions (39, 40). For example, studies in frontotemporal dementia, a
neurodegenerative diseases of socioemotional dysfunction, have shown that deficits in subscales of the
Interpersonal Reactivity Index are linked to structural deterioration of the aINS and regions connected to
the aINS such as the anterior cingulate and orbitofrontal cortices (40, 41).

Neuroimaging data acquisition. The cross-sectional and longitudinal cohorts from the Hillblom Aging
Network were scanned at the UCSF Neuroscience Imaging Center on a Siemens Trio 3T scanner. A T1-
weighted MP-RAGE structural scan was acquired with acquisition time=8 min 53 sec, sagittal orientation,
a field of view of 160 x 240 x 256 mm with an isotropic voxel resolution of 1 mm^3, TR=2300 ms, TE=2.98
ms, TI=900 ms, flip angle=9°. Task-free T2*-weighted echoplanar fMRI scans were acquired with an
acquisition time=8 min 6 sec, axial orientation with interleaved ordering, field of view=230 x 230 x 129
mm, matrix size=92 x 92, effective voxel resolution=2.5 x 2.5 x 3.0 mm, TR=2000 ms, TE=27 ms, for a
total of 240 volumes. During the 8-minute tf-fMRI acquisition protocol, participants were asked to close
their eyes and concentrate on their breathing.

Data from the Klinikum rechts der Isar was acquired on an integrated Siemens Biograph scanner capable
of simultaneously acquiring PET and MRI data (3T). FDG-PET activity and tf-fMRI was simultaneously
measured during the initial 10 min immediately after bolus injection of the FDG tracer. Scanning was
performed in a dimmed environment obtained by switching off all lights, including those in the scanner
bore. Subjects were instructed to keep their eyes closed or open, to relax, to not think of anything in
particular, and to not fall asleep. MRI data were acquired using the following scanning parameters: Task-
free echoplanar fMRI scans: TR, 2.000 ms; TE, 30 ms/angle, 90°; 35 slices (gap, 0.6 mm) aligned to AC/PC
covering the whole brain; sFOV, 192 mm; matrix size, 64 x 64; voxel size, 3.0 x 3.0 x 3.0 mm^3 (each
measurement consists of 300 acquisitions in interleaved mode with a total scan time of 10 min and 8 s);
MP-RAGE: TR, 2.300 ms; TE, 2.98 ms; angle, 9°; 160 slices (gap, 0.5 mm) covering the whole brain; FOV,
256 mm; matrix size, 256 x 256; voxel size, 1.0 x 1.0 x 1.0 mm^3; total length of 5 min and 3 s.

Neuroimaging data preprocessing. Before preprocessing, all images were visually inspected for quality
control. Images with excessive motion or image artifact were excluded. T1-weighted images underwent
segmentation using SPM12 (http://www.fil.ion.ucl.ac.uk/spm/software/spm12/). For each tf-fMRI scan,
the first five volumes were discarded. SPM12 and FSL (http://fsl.fmrib.ox.ac.uk/fsl) software were used
for subsequent tf-fMRI preprocessing. The remaining volumes were slice-time corrected, realigned to
the mean functional image and assessed for rotational and translational head motion. Volumes were
next co-registered to the MP-RAGE image, then normalized to the standard MNI-152 healthy adult brain
template using SPM segment, producing MNI-registered volumes with 2 mm^3 isotropic resolution. These
volumes were spatially smoothed with a 6 mm radius Gaussian kernel and temporally bandpass filtered
in the 0.008-0.15 Hz frequency range using fslmaths. Nuisance parameters in the preprocessed data
were estimated for the CSF using a mask in the central portion of the lateral ventricles and for the white
matter using a highest probability cortical white matter mask as labeled in the FSL tissue prior mask.
Additional nuisance parameters included the 3 translational and 3 rotational motion parameters, the

temporal derivatives of the previous 8 terms (white matter/CSF/6 motion), and the squares of the
previous 16 terms (42). Subjects were included only if they met all of the following criteria: no inter-
frame head translations greater than 3 mm, no inter-frame head rotations greater than 3 degrees, and
less than 24 motion spikes (defined as inter-frame head displacements > 1 mm), less than 10% of the
total number of frames. Nuisance parameters were regressed out from the filtered data using fslmaths,
and masked with a binarized, skullstripped MNI-152 brain mask. The WM/CSF/head movement
denoised data was used for the subsequent time-varying functional connectivity analysis. Findings with
global signal regression are not presented in the main body of the manuscript but were generated for
the longitudinal dataset and are reported as sensitivity analysis in the Supplement.

Time-varying functional connectivity analysis. For each individual, average blood oxygen level dependent
signal time courses were extracted from the right and left ventral and dorsal aINS using four regions-of-
interest from the Brainnetome Atlas (http://atlas.brainnetome.org/). In order to assess the impact of
seed region selection, in the cross-sectional dataset activity time courses were also extracted from a
language-relevant region centered on the left inferior frontal gyrus (IFG), using a region-of-interest
defined in a previous study (43). Using in-house custom scripts based on Python
(https://www.python.org/) and FSL, a sliding-window approach was implemented to generate time-
varying seed-to-whole-brain connectivity maps (Figure 1A). The derived time series from the aINS and
the entire tf-fMRI scan of each subject were divided into sliding-windows of 18 TRs (36 s) in steps of 1 TR
creating 218 (275 for eyes closed/open) untapered, rectangular windows. At each window, linear
regression was used to derive seed-to-whole-brain time-varying functional connectivity maps for each
aINS seed (Figure 1B). A window size of 36 seconds was chosen based on previous research showing that
window sizes between 30 and 60 seconds capture additional variations in functional connectivity not
found in larger window sizes (24, 25, 44). Ideal sliding-window size has been explored by additional
methodological work assessing the relationship between window length and cut-off frequencies,
supporting the use of sliding-windows between lengths of 30–60 seconds for tf-fMRI data preprocessed
using a low-pass filter set at 0.15 Hz (45). These studies are further supported by empirical findings
showing that cognitive states can be discerned within such window lengths (46, 47). Nevertheless, in
order to assess the impact of sliding-window length, control analyses were performed with window
lengths of 72 TR (144 s). The resulting findings did not significantly differ from the reported findings with
windows of 36 s (Supplementary Results and Figure S2); therefore 36 s windows were used throughout.
The code used to derive time-varying functional connectivity maps is provided in the Appendix of the
Supplement and in GitLab (https://gitlab.com/juglans/sliding-window-analysis/tree/master). The
derived time-varying functional connectivity maps were finally standardized to z-scores (see Video 1 for
left ventral aINS time-varying functional connectivity maps of a typical study participant), masked with a
binarized gray matter mask, vectorized, and concatenated across subjects resulting in windows x voxels
matrices for each aINS seed. K-means clustering was then applied to the concatenated window matrices
using a k = 4 to produce four clusters representing four time-varying functional connectivity modes of
aINS subregions present across the course of the functional scan (Figure 1C). The k-means algorithm
used Euclidean distance, and an optimal solution was selected after 100 iterations and 10 replications.
The optimal number of clusters, referred to henceforward as modes, was determined using elbow and
silhouette plots and by performing additional clustering solutions with k = 3, 5, and 6 (Supplementary
Results and Figure S3). With lower k solutions, the identified modes merged, resulting in information
loss. Using higher k solutions resulted in the generation of redundant sub-modes, exemplified by Mode

4 of the left ventral aINS that would split in anterior and posterior centered components
(Supplementary Figure S3B). A total of three clustering analyses per aINS subregion were independently
performed on time-varying functional connectivity windows: one on the cross-sectional dataset; one on
the longitudinal dataset; and one on the eyes closed/open dataset (see Supplementary Figure S1 for a
schematized summary of k-means clustering analyses performed).

Group-averaged maps of the four identified modes were generated, for each seed region, using the
mode-specific centroid maps rendered using k-means clustering. The spatial similarity of modes derived
from distinct aINS subregions and distinct groups was assessed by vectorizing the mode's template maps
and performing Pearson's correlation analyses. To assess the distinct spatial contribution of major large-
scale brain networks to time-varying connectivity modes, we used publicly available templates of major
brain networks from a study investigating the functional network organization of the human brain
(48)(https://www.jonathanpower.net/2011-neuron-bigbrain.html). Briefly, in this study subgraphs
corresponding to major brain systems were derived from tf-fMRI data of > 300 healthy adults by
retaining 2% of the strongest correlations. This procedure resulted in 12 binary templates spanning
cognitive, primary sensory and subcortical systems: the salience, default, cingulo-opercular task-control,
executive-control, ventral attention, dorsal attention, auditory, visual, ventral sensorimotor, dorsal
sensorimotor, medial temporal lobe ("memory retrieval") and subcortical networks (4, 48, 49)
(Supplementary Figure S4). Subsequently, for each aINS subregion the averaged z-score value enclosed
within these brain network templates was extracted from the mode-specific centroid maps generated in
the cross-sectional dataset.

Further, the assignment of each window to a specific mode was used to derive: (i) the number of
transitions from one mode to another; (ii) a fractional occupancy metric for each mode, defined as the
number of windows assigned to that mode divided by the total number of windows; and (iii) how often
aINS subregions simultaneously occupy distinct mode configurations in time. To identify meta-profiles of
aINS time-varying functional connectivity, individual subject fractional occupancy profiles, defined as a
vector of 16 elements summarizing the time fraction separately spent by individual aINS subregions on
the four identified modes, were derived from the cross-sectional dataset. Fractional occupancy profiles
were subsequently clustered using k-means with a clustering solution of k = 4 based on a silhouette
analysis (using Euclidean distance, 100 iterations, and 10 replication). On the cross-sectional sample,
static functional connectivity maps were also generated through voxel-wise regression analyses of each
aINS seed's time-course for the duration of the entire scan, following previous established methods (50,
51).

Statistical analysis. Statistical analyses were carried out using R (https://www.r-project.org/) and
Matlab-R2018 (https://www.mathworks.com/products/matlab.html). Pearson's correlation was used to
assess the spatial similarity of modes derived from different clustering analyses, using maps thresholded
for z-values higher than 0.3 and lower than -0.3.

In the cross-sectional dataset, the Shannon diversity index was used to analyze whether aINS subregions
of an individual subject transitioned repeatedly between distinct modes or showed instead a stereotypic
behavior spending most of the time only in a specific aINS mode:

$$\text{Shannon diversity index } = -\sum_{i=0}^{n-1} p_i log_2 p_i$$

wherein p is the fractional occupancy of an aINS subregion in a specific time-varying functional
connectivity mode. A Shannon diversity index of 1.4 indicates that all modes are equally occupied by the

aINS, while the lower the Shannon diversity index, the higher the fractional occupancy of the aINS on
one specific mode.

In the cross-sectional dataset, one-way ANOVAs were used to investigate whether aINS subregions
differed, within subjects, in terms of mode-specific fractional occupancy, number of transitions, and
Shannon diversity index. For descriptive purposes, averages and standard deviations derived over all
aINS subregions and modes are reported. Additionally, for each meta-profile cluster identified in the
cross-sectional dataset, one-way ANOVAs were used to assess differences in mode-specific fractional
occupancy averaged across aINS subregions. The assignment of participant to four meta-profiles of time-
varying aINS fractional occupancy was additionally used to test differences in aINS subregions' static
connectivity using ANOVA models and post-hoc t-tests implemented in SPM12 (height threshold
p<0.005; extent threshold p<0.05 FWE corrected for multiple comparisons).

Finally, in the cross-sectional dataset, four multiple linear regression models were used to test the
association of mode-specific fractional occupancies with subscales of the Interpersonal Reactivity Index
(39). Subscales of the Interpersonal Reactivity Index were used as dependent variables, and each model
contained the mode-specific fractional occupancies averaged across aINS subregions. Each model was
corrected for age, sex, and sum frame-wise head displacement (p < 0.05 uncorrected for multiple
comparisons). To assess specificity of the aINS results, four additional models were estimated, using
subscales of the Interpersonal Reactivity Index as dependent variables and fractional occupancy of time-
varying functional connectivity modes derived from the left IFG as predictors.

For the longitudinal datasets, cosine similarity was used to assess the similarity of fractional occupancy
profiles across distinct scanning dates.

265
$$\text{Cosine similarity} = 1 - \left(\frac{\sum_{i=1}^{n} A_i B_i}{\sqrt{\sum_{i=1}^{n} A_i^2} \sqrt{\sum_{i=1}^{n} B_i^2}} \right)$$

Where *A* and *B* are the vectorized fractional occupancy profiles of all aINS subregions at two different
scanning dates. Cosine similarity varies between 1 and 0, with 1 indicating identical fractional occupancy
profiles. Paired t-tests were performed to assess differences in fractional occupancy in the longitudinal
and in the eyes closed/open datasets. All statistical findings are reported at p < 0.03, two-tailed,
uncorrected for multiple comparisons except where specified otherwise.

Results

Bilateral ventral and dorsal aINS occupy four overlapping but distinct time-varying functional
connectivity modes

By combining seed-to-whole-brain functional connectivity, sliding-window analysis, and k-means
clustering, we found that aINS subregions adopt four overlapping but distinct large-scale configurations
or "modes" of time-varying functional connectivity (Figure 2A). All modes were characterized by
bilateral connectivity between the aINS and anterior cingulate cortices. Modes could be distinguished
from each other, however, by specific patterns of connectivity to other brain regions (Figure 2A and
Supplementary Table S1). In Mode 1, all four aINS subregions showed connectivity to the anterior

midcingulate/pre-supplementary motor area, right frontal operculum, and dorsal parietal and
dorsolateral prefrontal areas that together make up the task-control network. Mode 2 was
characterized by negative connectivity to primary visual and sensorimotor areas and prominent
connectivity to the ventral striatum and thalamus. Mode 3 showed inverted connectivity patterns to the
same regions. Finally, Mode 4 showed connectivity patterns aligned with more ventral salience network
regions, including pre- and sub-genual anterior cingulate and orbitofrontal cortices, with additional
connectivity to the temporal poles for the right ventral aINS. All four modes were identified in each aINS
subregion, as shown by the correlation matrix highlighting the spatial correspondence of equivalent
modes (Figure 2B). Important distinctions were found, however, when comparing dorsal and ventral
aINS connectivity patterns within modes. In Mode 1, the dorsal aINS showed prominent anti-
correlations to the precuneus, angular gyrus, and medial prefrontal cortex regions that make up the
default mode network (16). In Mode 4, the ventral aINS showed a more ventral connectivity pattern that
encompassed subgenual anterior cingulate cortex and temporal poles, when compared to the dorsal
aINS (Figure 2C, Supplementary Figure S6A). Modes derived from homologous left and right aINS regions
differed from each other mainly with regard to the extent of ipsilateral connectivity to neighboring
regions (Supplementary Figure S6B-C). Importantly, similar aINS time-varying functional connectivity
modes were identified using longer sliding-window lengths (Supplementary Figure S2), lower and higher
clustering solutions (Supplementary Figure S3), and data preprocessed using global signal regression
(Supplementary Figure S7).

aINS subregions coherently transition between time-varying functional connectivity modes

Average fractional occupancy, i.e. the proportion of time spent by aINS subregions in the four time-
varying functional connectivity modes ranged from 0.18 (left ventral aINS in Mode 2) to 0.30 (right
ventral aINS in Mode 1), with an overall average of 0.25 +/- 0.20 (Figure 3A). Importantly, no significant
differences in mode-specific fractional occupancy were found across the four aINS subregions (Mode 1:
F = 0.2, p = 0.82; Mode 2: F = 0.7, p = 0.56; Mode 3: F = 0.4, p = 0.77; Mode 4: F = 0.5, p = 0.71). In only a
few participants did aINS subregions spend disproportionate time in only one mode. The number of
transitions was also comparable across aINS subregions (F = 0.4, p = 0.759; Figure 3B), with an overall
average of 18.0 +/- 6.6 transitions over the task-free acquisition. In particular, our data shows that there
were no direct transitions between the primary sensory anticorrelated Mode 2 and the primary sensory
correlated Mode 3 (Supplementary Figure S8), suggesting that aINS subregions need to transition to
modes rooted in cognitive networks before occupying primary sensory-centered modes characterized by
opposing connectivity patterns. The Shannon diversity index was used to quantify the diversity of
fractional occupancy and to compare this diversity across subregions. On average, aINS subregions
showed high Shannon diversity, suggesting that each subregion moved between modes in most
subjects, rather than stacking on a single mode. The overall average Shannon diversity index of
fractional occupancy across aINS subregions was 1.0 +/- 0.2 and did not significantly differ between
subregions (F = 0.8; p = 0.503; Figure 3C). Subject-level fractional occupancy profiles are illustrated in
Figure 3D, which shows two subjects with contrasting signatures. We finally sought to assess whether
aINS subregions coherently occupy the same modes at the same time, or for instance, how often the
right ventral and dorsal aINS are simultaneously connected to Mode 1. For each subject and each aINS
subregion, we assessed how often possible mode configurations were jointly occupied in time. This
analysis revealed a tendency across aINS subregions to occupy the same modes in time (Figure 3E). On
average, the right ventral and dorsal aINS occupied the same modes 50% of the time, the left ventral

and dorsal aINS 53% of the time, the right and left ventral aINS in 51% of the time, and the right and left
dorsal aINS 61% of the time. These findings suggest an overall tendency for aINS subregions to
cohesively engage the same modes over time under task-free scanning conditions.

Individual aINS connectivity mode occupancy profiles are reproducible across samples and within
subjects over time

To assess the generalizability and reproducibility of our findings, we next turned to a longitudinal
healthy aging dataset, which consisted of 44 participants who were scanned twice over an interval
between 5-13 months. Clustering of time-varying connectivity windows identified modes that strongly
resembled the four modes derived from the larger (and non-overlapping) cross-sectional dataset (Figure
4A-D and Supplementary Figure S5 panels A, C and E). Average fractional occupancies of aINS subregions
in the four modes were comparable with the cross-sectional sample (comparing Figure 3A to 4A-D).
Across aINS subregions and modes, paired t-tests revealed no significant differences in fractional
occupancy when comparing the first scanning date with the second ($p < 0.05$ uncorrected for multiple
comparisons, for details on paired t-test statistics see Supplementary Table S2). Within subjects, cosine
similarity analysis showed that most participants had relatively stable fractional occupancy profiles over
time (mean cosine similarity = 0.70 +/- 0.19 s.d.), as further supported by the inverted bell-shaped
distribution of cosine similarity (Figure 5A). Stable fractional occupancies are exemplified in Subjects 10
(first scanning date in red; second scanning date in blue; cosine similarity = 0.98) and 41 (cosine
similarity = 0.92) (Figure 5B-C). A few participants, however, showed very different fractional occupancy
profiles when comparing the two scanning dates, as exemplified by Subjects 21 (cosine similarity = 0.46)
and 26 (cosine similarity = 0.24) (Figure 5D-E). Overall, however, the temporal stability of the individual
profiles suggests a major contribution from trait-level factors.

Eye opening decreases the time aINS spends anticorrelated with visual cortices

Independently performed clustering of time-varying connectivity data from the eyes closed/open
dataset revealed similar aINS modes to those identified in the cross-sectional and longitudinal aging
datasets (Figure 6A-B and Supplementary Figure S5 panels B, D and E). For all subregions except the
right ventral aINS, participants showed higher fractional occupancies in the anti-correlated primary
sensory/motor Mode 2 with eyes closed and higher fractional occupancies in the task-control Mode 1
with eyes open ($p < 0.05$ uncorrected, for details on paired t-test statistics see Supplementary Table S2).

Time-varying functional connectivity meta-profiles show that aINS subregions cohesively occupy the
same modes

Individual subjects were clustered into meta-profiles (see Methods "*Time-varying functional connectivity*
analysis"), based on the tendency of their aINS subregions to occupy specific time-varying functional
connectivity modes. Individual fractional occupancy profiles of the cross-sectional sample (n = 121) were
clustered into four clusters using k-means. Silhouette plots were used to choose the ideal number of
clusters, k = 4 (Supplementary Figure S9). This analysis revealed four meta-profiles, characterized by
preferred fractional occupancy in one of the four time-varying functional connectivity modes (Figure
7A). Meta-profile 1 (in green, 26 subjects) was characterized by a tendency to spend time in the task-
control Mode 1 (see Subject 22, Figure 7A); Meta-profile 2 (in blue, 31 subjects) was characterized by a
tendency to spend time in the sensory anticorrelated Mode 2 (see Subject 79, Figure 7A); Meta-profile 3
(in violet, 44 subjects) was characterized by a tendency to spend time in the sensory connected Mode 3

(see Subject 80, Figure 7A); while Meta-profile 4 (in red, 20 subjects) was characterized by a tendency to
spend time in the salience network Mode 4 (see Subject 88, Figure 7A). To evaluate statistical
differences in mode occupancy across meta-profiles, we averaged the time spent in each mode across
aINS subregions and used one-way ANOVAs to test for significant group differences in mode-specific
fractional occupancy across the distinct aINS meta-profiles (Meta-profiles 1, F = 50.1, p < 0.0001; Meta-
profile 2, F = 39.8, p < 0.0001; Meta-profile 3, F = 79.6, p < 0.0001; Meta-profile 4, F = 46.1, p < 0.0001,
see also Figure 7B). These differences were further statistically assessed via post-hoc t-tests. In
summary, these analyses revealed a preference for aINS subregions to collectively spend similar
fractions of time in a given mode, whereas distinct modes were primarily occupied by subjects in
different meta-profiles.

Next, we asked whether the constituents of a given meta-profile would show group-level differences in
terms of their static functional connectivity patterns. We therefore compared groups based on static
functional connectivity of aINS subregions across the four meta-profiles described above. This analysis
revealed that participants in Meta-profile 1 showed prominent static anticorrelation of the dorsal aINS
to the default mode network when compared to participants clustered on other meta-profiles.
Participants clustered on Meta-profile 2 and 3 showed marked visual and sensorimotor cortex static
hypoconnectivity and hyperconnectivity, across all aINS subregions, when compared to participants of
other meta-profiles; while participants in Meta-profile 4 showed prominent static connectivity of all
aINS subregions to more ventral frontal regions (Supplementary Figure S10). In summary, key
components of the time-varying connectivity modes were reflected in static connectivity differences
between meta-profile-based groups.

Comparison of aINS and left IFG time-varying functional connectivity

To control for the location of the seed, we derived time-varying functional connectivity modes by
seeding the left IFG. Time-varying functional connectivity modes of the left IFG shared some common
themes with modes identified using aINS subregions (Supplementary Figure S11A). The four identified
modes showed similar connectivity to regions of the language network, such as the left IFG, pre-
supplementary motor area, and superior parietal lobule (43, 52). This contrasts with the modes
identified in aINS subregions that showed overlapping time-varying functional connectivity to typical
task-control and salience network regions (Supplementary Figure S11B). Modes derived from both the
aINS and left IFG, however, showed similar patterns of default mode network anticorrelation (Mode 1),
primary sensory anticorrelation (Mode 2), and primary sensory hypercorrelation (Mode 3). A fourth
mode was identified using the left IFG, showing time-varying functional connectivity to regions of the
dorsal attention network (49) (Mode 4).

Time spent by the aINS in the task-control and salience networks correlates with metrics related to
dispositional empathy

A key question of this work concerns whether individual differences in aINS time-varying connectivity
relate to differences in aINS-associated traits and functions. Among the many tasks that activate and
depend on the anterior insula, empathy is one of the best documented (1, 18). The Interpersonal
Reactivity Index is an informant-based questionnaire widely used to assess distinct aspects of empathy
(39). Emotional aspects of empathy are covered by the empathic concern and personal distress
measures, with the first assessing another-centered emotional response, while the second reflects
general anxiety and self-oriented emotional reactivity. Perspective taking and fantasy scores measure

cognitive aspects of empathy, with the first assessing the tendency to spontaneously imagine the
cognitive perspective of another person, while the second measures the tendency to project oneself
into the experiences of fictional characters. Mode-specific fractional occupancies were averaged across
aINS subregions and used as regressors in four models using subscales of the Interpersonal Reactivity
Index as dependent variables. These multiple linear regression models were corrected for age, sex, and
sum frame-wise head displacement and identified two significant positive associations with time-varying
connectivity metrics. First, fractional occupancy in the task-control Mode 1 predicted higher levels of
personal distress, reflecting self-oriented feelings of personal anxiety and unease in tense interpersonal
settings (β = 7.1; $p < 0.05$ uncorrected for multiple comparisons; Supplementary Table S3). Second,
fractional occupancy in the salience Mode 4 correlated with higher scores on the fantasy subscale,
reflecting greater ability to transpose one's self imaginatively into the feelings and actions of fictitious
characters in books, movies, and plays (β = 9.2; $p < 0.05$ uncorrected for multiple comparisons;
Supplementary Table S3). These results remained significant when removing outliers (Mode 1 and
personal distress β = 6.9, $p < 0.05$ uncorrected for multiple comparisons; Mode 4 and fantasy score β =
9.2, $p < 0.05$ uncorrected for multiple comparisons). No significant associations were found between
fractional occupancy in any mode and the empathic concern or perspective taking subscales. As
hypothesized, fractional occupancy of left IFG time-varying connectivity modes was not significantly
associated with any measure of dispositional empathy ($p < 0.05$ uncorrected for multiple comparisons,
see Supplementary Table S3).

Discussion

To date, most efforts to relate large-scale networks to individual differences have relied on metrics that
capture the topology or strength of node-to-node static functional (or "intrinsic") connections. Here, we
developed a novel approach combining seed-to-whole-brain functional connectivity, sliding-window
analysis, and k-means clustering on tf-fMRI data to identify distinct time-varying functional connectivity
modes of the aINS, a brain region critical for many human socio-emotional functions (1, 18). These
robust modes were identified across methods and samples, showing both "state" and "trait"
characteristics. In particular, while aINS modes related to sensory processing were modulated by visual
input (eyes closed vs. open conditions during scanning), modes featuring connectivity to cognitive and
emotion processing regions were stable over time and related to measures of dispositional empathy.

Partially overlapping but distinct time-varying aINS functional connectivity modes

Our pipeline reliably identified four time-varying functional connectivity modes of the aINS across
different subregions, methods, and samples. These modes were characterized by common connectivity
of aINS subregions to the contralateral insula and to the anterior cingulate cortex, but could be
differentiated from each other based on recruitment of other brain regions. Mode 1 was characterized
by a more dorsal connectivity pattern that resembles previous characterizations of a cingulo-opercular
task-control network (4, 14, 53) involving the bilateral aINS and anterior cingulate cortex, with
additional connectivity to dorsomedial parietal and dorsolateral frontal areas. Mode 2 was characterized
by negative connectivity to primary sensory areas and increased connectivity to the thalamus while
Mode 3 showed an inverted connectivity pattern to these regions. Finally, Mode 4 was characterized by
fronto-insular connectivity to more rostral and ventral portions of the anterior cingulate cortex,
resembling the salience network (4, 8), with additional connectivity to the temporal pole and medial
orbitofrontal cortex, components of the "semantic-appraisal network" (18, 54). These findings are

consistent with those of a previous region-of-interest-based time-varying functional connectivity study
focusing on the insula, which reported time-varying visual and sensorimotor hyperconnectivity of the
ventral aINS. Our findings differ, however, in that we did not detect positive default mode network
connectivity to the dorsal aINS and that patterns of hypoconnectivity to primary visual and sensorimotor
areas were less apparent in the aforementioned study (15). While left and right aINS subregions
primarily differed with respect to connectivity to regions in the ipsilateral hemisphere, consistent
differences in time-varying functional connectivity at the system-level were identified when comparing
ventral to dorsal aINS subregions. Mirroring static functional connectivity studies and the
aforementioned time-varying connectivity study on the insula (7, 8, 15), the task-control Mode 1 of the
dorsal aINS showed stronger anti-correlations to the default mode network than did its ventral
counterpart. This observation is in line with salience network models proposing that the dorsal aINS
receives ventral aINS input streams regarding the moment-to-moment condition of the body and then,
based on these inputs, recruits task control (14) and executive control (4) network resources to maintain
cognitive set and guide behavior while inhibiting the default mode network (13). The salience network
Mode 4 showed stronger connectivity between ventral aINS and ventromedial prefrontal, orbitofrontal,
and temporopolar cortices when compared to its dorsal aINS counterpart. This finding is in line with
functional-anatomical models suggesting a close alliance between the salience and the semantic
appraisal networks (13, 18, 54), whose main components (the temporal pole, ventral striatum, medial
orbitofrontal cortex, and amygdala) have been proposed to interact with autonomic representations in
the aINS to construct the meaning and significance of social and non-social stimuli under prevailing
conditions (13, 18, 54). Importantly, we did not find any differences across aINS subregions in the
number of transitions or in the diversity of fractional occupancy profiles. Moreover, our analyses
exploring overlapping mode occupancy across aINS subregions revealed that distinct aINS subregions
tend to occupy the same mode at the same time. Further support for this conclusion comes from the
clustering analysis of individual fractional occupancy profiles, which led to the identification of four
meta-profiles characterized by occupancy of a dominant mode. Although our findings are not conclusive
on whether aINS subregions always engage the same mode in time, they comprehensively suggest a
tendency for coherent mode occupancy that could potentially underpin specific trait characteristics or
behavioral states. Future studies could leverage methods such as directed and effective connectivity to
assess the temporal relationships between intrinsic time-varying activity of distinct aINS subregions,
which could be lagged, phase-shifted, or anticorrelated over time. In order to assess the specificity of
aINS time-varying functional connectivity modes, we assessed and compared time-varying functional
connectivity of the left IFG, a region involved in language processing and syntax production (43, 52, 55).
The left IFG showed region-specific contributions from language processing regions but also displayed
pronounced anticorrelation to default mode network regions in Mode 1 and similar time-varying
connectivity to the same sensory regions found in Modes 2 and 3 derived from the aINS. These findings
suggest that distinct brain regions, although characterized by seed specific connectivity patterns, may
cohesively transition between broader modes characterized by shared regional time-varying functional
connectivity patterns, possibly revealing a fundamental property of functional brain organization.

"State" characteristics of aINS time-varying functional connectivity modes

Having determined that two of the four major modes of aINS connectivity involved varying connectivity
to sensory, and, most notably, visual cortices, we chose to evaluate how sensory input modifies aINS
connectivity dynamics. To this end, we studied the same participants under eyes closed and eyes open

scanning conditions. Compared to the eyes open condition, participants with eyes closed spent
significantly less time in the task-control network Mode 1 but more time in the primary sensory anti-
correlated Mode 2. This finding is consistent with recent static connectivity studies revealing occipital
and sensorimotor functional changes in eyes closed versus open scanning conditions (56, 57) and
suggests that environmental conditions impact time-varying aINS connectivity by shifting the major
modes occupied. Therefore, a measurable component of time-varying connectivity may reflect external
cues such as conditions of eyes closure (38, 58), mental states (59, 60), or internally driven physiological
conditions of the body (12). Finally, it is tempting to speculate that aINS occupancy of the visually
anticorrelated Mode 2 might reflect levels of drowsiness and fluctuations in alertness. Intriguingly, a
recent study investigated wakefulness fluctuations as a source of time-varying functional connectivity by
combining simultaneously acquired tf-fMRI and EEG data (61). This study revealed progressive whole-
brain hypoconnectivity during deeper sleep stages (N2 and N3) when compared to wakefulness, raising
the possibility that the increased aINS fractional occupancy in Mode 2 during eyes closed is related to
reduced levels of alertness and wakefulness.

"Trait" characteristics of aINS time-varying functional connectivity modes

To the extent that modes and profiles of time-varying aINS functional connectivity represent traits,
individuals should show stability of these features over time. To assess this stability, we studied a
longitudinal dataset of cognitively healthy older adults. The four aINS modes were stable over an
average of 9 months in this sample, whether assessed by comparing fractional occupancy at the group
level via parametric tests or at the individual level via cosine similarity. These findings converge with
several recent studies showing that individual topologies in static functional brain organization are
highly stable across time and show unique features with promise for use in precision functional mapping
of individual human brains (62–65). In line with previous work associating aINS function with social-
emotional functions such as empathy (15, 66, 67), we found that the time aINS subregions cohesively
spent in the task-control Mode 1 was positively associated with personal distress, while the time aINS
subregions spent in the salience Mode 4 was positively associated with the fantasy score from the
Interpersonal Reactivity Index. Caution is advised, however, in the interpretation of these findings since
the statistical associations were modest and did not survive correction for the four models used to
assess these relationships. Nevertheless, the identified associations were specific to the aINS (compared
to the left IFG) and survived when removing outliers, adding tentative support for the trait
characteristics of aINS time-varying functional connectivity. Nonetheless, the strongest evidence for a
trait component to the identified modes remains their reproducibility in individual subjects. The
personal distress subscale measures "self-oriented" feelings of personal anxiety and unease in tense
interpersonal settings. The level of personal distress was associated with greater time spent in a mode
characterized by dorsal cingulo-opercular regions overlapping with the task-control network and anti-
correlations to the default mode network, in line with findings associating connectivity of the anterior
cingulate cortex and insula with pre-scan anxiety levels, dispositional anxiety, and affective symptoms in
mood disorders (4, 20, 68). The fantasy score reflects the capacity of participants to transpose
themselves imaginatively into the feelings and actions of fictitious characters in books, movies, and
plays. This capacity requires a high level of social contextualization and involves semantic processes
typically associated with regions of the semantic appraisal network, such as the temporal lobes and
ventro-medial prefrontal areas (13, 18, 54), the same regions contributing to the salience Mode 4.
Importantly, fractional occupancies of time-varying functional connectivity modes derived from the left

IFG were not significantly associated with measures of dispositional empathy, suggesting a degree of
specificity of these findings to the aINS.

Limitations and future directions

Although time-varying functional connectivity analyses have been of increasing interest to the human
brain mapping community, consensus is still lacking about whether time-varying fluctuations in BOLD
signal coherence reflect brain physiology or are mainly due to noise stochastically occurring in the
scanner between the acquisition of distinct brain volumes (69). One limitation of this work relates to the
use of hard clustering techniques such as k-means to extract modes of time-varying functional
connectivity, since distinct modes are likely to reflect extremes in time-varying connectivity gradients
rather than discrete clusters of topographical connectivity. Therefore, the ideal number of derived time-
varying connectivity modes may vary depending on the temporal and spatial resolution of the used
dataset, on the investigated anatomical region, and on the specific question of the researcher. We make
no claim that the aINS transitions between only four modes, but since we were interested in large-scale
time-varying connectivity patterns, we deemed it impractical to use an overly fine-grained mode
decomposition by applying higher clustering solutions. Further, the sliding-window approach has been
scrutinized because of its moderate reliability (70) and since its output is prone to influence by head
movement, sliding-window length, and other methodical choices (71–73), such as global signal
regression. In our study, however, we did not find any significant association between time spent by
distinct aINS subregions in a specific mode and summary measures of head movement, as exemplified
by the regression analyses associating fractional occupancy and dispositional empathy, which were
corrected for mean frame-wise head displacement (42, 74). Importantly, our control analyses using the
longitudinal dataset preprocessed with global signal regression resulted in the identification of time-
varying functional connectivity modes that highly resembled the time-varying modes derived using data
without global signal regression (Supplementary results and Figure S6). The only exception was for the
primary sensory hyperconnected Mode 3, which was not identified in three out of four aINS subregions,
suggesting that this mode is highly modulated by the global signal. A recent report suggested that noise
sources (for instance motion and respiration) can have a temporally lagged effect on the BOLD signal,
which is greatly reduced by global signal regression (72). However, a recent study investigating time-
varying functional connectivity in mice reliably mapped distinct time-varying patterns of tf-fMRI co-
activation that occurred at specific phases of global signal fluctuations (75). These findings suggest that
the global signal may partially relate to oscillatory cycling of slowly propagating neural activity, as
observed in lag threads of propagated tf-fMRI signal in the human brain (76) and transient variation in
calcium co-activation patterns in mice (77). Slow oscillation in the global signal have been proposed to
coordinate fluctuating periods of brain network topologies characterized by heightened global
integration and shifts in arousal (30, 78).

Finally, the limitations in temporal resolution affecting the sliding-window approach may also explain
the inability to detect mode-specific fractional occupancy differences between the left and right aINS,
although both regions play a distinct role in autonomic outflow and behavior (11, 12). More
sophisticated methods that preserve the temporal richness of tf-fMRI data, such as hidden Markov
models and Topological Data Analysis (28, 31), may help elucidate lateralized differences in time-varying
connectivity of the left and right aINS. However, the detection of static functional connectivity
heterogeneity informed by time-varying fractional occupancy profiles and the reproducibility of our
findings across distinct preprocessing pipelines, sliding-window lengths, clustering choices, samples, and

time points (within individuals)(79) increase confidence in the biological relevance of findings. In the
future, our approach could be extended to analyze time-varying functional connectivity of other brain
regions, or to assess time-varying fractional occupancy profiles in psychiatric and neurological disorders
characterized by aINS dysfunction.

Acknowledgements
We thank the participants for their contributions to neuroscience research.

Conflicts of interest
The authors declare no competing conflicts of interest.

Data availability
The data that support the findings of this study are available on request from the corresponding author
W.W.S. The data are not publicly available due to them containing information that could compromise
the privacy of participants in the study.

Author's contributions
L.P and W.W.S conceived the study; all authors designed the experiments; L.P implemented the analysis
of the data; L.P. and W.W.S. wrote the manuscript; and all the authors edited the manuscript.

Funding
This work was supported by the Larry L. Hillblom Foundation grants 2014-A-004-NET and 2018-A-025-
FEL, The Bluefield Project to Cure FTD, the Tau Consortium, and NIH grants R01AG032289,
R01AG048234, and UCSF ADRC P50 AG023501.

References

1.   Craig AD (2009) How do you feel - now? The anterior insula and human awareness. *Nat Rev*
Neurosci 10(1):59–70.

2.   Smith SM, et al. (2009) Correspondence of the brain's functional architecture during activation
and rest. *Proc Natl Acad Sci U S A* 106(31):13040–5.

3.   Fox MD, et al. (2005) The human brain is intrinsically organized into dynamic , anticorrelated
functional networks. *Proc Natl Acad Sci U S A* 02(27):9673-8

4.   Seeley WW, et al. (2007) Dissociable intrinsic connectivity networks for salience processing and
executive control. *J Neurosci* 27(9):2349–2356.

5.   Deen B, Pitskel NB, Pelphrey KA (2011) Three systems of insular functional connectivity identified
with cluster analysis. *Cereb Cortex* 21(7):1498–1506.

6.   Kurth F, Zilles K, Fox PT, Laird AR, Eickhoff SB (2010) A link between the systems: functional
differentiation and integration within the human insula revealed by meta-analysis. *Brain Struct*
Funct 214:1–16.

7.   Touroutoglou A, Hollenbeck M, Dickerson BC, Feldman Barrett L (2012) Dissociable large-scale
networks anchored in the right anterior insula subserve affective experience and attention.
Neuroimage 60(4):1947–1958.

8.   Uddin LQ (2014) Salience processing and insular cortical function and dysfunction. *Nat Rev*

Neurosci 16(1):55–61.

9.   Critchley HD (2005) Neural mechanisms of autonomic, affective, and cognitive integration. *J*
617   *Comp Neurol* 493(1):154–166.

618   10.   Critchley HD, Harrison NA (2013) Visceral Influences on Brain and Behavior. *Neuron* 77(4):624–
619   638.

620   11.   Sturm VE, et al. (2018) Network architecture underlying basal autonomic outflow: Evidence from
621   frontotemporal dementia. *J Neurosci* (415):0347–18.

622   12.   Guo CC, et al. (2016) Dominant hemisphere lateralization of cortical parasympathetic control as
623   revealed by frontotemporal dementia. *Proc Natl Acad Sci*:201509184.

624   13.   Zhou J, Seeley WW (2014) Network Dysfunction in Alzheimer ' s Disease and Frontotemporal
625   Dementia⬚: Implications for Psychiatry. *Biol Psychiatry* 75(7):565–573.

626   14.   Dosenbach NUF, et al. (2006) A Core System for the Implementation of Task Sets. *Neuron*
627   50(5):799–812.

628   15.   Nomi JS, et al. (2016) Dynamic functional network connectivity reveals unique and overlapping
629   profiles of insula subdivisions. *Hum Brain Mapp* 37(5):1770–1787.

630   16.   Buckner RL, Andrews-Hanna JR, Schacter DL (2008) The brain's default network: Anatomy,
631   function, and relevance to disease. *Ann N Y Acad Sci* 1124:1–38.

632   17.   Menon V, Uddin LQ (2010) Saliency, switching, attention and control: a network model of insula
633   function. *Brain Struct Funct*:1–13.

634   18.   Seeley WW, Zhou J, Kim E-J (2012) Frontotemporal Dementia: What Can the Behavioral Variant
635   Teach Us about Human Brain Organization? *Neurosci* 18(4):373–385.

636   19.   Brandl F, et al. (2019) Specific Substantial Dysconnectivity in Schizophrenia: A Transdiagnostic
637   Multimodal Meta-analysis of Resting-State Functional and Structural Magnetic Resonance
638   Imaging Studies. *Biol Psychiatry* (10):1–11.

639   20.   Williams LM (2016) Precision psychiatry: A neural circuit taxonomy for depression and anxiety.
640   *The Lancet Psychiatry* 3(5):472–480.

641   21.   Sporns O, Tononi G, Kötter R (2005) The Human Connectome: A Structural Description of the
642   Human Brain. *PLoS Comput Biol* 1(4):e42.

643   22.   Lurie DJ, et al. (2018) On the nature of resting fMRI and time-varying connectivity. *Preprint*:1–33.
644   DOI: 10.31234/osf.io/xtzre

645   23.   Preti MG, Bolton TA, Van De Ville D (2017) The dynamic functional connectome: State-of-the-art
646   and perspectives. *Neuroimage* 160(December 2016):41–54.

647   24.   Allen E, Damaraju E, Plis S, Erhardt E (2014) Tracking whole-brain connectivity dynamics in the
648   resting state. *Cerebral. Cortex* 24(3):663-76.

649   25.   Damaraju E, et al. (2014) Dynamic functional connectivity analysis reveals transient states of
650   dysconnectivity in schizophrenia. *NeuroImage Clin* 5:298–308.

651   26.   Kaiser RH, et al. (2016) Dynamic Resting-State Functional Connectivity in Major Depression.

652   *Neuropsychopharmacology* 41(7):1822–1830.

653   27.   Sourty M, Thoraval L, Roquet D (2016) Identifying dynamic functional connectivity changes in
654   dementia with lewy bodies based on product hidden markov models. *Front Comput Neurosc*
655   10:60

656   28.   Saggar M, Sporns O, Gonzalez-castillo J, Bandettini PA, Carlsson G (2018) Towards a new
657   approach to reveal dynamical organization of the brain using topological data analysis. *Nature*
658   *Communication* 9:1–33

659   29.   Madhyastha TM, Askren MK, Boord P, Grabowski TJ (2015) Tara M. Madhyastha, Mary K. Askren,
660   Peter Boord, and Thomas J. Grabowski 1,2. 5(1):45–59.

661   30.   Shine JM, et al. (2016) The Dynamics of Functional Brain Networks: Integrated Network States
662   during Cognitive Task Performance. *Neuron* 92(2):544–554.

663   31.   Vidaurre D, Smith SM, Woolrich MW (2017) Brain network dynamics are hierarchically organized
664   in time. *Proc Natl Acad Sci*:201705120.

665   32.   Résibois M, et al. (2017) The neural basis of emotions varies over time: different regions go with
666   onset- and offset-bound processes underlying emotion intensity. *Soc Cogn Affect Neurosci*
667   12(8):1261–1271.

668   33.   Tobia MJ, Hayashi K, Ballard G, Gotlib IH, Waugh CE (2017) Dynamic functional connectivity and
669   individual differences in emotions during social stress. *Hum Brain Mapp* 38(12):6185–6205.

670   34.   Sadaghiani S, Poline J, Kleinschmidt A, Esposito MD (2015) Ongoing dynamics in large-scale
671   functional connectivity predict perception. *Proc Natl Acad Sci* 112(27):8463–8468.

672   35.   Ekman M, Derrfuss J, Tittgemeyer M, Fiebach CJ (2012) Predicting errors from reconfiguration
673   patterns in human brain networks. *Proc Natl Acad Sci* 109(41):16714–16719.

674   36.   Morris JC (1997) Clinical Dementia Rating: A reliable and valid diagnostic and staging measure for
675   dementia of the Alzheimer type. *Int Psychogeriatrics* 9(SUPPL. 1):173–176.

676   37.   Tombaugh TN, McIntyre NJ (1992) The mini-mental state examination: a comprehensive review. *J*
677   *Am Geriatr Soc* 40(9):922–35.

678   38.   Riedl V, et al. (2016) Metabolic connectivity mapping reveals effective connectivity in the resting
679   human brain. *Proc Natl Acad Sci* 113(8):6260–6266

680   39.   Davis MHA (1980) A Multidimensional Approach to Individual Differences in Empathy. *JSAS*
681   *Catalog of Selected Documents in Psychology* 10:85.

682   40.   Rankin KP, et al. (2006) Structural anatomy of empathy in neurodegenerative disease. *Brain*
683   129(11):2945–2956.

684   41.   Nana AL, et al. (2018) Neurons selectively targeted in frontotemporal dementia reveal early stage
685   TDP-43 pathobiology. *Acta Neuropathol* 137(1):27–46.

686   42.   Satterthwaite TD, et al. (2013) An improved framework for confound regression and filtering for
687   control of motion artifact in the preprocessing of resting-state functional connectivity data.
688   *Neuroimage* 64(1):240–256.

43.   Gorno-Tempini ML (2004) Cognition and Anatomy in Three Variants of Primary Progressive

Aphasia. *Ann Neurol* 119(11):2658–2666.

44.   Hutchison RM, et al. (2013) Dynamic functional connectivity: Promise, issues, and
interpretations. *Neuroimage* 80:360–378.

45.   Leonardi N, Van De Ville D (2015) On spurious and real fluctuations of dynamic functional
connectivity during rest. *Neuroimage* 104:430–436.

46.   Shirer WR, Ryali S, Rykhlevskaia E, Menon V, Greicius MD (2012) Decoding subject-driven
cognitive states with whole-brain connectivity patterns. *Cereb Cortex* 22(1):158–165.

47.   Wilson RS, et al. (2015) Influence of epoch length on measurement of dynamic functional
connectivity in wakefulness and behavioural validation in sleep. *Neuroimage* 112:169–179.

48.   Power JD, et al. (2011) Article Functional Network Organization of the Human Brain. *Neuron*
72(4):665–678.

49.   Dosenbach NUF, et al. (2007) Distinct brain networks for adaptive and stable task control in
humans. *Proc Natl Acad Sci* 104(26):11073–11078.

50.   Seeley WW, Crawford RK, Zhou J, Miller BL, Greicius MD (2009) Neurodegenerative Diseases
Target Large-Scale Human Brain Networks. *Neuron* 62(1):42–52.

51.   Toller G, et al. (2018) Individual differences in socioemotional sensitivity are an index of salience
network function. *Cortex* (March):1–13.

52.   Friederici AD, Gierhan SM (2013) The language network. *Curr Opin Neurobiol* 23(2):250–254.

53.   Sadaghiani S, D'Esposito M (2015) Functional Characterization of the Cingulo-Opercular Network
in the Maintenance of Tonic Alertness. *Cereb Cortex* 25(9):2763–2773.

54.   Guo CC, et al. (2013) Anterior temporal lobe degeneration produces widespread network-driven
dysfunction. *Brain* 136(10):2979–2991.

55.   Hickok G, Poeppel D (2007) The cortical organization of speech processing. *Nat Rev Neurosci*
8(5):393–402.

56.   Wei J, et al. (2018) Eyes-Open and Eyes-Closed Resting States With Opposite Brain Activity in
Sensorimotor and Occipital Regions: Multidimensional Evidences From Machine Learning
Perspective. *Front Hum Neurosci* 12:422.

57.   Agcaoglu O, Wilson T, Wang Y, Stephen J, Calhoun VD (2019) Resting state connectivity
differences in eyes open versus eyes closed conditions. *Hum Brain Mapp* 40(8):2488-2498.

58.   Hahn A, et al. (2018) Task-relevant brain networks identified with simultaneous PET/MR imaging
of metabolism and connectivity. *Brain Struct Funct* 223(3):1369–1378.

59.   Shine JM, Koyejo O, Poldrack RA (2016) Temporal metastates are associated with differential
patterns of time-resolved connectivity, network topology, and attention. *Proc Natl Acad Sci U S A*
113(35):9888–91.

60.   Kucyi A (2018) Just a thought: How mind-wandering is represented in dynamic brain connectivity.
Neuroimage 180:505–514.

61.   Haimovici A, Tagliazucchi E, Balenzuela P, Laufs H (2017) On wakefulness fluctuations as a source

of BOLD functional connectivity dynamics. *Sci Rep* 7(1):5908.

62.   Gordon EM, et al. (2017) Precision Functional Mapping of Individual Human NeuroResource
729   Precision Functional Mapping of Individual Human Brains. *Neuron* 95(4):791-807.

730   63.   Laumann TO, et al. (2015) Functional System and Areal Organization of a Highly Sampled
731   Individual Human Brain. *Neuron* 87(3):657–670.

732   64.   Poldrack RA, et al. (2015) Long-term neural and physiological phenotyping of a single human. *Nat*
733   *Commun* 6:8885.

734   65.   Guo CC, et al. (2012) One-year test–retest reliability of intrinsic connectivity network fMRI in
735   older adults. *Neuroimage* 61(4):1471–1483.

736   66.   Gu X, et al. (2012) Anterior insular cortex is necessary for empathetic pain perception. *Brain*
737   135(9):2726–2735.

738   67.   Leigh R, et al. (2013) Acute lesions that impair affective empathy. *Brain* 136(8):2539–2549.

739   68.   Menon V (2011) Large-scale brain networks and psychopathology: A unifying triple network
740   model. *Trends Cogn Sci* 15(10):483–506.

741   69.   Laumann TO, et al. (2017) On the Stability of BOLD fMRI Correlations. *Cereb Cortex* 27(10):4719-
742   4732.

743   70.   Choe AS, et al. (2017) NeuroImage Comparing test-retest reliability of dynamic functional
744   connectivity methods. 158:155–175.

745   71.   Hindriks R, et al. (2016) Can sliding-window correlations reveal dynamic functional connectivity in
746   resting-state fMRI? *Neuroimage* 127:242–256.

747   72.   Byrge L, Kennedy DP (2017) Identifying and characterizing systematic temporally-lagged BOLD
748   artifacts. *Neuroimage* 171:376–392.

749   73.   Nalci A, Rao BD, Liu TT (2018) Nuisance Effects and the Limitations of Nuisance Regression in
750   Dynamic Functional Connectivity fMRI. *bioRxiv*: doi.org/10.1101/285239

751   74.   Power JD, et al. (2014) NeuroImage Methods to detect , characterize , and remove motion
752   artifact in resting state fMRI. *Neuroimage* 84:320–341.

753   75.   Gutierrez-Barragan D, Basson MA, Panzeri S, Gozzi A (2018) Oscillatory brain states govern
754   spontaneous fMRI network dynamics. *bioRxiv*: doi.org/10.1101/393389.

755   76.   Mitra A, Snyder AZ, Constantino JN, Raichle ME (2015) The Lag Structure of Intrinsic Activity is
756   Focally Altered in High Functioning Adults with Autism. *Cereb Cortex* 27(2):1083-1093.

757   77.   Matsui T, Murakami T, Ohki K (2016) Transient neuronal coactivations embedded in globally
758   propagating waves underlie resting-state functional connectivity. *Proc Natl Acad Sci*
759   113(23):6556–6561.

760   78.   Liu X, et al. Subcortical evidence for a contribution of arousal to fMRI studies of brain activity. *Nat*
761   *Commun* (2018):1–10.

762   79.   Abrol A, et al. (2017) Replicability of time-varying connectivity patterns in large resting state fMRI
763   samples. *Neuroimage* 163:160–176.

771   **Tables, Figures and Legends**

773   **Table 1. Demographics and sample characteristics**

Dataset	Cross-sectional	Longitudinal	Eyes closed/open
Number of subjects	121	44	20
Age in years, mean (s.d.)	69.3 (3.6)	72.9 (7.6)	42.8 (6.4)
Sex (Female/Male)	73/48	25/19	9/11
Handedness (L/A/R)	0/0/121	6/0/38	0/0/20
Education in years, mean (s.d.)	17.6 (2.0)	17.5 (1.9)	14.4 (2.6)
IRI-EC, mean (s.d.)	27.4 (4.8)	NA	NA
IRI-PT, mean (s.d.)	24.3 (5.7)	NA	NA
IRI-PD, mean (s.d.)	13.1 (5.0)	NA	NA
IRI-FS, mean (s.d.)	18.3 (5.1)	NA	NA
Interval between scanning dates in months, mean (s.d.)	NA	9.2 (2.7)	1.0 (0.2)

774   A = ambidextrous; EC = empathic concern; FS = fantasy score; IRI = Interpersonal Reactivity Index; L =
775   left; NA = not applicable; PD = personal distress; PT = perspective taking; R = right; s.d. = standard
776   deviation

Figure 1. Time-varying functional connectivity pipeline. (A) For each individual S*i*, average BOLD activity time courses were extracted from subregions of the aINS, here the left ventral. **(B)** For each individual, 218 time-varying seed-to-whole-brain connectivity maps (S_iW_1-S_iW_{218}) were generated using a sliding-windows of 18 TRs (36 s) in steps of 1 TR. **(C)** The derived connectivity windows were vectorized and concatenated across subjects resulting in windows x voxels matrices for each aINS seed. K-means clustering was then applied to the concatenated window matrices using a value of k = 4 to produce four clusters representing four time-varying aINS modes present across the course of the functional scan. **(D)** Group averaged maps of the four modes were generated using the cluster-specific centroid maps generated through k-means (threshold at -0.5 > z > 0.5). The assignment of each window to a specific mode, was used to derive fractional occupancy profiles for each mode (i.e. the time spent by an aINS subregion in each mode), defined as the number of windows assigned to one mode divided through the total number of generated windows. aINS = anterior insula; BOLD = blood oxygen level dependent signal

Figure 2. Time-varying functional connectivity modes of the aINS. (A) Spatial maps of the four time-varying functional connectivity modes identified across the four aINS subregions in the cross-sectional dataset. Threshold at -0.5 > z > 0.5, negative connectivity is depicted in blue, positive in red. The right side of the brain is shown on the right side of the image. **(B)** Correlation matrix reflecting the spatial similarity of modes identified in distinct aINS subregions. **(C)** Differences in time-varying connectivity between correspondent modes of the right dorsal and ventral aINS (threshold at -0.3 > z > 0.3, higher connectivity of the dorsal aINS is depicted in red, higher connectivity of the ventral aINS in blue).

Figure 3. Temporal characteristics of the identified aINS time varying functional connectivity modes.
(A) Violin plots reflecting group fractional occupancy in the four identified time-varying functional connectivity modes across the four aINS subregions. **(B)** Violin plots representing the number of transitions between time-varying functional connectivity modes across the four aINS subregions. **(C)** Violin plots representing the diversity of fractional occupancy profiles – measured using the Shannon diversity index – across aINS subregions. This measure reflects whether an aINS subregion tends to spend similar amounts of time in different modes or spends most of its time in only one mode. **(D)** Polar plots schematizing the fractional occupancies of aINS subregions into the four time-varying functional connectivity modes. On the left side, the fractional occupancy profile of Subject 47 transitions between several modes, and is characterized by a high number of transitions and by high Shannon diversity index averaged across aINS subregions. On the right side, we can appreciate the highly stereotyped fractional occupancy profile of Subject 22, with aINS subregions spending most of their time in a single mode. Regardless of the subregion, aINS subregions in this participant tend to spend time only in Mode 1, showing hence a low number of transitions and low Shannon diversity index averaged across aINS subregions. **(E)** Heat maps reflecting how specific modes are jointly occupied at the same time by distinct aINS subregions. Scale bar reflects the average group probability that distinct aINS subregions occupy certain mode configurations.

Figure 4. Time-varying functional connectivity modes across time. In the longitudinal data sample, spatial patterns of time-varying functional connectivity modes where identified across the four aINS subregions bearing high similarity to the modes identified in the cross-sectional sample (maps averaged across both time points, sagittal plane only panels **A-D**). Fractional occupancy in the identified modes did not significantly differ between the first (red violin plots) and second scanning dates (blue violin plots) ~9 months apart (paired t-test; p < 0.05 uncorrected for multiple comparisons).

Figure 5. Individual fractional occupancy of aINS subregions across time. (A) Violin plot representing the similarity of aINS fractional occupancy profiles across scanning dates assessed using cosine similarity. Subjects 10, 41, 21 and 26 are highlighted with red circles in the individual data plot. The individual fractional occupancy profiles of these subjects and correspondent cosine similarity values are further schematized in panels **B-E**, where the fractional occupancies are shown at the first (red) and second (blue) scanning dates. aINS fractional occupancy in Subjects 10 and 41 did not substantially differ between scanning dates (note the overlap in the polar plots and the high cosine similarity index), while Subjects 21 and 26 showed very different aINS fractional occupancy profiles when assessed at different time points (note the little overlap in the polar plot and the low cosine similarity index).

Figure 6. Time-varying functional connectivity modes during scanning with eyes closed versus eyes open. In the eyes closed/open dataset, spatial patterns of time-varying functional connectivity modes where identified across the left dorsal, right dorsal, left ventral, and right ventral aINS bearing high similarity to the modes identified in the cross-sectional sample (maps averaged across eyes closed/open conditions, sagittal plane only panels **A-D**). With exception of the right ventral aINS, fractional occupancy in the visually anticorrelated Mode 2 was significantly higher during eyes closed (red violin plots) than during eyes open scanning (blue violin plots). On the other hand, in the eyes open condition participants spent significantly more time in the task-control Mode 1 (paired t-test; p < 0.05 uncorrected for multiple comparisons). *p < 0.05; **p < 0.005

Figure 7. Meta-profiles of time-varying dynamic connectivity modes. (A) Individual fractional occupancy profiles of aINS subregions were clustered into four clusters, reflecting four meta-profiles of time-varying aINS functional connectivity. 26 subjects were clustered into Meta-profile 1 (triangles, green), 31 into Meta-profile 2 (+ sign, blue), 44 into Meta-profile 3 (circles, violet), and 20 into Meta-profile 4 (x's, red). Panels show individual fractional occupancy profiles representative of the identified meta-profiles (see black circles in the scatterplots). Across aINS subregions, Subject 22 tended to spend time in Mode 1, Subject 79 in Mode 2, Subject 80 in Mode 3, and Subject 88 in Mode 4. **(B)** When averaged across aINS subregions, one-way ANOVAs and related post-hoc t-tests reveal that the aINS of participants clustered in Meta-profile 1 showed significantly higher fractional occupancy in Mode 1.

859 Similarly, the aINS of participants clustered on Meta-profiles 2, 3, or 4, spent more time on Modes 2, 3,
860 or 4, respectively.