Rapid degradation of *C. elegans* proteins at single-cell resolution with a synthetic

auxin

Michael A. Q. Martinez*, Brian A. Kinney[†], Taylor N. Medwig-Kinney*, Guinevere Ashley[‡],

James M. Ragle[‡], Londen Johnson[‡], Joseph Aguilera[‡], Christopher M. Hammell[†], Jordan

D. Ward[‡], David Q. Matus*

*Department of Biochemistry and Cell Biology, Stony Brook University, Stony Brook, NY

11794, USA.

[†]Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724, USA.

[‡]Department of Molecular, Cell, and Developmental Biology, University of California-

Santa Cruz, Santa Cruz, CA 95064, USA.

20   Running title: Protein degradation via a synthetic auxin

22   Keywords: *C. elegans*; AID system; NAA; Microfluidics; SCF complex; NHR-25

24   Corresponding author: David Q. Matus; Department of Biochemistry and Cell Biology,

25   Stony Brook University, Stony Brook, NY 11794, USA; david.matus@stonybrook.edu

ABSTRACT

As developmental biologists in the age of genome editing, we now have access to an

ever-increasing array of tools to manipulate endogenous gene expression. The auxin-

inducible degradation system, allows for spatial and temporal control of protein

degradation, functioning through the activity of a hormone-inducible *Arabidopsis* F-box

protein, transport inhibitor response 1 (TIR1). In the presence of auxin, TIR1 serves as a

substrate recognition component of the E3 ubiquitin ligase complex SKP1-CUL1-F-box

(SCF), ubiquitinating auxin-inducible degron (AID)-tagged proteins for proteasomal

degradation. Here, we optimize the *Caenorhabditis elegans* AID method, utilizing 1-

naphthaleneacetic acid (NAA), an indole-free synthetic analog of the natural auxin indole-

3-acetic acid (IAA). We take advantage of the photostability of NAA to demonstrate via

quantitative high-resolution microscopy that rapid degradation of target proteins can be

detected in single cells within 30 minutes of exposure. Additionally, we show that NAA

works robustly in both standard growth media and physiological buffer. We also

demonstrate that K-NAA, the water-soluble, potassium salt of NAA, can be combined with

microfluidics for targeted protein degradation in *C. elegans* larvae. We provide insight into

how the AID system functions in *C. elegans* by determining that TIR1 interacts with *C.*

elegans SKR-1/2, CUL-1, and RBX-1 to degrade target proteins. Finally, we present

highly penetrant defects from NAA-mediated degradation of the Ftz-F1 nuclear hormone

receptor, NHR-25, during *C. elegans* uterine-vulval development. Together, this work

provides a conceptual improvement to the AID system for dissecting gene function at the

single-cell level during *C. elegans* development.

INTRODUCTION

In situ techniques for targeted protein degradation enable a detailed analysis of

developmental events, mechanisms, and functions. RNAi and Cre or Flp-mediated

recombination (Qadota *et al.* 2007; Hubbard 2014; Shen *et al.* 2014) allow tissue-specific

study of gene products, but the persistence of the target protein following recombination

or RNA depletion can delay the manifestation of an otherwise acute phenotype. Several

methods have been described recently to enable tissue-specific protein degradation in

Caenorhabditis elegans, including ZF1 tagging (Armenti *et al.* 2014), a GFP nanobody

approach (Wang *et al.* 2017), sortase A (Wu *et al.* 2017), and auxin-mediated degradation

(Zhang *et al.* 2015).

The auxin-inducible degradation system allows for rapid and conditional

degradation of auxin-inducible degron (AID)-tagged proteins in *C. elegans* as well as in

other commonly used model systems including yeast (Nishimura *et al.* 2009), *Drosophila*

(Trost *et al.* 2016), zebrafish (Daniel *et al.* 2018), cultured mammalian cells (Nishimura *et*

al. 2009; Holland *et al.* 2012; Natsume *et al.* 2016), and mouse oocytes (Camlin and

Evans 2019). This protein degradation system relies on the expression of an *Arabidopsis*

F-box protein called transport inhibitor response 1 (TIR1). As a substrate-recognition

component of the SKP1-CUL1-F-box (SCF) E3 ubiquitin ligase complex, TIR1 carries out

its function only in the presence of the hormone auxin. Once bound to auxin, TIR1 targets

AID-tagged proteins for ubiquitin-dependent proteasomal degradation (**Figure 1A**).

84   The *C. elegans* version of the AID system is robust and specific with minimal off-

85 target effects (Zhang *et al.* 2015). However, re-evaluation of the system is needed to

86 assess its utility among *C. elegans* researchers conducting microscopy-based single-cell

87 biology within a narrow developmental time frame. Here, we use 1-naphthaleneacetic

88 acid (NAA) and its water-soluble potassium salt analog (K-NAA), indole-free synthetic

89 analogs of the natural auxin indole-3-acetic acid (IAA), to degrade target proteins at

90 single-cell resolution in *C. elegans* larvae in standard growth media and physiological

91 buffer. Given the ability to solubilize K-NAA solely in water or physiological buffer (M9),

92 we also demonstrated rapid degradation kinetics of an AID-tagged transgene in a *C.*

93 *elegans*-based microfluidics for the first time (Keil *et al.* 2017). Next, we sought to gain

94 insight into which SCF complex members interact with TIR1, identifying *skr-1/2*, *cul-1* and

95 *rbx-1* as putative TIR1 interactors through RNAi depletion experiments. Finally, we

96 demonstrate potent temporal effects on uterine and vulval development following targeted

97 degradation of endogenous NHR-25, the single *C. elegans* homolog of *Drosophila* Ftz-

98 F1 and human SF-1 and LRH-1 (Chen *et al.* 2004; Ward *et al.* 2013). It is our hope that

99 this synthetic auxin analog will be applied at all stages of *C. elegans* development,

100 allowing for precise, rapid degradation of target proteins in a high-resolution and

101 quantitative fashion.

MATERIALS AND METHODS

C. elegans strains and culture conditions

Animals were maintained using standard culture conditions at 25°C (Brenner 1974) and

were synchronized through alkaline hypochlorite treatment of gravid adults to isolate eggs

(Porta-de-la-Riva *et al.* 2012). In the main text and figure legends, we designate linkage

to a promoter using a greater than symbol (>) and fusion to a protein using a double colon

(::). The following alleles and transgenes were used in this manuscript for experimental

purposes: LG I: *kry61[nhr-23::AID];* LG II: *ieSi57[eft-3>TIR1::mRuby];* LG IV: *ieSi58[eft-*

3>AID::GFP], syIs49 [zmp-1>GFP]; LG X: *wrd10[nhr-25::GFP::AID].*

Constructs and microinjection

SapTrap was used to construct the 30xlinker::GFP^SEC^TEV::AID degron::3xFLAG

repair template (pJW1747) for generating the knock-in into the 3' end of the *nhr-25* gene

(Schwartz and Jorgensen 2016). DH10β competent *E. coli* cells, made in-house, were

used for generating the plasmid. The following reagents were used to assemble the final

repair template: pDD379 (backbone with F+E sgRNA), annealed oligos 3482+3483

(sgRNA), pJW1779 (5' homology arm), 3' homology arm PCR product, pJW1347 (30x

linker for CT slot), pDD372 (GFP for FP slot), pDD363 (SEC with LoxP sites), and

pJW1759 (TEV::AID degron::3xFLAG for NT slot).

The pJW1747 repair template was purified using the Invitrogen PureLink HQ Mini

Plasmid DNA Purification Kit (K210001). The optional wash step in the protocol using a

4 M guanidine-HCl + 40% isopropanol solution is highly recommended, as excluding it

dramatically reduced injection efficiency in our hands. N2 animals were injected with a

mix consisting of 10 ng/µl of pJW1747, 50 ng/µl of pDD121 (Cas9 vector), and co-injection

markers (10 ng/µl pGH8, 5 ng/µl pCFJ104, 2.5 ng/µl pCFJ90) as previously described

(Frøkjær-Jensen *et al.* 2012; Dickinson *et al.* 2013, 2015). Knock-ins were isolated as

previously described (Dickinson *et al.* 2015). Each knock-in junction was verified via PCR

using a primer that bound outside the homology arm paired with a primer binding within

pJW1747. The knock-in was backcrossed five times against wild-type N2 animals to

produce JDW58. The SEC was then excised by heat-shock (Dickinson *et al.* 2015) to

produce JDW59; the knock-in sequence was re-confirmed by PCR amplification and

sequencing, using the oligos flanking the homology arms. JDW58 was crossed to CA1200

(*eft-3>TIR1::mRuby*) to generate JDW70. The SEC was then excised (Dickinson *et al.*

2015) to produce JDW71.

pDD121, pDD363, pDD372, and pDD379 (Dickinson *et al.* 2018) were gifts from

Bob Goldstein (Addgene plasmid numbers are 91833, 91829, 91824, and 91834,

respectively). pJW1347 and pJW1759 will be deposited into Addgene's repository and

are also available upon request. pJW1347 and pJW1759 were generated by TOPO blunt

cloning of PCR products. pJW1779 was generated by Gateway cloning into pDONR221

(Invitrogen). Oligo sequences used to generate these plasmids, the sgRNA, the 3'

homology arm, and for genotyping are in the Reagent Table.

Auxin experiments

For all auxin experiments, synchronized L1 larval stage animals were first transferred to

standard nematode growth media (NGM) agar plates seeded with *E. coli* OP50 and then

transferred at the P6.p 2-cell stage (mid-L3 stage) to either OP50-seeded NGM agar

plates treated with IAA, NAA, or K-NAA, or M9 buffer treated with NAA in the absence of

bacteria, NAA plus *E. coli* NA22, or K-NAA plus *E. coli* NA22.

For IAA and NAA experiments on plates, a 250 mM stock solution in 95% ethanol

was prepared using powder IAA purchased from Alfa Aesar (A10556) and powder NAA

purchased from Sigma-Aldrich (317918) and stored at -20°C. IAA and NAA were then

diluted into the NGM agar (cooled to approximately 50°C) at the time of pouring plates.

Fresh OP50 was used to seed plates. For control experiments, 0.25% ethanol was used

as described previously (Zhang *et al.* 2015). For K-NAA experiments on plates, a 250 mM

stock solution in deionized water was prepared using powder K-NAA purchased from

PhytoTechnology Laboratories (N610) and stored at 4°C. For control experiments, OP50-

seeded NGM agar plates were used. Prior to each NAA experiment in M9 buffer, a fresh

1 mM (pH of 7.22) or 4 mM solution (pH of 8.14) in M9 buffer was prepared using 5.4 mM

NAA purchased in liquid form from Sigma-Aldrich (N1641). These pH levels are well

within the tolerance range of *C. elegans* for pH (Khanna *et al.* 1997). M9 buffer alone (pH

of 7.13) was used as a control. A detailed protocol for liquid-based NAA-mediated

degradation can be found in File S1. For experiments conducted in the microfluidic

platform, a 4 mM NAA or K-NAA solution in M9 buffer containing *E. coli* NA22 was

prepared and stored at 4°C for up to 2 weeks. M9 buffer containing NA22 was used as a

control. See File S2 for a detailed protocol describing the preparation of media for the

microfluidic device.

Brood size and viability assays

Brood size and viability assays were performed as described (Zhang _et al._ 2015). Briefly,

L4 hermaphrodites were picked onto individual MYOB plates containing 0% ethanol (K-

NAA control), 0.25% ethanol (IAA control), 4 mM K-NAA, or 4 mM IAA. Animals were then

transferred to new plates daily over 4 days. The eggs laid on each plate were counted

after removing the parent and viable progeny were quantified when the F1 reached L4 or

adult stages (2-3 days post egg-laying). At this point, we also scored for dead eggs. Brood

size is the sum of live progeny and dead eggs. Percent embryonic lethality was

determined by dividing dead eggs by total eggs laid.

RNAi experiments

RNAi targeting _cul-1_ was constructed by cloning 997 bp of synthetic DNA based on its

cDNA sequence available on WormBase (wormbase.org) into the highly efficient T444T

RNAi vector (Sturm _et al._ 2018). The synthetic DNA was generated by Integrated DNA

Technologies (IDT) as a gBlock gene fragment and cloned into the BglII/SalI restriction

digested T444T vector using the NEBuilder HiFi DNA Assembly Master Mix (E2621).

RNAi feeding strains silencing _skr-1/2_, _skr-7_, _skr-10_, and _rbx-1_ were obtained from the

Vidal RNAi library (Rual _et al._ 2004).

Scoring defects in anchor cell (AC) specification

Synchronized L1 stage *nhr-25::GFP::AID; eft-3*>TIR1::mRuby animals were plated onto

NGM agar plates containing either control or 4 mM NAA and grown for 24 hours at 25°C

until the early L3 stage (P6.p 1-cell stage), after the normal time of AC specification.

Images were acquired as specified below to score for the presence or absence of an AC,

visualized by characteristic morphology using DIC optics.

Scoring vulva precursor cell (VPC) arrest

Synchronized L1 stage *nhr-25::GFP::AID; eft-3*>TIR1::mRuby animals were plated onto

OP50 NGM agar plates and allowed to grow until the P6.p 1-cell stage. Animals were

then washed off plates with M9 and transferred onto NGM agar plates containing either

control or 4 mM NAA and grown at 25°C until the mid-L3 stage, after the normal time of

P6.p cell division. Images were acquired as specified below to score for P6.p divisions

using DIC optics. Remaining animals were scored for plate level adult phenotypes

approximately 24 hours later.

Image acquisition

Images were acquired using a Hamamatsu Orca EMCCD camera and a Borealis-

modified Yokagawa CSU-10 spinning disk confocal microscope (Nobska Imaging, Inc.)

with a Plan-APOCHROMAT x 100/1.4 oil DIC objective controlled by MetaMorph software

(version: 7.8.12.0). Animals were anesthetized on 5% agarose pads containing 10 mM

sodium azide and secured with a coverslip. Imaging on the microfluidic device was

performed on a Zeiss AXIO Observer.Z7 inverted microscope using a 40X glycerol

immersion objective and DIC and GFP filters controlled by ZEN software (version 2.5).

Images were captured using a Hamamatsu C11440 digital camera. For scoring plate level

phenotypes, images were acquired using a Moticam CMOS (Motic) camera attached to

a Zeiss dissecting microscope.

Image processing and analyses

All acquired images were processed using Fiji software (version: 2.0.0-rc-69/1.52p)

(Schindelin *et al.* 2012). To quantify AC- or VPC-specific degradation of AID::GFP,

images were captured at the P6.p 2-cell stage and 4-cell stage (mid-L3 stage) at time

points 0, 30, 60, 90, and 120 minutes in the absence or presence of auxin. Expression of

eft-3>AID::GFP was quantified by measuring the mean fluorescence intensity (MFI) of

ACs and VPCs subtracted by the MFI of a background region in the image to account for

camera noise. Cells were outlined using the freehand selection tool in Fiji. Data were

normalized by dividing the MFI in treated or untreated animals at time points 30, 60, 90,

and 120 minutes by the average MFI in untreated animals at 0 minutes. For experiments

utilizing RNAi, only ACs were measured due to the variable sensitivity of VPCs to RNAi

(Bourdages *et al.* 2014; Matus *et al.* 2014). To quantify AC-specific degradation of

AID::GFP in animals fed RNAi overnight, images were captured at the P6.p 2-cell stage

before auxin treatment and 60 minutes post-treatment. *eft-3*>AID::GFP expression in the

AC was quantified as described above. Data were normalized by dividing the MFI in auxin

treated animals by the average MFI in untreated animals. To analyze *nhr-25::GFP::AID*

degradation, GFP levels were quantified by measuring the MFI in individual GFP-

expressing nuclei in the AC/VU, AC, or VPCs subtracted by the MFI of a background

region in the image to account for background noise. Nuclei were outlined using the

threshold tool in Fiji or for animals with no detectable GFP signal, the corresponding DIC

image was utilized to identify the nucleus. Images of L3 larvae were captured in a *C.*

elegans larvae-specific microfluidic device (Keil *et al.* 2017). To quantify AID::GFP

degradation, animals were loaded into the microfluidic chamber and fed NA22 bacteria.

Images were captured at time points 0, 30, 60, 90, and 120 minutes with or without auxin.

Here, *eft-3*>AID::GFP expression was quantified by measuring the MFI in whole animals

subtracted by the MFI of a background region in the image to account for background

noise. Whole animals were outlined using the freehand selection tool in Fiji. Data were

normalized by dividing the MFI in treated or untreated animals at time points 30, 60, 90,

and 120 minutes by the average MFI in untreated animals at timepoints 30, 60, 90, and

120 minutes respectively to account for photobleaching from imaging the same animal.

Cartoons were created with BioRender (biorender.com) and ChemDraw software

(version: 18.0). Graphs were generated using Prism software (version: 8.1.2). Figures

were compiled using Adobe Photoshop (version: 20.0.6) and Illustrator (version: 23.0.26).

Statistical analyses

A power analysis was performed to determine the sample size (*n*) needed per experiment

to achieve a power level of 0.80 or greater (Cohen 1992; Pollard *et al.* 2019). Statistical

significance was determined using either a two-tailed unpaired Student's t-test or Mann

Whitney U test. *P* < 0.05 was considered statistically significant. The figure legends

specify when error bars represent the standard deviation (SD) or interquartile range

(IQR).

Data availability

Supplemental data and key reagents can be found at Figshare. Worm strains CA1202,

CA1204, and PS3239 are available to order from the *Caenorhabditis* Genetics Center. All

other strains are available upon request. The data that support the findings of this study

are available upon reasonable request.

RESULTS AND DISCUSSION

NAA is a synthetic alternative to the natural auxin IAA

Given the recent advances in CRISPR/Cas9-genome editing technology (Dickinson and

Goldstein 2016; Dokshin *et al.* 2018), the auxin-inducible degron (AID) with or without a

fluorescent reporter (e.g., GFP or its derivatives) can be inserted into a genomic locus of

interest (Röth *et al.* 2019). Though this technology can be applied with ease, there are

certain limitations that exist with the use of the natural auxin indole-3-acetic acid (IAA),

including its limited solubility in water. While levels of the ethanol solvent used to dissolve

IAA (0.25–1.52%) are well below the threshold for causing a physiologic response

(Morgan and Sedensky 1995; Kwon *et al.* 2004), higher percentages of ethanol (7%) have

been shown to cause rapid changes in *C. elegans* gene expression (Kwon *et al.* 2004).

A potentially more problematic limitation of IAA for live-cell imaging-based applications is

cytotoxicity related to excitation with UV and blue light. Specifically, IAA has been shown

in yeast (Papagiannakis *et al.* 2017) and mouse oocytes (Camlin and Evans 2019) to

cause cytotoxicity, likely due to acceleration of the oxidative decarboxylation of IAA to

methylene-oxindole (Srivastava 2002). In yeast, IAA exposure during live-cell imaging

suppressed cell proliferation (Papagiannakis *et al.* 2017) and mammalian oocytes failed

to complete meiotic maturation (Camlin and Evans 2019). In both systems, the use of a

synthetic auxin, 1-naphthaleneacetic acid (NAA), rescued these cytotoxic responses. For

these reasons, we chose to examine whether NAA would also function in *C. elegans* to

degrade AID-tagged proteins in the presence of TIR1. Ultimately, we wished to evaluate

AID-mediated degradation (**Figure 1A**) in single cells and tissues using live-cell imaging.

Thus, we determined the kinetics of protein degradation using spinning disk confocal

microscopy, rather than using low-magnification microscopy and Western blot analysis to

measure protein loss in whole animals, as performed previously (Zhang *et al.* 2015). We

chose to focus primarily on the L3 stage of post-embryonic development due to many of

the dynamic cellular behaviors occurring over relatively short time scales (within minutes

to hours) in this developmental window, including uterine-vulval attachment and vulval

morphogenesis (**Figure 1B**) (Gupta *et al.* 2012).

To analyze the kinetics of AID-mediated degradation in the uterine anchor cell (AC)

and underlying vulva precursor cells (VPCs) (**Figure 1B**), we utilized a previously

published strain expressing AID::GFP and TIR1::mRuby under the same ubiquitously

expressed *eft-3* promoter (Zhang *et al.* 2015). The single-cell abundance of AID::GFP

was measured over time in mid-L3 stage animals exposed to different concentrations of

auxin incorporated into standard *C. elegans* solid culture media (**Figure 2A**). In addition

to testing the natural auxin IAA, also tested whether it was possible to perform auxin-

inducible degradation in the AC and VPCs using the synthetic auxin analog NAA (**Figure**

2B). In the presence of ≥ 1 mM IAA or NAA, AID::GFP abundance in the AC and VPCs

was reduced by approximately 80% of its initial level within 30 minutes (**Figure 2C-D**).

Within 60 minutes, AID::GFP was virtually undetectable (**Figure 2C-D**). These results

indicate that NAA can serve as a viable substitute to IAA. These results indicate that NAA

can serve as a viable substitute for IAA. Similar to (Zhang *et al.* 2015), we observed that

growth on IAA resulted in similar brood sizes compared to control (**Table S1**). Growth on

the potassium salt of NAA (K-NAA) resulted in similar brood sizes compared to control,

and also produced a modest but significant (P = 0.02) reduction in embryonic lethality

compared to IAA treatment (**Table S1**). Consistent with this result, higher levels of toxicity

was observed when using IAA over NAA in studies investigating circadian rhythm biology

in *Drosophila* (Chen *et al.* 2018). Zhang *et al.* (2015) reported inhibited bacterial growth

at high concentrations of auxin. Compared to bacterial growth on IAA and NAA, we

observed more robust OP50 growth on K-NAA with no trade-off in degradation rate

(**Figure S1A).** Together, these results demonstrate that NAA is a viable alternative to IAA

for targeted protein degradation in *C. elegans* larvae.

NAA is soluble in physiological buffer

To compare degradation kinetics on plate-based growth to depletion in liquid culture

during imaging, we measured AID::GFP abundance in the AC and VPCs in mid-L3 stage

animals exposed to different concentrations of NAA solubilized in M9 buffer (**Figure 3A-**

D). In the presence of ≥ 1 mM liquid NAA, AID::GFP in the AC and VPCs was reduced by

80%, as compared to initial levels, within 30 minutes and was nearly undetectable within

60 minutes (**Figure 3B-D**). These results show that NAA can induce auxin-dependent

degradation in liquid culture in *C. elegans*, reducing the need to rear animals on auxin

plates and transfer to slides for imaging. This finding raises the possibility of depleting

proteins and imaging the resulting developmental consequences at single-cell resolution.

K-NAA is an option for C. elegans researchers employing microfluidics

The ability to easily solubilize NAA in physiological buffer raises the possibility of

performing protein degradation experiments paired with microfluidics where individual

animals can be imaged over long periods at cellular resolution. *C. elegans* lifespan and

behavioral assays can involve, subtle phenotypes sensitive to environmental

perturbations. Accordingly, the low level of ethanol present in IAA plates is not optimal for

these types of assays. The reduced bacterial growth on ethanol and IAA could also affect

nutrition (**Figure S1A**) (Cabreiro *et al.* 2013), making water soluble K-NAA an attractive

alternative. To compare degradation kinetics between NAA and K-NAA, we time-lapsed

L3 stage animals using a microfluidic device optimized for long-term imaging of *C.*

elegans larvae (Keil *et al.* 2017), assessing depletion of ubiquitously expressed AID::GFP

in trapped animals (**Figure 3A**). At the L3 stage, animals were loaded into the microfluidic

chamber in M9 and flushed with a mixture of M9, 4 mM NAA or K-NAA, and NA22 *E. coli*

as a bacterial food source (Keil *et al.* 2017). The animals were imaged every 30 minutes

for 2 hours. During image acquisition, animals were temporarily immobilized by manually

increasing the negative pressure on the compression layer of the device (Keil *et al.* 2017).

Between timepoints, animals were allowed to move and feed freely in 4 mM NAA or K-

NAA combined with NA22 in M9. Although degradation kinetics were slower than those

observed in NAA solubilized in M9 alone (**Figure 3B-D**) or NGM plates containing NAA

or K-NAA (**Figure S1B-C**), we still observed approximately 60–70% reduction of

AID::GFP expression within the first 30 minutes of NAA or K-NAA exposure, and nearly

80% depletion of whole animal AID::GFP within 1 hour (**Figure 3E-F**). Our results may

be under-representing the overall loss of AID::GFP as we did not account for gut

autofluorescence in our quantification of fluorescence intensity in whole animals

(Teuscher and Ewald 2018). Nonetheless, our results demonstrate that AID-tagged

proteins can be depleted in a microfluidic platform which, when combined with long-term

high-resolution imaging, provides a powerful tool for studying post-embryonic *C. elegans*

development at cellular resolution.

The AID system functions through specific components of the C. elegans SCF

complex to degrade target proteins

Our work demonstrates that the AID system functions rapidly to degrade target proteins

in *C. elegans* as described previously (Zhang *et al.* 2015). As a heterologous system,

researchers have shown in yeast that *Arabidopsis* TIR1 interacts with the *S. cerevisiae*

Cul1 homolog, Cdc53 (Nishimura *et al.* 2009). Whether *Arabidopsis* TIR1 also functions

through *C. elegans* proteins homologous to yeast SCF proteins is unknown. Thus, to

examine interactions between TIR1 and SCF complex proteins in *C. elegans*, we used

RNAi technology directed against components of the SCF complex and quantified

AID::GFP in the presence and absence of NAA (**Figure 4 and Figures S2-S3**). This

experiment was designed to provide insight into the mechanism through which the AID

system depletes target proteins in *C. elegans* and as an intersectional proof-of-concept

test of combining auxin-based depletion with a RNAi feeding approach. Briefly, the SCF

complex consists of three components: SKP1, CUL1, and RBX1. In contrast to yeast and

humans, which contain only one functional SKP1 protein, the scaffold protein CUL-1 is

known to interact with eight of the Skp1-related adaptor proteins in *C. elegans*, including

SKR-1, -2, -3, -4, -7, -8, -9 and -10 (Nayak *et al.* 2002; Yamanaka *et al.* 2002). We first

perturbed *cul-1* expression. To deplete CUL-1, we generated a new RNAi construct

targeting *cul-1* in the upgraded T444T RNAi targeting vector (Sturm *et al.* 2018). Notably,

this vector contains T7 terminator sequences, which prevents non-specific RNA

fragments from being synthesized from the vector backbone (Sturm *et al.* 2018). This

vector modification increases the efficiency of mRNA silencing over the original L4440

vector (Sturm *et al.* 2018).

We hypothesized that depleting CUL-1 would strongly interfere with the

proteasomal machinery and thus protein turnover. To assess the abundance of AID::GFP

in the *C. elegans* AC, we treated animals with either *control(RNAi)* or *cul-1(RNAi)*. As

before, we made use of a strain expressing AID::GFP and TIR1::mRuby from an *eft-3*

driver (Zhang *et al.* 2015), and we examined animals at the P6.p 2-cell stage (**Figure 4A**).

RNAi knockdown of *cul-1* resulted in a modest but statistically significant increase in the

abundance of AID::GFP in the AC compared to *control(RNAi)* treatment (+19%, *n* = 31

and 33, respectively, *P* < 0.0001) (**Figure 4C**). This result suggests that there is TIR-

dependent, auxin-independent depletion of AID::GFP, similar to reports in other systems

(Morawska and Ulrich 2013; Nishimura and Fukagawa 2017; Zasadzińska *et al.* 2018).

To further test this notion, we assessed GFP abundance in the AC in animals lacking

TIR1::mRuby.   AID::GFP and GFP were driven by *eft-3* and *zmp-1* promoters,

respectively, and we performed the experiment on animals at the P6.p 2-cell stage. We

did not assess AID::GFP abundance in the VPCs due to the variable sensitivity of this

tissue to RNAi (compare **Figure 4A and S2**) (Bourdages *et al.* 2014; Matus *et al.* 2014)**.**

Depletion of CUL-1 in animals expressing AID::GFP (-1.8%, *n* = 29 for both treatments,

P = 0.1168; **Figure S3A-B**) or GFP (-2.4%, *n* = 20 for both RNAi treatments, *P* = 0.7682;

Figure S3C-D) resulted in a slight decrease in GFP abundance, though it was not

statistically significant in either case compared to treatment with *control(RNAi)*. The

modest decrease in protein abundance in animals lacking TIR1::mRuby suggests that

knockdown of *cul-1* might mildly perturb protein homeostasis, but TIR1-mediated

proteosomal degradation of AID-tagged proteins independent of auxin exposure

requires endogenous levels of CUL-1 to function robustly.

Next, we tested whether depletion of *cul-1* would inactivate AID-mediated protein

degradation. We fed synchronized L1 stage animals with RNAi targeting *cul-1*, and then

treated animals at the P6.p 2-cell stage with 1 mM NAA for 60 minutes before quantifying

AID::GFP degradation in the AC (**Figure 4A**). For *control(RNAi)*-treated animals, the

abundance of AID::GFP in the AC was nearly undetectable within 60 minutes (-94%, *n* =

33; **Figure 4D**). Remarkably, the abundance of AID::GFP in the AC was reduced by only

29% within 60 minutes for animals treated with *cul-1(RNAi)* (*n* = 31, *P* < 0.0001; **Figure**

4D).

We next wanted to determine if any of the Skp1-related proteins in *C. elegans*

function as adaptors that link CUL-1 to the F-box protein TIR1 to mediate degradation of

AID-tagged target proteins. Based on the availability of RNAi clones, we fed synchronized

L1 stage animals with RNAi targeting four of the eight Skp1-related adaptors known to

interact with CUL-1; *skr-1*, *skr-2*, *skr-7*, and *skr-10* (Nayak *et al.* 2002; Yamanaka *et al.*

2002). Owing to the 83% sequence homology between *skr-1* and *skr-2* likely stemming

from a gene duplication event and predicted cross-RNAi effects, their gene names are

unified in this report as *skr-1/2* similar to (Nayak *et al.* 2002). Of all the Skp1 homologs,

C. elegans skr-1 and human Skp1 share the greatest sequence homology (Yamanaka *et*

al. 2002). We also fed animals RNAi targeting *rbx-1*, which encodes the RING finger

protein in the SCF E3 ubiquitin ligase (Yamanaka *et al.* 2002). To assess AID::GFP

abundance in the AC, we again used animals expressing *eft*-3>AID::GFP and *eft*-

3>TIR1::mRuby and examined animals at the P6.p 2-cell stage (**Figure 4B**). RNAi

knockdown of *skr-1/2* compared to *control(RNAi)* led to differences in AID::GFP

abundance that were not statistically significant (*n* = 23 and 20, respectively, *P* = 0.3522).

However, similar to *cul-1(RNAi)*, RNAi silencing of *rbx-1* (*n* = 22) resulted in a statistically

significant increase in the abundance of AID::GFP in the AC compared to *control(RNAi)*

treatment (*P* = 0.0283) (**Figure 4E**). Interestingly, both *skr-7(RNAi)* (*n* = 23) and *skr-*

10(RNAi) (*n* = 21) resulted in a statistically significant decrease in AID::GFP abundance

compared to control (*P* < 0.0001).

We also wanted to determine whether depletion of *skr-1/2*, *skr-7*, *skr-10,* and *rbx-*

1 could inactivate AID-mediated protein degradation We fed synchronized L1 stage

animals with RNAi targeting these SCF complex components. We treated animals at the

P6.p 2-cell stage with 1 mM NAA for 60 minutes and quantified AID::GFP degradation in

the AC (**Figure 4F**). For *control(RNAi)*-treated animals, the abundance of AID::GFP in the

AC was once again nearly undetectable within 60 minutes (-96%, *n* = 21) (**Figure 4F**).

Similarly, the AID::GFP abundance in animals treated with *skr-7(RNAi)* (*n* = 21) and *skr-*

10(RNAi) (*n* = 21) was undetectable within 60 minutes of NAA exposure (-94% and -93%,

respectively). For animals treated with *skr-1/2(RNAi)* (*n* = 21, *P* < 0.0001), the abundance

of AID::GFP in the AC was reduced by 57% within 60 minutes (**Figure 4F**). For animals

treated with *rbx-1(RNAi)* (*n* = 23, *P* < 0.0001), the abundance of AID::GFP in the AC was

reduced by 65% within 60 minutes (**Figure 4F**). These results suggest that: 1)

suppression of *cul-1*, *skr-1/2*, or *rbx-1* is sufficient to block TIR1-mediated degradation,

while suppression of *skr-7* or *skr-10* is not; 2) TIR1 functions as a substrate recognition

component of the *C. elegans* CUL-1-based SCF complex, which was also previously

shown in yeast (Nishimura *et al.* 2009); and 3) it is possible to deplete multiple targets

simultaneously using both AID and RNAi technology.

Inhibiting the expression of *cul-1, skr-1/2, or rbx-1* is a valid approach for reversing

AID-mediated degradation in *C. elegans*. We suggest using *cul-1, skr-1/2 or rbx-1(RNAi)*

for this purpose with caution, as they have known cell cycle-dependent functions and

therefore silencing them may conflate the recovery of AID-tagged proteins with a cell

cycle phenotype (Kipreos *et al.* 1996). As an alternative approach to achieving recovery

of AID-tagged proteins, we propose the use of RNAi targeting TIR1 or simply using

auxinole, a commercially available inhibitor of TIR1 (Hayashi *et al.* 2012; Yesbolatova *et*

al. 2019). One caveat to this approach is that auxinole is expensive and thus it may be

difficult to obtain stoichiometrically equivalent amounts of auxin and auxinole to truly

achieve recovery of one's protein of interest. However, for *C. elegans* researchers

requiring tighter temporal control, these may be avenues worth exploring. Presently,

recovery from degradation with 1 mM auxin takes up to 24 hours to fully recover

expression of the target protein (Zhang *et al.* 2015). Such protein recovery kinetics are

insufficient for studying events in the nematode that occur within minutes to hours such

as uterine-vulval attachment, vulval morphogenesis, or many other developmental events

occurring post-embryonically.

NAA as a tool for exploring phenotypes during development and beyond

As our previous results demonstrate that we could effectively deplete a non-functional

AID::GFP reporter expressed in the uterine AC and VPCs, we next tested whether NAA-

mediated depletion of target proteins could be utilized to study post-embryonic

developmental events occurring over a tight temporal window. We focused on a well-

studied system of organogenesis, *C. elegans* uterine-vulval cell specification and

morphogenesis (**Figure 1B**) (Schindler and Sherwood 2013). As a proof-of-principle, we

chose to deplete the nuclear hormone receptor, *nhr-25,* a homolog of arthropod Ftz-F1

and vertebrate SF-1/NR5A1 and LRH-1/NR5A2, which has been shown to function

pleiotropically in a wide array of developmental events, from larval molting (Asahina *et al.*

2000; Gissendanner and Sluder 2000; Frand *et al.* 2005), heterochrony (Hada *et al.*

2010), and uterine-vulval morphogenesis (Chen *et al.* 2004; Hwang and Sternberg 2004;

Asahina *et al.* 2006; Hwang *et al.* 2007; Ward *et al.* 2013). It was this pleiotropy that made

targeting NHR-25 an attractive target, as RNAi and mutant analyses have shown

previously that it is initially required in the AC during the AC/VU decision for proper

specification of AC fate (Hwang and Sternberg 2004; Asahina *et al.* 2006) and

approximately 7 hours later it is required in the underlying VPCs for cell division (Chen *et*

al. 2004; Hwang *et al.* 2007; Ward *et al.* 2013).

First, we examined the *nhr-25::GFP::AID* expression pattern, and observed GFP

localization to the nuclei of the AC/VU cells during the mid-L2 stage, enrichment in the

AC following specification, and nuclear localization in the 1˚ and 2˚ VPCs during all stages

of vulval division and morphogenesis (**Figure 5A**). We quantified GFP fluorescence over

developmental time. Consistent with previous reports based on transgene analyses

(Gissendanner and Sluder 2000; Ward *et al.* 2013), endogenous *nhr-25::GFP::AID* AC

expression peaks after AC specification in the early L3 at the P6.p 1-cell stage and is

undetectable above background by the P6.p 4-cell stage at the time of AC invasion.

Conversely, *nhr-25::GFP::AID* increases in intensity in the VPCs at the P6.p 4-cell stage,

peaking during the morphogenetic events following AC invasion (**Figure 5B**). Given this

temporally-driven expression pattern and based on previous experimental results from

RNAi and mutant analyses (Chen *et al.* 2004; Hwang and Sternberg 2004; Asahina *et al.*

2006; Hwang *et al.* 2007; Ward *et al.* 2013), we hypothesized that depleting AID-tagged

NHR-25 prior to AC specification should interfere with the AC/VU decision. To test this

hypothesis, we used synchronized L1 stage animals expressing *eft-3*>TIR1::mRuby and

endogenously tagged NHR-25::GFP::AID. We exposed these larvae to 4 mM NAA or a

buffer control and examined animals in the early L3 stage, after the normal time of AC

specification. Strikingly, all 36 animals examined showed a failure to specify the AC fate,

with the presence of either one (10/36) or two (26/36) small AC/VU-like cells in the central

gonad as compared to control animals (**Figure 5C**). Next, we repeated the experiment

but waited until after AC specification, in the early L3 stage, to expose animals to buffer

control or 4 mM NAA. Here, in all animals, we detected the presence of an AC situated

over P6.p, but in 34 of the 36 animals, P6.p failed to divide as compared to controls at

the mid-L3 stage (**Figure 5E**). Quantification of *nhr-25::GFP::AID* in AC/VU cells (**Figure**

5D) and VPCs (**Figure 5F**) demonstrated the 4 mM NAA treatment robustly depleted

endogenous protein by 95% in the AC/VU and 81% in the VPCs, respectively. Finally, we

waited until treated animals (early L3 stage) became adults (approximately 24 hours later)

and examined them for plate level phenotypes. We saw a 100% Egg-laying defect (Egl)

in 4 mM NAA treated animals as compared to control treated plates (**Figure 5G**).

Together, these results indicate that the synthetic auxin, NAA, can robustly deplete target

endogenous proteins in a facile, high-throughput fashion during uterine-vulval

development. This method should prove valuable in dissecting pleiotropic gene function

in the future.

Important caveats of the C. elegans AID system

Our results reported here with a synthetic auxin, NAA, as well as the initial report

developing the *C. elegans* AID system (Zhang *et al.* 2015) clearly demonstrate the

effectiveness of auxin-induced targeted protein depletion. Several other recent reports

have also effectively used the AID system in *C. elegans* to control protein function,

including controlling spermatogenesis by manipulating *spe-44* levels (Kasimatis *et al.*

2018), depleting a mediator component to modulate longevity (Lee *et al.* 2019), examining

chromosome segregation during oogenesis (Ferrandiz *et al.* 2018), meiotic crossover

(Zhang *et al.* 2018), and revealing novel roles of neuronal gene function through

conditional depletion (Serrano-Saiz *et al.* 2018). Despite the increasing frequency of AID

system usage in the *C. elegans* community, there are only a handful of TIR1 driver lines

published, and the importance of copy number and promoter strength has not been

systematically assessed.

While we are optimistic that the use of the synthetic analog of auxin presented

here will allow for even more widespread utility of the AID system in the *C. elegans*

research community, there are still are some areas open to improvement for the

technology. A recent report in mammalian cell culture identified that AID-tagged proteins

are depleted in an auxin-independent fashion in the presence of TIR1, relative to wild-

type levels (Li *et al.* 2019; Sathyan *et al.* 2019). We examined if this was also occurring

in *C. elegans* strains in our laboratory expressing AID-tagged proteins and TIR1. We were

able to detect statistically significant auxin-independent depletion of both a ubiquitously

expressed AID::GFP transgene under the *eft-3* promoter in ACs (-22%, $n = 24$, $P <$

0.0001) and VPCs (-24%, $n = 24$, $P < 0.0001$; **Figure S4A-B**) and an endogenously

tagged *nhr-25::GFP::AID* allele in ACs (-22%, $n = 26$, $P = 0.0022$) and VPCs (-35%, $n =$

26, $P < 0.0001$; **Figure S4C-D**). As partial loss of an endogenous protein could generate

a hypomorphic condition, both from the placement of the AID tag and apparent triggering

of the degradation machinery, we urge caution in carefully evaluating AID-tagged alleles

paired with TIR1, independent of auxin delivery. Further optimization of the AID system

in *C. elegans* will hopefully ameliorate this concern, as researchers recently used the

heterologous co-expression of an auxin response factor (ARF) with TIR1 to rescue auxin-

independent degradation in mammalian cell culture (Sathyan *et al.* 2019).

Conclusion

The ease of editing the *C. elegans* genome using CRISPR/Cas9-based approaches

(Calarco and Friedland 2015) and heterologous gene manipulation tools is ushering in a

new era of cellular and developmental biology. Several new tools available to *C. elegans*

researchers require the insertion of small amino acid tags into target loci, including ZF1

tagging (Armenti *et al.* 2014), sortase A (Wu *et al.* 2017), and the AID system (Zhang *et*

al. 2015). Alternatively, any GFP fusion can be targeted via a GFP nanobody tethered to

ZIF1 (Wang *et al.* 2017). These genomic edits are then paired with single transgene

expression to allow for targeted spatial and temporal loss-of-function approaches through

manipulation of endogenous loci. Prior to their advent, spatial and temporal control of

protein function was largely missing from the *C. elegans* genomic toolkit. With an ever-

increasing set of these tools being optimized for *C. elegans*, it is clear that different tools

will have strengths and weaknesses depending on multiple variables, including

subcellular localization of target protein, availability of tissue- and cell-type specific

drivers, and inducibility of depletion. Here, we optimize a powerful heterologous system,

the auxin-inducible degradation system. We demonstrate that a synthetic auxin analog,

NAA, and its water-soluble, potassium salt, K-NAA, can function equivalently to natural

auxin. The water solubility permits easier preparation of media and allows researchers to

perform experiments in liquid culture and microfluidics. Importantly, the use of ethanol

free K-NAA may be beneficial to *C. elegans* researchers studying behavior and aging,

where introduction of ethanol may lead to confounding results. We also demonstrate the

strength of the AID system for studying developmental cell biology by examining multiple

spatial and temporal roles of the Ftz-F1 homolog *nhr-25* during uterine and vulval

morphogenesis. It is our hope that the use of the synthetic auxin NAA will complement

the AID system in *C. elegans* when examining targeted protein depletion phenotypes in

tissues and developmental stages of interest. As the library of tissue-specific TIR1 drivers

continues to grow, we envision researchers being able to rapidly degrade proteins of

interest in specific tissues and visualize the outcome at single-cell resolution.

ACKNOWLEDGEMENTS

We thank R. Adikes for the suggestion to look into a water-soluble form of auxin and B.

Lacroix of the Vitaly Citovsky lab for providing the initial aliquot of NAA. We thank R.

Adikes, J. Smith and N. Palmisano for insightful discussions while developing the liquid

based NAA protocol and R. Adikes for helpful advice regarding image acquisition

parameters and quantification. We thank N. Stec of the Christopher Hammell lab for help

with the microfluidic device. We thank M. Li and N. Kim of the Jessica Seeliger lab for

help with the *E. coli* OP50 growth curve. We thank R. Adikes, N. Palmisano, A. Kohrman

and A. Nishimura for advice and comments on the manuscript. We appreciate the help of

W. Zhang for preparing media and plates.

This work was funded by the National Institute of Health (NIH) National Institute of

General Medical Sciences (NIGMS) [1R01GM121597-01 to D.Q.M.]. D.Q.M. is also a

Damon Runyon-Rachleff Innovator supported (in part) by the Damon Runyon Cancer

Research Foundation [DRR-47-17]. M.A.Q.M. is supported by the NIH NIGMS

[2T32GM007518-41] and [3R01GM121597-02S2]. T.N.M. is supported by the NIH

NICHD [1F31HD100091-01]. C.M.H and B.A.K. are supported by the NIH NIGMS

[R01GM117406]. J.D.W. is supported by the NIH NIGMS [R00GM-107345]. Some strains

were provided by the *Caenorhabditis* Genetics Center, which is funded by the NIH Office

of Research Infrastructure Programs [P40 OD010440].

AUTHOR CONTRIBUTIONS

M.A.Q.M. and D.Q.M conceived and designed the experiments. G.A. and J.D.W.

designed the constructs. J.M.R. performed the microinjections. G.A. performed the

crosses and characterized strains. M.A.Q.M, B.A.K., T.N.M., L.J. and J.A. performed the

experiments. M.A.Q.M. and D.Q.M. analyzed and quantified the data. M.A.Q.M. and

D.Q.M. wrote the manuscript with contributions from the other authors. The authors

declare no competing interests.

Martinez, et al. (2019)

LITERATURE CITED

Armenti, S. T., L. L. Lohmer, D. R. Sherwood, and J. Nance, 2014 Repurposing an endogenous degradation system for rapid and targeted depletion of C. elegans proteins. Development 141: 4640–4647.

Asahina, M., T. Ishihara, M. Jindra, Y. Kohara, I. Katsura *et al.*, 2000 The conserved nuclear receptor Ftz-F1 is required for embryogenesis, moulting and reproduction in Caenorhabditis elegans. Genes to Cells 5: 711–723.

Asahina, M., T. Valenta, M. Silhankova, V. Korinek, and M. Jindra, 2006 Crosstalk between a Nuclear Receptor and β-Catenin Signaling Decides Cell Fates in the C. elegans Somatic Gonad. Dev. Cell 11: 203–211.

Bourdages, K. G., B. Lacroix, J. F. Dorn, C. P. Descovich, and A. S. Maddox, 2014 Quantitative analysis of cytokinesis In Situduring C. Elegans postembryonic development. PLoS One 9:.

Brenner, S., 1974 The genetics of Caenorabditis elegans. Genetics 77: 71–94.

Cabreiro, F., C. Au, K. Y. Leung, N. Vergara-Irigaray, H. M. Cochemé *et al.*, 2013 Metformin retards aging in C. elegans by altering microbial folate and methionine metabolism. Cell 153: 228–239.

Calarco, J. A., and A. E. Friedland, 2015 *Creating Genome Modifications in C. elegans Using the CRISPR/Cas9 System*.

Camlin, N. J., and J. P. Evans, 2019 Auxin-inducible protein degradation as a novel approach for protein depletion and reverse genetic discoveries in mammalian

oocytes. Biol. Reprod.

Chen, Z., D. J. Eastburn, and M. Han, 2004 The Caenorhabditis elegans Nuclear

Receptor Gene nhr-25 Regulates Epidermal Cell Development. Mol. Cell. Biol. 24:

7345–7358.

Chen, W., M. Werdann, and Y. Zhang, 2018 The auxin-inducible degradation system

enables conditional PERIOD protein depletion in the nervous system of Drosophila

melanogaster. FEBS J. 285: 4378–4393.

Cohen, J., 1992 A Power Primer. Psychol. Bull. 112: 155–159.

Daniel, K., J. Icha, C. Horenburg, D. Müller, C. Norden *et al.*, 2018 Conditional control of

fluorescent protein degradation by an auxin-dependent nanobody. Nat. Commun.

9:.

Dickinson, D. J., and B. Goldstein, 2016 CRISPR-based methods for Caenorhabditis

elegans genome engineering. Genetics 202: 885–901.

Dickinson, D. J., A. M. Pani, J. K. Heppert, C. D. Higgins, and B. Goldstein, 2015

Streamlined Genome Engineering with a Self-Excising Drug Selection Cassette.

200: 1035–1049.

Dickinson, D. J., M. M. Slabodnick, A. H. Chen, and B. Goldstein, 2018 SapTrap

assembly of repair templates for Cas9-triggered homologous recombination with a

self-excising cassette. microPublication Biol.

Dickinson, D. J., J. D. Ward, D. J. Reiner, and B. Goldstein, 2013 Engineering the

Caenorhabditis elegans genome using Cas9-triggered homologous recombination.

Martinez, et al. (2019)

   Nat. Methods 10: 1028–1034.

Dokshin, G. A., K. S. Ghanta, K. M. Piscopo, and C. C. Mello, 2018 Robust genome

   editing with short single-stranded and long, partially single-stranded DNA donors in

   caenorhabditis elegans. Genetics 210: 781–787.

Ferrandiz, N., C. Barroso, O. Telecan, N. Shao, H. M. Kim *et al.*, 2018 Spatiotemporal

   regulation of Aurora B recruitment ensures release of cohesion during C. Elegans

   oocyte meiosis. Nat. Commun. 9: 1–16.

Frand, A. R., S. Russel, and G. Ruvkun, 2005 Functional genomic analysis of c.

   elegans molting. PLoS Biol. 3:.

Frøkjær-Jensen, C., M. W. Davis, M. Ailion, and E. M. Jorgensen, 2012 Improved

   Mos1-mediated transgenesis in C. elegans. Nat. Methods 9: 117–118.

Gissendanner, C. R., and A. E. Sluder, 2000 nhr-25, the Caenorhabditis elegans

   ortholog of ftz-f1, is required for epidermal and somatic gonad development. Dev.

   Biol. 221: 259–272.

Gupta, B. P., W. Hanna-Rose, and P. W. Sternberg, 2012 Morphogenesis of the vulva

   and the vulval-uterine connection. WormBook 1–20.

Hada, K., M. Asahina, H. Hasegawa, Y. Kanaho, F. J. Slack *et al.*, 2010 The nuclear

   receptor gene nhr-25 plays multiple roles in the Caenorhabditis elegans

   heterochronic gene network to control the larva-to-adult transition. Dev. Biol. 344:

   1100–1109.

Hayashi, K. I., J. Neve, M. Hirose, A. Kuboki, Y. Shimada *et al.*, 2012 Rational design of

an auxin antagonist of the SCF TIR1 auxin receptor complex. ACS Chem. Biol. 7: 590–598.

Holland, A. J., D. Fachinetti, J. S. Han, and D. W. Cleveland, 2012 Inducible, reversible system for the rapid and complete degradation of proteins in mammalian cells. Proc. Natl. Acad. Sci. 109: E3350–E3357.

Hubbard, E. J. A., 2014 FLP/FRT and Cre/lox recombination technology in C. elegans. Methods 68: 417–424.

Hwang, B. J., A. D. Meruelo, and P. W. Sternberg, 2007 C. elegans EVI1 proto-oncogene, EGL-43, is necessary for Notch-mediated cell fate specification and regulates cell invasion. Development 134: 669–679.

Hwang, B. J., and P. W. Sternberg, 2004 A cell-specific enhancer that specifies lin-3 expression in the C. elegans anchor cell for vulval development. Development 131: 143–151.

Kasimatis, K. R., M. J. Moerdyk-Schauwecker, and P. C. Phillips, 2018 Auxin-mediated sterility induction system for longevity and mating studies in Caenorhabditis elegans. G3 Genes, Genomes, Genet. 8: 2655–2662.

Keil, W., L. M. Kutscher, S. Shaham, and E. D. Siggia, 2017 Long-Term High-Resolution Imaging of Developing C. elegans Larvae with Microfluidics. Dev. Cell 40: 202–214.

Khanna, N., C. P. C. Iii, C. P. Tatara, and P. L. Williams, 1997 Environmental Contamination and Toxicology Tolerance of the Nematode Caenorhabditis elegans

to pH , Salinity , and Hardness in Aquatic Media. Arch. Environ. Contam. Toxicol. 114: 110–114.

Kipreos, E. T., L. E. Lander, J. P. Wing, W. W. He, and E. M. Hedgecock, 1996 cul-1 is required for cell cycle exit in C. elegans and identifies a novel gene family. Cell 85: 829–839.

Kwon, J. Y., M. Hong, M. S. Choi, S. Kang, K. Duke *et al.*, 2004 Ethanol-response genes and their regulation analyzed by a microarray and comparative genomic approach in the nematode Caenorhabditis elegans. Genomics 83: 600–614.

Lee, D., S. W. A. An, Y. Jung, Y. Yamaoka, Y. Ryu *et al.*, 2019 MDT-15/MED15 permits longevity at low temperature via enhancing lipidostasis and proteostasis. PLOS Biol. 17: e3000415.

Li, S., X. Prasanna, V. T. Salo, I. Vattulainen, and E. Ikonen, 2019 An efficient auxin-inducible degron system with low basal degradation in human cells. Nat. Methods 16: 866–869.

Matus, D. Q., E. Chang, S. C. Makohon-Moore, M. A. Hagedorn, Q. Chi *et al.*, 2014 Cell division and targeted cell cycle arrest opens and stabilizes basement membrane gaps. Nat. Commun. 5: 1–13.

Morawska, M., and H. D. Ulrich, 2013 An expanded tool kit for the auxin-inducible degron system in budding yeast. Yeast 30: 341–351.

Morgan, P. G., and M. M. Sedensky, 1995 Mutations Affecting Sensitivity to Ethanol in the Nematode, Caenorhabditis elegans. Alcohol. Clin. Exp. Res. 19: 1423–1429.

Martinez, et al. (2019)

Natsume, T., T. Kiyomitsu, Y. Saga, and M. T. Kanemaki, 2016 Rapid Protein Depletion
in Human Cells by Auxin-Inducible Degron Tagging with Short Homology Donors.
Cell Rep. 15: 210–218.

Nayak, S., F. E. Santiago, H. Jin, D. Lin, T. Schedl *et al.*, 2002 The Caenorhabditis
elegans Skp1-related gene family: Diverse functions in cell proliferation,
morphogenesis, and meiosis. Curr. Biol. 12: 277–287.

Nishimura, K., and T. Fukagawa, 2017 An efficient method to generate conditional
knockout cell lines for essential genes by combination of auxin-inducible degron tag
and CRISPR/Cas9. Chromosom. Res. 25: 253–260.

Nishimura, K., T. Fukagawa, H. Takisawa, T. Kakimoto, and M. Kanemaki, 2009 An
auxin-based degron system for the rapid depletion of proteins in nonplant cells.
Nat. Methods 6: 917–922.

Papagiannakis, A., J. J. De Jonge, Z. Zhang, and M. Heinemann, 2017 Quantitative
characterization of the auxin-inducible degron: A guide for dynamic protein
depletion in single yeast cells. Sci. Rep. 7: 1–13.

Pollard, D. A., T. D. Pollard, and K. S. Pollard, 2019 Empowering statistical methods for
cellular and molecular biologists. Mol. Biol. Cell 30: 1359–1368.

Porta-de-la-Riva, M., L. Fontrodona, A. Villanueva, and J. Cerón, 2012 Basic
Caenorhabditis elegans Methods: Synchronization and Observation. J.
Vis. Exp. 1–9.

Qadota, H., M. Inoue, T. Hikita, M. Köppen, J. D. Hardin *et al.*, 2007 Establishment of a

Martinez, et al. (2019)

tissue-specific RNAi system in C. elegans. Gene 400: 166–173.

Röth, S., L. J. Fulcher, and G. P. Sapkota, 2019 Advances in targeted degradation of endogenous proteins. Cell. Mol. Life Sci.

Rual, J.-F., J. Ceron, J. Koreth, T. Hao, A.-S. Nicot *et al.*, 2004 Toward Improving Caenorhabditis elegans Phenome Mapping With an ORFeome-Based RNAi Library. Genome Res. 14: 2162–2168.

Sathyan, K. M., B. D. McKenna, W. D. Anderson, F. M. Duarte, L. Core *et al.*, 2019 An improved auxin-inducible degron system preserves native protein levels and enables rapid and specific protein depletion. Genes Dev. 1–15.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair *et al.*, 2012 Fiji: an open-source platform for biological-image analysis. Nat. Methods 9: 676–82.

Schindler, A. J., and D. R. Sherwood, 2013 Morphogenesis of the Caenorhabditis elegans vulva. Wiley Interdiscip. Rev. Dev. Biol. 2: 75–95.

Schwartz, M. L., and E. M. Jorgensen, 2016 SapTrap, a toolkit for high-throughput CRISPR/Cas9 gene modification in Caenorhabditis elegans. Genetics 202: 1277–1288.

Serrano-Saiz, E., E. Leyva-Díaz, E. De La Cruz, and O. Hobert, 2018 BRN3-type POU Homeobox Genes Maintain the Identity of Mature Postmitotic Neurons in Nematodes and Mice. Curr. Biol. 28: 2813-2823.e2.

Shen, Z., X. Zhang, Y. Chai, Z. Zhu, P. Yi *et al.*, 2014 Conditional knockouts generated by engineered CRISPR-Cas9 endonuclease reveal the roles of coronin in c.elegans

Martinez, et al. (2019)

neural development. Dev. Cell 30: 625–636.

Srivastava, L. M., 2002 *CHAPTER 6 - Auxins*. Academic Press.

Sturm, Á., É. Saskoi, K. Tibor, N. Weinhardt, and T. Vellai, 2018 Highly efficient RNAi
and Cas9-based auto-cloning systems for C. elegans research. Nucleic Acids Res.
46: e105.

Teuscher, A., and C. Ewald, 2018 Overcoming Autofluorescence to Assess GFP
Expression During Normal Physiology and Aging in Caenorhabditis elegans. Bio-
Protocol 8:.

Trost, M., A. C. Blattner, and C. F. Lehner, 2016 Regulated protein depletion by the
auxin-inducible degradation system in Drosophila melanogaster. Fly (Austin). 10:
35–46.

Wang, S., N. H. Tang, P. Lara-Gonzalez, Z. Zhao, D. K. Cheerambathur *et al.*, 2017 A
toolkit for GFP-mediated tissue-specific protein degradation in C. elegans .
Development 144: 2694–2701.

Ward, J. D., N. Bojanala, T. Bernal, K. Ashrafi, M. Asahina *et al.*, 2013 Sumoylated
NHR-25/NR5A Regulates Cell Fate duringC. elegans Vulval Development. PLoS
Genet. 9:.

Wu, Q., H. L. Ploegh, and M. C. Truttmann, 2017 Hepta-Mutant Staphylococcus aureus
Sortase A (SrtA7m) as a Tool for in Vivo Protein Labeling in Caenorhabditis
elegans. ACS Chem. Biol. 12: 664–673.

Yamanaka, A., M. Yada, H. Imaki, M. Koga, Y. Ohshima *et al.*, 2002 Multiple Skp1-

related proteins in Caenorhabditis elegans: Diverse patterns of interaction with Cullins and F-box proteins. Curr. Biol. 12: 267–275.

Yesbolatova, A., T. Natsume, K. ichiro Hayashi, and M. T. Kanemaki, 2019 Generation of conditional auxin-inducible degron (AID) cells and tight control of degron-fused proteins using the degradation inhibitor auxinole. Methods 0–1.

Zasadzińska, E., J. Huang, A. O. Bailey, L. Y. Guo, N. S. Lee *et al.*, 2018 Inheritance of CENP-A Nucleosomes during DNA Replication Requires HJURP. Dev. Cell 47: 348-362.e7.

Zhang, L., S. Köhler, R. Rillo-Bohn, and A. F. Dernburg, 2018 A compartmentalized signaling network mediates crossover control in meiosis. Elife 7: 1–32.

Zhang, L., J. D. Ward, Z. Cheng, and A. F. Dernburg, 2015 The auxin-inducible degradation (AID) system enables versatile conditional protein depletion in C. elegans. Development 142: 4374–4384.

Figure 1. Overview of the auxin-inducible degradation system and *C. elegans* uterine-vulval development. (A) In this system, a target protein is fused to an auxin-inducible degron (AID). Heterologous expression of *Arabidopsis* TIR1 mediates robust auxin-dependent proteasomal degradation of AID-tagged proteins through the SKP1-CUL1-F-box (SCF) E3 ubiquitin ligase complex. (B) Schematic of uterine-vulval morphogenesis during *C. elegans* larval development. In *C. elegans*, AC specification and morphogenesis of uterine-vulval attachment occurs from the mid-L2 through the early L4 stage (Schindler and Sherwood 2013). The AC is specified in a stochastic reciprocal Notch-Delta signaling event in the mid-L2 stage (top panel). Following AC specification, the AC specifies the 1° fate of the underlying vulval precursor cell, P6.p in the early L3 (second panel), which then divides three times to ultimately give rise to eight of the 22 cells of the adult vulva (bottom three panels).

Figure 2. Comparison of IAA- and NAA-mediated degradation in the *C. elegans* AC and VPCs. (A) DIC and corresponding GFP images of ACs (arrowheads) and underlying 1° fated VPCs (brackets) from mid-L3 stage animals at the P6.p 2-cell stage. Animals expressing AID::GFP and TIR1::mRuby under the same *eft-3* promoter were treated with natural auxin indole-3-acetic acid (IAA) and synthetic auxin 1-naphthaleneacetic acid (NAA) in NGM agar containing OP50. (B) Chemical structure of IAA and NAA. (C, D) Rates of degradation determined by quantifying AID::GFP in (C) ACs and (D) VPCs

Martinez, et al. (2019)

following auxin treatment. Data presented as the mean±SD ($n \geq 30$ animals examined for each time point).

Figure 3. Solubility of NAA/K-NAA in physiological buffer enhances utility. (A) Schematic representation of the liquid NAA-based degradation protocol for use in high-resolution microscopy or microfluidics-based approaches. (B) DIC and corresponding GFP images of ACs (arrowheads) and underlying VPCs (brackets) from mid-L3 stage animals at the P6.p 2-cell stage. Animals expressing AID::GFP and TIR1::mRuby under the same *eft-3* promoter were treated with NAA in M9. (C, D) Rates of degradation were determined by quantifying AID::GFP in (C) ACs and (D) VPCs following auxin treatment. Data presented as the mean±SD ($n \geq 30$ animals examined for each time point). (E) Images of AID::GFP expression from mid-L3 stage animals in control conditions (M9 buffer containing NA22 only, left) or conditions where a 4 mM NAA (middle) or K-NAA (right) solution in M9 buffer containing NA22 was perfused through the microfluidic chamber for the time indicated (Keil et al. 2017). Anterior is left and ventral is down. (F) Rates of degradation were determined by quantifying AID::GFP in whole animals following auxin treatment. Data presented as the mean±SD ($n \geq 4$ animals examined for each time point).

Figure 4. Suppression of SCF complex member expression inhibits TIR1-dependent degradation in the *C. elegans* AC. (A-B) DIC and corresponding GFP images of ACs (arrowheads) and underlying VPCs (brackets) from mid-L3 stage animals at the P6.p 2-cell stage. Animals expressing AID::GFP and TIR1::mRuby under the same *eft-3* promoter were treated with (A) *cul-1(RNAi)* and (B) *skr-1/2, skr-7, skr-10* and *rbx-1(RNAi)*. (C) Quantification of AID::GFP in ACs following *cul-1(RNAi)* treatment. Data presented as the mean±SD ($n \geq 30$ animals examined for each, and $P < 0.0001$ by a Student's t-test). (D) Quantification of AID::GFP in ACs following treatment with NAA. Data presented as the median+IQR ($n \geq 30$ animals examined for each, and $P < 0.0001$ by a Mann Whitney U test). (E) Quantification of AID::GFP in ACs following RNAi knockdown of Skp1-related (*skr*) genes and *rbx-1*. Data presented as the median+IQR ($n \geq 20$ animals examined for each, and P values examined by a Mann Whitney U test). (F) Quantification of AID::GFP in ACs following treatment with NAA. Data presented as the median+IQR ($n \geq 21$ animals examined for each, and P values examined by a Mann Whitney U test).

Figure 5. NAA-mediated degradation of NHR-25 causes AC specification and VPC division defects. (A) *nhr-25::GFP::AID* localizes to the nuclei of the AC/VU (black arrowheads), the AC (white arrowheads) and VPCs (brackets). At P6.p 8-cell stage (far right) the AC is not in the same focal plane as the 1° VPCs. (B) Quantification of *nhr-*

25::GFP::AID over developmental time, from the AC/VU decision to the P6.p 8-cell stage. The curve is connected by the mean at each developmental stage (n = 20, 31, 20, 21, and 12 animals quantified, respectively). (C) DIC and corresponding GFP images of ACs (arrowheads) and underlying VPCs (brackets) from early L3 stage animals. Animals expressing *nhr-25::AID::GFP* and *eft-3*>TIR1::mRuby were treated with control and 4 mM NAA. (D) Quantification of *nhr-25::GFP::AID* in AC/VUs following NAA treatment. Data presented as the median+IQR ($n \geq 20$ animals examined for each, and $P < 0.0001$ by a Mann Whitney U test). (E) DIC and corresponding GFP images of ACs (arrowheads) and underlying VPCs (brackets) from mid-L3 stage animals. Animals expressing *nhr-25::AID::GFP* and *eft-3*>TIR1::mRuby were treated with control and 4 mM NAA. (F) Quantification of *nhr-25::GFP::AID* in VPCs following NAA treatment. Data presented as the median+IQR ($n \geq 30$ animals examined for each, and $P < 0.0001$ by a Mann Whitney U test). (G) Representative images of adult plate level phenotypes following control and 4 mM NAA treatments added at the L3 stage ($n \geq 30$ animals examined). Scale bar in (G), 500 µm.