TITLE

Low coverage whole genome sequencing enables accurate assessment of common variants

and calculation of genome-wide polygenic scores

AUTHORS AND AFFILIATIONS

Julian R. Homburger,[1] Cynthia L. Neben,[1] Gilad Mishne,[1] Alicia Y. Zhou,[1] Sekar Kathiresan,[2-4]

Amit V. Khera[2-4] *

[1]Color Genomics, 831 Mitten Road, Suite 100, Burlingame, CA, 94010 USA

[2]Center for Genomic Medicine and Cardiology Division, Department of Medicine, Massachusetts

General Hospital, Boston, MA, 02114 USA

[3]Cardiovascular Disease Initiative of the Broad Institute of MIT and Harvard, Cambridge, MA,

02142 USA

[4]Harvard Medical School, Boston, MA, 02115 USA

CORRESPONDING AUTHOR

*Amit V. Khera, MD, MSc

Center for Genomic Medicine

Massachusetts General Hospital

Simches Research Building | CPZN 6.256

Boston, MA 02114 USA

Tel: 617.726.7876

Email: avkhera@mgh.harvard.edu

ABSTRACT

Background: The inherited susceptibility of common, complex diseases may be caused by

rare, 'monogenic' pathogenic variants or by the cumulative effect of numerous common,

'polygenic' variants. As such, comprehensive genome interpretation could involve two distinct

genetic testing technologies -- high coverage next generation sequencing for known genes to

detect pathogenic variants and a genome-wide genotyping array followed by imputation to

calculate genome-wide polygenic scores (GPSs). Here we assessed the feasibility and

accuracy of using low coverage whole genome sequencing (lcWGS) as an alternative to

genotyping arrays to calculate GPSs.

Methods: First, we performed downsampling and imputation of WGS data from ten individuals

to assess concordance with known genotypes. Second, we assessed the correlation between

GPSs for three common diseases -- coronary artery disease (CAD), breast cancer (BC), and

atrial fibrillation (AF) -- calculated using lcWGS and genotyping array in 184 samples. Third, we

assessed concordance of lcWGS-based genotype calls and GPS calculation in 120 individuals

with known genotypes, selected to reflect diverse ancestral backgrounds. Fourth, we assessed

the relationship between GPSs calculated using lcWGS and disease phenotypes in 11,502

European individuals seeking genetic testing.

Results: We found imputation accuracy r^2 values of greater than 0.90 for all ten samples --

including those of African and Ashkenazi Jewish ancestry -- with lcWGS data at 0.5X. GPSs

calculated using both lcWGS and genotyping array followed by imputation in 184 individuals

were highly correlated for each of the three common diseases (r^2 = 0.93 - 0.97) with similar

score distributions. Using lcWGS data from 120 individuals of diverse ancestral backgrounds,

including South Asian, East Asian, and Hispanic individuals, we found similar results with

respect to imputation accuracy and GPS correlations. Finally, we calculated GPSs for CAD, BC,

and AF using lcWGS in 11,502 European individuals, confirming odds ratios per standard

deviation increment in GPSs ranging 1.28 to 1.59, consistent with previous studies.

Conclusions: Here we show that lcWGS is an alternative approach to genotyping arrays for

common genetic variant assessment and GPS calculation. lcWGS provides comparable

imputation accuracy while also overcoming the ascertainment bias inherent to variant selection

in genotyping array design.

KEYWORDS

Genome-wide polygenic score; low coverage whole genome sequencing; coronary artery

disease; breast cancer; atrial fibrillation

BACKGROUND

Cardiovascular disease and cancer are common, complex diseases that remain leading causes of global mortality [1]. Long recognized to be heritable, recent advances in human genetics have led to consideration of DNA-based risk stratification to guide prevention or screening strategies. In some cases, such conditions can be caused by rare, 'monogenic' pathogenic variants that lead to a several-fold increased risk -- important examples are pathogenic variants in *LDLR* that cause familial hypercholesterolemia and pathogenic variants in *BRCA1* and *BRCA2* that underlie hereditary breast and ovarian cancer syndrome. However, the majority of individuals afflicted with these diseases do not harbor any such pathogenic variants. Rather, the inherited susceptibility of many complex traits and diseases is often 'polygenic,' driven by the cumulative effect of numerous common variants scattered across the genome [2].

Genome-wide polygenic scores (GPSs) provide a way to integrate information from numerous sites of common variation into a single metric of inherited susceptibility and are now able to identify individuals with a several-fold increased risk of common, complex diseases, including coronary artery disease (CAD), breast cancer (BC), and atrial fibrillation (AF) [3]. For example, for CAD, we noted that 8% of the population inherits more than triple the normal risk on the basis of polygenic variation, a prevalence more than 20-fold higher than monogenic familial hypercholesterolemia variants in *LDLR* that confer similar risk [3].

Comprehensive genome interpretation for common, complex disease therefore could involve both high-fidelity sequencing of important driver genes to identify potential monogenic risk pathogenic variants and a survey of all common variants across the genome to enable GPS calculation. High coverage whole genome sequencing (hcWGS; for example, 30X coverage) will likely emerge as a single genetic testing strategy, but current prices remain a barrier to large-scale adoption. Instead, the traditional approach has mandated use of two distinct genetic

testing technologies -- high coverage next generation sequencing (NGS) of important genes to

detect pathogenic variants and a genome-wide genotyping array followed by imputation to

calculate GPSs.

Low coverage whole genome sequencing (lcWGS; for example, 0.5X coverage) followed by

imputation is a potential alternative approach to genotyping arrays for assessing the common

genetic variants needed for GPS calculations. Several recent studies have demonstrated the

efficiency and accuracy of lcWGS for other applications of statistical genetics, including local

ancestry deconvolution, complex trait association studies, and detection of rare genetic variants

[4–7].

We developed a pipeline for common genetic variant imputation using lcWGS data on samples

from the 1000 Genomes Project (1KGP) and Genome in a Bottle (GIAB) Consortium and herein

demonstrate imputation accuracy for lcWGS similar to genotyping arrays. Using three recently

published GPSs for CAD [3], BC [8], and AF [3], we show high technical concordance in GPSs

calculated from lcWGS and genotyping arrays. Finally, using our pipeline in a large European

population seeking genetic testing, we observe similar GPS risk stratification performance as

previously published array-based results [3,8].

METHODS

Study design

The study design is summarized in Figure 1 and described in detail below. The pipeline

validation data set (n = 10) was used to assess imputation accuracy for common genetic

variants (Figure 1A). The technical concordance cohort (n = 184) was used to assess the

correlation between three previously published GPSs for CAD [3], BC [8], and AF [3] from

lcWGS and genotyping arrays (Figure 1B). The diverse ancestry data set (n = 120) was used to

assess imputation accuracy for common genetic variants and performance of GPS_{CAD}, GPS_{BC},

and GPS_{AF} (Figure 1B). The clinical cohort (n = 11,502) was used to assess performance of

GPS_{CAD}, GPS_{BC}, and GPS_{AF} in a large European population seeking genetic testing (Figure 1B).

Data set and cohort selection

The pipeline validation data set included seven globally representative samples from 1KGP

populations (HG02155, NA12878, HG00663, HG01485, NA21144, NA20510, and NA19420;

see Supplementary Table 1, Additional File 1) and a trio of Ashkenazi samples (NA24385,

NA24143, and NA24149) from the GIAB Consortium (Figure 1A).

The technical concordance cohort included DNA samples from 184 individuals whose

healthcare provider had ordered a Color multi-gene panel test (Figure 1B). All individuals 1) had

85% or greater European genetic ancestry calculated using fastNGSadmix [9] using 1KPG as

the reference panel, 2) self-identified as 'Caucasian', and 3) did not have pathogenic or likely

pathogenic variants in the multi-gene NGS panel test, as previously described [10] (see

Supplementary Methods, Additional File 2). Demographics are provided in Supplementary Table

2, Additional File 1. All phenotypic information was self-reported by the individual through an

online, interactive health history tool. Of the 184 individuals, 61 individuals reported having a

personal history of CAD (defined here as a myocardial infarction or coronary artery bypass

surgery), 62 individuals reported no personal history of CAD, and 61 individuals reported no

personal history of CAD but were suspected to have a high GPS_{CAD} based on preliminary

analysis. This preliminary analysis included imputation from multi-gene panel and off-target

sequencing data, which has been shown to have similar association statistics and effect sizes

compared to genotyping arrays [4]. These individuals were included in the technical

concordance cohort to artificially create a relatively uniform distribution of GPS_{CAD} in the data

set. Correlation coefficients between GPS_{CAD} from lcWGS and genotyping array were calculated

after removing the 61 individuals who were suspected to have a high GPS_{CAD} based on multi-

gene panel and off-target sequencing data to avoid artificial inflation of the correlation

coefficient. Two individuals who reported no personal history of CAD but were suspected to

have a high GPS_{CAD} failed genotyping (quality control call rate of < 97%) and lcWGS (overall

coverage of < 0.5X), leaving a total of 182 individuals for analyses.

The diverse ancestry data set included a total of 120 samples from the following populations

from 1KGP: Han Chinese in Beijing, China (CHB); Yoruba in Ibadan, Nigeria (YRI); Gujarati

Indian from Houston, Texas (GIH); Americans of African Ancestry in Southwest USA (ASW);

Mexican Ancestry from Los Angeles, USA (MXL); and Puerto Ricans from Puerto Rico (PUR)

(see Supplementary Table 3, Additional File 1; Figure 1B). Four samples, including NA18917

and NA19147 from the YRI population and NA19729 and NA19785 from the MXL population,

were below the target 0.5X coverage and removed from analyses.

The clinical cohort included DNA samples from 11,502 individuals whose healthcare provider

had ordered a Color multi-gene panel test (Figure 1B). All individuals 1) had 90% or greater

European genetic ancestry calculated using fastNGSadmix [9] using 1KPG as the reference

panel, 2) self-identified as 'Caucasian', 3) provided history of whether they had a clinical

diagnosis of CAD, BC, or AF, and 4) did not have pathogenic or likely pathogenic variants

detected in the multi-gene NGS panel test, as previously described [10] (see Supplementary

Methods, Additional File 2). Demographics are provided in Supplementary Table 2, Additional

File 1. All phenotypic information was self-reported by the individual through an online,

interactive health history tool.

Whole genome sequencing

DNA was extracted from blood or saliva samples and purified using the Perkin Elmer Chemagic

DNA Extraction Kit (Perkin Elmer, Waltham, MA) automated on the Hamilton STAR (Hamilton,

Reno, NV) and the Chemagic Liquid Handler (Perkin Elmer, Waltham, MA). The quality and

quantity of the extracted DNA were assessed by UV spectroscopy (BioTek, Winooski, VT). High

molecular weight genomic DNA was enzymatically fragmented and prepared using the Kapa

HyperPlus Library Preparation Kit (Roche Sequencing, Pleasanton, CA) automated on the

Hamilton Star liquid handler and uniquely tagged with 10 bp dual-unique barcodes (IDT,

Coralville, IA). Libraries were pooled together and loaded onto the NovaSeq 6000 (Illumina, San

Diego, CA) for 2 x 150 bp sequencing.

For the pipeline validation data set, all samples underwent WGS with mean coverage of 13.22X

(range 7.82X to 17.30X); downsampling was then performed using SAMtools to simulate

lcWGS. For the technical concordance cohort, all samples underwent lcWGS with mean

coverage of 1.24X (range 0.54X to 1.76X). Imputed genotypes were compared with published,

high-confidence known genotypes from 1KGP and the GIAB Consortium. For the diverse

ancestry data set, all samples underwent lcWGS with mean coverage of 0.89X (range 0.68X to

1.24X). For the clinical cohort, all samples underwent lcWGS with mean coverage of 0.95X

(range 0.51X to 2.57X).

Downsampling

For the pipeline validation data set, aligned reads were downsampled using SAMtools [11] to

2.0X, 1.0X, 0.75X, 0.5X, 0.4X, 0.25X, and 0.1X coverage. For the technical concordance cohort,

aligned reads were downsampled to 1.0X, 0.75X, 0.5X, 0.4X, 0.25X, and 0.1X coverage. In a

few cases in the technical concordance cohort, the primary samples had fewer reads than the

target downsample. In those situations, all of the reads were retained. For example, if the

primary sample only had 0.8X coverage, when downsampled to 1.0X, all reads were retained.

Downsampling was repeated using two independent seeds in SAMtools. Once the

downsampled data was generated, the imputation was repeated to generate imputed genotypes

using only the downsampled reads.

Imputation site selection

All data sets and cohorts were imputed to a set of autosomal SNP and insertion-deletion (indel)

sites from 1KGP with greater than 1% allele frequency in any of the five 1KGP super

populations (African, American, East Asian, European, and South Asian), for a total of

21,770,397 sites. This is hereafter referred to as the 'imputation SNP loci.' Multi-allelic SNPs

and indels were represented as two biallelic markers for imputation.

Genotype likelihood calculations and imputation

Genotype likelihood calculations and imputation were performed independently for each

sample. Sequence reads were aligned with the human genome reference GRCh37.p12 using

the Burrows-Wheeler Aligner (BWA) [12], and duplicate and low quality reads were removed.

Genotype likelihoods were then calculated at each of the biallelic SNP loci in the imputation

SNP loci that were covered by one or more sequencing reads called using the mpileup

command implemented in bcftools version 1.8 [13]. Indels or multi-allelic sites were not included

in this first genotype likelihood calculation. Reads with a minimum mapping alignment quality of

10 or greater and bases with a minimum base quality of 10 or greater were included. Genotype

likelihoods at each observed site were then calculated using the bcftools call command with

allele information corresponding to the imputation SNP loci. This procedure discarded calls with

indels or calls where the observed base did not match either the reference or expected alternate

allele for the SNP locus.

Imputation was performed using the genotype likelihood imputation option implemented in

BEAGLE 4.1 [14]. This imputation used default parameters except with a model scale

parameter of 2 and the number of phasing iterations to 0. A custom reference panel was

constructed for each sample being imputed by selecting the 250 most similar samples to that

sample from 1KGP Phase 3 release using Identity-by-State (IBS) comparison. A reference

panel size of 250 was selected to best balance imputation run time and accuracy (see

Supplementary Figure 1, Additional File 2). To ensure that IBS values were comparable across

samples, a set of regions consistently sequenced at high depth (> 20X) across all samples was

utilized. When imputation was performed on samples included in 1KGP Phase 3 release, that

sample and any related samples were excluded from the custom reference panel.

To generate genotypes at all of the remaining untyped sites, a second round of imputation was

performed using BEAGLE 5.0 [15]. This imputation used default settings and included the full

1KGP as the imputation reference panel. To note, when performing analysis using 1KGP

samples, any related individuals were removed. Each sample then had imputed genotype calls

at each of the imputation SNP loci. Indels and multiallelic sites were included in this second

genotype likelihood calculation.

Genotyping array

DNA was extracted from blood or saliva samples and purified using the Perkin Elmer Chemagic

DNA Extraction Kit (Perkin Elmer, Waltham, MA) automated on the Hamilton STAR (Hamilton,

Reno, NV) and the Chemagic Liquid Handler (Perkin Elmer, Waltham, MA). The quality and

quantity of the extracted DNA were assessed by UV spectroscopy (BioTek, Winooski, VT).

DNA was genotyped on the Axiom UK Biobank Array by Affymetrix (Santa Clara, CA).

Genotypes were filtered according to the manufacturer's recommendations, removing loci with

greater than 5% global missingness and those that significantly deviated from Hardy-Weinberg

equilibrium. In addition, all A/T and G/C SNPs were removed due to potential strand

inconsistencies. Each of the remaining SNPs were aligned with the hg19 reference sequence to

correctly code the reference alleles as allele 1, matching the sequencing data.

To generate genotypes at all of the remaining untyped sites, imputation was performed using

BEAGLE 5.0 [15]. This imputation used default settings and included the full 1KGP as the

imputation reference panel. To note, when performing analysis using 1KGP samples, any

related individuals were removed. Each sample then had imputed genotype calls at each of the

imputation SNP loci.

Imputation accuracy and quality assessment

Imputation accuracy for 1KGP and GIAB samples was calculated by comparing imputation

results with previously released genotypes, excluding regions marked as low confidence by

GIAB.

Imputation accuracy on the genotyped samples was assessed on 470,363 sites that were

included on the genotyping array at different allele frequency buckets: 257,362 sites with greater

than 5% allele frequency, 119,978 sites between 1-5% allele frequency, and 93,022 sites with

less than 1% allele frequency. Imputation quality was assessed through site-specific dosage r^2

comparing with genotype values from the genotyping array.

GPS selection

The GPSs for CAD [3], BC [8], and AF [3] were previously published and selected based on

their demonstrated ability to accurately predict and stratify disease risk as well as identify

individuals at risk comparable to monogenic disease. GPS_{CAD} contained 6,630,150

polymorphisms, GPS_{BC} contained 3,820 polymorphisms, and GPS_{AF} contained 6,730,541

polymorphisms. All loci included in these scores were included in the imputation SNP loci.

GPS normalization

In the clinical cohort, raw GPSs were normalized by taking the standardized residual of the

predicted score after correction for the first 10 principal components (PC) of ancestry [16]. PCs

were calculated by projecting lcWGS samples into 10 dimensional PC analysis (PCA) space

using the LASER program [17]. A combination of samples from 1KGP and the Human Origins

[18] project were used as a reference for the projection.

RESULTS

Development and validation of imputation pipeline for lcWGS

Previous studies have evaluated the potential use of lcWGS in local ancestry deconvolution,

complex trait association studies, and detection of rare genetic variants [4–6]. To assess the

feasibility and accuracy of this approach for GPSs, we first developed an imputation pipeline

that reads raw fastq sequence data and generates a vcf with imputed site information at 21.7

million sites (imputation SNP loci) (Figure 1A, B). Briefly, reads are aligned to the reference

genome and filtered for duplicates and low quality. Using this BAM file, we then calculate

genotype likelihoods and impute expected genotypes using 1KGP as the imputation reference

panel.

To validate this imputation pipeline, we performed hcWGS and downsampling on seven

samples from different 1KGP populations and a trio of Ashkenazi Jewish GIAB samples

(pipeline validation data set) to varying depths of coverage from 2.0X to 0.1X (See

Supplementary Table 1, Additional File 2). We used the published genotype calls for each of

these samples as truth data and found that imputation accuracy was above 0.90 r^2 for all

samples at 0.5X and higher (Figure 2). As expected, this was correlated with sequencing depth,

with diminishing gains observed at coverages above 1.0X. While imputation accuracy was

similar across diverse populations, it was slightly reduced in the Colombian sample (HG01485),

likely due to complex local ancestry related to admixture, and in the Yoruban sample

(NA19240), likely due to the shorter blocks of linkage disequilibrium and higher genetic diversity

in Africa [19]. Taken together, these data suggest that at sequencing depth at or above 0.5X,

our pipeline has similar imputation accuracy to genotyping array-based imputation across

individuals from multiple populations. As such, we set 0.5X as a quality control for success and

removed samples with coverage below this threshold in subsequent analyses.

Technical concordance between GPSs calculated from lcWGS and genotyping array

To assess the technical concordance of using lcWGS to calculate GPSs, we performed low

coverage sequencing and used genotyping arrays on DNA from 184 individuals (technical

concordance cohort) (Figure 1B). This concordance assessment was restricted to individuals of

European ancestry to most closely align with the populations used for GPS training and

validation.

We first compared the lcWGS genotype dosages with a subset of variants directly genotyped (n

= 470,362) on the genotyping array to assess imputation performance. Assuming the typed loci

called on the genotyping array as 'true', we observed an average imputation r^2 > 0.90 at 0.5X

depth for variants with global minor allele frequency (MAF) greater than 5% (see Supplementary

Figure 2, Additional File 3). As expected, imputation accuracy was highest for variants with

higher MAF. For lower frequency variants, we saw a reduction in imputation accuracy, as

expected, with r^2 > 0.85 for variants at 1% to 5% MAF and r^2 > 0.80 for variants less than 1%

global MAF. Taken together, this demonstrates that lcWGS has high accuracy in this test

setting.

We then calculated previously published GPSs for CAD [3], BC [8], and AF [3] on each sample

using genotyping array data or lcWGS data. We found that GPS_{CAD}, GPS_{BC}, and GPS_{AF} were

highly correlated (Figure 3A-C), with the score mean (Student t-test p = 0.17) and variance (F

test p = 0.91) equivalent between lcWGS and the genotyping array. The correlations of GPS_{CAD}

and GPS_{AF} (r^2 = 0.98 and r^2 = 0.97, respectively) were slightly higher than that of GPS_{BC} (r^2 =

0.93), which could be due to 1) the smaller number of loci in GPS_{BC} (6.6 million compared to

3820 SNPs), 2) differences in allele frequencies between SNPs with high weights, and/or 3) the

fact that GPS_{BC} was trained and validated on a different genotyping array, the OncoArray, than

the Axiom UK Biobank Array used in this study [8].

The technical concordance cohort ranged in coverage from 0.54X to 1.76X with a mean

coverage of 1.24X, and we have shown that depth can impact imputation performance -- depth

increases above 0.5X have a smaller but measurable effect on imputation performance (Figure

2; see Supplementary Figure 2, Additional File 3). To determine the low coverage sequencing

depth required for GPS accuracy, we used SAMtools to downsample the lcWGS data in this

cohort to 1.0X, 0.75X, 0.5X, 0.4X, 0.25X, and 0.1X. We found that GPS_{CAD}, GPS_{BC}, and GPS_{AF}

are robust to lcWGS sequencing depth 0.5X and that coverages do not systematically bias GPS

calculations in a specific direction (see Supplementary Figure 3 and Supplementary Figure 4,

Additional File 3), indicating that samples above 0.5X with small changes in coverage variation

can be combined for downstream analysis. In addition, the correlation increases logarithmically

as coverage increases (see Supplementary Figure 5, Additional File 3). These data

demonstrate high correlation between GPSs from lcWGS data and genotyping array in a

randomly selected sample. Interestingly, correlation at 0.1X was still high enough that GPSs at

this coverage may have research utility, suggesting that significant amounts of data regarding

common genetic variation could be recovered from off-target reads in exome and multi-gene

panel sequencing studies to allow for GPS calculation. Taken together, these data demonstrate

that lcWGS provides equivalent accuracy for calculation of GPSs, with sequencing coverage as

low as 0.5X.

Assessment of imputation performance and technical concordance across diverse

populations

To further assess the performance of our imputation pipeline across diverse populations, we

performed lcWGS on 120 additional samples from six 1KGP populations (CHB, GIH, YRI, ASW,

MXL, and PUR; see Supplementary Table 3, Additional File 1) that represent the range of

ancestry observed in admixed populations (diverse ancestry data set). We compared genotypes

imputed using our lcWGS pipeline to known 1KGP WGS data and found that imputation

accuracy was above 0.90 r^2 for all samples (range 0.94 - 0.97) (Figure 4A). In addition, we

found that GPS calculated from lcWGS data and GPS calculated from the Phase 3 1KGP WGS

data release have a high correlation, with an r^2 value of 0.98, 0.91, and 0.98 for CAD, BC, and

AF, respectively (Figure 4B-D). These results suggest that lcWGS can enable accurate

imputation and calculation of GPSs in diverse populations.

Association of lcWGS-calculated GPSs with disease phenotypes in a clinical cohort

Previous studies have demonstrated the association of GPSs with prevalent disease using

genotyping arrays [3,8,20–22] and hcWGS [16]. To observe the performance of lcWGS-

calculated GPSs in a large population, we performed low coverage sequencing on 11,502

European individuals (clinical cohort) (See Supplementary Table 2, Additional File 1) and

calculated GPS_{CAD}, GPS_{BC}, and GPS_{AF} for each individual. Raw GPSs were normalized by

taking the standardized residual of the predicted score after correction for the first 10 PCAs (see

Supplementary Figure 6, Additional File 3) [16,23]. First, we note that there are no major outliers

(defined as a z-score greater than 5) in GPS_{CAD}, GPS_{BC}, and GPS_{AF} and that the normalized

scores formed an approximately normal distribution for each (see Supplementary Figure 7,

Additional File 3). Each of the GPSs were strongly associated with self-reported history of

disease, with effect estimates comparable to prior reports using genotyping arrays to calculate

GPS -- GPS_{CAD} (OR per standard deviation = 1.59 (1.32 - 1.92), n = 11,010), GPS_{BC} (OR per

standard deviation = 1.56 (1.45 - 1.68); n = 8722), and GPS_{AF} (OR per standard deviation =

1.28 (1.12 - 1.46); n = 10,303) (Figure 5).

Previous studies have noted significantly increased disease prevalence among individuals in the

extreme tails of the GPS distribution when compared to the remainder of the population [3,8].

We replicated this observation by assessing the prevalence of disease in the highest 5% of the

GPS distribution for each of the three diseases, noting odds ratios of 4.5 (2.62 - 7.74), 2.62

(2.04 - 3.36), and 1.96 (1.24 - 3.11) for GPS_{CAD}, GPS_{BC}, and $GPS_{AF,}$ respectively.

Area under the curve (AUC) is an additional metric used to assess the ability of a given risk

factor to discriminate between affected cases and disease-free controls. When only the GPS

was included in the prediction model, GPS_{CAD} had an AUC of 0.60, GPS_{BC} had an AUC of 0.63,

and GPS_{AF} had an AUC of 0.57. The additional inclusion of age and sex increased the AUCs to

0.86 for GPS_{CAD}, 0.78 for GPS_{BC}, and 0.78 for GPS_{AF}. For each of these three common,

complex diseases, the magnitude of associations with clinical disease and AUC metrics were

consistent with previous publications [3,8]. Taken together, these results suggest that lcWGS-

calculated GPSs can accurately stratify risk with comparable accuracy to previously published

GPS-disease associations calculated on the basis of genotyping array data.

DISCUSSION

For the past two decades, genotyping array-based GWAS and imputation have been the driving

force in our discovery of genetic loci predictive of disease and derivation and calculation of

GPSs. In this study, we developed and validated an imputation pipeline to calculate GPSs from

variably downsampled hcWGS and lcWGS data sets. While the efficiency of lcWGS has been

reported for other applications of statistical genetics [4–6], we demonstrate that lcWGS achieves

similar technical concordance as the Axiom UK Biobank Array by Affymetrix for determining

GPSs. Furthermore, the imputation r^2 from lcWGS was greater than 90%, which is similar to the

imputation accuracy reported from other commercially-available genotyping arrays [24]. Taken

together, these data suggest that lcWGS has comparable accuracy to genotyping arrays for

assessment of common variants and subsequent calculation of GPSs.

Our finding that lcWGS can be used for accurate genotyping and imputation of common genetic

variants has implications for the future of genomic research and medicine. Currently, disease

GWAS are performed using a variety of genotyping arrays that are designed to target specific

sets of genes or features, reducing imputation quality in regions that are not targeted [25].

lcWGS enables less biased imputation than genotyping arrays by not pre-specifying the genetic

content that is included for assessment, as is necessary for genotyping arrays. Because initial

GWAS focused on populations with high homogeneity to reduce noise and increase fit of risk

stratification, many genotyping arrays were designed to capture common genetic variants based

on the linkage disequilibrium structure in European populations [26]. However, this

ascertainment bias reduces the imputation performance from genotyping array data in diverse

populations [27–29]. Imputation from lcWGS data reduces this bias by including all SNPs

observed in 1KGP populations as potential predictors. The effects of SNP selection bias are

also not equivalent across genotyping arrays, and therefore variants included in a GPS trained

and validated on one genotyping array may not be as predictive on another genotyping array

[30]. lcWGS systematically surveys variants independent of SNP selection bias and thus

provides one approach to overcome this issue. Our findings here demonstrate that GPSs

trained and validated on different genotyping arrays are transferable to lcWGS-calculated GPS.

Furthermore, as new genetic associations are discovered, lcWGS can be re-analyzed with ever

more inclusive sets of known SNPs, further reducing SNP selection bias and advancing the

study and understanding of the genetic contributions to disease. In contrast, genotyping arrays

are static and cannot be easily updated or changed without designing a *de novo* platform.

lcWGS also has the potential to easily integrate into current clinical sequencing pipelines. In

contrast to genotyping arrays, which require investment in separate laboratory technology,

lcWGS can be performed on the same platform as current hcWGS or targeted multi-gene panel

clinical testing. The ease of combining these two pathways could help to drive GPS adoption

into clinical practice and can likely be achieved at a cost comparable to genotyping arrays [4].

As the cost of next generation sequencing continues to decrease, the cost of lcWGS will also

continue to decrease.

This study should be interpreted in the context of potential limitations. First, the imputation

accuracy observed in our analysis may have been limited by the reference panel size. Future

efforts using an even larger reference panel may lead to further improved imputation accuracy,

particularly for variants with allele frequency less than 1% [24]. Second, while lcWGS may

ultimately enable derivation of GPSs with improved predictive accuracy or ethnic transferability,

this was not explicitly explored here. Rather, we demonstrate the feasibility and accuracy of

using lcWGS of calculating GPSs published in previous studies. Third, disease phenotypes in

our clinical cohort were based on individual self-report rather than review of health records.

However, several studies have shown that self-reported personal history data have high

concordance with data reported by a healthcare provider or electronic health records [31–34],

and any inaccuracies would be expected to bias GPS-disease associations to the null.

CONCLUSIONS

In conclusion, this work establishes lcWGS as an alternative approach to genotyping arrays for

common genetic variant assessment and GPS calculation -- providing comparable accuracy at

similar cost while also overcoming the ascertainment bias inherent to variant selection in

genotyping array design.

LIST OF ABBREVIATIONS

GPS, genome-wide polygenic score

lcWGS, low coverage whole genome sequencing

CAD, coronary artery disease

BC, breast cancer

AF, atrial fibrillation

1KGP, 1000 Genomes Project

GIAB, Genome in a Bottle

Indel, insertion-deletion

BWA, Burrows-Wheeler Aligner

IBS, Identity-by-State

PC, principal components

PCA, PC analysis

MAF, minor allele frequency

AUC, area under the curve

DECLARATIONS

Ethics approval and consent to participate

All individuals in the technical concordance cohort and clinical cohort gave electronic informed

consent to have their de-identified information and sample used in anonymized studies

(Western Institutional Review Board, #20150716).

Consent for publication

All individuals in the technical concordance cohort and clinical cohort gave electronic informed

consent that Color may author publications using non-aggregated, de-identified information,

either on its own or in collaboration with academic or commercial third parties.

Availability of data and material

The technical concordance and clinical cohort data are not publicly available given the potential

to compromise research participant privacy or consent.

1KGP, http://www.internationalgenome.org/

GIAB, ftp://ftp-trace.ncbi.nlm.nih.gov/giab/ftp/release/AshkenazimTrio/

Samtools/Bcftools, http://www.htslib.org/

BEAGLE, https://faculty.washington.edu/browning/beagle/beagle.html

FastNGSAdmix, http://www.popgen.dk/software/index.php/FastNGSadmix

Competing interests

JRH, CLN, and AYZ are currently employed by and have equity interest in Color Genomics.

JRH has previously consulted for Twist Bioscience and Etalon Diagnostics. GM was previously

employed at Color Genomics and Operator. JRH and GM report a patent application related to

low coverage whole genome sequencing. SK is an employee of Verve Therapeutics and holds

equity in Verve Therapeutics, Maze Therapeutics, Catabasis, and San Therapeutics. He is a

member of the scientific advisory boards for Regeneron Genetics Center and Corvidia

Therapeutics; he has served as a consultant for Acceleron, Eli Lilly, Novartis, Merck, Novo

Nordisk, Novo Ventures, Ionis, Alnylam, Aegerion, Haug Partners, Noble Insights, Leerink

Partners, Bayer Healthcare, Illumina, Color Genomics, MedGenome, Quest, and Medscape; he

reports patents related to a method of identifying and treating a person having a predisposition

to or afflicted with cardiometabolic disease (20180010185) and a genetic risk predictor

(20190017119). AVK has served as a consultant for Color Genomics and reports a patent

related to a genetic risk predictor (20190017119).

Funding

This work was supported by Color Genomics.

Authors' contributions

JRH, GM, AYZ, and AVK designed the overall study. JRH, CLN, GM, AYZ, SK, and AVK

contributed to data acquisition and analysis. JRH, CLN, GM, AYZ, SK, and AVK drafted or

critically revised the manuscript for important intellectual content. AYZ and AVK are the

guarantors of this work and, as such, have full access to all of the data in the study and take

responsibility for the integrity of the data and the accuracy of the data analysis.

Acknowledgements

We would like to thank Will Stedden, Carmen Lai, and Anjali D. Zimmer for helpful discussions

and Justin Lock, Alok Sabnis, and Valerie Ngo for laboratory support and sample processing.

REFERENCES

1. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional

mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis

for the Global Burden of Disease Study 2010. Lancet. 2012;380:2095–128.

2. Boyle EA, Li YI, Pritchard JK. An Expanded View of Complex Traits: From Polygenic to

Omnigenic. Cell. 2017;169:1177–86.

3. Khera AV, Chaffin M, Aragam KG, Haas ME, Roselli C, Choi SH, et al. Genome-wide

polygenic scores for common diseases identify individuals with risk equivalent to monogenic

mutations. Nat Genet [Internet]. 2018; Available from: http://dx.doi.org/10.1038/s41588-018-

0183-z

4. Pasaniuc B, Rohland N, McLaren PJ, Garimella K, Zaitlen N, Li H, et al. Extremely low-

coverage sequencing and imputation increases power for genome-wide association studies. Nat

Genet. 2012;44:631–5.

5. Gilly A, Southam L, Suveges D, Kuchenbaecker K, Moore R, Melloni GEM, et al. Very low

depth whole genome sequencing in complex trait association studies. Bioinformatics [Internet].

2018; Available from: http://dx.doi.org/10.1093/bioinformatics/bty1032

6. Liu S, Huang S, Chen F, Zhao L, Yuan Y, Francis SS, et al. Genomic Analyses from Non-

invasive Prenatal Testing Reveal Genetic Associations, Patterns of Viral Infections, and

Chinese Population History. Cell. 2018;175:347–59.e14.

7. Navon O, Sul JH, Han B, Conde L, Bracci PM, Riby J, et al. Rare variant association testing

under low-coverage sequencing. Genetics. 2013;194:769–79.

8. Mavaddat N, Michailidou K, Dennis J, Lush M, Fachal L, Lee A, et al. Polygenic Risk Scores

for Prediction of Breast Cancer and Breast Cancer Subtypes. Am J Hum Genet. 2019;104:21–

34.

9. Jørsboe E, Hanghøj K, Albrechtsen A. fastNGSadmix: admixture proportions and principal

component analysis of a single NGS sample. Bioinformatics. 2017;33:3148–50.

10. Neben CL, Zimmer AD, Stedden W, van den Akker J, O'Connor R, Chan RC, et al. Multi-

Gene Panel Testing of 23,179 Individuals for Hereditary Cancer Risk Identifies Pathogenic

Variant Carriers Missed by Current Genetic Testing Guidelines. J Mol Diagn [Internet]. Elsevier;

2019 [cited 2019 Jun 11];0. Available from: https://jmd.amjpathol.org/article/S1525-

1578(18)30334-9/fulltext

11. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9.

12. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM

[Internet]. arXiv [q-bio.GN]. 2013. Available from: http://arxiv.org/abs/1303.3997

13. Li H. A statistical framework for SNP calling, mutation discovery, association mapping and

population genetical parameter estimation from sequencing data. Bioinformatics. 2011;27:2987–

93.

14. Browning BL, Browning SR. Genotype Imputation with Millions of Reference Samples. Am J

Hum Genet. 2016;98:116–26.

15. Browning BL, Zhou Y, Browning SR. A One-Penny Imputed Genome from Next-Generation

Reference Panels. Am J Hum Genet. 2018;103:338–48.

16. Khera AV, Chaffin M, Zekavat SM, Collins RL, Roselli C, Natarajan P, et al. Whole Genome

Sequencing to Characterize Monogenic and Polygenic Contributions in Patients Hospitalized

with Early-Onset Myocardial Infarction. Circulation [Internet]. American Heart Association

Bethesda, MD; 2018 [cited 2018 Nov 27]; Available from:

https://www.ahajournals.org/doi/abs/10.1161/CIRCULATIONAHA.118.035658

17. Wang C, Zhan X, Liang L, Abecasis GR, Lin X. Improved ancestry estimation for both

genotyping and sequencing data using projection procrustes analysis and genotype imputation.

Am J Hum Genet. 2015;96:926–37.

18. Lazaridis I, Nadel D, Rollefson G, Merrett DC, Rohland N, Mallick S, et al. Genomic insights

into the origin of farming in the ancient Near East. Nature. 2016;536:419–24.

19. 1000 Genomes Project Consortium, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang

HM, et al. A global reference for human genetic variation. Nature. 2015;526:68–74.

20. Inouye M, Abraham G, Nelson CP, Wood AM, Sweeting MJ, Dudbridge F, et al. Genomic

Risk Prediction of Coronary Artery Disease in 480,000 Adults: Implications for Primary

Prevention. J Am Coll Cardiol. 2018;72:1883–93.

21. Richardson TG, Harrison S, Hemani G, Smith GD. An atlas of polygenic risk score

associations to highlight putative causal relationships across the human phenome [Internet].

bioRxiv. 2018 [cited 2018 Nov 27]. p. 467910. Available from:

https://www.biorxiv.org/content/early/2018/11/11/467910

22. Mavaddat N, Pharoah PDP, Michailidou K, Tyrer J, Brook MN, Bolla MK, et al. Prediction of

breast cancer risk based on profiling with common genetic variants. J Natl Cancer Inst [Internet].

2015;107. Available from: http://dx.doi.org/10.1093/jnci/djv036

23. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal

components analysis corrects for stratification in genome-wide association studies. Nat Genet.

2006;38:904–9.

24. McCarthy S, Das S, Kretzschmar W, Delaneau O, Wood AR, Teumer A, et al. A reference

panel of 64,976 haplotypes for genotype imputation. Nat Genet. 2016;48:1279–83.

25. Voight BF, Kang HM, Ding J, Palmer CD, Sidore C, Chines PS, et al. The metabochip, a

custom genotyping array for genetic studies of metabolic, cardiovascular, and anthropometric

traits. PLoS Genet. 2012;8:e1002793.

26. Lachance J, Tishkoff SA. SNP ascertainment bias in population genetic analyses: why it is

important, and how to correct it. Bioessays. 2013;35:780–6.

27. Wojcik GL, Fuchsberger C, Taliun D, Welch R, Martin AR, Shringarpure S, et al. Imputation-

Aware Tag SNP Selection To Improve Power for Large-Scale, Multi-ethnic Association Studies.

G3 . 2018;8:3255–67.

28. Nelson SC, Doheny KF, Pugh EW, Romm JM, Ling H, Laurie CA, et al. Imputation-Based

Genomic Coverage Assessments of Current Human Genotyping Arrays [Internet]. G3:

Genes|Genomes|Genetics. 2013. p. 1795–807. Available from:

http://dx.doi.org/10.1534/g3.113.007161

29. Carlson CS, Matise TC, North KE, Haiman CA, Fesinmeyer MD, Buyske S, et al.

Generalization and dilution of association results from European GWAS in populations of non-

European ancestry: the PAGE study. PLoS Biol. 2013;11:e1001661.

30. Johnson EO, Hancock DB, Levy JL, Gaddis NC, Saccone NL, Bierut LJ, et al. Imputation

across genotyping arrays for genome-wide association studies: assessment of bias and a

correction strategy. Hum Genet. 2013;132:509–22.

31. Gentry-Maharaj A, Fourkala E-O, Burnell M, Ryan A, Apostolidou S, Habib M, et al.

Concordance of National Cancer Registration with self-reported breast, bowel and lung cancer

in England and Wales: a prospective cohort study within the UK Collaborative Trial of Ovarian

Cancer Screening. Br J Cancer. 2013;109:2875–9.

32. D'Aloisio AA, Nichols HB, Hodgson ME, Deming-Halverson SL, Sandler DP. Validity of self-

reported breast cancer characteristics in a nationwide cohort of women with a family history of

breast cancer. BMC Cancer. 2017;17:692.

33. Kehoe R, Wu SY, Leske MC, Chylack LT Jr. Comparing self-reported and physician-

reported medical history. Am J Epidemiol. 1994;139:813–8.

34. Malmo V, Langhammer A, Bønaa KH, Loennechen JP, Ellekjaer H. Validation of self-

reported and hospital-diagnosed atrial fibrillation: the HUNT study. Clin Epidemiol. 2016;8:185–

93.

ADDITIONAL FILES

Additional File 1

Homburger et al Additional File 1, PDF

Supplementary tables and legends

Additional File 2

Homburger et al Additional File 2, PDF

Supplementary Methods

Additional File 3

Homburger et al Additional File 3, PDF

Supplementary figures and legends

 FIGURE TITLES AND LEGENDS

Figure 1. Study design and imputation pipelines. The study design has four groups: (A) pipeline

validation data set and (B) technical concordance cohort, diverse ancestry data set, and clinical

cohort. The imputation pipeline for each group is depicted. hcWGS, high coverage whole

genome sequencing. lcWGS, low coverage whole genome sequencing. HWE, Hardy–Weinberg

equilibrium. GPS, genome-wide polygenic score. CAD, coronary artery disease. BC, breast

cancer. AF, atrial fibrillation.

Figure 2. Assessment of imputation performance in the pipeline validation data set.

Downsampling from 30X to 0.1X showed that lcWGS accuracy was above 0.90 r^2 for all

samples at 0.5X (n = 4 independent random seeds for each sample and coverage value; error

bars are 95% confidence intervals). The thick brown dashed line is a smoothed trendline of the

average imputation quality while the thin grey dashed line demonstrates previously reported

imputation quality from a genotyping array (r^2 = 0.90) [4]. AJ, Ashkenazi Jewish. CDX, Chinese

Dai in Xishuangbanna, China. CEU, Utah Residents with Northern and Western European

Ancestry. CHB, Han Chinese in Beijing, China. CLM, Colombians from Medellin, Colombia.

GIH, Gujarati Indian from Houston, Texas. TSI, Toscani in Italia. YRI, Yoruba in Ibadan, Nigeria.

Figure 3. Correlation of GPSs between genotyping array and lcWGS in the technical

concordance cohort. (A) GPS_{CAD} calculated using lcWGS was highly correlated (r^2 = 0.98) with

those calculated using genotyping array (n = 182). (B) GPS_{BC} calculated using lcWGS was

highly correlated (r^2 = 0.93) with those calculated using genotyping array (n = 182). (C) GPS_{AF}

was highly correlated (r^2 = 0.97) with those calculated using genotyping arrays (n = 182). x-axis

is the raw GPS calculated from the genotyping array, and y-axis is the raw GPS calculated from

the lcWGS data; raw GPS values are unitless. lcWGS, low coverage whole genome

sequencing. GPS, genome-wide polygenic score. CAD, coronary artery disease. BC, breast

cancer. AF, atrial fibrillation.

Figure 4. Assessment of imputation performance and technical concordance across diverse

populations. (A) GPS_{CAD} calculated using lcWGS data was highly correlated with those

calculated using known 1KGP data (n = 116), with all samples having a correlation coefficient

above 0.90. The thin grey dashed line demonstrates previously reported imputation quality from

a genotyping array (r^2 = 0.90) [4]. (B) GPS_{CAD} calculated using lcWGS data was highly

correlated (r^2 = 0.98) with those calculated using known 1KGP data (n = 116). (C) GPS_{BC}

calculated using lcWGS data was highly correlated (r^2 = 0.91) with those calculated using

known 1KGP data (n = 116). (D) GPS_{AF} was highly correlated (r^2 = 0.98) with those calculated

using known 1KGP data (n = 116). 1KGP, 1000 Genomes Project. lcWGS, low coverage whole

genome sequencing. GPS, genome-wide polygenic score. CAD, coronary artery disease. BC,

breast cancer. AF, atrial fibrillation.

Figure 5. Association of lcWGS-calculated GPSs with disease phenotypes in the clinical cohort.

lcWGS-calculated GPS_{CAD} was associated with personal history of CAD (OR = 1.589 (1.32 -

1.92), n = 11,010, p = 1.32 x 10^{-6}). GPS_{CAD} was adjusted for age and sex. lcWGS-calculated

GPS_{BC} was associated with personal history of BC (OR = 1.56 (1.45 - 1.68); n = 8,722, p = 1.0 x

10^{-16}). GPS_{BC} was calculated only for females and adjusted for age at menarche. lcWGS-

calculated GPS_{AF} was associated with personal history of AF (OR = 1.277 (1.12 - 1.46); n =

10,303, p = 0.000292). GPS_{AF} was adjusted for age and sex. lcWGS, low coverage whole

genome sequencing. GPS, genome-wide polygenic score. CAD, coronary artery disease. BC,

breast cancer. AF, atrial fibrillation.

Figure 1

Figure 4

Figure 5

Odds Ratio per SD Increase in GPS