Molecular and genetic regulation of pig pancreatic islet cell development

Seokho Kim[1,9], Robert L. Whitener[1,9], Heshan Peiris[1], Xueying Gu[1], Charles A. Chang[1],

Jonathan Y. Lam[1], Joan Camunas-Soler[2], Insung Park[3], Romina J. Bevacqua[1], Krissie Tellez[1],

Stephen R. Quake[2,4], Jonathan R. T. Lakey[5], Rita Bottino[6], Pablo J. Ross[3], Seung K. Kim[1,7,8]

[1]Department of Developmental Biology, Stanford University School of Medicine,

Stanford, CA, 94305 USA

[2]Department of Bioengineering, Stanford University, Stanford, CA, 94305 USA

[3]Department of Animal Science, University of California Davis, Davis, CA, 95616 USA

[4]Chan Zuckerberg Biohub, San Francisco, CA 94518, USA.

[5]Department of Surgery, University of California at Irvine, Irvine, CA, 92868 USA

[6]Institute of Cellular Therapeutics, Allegheny Health Network, Pittsburgh, PA, 15212 USA

[7]Department of Medicine, Stanford University School of Medicine, Stanford, CA, 94305 USA

[8]Stanford Diabetes Research Center, Stanford University School of Medicine,

Stanford, CA, 94305 USA

[9]These authors contributed equally

Correspondence and requests for materials should be addressed to S.K.K (email:

seungkim@stanford.edu)

Key Words: pancreas; metabolism; organogenesis; β-cell; α-cell; δ-cell; diabetes mellitus

Summary Statement: This study reveals transcriptional, signaling and cellular programs

governing pig pancreatic islet development, including striking similarities to human islet

ontogeny, providing a novel resource for advancing human islet replacement strategies.

Abstract

Reliance on rodents for understanding pancreatic genetics, development and islet function could limit progress in developing interventions for human diseases like diabetes mellitus. Similarities of pancreas morphology and function suggest that porcine and human pancreas developmental biology may have useful homologies. However, little is known about pig pancreas development. To fill this knowledge gap, we investigated fetal and neonatal pig pancreas at multiple, crucial developmental stages using modern experimental approaches. Purification of islet β-, α- and δ-cells followed by transcriptome analysis (RNA-Seq) and immunohistology identified cell- and stage-specific regulation, and revealed that pig and human islet cells share characteristic features not observed in mice. Morphometric analysis also revealed endocrine cell allocation and architectural similarities between pig and human islets. Our analysis unveiled scores of signaling pathways linked to native islet β-cell functional maturation, including evidence of fetal α-cell GLP-1 production and signaling to β-cells. Thus, the findings and resources detailed here show how pig pancreatic islet studies complement other systems for understanding the developmental programs that generate functional islet cells, and that are relevant to human pancreatic diseases.

Introduction

Progress from studies of pancreas biology in humans (Hart and Powers, 2019; Hrvatin et al., 2014b) has advanced our understanding of post-natal islet regulation and function. The recognition that mouse pancreas biology – while essential – has limitations for understanding human pancreas formation or diseases (Hattersley and Patel, 2017; Maestro et al., 2007), has intensified interest in experimental systems that more closely reflect human pancreas development (McKnight et al., 2010; Pan and Brissova, 2014). There is a specific, crucial knowledge gap in our understanding of human islet and pancreas development from mid-gestation through neonatal and juvenile stages, a period when critical aspects of islet development are known to occur (Arda et al., 2016). The use of human primary tissues to address such questions is limited by inter-individual heterogeneity, unpredictable and restricted access to tissue from key developmental stages, and challenges implementing high-throughput molecular approaches. Likewise, reconstitution of islet development from stem cell sources or fetal human cell lines remains imperfect (Sneddon et al., 2018), thereby limiting interpretation of developmental studies in those systems.

Humans and pigs are omnivorous mammals with similar physiology and metabolic diseases, including diabetes mellitus and its complications, from diet-induced obesity, insulinopathy and β-cell stress (Dyson et al., 2006; Lim et al., 2018; Renner et al., 2013). Recent experimental advances expand possible studies of pig pancreas development to include gain- and loss-of-function genetics (Kemter et al., 2017; Matsunari et al., 2013; Sheets et al., 2018; Wu et al., 2017), fetal surgery and immunomodulation (Fisher et al., 2013), and primary islet cell genetics (Peiris et al., 2018). Prior studies of pig pancreas development have largely relied on immunohistological survey of tissue cell types (Carlsson et al., 2010; Ferrer et al., 2008; Hassouna et al., 2018; Nagaya et al., 2019). Moreover, studies linking hallmark islet functions, like regulated insulin secretion and dynamic changes in gene expression at advancing developmental stages, have not been previously reported (Mueller et al., 2013). Thus, pancreas and metabolic research could benefit from systematic application of powerful methods like cell purification by flow cytometry, high-throughput transcriptome analysis, and islet physiological studies across a comprehensive range of pig developmental stages.

Here we apply these and other modern approaches to delineate pig pancreas development, with a focus on islet β-cells, the sole source of insulin, and α-cells, the principal systemic source of the hormone glucagon. Dysregulation of hormone secretion by these cells is thought to be a leading cause of both type 1 diabetes (T1D) and type 2 diabetes (T2D) in humans (Holst et al., 2017; Lee et al., 2016). Our findings provide evidence of greater similarity between pig and

human β-cells and α-cells than to cognate mouse cells, and demonstrate how studies of pig

pancreatic cells can advance our understanding of the genetic, molecular, signaling and

physiological programs that generate functional pancreatic islet cells.

Results

Pancreas dissociation and FACS purification of islet cells from fetal and neonatal pigs

To measure gene expression and functional changes in the developing pig pancreas, we

established a reliable infrastructure to procure pancreata from fetal or neonatal pigs using

rigorous criteria that optimized tissue quality and gene expression analysis (Fig. 1A). To

investigate the crucial period of ontogeny culminating in functional islet cells, we systematically

isolated pancreata from gestational days (d) 40, 70, 90 and post-natal days (P) 8 and 22 (Fig.

1A). We surveyed sorting methods based on cell surface epitopes, like in prior work with mouse

and human cells (Arda et al., 2016; Arda et al., 2018; Sugiyama et al., 2013), but have not yet

achieved reliable cell fractionation based with this strategy (R. Whitener and S. K. Kim, unpubl.

results). Instead, we used intracellular sorting with specific antibodies against insulin-, glucagon-

and somatostatin- to isolate β-, α- and δ-cells, respectively, from fetal or neonatal pancreata

(Fig. S1A), like in prior work on human islets (Arda et al., 2016; Hrvatin et al., 2014a; Peiris et

al., 2018). Quantitative RT-PCR (qRT-PCR: Fig. 1B) confirmed enrichment or depletion of

expected marker genes for β- (*INS*), α- (*GCG*), δ- (*SST*), ductal (*KRT19*) and acinar cells

(*AMY2*).

To obtain comprehensive gene expression profiles, we performed RNA-Seq on 37 libraries

generated from fetal and neonatal pancreatic cells (Table S1). An average of 7 million reads per

sample library uniquely mapped to a current pig reference transcriptome (Sscrofa11.1).

Unsupervised hierarchical clustering of transcriptome data with Pearson correlation analysis

revealed distinct expression profiles that changed with advancing developmental stage within

specific cell types (Fig. S1B). Principal component analysis with these samples showed that

transcriptomes clustered in the first principal component according to cell type (β-, α- or δ-cells:

PC1, 35% of variance), and according to advancing developmental stage in the second principal

component (PC2, 25% of variance: Fig. S1C). Samples from 'late fetal' stages d70 and d90

clustered closely and separately from d40 samples, and were grouped together for subsequent

analysis (pig development *in utero* averages 114 days). In summary, our cell purification

strategy generated high-quality gene expression profiles of developing pig β-, α- and δ-cells.

High-depth transcriptome maps in pig β-, α- and δ-cells.

To evaluate stage-specific differential gene expression, we analyzed transcriptome data in
early fetal (d40), late fetal (d70 and d90) and neonatal stages (P8-P22) (Fig. 1C-H). We used
the DESeq2 algorithm (Love et al., 2014) to find differentially expressed genes (DEGs)
encoding transcripts whose level changed at least 1.5-fold (adjusted P<0.05) between these
stages in β-, α- or δ-cells. Between early and late fetal stages, we identified 2696 DEGs in β-
cells, with transcripts from 1270 genes increased and 1426 decreased. Between late fetal and
neonatal stages, we identified 1463 DEGs: transcripts from 785 genes increased, and 678
genes decreased (Fig. 1C and Table S2). In α-cells between early and late fetal stages, we
identified 2823 DEGs with transcripts from 1311 genes increased and 1512 genes decreased.
Between late fetal and neonatal stages, we identified 2291 DEGs: transcripts from 1200 genes
increased, and 1091 genes decreased (Fig. 1E and Table S2). In δ-cells between late fetal and
neonatal stages, we identified 1698 DEGs with transcripts from 985 genes increased, and 713
genes decreased (Fig. 1G and Table S2). Among DEGs, we observed subsets of genes
enriched in either fetal or neonatal stages in β-, α- or δ-cells, confirming the presence of
regulated developmental gene expression programs in specific pig islet cell types. Transcript
abundance data from β-, α- and δ-cells are also provided in a searchable dataset (Table S3).
To evaluate coherent patterns of differential gene expression, and effects on signaling
pathways at specific stages, we performed gene ontology (GO) term analysis for biological
processes with Clusterprofiler (Yu et al., 2012). In the transition from the early to the late fetal
stages, we observed enrichment of specific terms linked to islet β- or α-cell development, like
glucose homeostasis, endocrine pancreas development, store-operated calcium entry,
response to glucose, and regulation of hormone secretion (Fig. 1D,F). In the transition from late
fetal and neonatal stages in β-, α- or δ-cells, we observed enrichment of biological processes
distinct from those at earlier developmental stages. These included MAP kinase signaling,
hormone secretion, cAMP metabolic process, and Rho protein signaling transduction (Fig.
1D,F,H).
140
Cell type- and stage-specific gene expression in pig islet β-, α- and δ-cells
Next, we assessed expression of genes thought to regulate development of β-, α- and δ-cells
(Fig. 2A-H). Since transcription factors, including PDX1, NKX6.1 and ARX, are known to govern
β- and α-cell differentiation in mouse and human islet cells (Aguayo-Mazzucato et al., 2011;
Artner et al., 2007; Artner et al., 2010; Collombat et al., 2003; Schaffer et al., 2013), we
assessed expression of transcription factors in pigs. Transcripts encoding the factors PDX1,

MAFA and NKX6.1 were highly enriched in pig β-cells throughout development, while genes

encoding ARX and IRX2 were exclusively expressed in pig α-cells (Fig. 2A, B). Unlike in mice,

MAFB is known to be expressed in both human β- and α-cells (Arda et al., 2016; Blodgett et al.,

2015). Notably, we observed that *MAFB* was expressed in both pig β- and α-cells (Fig. 2A). As

expected, pig δ-cells expressed genes encoding transcription factors such as *HHEX*, *PDX1*, and

the transporter *RBP4*, consistent with previous findings in other systems (Muraro et al., 2016)

(Fig. 2A,C). GLP1R is known to be highly expressed in human β-cells and involved in β-cell

function (Dai et al., 2017). Similar to human, *GLP1R* mRNA was highly expressed in pig β-cells

(Fig. 2D). Urocortin3 (UCN3) secreted by β-cells binds to corticotropin-releasing hormone

receptor 2 (CRHR2) in δ-cells, stimulating somatostatin secretion (van der Meulen et al., 2015).

In accordance with these previous findings, we observed that *UCN3* and *CRHR2* mRNA are

highly expressed in pig β-cells and δ-cells, respectively, raising the possibility that this signaling

axis is conserved in the pig (Fig. 2D).

We also observed dynamic expression of genes regulating islet cell proliferation and function

in developing pig β- and α-cells (Fig. 2E-H). For example, *MKI67*, *BUB1B* and *NUSAP1*, which

encode products known to regulate mitosis and cell cycle progression, declined as β- and α-cell

development advanced (Fig. 2E). To validate these findings, we used immunohistology to

quantify the nuclear factor Ki67, a marker of cell proliferation encoded by *MKI67* in multiple

species, including pigs, humans and mice (Arda et al., 2016; Georgia and Bhushan, 2004; Meier

et al., 2008; Teta et al., 2007). In pigs, pancreas immunostaining revealed a high percentage

(10%) of Ki67[+] early fetal islet cells (d40), that declined to 2% by late fetal and neonatal stages,

then further declined to <0.1% by adulthood (Fig. 2F and Fig. S2). These changes were

mirrored by mRNA levels encoding *MKI67* (Fig. 2E). Thus, we observed evidence of vigorous

fetal-stage proliferation of islet cells that, like in mice and humans, declined further after birth.

Likewise, we found that expression of 'disallowed' genes like *COX5A, SLC16A1*, *MGST3*

and *LDHA*, originally identified in β-cells and thought to restrain β-cell function and maturation

(Lemaire et al., 2016; Pullen et al., 2017; Pullen et al., 2010; Sekine et al., 1994), declined

during development (Fig. 2G). We also detected dynamic or cell type-specific expression of

genes encoding established effectors of islet hormone secretion, like the proprotein convertase

PCSK2, the ion transporter *SLC30A8*, and the glucose transporter *SLC2A2* (Fig. 2H). These

data delineate gene expression changes at genome-scale that orchestrate age-dependent

changes in proliferation and maturation of islet β-, α- and δ-cells. Thus, our studies revealed

both similarities of (1) *cell type-specific* gene expression between pig and human islet β-, α- and

δ-cells, and (2) conserved *stage-specific* gene expression dynamics in pigs.

Stage-specific islet cell allocation and production of transcription factors

To verify further our RNA-Seq findings in developing islets, we performed immunostaining of

pig pancreata at specific developmental stages. At fetal stages, islet β-, α- and δ-cells formed

small clusters, which enlarged subsequently in neonatal and adult pancreas (Fig. 3A-C and Fig.

S3A). Thus, like in other species (Carlsson et al., 2010; Jeon et al., 2009; Jorgensen et al.,

2007), islet cells formed clusters at the inception of their ontogeny, then continued their

morphogenesis in multicellular structures through post-natal stages. In fetal, neonatal and adult

islets, standard morphometry revealed that differentiation of islet cells into specific lineages was

dynamic (Fig. S3A,B). In early fetal pancreas, the number of α-cells is slightly higher than that of

β-cells, while the δ-cell fraction was very low (<4%). However, from late fetal to neonatal stages,

the number of β-cells is higher than that of α-cells, while the number of δ-cells is similar to that

of α-cells (Fig. S3B). Consistent with our RNA-Seq data, antibodies that detect the transcription

factors (TFs) PDX1, NKX6.1 and MAFB showed restricted production in specific islet cell

subsets. In β-cells, we detected nuclear PDX1, NKX6.1. By contrast, we detected MAFB protein

in both β- and α-cells (Fig. 3D-F). Unlike in mice, human β-cells express the transcription factors

SIX3 and SIX2 in an age-dependent manner; moreover human α-cells do not express *SIX3* or

SIX2 (Arda et al., 2016). Immunohistology (Fig. 3G) demonstrated SIX3 production specifically

in a subset of adult β-cells, and qRT-PCR (Fig. 3H) of pig islets revealed adult-specific

production of SIX3. We observed a similar age-dependent expression of *SIX2* (Fig. 3I); the

absence of antibodies for pig SIX2 precluded immunohistology studies. In summary, stage-

specific and cell-specific expression of multiple TFs in pig islets resembled that observed

previously in human islet cell subsets.

Similarities of pig and human β-cells and α-cells

Beyond assessment of TFs, we sought independent evidence of similarities between pig,

human and mouse β-cells. We used unsupervised hierarchical clustering with distance matrix

analysis of our RNA-Seq data, and extant human and mouse islet cell bulk RNA-Seq data (Arda

et al., 2016; Blodgett et al., 2015; Qiu et al., 2017). This analysis suggested that neonatal β-cell

transcriptomes in pig and human were globally more similar than to the β-cell transcriptome of

mice (Fig. 4A). The absence of sufficient fetal or neonatal mouse α- or δ-cell transcriptome data

precluded analogous global comparisons of human, pig and mouse α-cells or δ-cells. GO term

enrichment analysis also revealed differential molecular signatures between neonatal pig and

juvenile human β-cells. Significantly enriched pig pathways were associated with proliferation,

while pathways enriched in human juvenile islet β-cells included those regulating β-cell

secretory function (Fig S4 and Table S5). This is consistent with prior observations, including

relatively higher proliferation in neonatal pig β-cells compared to juvenile human β-cells (Arda et

al., 2016).

Next, we performed pairwise comparison of pig and human (Blodgett et al., 2015) cells from

fetal and postnatal stages (Fig. 4B-G and Table S4). This analysis revealed conserved cell type-

specific and stage-specific regulation of multiple factors. For example, *GLP1R* encodes a G

protein-coupled receptor for the incretin hormone GLP-1. Like in humans (Dai et al., 2017),

GLP1R is expressed in pig β-cells but not α-cells and this expression increases from fetal to

neonatal stages (Fig. 4B). By contrast the G protein coupled receptor encoded by *GPR87* is

expressed in pig and human α-cells but not β-cells and has increasing expression during

postnatal islet maturation (Fig. 4C). Similarities in cell type-specificity or stage-specific

expression for other regulators, including *DNMT1*, *PPID*, *KLF9*, and *DCHS1,* provide further

evidence of conserved genetic programs underlying pig and human β- and α-cell development

(Fig. 4D-G). Taken together, our analysis of specific islet TF's, global gene expression

similarities in β-cells, similarities of developmental islet cell allocation, and of islet morphology

provide evidence of remarkable similarities in pig and human β-cell development.

233   Dynamic gene regulation during pig β- and α-cell development

Our studies of pig islet development provided an unprecedented opportunity to assess gene

expression dynamics throughout fetal and postnatal stages. From the early fetal to neonatal

stage, we identified 3,111 genes in β-cells and 3,668 genes in α-cells with dynamic expression

(Fig. 5A,B and Table S6). Thus, our analysis permitted grouping of genes in 'clusters', based on

the sequence of changes observed between three discrete developmental periods. For example,

we observed genes in β- and α-cells that increased through late fetal stages, then did n̲ot

c̲hange thereafter (cluster 2, 'Up_NC') and other genes whose transcript levels declined through

late fetal stages then did not change thereafter (cluster 7, 'Down_NC') (Table S6). We grouped

genes in 8 clusters (Up_Down, Down_Up, etc.), and GO term analysis of each cluster revealed

dynamic biological processes (Fig. 5A,B and Fig. S5A,B). For example, we observed that the β-

and α-cell "Down_NC" clusters were significantly enriched for genes that regulate cell
proliferation and cell cycle progression (like *CDK2, CDK14* and *CDC20*; Fig. 5A,B and Table
S6). The "Up_NC" cluster in both cell types was enriched for genes associated with multiple
signaling pathways involved in glucose and steroid hormone signaling (Fig. S5A,B). Thus, the
data identified multiple biological processes involved in islet differentiation and maturation that
are dynamically regulated in development.

The frequency of transcript measurement in our workflow also identified unanticipated gene
expression dynamics in developing β- and α-cells, particularly in gene clusters with "Up_Down"
or "Down_Up" trajectories (clusters 3 and 6: Fig. 5C,D). For example, from fetal to neonatal
stages in islet β-cells, we observed increased levels of transcript encoding pyruvate
dehydrogenase kinase 4 (PDK4), a factor previously characterized (Pullen et al., 2017) as
'disallowed' in β-cells (Fig. 5C). Consistent with this, we observed that expression of
PPARGC1A, a known positive regulator of *PDK4* transcription (Wende et al., 2005) also
increased at this stage in β-cell development ("NC_Up" cluster; Table S6). Thus, unlike other
disallowed genes whose expression continuously declined in developing β- and α-cells, *PDK4*
expression appeared to increase after late fetal stages, raising the possibility of as-yet
unidentified functions for PDK4 in pig β-cell development.

In human diabetes, genetic risk may reflect monogenic or polygenic mechanisms
(Cerolsaletti et al., 2019; Hattersley and Patel, 2017; Pearson, 2019; Sanyoura et al., 2018). We
hypothesized that causal genes in monogenic forms of diabetes including neonatal diabetes
mellitus (NDM) and maturity onset diabetes of the young (MODY) might be developmentally
regulated. Thus, we compared expression dynamics of 33 genes, linked by prior studies to
human NDM and MODY (Velayos et al., 2017; Yang and Chan, 2016), and with known pig
orthologues (Table S7). We found ten genes dynamically expressed between fetal and neonatal
stages in pig β-cells. Of those, six genes - *INS, STAT3, NEUROG3, GATA4*, *SLC2A2*, and
PCBD1 - are known to have dynamic expression in human β-cells. In pig α-cells, 13 causal
genes are dynamically expressed, with four - *WFS1, NEUROG3, CEL*, and *GATA6* - known to
have changing expression in human α-cells. We also identified seven genes which are
differentially expressed between the late fetal and neonatal stages in pig δ-cells. Thus our work
reveals pig orthologues of causal genes for MODY and NDM whose expression changes in
development of multiple islet cell types, identifying opportunities for investigating native dynamic
regulation of these genes.

PCSK1 encodes an endopeptidase normally expressed in intestinal L-cells to regulate GLP-1
production (Rouille et al., 1995; Steiner, 1998), and in β-cells for proinsulin processing.
Unexpectedly, we observed a transient increase of *PCSK1* mRNA in fetal α-cells (Fig. 5E);
immunostaining confirmed that PCSK1 protein was produced in a subset of late fetal d70 α-cells
but not in early fetal or adult α-cells (Fig. 5F). Consistent with these findings, ELISA
quantification of bioactive GLP-1 revealed abundant production in d70 islets that decreased
thereafter and was undetectable in adult islets (Fig. 5G). Likewise, *DPP4* encodes a protease
that inactivates GLP-1 and whose expression is extinguished in postnatal human β-cells (Arda
et al., 2016; Blodgett et al., 2015). However, we observed a significant *increase* of *DPP4* mRNA
and protein levels in β-cells at late fetal stages, when β-cells also express *GLP1R* (Fig. 5H,I and
Table S3). Together, these findings provide evidence that fetal α-cells transiently produce GLP-
1, and that GLP-1/GLP1R/DPP4 signaling interactions could regulate fetal β-cell development.
288
Genetic regulation of β-cell functional maturation
Age-dependent enhancement of glucose-regulated insulin secretion has been described in
rodents and humans (Aguayo-Mazzucato et al., 2011; Arda et al., 2016; Avrahami et al., 2015;
Blum et al., 2012; Rorsman et al., 1989). In pigs, prior studies of islet insulin secretion have
focused on postnatal stages (Mueller et al., 2013), but did not compare fetal to neonatal stages.
We compared glucose-stimulated insulin secretion in islets isolated from late fetal (d70) or
weaning-age donors (P22). Glucose or IBMX, a potentiator of intracellular cAMP levels and
insulin secretion, did not significantly increase insulin output by fetal islets in static batch
cultures (Fig. 6A). Moreover, insulin secretion by fetal islets in the 'basal' 2.8 mM glucose
condition was higher than that observed from P22 islets (Fig. 6A). In P22 pig islets, glucose or
glucose + IBMX significantly increased insulin secretion (Fig. 6A), though to a lesser degree
than in human juvenile islets (Arda et al 2016). The pattern of declining basal secretion
accompanied by enhanced glucose-regulated insulin secretion in pig islets is similar to findings
from prior studies of isolated late-fetal and neonatal rodent islets (Sodoyez-Goffaux et al., 1979),
and provides evidence of post-natal pig β-cell functional maturation.
To assess the genetic basis of this maturation, we compared transcriptomes from late fetal
and P22 β-cells (Fig. 6B). DESeq2 analysis revealed significantly increased expression of 924
genes and decreased expression of 895 genes between late fetal (both E70 and E90) and P22
(Table S8). GO term analysis revealed that up-regulated transcripts were encoded by genes
associated with pathways involved in insulin secretion (Fig. 6C and Table S9). Intracellular

cAMP and Ca^{2+} levels are crucial regulators of insulin exocytosis (Tengholm and Gylfe, 2017).

We observed that transcripts of genes encoding channels activated by cAMP, such as HCN4

and KCNQ1 were up-regulated (Fig. 6D). We also found increased expression of genes

regulating cytoplasmic Ca^{2+} concentration or genes involved in Ca^{2+}-mediated signaling

pathways such as CRACR2A and CAMK2D (Fig. 6D). Phosphodiesterases (PDEs), which

induce hydrolysis of cAMP to 5'-AMP, were also up-regulated by stage P22 (Fig. 6C). These

findings are consistent with the view that cAMP signaling 'tone' may be relatively high in fetal

islets, then reduced in the post-natal pig β-cell, a view supported by our quantification of

dynamic basal and glucose + IBMX regulation of insulin secretion at these stages (Fig. 6A).

Consistent with our observation that insulin secretion at basal glucose levels declined during

fetal to postnatal islet development, we observed that pathways involved in glucose processing

such as "glucose catabolic process", "ATP generation from ADP" and "calcium ion regulated

exocytosis" were down-regulated in β-cells (Fig. 6E). Compared to a relative peak in late fetal

development, transcripts of genes regulating glycolysis such as *ENO2, ALDOB, GPD1* and

GCK were down regulated by stage P22 (Fig. 6F). Thus, our data suggests that multiple inter-

related signaling pathways regulating insulin exocytosis continue to mature through weaning-

age (P22). In summary, our combined developmental, functional and molecular investigations

here revealed molecular and genetic factors governing maturation of neonatal β-cells and

unveiled conserved mechanisms underlying islet cell development and function in pigs and

humans.

Discussion

Based on the crucial role of pancreatic islets in human diseases and the expanding catalog of

differences between human and mouse islet developmental biology or regulation (Arda et al.,

2013; Arda et al., 2016; Brissova et al., 2005; Hart and Powers, 2019), there is intense interest in

developing complementary experimental systems for investigating islet development and

maturation. To address this need, we used an integrated approach to delineate pancreas

development in pigs. This multimodal assessment of genetic and developmental phenotypes

across a comprehensive range of fetal and post-natal stages of pig pancreas ontogeny and

maturation revealed multiple similarities in formation and regulation of pig and human islet β-, α-

and δ-cells, including features not observed in mice. Systematic phenotyping of pig pancreas

development - particularly gene expression and identification of developmental signaling

pathways - revealed unexpected dynamic gene regulation, and evidence of native intra-islet

GLP-1 signaling. Moreover, studies in pigs are cost-efficient (see Methods). Our experimental

approaches and findings provide a unique roadmap for diabetes and metabolic research,
including attempts to direct differentiation and maturation of replacement islet cell types from
renewable human cell sources.

Here, we established cell purification strategies based on flow cytometry to investigate
developmental genetics in fetal and neonatal pig β-, α- and δ-cells, an approach we and others
have successfully used in pancreas from mice and humans (Arda et al., 2016; Blodgett et al.,
2015; DiGruccio et al., 2016; Hrvatin et al., 2014a; Qiu et al., 2017). Like in prior studies (Arda
et al., 2016; Hrvatin et al., 2014a), we isolated primary β-, α- and δ-cells throughout fetal and
postnatal pancreas development using antibodies against hormones specific to these cell types,
an approach permitting comprehensive, high-quality stage-specific transcriptome analysis.
Similar approaches for primary ductal and acinar cells should provide opportunities in the future
to delineate the transcriptome of these important components of the exocrine pancreas.
Moreover, discovery of flow cytometry-based methods for purifying live primary endocrine or
exocrine cells, with antibodies that recognize cell-surface epitopes, should expand our capacity
to investigate other important aspects of pancreatic gene regulation and function, using
approaches like ATAC-Seq, promoter capture Hi-C, and Patch-Seq (Arda et al., 2018;
Camunas-Soler et al., 2019; Miguel-Escalada et al., 2019).

Transcriptome analysis described here provides evidence of genetic regulatory mechanisms
and cell development in pigs that are also conserved in other mammals, including (1) cell type-
specific expression of transcription factors (*PDX1, MAFA, ARX, IRX, HHEX*) and functional
regulators (*PCSK1, PCSK2, SLC2A2, SLC30A8, GLP1R, GCGR*), (2) transient β-, α- and δ-cell
proliferation, peaking in fetal/perinatal stages and regulated by conserved factors like *MKI67,*
BUB1B and *NUSAP1* (Georgia and Bhushan, 2004; Meier et al., 2008; Teta et al., 2007), (3)
dynamic expression of histone or chromatin regulators like *EZH2* and *DNMT1* (Chakravarthy et
al., 2017; Dai et al., 2017) and (4) reduction or elimination of transcripts encoding 'disallowed'
genes (Lemaire et al., 2017; Pullen et al., 2017; Pullen et al., 2010; Thorrez et al., 2011). These
findings from transcriptome analysis also correspond well with observations from our other
stage-specific tissue phenotyping (see below). Stage-specific gene expression patterns
revealed that pig and human β-cells and α-cells shared characteristic molecular and
developmental features not observed in mouse β-cells or α-cells. Like in human islets (Benner
et al., 2014), the transcription factor MAFB is expressed in both pig β- and α-cells (Figs. 2,3).
Similarly, we found that the transcription factor SIX3 is expressed in pig β-cells in an age-
dependent manner, like in human islets (Arda et al., 2016) (Fig. 3). Thus, studies of
mechanisms governing cell type- and age-dependent expression of genes like pig *MAFB* and

SIX3 could reveal the basis of β-cell functional maturation. Likewise, we observed features of
pig fetal islets not yet noted in humans. *MAFA* expression in pig β-cells increased in fetal stages,
and continued through postnatal stages. In humans, β-cell *MAFA* expression begins in fetal
stages (Blodgett et al., 2015) and also increases after birth (Arda et al., 2016), but its fetal
regulation remains unreported. Our studies also provide a reference transcriptome analysis of
developing islet δ-cells, whose recognized role in health and diabetes is growing (Rorsman and
Huising, 2018). Together, the resources and methods described here provide evidence that
developmental genetic studies of pancreas in pigs could complement similar studies in humans
and mice.

In addition to the resemblance of pancreatic gene regulation in pigs and humans (Figs. 2,4),
we observed similarities of islet morphogenesis, cell specification, regulation and islet cell
allocation. Scattered islet β-, α- and δ-cells coalesced into small clusters at late fetal stages,
which contained proliferating cells (Figs. 2,3). At birth, the proportion of α- and δ-cells in pig
islets was ~50%, like that of neonatal human islets , and unlike in mice. In pig pancreas
development, a large portion of fetal hormone-expressing islet cells appeared to be contained
within small, dispersed clusters of varying cell composition. After birth, these clusters enlarged,
and the geometry of β-, α- and δ-cells in islets appeared intermingled, like in human islets
(Brissova et al., 2005) (Fig. 3). Studies of insulin secretion further demonstrated that β-cell
function matures from fetal to post-natal stages like in humans (Arda et al., 2016) and other
mammals. Thus, our study reveals multiple features in pig islet development that are closely
conserved in human pancreas, including a subset of phenotypes not observed in rodent islets,
suggesting that pigs could be a reliable surrogate animal model to study the islet function of
human gene orthologs. This includes studies of dominant forms of diabetes like MODY and
NDM, whose genetics have not been reconstituted in mice (Hattersley and Patel, 2017; Maestro
et al., 2007). By contrast, dominant monogenic diabetes resulting from mutations in *INS* or the
MODY3 gene *HNF1α* has been reconstituted in pigs (Renner et al., 2013; Umeyama et al.,
2009). Thus, studies of gene function and pathogenesis in diseases like MODY and NDM could
be advanced through investigations in pigs.

Here, we evaluated pancreas phenotypes through a full range of fetal and peri-natal stages.
In β-, α- and δ-cells, this 'granularity' of phenotyping revealed changes in >6000 genes during
development, and identified developmental dynamics not previously noted in studies of human
islet gene expression with comparatively limited sampling during developmental stages (Arda et
al., 2016; Blodgett et al., 2015; Bramswig et al., 2013), or in one study, a single stage (Ramond

et al., 2018). For example, in β-cells, genes involved in glucose processing, including *G6PC2*

and *PDK3*, are in the "Up_NC" cluster whereas genes regulating insulin secretion, such as

SLC2A2, are in the "NC_Up" cluster (Table S6), indicating that mechanisms governing glucose

sensing and metabolism, or insulin secretion, are established at different developmental stages

in pig β-cells. Moreover, we noted 366 β-cell or α-cell transcripts whose levels changed with a

'discontinuous' trajectory in development (Fig. 5 and Table S6). That is, we observed

heterogeneous patterns of expression, including early rise then fall (like *DPP4, FEV* in β-cells,

and *FOXO1* or *KIT* in α-cells), or initial decline followed by a later rise in transcript levels,

including those encoding a subset of disallowed gene products like PDK4 in β-cells (Fig. 5 and

Table S6). Thus, our ability to detect dynamic gene expression trajectories in β-cell or α-cells

was enhanced by the comprehensiveness of developmental phenotyping afforded in pigs.

Cell- and stage-specific phenotyping here also revealed links between our data to human

diabetes risk and islet replacement. Prior studies have noted a significant enrichment of

candidate diabetes risk genes among those with dynamic developmental expression (Arda et al.,

2016; Perez-Alcantara et al., 2018). Similarly, in our recent work, prioritization of candidate

diabetes risk genes with dynamic fetal or post-natal expression led to identification of SIX2,

SIX3 and BCL11A as unrecognized regulators of human β-cell development and function (Arda

et al., 2016; Peiris et al., 2018). Here, we found significant enrichment of β- and α-cell gene sets

involved in specific developmental signaling pathways (Fig. 5). Of these, only a subset, such as

mTOR, retinoic acid receptor and TGF-β signaling, have previously been linked to β- and α-cell

biology (Blandino-Rosano et al., 2017; Brun et al., 2015; Lin et al., 2009; Yokoi et al., 2016).

Thus, our datasets provide a foundation for systematic investigations of *cis*- and *trans*-

regulatory elements governing the dynamic expression of diabetes risk genes in the native,

physiological setting of pig islet development.

Identifying signals controlling islet development, including detection of signaling modulation,

should enhance efforts to control and direct differentiation of functional replacement islet cells

from stem cells and other renewable sources. Recent studies have explored use of neonatal pig

islet cultures for identifying signals that promote β-cell functional maturation (Hassouna et al.,

2018). In their work with stepwise directed differentiation of human stem cells into distinct

lineages, Loh and colleagues have demonstrated the value of inhibiting signaling pathways –

not merely through signal withdrawal, but by exposure of cells to potent signaling inhibitors – to

prevent development along undesired fates or lineages (Ang et al., 2018; Loh et al., 2014). The

identification of dynamically regulated genes and pathways, including those with discontinuous

trajectories (Up then Down, etc.), in β-cell and α-cell development provides a new roadmap for
modulating islet cell replacement efforts. Fetal and neonatal β-cells and α-cells develop in close
proximity, and our work likely identifies intercellular signaling pathways (like those regulated by
PCSK1, *GLP1R* and *DPP4*) that could regulate differentiation of these cells. PCSK1 expression
in mouse fetal α-cells was reported previously (Wilson et al., 2002) but active GLP-1 production
from fetal islets was not assessed; likewise, to our knowledge, human fetal islet GLP-1
production has not been reported. Recent studies have demonstrated that GLP1R can be
stimulated by glucagon, in addition to GLP-1 (Capozzi et al., 2019; Svendsen et al., 2018; Zhu
et al., 2019); however, it remains unknown if all signaling downstream of GLP1R is activated by
glucagon, or if islet cells are the sole target of GLP-1 signaling in the fetal pancreas. Further
studies of endogenous GLP-1 signaling in pig fetal islet development could also clarify the
emerging concept of intra-islet GLP-1 signaling in metabolically-stressed adult islets (Capozzi et
al., 2019; Drucker, 2013). Thus, our findings identify multiple pathways associated with fetal and
post-natal β-cell maturation or function (Table S9), including many not directly modulated in
prior studies (Nair et al., 2019; Pagliuca et al., 2014; Rezania et al., 2014; Russ et al., 2015). In
summary, data, conceptual advances and resources detailed here provide an unprecedented
molecular, cellular and developmental framework for investigating pancreas development that
could inform our understanding of diabetes and obesity risk, and advance efforts to produce
replacement human islets for diabetes.
462

Materials and methods

Animals and pancreas harvest

Pigs used in this study were outbred and used in accordance to protocols approved by the Institutional Animal Care and Use Committees at University of California Davis and Stanford University. For collection of fetal pig pancreata, pregnancies were confirmed via ultrasound 30 days after mating and pregnant gilts were humanely euthanized at 40, 70, or 90 days. The reproductive tracts were removed after mid-ventral laparotomy, cooled on ice, and fetuses were collected. Fetuses remained on ice until dissection. Pancreas was identified according to anatomical location and harvested under a stereoscopic microscope. Neonatal pancreata were harvested from neonatal piglets at 8 and 22 days of age, while adult pancreata were collected from pigs 6 months of age or older. Briefly, after euthanasia, a midline incision was performed to provide access to the peritoneum and thoracic cavity. The pancreas was rapidly chilled via either perfusion of sterile ice-cold saline solution into the descending aorta or by pouring the same solution directly into the peritoneal cavity and then surgically removed. Harvested pancreatic tissue was maintained at a temperature of 2-10°C in saline solution until further downstream processing.

Generation of pancreatic single cell suspensions

Morphological differences of the pancreas between developmental stages necessitated customized protocols to generate single cell suspensions suitable for downstream assays. Embryonic day 40 pancreata were diced finely with a razor blade and transferred into a 4°C solution of Collagenase D (Millipore-Sigma, Cat# C9263) at a concentration of 1 mg/ml in HBSS, and digested to single cells overnight at 4°C. Embryonic day 70 and 90 pancreata were finely diced with a razor blade and digested for 5-8 minutes at 37°C using the same Collagenase D solution above. The reaction was stopped with the addition of equal volume HBSS containing 10% porcine serum (Millipore-Sigma, Cat# P9783). If necessary, an additional digestion using TrypLE (Gibco, Cat#12605-010) for 3 minutes at 37°C was used to fully disperse any remaining cell clusters. Post-natal pancreata were digested following published methods (Lamb et al., 2014). (Logistical and technical challenges related to the mass of adult pigs precluded isolation of single cell islet suspensions of sufficient quality suitable for molecular analysis of adult pig islets.) Briefly, pancreata were trimmed to remove lymphatic, connective, and fat tissues, and chopped into 1-5 mm pieces. The tissue was digested using 2 mg/ml Collagenase D in HBSS at 37°C for 12-18 minutes. 1.5 volumes of HBSS with 10% porcine serum and 0.5% Antibiotic/Antimycotic solution was added to stop digestion. Neonatal Islet Cell clusters (NICCs)

of the desired size were obtained after straining digest through a 500 μm mesh (PluriSelect,
Cat# 43-50500-03) to remove large undigested tissue, washed, and allowed to gravity settle two
times. To generate single cell suspensions, some NICCs were further dispersed with TrypLE for
3 minutes at 37°C. Remaining NICCs were spun down and resuspended in RPMI 1640 (Gibco,
Cat# 11879-020) containing 5 mM glucose, 10% porcine serum, 10 mM HEPES (Caisson Labs,
HOL06-6X100ML) and 1% Pen/Strep (Life Technologies, Cat# 15140-122) and placed in a
humidified, 5% CO_2 tissue culture incubator.

The cost basis of pancreas procurement and islet isolation from pigs is currently lower than
for humans. In 2019, reimbursement for a single human cadaveric donor pancreas is between
$4,500 and $7,000 (US dollars). For human islets, investigators are charged $0.12 per islet
equivalent using the NIH-supported Integrated Islet Distribution Program (https://iidp.coh.org).
By contrast, the cost of a P22 pig pancreas is approximately $500, and after processing, a P22
pancreas yields approximately 90- to 100,000 NICCs. The total cost per pig NICC is <$0.008.
510

Cell flow cytometry and purification
Single cell suspensions were washed twice with cold PBS and filtered with a 70 µm filter (BD
Falcon, Cat# 352350). Prior to fixation, cells were stained with LIVE/DEAD fixable Aqua
(ThermoFisher, Cat# L34976) or LIVE/DEAD fixable Near-IR (ThermoFisher, Cat#L10119) dead
cell stains according to manufacturer's protocol. Cells were fixed with 4% PFA in PBS (Electron
Microscopy Sciences) for 10 minutes at room temperature. After fixation, cells were
permeabilized with Permeabilization Buffer: 1X PBS containing permeabilization solution
(BioLegend, Cat# 421002), 0.2% BSA (Sigma-Aldrich, Cat#A9576), mouse and rat IgG
(Jackson Labs, Cat# 015-000-003 and 012-000-003), and 1:40 Ribolock RNase Inhibitor (Fisher
Scientific, Cat# FEREO0384), for 30 minutes at 4°C. Cells were pelleted and resuspended with
Wash Buffer: 1X PBS containing permeabilization solution, 0.2% BSA, and 1:100 Ribolock.
Cells were pelleted again, supernatant removed, and stained with Alexa Fluor 488 labeled anti-
insulin (R&D Systems, IC1417G-100UG), Alexa Fluor 647 labeled anti-glucagon (Santa Cruz,
sc-57171 AF647) and PE labeled anti-somatostatin (Santa Cruz, sc-55565 PE) antibodies in
Staining Buffer: 1X PBS containing permeabilization solution, 1% BSA, and 1:40 Ribolock, for
30 minutes at 4°C. Cells were washed three times in Wash Buffer and resuspended in Sort
Buffer: 1X PBS containing 0.5% BSA, and 1:20 Ribolock. Labeled cells were sorted on a
special-order, 5-laser FACS Aria II using FACSDiva software (BD Biosciences). After excluding
doublets and dead cells, desired cell populations were identified using fluorescence minus one
(FMO) controls. Cells were sorted using the purity mask in chilled low retention tubes containing

50 μl of Sorting Buffer. We attempted to isolate cells producing more than one islet hormone

(insulin, glucagon or somatostatin) simultaneously, which have been reported in both human

and pig fetal pancreas (Lukinius et al., 1992; Riopel et al., 2014). In contrast to these studies,

however, the relative frequency of these cells after FACS was quite low (<0.001% of live cells)

and precluded downstream analysis. Similarly, the relative paucity of δ cells in the early fetal

porcine pancreas has proven a significant hurdle to collection of sufficient numbers for

downstream analysis. Post-hoc visualization of recorded data was performed using FlowJo vX

software (FlowJo, LLC).

Total RNA extraction and quantitative Reverse Transcription PCR (qRT-PCR)

Total RNA was extracted from the sorted cells as described (Hrvatin et al., 2014a) and

concentrated using the RNA Clean & Concentrator-5 kit (Zymo Research, Cat# R1013).

Complementary DNA (cDNA) was generated using the Maxima First Strand cDNA Synthesis kit

(Thermo Scientific, Cat# K1642) according to the manufacturer's instructions. Purity of sorted

populations was confirmed via Quantitative PCR using an Applied Biosystems 7500 Real-Time

PCR System. The following TaqMan probes were used: INS (Ss03386682_u1), GCG

(Ss03384069_u1), SST (Ss03391856_m1), AMY2 (Ss03394345_m1), ACTB (Ss03376081_u1),

KRT19 (Forward: 5'-GAAGAGCTGGCCTACCTG-3', Reverse: 5'-ATA

CTGGCTTCTCATGTCGC-3'), SIX2 (Forward: 5'-TCAAGGAAAAGAGTCGCAGC-3', Reverse:

5'-TGAACCAGTTGCTGACCTG-3') and SIX3 (Forward: 5'-AACAAGCACGAGTCGATCC-3',

Reverse: 5'-CCACAAGTTCACCAAGGAGT-3'). Relative mRNA abundance was calculated by

the Comparative Ct (ΔΔCt) relative quantitation method.

RNA-Seq Library preparation

Multiplexed libraries were prepared using the SMARTer Stranded Total RNA-Seq Kit v2 - Pico

Input Mammalian (TakaraBio, Cat# 634413) following manufacturer's instructions. Library

fragments of approximately 350 bp were obtained, and the quality was assessed using a 2100

Bioanalyzer (Agilent). Barcoded libraries were multiplexed and sequenced on an Illumina

NextSeq sequencer.

Bioinformatic and statistical analysis of RNA-Seq datasets

RNA-Seq libraries were sequenced as 75 bp pair-end reads to a depth of 30 to 60 million reads.

To estimate transcriptome abundance, Salmon algorithm (v0.11.3) was used (Patro et al.,

2017). An index was built on the Sus Scrofa reference transcriptome (Sscrofa 11.1, version 96)

(Zerbino et al., 2018) using Salmon with parameters "salmon index --keepDuplicates -t

transcripts -i transcripts_index --type quasi -k 31". The reads were aligned to the index and

quantified using Salmon with parameters "salmon quant -i $salmon_index -l A \-1 $Input1 \-2

$Input2 \ p 8 -o quants/${Input1}_quant \ --gcBias –seqBias". After getting the transcript counts,

differentially expressed gene analysis was performed using the DEseq2 analysis pipeline (Love

et al., 2014; Soneson et al., 2015). Differentially expressed genes (DEGs) were obtained by

comparisons of any two sequential stages (i.e. Early fetal versus Late fetal and Late fetal versus

Neonatal) in a cell type specific manner. Genes were considered differentially expressed if the

fold change between samples was greater than or equal to 1.5 with setting of Benjamini-

Hochberg (BH) false discovery rate (FDR) adjusted P value of <0.05 in the package.

Hierarchical clustering of samples were done with the cor() functions of R. Principal component

analysis (PCA) was performed with the function *plotPCA* that comes with DEseq2 package.

Visualization of RNA-seq data examined in this study were produced with pheatmap 1.0.12

(Kolde, 2012) and ggplot2 (Wickham, 2009). GO term analysis was carried out using the R

package clusterProfiler as described (Yu et al., 2012) with a Benjamini-Hochberg adjusted P

value of <0.1.

Comparison to extant human and mouse β-cell transcriptomes

Human (Arda et al., 2016; Blodgett et al., 2015) (GEO accession numbers GSE67543 and

GSE79469) and mouse (Qiu et al., 2017) (GEO accession number GSE87375) datasets were

obtained from the NCBI Gene Expression Omnibus (GEO). Indices for human and mouse

libraries were built from reference transcriptomes GRCh38 cDNA and GRCm38 cDNA,

respectively (Homo sapiens, version 96; Mus musculus, version 96; (Zerbino et al., 2018)). Pig

and mouse Ensembl IDs were converted to their corresponding human Ensembl ID using

predicted ortholog tables across these species obtained from Ensembl BioMart (Zerbino et al.,

2018). Sample-wise hierarchical clustering analysis (Fig. 4A) was performed using dist()

function of R.

Identifying genes with dynamic expression

To create lists of dynamically expressed genes, DEGs from comparisons between early fetal

and late fetal stages and between late fetal and neonatal stage were intersected, which created

lists of genes showing patterns of gene expression with up, down or no change directions as

development proceeds in islets development. Gene TPM values shifted by +1, log based 2

transformed were averaged across biological replicates and subsequently standardized to their
Z-scores.
600

Identification of developmentally regulated MODY and NDM genes

602   We queried 33 genes previously linked to MODY and/or NDM (Velayos et al., 2017; Yang
603   and Chan, 2016), and identified 19 dynamically expressed genes during the transition from fetal
604   to neonatal stages in pig β-, α- or δ-cells. Among these 19 candidates, we then used prior data
605   sets (Blodgett et al., 2015) to identify a subset that also have dynamic expression in human islet
606   β- or α-cells – a lack of available human δ-cell datasets precludes this comparison for δ-cells.
607

Immunohistochemistry

Biopsies of harvested pancreata were fixed in 4% paraformaldehyde (w/v) in PBS at 4°C
overnight, washed twice in PBS, and cryoprotected via sucrose saturation. The tissues were
imbedded in Tissue-Tek OCT compound (VWR, Cat# 25608-930) and stored at -80°C until
sectioning. 10μm thick sections were collected on ThermoFisher Superfrost Plus slides. For
staining, frozen slides were thawed at room temperature for 30 min and washed with PBS.
Antigen retrieval was performed by boiling slides immersed in Target Retrieval Solution (DAKO,
S169984-2) for 30 minutes, as needed. Subsequently, sections were incubated with
streptavidin/biotin blocking solution (Vector Labs, SP-2002) according to manufacturer's
instruction. Then, sections were incubated in permeabilization/blocking buffer (5% normal
donkey serum, 0.3% Triton X-100 in PBS) at room temperature for 1 hour followed by overnight
incubation at 4°C with primary antibodies diluted in Antibody dilution buffer (1% bovine serum
albumin, 0.3% Triton X-100 in PBS). Slides were washed and stained with appropriate
secondary antibodies for 1 hour at room temperature. Primary antibodies and dilutions used are
as follows: guinea pig anti-insulin (DAKO, A0564, 1:1000), mouse anti-glucagon (Sigma-Aldrich,
G2654, 1:1000), rabbit anti-SIX3 (LifeSpan Biosciences, LS-B9336, 1:100), goat anti-PDX1
(R&D systems, AF2419, 1:200), mouse anti-human NKX6.1 (R&D systems, AF5857, 1:200,)
rabbit anti-MAFB (Bethyl, IHC-00351, 1:200), rabbit anti-MAFA (AVIVA systems biology,
ARP47760_P050, 1:200), goat anti-somatostatin (Santa Cruz biotechnology, 1:500), rabbit anti-
Chromogranin A (Immunostar, 20085, 1:500), rat anti-E-cadherin (Life Technologies, 13-1900
1:500).   Secondary antibodies used are donkey anti-isotype conjugated with Alexa Fluor
488/555/647 (1:500; Molecular Probes). After a final wash with PBS, slides were preserved with
a mounting medium containing DAPI (Vector labs, H-1200) and coverslipped. Image acquisition
was performed on a Leica SP8 confocal microscope. Islet cells were quantified by counting

insulin positive β-cells, glucagon positive α-cells, and somatostatin positive δ-cells using

pancreas sections from different stages.

Morphometric Analysis

Tissue sections were prepared as above. Single-channel images were captured using a Zeiss

Axio Imager M2 microscope using a 20X objective with the AxioVision version 4.8 software.

Respective channel images were then imported into the Image-Pro Plus version 5 software

(Media Cybernetics, Inc.). Quantification of hormone$^+$ cells was performed by using the count

function within the Measure menu. The number of individual hormone$^+$ cells were then

summated for each developmental stage, and the fractional percentage comprising each of the

three cell types was calculated. To determine the percentage of β-, α-, and δ-cells which were

undergoing cell division at each developmental stage, slides were stained for a combination of

insulin, glucagon, and Ki67 or insulin, somatostatin, and Ki67. Single-channel images were

collected as above and then merged into a multi-channel image using the Image-Pro Plus

software. Cells which were Ki67$^+$ and hormone$^+$, as well as the total number of hormone$^+$ cells,

were then counted manually.

In vitro insulin secretion assay

Functional maturity of NICCs isolated from late fetal or 22-day old piglets was assessed by in

vitro glucose stimulated insulin secretion (GSIS) assay performed after 5 days of culture. In brief,

100 NICCs were washed and incubated at 2.8 mM glucose for 1 hour. After the initial

preincubation, clusters were serially incubated in media above containing 2.8 mM, 20 mM, and

20 mM plus IMBX glucose concentrations for 1 hour each. Finally, clusters were lysed in acid-

ethanol solution to extract the total insulin content. Insulin from supernatants and the NICC

lysates was quantified using the pig insulin ELISA kit (Mercodia, 10-12000-01). Secreted insulin

is presented as a percent of total insulin detected in lysate. All steps were performed in triplicate

at 37°C using RPMI 1640 supplemented with 2% porcine serum adjusted to the respective

glucose concentrations.

In vitro active GLP-1 assay

Islet cell clusters were handpicked immediately after isolation, washed twice in ice cold PBS,

then resuspended in 4°C lysis buffer containing DPP4 inhibitor (EMD Millipore, DPP4). Clusters

were sonicated at 4°C and immediately stored at -80°C. Active GLP-1 (7-36) was measured

using the Mouse / Rat GLP-1 Active (7-36) ELISA Kit (Eagle Biosciences, GP121-K01).

Statistical analysis

667 For qRT-PCR, GSIS and quantification of immunohistology experiments, the number of
668 biological or technical replicates (n), measure of central tendency (e.g. average), standard
669 deviation (SD), and statistical analysis is detailed in the Figure legend. Graphs and statistical
670 analysis were produced and performed by using GraphPad Prism (version 8) software.

Acknowledgements

We thank past and current members of the Kim group for advice and encouragement, Dr. Sam
Baker (Stanford, Dept. of Comparative Medicine, Veterinary Service Center), Dr. Jing Wang
(Stanford Diabetes Research Center and Stanford Islet Research Core) and Michael Alexander
(U.C. Irvine) for assistance with pig pancreas work, Serena Whitener for help developing the pig
pancreas gene expression browser, Dr. Kyle Loh for stimulating discussions, and Drs. Dan Lu
and Ramesh Nair for help on bioinformatics.

Competing interests: Nothing to declare

Funding

This work was supported by a Deans Innovation Fund from the Dept. of Developmental Biology
(Stanford), and by fellowship awards from the L.L. Hillblom Foundation (2017-D-008-FEL) to
S.K., from the Division of Endocrinology T32 training grant (DK007217-41, A. Hoffman and F.
Kraemer) to R.L.W., and from the Stanford Child Health Research Institute (UL1 TR001085)
and the American Diabetes Association (1-16-PDF-086) to H.P. Work in the Kim lab was also
supported by grants from the U.S. National Institutes of Health P30 DK116074, and by the
Stanford Islet Research Core and Diabetes Genomics and Analysis Core of Stanford Diabetes
Research Center.

Data Availability

694 The data discussed in this publication have been deposited in NCBI's Gene Expression
695 Omnibus (Edgar et al., 2002) and are accessible through GEO Series accession number
696 GSE143889 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc= GSE143889).

698 **Requests for materials & correspondence** should be addressed to Seung K. Kim.

Author contributions

S.K., R.L.W. and S.K.K. designed the study; S.K, R.L.W, X.G., C.A.C., J.Y.L., I.P., R.J.B., and K.T. performed the experiments; J.C., S.R.Q., J.R.T.L., R.B., and P.J.R. shared materials; S.K., R.L.W., H.P., and S.K.K analyzed the data and wrote the manuscript.

References

Aguayo-Mazzucato, C., Koh, A., El Khattabi, I., Li, W. C., Toschi, E., Jermendy, A., Juhl, K., Mao, K., Weir, G. C., Sharma, A., et al. (2011). Mafa expression enhances glucose-responsive insulin secretion in neonatal rat beta cells. *Diabetologia* **54**, 583-593.

Ang, L. T., Tan, A. K. Y., Autio, M. I., Goh, S. H., Choo, S. H., Lee, K. L., Tan, J., Pan, B., Lee, J. J. H., Lum, J. J., et al. (2018). A Roadmap for Human Liver Differentiation from Pluripotent Stem Cells. *Cell Rep* **22**, 2190-2205.

Arda, H. E., Benitez, C. M. and Kim, S. K. (2013). Gene regulatory networks governing pancreas development. *Dev Cell* **25**, 5-13.

Arda, H. E., Li, L., Tsai, J., Torre, E. A., Rosli, Y., Peiris, H., Spitale, R. C., Dai, C., Gu, X., Qu, K., et al. (2016). Age-Dependent Pancreatic Gene Regulation Reveals Mechanisms Governing Human beta Cell Function. *Cell metabolism* **23**, 909-920.

Arda, H. E., Tsai, J., Rosli, Y. R., Giresi, P., Bottino, R., Greenleaf, W. J., Chang, H. Y. and Kim, S. K. (2018). A Chromatin Basis for Cell Lineage and Disease Risk in the Human Pancreas. *Cell Syst* **7**, 310-322 e314.

Artner, I., Blanchi, B., Raum, J. C., Guo, M., Kaneko, T., Cordes, S., Sieweke, M. and Stein, R. (2007). MafB is required for islet beta cell maturation. *Proc Natl Acad Sci U S A* **104**, 3853-3858.

Artner, I., Hang, Y., Mazur, M., Yamamoto, T., Guo, M., Lindner, J., Magnuson, M. A. and Stein, R. (2010). MafA and MafB regulate genes critical to beta-cells in a unique temporal manner. *Diabetes* **59**, 2530-2539.

Avrahami, D., Li, C., Zhang, J., Schug, J., Avrahami, R., Rao, S., Stadler, M. B., Burger, L., Schubeler, D., Glaser, B., et al. (2015). Aging-Dependent Demethylation of Regulatory Elements Correlates with Chromatin State and Improved beta Cell Function. *Cell metabolism* **22**, 619-632.

Benner, C., van der Meulen, T., Caceres, E., Tigyi, K., Donaldson, C. J. and Huising, M. O. (2014). The transcriptional landscape of mouse beta cells compared to human beta cells reveals notable species differences in long non-coding RNA and protein-coding gene expression. *BMC Genomics* **15**, 620.

Blandino-Rosano, M., Barbaresso, R., Jimenez-Palomares, M., Bozadjieva, N., Werneck-de-Castro, J. P., Hatanaka, M., Mirmira, R. G., Sonenberg, N., Liu, M., Ruegg, M. A., et al. (2017). Loss of mTORC1 signalling impairs beta-cell homeostasis and insulin processing. *Nat Commun* **8**, 16014.

Blodgett, D. M., Nowosielska, A., Afik, S., Pechhold, S., Cura, A. J., Kennedy, N. J., Kim, S., Kucukural, A., Davis, R. J., Kent, S. C., et al. (2015). Novel Observations From Next-Generation RNA Sequencing of Highly Purified Human Adult and Fetal Islet Cell Subsets. *Diabetes* **64**, 3172-3181.

Blum, B., Hrvatin, S., Schuetz, C., Bonal, C., Rezania, A. and Melton, D. A. (2012). Functional beta-cell maturation is marked by an increased glucose threshold and by expression of urocortin 3. *Nat Biotechnol* **30**, 261-264.

Bramswig, N. C., Everett, L. J., Schug, J., Dorrell, C., Liu, C., Luo, Y., Streeter, P. R., Naji, A., Grompe, M. and Kaestner, K. H. (2013). Epigenomic plasticity enables human pancreatic alpha to beta cell reprogramming. *J Clin Invest* **123**, 1275-1284.

Brissova, M., Fowler, M. J., Nicholson, W. E., Chu, A., Hirshberg, B., Harlan, D. M. and Powers, A. C. (2005). Assessment of human pancreatic islet architecture and composition by laser scanning confocal microscopy. *J Histochem Cytochem* **53**, 1087-1097.

Brun, P. J., Grijalva, A., Rausch, R., Watson, E., Yuen, J. J., Das, B. C., Shudo, K., Kagechika, H., Leibel, R. L. and Blaner, W. S. (2015). Retinoic acid receptor signaling is required to maintain glucose-stimulated insulin secretion and beta-cell mass. *FASEB J* **29**, 671-683.

Camunas-Soler, J., Dai, X., Hang, Y., Bautista, A., Lyon, J., Suzuki, K., Kim, S. K., Quake, S. R. and MacDonald, P. E. (2019). Pancreas patch-seq links physiologic dysfunction in diabetes to single-cell transcriptomic phenotypes. *bioRxiv*, 555110.

Capozzi, M. E., Svendsen, B., Encisco, S. E., Lewandowski, S. L., Martin, M. D., Lin, H., Jaffe, J. L., Coch, R. W., Haldeman, J. M., MacDonald, P. E., et al. (2019). beta Cell tone is defined by proglucagon peptides through cAMP signaling. *JCI Insight* **4**.

Carlsson, G. L., Heller, R. S., Serup, P. and Hyttel, P. (2010). Immunohistochemistry of pancreatic development in cattle and pig. *Anat Histol Embryol* **39**, 107-119.

Cerolsaletti, K., Hao, W. and Greenbaum, C. J. (2019). Genetics Coming of Age in Type 1 Diabetes. *Diabetes Care* **42**, 189-191.

Chakravarthy, H., Gu, X., Enge, M., Dai, X., Wang, Y., Damond, N., Downie, C., Liu, K., Wang, J., Xing, Y., et al. (2017). Converting Adult Pancreatic Islet alpha Cells into beta Cells by Targeting Both Dnmt1 and Arx. *Cell metabolism* **25**, 622-634.

Collombat, P., Mansouri, A., Hecksher-Sorensen, J., Serup, P., Krull, J., Gradwohl, G. and Gruss, P. (2003). Opposing actions of Arx and Pax4 in endocrine pancreas development. *Genes Dev* **17**, 2591-2603.

Dai, C., Hang, Y., Shostak, A., Poffenberger, G., Hart, N., Prasad, N., Phillips, N., Levy, S. E., Greiner, D. L., Shultz, L. D., et al. (2017). Age-dependent human beta cell proliferation induced by glucagon-like peptide 1 and calcineurin signaling. *J Clin Invest* **127**, 3835-3844.

DiGruccio, M. R., Mawla, A. M., Donaldson, C. J., Noguchi, G. M., Vaughan, J., Cowing-Zitron, C., van der Meulen, T. and Huising, M. O. (2016). Comprehensive alpha, beta and delta cell transcriptomes reveal that ghrelin selectively activates delta cells and promotes somatostatin release from pancreatic islets. *Mol Metab* **5**, 449-458.

Drucker, D. J. (2013). Incretin action in the pancreas: potential promise, possible perils, and pathological pitfalls. *Diabetes* **62**, 3316-3323.

Dyson, M. C., Alloosh, M., Vuchetich, J. P., Mokelke, E. A. and Sturek, M. (2006). Components of metabolic syndrome and coronary artery disease in female Ossabaw swine fed excess atherogenic diet. *Comp Med* **56**, 35-45.

Edgar, R., Domrachev, M. and Lash, A. E. (2002). Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res* **30**, 207-210.

Ferrer, J., Scott, W. E., 3rd, Weegman, B. P., Suszynski, T. M., Sutherland, D. E., Hering, B. J. and Papas, K. K. (2008). Pig pancreas anatomy: implications for pancreas procurement, preservation, and islet isolation. *Transplantation* **86**, 1503-1510.

Fisher, J. E., Lillegard, J. B., McKenzie, T. J., Rodysill, B. R., Wettstein, P. J. and Nyberg, S. L. (2013). In utero transplanted human hepatocytes allow postnatal engraftment of human hepatocytes in pigs. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society* **19**, 328-335.

Georgia, S. and Bhushan, A. (2004). Beta cell replication is the primary mechanism for maintaining postnatal beta cell mass. *J Clin Invest* **114**, 963-968.

Hart, N. J. and Powers, A. C. (2019). Use of human islets to understand islet biology and diabetes: progress, challenges and suggestions. *Diabetologia* **62**, 212-222.

Hassouna, T., Seeberger, K. L., Salama, B. and Korbutt, G. S. (2018). Functional Maturation and In Vitro Differentiation of Neonatal Porcine Islet Grafts. *Transplantation* **102**, e413-e423.

Hattersley, A. T. and Patel, K. A. (2017). Precision diabetes: learning from monogenic diabetes. *Diabetologia* **60**, 769-777.

Holst, J. J., Holland, W., Gromada, J., Lee, Y., Unger, R. H., Yan, H., Sloop, K. W., Kieffer, T. J., Damond, N. and Herrera, P. L. (2017). Insulin and Glucagon: Partners for Life. *Endocrinology* **158**, 696-701.

Hrvatin, S., Deng, F., O'Donnell, C. W., Gifford, D. K. and Melton, D. A. (2014a). MARIS:
method for analyzing RNA following intracellular sorting. *PLoS One* **9**, e89459.
Hrvatin, S., O'Donnell, C. W., Deng, F., Millman, J. R., Pagliuca, F. W., Dilorio, P., Rezania,
A., Gifford, D. K. and Melton, D. A. (2014b). Differentiated human stem cells resemble
fetal, not adult, β cells. *Proceedings of the National Academy of Sciences* **111**, 3038.
Jeon, J., Correa-Medina, M., Ricordi, C., Edlund, H. and Diez, J. A. (2009). Endocrine cell
clustering during human pancreas development. *J Histochem Cytochem* **57**, 811-824.
Jorgensen, M. C., Ahnfelt-Ronne, J., Hald, J., Madsen, O. D., Serup, P. and Hecksher-
Sorensen, J. (2007). An illustrated review of early pancreas development in the mouse.
Endocr Rev **28**, 685-705.
Kemter, E., Cohrs, C. M., Schafer, M., Schuster, M., Steinmeyer, K., Wolf-van Buerck, L.,
Wolf, A., Wuensch, A., Kurome, M., Kessler, B., et al. (2017). INS-eGFP transgenic pigs:
a novel reporter system for studying maturation, growth and vascularisation of neonatal
islet-like cell clusters. *Diabetologia* **60**, 1152-1156.
Kolde, R. (2012). Pheatmap: pretty heatmaps. *R package version* **61**, 617.
Lamb, M., Laugenour, K., Liang, O., Alexander, M., Foster, C. E. and Lakey, J. R. (2014). In
vitro maturation of viable islets from partially digested young pig pancreas. *Cell Transplant*
23, 263-272.
Lee, Y. H., Wang, M. Y., Yu, X. X. and Unger, R. H. (2016). Glucagon is the key factor in the
development of diabetes. *Diabetologia* **59**, 1372-1375.
Lemaire, K., Granvik, M., Schraenen, A., Goyvaerts, L., Van Lommel, L., Gomez-Ruiz, A.,
In 't Veld, P., Gilon, P. and Schuit, F. (2017). How stable is repression of disallowed
genes in pancreatic islets in response to metabolic stress? *PLoS One* **12**, e0181651.
Lemaire, K., Thorrez, L. and Schuit, F. (2016). Disallowed and Allowed Gene Expression:
Two Faces of Mature Islet Beta Cells. *Annu Rev Nutr* **36**, 45-71.
Lim, H., Lim, Y. M., Kim, K. H., Jeon, Y. E., Park, K., Kim, J., Hwang, H. Y., Lee, D. J.,
Pagire, H., Kwon, H. J., et al. (2018). A novel autophagy enhancer as a therapeutic agent
against metabolic syndrome and diabetes. *Nat Commun* **9**, 1438.
Lin, H. M., Lee, J. H., Yadav, H., Kamaraju, A. K., Liu, E., Zhigang, D., Vieira, A., Kim, S. J.,
Collins, H., Matschinsky, F., et al. (2009). Transforming growth factor-beta/Smad3
signaling regulates insulin gene transcription and pancreatic islet beta-cell function. *J Biol*
Chem **284**, 12246-12257.
Loh, K. M., Ang, L. T., Zhang, J., Kumar, V., Ang, J., Auyeong, J. Q., Lee, K. L., Choo, S.
H., Lim, C. Y., Nichane, M., et al. (2014). Efficient endoderm induction from human
pluripotent stem cells by logically directing signals controlling lineage bifurcations. *Cell Stem*
Cell **14**, 237-252.
Love, M. I., Huber, W. and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550.
Lukinius, A., Ericsson, J. L., Grimelius, L. and Korsgren, O. (1992). Ultrastructural studies of
the ontogeny of fetal human and porcine endocrine pancreas, with special reference to
colocalization of the four major islet hormones. *Dev Biol* **153**, 376-385.
Maestro, M. A., Cardalda, C., Boj, S. F., Luco, R. F., Servitja, J. M. and Ferrer, J. (2007).
Distinct roles of HNF1beta, HNF1alpha, and HNF4alpha in regulating pancreas
development, beta-cell function and growth. *Endocr Dev* **12**, 33-45.
Matsunari, H., Nagashima, H., Watanabe, M., Umeyama, K., Nakano, K., Nagaya, M.,
Kobayashi, T., Yamaguchi, T., Sumazaki, R., Herzenberg, L. A., et al. (2013). Blastocyst
complementation generates exogenic pancreas in vivo in apancreatic cloned pigs. *Proc Natl*
Acad Sci U S A **110**, 4557-4562.
McKnight, K. D., Wang, P. and Kim, S. K. (2010). Deconstructing pancreas development to
reconstruct human islets from pluripotent stem cells. *Cell Stem Cell* **6**, 300-308.

Meier, J. J., Butler, A. E., Saisho, Y., Monchamp, T., Galasso, R., Bhushan, A., Rizza, R. A. and Butler, P. C. (2008). Beta-cell replication is the primary mechanism subserving the postnatal expansion of beta-cell mass in humans. *Diabetes* **57**, 1584-1594.

Miguel-Escalada, I., Bonas-Guarch, S., Cebola, I., Ponsa-Cobas, J., Mendieta-Esteban, J., Atla, G., Javierre, B. M., Rolando, D. M. Y., Farabella, I., Morgan, C. C., et al. (2019). Human pancreatic islet three-dimensional chromatin architecture provides insights into the genetics of type 2 diabetes. *Nat Genet* **51**, 1137-1148.

Mueller, K. R., Balamurugan, A. N., Cline, G. W., Pongratz, R. L., Hooper, R. L., Weegman, B. P., Kitzmann, J. P., Taylor, M. J., Graham, M. L., Schuurman, H. J., et al. (2013). Differences in glucose-stimulated insulin secretion in vitro of islets from human, nonhuman primate, and porcine origin. *Xenotransplantation* **20**, 75-81.

Muraro, M. J., Dharmadhikari, G., Grun, D., Groen, N., Dielen, T., Jansen, E., van Gurp, L., Engelse, M. A., Carlotti, F., de Koning, E. J., et al. (2016). A Single-Cell Transcriptome Atlas of the Human Pancreas. *Cell Syst* **3**, 385-394 e383.

Nagaya, M., Hayashi, A., Nakano, K., Honda, M., Hasegawa, K., Okamoto, K., Itazaki, S., Matsunari, H., Watanabe, M., Umeyama, K., et al. (2019). Distributions of endocrine cell clusters during porcine pancreatic development. *PLoS One* **14**, e0216254.

Nair, G. G., Liu, J. S., Russ, H. A., Tran, S., Saxton, M. S., Chen, R., Juang, C., Li, M. L., Nguyen, V. Q., Giacometti, S., et al. (2019). Recapitulating endocrine cell clustering in culture promotes maturation of human stem-cell-derived beta cells. *Nat Cell Biol* **21**, 263-274.

Pagliuca, F. W., Millman, J. R., Gurtler, M., Segel, M., Van Dervort, A., Ryu, J. H., Peterson, Q. P., Greiner, D. and Melton, D. A. (2014). Generation of functional human pancreatic beta cells in vitro. *Cell* **159**, 428-439.

Pan, F. C. and Brissova, M. (2014). Pancreas development in humans. *Current opinion in endocrinology, diabetes, and obesity* **21**, 77-82.

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. and Kingsford, C. (2017). Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods* **14**, 417-419.

Pearson, E. R. (2019). Type 2 diabetes: a multifaceted disease. *Diabetologia* **62**, 1107-1112.

Peiris, H., Park, S., Louis, S., Gu, X., Lam, J. Y., Asplund, O., Ippolito, G. C., Bottino, R., Groop, L., Tucker, H., et al. (2018). Discovering human diabetes-risk gene function with genetics and physiological assays. *Nat Commun* **9**, 3855.

Perez-Alcantara, M., Honore, C., Wesolowska-Andersen, A., Gloyn, A. L., McCarthy, M. I., Hansson, M., Beer, N. L. and van de Bunt, M. (2018). Patterns of differential gene expression in a cellular model of human islet development, and relationship to type 2 diabetes predisposition. *Diabetologia* **61**, 1614-1622.

Pullen, T. J., Huising, M. O. and Rutter, G. A. (2017). Analysis of Purified Pancreatic Islet Beta and Alpha Cell Transcriptomes Reveals 11beta-Hydroxysteroid Dehydrogenase (Hsd11b1) as a Novel Disallowed Gene. *Front Genet* **8**, 41.

Pullen, T. J., Khan, A. M., Barton, G., Butcher, S. A., Sun, G. and Rutter, G. A. (2010). Identification of genes selectively disallowed in the pancreatic islet. *Islets* **2**, 89-95.

Qiu, W.-L., Zhang, Y.-W., Feng, Y., Li, L.-C., Yang, L. and Xu, C.-R. (2017). Deciphering Pancreatic Islet β Cell and α Cell Maturation Pathways and Characteristic Features at the Single-Cell Level. *Cell metabolism* **25**, 1194-1205.e1194.

Ramond, C., Beydag-Tasoz, B. S., Azad, A., van de Bunt, M., Petersen, M. B. K., Beer, N. L., Glaser, N., Berthault, C., Gloyn, A. L., Hansson, M., et al. (2018). Understanding human fetal pancreas development using subpopulation sorting, RNA sequencing and single-cell profiling. *Development* **145**.

Renner, S., Braun-Reichhart, C., Blutke, A., Herbach, N., Emrich, D., Streckel, E., Wunsch, A., Kessler, B., Kurome, M., Bahr, A., et al. (2013). Permanent neonatal diabetes in INS(C94Y) transgenic pigs. *Diabetes* **62**, 1505-1511.

Rezania, A., Bruin, J. E., Arora, P., Rubin, A., Batushansky, I., Asadi, A., O'Dwyer, S.,
Quiskamp, N., Mojibian, M., Albrecht, T., et al. (2014). Reversal of diabetes with insulin-
producing cells derived in vitro from human pluripotent stem cells. *Nat Biotechnol* **32**, 1121-
1133.
Riopel, M., Li, J., Fellows, G. F., Goodyer, C. G. and Wang, R. (2014). Ultrastructural and
immunohistochemical analysis of the 8-20 week human fetal pancreas. *Islets* **6**, e982949.
Rorsman, P., Berggren, P. O., Bokvist, K., Ericson, H., Mohler, H., Ostenson, C. G. and
Smith, P. A. (1989). Glucose-inhibition of glucagon secretion involves activation of GABAA-
receptor chloride channels. *Nature* **341**, 233-236.
Rorsman, P. and Huising, M. O. (2018). The somatostatin-secreting pancreatic delta-cell in
health and disease. *Nat Rev Endocrinol* **14**, 404-414.
Rouille, Y., Martin, S. and Steiner, D. F. (1995). Differential processing of proglucagon by the
subtilisin-like prohormone convertases PC2 and PC3 to generate either glucagon or
glucagon-like peptide. *J Biol Chem* **270**, 26488-26496.
Russ, H. A., Parent, A. V., Ringler, J. J., Hennings, T. G., Nair, G. G., Shveygert, M., Guo,
T., Puri, S., Haataja, L., Cirulli, V., et al. (2015). Controlled induction of human pancreatic
progenitors produces functional beta-like cells in vitro. *The EMBO journal* **34**, 1759-1772.
Sanyoura, M., Philipson, L. H. and Naylor, R. (2018). Monogenic Diabetes in Children and
Adolescents: Recognition and Treatment Options. *Curr Diab Rep* **18**, 58.
Schaffer, A. E., Taylor, B. L., Benthuysen, J. R., Liu, J., Thorel, F., Yuan, W., Jiao, Y.,
Kaestner, K. H., Herrera, P. L., Magnuson, M. A., et al. (2013). Nkx6.1 controls a gene
regulatory network required for establishing and maintaining pancreatic Beta cell identity.
PLoS Genet **9**, e1003274.
Sekine, N., Cirulli, V., Regazzi, R., Brown, L. J., Gine, E., Tamarit-Rodriguez, J., Girotti, M.,
Marie, S., MacDonald, M. J., Wollheim, C. B., et al. (1994). Low lactate dehydrogenase
and high mitochondrial glycerol phosphate dehydrogenase in pancreatic beta-cells.
Potential role in nutrient sensing. *J Biol Chem* **269**, 4895-4902.
Sheets, T. P., Park, K. E., Park, C. H., Swift, S. M., Powell, A., Donovan, D. M. and Telugu,
B. P. (2018). Targeted Mutation of NGN3 Gene Disrupts Pancreatic Endocrine Cell
Development in Pigs. *Sci Rep* **8**, 3582.
Sneddon, J. B., Tang, Q., Stock, P., Bluestone, J. A., Roy, S., Desai, T. and Hebrok, M.
(2018). Stem Cell Therapies for Treating Diabetes: Progress and Remaining Challenges.
Cell Stem Cell **22**, 810-823.
Sodoyez-Goffaux, F., Sodoyez, J. C., De Vos, C. J. and Foa, P. P. (1979). Insulin and
glucagon secretion by islets isolated from fetal and neonatal rats. *Diabetologia* **16**, 121-123.
Soneson, C., Love, M. I. and Robinson, M. D. (2015). Differential analyses for RNA-seq:
transcript-level estimates improve gene-level inferences. *F1000Res* **4**, 1521.
Steiner, D. F. (1998). The proprotein convertases. *Curr Opin Chem Biol* **2**, 31-39.
Sugiyama, T., Benitez, C. M., Ghodasara, A., Liu, L., McLean, G. W., Lee, J., Blauwkamp,
T. A., Nusse, R., Wright, C. V., Gu, G., et al. (2013). Reconstituting pancreas development
from purified progenitor cells reveals genes essential for islet differentiation. *Proc Natl Acad*
Sci U S A **110**, 12691-12696.
Svendsen, B., Larsen, O., Gabe, M. B. N., Christiansen, C. B., Rosenkilde, M. M., Drucker,
D. J. and Holst, J. J. (2018). Insulin Secretion Depends on Intra-islet Glucagon Signaling.
Cell Rep **25**, 1127-1134 e1122.
Tengholm, A. and Gylfe, E. (2017). cAMP signalling in insulin and glucagon secretion.
Diabetes Obes Metab **19 Suppl 1**, 42-53.
Teta, M., Rankin, M. M., Long, S. Y., Stein, G. M. and Kushner, J. A. (2007). Growth and
regeneration of adult beta cells does not involve specialized progenitors. *Dev Cell* **12**, 817-
826.

Thorrez, L., Laudadio, I., Van Deun, K., Quintens, R., Hendrickx, N., Granvik, M., Lemaire, K., Schraenen, A., Van Lommel, L., Lehnert, S., et al. (2011). Tissue-specific disallowance of housekeeping genes: the other face of cell differentiation. *Genome Res* **21**, 95-105.

Umeyama, K., Watanabe, M., Saito, H., Kurome, M., Tohi, S., Matsunari, H., Miki, K. and Nagashima, H. (2009). Dominant-negative mutant hepatocyte nuclear factor 1alpha induces diabetes in transgenic-cloned pigs. *Transgenic Res* **18**, 697-706.

van der Meulen, T., Donaldson, C. J., Caceres, E., Hunter, A. E., Cowing-Zitron, C., Pound, L. D., Adams, M. W., Zembrzycki, A., Grove, K. L. and Huising, M. O. (2015). Urocortin3 mediates somatostatin-dependent negative feedback control of insulin secretion. *Nat Med* **21**, 769-776.

Velayos, T., Martinez, R., Alonso, M., Garcia-Etxebarria, K., Aguayo, A., Camarero, C., Urrutia, I., Martinez de LaPiscina, I., Barrio, R., Santin, I., et al. (2017). An Activating Mutation in STAT3 Results in Neonatal Diabetes Through Reduced Insulin Synthesis. *Diabetes* **66**, 1022-1029.

Wende, A. R., Huss, J. M., Schaeffer, P. J., Giguere, V. and Kelly, D. P. (2005). PGC-1alpha coactivates PDK4 gene expression via the orphan nuclear receptor ERRalpha: a mechanism for transcriptional control of muscle glucose metabolism. *Mol Cell Biol* **25**, 10684-10694.

Wickham, H. (2009). ggplot2: Elegant Graphics for Data Analysis. *Use R*, 1-212.

Wilson, M. E., Kalamaras, J. A. and German, M. S. (2002). Expression pattern of IAPP and prohormone convertase 1/3 reveals a distinctive set of endocrine cells in the embryonic pancreas. *Mech Dev* **115**, 171-176.

Wu, J., Platero-Luengo, A., Sakurai, M., Sugawara, A., Gil, M. A., Yamauchi, T., Suzuki, K., Bogliotti, Y. S., Cuello, C., Morales Valencia, M., et al. (2017). Interspecies Chimerism with Mammalian Pluripotent Stem Cells. *Cell* **168**, 473-486 e415.

Yang, Y. and Chan, L. (2016). Monogenic Diabetes: What It Teaches Us on the Common Forms of Type 1 and Type 2 Diabetes. *Endocr Rev* **37**, 190-222.

Yokoi, N., Gheni, G., Takahashi, H. and Seino, S. (2016). beta-Cell glutamate signaling: Its role in incretin-induced insulin secretion. *J Diabetes Investig* **7 Suppl 1**, 38-43.

Yu, G., Wang, L. G., Han, Y. and He, Q. Y. (2012). clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* **16**, 284-287.

Zerbino, D. R., Achuthan, P., Akanni, W., Amode, M. R., Barrell, D., Bhai, J., Billis, K., Cummins, C., Gall, A., Giron, C. G., et al. (2018). Ensembl 2018. *Nucleic Acids Res* **46**, D754-D761.

Zhu, L., Dattaroy, D., Pham, J., Wang, L., Barella, L. F., Cui, Y., Wilkins, K. J., Roth, B. L., Hochgeschwender, U., Matschinsky, F. M., et al. (2019). Intra-islet glucagon signaling is critical for maintaining glucose homeostasis. *JCI Insight* **5**.

 Figure legends

Figure 1. Pancreas dissociation and FACS purification of islet cells from fetal and neonatal pigs.

(A) Schematic representation of the study design. (B) qRT-PCR analysis of *INS*, *GCG*, *SST*,

KRT, and *AMY2* of FACS sorted β-, α- and δ-cell populations from P22 neonatal pigs. The fold

change (2^-ΔΔCT) relative to the unsorted cell population is shown on the y-axis. (C) Heatmap

of differentially expressed genes (DEGs) in pig β-cells between early fetal, late fetal and

neonatal stages. (D) GO Term enrichment analysis of genes increasing with age in pig β-cells.

(E) Heatmap of DEGs in pig α-cells between early fetal, late fetal and neonatal stages. (F) GO

Term enrichment analysis of genes increasing with age in pig α-cells. (G) Heatmap of DEGs in

pig δ-cells between late fetal and neonatal stages. (H) GO Term enrichment analysis of genes

increasing with age in pig δ-cells. The Z-score scale represents log_2(X+1) transformed

transcripts per million (TPM) counts. Red and blue color intensity of the Z-score indicates up-

regulation and down-regulation, respectively. Adjusted *P* value threshold for GO term analysis

was 0.1.

1009

Figure 2. Analysis of gene expression in developing pig islet β-, α- and δ-cells**.** (A-E,G,H)

Boxplots displaying normalized TPM (transcripts per million) counts of β-, α-, and δ-cell specific

genes (A-D), markers associated with proliferation (E), disallowed genes (G) and β-cell

functional regulators (G). (F) Quantification showing average percentage of INS$^+$, GCG$^+$ or SST$^+$

cells that are Ki67$^+$ in each developmental stage (n = 40 images per group, from 3 pigs per

group; bars show SD). Adjusted *P* value * *P≤0.05; ** P≤0.01; *** P≤0.001*

1016

Figure 3. Stage-specific expression of islet factors during development. (A-C) Immunostaining

of INS (green) and GCG (white) paired with Somatostatin (SST; red) (A), E-Cadherin (ECad;

white) (B), and Chromogranin-A (CHGA; white) (C) in the early fetal, neonatal, and adult pig

pancreas. (D-F) Immunostaining of the transcription factors PDX1 (D), NKX6.1 (E), and MAFB

(F) in the early fetal, neonatal, and adult pig pancreas. (**G**) Immunostaining of SIX3 with INS and

GCG in adult pig pancreas. Insets indicated by the dashed yellow box. Scale bars, 50 μm. (**H, I**)

qRT-PCR measures of relative mRNA levels encoding *SIX3* **(H)** and *SIX2* **(I)** in isolated

neonatal and adult pig islets (n=2, t-test; Error bars indicate SD). ΔCT values for neonatal and

adult *SIX3* expression were 5.1 and 2.0, respectively (compared to β–Actin). ΔCT values for

neonatal and adult *SIX2* (compared to β–Actin) were 7.2 and 2.6, respectively.

1027

1028   **Figure 4.** Comparison of gene expression in pig and human β- and α-cell development. (A)

1029   Unsupervised hierarchical clustering using distance matrices of neonatal pig, human, and

1030   mouse β-cells. (B-G) Boxplots displaying normalized TPM (transcripts per million) counts of

1031   select genes whose expression changes in a conserved manner between human and pig

1032   development. Adjusted *P* value * *P≤0.05; ** P≤0.01; *** P≤0.001.*

1033

Figure 5. Dynamic gene regulation during pig β- and α-cell development. (A-B) Heat maps of

genes with altered expression from early fetal to neonatal stages in β-cells (A) and α-cells (B),

showing 8 distinct clusters. Cluster 2 includes genes whose mRNA increased up to late fetal

stages then did not change thereafter. Cluster 7 includes genes whose mRNA declined then did

not change thereafter. Enriched GO terms in each cluster are shown to the right. Adjusted *P*

value threshold was 0.1. The Z-score scale represents $\log_2(X+1)$ transformed transcripts per

million (TPM) counts. Red and blue color intensity of the Z-score indicates up-regulation and

down-regulation, respectively. (C-D) The Z-score scale represents $\log_2(X+1)$ transformed TPM

counts with total genes in β-cells (C) and α-cells (D). Cluster 3 (Up-Down) or Cluster 6

(Down_Up). Boxplots displaying normalized TPM (transcripts per million) counts of genes

shown. (E) Boxplots displaying normalized TPM (transcripts per million) counts of PCSK1

mRNA. (F) Immunostaining of PCSK1 (white) with insulin (INS; green) and glucagon (GCG; red)

in early fetal, late fetal and adult pig pancreas. Insets indicated by the dashed yellow box. White

arrows indicate PCSK1[+] GCG[+] (double positive) cells. (G) Concentrations of active GLP-1 in

lysates from pig islet cell clusters at different developmental stages (ANOVA, ** *P≤0.01*; bars

indicate SD). (H) Boxplots displaying normalized TPM (transcripts per million) counts of *DPP4*

mRNA. (I) Immunostaining of DPP4 (white) with insulin (INS; green) and glucagon (GCG; red) in

early fetal, neonatal and adult pig pancreas. Insets indicated by the dashed yellow box. White

arrows indicate DPP4[+] INS[+] (double positive) cells. Adjusted *P* value * *P≤0.05; ** P≤0.01; ***

P≤0.001*, Scale bars, 50 µm.

1054

Figure 6. Maturation of β-cell function in postnatal pigs. (A) Glucose-stimulated insulin secretion

(GSIS) assay measured insulin secretion in response to 2.8mM (low), 20mM (high) glucose and

20mM glucose + IBMX. Data show average secreted insulin as a percentage of insulin content

in the isolated islets from late fetal (n=4) and P22 (n=3) piglets (t-test, ** *P≤0.01; *** P≤0.001*;

bars indicate SD). (B) Heat map of DEGs between late fetal and P22 β-cells. The Z-score scale

represents $\log_2(X+1)$ transformed transcripts per million (TPM) counts. Red and blue color

intensity of the Z-score indicates up-regulation and down-regulation, respectively. (C,E) GO

term analysis for biological processes with up-regulated (C) and down-regulated (E) genes

between late fetal and P22 β-cells. Adjusted *P* value threshold was 0.1. (D,F) Boxplots

displaying normalized TPM (transcripts per million) counts of up-regulated (D) or down-

regulated (F) genes between late fetal and P22. Adjusted *P* value * *P≤0.05; ** P≤0.01; ***

P≤0.001.

Fig. 1

	Fetal	Neonatal	Adult

A INS GCG SST

B INS GCG ECad

C INS GCG CHGA

D INS GCG PDX1

E INS GCG NKX6.1

F INS GCG MAFB

G Adult

INS GCG SIX3

H SIX3 expression

Fold change (relative to Neonatal)

Neonatal Adult

I SIX2 expression

Fold change (relative to Neonatal)

Neonatal Adult

Fig. 3

Fig. 4

Fig. 5

Fig. 6