Phenotypic and functional characterization of corneal endothelial cells during in vitro expansion

Ricardo F. Frausto[1], Vinay S. Swamy[1], Gary S. L. Peh[2], Payton M. Boere[1], E. Maryam Hanser[1], Doug. D.

Chung[1], Marco Morselli[3,4], Liyo Kao[5], Rustam Azimov[5], Matteo Pellegrini[3,4,6,7], Ira Kurtz[5,8], Jodhbir S.

Mehta[2] and Anthony J. Aldave[1]

[1]Stein Eye Institute, David Geffen School of Medicine at UCLA, Los Angeles, California, USA; [2]Tissue

Engineering and Stem Cell Group, Singapore Eye Research Institute, Singapore; [3]Department of

Molecular, Cell and Developmental Biology, UCLA, Los Angeles, California 90095, USA; [4]Institute for

Quantitative and Computational Biology, UCLA, Los Angeles, California 90095, USA; [5]Division of

Nephrology, David Geffen School of Medicine at UCLA, Los Angeles, California 90095; [6]Molecular

Biology Institute, UCLA, Los Angeles, California 90095, USA; [7]Jonsson Comprehensive Cancer Center,

UCLA, Los Angeles, California 90095, USA; [7]Brain Research Institute, UCLA, Los Angeles, California

90095

Support provided by National Eye Institute Grants R01 EY022082 (A.J.A.), P30 EY000331 (core grant),

the Walton Li Chair in Cornea and Uveitis (A.J.A.), Eye Bank Association of America Richard Lindstrom

Research Grant (AJA), the Stotter Revocable Trust (SEI Cornea Division), an unrestricted grant from

Research to Prevent Blindness (A.J.A.), National Institute of Diabetes and Digestive and Kidney Diseases

1R01 DK077162 (I.K.), the Allan Smidt Charitable Fund (I.K.), Ralph Block Family Foundation (I.K.),

and U.S. Department of Energy Office of Science, Office of Biological and Environmental Research

Program DE-FC02-02ER63421 (M.M).

Correspondence:

Anthony J. Aldave, M.D.

Professor of Ophthalmology

Stein Eye Institute, 200 Stein Plaza, UCLA, Los Angeles, CA 90095-7003

Tele: 310.206.7202   Fax: 310.794.7906   Email: aldave@jsei.ucla.edu

27   **SUMMARY**

The advent of cell culture-based methods for the establishment and expansion of human corneal

endothelial cells (CEnC) has provided an available source of transplantable corneal endothelium, with a

significant potential to challenge the one donor-one recipient paradigm. However, concerns over cell state

identity remain, and a comprehensive characterization of the cultured CEnC has not been performed. To

this end, we compared two established CEnC culture methods by assessing cell function and

characterizing the transcriptomic changes that occur during in vitro expansion. In confluent monolayers,

low mitogenic culture conditions preserved corneal endothelial cell state identity better than culture in

high mitogenic conditions. Expansion by continuous passaging induced replicative cell senescence.

Transcriptomic analysis of the senescent phenotype identified a cell senescence signature distinct for

CEnC. We identified activation of both classic and new cell signaling pathways that may be targeted to

prevent senescence, a significant barrier to realizing the potential clinical utility of in vitro expansion.

Keywords: CEnC, cornea, endothelium, endothelial cells, senescence, transcriptomics, transplantation,

cell barrier, cell migration, pump function

INTRODUCTION

The worldwide shortage of donor corneal tissue for the treatment of corneal endothelial dysfunction

necessitates the development of viable alternatives to the paradigm of one donor cornea being used for

only one recipient (Mehta et al., 2019; Okumura et al., 2014b; Soh et al., 2017). Corneal endothelial cell

(CEnC) dysfunction is the primary indication for corneal transplantation both in the United States and

worldwide, and while endothelial keratoplasty represents a significant advance in the surgical

management of corneal endothelial dysfunction, the worldwide shortage of surgical-grade donor corneas

and the lack of adequately trained surgeons in the majority of countries, as well as a variety of associated

intraoperative and postoperative complications, have significantly limited the impact of endothelial

keratoplasty on visual impairment worldwide due to corneal endothelial dysfunction (Deng et al., 2015;

Lass et al., 2017; Van den Bogerd et al., 2018). The in vitro generation of stem cell-derived corneal

endothelial-like cells (Yamashita et al., 2018), immortalized CEnC lines (Schmedt et al., 2012; Valtink et

al., 2008) and expansion of primary CEnC from cadaveric donor corneal tissue (Okumura et al., 2014b;

Parekh et al., 2016; Peh et al., 2019) have challenged the one donor-one recipient paradigm of corneal

transplantation. Nevertheless, in vitro culture poses its own challenges, including unwanted changes in

cell phenotype (e.g., endothelial to fibroblastic) and progression towards replicative senescence that limits

cell numbers (Sheerin et al., 2012; Soh et al., 2017). In addition, the quality of the donor tissue from

which the CEnC are derived is critical in the successful establishment of an in vitro CEnC culture. Donor

age significantly impacts culture success rate, with the optimal age being less than 40 years old. Reduced

success rates from older donors are correlated with an appearance of senescence-associated markers ex

vivo (Mimura and Joyce, 2006), and this is believed to significantly limit re-entry into the cell cycle even

in the presence of potent mitogenic factors. While senescence is generally an irreversible cell state

(Campisi, 2013), CEnC from younger donors are characterized by a quiescent cell state (G1 cell cycle

arrest), which is a reversible mitotic arrest that enables these cells to undergo mitogen-induced cell cycle

re-entry (Joyce, 2005). Donor age and other factors (e.g., days in preservation medium, donor medical

history, cell count) dictate the success of establishing in vitro cultures (Soh et al., 2017). This makes

identifying an optimal in vitro culture protocol essential for ensuring consistent establishment and

expansion of CEnC .

The corneal endothelium is a neuroectoderm-derived tissue that is located on the posterior surface

of the cornea and is a semipermeable monolayer of mitotically inactive (i.e., quiescent) CEnC. A critical

functional property of the corneal endothelium is to maintain the corneal stroma in a relatively dehydrated

state. This process ensures that the collagen fibers of the stroma retain an ultrastructural organization

essential for corneal transparency. The pump-leak hypothesis is believed to best explain the role that the

endothelium plays in maintaining a relatively dehydrated stroma. Passive movement of water from the

aqueous humor to the stroma (i.e., leak) and active transport of solutes (i.e., pump) in the opposite

direction are regulated by the CEnC barrier formation (i.e., tight junctions, cell adhesion) and expression

of solute transporters (e.g., Na/K ATPases, SLC4A11) (Gottsch et al., 2003). Mutations in genes

associated with transporter function (e.g., *SLC4A11*) or CEnC identity (e.g., *ZEB1, TCF4*) lead to stromal

edema and loss of corneal clarity (Baratz et al., 2010; Krafchak et al., 2005; Vithana et al., 2006; Wieben

et al., 2012). In general, loss of barrier integrity, dysfunction of solute transporters or a significant

decrease in CEnC density leads to loss of corneal clarity that necessitates corneal transplantation.

Endothelial cell failure constitutes the primary indication for corneal transplantation in the U.S., serving

at the indication for 55% of all keratoplasty procedures performed in 2018 (2018). While endothelial

keratoplasty remains the primary method for managing endothelial cell dysfunction in the U.S., the

aforementioned factors that have limited the impact of endothelial keratoplasty on decreasing the global

burden of vision loss from endothelial dysfunction necessitates the development of alternative therapeutic

interventions.

To achieve this goal, we assessed two previously reported methods for establishing cultures of

primary CEnC (Bednarz et al., 1998; Peh et al., 2015) by using a multipronged approach. We determined

the impact of in vitro expansion on CEnC gene expression by performing a transcriptomics analysis, and

identified gene expression features of replicative senescence. In addition, we performed a variety of

assays to determine the impact of these two methods on essential CEnC functions. We identified new

potential targets for suppressing cellular senescence, and confirmed that a relatively low mitogenic

environment is better at maintaining the CEnC phenotype in vitro (Peh et al., 2015). These findings form

the basis for continued development of in vitro culture and expansion of primary CEnC for their eventual

use in cell replacement therapy for the management of corneal endothelial loss or dysfunction.

METHODS

Primary corneal endothelial cell cultures

Corneas used in this study were obtained from commercial eye banks (Table S1). Criteria used for

selection of high quality donor corneal tissue were: 1) donor younger than 40 years (mean: 17.6; range: 2-

35); 2) no donor history of diabetes or corneal disease; 3) endothelial cell density greater than 2300

cells/mm^2 (mean: 3019; range: 2387-3436); 4) death to preservation less than 12 hours, if body not

cooled, or less than 24 hours, if body cooled; and 4) death to culture less than 15 days (mean: 6; range: 2-

14). Descemet membrane with attached endothelium was stripped from the stroma using the method

commonly employed in preparation of the donor cornea for Descemet membrane endothelial keratoplasty.

Seven independent CEnC cultures were established using two previously described protocols with minor

modifications (Bednarz et al., 1998; Peh et al., 2015). One method utilizes trypsin for dissociation of

endothelial cells from Descemet membrane, followed by seeding on laminin coated cell culture plastic,

and culturing in a 1:1 mixture of F12-Ham's and M199 (F99) medium. The second method utilizes

collagenase A for dissociation of endothelial cells from Descemet membrane. This is followed by seeding

on collagen IV coated cell culture plastic, and culturing initially in Endothelial SFM (M5) followed by

culturing in a 1:1 mixture of F12-Ham's and M199 (M4) medium. When cells reach confluence, the

medium is changed back to M5 medium for establishment and maintenance of the CEnC phenotype. Cell

passaging is performed with TrypLE Select (Thermo Fisher Scientific). Cells isolated using each protocol

are referred to as F99 cells or M5 (M4/M5 or dual media) cells to indicate the method used to establish

the cultures.

Total RNA isolation

Primary CEnC were lysed in TRI Reagent (Thermo Fisher) and total RNA was prepared as per the

manufacturer's instructions. RNA preparations were subsequently purified using the RNeasy Clean-Up

Kit (Qiagen, Valencia, CA). The quality of the total RNA was assessed with both the Agilent 2100

Electrophoresis Bioanalyzer System (Agilent Technologies, Inc., Santa Clara, CA) and the Agilent

TapeStation 2200 (Agilent Technologies, Inc.).

RNA-sequencing and data processing

RNA was isolated and RNA-seq libraries were prepared using the KAPA mRNA HyperPrep Kit with an

automated liquid handler (Janus G3 – PerkinElmer) according to the manufacturer's instructions. Library

preparation was performed at the UCLA Institute for Quantitative and Computational Biology. DNA

libraries were submitted to the UCLA Technology Center for Genomics and Bioinformatics for

sequencing, which was performed on the Illumina HiSeq 3000 platform. All RNA-seq data were single-

end 50 base reads. Reads were aligned to the human GRCh38.p12 genome, and transcripts were

quantified using the kallisto (v0.44.0) program (Bray et al., 2016) with the Ensembl Annotation Release

version 92. Quantities were given in transcripts per million (TPM), and differential gene expression

analysis was performed with the Sleuth (v0.30.0) R-package (Pimentel et al., 2017). Differential

expression was tested using a likelihood ratio test (negative binomial test), and corrected for multiple

testing using the Benjamini-Hochberg correction. Given the sporadic availability of donor corneas, each

of the seven cultures was established as an individual batch. To account for batch effects in the data, we

included batch number (i.e., culture number) as a covariate in the model used to test for differential

expression. The following thresholds defined differential expression: fold change>1.5, TPM>11.25 and q-

value<0.05. Of note, our background threshold was 7.5 TPM, and the threshold of 11.25 TPM (stated

above) accounted for the 1.5 fold change criterion. With the goal of achieving a balance between Type I

and Type II errors, the background threshold was selected to retain genes with low expression (e.g., *ZEB1*

with a TPM of ~25 TPM), but was sufficiently robust to exclude many genes with low (<7.5 TPM)

abundance values. RNA-seq data generated from ex vivo corneal cell types were obtained from the GEO

DataSets database (accession GSE121922). RNA-seq data generated for this study were submitted to

GEO DataSets and assigned accession number GSE132204.

Gene ontology and pathway analysis

Gene ontology (GO) and pathway analysis on gene symbols only was performed using the web-based tool

gProfiler (version: e94_eg41_p11_04285a3). The three main GO categories are Molecular Function

(MF), Biological Process (BP) and Cellular Component (CC) (data version: release/2018-12-28).

Databases used for pathway analysis were Kyoto Encyclopedia of Genes and Genomes (KEGG)

pathways (version: KEGG FTP Release 2019-01-07), REACTOME (REAC) pathways (classes: 2019-1-

24), and WikiPathways (WP)(version: 20190110). The g:SCS method for computing multiple testing

correction was used for selecting significantly enriched GO terms and pathways identified by gProfiler.

Ingenuity Pathway Analysis software (Qiagen; build: 486617M) was used to determine the pathways

enriched and activated in our lists of differentially expressed genes.

Quantitative PCR

Quantitative polymerase chain reaction (qPCR) was performed to validate gene transcript abundances

observed by RNA-seq. Total RNA (100 ng) was subjected to first-strand cDNA synthesis using the

SuperScript III First-Strand Synthesis kit (Thermo Fisher Scientific) with oligo-dT primers. Reactions

were performed on the LightCycler 480 System (Roche) using the KAPA SYBR FAST qPCR Kit

(Roche) and gene-specific oligonucleotide primers obtained from the Harvard Primer Bank database

(Spandidos et al., 2008; Spandidos et al., 2010; Wang and Seed, 2003) or designed using the NCBI

Primer-BLAST tool (Table S2). Relative transcript abundances were calculated in comparison with the

housekeeping gene *RAB7* using the comparative Ct ($2^{-\Delta Ct}$) method (Livak and Schmittgen, 2001).

Transcript quantities were graphed as $2^{-\Delta Ct}$.

Capillary-based, automated protein immunodetection assay

Cells were lysed with RIPA buffer and whole-cell lysates were prepared and processed for protein

detection using the Wes separation 12-230 kDa capillary cartridges (Protein Simple). Separation and

detection were performed as per the manufacturer's instructions. Quantification and data analysis were

performed using the Compass for SW software (version 3.1.7; build ID: 1205). Primary antibodies are

listed in Table S3).

Morphometric analysis

At each passage, we acquired phase contrast images of confluent monolayers established in either F99 or

M5. Image acquisition was performed with the Leica DMIL LED inverted microscope (Leica

Microsystems), the N PLAN L 20x/0.35 PH1 objective, and the SPOT Insight color camera

(Diagnostic Instruments, Inc.). Images were captured with the SPOT software (version 4.6). Image

analysis was performed using ImageJ 1.51h (National Institutes of Health). We created regions of

interest (ROI) by outlining the periphery of at least 20 representative cells using the freehand tool.

ImageJ output the values for area (um2) and circularity (ratio) for the selected ROI.

Cell barrier assay

The electrode array 8W10E+ ECIS (Applied BioPhysics) was stabilized with F99 or M5 medium. The

array surface was coated with 40 µg/cm^2 chondroitin sulfate A (Sigma-Aldrich) and 400 ng/cm^2 laminin

(Sigma-Aldrich) in phosphate-buffered saline (PBS) for two hours (for F99 cells) or with 7.5 µg/cm^2

collagen IV (Sigma-Aldrich) in 1x HEPES for 30 min (for M5 cells). Cells were seeded at a density to

achieve 100% confluence shortly after seeding, and were incubated in the arrays at room temperature for

one hour to facilitate even distribution. After seeding and preliminary cell attachment, arrays were

positioned into a 16-well array station and connected to the ECIS Zθ instrument to measure electric

impedance (Ω at 4000 Hz) for 4 days. Cell-cell (R_b, $\Omega \cdot cm^2$) and cell-substrate (α, $\Omega^{1/2} \cdot cm$) adhesion

were modeled from the electric impedance data obtained at 4000 Hz (Stolwijk et al., 2015).

Transporter assays

Lactate (SLC16A1), bicarbonate (SLC4A4) and proton (SLC4A11) transport were monitored by

measuring free H^+ concentration (pHi) with a microscope fluorometer (Kao et al., 2016) using the

fluorescence-based (dual-excitation 500 nm and 440 nm) ratiometric pH indicator BCECF (Thermo

Fisher Scientific), which was pre-loaded into the cells prior to substrate exposure.

SLC16A1 transporter: BCECF loading was performed in lactate-free solution (20mM Na gluconate,

120mM NaCl, 1mM $CaCl_2$, 1mM $MgCl_2$, 2.5mM K_2HPO_4, 5mM dextrose and 5mM HEPES, pH 7.4).

Fluorescence was monitored until a stable pH_i was achieved. Subsequently, the lactate-free buffer was

replaced by perfusion with lactate-containing solution (20mM Na lactate in place of 20mM Na

gluoconate) for about 200 seconds and then switched back to the lactate-free solution.

SLC4A4 transporter: BCEFC loading was performed in bicarbonate-free solution (140mM TMACl,

2.5mM K2HPO4, 1mM CaCl2, 1mM MgCl2, 5mM Dextrose and 5mM HEPES, pH 7.4). After pH_i

stabilization was attained, the solution was replaced by perfusion with a solution containing

tetramethylammonium bicarbonate (115mM TMACl and 25mM TMAHCO3 in place of 140mM TMACl

and 5mM HEPES). Perfusion was performed for about 200 seconds and then switched to a solution

containing sodium bicarbonate (115mM NaCl, 2.5mM K2HPO4, 1mM CaCl2, 1mM MgCl2, 5mM

Dextrose and 25mM NaHCO3, pH 7.4), and fluorescence was monitored for an additional 400-600

seconds.

SLC4A11 transporter: BCECF loading was performed in HEPES buffer (110mM KCl, 35mM TMACl

and 5mM HEPES, pH 7.4). Once pH_i stabilized, a second HEPES buffer (110 mM KCl, 35 mM TMACl

and 5 mM HEPES pH 6.2) was replaced by perfusion and pH_i was monitored for approximately 800

seconds.

Cell migration assay

A non-wounding method to assess cell migration was used. Two-well silicone inserts (ibidi GmbH), each

creating a 500um gap, were placed onto cell culture treated plastic. Wells were coated as described above

for F99 and M5 cells. After cell isolation was performed, cells were seeded into each well and allowed to

grow to confluence. Cell migration was initiated by removing the insert and was monitored for up to 60

hours by phase-contrast microscopy. Images were acquired using the Leica DMIL LED inverted

microscope (Leica Microsystems) and the N PLAN L 20x/0.35 PH1 objective. Image capture was

performed with the Leica DFC3000 G monochrome camera controlled with the Leica Application

Suite X software (version 3.0.3.16319). Gap closure was measured at 22 hours and cell size was

measured at 60 hours using ImageJ 1.51h software (National Institutes of Health).

230   **RESULTS**

In vitro expansion of CEnC induces senescence-associated morphogenesis

The morphogenic effects of culture in high mitogenic (F99) and low mitogenic (M5) conditions on

primary CEnC were examined (Fig. 1). Phase contrast images were acquired at each passage when

confluent monolayers were established (Fig. 1B and Fig. S1). Morphometric analysis was performed at

each passage (Fig. 1C). Up to passage 3, the area occupied by each cell was greater in F99, compared

with cells in M5, but the effect of medium on the curves was not statistically significant (p=0.065). Cell

circularity, which measures the degree to which a cell shape resembles a circle (1.0 is a perfect circle),

was greater at all passages for cells in M5 medium, compared with cells in F99. The effect of medium on

the curves for circularity was statistically significant (p=0.042). As the value approaches 0, cell shape is

increasingly irregular and/or elongated.

Figure 1. M5 medium delays morphologic features associated with a senescent phenotype. (A) Two protocols for the isolation and culture of primary CEnC. After detachment from the cornea, Descemet membrane (DM) with the attached endothelium was either incubated overnight in F99 medium at 37C (left panel) or subjected directly to collagenase digestion (middle panel). Cells incubated overnight were detached from DM with trypsin and seeded onto laminin-coated plastic. Cells dissociated from DM using collagenase were seeded onto collagen-coated plastic. Images show cells 1-day after seeding (right panel). (B) Images show confluent CEnC cultures at five passages using two culture methods (F99 or M5). (C) Line graph shows mean cell area (μm) at each passage. (D) Line graph shows mean circularity at each passage. Data in (C) and (D) are represented as the mean ±SEM (n=6). Statistical comparisons were performed using two-way ANOVA, with passage and medium defining the variables for this comparison. Scale bars, 100 μm.

A low mitogenic environment maintains a robust CEnC-specific gene expression profile in primary

CEnC

To examine the ability of the cultured cells to maintain a CEnC-specific gene expression profile in low-

or high-mitogenic environments, we compared the expression of 97 genes, previously identified as

specific to ex vivo corneal endothelium (evCEnC), in primary CEnC in M5 versus F99 at each passage

(Fig. 2) (Frausto et al., 2016). At P0, the cells in M5 expressed 83 of the 97 (85.6%) evCEnC-specific

genes, while cells in F99 expressed 76 (78.4%) (Fig. 2A). By P4, the percentages decreased to 75.3%

(73/97) in cells cultured in M5, and 66% (64/97) in cells cultured in F99 (Fig. 2A). We examined the 97

evCEnC-specific genes in the passaged cells to identify those that may be suitable positive selection

markers for assessing the quality of CEnC cultures using the following criteria: 1) expression in 2 or more

sequential passages, starting with P0; 2) decreasing expression over increasing passages (Fig. 2B); and 3)

encoding a cell surface protein. Only *TMEM178A* demonstrated these characteristics (Fig. 2C), and its

expression was validated by qPCR at P0 and P3 (Fig. 2D).

Figure 2. M5 medium maintains CEnC-specific gene expression in cultured CEnC. (A) Line graph shows the number of evCEnC-specific genes expressed in CEnC cultured in either F99 or M5 medium. Dashed line represents the number (97) of evCEnC-specific genes identified in a previous study. (B) Venn diagrams demonstrating the expression of evCEnC-specific genes at passages P0-P3 (P4 not shown due to difficulty visualizing Venn diagrams of greater than 5 sets). *TMEM178A* was identified as a cell surface marker expressed in early passages (at least P0 and P1) in CEnC cultured in either F99 or M5. (C) Line graphs show expression pattern of *TMEM178A* in CEnC cultured in F99 or M5. Gene was considered expressed if TPM value was greater than 7.5 (dashed line). (D) Bar graph shows *TMEM178A* transcript abundances in P0 and P3 CEnC using qPCR. Data in (D) are represented as the mean ±SEM (n=4). Statistical comparisons were performed using one-way ANOVA with post-hoc Tukey test. **, $P<0.01$.

An analysis of the expression of genes purported to be markers for CEnC identity and/or to determine the quality of CEnC in culture revealed that many were neither specific nor highly expressed in corneal endothelium (e.g., *ATP1A1*, *TJP1*, *VIM*, *PRDX6*, *SLC3A2*), or did not correlate well with other quality metrics, such as cell morphology (e.g., *ALCAM1*, *ERBB2*, *CD248*, *SLC25A11*)(Fig. S2). We should note that our data represents transcript abundance, and these markers may prove to be adequate based on protein abundance, but in our study we did not examine the majority of the reported markers for

protein abundance. Of these markers, we found that *SLC4A11* and *CD44* may represent the optimal

markers for selection of high quality cultured CEnC, with *SLC4A11* classified as a positive selection

marker and *CD44* classified as a negative selection marker (Fig. 3). *SLC4A11* and *CD44* transcript

abundances at P0 and P3 in CEnC cultured in both F99 and M5 was determined by RNA-seq and

validated by both qPCR and a Western assay (Fig. 3A). Expression of *SLC4A11*, which was high in

evCEnC, decreased with increasing passage, while expression of *CD44*, which was high in evCEpC,

increased with increasing passage (Fig. 3B).

Figure 3. SLC4A11 and CD44 as cell surface markers for the selection of high quality cultured
CEnC. Assessment of previously published markers and results obtained in this study provide evidence
that SLC4A11 and CD44 represent the most informative markers for the selection of high quality cultured
cells, and represent a minimum for selection. (A) Bar graphs show transcript (RNA-seq and qPCR)
abundance of *SLC4A11* and *CD44* in P0 and P3 CEnC cultured in F99 and M5. Detection and
quantification of SLC4A11 and CD44 was performed using a Western assay (Wes). Bar graphs show
protein quantification. (B) Line graphs show *SLC4A11 and CD44* transcript quantities in CEnC at each
passage (P0 – P4). Bar graphs show transcript abundance in the three main cell types of the cornea
((epithelial cells (evCEnC), keratocytes (evK) and endothelial cells (evCEnC)). Data in bar graphs in (A)
are represented as the mean ±SEM (n=4, mRNA; n=3, protein). Data in line graphs are represented as the
mean TPM at each passage (n=7). Data in bar graphs in (B) are represented as the mean TPM ,±SEM
(n=3).
307

CEnC grown in a low-mitogenic environment possess a robust cellular respiration phenotype

We performed bioinformatics analyses using differentially expressed genes to determine the biological

characteristics of cells cultured with F99 and M5. We identified the genes that were significantly

differentially expressed in cells at each passage between the two growth conditions (Table S4). With

progressive passaging, the number of genes showing differential expression (M5 compared to F99 at each

passage) decreased for cells in each media condition, with the transcriptomes of the cells grown in F99

and cells grown in M5 becoming increasingly more similar (Fig. S3A). As the gene expression difference

was greatest at P0 (between F99 and M5), we performed bioinformatics analyses on the data set obtained

at P0. The genes upregulated in cells with each media at P0 were subsequently analyzed to identify gene

ontology (GO) and pathway terms enriched in our data sets (Table S5). GO terms significantly (q-value

<0.05) enriched in the F99 (P0) upregulated genes data set were associated with cell differentiation and

tissue development (e.g., developmental process, system development, and multicellular organism

development)(Table 1). In addition, significantly enriched pathway terms were associated with cell state

transitions (e.g., TGF-beta signaling pathway and epithelial to mesenchymal transition in colorectal

cancer) and cell senescence (e.g., senescence and autophagy in cancer). The significantly enriched GO

terms in the M5 (P0) upregulated genes data set were associated with cellular respiration (e.g.,

oxidoreductase activity and electron transfer activity) and lipid metabolism (e.g., lipid oxidation and fatty

acid oxidation). Similarly, pathway terms enriched in this data set were also associated with cellular

respiration (e.g., electron transport chain (OXPHOS system in mitochondria)) and lipid metabolism (e.g.,

fatty acid metabolism). GO and pathway enrichment results were similar for the other passages within the

same media group (Table S5).

Senescence of primary CEnC is not dependent on mitogen concentration

To determine whether the observed senescent-associated morphogenic changes coincide with gene

expression changes, we assessed the expression changes for genes previously associated with cell

senescence (Table S6). Many of these genes demonstrated marked expression changes and trends in

expression that are consistent with the senescent phenotype observed for many distinct cell systems. To

determine the functional consequences of changes in gene expression, we identified genes that were

significantly differentially expressed at each passage (P0 was the reference), and for each medium (Table

S7). Passage 3 and 4 showed the greatest number of differentially expressed genes (Fig. S3B), which

coincided with marked changes in morphometric measures. We subsequently performed GO and pathway

analysis to determine the GO and pathway terms enriched in this data set (Table S8). GO terms

significantly (q-value<0.05) enriched in the F99 (P3 and P4) differentially expressed genes data set were

associated with cell senescence-associated GO terms (e.g., mitotic cell cycle, response to stress and

MAPK cascade), and enriched pathway terms also associated with senescence (e.g., p53 signaling

pathway and cellular senescence)(Table 2). Similarly, the significantly enriched GO and pathway terms in

the M5 (P3 and P4) differentially expressed genes data set were also associated with cellular senescence,

but showed significant enrichment for pathway terms associated with cellular energy processing (e.g.,

oxidative phosphorylation and the citric acid (TCA) cycle and respiratory electron transport).

The gene ontology and pathway analyses described above identified biological features in CEnC

as a consequence of differentially expressed genes. However, these analyses are performed in the absence

of the known direction of differential expression (i.e., downregulated and upregulated) for each gene. To

utilize the direction of differential gene expression for the purpose of informing not only pathway

enrichment, but also pathway activation state, we used the Ingenuity Pathway Analysis software (Table

S9). Significantly activated (z-score>1) pathways in all (P3 (F99 and M5) and P4 (F99 and M5)) data sets

were associated with an immune-like response (Table 3). Significantly deactivated (z-score<-1.0)

pathways for all data sets were generally associated with metabolic pathways, which included pathways

associated with lipid metabolism, nucleotide biosynthesis, glycolysis and cellular respiration.

Cell senescence is characterized by an irreversible cell cycle arrest. To determine the impact of

passaging (i.e., expansion) on the cell cycle, we examined the intra-media data sets by focusing on the

predicted activation state (as predicted by IPA software) of cell cycle functions (Table S10). Generally,

this analysis demonstrated activation (z-score>0.5) of cell senescence-associated functions (e.g.,

senescence of cells) and deactivation (z-score<-1.0) of cell cycle functions that have a positive effect on

the cell cycle (e.g., mitosis and cell and cell cycle progression) (Table 4).

Senescence of primary CEnC involves the activation of the p53 and p38-MAPK pathways

To determine whether senescence of primary CEnC involves activation of the p53 and p38-MAPK

pathways and regulation of the cell cycle, we examined expression and activation of genes/proteins

known to regulate cell cycle arrest due to senescence (Fig. 4). We created a custom pathway network that

included p53 (TP53) and p38-MAPK (MAPK14), the cyclin-dependent kinase inhibitors p21^{CIP1}

(CDKN1A) and p16^{INK4} (CDKN2A), and well-established markers of cell senescence (LMNB1,

GADD45A and CD44) (Fig. 4A). In addition, we included the "senescence of cells" as the phenotype

connecting these factors. Initially, we superimposed the differential gene expression results (fold-change)

for genes encoding these proteins, and then we applied an algorithm that predicted the "activation" state

of the other proteins and the phenotype. Applying the prediction algorithm demonstrated activation of

both p53 and p38-MAPK, and activation of "senescence of cells" phenotype in primary CEnC at P3 in

both F99 and M5. Detection of phosphorylation of p53 and p38-MAPK at P3 by Western blotting

provided confirmation of their activation (Fig. 4B). Although *TP53* and *MAPK14* gene expression were

not markedly different between P0 and P3 CEnC, phosphorylation of the encoded proteins was

significantly greater at P3 than P0 ($p < 0.05$)(Fig. 4C). In addition, significant ($p < 0.05$) differential gene

expression of the senescence markers *LMNB1*, *GADD45A* and *CD44* was observed at P3 in either F99 or

M5 or both, a result consistent with cell senescence (Fig. 4D). In addition, we confirmed significantly

increased expression of the CD44 protein at P3, which was consistent with *CD44* transcript levels.

Figure 4. Cellular senescence of cultured CEnC is associated with activation of p53 and p38-MAPK
pathways. (A) Classic cell senescence pathway involving p53 (TP53) and p38 (MAPK14). Pathway
schematics show expression of senescence-associated genes that were differentially expressed in P3
CEnC (left panel). Shades of green, decreased expression; shades of red, increased expression. Pathway
schematics show differentially expressed genes together with activity prediction results (orange,
activated; blue, repressed) for upstream and downstream factors (right panel). (B) Western results show
phosphorylation of TP53 at Serine 15 and MAPK14 at Threonine 180 and Tyrosine 182 in CEnC at P0
and P3, cultured in either F99 or M5. (C) Bar graphs show *TP53* and *MAPK14* transcript levels in P0 and
P3 CEnC using RNA-seq and validated by qPCR. Bar graphs show the quantification of TP53 and
MAPK14 phosphorylation in P0 and P3 CEnC. (D) Bar graphs show the transcript abundances of
LMNB1, *GADD45A* and *CD44*, and validated using qPCR. Bar graphs show CD44 protein abundances
detected using a Western assay. (E) Diagram of the cell cycle including factors that positively (cyclins
and CDK) and negatively (INK4 and KIP/CIP) regulate cell cycle progression. (F) The p21^{CIP1}
(CDKN1A) and p 16^{INK4} (CDKN2A) are inhibitors of cyclin-dependent kinases, and are classically
associated with cellular senescence. Western results show increased expression of CDKN1A and
CDKN2A concomitant with increased expression of the CDK4 and CDK6 in CEnC at P3, cultured in
either F99 or M5. (G) Bar graphs show: *CDKN1A*, *CDKN2A*, *CDK4* and *CDK6* transcript levels in P0
and P3 CEnC using RNA-seq and validated by qPCR; the quantification of the proteins encoded by these
genes in P0 and P3 CEnC. (H) Line graphs show the transcript abundances of the cyclins at each passage

401 in CEnC cultured in F99 or M5. TUBA was used as a loading control in the Westerns. Data in bar graphs
402 are represented as the mean ±SEM (n=4, mRNA; n=3, protein). Statistical comparisons were performed
403 using one-way ANOVA with post-hoc Tukey test. *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$.
404
The cell cycle includes various phases (G1, S, G2 and M) that involve cyclin and cyclin-

dependent kinases (CDK) (Fig. 4E). In the absence of inhibitors of the CDK, cell cycle progression is

dependent on the availability and interaction of cyclins and CDK. However, two classes of inhibitors,

INK4 and KIP/CIP, are able to inhibit cell cycle progression via interaction of the inhibitors (e.g., $p21^{CIP1}$

and/or $p16^{INK4}$) with the cyclin/CDK complexes, an interaction that ultimately blocks the complexes'

downstream actions and results in cell cycle arrest. We assessed protein expression of both $p21^{CIP1}$

(CDKN1A) and $p16^{INK4}$ (CDKN2A), which are upregulated in senescent cells (Fig. 4F). Marked increases

in both of these inhibitors were observed in P3 CEnC cultured in F99 or M5. In addition, accumulation of

CDK4 and CDK6, CDK important in progression through G1/S of the cell cycle, was observed in P3 cells

cultured in F99 and M5 (Fig. 4F). While transcript quantification demonstrated a significant increase in

gene expression in P3 cells only for CDKN2A, the expression of the encoded proteins were either

significantly ($p<0.05$; CDKN1A, CDK4) or markedly (CDKN2A, CDK4 and CDK6) increased in P3

cells cultured in F99 and M5 (Fig. 4G)., As an increase in cyclin D, in conjunction with increases in

$p16^{INK}$ (CDKN2A), CDK4 and CDK6, is a hallmark of G1 cell cycle arrest, we measured the expression

of the four cyclins (cyclins A, B, D and E). A marked and progressive increase in cyclin D was observed

in cells cultured in both F99 and M5, in contrast to the expression of the other three cyclins, which was

unchanged through P4 (Fig. 4H).

Low mitogenic conditions establish a robust functional barrier in CEnC

To determine the impact of cell culture conditions on the establishment of a functional barrier, we

measured resistance of the CEnC to an electrical current (Fig. 5). Initially, we measured barrier function

of cells isolated/grown using unmodified protocols for each of the two methods employed in this study

(Tryp-LN-F99 and ColA-ColIV-M4M5) CEnC cultured using the ColA-ColIV-M4M5 method

demonstrated markedly greater impedance compared with CEnC cultured using the Tryp-LN-F99 method

(Fig. 5A). Impedance data was then used to model cell-cell (R_b) and cell-substrate (α) interactions, which

demonstrated that both contribute to the greater barrier function observed in cells cultured using the

ColA-ColIV-M4M5 method. To identify a potential effect of the dissociation enzymes (trypsin and

collagenase A) on the establishment of a functional barrier, we modified the isolation/growth protocols to

exchange one enzyme for the other. Exchanging the enzymes did not have a marked effect on the

establishment of a functional barrier, with the cells cultured using the protocol containing ColIV-M4M5

establishing a more robust functional barrier compared with cultures established with the protocol

containing LN-F99 (Fig. 5B). To determine the contribution, if any, of the substrate (laminin or collagen)

to which the cells adhere, or of the media (F99 or M5) in which the cells are cultured, we exchanged the

substrates for each other and used F99 and M5 for each (Fig. 5C). Due to the limitation in the number of

replicates we could assay, we chose trypsin (Tryp) as the dissociation enzyme, since use of either trypsin

or collagenase A did not make a significant difference on establishment of a functional barrier. While

only marginal differences in the electrical impedance were observed up to approximately 48 hours, after

that time point the cells that were cultured using the Tryp-LN-M4M5 method demonstrated a

progressively weakening barrier. Additionally, the cells that were cultured using the Tryp-LN-M4M5

method demonstrated a marked decrease in cell-cell and cell-substrate adhesion after 48 hours.

To determine whether the differences observed in barrier function between the two culturing

methods (F99 versus M5) could be explained by differences in barrier-associated proteins or in the

organization of the adhesive interactions between cells, we examined the expression and localization of

the cadherin protein CDH2. While *CDH2* transcript levels measured by qPCR were significantly (p<0.01)

lower in cells cultured in M5, the CDH2 protein levels were not different (Fig. 5D). However, the

organization of the adhesive interaction between cells was markedly different between cells grown in F99

and M5 (Fig. 5E). In F99, the cell-cell interactions appeared diffuse and frayed, although these features

were not consistent for all cell-cell interactions, and lacked features of a well-formed lateral membrane. In

contrast, the cells grown in M5 appeared to have consistent cell-cell interactions with what appears as

prominent lateral membranes, which give the image the impression of depth.

Figure 5. Cultured CEnC establish and maintain robust barrier function in M5 medium. (A) Line
graph shows electrical impedance (Ω at 4000 Hz) for P0 CEnC monolayers established in F99 and M5
media. The contribution of cell-cell adhesion to the resistance of electrical current was modeled from
impedance data and given as R_b ($\Omega \cdot cm^2$). The contribution of cell-substrate adhesion to resistance of
electrical current was modeled from impedance data and given as (α, $\Omega^{\frac{1}{2}} \cdot cm$). (B) Line graph shows
electrical impedance for CEnC monolayers established from cells dissociated with trypsin (Tryp) or
collagenase (ColA), and each cultured in F99 and M5 medium. The contribution of cell-cell (R_b) and cell-
substrate (α) to electrical resistance were modeled from the impedance data. (C) Line graph shows
electrical impedance for CEnC monolayers established from cells dissociated with trypsin (Tryp), seeded
on either laminin (LN) or collagen (ColIV), and cultured in either F99 and M5 medium. The contribution
of cell-cell (R_b) and cell-substrate (α) to electrical resistance was modeled from the impedance data. (D)
Bar graphs show abundance of *CDH2*, a contributor to cell-cell adhesion, in CEnC cultured in F99 and
M5 (top panel). Western results for CDH2 protein and bar graph shows protein quantification (bottom
panel). TUBA was used as a loading control in the Western. (E) Immunofluorescence results show
localization of CDH2 protein in CEnC cultured in F99 and M5. Red, Alexafluor 594; blue, DAPI. Data in
line graphs are represented as the mean ±SD (n=4), and data in bar graphs are represented as the mean
±SEM (n=4, mRNA; n=3, protein). Statistical comparisons were performed using one-way ANOVA with
post-hoc Tukey test. **, P<0.01.
474

A strong functional barrier in primary CEnC is associated with robust expression of adhesion and

glycocalyx proteins

To determine the extent to which barrier-associated proteins may be involved in the more robust

establishment of a functional barrier of CEnC in M5 media, we examined several proteins associated with

either cell-cell adhesion or glycocalyx formation (Fig. 6). We first measured the expression of genes that

encode adhesion-associated proteins (CLDN11, AJAP1, TMEM204 and TMEM178A), and quantified

protein levels for each (Fig. 6A). The expression of each of the four genes was higher in CEnC cultured in

M5 compared to F99, reaching statistical significance (p<0.05) for all four when measured at the

transcript level by qPCR, and for two (CLDN11 and TMEM178A) when measured at the protein level

(p<0.01). We then measured the expression of genes that encode glycocalyx-associated proteins (APOE,

MYOC, DCN, LUM and APOD), and quantified protein levels for each (Fig. 6B). Significantly (p<0.05)

higher transcript levels were observed for all five genes in CEnC cultured in M5, but only two of the

proteins (MYOC and APOD) demonstrated significantly higher levels. The APOE (p=0.05) and LUM

(p=0.14) proteins were markedly higher in M5, but did not achieve statistical significance. DCN did not

demonstrate a difference in protein levels between the two media. Of note, APOD is known to exist in

three forms (pre-modified, post-modified, and complexed), each of progressively greater molecular

weight, all of which demonstrated significantly (p<0.05) higher expression in M5 compared with F99.

Figure 6. M5 medium increases the expression of barrier-associated genes and proteins. (A) Bar
graphs show the expression levels (RNA-seq and qPCR) of genes associated with cell adhesion
(*CLDN11*, *AJAP1*, *TMEM204* and *TMEM178A*) in P0 CEnC. Detection and quantification of the proteins
encoded by these genes were performed using a Western assay (Wes). (B) Bar graphs show the
expression levels (RNA-seq and qPCR) of genes associated with the glycocalyx (*APOE*, *MYOC*, *DCN*,
LUM and APOD). Detection and quantification of the proteins encoded by these genes were performed
using a Western assay (Wes). Three forms of APOD were observed, and each was quantified and
graphed. The protein forms observed were a pre-modified form (1), a post-modified form (2) and a high
molecular weight complex form (3). TUBA was used as a loading control in the Westerns. Data in bar
graphs are represented as the mean ±SEM (n=4, mRNA; n=3, protein). Statistical comparisons were
performed using one-way ANOVA with post-hoc Tukey test. *, P<0.05; **, P<0.01; ***, P<0.001; ****,
P<0.0001.
505

Robust CEnC pump function is established by culture in low mitogenic conditions

The SLC4A4, SLC4A11 and SLC16A1 transporters are highly expressed in in vivo corneal endothelium,

and play a prominent role in maintaining the solute gradients necessary for regulation of water transport

from the stroma to the anterior chamber. To determine the level of expression and functional capacity of

each in F99 and M5 media, we measured transcript levels by RNA-seq and qPCR, protein levels by

Western, and transporter activity with a fluorescence-based transporter assay (Fig. 7). *SLC4A11*

demonstrated significantly higher (p<0.01) and *SLC4A4* demonstrated non-significantly (p=0.081) higher

transcript levels in M5 medium, while the proteins encoded by all three genes demonstrated significantly

(p<0.05) higher levels in M5 compared with F99 (Fig. 7A).

To assess transporter function, we assayed for intracellular proton concentration (i.e., pH_i) over

time, including after changes in buffer perfusion, since each of the transporters transfer protons across the

plasma membrane (Fig. 7B). Traces showing intracellular pH demonstrate sensitivity to buffer changes

for each transporter. Qualitatively, the rate of change in intracellular proton concentration mediated by

SLC4A4 and SLC4A11 following a change in buffer appears greater in M5 compared with F99 medium.

In contrast, the rate of change in intracellular proton concentration mediated by SLC16A1 following a

change in buffer appears greater in F99 compared with M5 medium. Quantitatively, and as an indirect

measure of transporter activity, we calculated the change in intracellular proton concentration over time

$(d[H_i]/dt)$ for each of the assays, and observed significantly (p<0.01) higher SLC4A4 and SLC4A11 and

significantly (p<0.01) lower activity for SLC16A1 for cells cultured in M5.

Figure 7. SLC4A4 and SLC4A11 transporter activity is increased in M5 medium. (A) Bar graphs
show transcript abundances (RNA-seq and qPCR) in P0 CEnC for genes that encode membrane bound
transporters (*SLC4A4*, *SLC4A11* and *SLC16A1*). Detection and quantification of the encoded proteins was
performed using a Western assay (Wes). (B) Traces show change in pH levels before and after perfusion
with buffer containing indicated substrate specified for the transporter (SLC4A4, Na^+; SLC4A11, H^+;
SLC16A1, lactate). Arrows indicate time at which buffer was changed. Bar graphs show the rate of
change in intracellular proton concentration [Hi], which was calculated as a proxy for transporter activity.
Data in bar graphs are represented as the mean ±SEM (n=4, mRNA; n=3, protein; n=6, transporter
activity). Statistical comparisons were performed using one-way ANOVA with post-hoc Tukey test. *,
$P<0.05$; **, $P<0.01$; ****, $P<0.0001$.
537

Low mitogenic conditions cause decreased cell migration capacity

We measured cell migration using a non-wounding method (Fig. 8). Phase contrast imaging of cell

migration demonstrated significantly (p<0.001) less gap closure for CEnC in M5 compared with F99

medium (Fig. 8A and B). We observed that cells cultured in M5, while possessing lower migration

capacity, also appear to markedly cover the gap primarily by increasing the cell size rather than by

migration (Fig. 8C). We calculated the ratio of the area occupied by the cells that "migrated" into the gap

versus cells more distant from the gap, and which have not substantially changed their size compared with

prior to when migration was initiated (Fig. 8D). Cell size for cells cultured in M5 and occupying any

portion of the gap cover a significantly (p<0.05) larger area compared with cells cultured in F99.

Figure 8. CEnC cultured in M5 medium possess attenuated migration capacity. (A) Phase contrast
images show initial gap (0 hours) and gap closure (22 hours) of P0 CEnC cultured in F99 and M5. (B)
Bar graph shows quantification of gap closure at 22 hours as a percent of the initial gap . (C) Phase
contrast images show gap closure at 60 hours. (D) Bar graph shows the ratio of the mean area at 60 hours
of the cells near or populating the gap divided by the mean area of the cells not near the gap. Data in bar
graphs are represented as the mean ±SEM (n=4). Statistical comparisons were performed using one-way
ANOVA with post-hoc Tukey test. *, P<0.05; ***, P<0.001.

DISCUSSION

Corneal clarity depends on the corneal endothelium, and disease or injury to this tissue causes loss of

visual acuity that may necessitate corneal transplantation. The development and widespread adoption of

endothelial keratoplasty techniques has significantly improved outcomes of the surgical management of

corneal endothelial cell dysfunction although postoperative complications remain (Deng et al., 2015; Lass

et al., 2017; Van den Bogerd et al., 2018). However, The current paradigm of one donor cornea for one

recipient requires cornea donation, recovery and utilization rates that are present in only a few countries

in the world that are able to meet the domestic need for transplantable corneas. Thus, scientist and

clinicians have developed various methods for isolation and in vitro culturing of CEnC, with the aim of

achieving sufficient cell numbers of transplantable high quality CEnC to challenge the one donor-one

recipient paradigm (Soh et al., 2017). While several metrics have been used to assess the identity (e.g.,

morphology, biomarker expression) and function (e.g., transcellular resistance, intracellular proton

concentration) and of cultured CEnC, a comprehensive study of the effects of in vitro expansion on the

CEnC transcriptome has not been performed. The transcriptomic work performed as part of this study

may be used as one of the criteria for assessing the quality and viability of in vitro expanded CEnC for

use in cell-replacement therapy. In particular, because cell senescence is a feature of primary cell culture,

it is important to identify all the potential signaling and metabolic pathways involved in CEnC senescence

in the hope of delaying senescence by targeting these pathways, and therefore producing a larger number

of therapeutically suitable cells.

In 1978, the first human CEnC cultures were described, but given a lack of human-specific

protocols, the researchers used culture conditions optimized for growth of rabbit CEnC (Jumblatt et al.,

1978). Over the next decade, techniques for the isolation of human CEnC and growth media for the in

vitro culture of human CEnC were optimized. In 1989, Engelmann et al. determined that a 1:1 mixture of

M199 and F12 media was the optimal base medium for culture of primary CEnC (Engelmann and Friedl,

1989). Two decades later, the formulation of the medium was modified to induce cell cycle re-entry (due

to relatively mitogen-rich conditions) and establishment of contact inhibition, leading to quiescence, a

non-proliferative cell state that characterize CEnC in vivo (Valtink et al., 2008). We started culturing

primary CEnC in 2011, and chose the medium formulation described by Valtink et al. as it was at the time

the most effective formulation for the culture of both primary and immortalized CEnC. Nevertheless,

about 40% (3/7 at P0 in this study) of the cultures we established did not possess CEnC-like morphology,

and were generally characterized by a mixture of morphologies (some resembling fibroblasts), an

observation consistent with previous findings by the developers of the formulation (personal

communication). In 2015, Peh et al. described a novel approach to in vitro culturing of CEnC in which

cell proliferation was accomplished by growth in a mitogenic-rich medium, followed by maintenance in a

medium with reduced mitogens (Peh et al., 2015). Coincidently, a recent report independently

demonstrated the superiority of the dual media approach (Bartakova et al., 2018). Taken together, these

studies indicated that the low mitogenic medium established cultures with better morphology, marker

expression and function compared with growth in a mitogen-rich medium alone. Herein, we compared the

two culture methods published by Valtnik et al. and Peh et al. using extensive bioinformatics analyses and

additional functional techniques to determine which method performs best in maintaining the CEnC

phenotype throughout in vitro expansion.

597   Cell morphology is an initial metric for assessing the quality of CEnC cultures. We observed that

the low mitogenic environment was better at maintaining a CEnC-like morphology, but continued

passaging caused a distinct morphogenic transformation consistent with cellular senescence. This change

in morphology correlated with a progressive loss of CEnC identity as defined by the expression of

evCEnC-specific genes. Temporal changes in evCEnC-specific gene expression as a consequence of

passaging allowed us to identify a novel cell surface marker, encoded by *TMEM178A*, as a potential

positive selection marker for high quality cultured CEnC. In addition, after a reassessment of reported

selection markers, we propose SLC4A11 (positive), TMEM178A (positive) and CD44 (negative) as the

minimum for selection of high quality in vitro CEnC (Bartakova et al., 2018; Cheong et al., 2013; Chng et

al., 2013; Ding et al., 2014; Frausto et al., 2016; Okumura et al., 2014a; Toda et al., 2017; Yoshihara et

al., 2015). SLC4A11 is a highly expressed transporter protein in corneal endothelium that plays a

significant role in CEnC function, and biallelic mutation of this protein leads to congenital hereditary

endothelial dystrophy (Aldave et al., 2013; Jiao et al., 2007; Vithana et al., 2006). Conversely, CD44 is

not expressed in the corneal endothelium, and its expression correlates with a senescent phenotype in

vascular and corneal endothelial cells (Mun and Boo, 2010; Ueno et al., 2016).

612   A prominent barrier to expansion of CEnC in vitro for use in a cell-based therapy is the limited

proliferative capacity of primary CEnC. In vivo, CEnC are in a state of cellular quiescence, which in this

cell type is specifically a cell cycle arrest at the G1 phase that is established by contact-inhibition,

maintained by TGFβ2, and regulated by p27^{KIP1} (Joyce, 2003). However, in vivo CEnC have been shown

to express senescent markers in older donors, which is postulated to contribute to the lower proliferative

capacity of primary CEnC isolated from older donors compared with those from younger donors.

Nevertheless, CEnC are able to re-engage the cell cycle by loosening of cell-cell contacts and/or exposure

to potent mitogenic factors. As with most primary cells, primary CEnC have a limited life span under

proliferative (i.e., passaging and growth in mitogen-rich) conditions. This limitation is caused by cellular

senescence of CEnC (in vitro and ex vivo), and may be, in part, a consequence of oxidative stress (Joyce

et al., 2009). This is supported by our observations in late passage CEnC (P3 and/or P4) that p38MAPK

was phosphorylated (stress response), that expression of p16^{INK4} (gene and protein) was increased, and

that the set of differentially expressed genes was enriched for genes implicated in oxidative stress-

associated pathways (e.g., NRF2-mediated oxidative stress response, DNA damage checkpoint

regulation). In addition, there is also an apparent repression of oxidative phosphorylation (i.e.,

mitochondrial activity), which was observed in our pathway analysis. Moreover, we also identified

enrichment for genes associated with p53 signaling, an observation made previously (Sheerin et al.,

2012), and this coincided with phosphorylation of p53 and increased expression of p21^{CIP1} (CDKN1A)

(gene and protein) in senescent CEnC (i.e., P3 and P4). Cell senescence in primary CEnC in this study

was also associated with increased expression of CDK4 and CDK6 proteins (but not gene expression),

and increased *cyclin D* (*CCND1*) gene expression. Taken together, this represents a gene/protein

expression profile consistent with senescence-induced G1 cell cycle arrest. Overall, activation of the

p38MAPK and p53 pathways did not appear significantly influenced by growth in a low- or high-

mitogenic environment.

Achieving and maintaining characteristic CEnC functional properties is essential for the clinical

utility of in vitro CEnC in management of CEnC dysfunction. The corneal endothelium is a monolayer

mosaic of CEnC, and its primary function is to maintain the corneal stroma in a partially dehydrated state

to preserve its optical clarity. It does this by establishing a robust barrier conferred by cell-cell and cell-

substrate interactions, which allow for passive leak of water into the stroma, but not most solutes

(Srinivas, 2010). Through the combined action of various solute transporters, unidirectional translocation

of water from the stroma into the aqueous humor occurs (Bonanno, 2012). Taken together, this describes

the "pump-leak" hypothesis of water homeostasis in the cornea. We demonstrated that a low mitogenic

environment combined with culture on collagen represented better in vitro conditions for establishing a

robust CEnC barrier (as measured by electrical impedance) compared with a high mitogenic environment

and culture on laminin. Recent studies showed that collagen, compared with various other extracellular

matrix proteins, including laminin, and low mitogen-containing media were optimal for establishing

robust barrier function in a CEnC line (Santander-Garcia et al., 2016) and primary CEnC (Bartakova et

al., 2018). Detection of CDH2, a CEnC-specific marker, in CEnC cultures in low mitogenic conditions

demonstrated that cell adhesions (cell-cell and cell-substrate) are considerably more complex, and showed

striking resemblance to the organization of adhesions in ex vivo human corneas (He et al., 2016). We also

demonstrated that this increase in barrier function is associated with an increase in adhesion/glycocalyx-

associated protein expression. To our knowledge, this represents the first report implicating glycocalyx

proteins in corneal endothelial biology. Although some of the barrier-associated proteins we examined in

this study have been associated with the glycocalyx of the vascular endothelium (Friden et al., 2011;

Reitsma et al., 2007), it remains to be determined if the increase in glycocalyx-associated proteins

coincides with the formation of a glycocalyx in corneal endothelium either in vitro or in vivo. However,

early reports suggest that the human corneal endothelium may possess a glycocalyx-like layer composed

polysaccharides, glycosaminoglycans and other, as yet, unidentified molecules (Hornung and Wollensak,

1979; Jacobsen and Sperling, 1978; Schroder and Sperling, 1977; Sperling and Jacobsen, 1980). We also

showed that at least two transporters critical for CEnC function (SLC4A4 and SLC4A11) are expressed at

significantly higher levels and show increased transporter activity in low mitogenic conditions compared

with high mitogenic conditions. As such, CEnC expanded in mitogenic-rich medium and maintained in

low mitogenic conditions show functional features consistent with the CEnC requirement for regulating

water homeostasis in the cornea. These results are consistent with previous studies demonstrating an

intact pump function (i.e., prevention/mitigation of edema) in in vitro expanded corneal endothelial cells

in an animal model of bullous keratopathy (Peh et al., 2017; Peh et al., 2019).

668   A long known characteristic of in vivo CEnC is that when mild cell loss occurs, the vacated area

is covered by migration and increase in size of adjacent cells (Sherrard, 1976). To determine if this

phenomenon is recapitulated in vitro, we performed a non-wounding cell migration assay. Cells in a high

mitogenic environment migrated and/or proliferated (not distinguished in our study) to cover the gap with

no significant increase in cell size. In contrast, CEnC in low mitogenic conditions behaved in a manner

consistent with their behavior in vivo, so that these cells showed low migration capacity, but a significant

propensity to increase cell size to cover the gap. In addition, these results are consistent with the

differences observed for cell barrier formation, because increased cell-cell adhesion is antithetical to

robust cell migration (and also to robust cell proliferation capacity).

677   In summary, we provide evidence that CEnC proliferation in a high mitogenic medium, followed

by maintenance of contact-inhibition of confluent monolayers in a low mitogenic medium, supports

establishment of CEnC in which in vivo function is significantly recapitulated and permits sufficient

expansion of CEnC to significantly challenge the "one donor-one recipient" paradigm. In our current

study, we can theoretically achieve a 1:5-1:15 donor:recipient ratio (donor:recipient, 1:5 (P0) or 1:15 (P1)

– at approximately 3000 cells/mm^2), as these are the passages at which the cells possess the most robust

CEnC phenotype as determined by gene (or marker) expression and cell morphology. Maintaining the

CEnC morphologic and functional phenotype at higher passage numbers remains an active area of

research. Also, while we examined cell function at an early passage (typically at P0), we recognize that

functional analysis of cells will need to be performed at later passages to ensure viability as a therapeutic

modality. The progression to senescence in vitro remains a significant barrier to the expansion of

cultured CEnC. Identification of oxidative stress as a component of CEnC senescence has prompted

investigators to use p38MAPK inhibitors to delay the onset of senescence, but this approach has shown

mixed results (Hongo et al., 2017; Nakahara et al., 2018; Sheerin et al., 2012). Similarly, ascorbic acid

was a supplement added to early media formulations (added to F99 and M4) to reduce/inhibit oxidant-

induced stress/apoptosis (Serbecic and Beutelspacher, 2005; Shima et al., 2011). In addition, senescent

CEnC in vitro are characterized by the Senescence-Associated Secretory Phenotype (SASP), which

includes the secretion of immune-regulating factors (cytokines, chemokines and growth factors) that have

been demonstrated to support tumorigenesis in adjacent epithelial tissues (Georgilis et al., 2018; Laberge

et al., 2015; Lau and David, 2019; Lopes-Paciencia et al., 2019). As a remedy to senescence, we identify

both classical and novel pathways associated with CEnC senescence that can be manipulated

experimentally to potentially achieve significant CEnC expansion with minimal senescence. In addition,

we identified which previously reported markers represent the optimal markers for selection of high

quality cultured CEnC, and propose *TMEM178A* as a novel positive selection marker. We propose that

because senescent CEnC can be present in cultured CEnC preparations and that senescent cells may

support tumorigenesis, a Descemet membrane biomimetic carrier is preferred to injection of a suspension

of cells, to both maximize cell count and minimize their tumorigenic potential (Gutermuth et al., 2019;

Kinoshita et al., 2018; Teichmann et al., 2013). In conclusion, while additional changes to the dual media

method may improve the culturing of CEnC in vitro, we provide evidence that this method is preferred to

continuous culture in mitogen-rich media.

REFERENCES

Eye Bank Association of America. 2018 Eye Banking Statistical Report. 2018.

Aldave, A.J., Han, J., and Frausto, R.F. (2013). Genetics of the corneal endothelial dystrophies: an
evidence-based review. Clin Genet *84*, 109-119.

Baratz, K.H., Tosakulwong, N., Ryu, E., Brown, W.L., Branham, K., Chen, W., Tran, K.D., Schmid-
Kubista, K.E., Heckenlively, J.R., Swaroop, A., Abecasis, G., Bailey, K.R., and Edwards, A.O. (2010).
E2-2 protein and Fuchs's corneal dystrophy. N Engl J Med *363*, 1016-1024.

Bartakova, A., Kuzmenko, O., Alvarez-Delfin, K., Kunzevitzky, N.J., and Goldberg, J.L. (2018). A Cell
Culture Approach to Optimized Human Corneal Endothelial Cell Function. Invest Ophthalmol Vis Sci *59*,
1617-1629.

Bednarz, J., Rodokanaki-von Schrenck, A., and Engelmann, K. (1998). Different characteristics of
endothelial cells from central and peripheral human cornea in primary culture and after subculture. In
Vitro Cell Dev Biol Anim *34*, 149-153.

Bonanno, J.A. (2012). Molecular mechanisms underlying the corneal endothelial pump. Exp Eye Res *95*,
2-7.

Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-seq
quantification. Nat Biotechnol *34*, 525-527.

Campisi, J. (2013). Aging, cellular senescence, and cancer. Annu Rev Physiol *75*, 685-705.

Cheong, Y.K., Ngoh, Z.X., Peh, G.S., Ang, H.P., Seah, X.Y., Chng, Z., Colman, A., Mehta, J.S., and Sun,
W. (2013). Identification of cell surface markers glypican-4 and CD200 that differentiate human corneal
endothelium from stromal fibroblasts. Invest Ophthalmol Vis Sci *54*, 4538-4547.

Chng, Z., Peh, G.S., Herath, W.B., Cheng, T.Y., Ang, H.P., Toh, K.P., Robson, P., Mehta, J.S., and
Colman, A. (2013). High throughput gene expression analysis identifies reliable expression markers of
human corneal endothelial cells. PLoS One *8*, e67546.

Deng, S.X., Sanchez, P.J., and Chen, L. (2015). Clinical outcomes of Descemet membrane endothelial
keratoplasty using eye bank-prepared tissues. Am J Ophthalmol *159*, 590-596.

Ding, V., Chin, A., Peh, G., Mehta, J.S., and Choo, A. (2014). Generation of novel monoclonal antibodies
for the enrichment and characterization of human corneal endothelial cells (hCENC) necessary for the
treatment of corneal endothelial blindness. MAbs *6*, 1439-1452.

Engelmann, K., and Friedl, P. (1989). Optimization of culture conditions for human corneal endothelial
cells. In Vitro Cell Dev Biol *25*, 1065-1072.

Frausto, R.F., Le, D.J., and Aldave, A.J. (2016). Transcriptomic Analysis of Cultured Corneal Endothelial
Cells as a Validation for Their Use in Cell Replacement Therapy. Cell Transplant *25*, 1159-1176.

Friden, V., Oveland, E., Tenstad, O., Ebefors, K., Nystrom, J., Nilsson, U.A., and Haraldsson, B. (2011).
The glomerular endothelial cell coat is essential for glomerular filtration. Kidney Int *79*, 1322-1330.

Georgilis, A., Klotz, S., Hanley, C.J., Herranz, N., Weirich, B., Morancho, B., Leote, A.C., D'Artista, L.,
Gallage, S., Seehawer, M., Carroll, T., Dharmalingam, G., Wee, K.B., Mellone, M., Pombo, J., Heide, D.,
Guccione, E., Arribas, J., Barbosa-Morais, N.L., Heikenwalder, M., Thomas, G.J., Zender, L., and Gil, J.
(2018). PTBP1-Mediated Alternative Splicing Regulates the Inflammatory Secretome and the Pro-
tumorigenic Effects of Senescent Cells. Cancer Cell *34*, 85-102 e109.

Gottsch, J.D., Bowers, A.L., Margulies, E.H., Seitzman, G.D., Kim, S.W., Saha, S., Jun, A.S., Stark,
W.J., and Liu, S.H. (2003). Serial analysis of gene expression in the corneal endothelium of Fuchs'
dystrophy. Invest Ophthalmol Vis Sci *44*, 594-599.

Gutermuth, A., Maassen, J., Harnisch, E., Kuhlen, D., Sauer-Budge, A., Skazik-Voogt, C., and
Engelmann, K. (2019). Descemet's Membrane Biomimetic Microtopography Differentiates Human
Mesenchymal Stem Cells Into Corneal Endothelial-Like Cells. Cornea *38*, 110-119.

He, Z., Forest, F., Gain, P., Rageade, D., Bernard, A., Acquart, S., Peoc'h, M., Defoe, D.M., and Thuret,
G. (2016). 3D map of the human corneal endothelial cell. Sci Rep *6*, 29047.

Hongo, A., Okumura, N., Nakahara, M., Kay, E.P., and Koizumi, N. (2017). The Effect of a p38
Mitogen-Activated Protein Kinase Inhibitor on Cellular Senescence of Cultivated Human Corneal
Endothelial Cells. Invest Ophthalmol Vis Sci *58*, 3325-3334.

Hornung, M., and Wollensak, J. (1979). [Proof of glycosaminoglycans in the corneal endothelium].
Albrecht Von Graefes Arch Klin Exp Ophthalmol *211*, 67-84.

Jacobsen, S.R., and Sperling, S. (1978). Scanning electron microscopic observations of a
mucopolysaccharide coating on human corneal endothelium. Acta Ophthalmol (Copenh) *56*, 161-167.

Jiao, X., Sultana, A., Garg, P., Ramamurthy, B., Vemuganti, G.K., Gangopadhyay, N., Hejtmancik, J.F.,
and Kannabiran, C. (2007). Autosomal recessive corneal endothelial dystrophy (CHED2) is associated
with mutations in SLC4A11. J Med Genet *44*, 64-68.

Joyce, N.C. (2003). Proliferative capacity of the corneal endothelium. Prog Retin Eye Res *22*, 359-389.

Joyce, N.C. (2005). Cell cycle status in human corneal endothelium. Exp Eye Res *81*, 629-638.

Joyce, N.C., Zhu, C.C., and Harris, D.L. (2009). Relationship among oxidative stress, DNA damage, and
proliferative capacity in human corneal endothelium. Invest Ophthalmol Vis Sci *50*, 2116-2122.

Jumblatt, M.M., Maurice, D.M., and McCulley, J.P. (1978). Transplantation of tissue-cultured corneal
endothelium. Invest Ophthalmol Vis Sci *17*, 1135-1141.

Kao, L., Azimov, R., Shao, X.M., Frausto, R.F., Abuladze, N., Newman, D., Aldave, A.J., and Kurtz, I.
(2016). Multifunctional ion transport properties of human SLC4A11: comparison of the SLC4A11-B and
SLC4A11-C variants. Am J Physiol Cell Physiol *311*, C820-C830.

Kinoshita, S., Koizumi, N., Ueno, M., Okumura, N., Imai, K., Tanaka, H., Yamamoto, Y., Nakamura, T.,
Inatomi, T., Bush, J., Toda, M., Hagiya, M., Yokota, I., Teramukai, S., Sotozono, C., and Hamuro, J.
(2018). Injection of Cultured Cells with a ROCK Inhibitor for Bullous Keratopathy. N Engl J Med *378*,
995-1003.

Krafchak, C.M., Pawar, H., Moroi, S.E., Sugar, A., Lichter, P.R., Mackey, D.A., Mian, S., Nairus, T.,
Elner, V., Schteingart, M.T., Downs, C.A., Kijek, T.G., Johnson, J.M., Trager, E.H., Rozsa, F.W.,
Mandal, M.N., Epstein, M.P., Vollrath, D., Ayyagari, R., Boehnke, M., and Richards, J.E. (2005).
Mutations in TCF8 cause posterior polymorphous corneal dystrophy and ectopic expression of COL4A3
by corneal endothelial cells. Am J Hum Genet *77*, 694-708.

Laberge, R.M., Sun, Y., Orjalo, A.V., Patil, C.K., Freund, A., Zhou, L., Curran, S.C., Davalos, A.R.,
Wilson-Edell, K.A., Liu, S., Limbad, C., Demaria, M., Li, P., Hubbard, G.B., Ikeno, Y., Javors, M.,
Desprez, P.Y., Benz, C.C., Kapahi, P., Nelson, P.S., and Campisi, J. (2015). MTOR regulates the pro-
tumorigenic senescence-associated secretory phenotype by promoting IL1A translation. Nat Cell Biol *17*,
1049-1061.

Lass, J.H., Benetz, B.A., Verdier, D.D., Szczotka-Flynn, L.B., Ayala, A.R., Liang, W., Aldave, A.J.,
Dunn, S.P., McCall, T., Mian, S.I., Navarro, L.C., Patel, S.V., Pramanik, S., Rosenwasser, G.O., Ross,
K.W., Terry, M.A., Kollman, C., Gal, R.L., Beck, R.W., and Cornea Preservation Time Study, G. (2017).
Corneal Endothelial Cell Loss 3 Years After Successful Descemet Stripping Automated Endothelial
Keratoplasty in the Cornea Preservation Time Study: A Randomized Clinical Trial. JAMA Ophthalmol
135, 1394-1400.

Lau, L., and David, G. (2019). Pro- and Anti-Tumorigenic Functions of the Senescence-Associated
Secretory Phenotype. Expert Opin Ther Targets.

Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods *25*, 402-408.

Lopes-Paciencia, S., Saint-Germain, E., Rowell, M.C., Ruiz, A.F., Kalegari, P., and Ferbeyre, G. (2019).
The senescence-associated secretory phenotype and its regulation. Cytokine *117*, 15-22.

Mehta, J.S., Kocaba, V., and Soh, Y.Q. (2019). The future of keratoplasty: cell-based therapy,
regenerative medicine, bioengineering keratoplasty, gene therapy. Curr Opin Ophthalmol *30*, 286-291.

Mimura, T., and Joyce, N.C. (2006). Replication competence and senescence in central and peripheral
human corneal endothelium. Invest Ophthalmol Vis Sci *47*, 1387-1396.

Mun, G.I., and Boo, Y.C. (2010). Identification of CD44 as a senescence-induced cell adhesion gene
responsible for the enhanced monocyte recruitment to senescent endothelial cells. Am J Physiol Heart
Circ Physiol *298*, H2102-2111.

Nakahara, M., Okumura, N., Nakano, S., and Koizumi, N. (2018). Effect of a p38 Mitogen-Activated
Protein Kinase Inhibitor on Corneal Endothelial Cell Proliferation. Invest Ophthalmol Vis Sci *59*, 4218-
4227.

Okumura, N., Hirano, H., Numata, R., Nakahara, M., Ueno, M., Hamuro, J., Kinoshita, S., and Koizumi,
N. (2014a). Cell surface markers of functional phenotypic corneal endothelial cells. Invest Ophthalmol
Vis Sci *55*, 7610-7618.

Okumura, N., Kinoshita, S., and Koizumi, N. (2014b). Cell-based approach for treatment of corneal
endothelial dysfunction. Cornea *33 Suppl 11*, S37-41.

Parekh, M., Ferrari, S., Sheridan, C., Kaye, S., and Ahmad, S. (2016). Concise Review: An Update on the
Culture of Human Corneal Endothelial Cells for Transplantation. Stem Cells Transl Med *5*, 258-264.

Peh, G.S., Chng, Z., Ang, H.P., Cheng, T.Y., Adnan, K., Seah, X.Y., George, B.L., Toh, K.P., Tan, D.T.,
Yam, G.H., Colman, A., and Mehta, J.S. (2015). Propagation of human corneal endothelial cells: a novel
dual media approach. Cell Transplant *24*, 287-304.

Peh, G.S.L., Ang, H.P., Lwin, C.N., Adnan, K., George, B.L., Seah, X.Y., Lin, S.J., Bhogal, M., Liu,
Y.C., Tan, D.T., and Mehta, J.S. (2017). Regulatory Compliant Tissue-Engineered Human Corneal
Endothelial Grafts Restore Corneal Function of Rabbits with Bullous Keratopathy. Sci Rep *7*, 14149.

Peh, G.S.L., Ong, H.S., Adnan, K., Ang, H.P., Lwin, C.N., Seah, X.Y., Lin, S.J., and Mehta, J.S. (2019).
Functional Evaluation of Two Corneal Endothelial Cell-Based Therapies: Tissue-Engineered Construct
and Cell Injection. Sci Rep *9*, 6087.

Pimentel, H., Bray, N.L., Puente, S., Melsted, P., and Pachter, L. (2017). Differential analysis of RNA-
seq incorporating quantification uncertainty. Nat Methods *14*, 687-690.

Reitsma, S., Slaaf, D.W., Vink, H., van Zandvoort, M.A., and oude Egbrink, M.G. (2007). The
endothelial glycocalyx: composition, functions, and visualization. Pflugers Arch *454*, 345-359.

Santander-Garcia, D., Ortega, M.C., Benito-Martinez, S., Barroso, S., Jimenez-Alfaro, I., and Millan, J.
(2016). A human cellular system for analyzing signaling during corneal endothelial barrier dysfunction.
Exp Eye Res *153*, 8-13.

Schmedt, T., Chen, Y., Nguyen, T.T., Li, S., Bonanno, J.A., and Jurkunas, U.V. (2012). Telomerase
immortalization of human corneal endothelial cells yields functional hexagonal monolayers. PLoS One *7*,
e51427.

Schroder, H.D., and Sperling, S. (1977). Polysaccharide coating of human corneal endothelium. Acta
Ophthalmol (Copenh) *55*, 819-826.

Serbecic, N., and Beutelspacher, S.C. (2005). Vitamins inhibit oxidant-induced apoptosis of corneal
endothelial cells. Jpn J Ophthalmol *49*, 355-362.

Sheerin, A.N., Smith, S.K., Jennert-Burston, K., Brook, A.J., Allen, M.C., Ibrahim, B., Jones, D., Wallis,
C., Engelmann, K., Rhys-Williams, W., Faragher, R.G., and Kipling, D. (2012). Characterization of
cellular senescence mechanisms in human corneal endothelial cells. Aging Cell *11*, 234-240.

Sherrard, E.S. (1976). The corneal endothelium in vivo: its response to mild trauma. Exp Eye Res *22*,
347-357.

Shima, N., Kimoto, M., Yamaguchi, M., and Yamagami, S. (2011). Increased proliferation and replicative
lifespan of isolated human corneal endothelial cells with L-ascorbic acid 2-phosphate. Invest Ophthalmol
Vis Sci *52*, 8711-8717.

Soh, Y.Q., Peh, G.S.L., and Mehta, J.S. (2017). Translational issues for human corneal endothelial tissue
engineering. J Tissue Eng Regen Med *11*, 2425-2442.

Spandidos, A., Wang, X., Wang, H., Dragnev, S., Thurber, T., and Seed, B. (2008). A comprehensive
collection of experimentally validated primers for Polymerase Chain Reaction quantitation of murine
transcript abundance. BMC Genomics *9*, 633.

Spandidos, A., Wang, X., Wang, H., and Seed, B. (2010). PrimerBank: a resource of human and mouse
PCR primer pairs for gene expression detection and quantification. Nucleic Acids Res *38*, D792-799.

Sperling, S., and Jacobsen, S.R. (1980). The surface coat on human corneal endothelium. Acta
Ophthalmol (Copenh) *58*, 96-102.

Srinivas, S.P. (2010). Dynamic regulation of barrier integrity of the corneal endothelium. Optom Vis Sci
87, E239-254.

Stolwijk, J.A., Matrougui, K., Renken, C.W., and Trebak, M. (2015). Impedance analysis of GPCR-
mediated changes in endothelial barrier function: overview and fundamental considerations for stable and
reproducible measurements. Pflugers Arch *467*, 2193-2218.

Teichmann, J., Valtink, M., Nitschke, M., Gramm, S., Funk, R.H., Engelmann, K., and Werner, C.
(2013). Tissue engineering of the corneal endothelium: a review of carrier materials. J Funct Biomater *4*,
178-208.

Toda, M., Ueno, M., Hiraga, A., Asada, K., Montoya, M., Sotozono, C., Kinoshita, S., and Hamuro, J.
(2017). Production of Homogeneous Cultured Human Corneal Endothelial Cells Indispensable for
Innovative Cell Therapy. Invest Ophthalmol Vis Sci *58*, 2011-2020.

Ueno, M., Asada, K., Toda, M., Nagata, K., Sotozono, C., Kosaka, N., Ochiya, T., Kinoshita, S., and
Hamuro, J. (2016). Concomitant Evaluation of a Panel of Exosome Proteins and MiRs for Qualification
of Cultured Human Corneal Endothelial Cells. Invest Ophthalmol Vis Sci *57*, 4393-4402.

Valtink, M., Gruschwitz, R., Funk, R.H., and Engelmann, K. (2008). Two clonal cell lines of
immortalized human corneal endothelial cells show either differentiated or precursor cell characteristics.
Cells Tissues Organs *187*, 286-294.

Van den Bogerd, B., Dhubhghaill, S.N., Koppen, C., Tassignon, M.J., and Zakaria, N. (2018). A review
of the evidence for in vivo corneal endothelial regeneration. Surv Ophthalmol *63*, 149-165.

Vithana, E.N., Morgan, P., Sundaresan, P., Ebenezer, N.D., Tan, D.T., Mohamed, M.D., Anand, S.,
Khine, K.O., Venkataraman, D., Yong, V.H., Salto-Tellez, M., Venkatraman, A., Guo, K., Hemadevi, B.,
Srinivasan, M., Prajna, V., Khine, M., Casey, J.R., Inglehearn, C.F., and Aung, T. (2006). Mutations in
sodium-borate cotransporter SLC4A11 cause recessive congenital hereditary endothelial dystrophy
(CHED2). Nat Genet *38*, 755-757.

Wang, X., and Seed, B. (2003). A PCR primer bank for quantitative gene expression analysis. Nucleic
Acids Res *31*, e154.

Wieben, E.D., Aleff, R.A., Tosakulwong, N., Butz, M.L., Highsmith, W.E., Edwards, A.O., and Baratz,
K.H. (2012). A common trinucleotide repeat expansion within the transcription factor 4 (TCF4, E2-2)
gene predicts Fuchs corneal dystrophy. PLoS One *7*, e49083.

Yamashita, K., Inagaki, E., Hatou, S., Higa, K., Ogawa, A., Miyashita, H., Tsubota, K., and Shimmura, S.
(2018). Corneal Endothelial Regeneration Using Mesenchymal Stem Cells Derived from Human
Umbilical Cord. Stem Cells Dev *27*, 1097-1108.

Yoshihara, M., Ohmiya, H., Hara, S., Kawasaki, S., consortium, F., Hayashizaki, Y., Itoh, M., Kawaji, H.,
Tsujikawa, M., and Nishida, K. (2015). Discovery of molecular markers to discriminate corneal
endothelial cells in the human body. PLoS One *10*, e0117581.
892

893   Table 1. Gene ontology and pathway enrichment analysis of the P0 inter-media comparison.

Source	Term Name	Term ID	FDR p-value
F99 Passage 0			
GO:BP	developmental process	GO:0032502	4.2E-18
GO:BP	system development	GO:0048731	1.3E-16
GO:BP	multicellular organism development	GO:0007275	1.1E-15
GO:BP	anatomical structure morphogenesis	GO:0009653	3.6E-15
GO:BP	tissue development	GO:0009888	2.1E-14
GO:BP	cellular developmental process	GO:0048869	7.8E-14
GO:BP	cell differentiation	GO:0030154	2.3E-13
GO:BP	cell development	GO:0048468	3.6E-09
GO:BP	tube development	GO:0035295	3.8E-09
GO:BP	blood vessel development	GO:0001568	4.4E-09
GO:BP	circulatory system development	GO:0072359	4.6E-09
GO:BP	regulation of cell differentiation	GO:0045595	6.0E-09
GO:BP	nervous system development	GO:0007399	1.0E-07
GO:BP	epithelium development	GO:0060429	3.7E-07
KEGG	TGF-beta signaling pathway	KEGG:04350	3.2E-02
WP	miR-509-3p alteration of YAP1/ECM axis	WP:WP3967	5.1E-04
WP	Epithelial to mesenchymal transition in colorectal cancer	WP:WP4239	2.1E-02
WP	Senescence and Autophagy in Cancer	WP:WP615	3.2E-02
M5 Passage 0			
GO:MF	oxidoreductase activity	GO:0016491	1.0E-18
GO:MF	electron transfer activity	GO:0009055	7.9E-16
GO:MF	proton transmembrane transporter activity	GO:0015078	1.3E-12
GO:MF	cytochrome-c oxidase activity	GO:0004129	2.3E-07
GO:MF	NADH dehydrogenase activity	GO:0003954	3.4E-07
GO:BP	oxidative phosphorylation	GO:0006119	4.9E-29
GO:BP	cellular respiration	GO:0045333	5.1E-25
GO:BP	lipid oxidation	GO:0034440	4.0E-05
GO:BP	fatty acid oxidation	GO:0019395	2.1E-04
GO:BP	lipid catabolic process	GO:0016042	9.2E-04
GO:BP	fatty acid metabolic process	GO:0006631	1.1E-03
GO:CC	mitochondrial part	GO:0044429	1.6E-47
GO:CC	mitochondrion	GO:0005739	3.8E-37
KEGG	Oxidative phosphorylation	KEGG:00190	1.5E-31
KEGG	Citrate cycle (TCA cycle)	KEGG:00020	7.0E-05
KEGG	Fatty acid metabolism	KEGG:01212	1.6E-03
WP	Electron Transport Chain (OXPHOS system in mitochondria)	WP:WP111	2.3E-38
WP	Fatty Acid Beta Oxidation	WP:WP143	2.2E-04

894   Note: See Figure S3A for complete set of inter-media comparisons.
895   FDR, false discovery rate; GO, gene ontology; BP, biological process; MF, molecular function; CC,
896   cellular component; KEGG, Kyoto encyclopedia of genes and genomes; WP, WikiPathways.

897 Table 2. Gene ontology and pathway enrichment analysis of the P3 and P4 intra-media comparisons.

Source	Term Name	Term ID	FDR p-value
F99 Passage 3			
GO:BP	system development	GO:0048731	5.1E-06
GO:BP	animal organ development	GO:0048513	1.2E-05
GO:BP	cellular response to stress	GO:0033554	3.4E-04
GO:BP	mitotic cell cycle	GO:0000278	2.9E-03
GO:BP	response to stress	GO:0006950	3.6E-03
GO:BP	regulation of mitotic cell cycle	GO:0007346	1.4E-02
GO:BP	cell differentiation	GO:0030154	2.0E-02
GO:BP	MAPK cascade	GO:0000165	4.0E-02
GO:CC	vesicle	GO:0031982	6.3E-09
GO:CC	endomembrane system	GO:0012505	1.1E-08
KEGG	p53 signaling pathway	KEGG:04115	4.4E-04
KEGG	Focal adhesion	KEGG:04510	1.1E-02
KEGG	Apelin signaling pathway	KEGG:04371	3.3E-02
KEGG	Cellular senescence	KEGG:04218	4.8E-02
REAC	Senescence-Associated Secretory Phenotype (SASP)	REAC:R-HSA-2559582	3.0E-02
REAC	Cell Cycle, Mitotic	REAC:R-HSA-69278	4.2E-02
WP	Wnt Signaling	WP:WP428	1.1E-02
F99 Passage 4			
GO:MF	cyclin-dependent protein serine/threonine kinase inhibitor activity	GO:0004861	3.6E-03
GO:BP	nervous system development	GO:0007399	1.0E-08
GO:BP	response to oxidative stress	GO:0006979	1.2E-07
GO:BP	regulation of programmed cell death	GO:0043067	7.1E-07
GO:BP	apoptotic process	GO:0006915	1.1E-06
GO:BP	cell differentiation	GO:0030154	2.2E-06
GO:BP	mitotic cell cycle	GO:0000278	1.3E-05
GO:BP	aging	GO:0007568	2.4E-02
GO:BP	MAPK cascade	GO:0000165	4.5E-02
GO:CC	vesicle	GO:0031982	1.8E-15
GO:CC	endomembrane system	GO:0012505	3.9E-15
KEGG	Focal adhesion	KEGG:04510	1.0E-06
KEGG	p53 signaling pathway	KEGG:04115	5.8E-05
KEGG	Cellular senescence	KEGG:04218	4.2E-03
KEGG	PI3K-Akt signaling pathway	KEGG:04151	8.5E-03
REAC	Developmental Biology	REAC:R-HSA-1266738	1.5E-02
REAC	Cell Cycle, Mitotic	REAC:R-HSA-69278	3.5E-02
M5 Passage 3			
GO:MF	growth factor binding	GO:0019838	9.6E-04
GO:MF	cyclin-dependent protein serine/threonine kinase regulator activity	GO:0016538	3.2E-03
GO:BP	cell cycle	GO:0007049	1.7E-11

GO:BP	regulation of programmed cell death	GO:0043067	8.0E-07
GO:BP	cell cycle phase transition	GO:0044770	2.4E-06
GO:BP	cell differentiation	GO:0030154	2.0E-05
GO:BP	programmed cell death	GO:0012501	5.2E-05
GO:BP	MAPK cascade	GO:0000165	9.8E-05
GO:BP	response to oxidative stress	GO:0006979	2.1E-04
GO:BP	aging	GO:0007568	1.4E-02
GO:CC	vesicle	GO:0031982	2.2E-18
GO:CC	endomembrane system	GO:0012505	2.8E-08
KEGG	Cell cycle	KEGG:04110	2.9E-06
KEGG	p53 signaling pathway	KEGG:04115	1.1E-03
KEGG	Focal adhesion	KEGG:04510	1.6E-03
KEGG	MAPK signaling pathway	KEGG:04010	3.6E-03
REAC	Cell Cycle, Mitotic	REAC:R-HSA-69278	3.9E-06
M5 Passage 4			
GO:MF	cyclin-dependent protein serine/threonine kinase regulator activity	GO:0016538	3.6E-02
GO:BP	cell division	GO:0051301	2.1E-09
GO:BP	mitotic cell cycle	GO:0000278	1.8E-08
GO:BP	positive regulation of cell death	GO:0010942	4.2E-08
GO:BP	response to oxidative stress	GO:0006979	5.3E-06
GO:BP	cellular response to stress	GO:0033554	7.3E-06
GO:BP	regulation of MAPK cascade	GO:0043408	6.8E-05
GO:BP	electron transport chain	GO:0022900	3.7E-04
GO:BP	exocytosis	GO:0006887	1.4E-03
GO:BP	secretion by cell	GO:0032940	5.8E-03
GO:CC	vesicle	GO:0031982	5.7E-21
KEGG	Focal adhesion	KEGG:04510	2.6E-05
KEGG	MAPK signaling pathway	KEGG:04010	1.0E-03
KEGG	p53 signaling pathway	KEGG:04115	1.5E-03
KEGG	Oxidative phosphorylation	KEGG:00190	3.8E-02
REAC	The citric acid (TCA) cycle and respiratory electron transport	REAC:R-HSA-1428517	2.2E-04
WP	Electron Transport Chain (OXPHOS system in mitochondria)	WP:WP111	6.3E-03

898 Note: Note: See Figure S3B for complete set of intra-media comparisons.
899 FDR, false discovery rate; GO, gene ontology; BP, biological process; MF, molecular function; CC,
900 cellular component; KEGG, Kyoto encyclopedia of genes and genomes; REAC, REACTOME; WP,
901 WikiPathways.
902

903 Table 3: Activation state of Ingenuity canonical pathways in P3 and P4 intra-media comparisons.

Ingenuity Canonical Pathways	z-score
F99 Passage 3	
NRF2-mediated Oxidative Stress Response	3.36
IL-8 Signaling	2.89
Apelin Endothelial Signaling Pathway	2.71
Cell Cycle: G2/M DNA Damage Checkpoint Regulation	2.65
Chemokine Signaling	2.33
IL-3 Signaling	2.33
ERK/MAPK Signaling	1.90
VEGF Family Ligand-Receptor Interactions	1.89
IL-6 Signaling	1.89
IGF-1 Signaling	1.67
Acute Phase Response Signaling	1.67
JAK/Stat Signaling	1.41
p53 Signaling	1.13
FGF Signaling	1.00
Estrogen-mediated S-phase Entry	-1.00
Cyclins and Cell Cycle Regulation	-1.13
RhoGDI Signaling	-1.41
LXR/RXR Activation	-2.24
F99 Passage 4	
Cell Cycle: G2/M DNA Damage Checkpoint Regulation	3.46
NRF2-mediated Oxidative Stress Response	3.27
IL-8 Signaling	3.09
Chemokine Signaling	2.89
p53 Signaling	2.83
UVC-Induced MAPK Signaling	2.33
IGF-1 Signaling	1.41
Inflammasome pathway	1.34
Apelin Endothelial Signaling Pathway	1.28
IL-6 Signaling	1.21
IL-1 Signaling	1.13
HGF Signaling	1.00
JAK/Stat Signaling	1.00
Estrogen-mediated S-phase Entry	-1.13
Chondroitin Sulfate Degradation (Metazoa)	-1.34
Pyrimidine Deoxyribonucleotides De Novo Biosynthesis I	-1.34
NER Pathway	-1.90
LXR/RXR Activation	-3.00
M5 Passage 3	
NRF2-mediated Oxidative Stress Response	3.27
Cell Cycle: G2/M DNA Damage Checkpoint Regulation	3.16

Chemokine Signaling	2.71
IL-8 Signaling	2.69
Acute Phase Response Signaling	2.50
IL-6 Signaling	2.32
TGF-β Signaling	2.11
Inflammasome pathway	2.00
IL-1 Signaling	1.89
JAK/Stat Signaling	1.51
p53 Signaling	1.39
p38 MAPK Signaling	1.00
Glycolysis I	-1.00
Fatty Acid β-oxidation I	-1.00
Estrogen Biosynthesis	-1.00
Pyrimidine Deoxyribonucleotides De Novo Biosynthesis I	-2.00
NER Pathway	-2.33
LXR/RXR Activation	-3.00
M5 Passage 4	
NRF2-mediated Oxidative Stress Response	4.60
IL-8 Signaling	3.51
Chemokine Signaling	3.00
IL-6 Signaling	2.84
Cell Cycle: G2/M DNA Damage Checkpoint Regulation	2.67
Inflammasome pathway	2.24
p53 Signaling	2.07
IL-1 Signaling	1.90
Acute Phase Response Signaling	1.88
p38 MAPK Signaling	1.70
JAK/Stat Signaling	1.41
IGF-1 Signaling	1.34
Fatty Acid β-oxidation I	-1.63
Acetyl-CoA Biosynthesis I (Pyruvate Dehydrogenase Complex)	-2.00
Cholesterol Biosynthesis I	-2.00
Superpathway of Cholesterol Biosynthesis	-2.45
TCA Cycle II (Eukaryotic)	-2.65
Oxidative Phosphorylation	-5.11

904 Notes: 1) See Figure S3B for complete set of intra-media comparisons. 2) A positive z-score denotes
905 activation and a negative z-score denotes deactivation.
906

907 Table 4: Activation state of cell cycle functions in P0 and P4 intra-media comparisons.

Function Annotation	z-score
F99 Passage 3	
Senescence of cells	0.99
Replicative senescence of cells	0.60
Cell cycle progression	-0.86
Re-entry into interphase	-1.34
Mitosis	-1.72
Initiation of interphase	-1.73
F99 Passage 4	
Senescence of tumor cell lines	1.51
G1 phase	1.35
Senescence of cells	0.69
M phase	-0.53
M phase of cervical cancer cell lines	-0.79
M phase of tumor cell lines	-1.30
M5 Passage 3	
Senescence of cells	1.14
G1 phase	0.90
Cell cycle progression	-0.81
M phase	-1.19
Mitosis	-1.69
Segregation of chromosomes	-2.13
M5 Passage 4	
G1 phase	2.24
Senescence of cells	1.70
Cell cycle progression	-0.88
Cytokinesis	-1.06
M phase	-1.18
Segregation of chromosomes	-1.69

908 Notes: 1) See Figure S3B for complete set of intra-media comparisons. 2) A positive z-score denotes
909 activation and a negative z-score denotes deactivation.

910   **SUPPLEMENTARY TABLES**
911
Table S1. Donor cornea information

Table S2. Quantitative PCR primers.

Table S3. Antibodies.

Table S4. Differential gene expression lists for inter-media comparisons.

Table S5. Full results of gene ontology and pathway enrichment in inter-media data sets.

Table S6. Expression of cell senescence-associated genes in primary CEnC.

Table S7. Differential gene expression lists for intra-media comparisons.

Table S8. Full results of gene ontology and pathway enrichment in intra-media data sets.

Table S9. Full results of activation of Ingenuity canonical pathway in intra-media data sets.

Table S10. Full results of activation of Ingenuity cell cycle functions in intra-media data sets.