Assigning metabolic rate measurements to torpor and euthermia in heterothermic endotherms: "torpor", a new package for R.

Nicolas J. Fasel[1], Colin Vullioud[2], Michel Genoud[1]

1. Department of Ecology and Evolution, University of Lausanne, CH–1015 Lausanne,
Switzerland.
2. Department Evolutionary Genetics, Leibniz Institute for Zoo and Wildlife Research, D-10315
Berlin, Germany.

Corresponding author:
Nicolas J. Fasel
Department of Ecology and Evolution
University of Lausanne
CH–1015 Lausanne
Switzerland
fasel.nicolas@gmail.com

Running head: A R-package for thermal biology
Key words: Energetics, metabolism, Bayesian, Mixture models, Energy saving

Abstract

1.   Torpor is a state of controlled reduction of metabolic rate (M) and body temperature in endotherms. Assigning measurements of M to torpor or euthermia can be challenging. All current techniques available to distinguish between those two states have their limitations and are at least partly arbitrary.

2.   Here, we firstly propose a novel method for distinguishing torpor from euthermia. We use the variation in M measured during euthermic rest and torpor at different ambient temperatures (T_a) to estimate the lower critical temperature (T_{lc}) of the thermoneutral zone (TNZ) and determine physiological state membership using a mixture model. In addition, this method allows the prediction of M during euthermic rest and torpor along T_a, including resting M within the TNZ. Secondly, we test this method on previously published datasets (N=28), covering a diversity of metabolic and thermal strategies.

3.   The method's results match the original assignment with very high accuracy (mean=0.95). Most of the conflicts concerned M measured around T_{lc}.

4.   Our method provides an objective approach to estimate T_{lc} and assign measurements of M to euthermia or torpor. Our method further enables the prediction of M during euthermic rest and torpor at any given T_a below the upper limit of the TNZ. The present method is applicable to species conforming to the "Scholander-Irving model" and its later extensions to include torpor. This model has proven to be an appropriate description of the variation of M along T_a in most endotherms studied to date.

5.   Our method aims ultimately at improving the standardization of respirometry analyses in heterotherms. Finally, the development of the associated R-package (torpor), should spread and ease the use of this method.

Introduction

Torpor is a state of controlled reduction of metabolic rate and body temperature observed in

numerous mammals and birds (Ruf & Geiser, 2015). This physiological state tends to be either

shallow and short (< 24h, daily torpor) or deep and long-lasting (> 24h, hibernation)

(Schleucher & Withers, 2001; McKechnie & Lovegrove, 2002; Ruf & Geiser, 2015). However, a

continuity between these types of torpor as well as between torpor and euthermic rest has

been suggested (McKechnie & Lovegrove, 2002; Boyles et al., 2013; van Breukelen & Martin,

2015). Torpor has profound implications on energy use and allocation (Lyman, Willis, Malan,

& Wang, 1982) and potentially affects many biological functions (Geiser & Brigham, 2012;

Nowack, Stawski, & Geiser, 2017). Accordingly, a vast literature describes the occurrence of

torpor and its associated energy savings (Geiser & Ruf, 1995; Lovegrove, 2012; Ruf & Geiser,

2015; Nowack, Levesque, Reher, & Dausmann, 2020).

Assigning measurements of metabolic rate (M) or body temperature (T_b) to torpor or

euthermic rest can be challenging. The distinction between the two states is usually

straightforward in typical hibernators, which undergo deep torpor bouts at low ambient

temperature (T_a). But even in such species, that distinction can become problematic at

ambient temperatures slightly below the thermoneutral zone (TNZ), i.e. the range of T_a where

temperature regulation can be achieved without regulatory changes in metabolic heat

production or evaporative heat loss (IUPS Thermal Commission, 2003). Indeed, at T_a slightly

below the TNZ, the differences in state-specific average M and T_b are small compared to the

state-specific variability in each of these variables. For the same reason additional difficulties

can be encountered in species whose minimum T_b in torpor lies only slightly below the

euthermic T_b (McNab, 1980a; Genoud, Bonaccorso, & Anends, 1990; Bonaccorso & McNab,

1997; Bartels, Law, & Geiser, 1998) and in species exhibiting a large variability in metabolic

rate and body temperature at rest (Geiser, Coburn, Kourtner, & Law, 1996; Coburn & Geiser,

1998).

Barclay, Lausen, & Hollis (2001) reviewed the variety of criteria that have been chosen in the

past to distinguish torpor from euthermia. Apart from a minority of reports that identified

torpor on the basis of behavioural features (Geiser & Kenagy, 1988; Geiser & Masters, 1994;

Brice, Grigg, Beard, & Donovan, 2002), the vast majority of studies used patterns of variation

in body (or skin) temperature and/or rate of metabolism to separate the two states in the field

or in the laboratory. Animals were usually deemed to be in torpor below a threshold T_b or M

(Geiser et al., 1996; Coburn & Geiser, 1998; Hosken & Withers, 1999; Mzilikazi & Lovegrove,

2002; Kelm & von Helversen, 2007; Levesque, 2008), below some threshold temperature

differential between the body and air (Levesque & Lovegrove, 2014), or below a threshold

percentage of the euthermic rate of metabolism (Hudson & Scott, 1979; Geiser, 1988a). The

diagnostic threshold value was sometimes calculated on the basis of the parameter's variation

in the euthermic state (e.g. Lovegrove & Raman, 1998; McKechnie, Ashdown, Christian, &

Brigham, 2007). The diagnostic threshold can also be predicted on the basis of body mass and

T_a (Willis, 2007). Despite the efforts made to render them more objective, all available

techniques have their limitations and are at least partly arbitrary (Barclay et al., 2001; Boyles,

Smit, & McKechnie, 2011).

With this paper, we firstly propose an automatized method for distinguishing torpor from

euthermic rest in species undergoing daily or seasonal torpor, hereafter referred to as

heterotherms following the terminology of Geiser & Ruf (1995) or Lovegrove (2012).

Assignment to an either euthermic or torpid state is based on the variation observed among

measurements of M. We assume that the relationship between M of resting animals and T_a

follows the classical "Scholander-Irving model" (Scholander, Hock, Walters, Johnson, & Irving,

1950; for a discussion see McNab, 2002), which was later extended to include torpor

(Hainsworth & Wolf, 1970; Humphries, Thomas, & Speakman, 2002; Speakman & Thomas,

2003; Geiser, 2011, Fig. 1). That model predicts the vast majority of patterns observed among

endotherms entering daily torpor or hibernation (but see discussion). We explicitly consider

that metabolic inhibition may occur during torpor (Geiser, 1988b, 2004; Geiser & Kenagy,

1988; Withers, Cooper, Maloney, Bozinovic, & Cruz-Neto, 2016), by allowing the curve of

torpor M to reach the lower limit of the thermoneutral zone (or lower critical temperature,

T_{lc}) at a level equal to – or lower than the resting metabolic rate within the TNZ (M_{TNZ}). Our

method is specifically intended to facilitate the discrimination between torpor and euthermia

in laboratory experiments using respirometry, although it can also be useful to model T_{lc}, M_{TNZ},

the minimum rate of metabolism in torpor (TMR), as well as T_t, the threshold T_a below which

a minimum T_b in torpor is maintained by varying M. Secondly, we assess the performance of

our method, by applying it to previously published data on the metabolic rate of various

mammals. Finally, we provide a new package running under R (R Development Core Team,

2012) allowing researchers to apply the present method to their own data. This package

("torpor") comprises several useful functions that will improve standardization of the analyses

of metabolic measures for thermal biology.

Material and Methods

The three-steps Method

Our present method is based on three steps and requires measurements of the metabolic rate

(M) of resting or torpid animals at ambient temperatures (T_a) ranging from below – to within

the TNZ. Animals are assumed to be in one of three "states": euthermic rest below the TNZ,

torpor below the TNZ or rest within the TNZ. We do not consider here the relation between

M and T_a above the TNZ. Consequently, metabolic rates measured at T_a higher than the upper

limit of the TNZ should be excluded. During the different steps, a mixture model based on

Bayesian inference is run under varying conditions.

The model

The Scholander-Irving model and its extensions consider that resting M measured within the

TNZ is independent of T_a. This rate is hereafter referred to as M_{TNZ}, although it corresponds to

the basal metabolic rate (BMR) provided that the specific criteria for the BMR are met (see

discussion). Below T_{lc}, the M of euthermic animals (M_e) increases linearly with decreasing T_a

(Fig. 1):

$M_e \sim \alpha_e + \beta_e * T_a \mid T_a < T_{lc}$

The model assumes that M_e equals M_{TNZ} at T_{lc} and therefore:

$\alpha_e = M_{TNZ} - \beta_e * T_{lc}$

Where:

$\beta_e = \dfrac{M_{TNZ}}{T_{lc} - T_{be}}$

T_{be} represents the hypothetical T_a where M_e equals 0. It would correspond to the euthermic

body temperature, provided that thermal conductance and body temperature do not vary

with T_a below the TNZ (McNab, 1980b).

Below some threshold ambient temperature (T_t), the metabolic rate in torpor (M_t) increases

linearly with decreasing T_a to maintain a minimal body temperature (Fig. 1). This state is

usually referred to as "regulated torpor":

$M_t \sim \alpha_r + \beta_r * T_a \mid T_a < T_t$

Where:

$T_t = \dfrac{(TMR - \alpha_r)}{\beta_r}$

and

$\alpha_r = -T_{bt} * \beta_r$

TMR represents the minimal M_t at T_t. T_{bt} is the hypothetical T_a at which M_t of the regulated

torpor would equal 0. T_{bt} would correspond to the minimal body temperature in torpor

provided that thermal conductance and body temperature in regulated torpor do not vary

with T_a below T_t. Between T_t and T_{lc}, M_t follows an exponential curve (Fig. 1). In this T_a range,

torpor is referred to as "conforming torpor".

$M_t \sim \alpha_c * \exp(\beta_c * T_a) \mid T_t \leq T_a \leq T_{lc}$

Where:

$\beta_c = \dfrac{\log(M_r) - \log(TMR)}{T_{lc} - T_t}$

and

$\alpha_c = \dfrac{M_r}{\exp(\beta_c * T_{lc})}$

M_r represents M_t at T_{lc}. In accordance with the general trend observed (Geiser, 2004), we

assume that

$\beta_r = \beta_e$

The following parameters are modelled: the fractions of measurements belonging to each

state, which provides the value specific state membership probabilities for each measurement

(see Second step: Metabolic rate (M) measurements pre-assignment, below), T_{be}, T_{bt}, TMR,

and the standard deviations for M_{TNZ} and regulated/conforming metabolic rates (SD_{TNZ}, SD_r

and SD_c).

First step: Estimation of M_{TNZ} and T_{lc}

The method initially defines T_{lc_low} as the highest possible T_a within the data set that still

underestimates T_{lc}. Above T_{lc_low}, a linear regression between M and T_a should neither result

in a significant negative slope nor be affected by heteroscedasticity. Beginning with the ten M

values measured at the highest T_a, linear regressions are performed on sets of M values

progressively including values at lower T_a. The inclusion of additional M values obtained below

T_{lc} eventually leads to a significantly negative slope in the case of animals not entering torpor

and/or to heteroscedasticity in the case of heterotherms. A significantly positive slope would

reveal an absence or insufficient number of values for euthermic rest below the TNZ, hence it

automatically stops the analysis as T_{lc_low} is undefinable. Heteroscedasticity is assessed with a

Breusch-Pagan test with the function "bptest" from package "lmtest" (Zeileis & Hothorn,

2002). Significance level for the Breusch-Pagan test is set at 0.05. The significance of the

negative regression is assessed with a linear regression, function "lm" and significance level is

set at 0.01.

Then, in order to get T_{lc}, the model described in the second step (see Second step: Metabolic

rate (M) measurements pre-assignment) is first run without data points measured above

T_{lc_low}. Moreover, within this step, the parameter T_{lc} is as well modelled. M_{TNZ} provided in that

analysis is the mean of the M measured at T_a higher than T_{lc_low}. Once T_{lc} has been estimated,

M_{TNZ} is recalculated as the mean of the M measured at T_a higher than T_{lc}.

Second step: Metabolic rate (M) measurements pre-assignment

The assignment of M measurements to one of the three physiological states is based on a

probabilistic process. Below T_{lc}, a posterior categorical distribution is generated for each

coupled data (T_a and M), providing the value specific state membership probabilities to belong

to each of the possible states (torpor or euthermic rest). Independent of the pre-assignment,

M values higher than the predicted M_e or lower than the predicted M_t are automatically

assigned to M_e and M_t, respectively. This latter automatic procedure is, however, only

performed if at least one M measurement has been assigned to torpor during that second

step. As defined previously, M values located above T_{lc} are automatically assigned to M_{TNZ}.

Automatically assigned values get a state membership probability of one.

Third step: Metabolic rate (M) measurements final assignment and estimation

of the functions parameters

In the third step, only M values situated between the predicted M_e and M_t are assigned.

Measurements' states that are automatically assigned during the second step are provided

along with the coupled M and T_a values. During that final step, the parameters of the functions

relating M to T_a below the T_{lc}, as well as the standard deviation for the different physiological

states are modelled.

Bayesian parametrization

Prior distributions are defined either uninformatively or with biologically relevant limits.

Except when specified, all prior distributions are Gaussian with a mean of 0 and a precision

(i.e. $1/SD^2$) of 0.001 and are truncated based on the specified limits. Specifically, the prior

distribution of T_{lc} during the first step is constrained between T_{lc_low} and the maximal T_a

recorded within the dataset. T_{be} prior distribution is constrained between T_{lc} (T_{lc_low} for the

first step) and 50°C. The upper limit of the T_{bt} prior distribution satisfies two conditions. First,

T_t should be inferior to the lower 95% credible interval (CI) of T_{lc}. Second, the ratio of

conforming M_t values corresponding to body temperatures differing by 10°C (Q_{10}) should not

exceed a value of 5 (Geiser, 1988b). Those conditions are verified by fixing the upper range of

the prior distribution of T_{bt} as:

$$\min \left(\frac{\text{lower 95CI } T_{lc} * \beta_t - TMR}{\beta_t} , T_{be} - \frac{M_{TNZ} * 2}{TMR} \right)$$

The lower limit of the T_{bt} prior distribution is set at -5°C (Barnes, 1989). The upper limit of the

prior distribution of TMR is defined as 80% of the M_{TNZ} (Ruf & Geiser, 2015), while its lower

one is fixed at 0. For M_r, the prior distribution ranges from TMR to M_{TNZ}. An uninformative

Dirichlet distribution (i.e. two concentration values of 1) is used for the priors of the data set

specific state membership fractions, which provide the value specific state membership

probabilities of each measurement (see Second step: Metabolic rate (M) measurements pre-

assignment, above). Finally, the prior distributions of the standard deviations for M_{TNZ} and

regulated/conforming metabolic rates (SD_{TNZ}, SD_r and SD_c) are uniform, with that of SD_r

constrained between 0 and 3, that of SD_c constrained between one fifth of the value of SD_r

and SD_r and that of SD_{TNZ} constrained between half the value of SD_r and SD_r.

Three different Markov chains are run during 50'000 iterations starting at initial values within

the range of parameter space. The initial convergence phase is excluded by dropping the first

30'000 iterations. Markov chains are thinned by a factor of 10 and the Brooks–Gelman–Rubin

criterion $\hat{R}$ (Brooks & Gelman, 1998) is used to assess the convergence of chains (< 1.1). To

ease parameters' estimation, metabolic rate measurements are divided by their mean.

Identifiability of the parameters

The prior and posterior distribution overlap (PPO) can enable the evaluation of the modelled

some parameters' relevance. Overlap values are obtained with the function "MCMCtrace"

from package "MCMCvis" (Youngflesh, 2018). Truncated Gaussian prior distributions are

provided with the function "rtruncnorm" from package "truncnorm" (Mersmann, Trautmann,

Steuer, & Bornkamp, 2018).

Assignment confidence

The M value is assigned to the physiological state having the highest specific state membership

probability. The assignment confidence represents the product of the highest specific state

membership probability and of the probability that T_a is above (for M_{TNZ}) or below (for M_t and

M_e) T_{lc}. That latter probability is calculated from the variation in the estimation of T_{lc} (cf. First

step: Estimation of M_{TNZ} and T_{lc}), as the value at T_a of the T_{lc} cumulative posterior distribution

function, or 1 minus that function respectively. A threshold proportion is then selected to

consider as invalid any assignment with a significantly lower assignment confidence. That

hypothesis is tested with a binomial test, function: "binom.test", with the alternative

hypothesis set as "greater". The significance level for the binomial test is set at 0.05. For the

present study, assignments with a confidence higher than 0.8 are considered as valid.

Data collection and method evaluation

We evaluate our method by applying it on 28 published sets of metabolic data obtained by

respirometry on 26 species of small or medium-sized mammals at rest or in torpor (Table 1).

This assemblage of species covers a diversity of metabolic and thermal strategies along the

continuum from permanent homeothermy to heterothermy including shallow, daily torpor

and deep, long-term hibernation. For each published set, we consider all metabolic data

provided, except those made above the upper critical temperature defined by the authors

(see "the three-steps Method" above). In 20 species (here considered as "heterotherms",

Table 1), rates of metabolism measured at T_a's below the described thermoneutral zone were

originally assigned to either of two states, which were usually referred to as euthermia (or

normothermia) and torpor. In *Cynopterus brachyotis*, values corresponding to particularly low

T_b were distinguished but not explicitly referred to as torpor (McNab, 1989). In *Lasiurus*

seminolus, some of the values measured during torpor were originally distinguished as the

bats displayed "irregular fluctuations" in rate of metabolism (Genoud, 1993). For simplicity,

we hereafter treat these values as if the authors assigned them to torpor. The remaining six

species are hereafter referred to as "homeotherms" (Table 1), as none of the values in the

corresponding data sets were assigned to torpor by the original authors. Chosen studies

reported M_{TNZ} (mostly referred to as *BMR*) and T_{lc} estimates, although the latter had to be

extracted from a graph in two cases and calculated as the intercept between the euthermic

line below thermal neutrality and M_{TNZ} provided by the authors in another case (Table 1). They

all provided a sufficiently precise graph of metabolic rate values as a function of T_a and several

of them also estimated T_t. Plotted data were digitized using the software Plot Digitizer

(Huwaldt & Steinhorst, 2015). Sample sizes of the digitized data range between 27 and 204

(Table 1). We test the correlation between the M_{TNZ}, T_{lc} and T_t values modelled with the

present method and those provided by the authors with Pearson's paired-samples correlation

test, function "cor.test". Cases in which the estimation of T_t falls outside the range of

measured T_a are excluded from this analysis as T_t is considered unidentifiable. Then, we

examine whether the assignment of the metabolic values to torpor or euthermia made by the

authors and provided by the method coincides, and we calculate a corroboration index as the

fraction of the matched assignments. Only measures with valid assignments are considered

for the correlation tests and the corroboration index calculations. Finally, in order to define a

PPO range highlighting an identifiable parameter, the PPOs of *TMR* from studies where *M*

measurements are assigned to torpor are compared with those from studies where no *M*

measurement is assigned to torpor and *TMR* is consequently unidentifiable.

Results

In studies where no M is assigned to torpor (i.e. "homeotherms"), PPOs of TMR range from

76.80 to 95.60%. In identified heterotherms, PPOs of TMR values range from 4.60 to 64.60%

(Table 1).

Values of T_{lc}, M_{TNZ} and T_t modelled by the model and values obtained from the original studies

are significantly correlated (T_{lc}: N=28, Pearson's coefficient: 0.65, p<0.001, M_{TNZ}: N=28,

Pearson's coefficient: 1.00, p<0.001, T_t: N=14, Pearson's coefficient: 0.97, p<0.001).

Over all data sets, the proportions of valid assignments (i.e. assignment confidence > 0.8)

range from 0.63 to 1.00 (median=0.87, Table 1). The corroboration indexes range between

0.66 and 1.00 (median=1.00, Table 1). Complete matches between the method assignments

and the authors' descriptions are found for 15 data sets (e.g. *Nyctophilus geoffroyi* (Hosken &

Withers, 1999), Fig. 2A). Mismatches mostly occur close to T_{lc} but can also be found as far as

41.0°C from T_{lc} (Fig. 3). Among homeotherms, no M measurement is assigned to torpor (e.g.

Sorex minutus (Sparti & Genoud, 1989), Fig. 2B).

343
344

Discussion

The present method aims at standardizing the assignment of M measurements to physiological states and at estimating the parameters' functions which link M to T_a. The use of published data sets enable us to assess our method's performance, although it is not our goal to challenge the original analyses. Authors of the published data sets may have benefited from additional information (e.g. body temperature or behaviour) to assign M to the various physiological states, while our method is based on statistics exploiting only T_a and M.

Our Bayesian inference-based method models all parameters whatever the available data. It remains therefore crucial to consider the parameters' identifiability. The prior and posterior distribution overlaps (PPO) of T_{bt} and T_{lc} are not reliable for that purpose, because the prior distribution of those parameters largely depends on the data provided. Considering only the PPO of TMR, we can define that PPO higher than 75% clearly reveals that the parameter is not identifiable.

The parameters estimated by our method correlate significantly with those extracted from the original studies; T_{lc} values modelled with our method differ appreciably from those provided originally, while remaining within an acceptable range (Table 1). Our estimation of T_{lc} requires numerous M values of euthermic rest to precisely model the T_a at which M_e reaches M_{TNZ}. It is possible that the amount of M_e values is sometimes not sufficient to model T_{lc} adequately. Additionally, a statistical estimation of T_{lc} (e.g. Song, Körtner, & Geiser, 1995, 1997; Bonaccorso & McNab, 1997; Genoud, Martin, & Glaser, 1997; Hosken & Withers, 1999; Willis, Lane, Liknes, Swanson, & Brigham, 2005; Genoud, 2014) was hitherto often lacking,

leaving potentially space for some misleading subjectivity on the part of the original authors.

Finally, an overestimation of T_{lc} by our method remains possible if an appreciable metabolic

inhibition occurs during torpor (see below). The difference in the estimations of T_t is

consistently smaller than that of the estimations of T_{lc} (Table 1). This could be linked to the

smaller variation in M exhibited during steady conforming torpor in comparison to that

measured during euthermic rest, hence to the higher precision in the estimation of T_t. Finally,

it is not surprising that the M_{TNZ} obtained by our method and by the original authors are close

to identical (Table 1), since both are largely based on the same M values.

The assignments made by our method and by the original authors agree well, as illustrated by

the corroboration index (≥ 0.9 for 22 out of 28 data sets, Table 1). Many of the conflicting

assignments in the region of T_{lc} are a mere consequence of the difference between the

estimated T_{lc} and that of authors (Table 1, Figs. 2C & 3). In particular, differences in modelled

T_{lc} explain all homeotherm species mismatches, as no M value is assigned to torpor for these

species. The remaining mismatches can generally be explained by intra-state frequency

distributions of M that do not segregate clearly. This condition is often found close to the

thermoneutral zone, but also occurred in species where T_t is relatively close to T_{lc} (e.g. McNab,

1980a, 1989). Such discrepancies illustrate the difficulty of the assignment process when

torpor and euthermia have to be distinguished solely on the basis of measurements of M. In

this study, we consider assignments with a confidence lower than 0.8 as invalid. Obviously, an

increase of this threshold value would lead to fewer assignments' mismatches. While the use

of a statistical method enables an objective assignment, it is nevertheless worthwhile to recall

that some authors also used patterns of T_b to assign their data (e.g. McNab, 1980a, 1989).

These specific cases should deserve further attention as they can highlight mechanisms

decoupling M and T_b control (e.g. Daniels, 1984; McNab, 1988; Heldmaier, Ortmann, & Elvert,

2004).

The present method furthermore aims at providing the parameters of the functions relating

M in resting and torpid animals to T_a. These functions enable the estimation of TMR and T_t.

Predicting M at any T_a is crucial to model energy costs related to torpor expression (e.g. Boyles,

Johnson, Blomberg, & Lilley, 2020). Another central aspect lies in that our method allows the

M function for torpor to be lower than M_{TNZ} at T_{lc}. We thus explicitly admit that metabolic

inhibition could occur in some species during torpor (Geiser, 2004; Withers et al., 2016).

Caution should be kept in the interpretation of a difference between the estimated M_r and

M_{TNZ}, as the latter will often overestimate metabolic inhibition in torpor. Indeed, insufficient

amount of M_t values corresponding to conforming torpor leads to the unidentifiability of M_r

(e.g. *Melonycteris melanops* (Bonaccorso & McNab, 1997), Fig. 2D). Consequently, metabolic

inhibition should only be considered when sufficient M_t values corresponding to the

conforming torpor are provided and M_r PPO indicates adequate parameter's identifiability.

Considering only identifiable parameters (i.e. Mr PPO < 75%), we quantify metabolic inhibition

as M_r over M_{TNZ}. Values range between 0.27 and 0.93 (N=18, median=0.77, Table 1). By

extrapolation, we should consider the possibility that a hypometabolic state might occur

within the TNZ as well (e.g. Grimpo, Legler, Heldmaier, & Exner, 2013; Reher, Ehlers,

Rabarison, & Dausmann, 2018). With our method, we have not implemented an assignment

to a state of torpor within the thermoneutral zone, hence M_{TNZ} corresponds to the average of

all values from the TNZ. Our method might underestimate the eumetabolic M_{TNZ} if animals

occasionally enter a hypometabolic state in the TNZ, and might consequently overestimate

T_{lc}. A predicted M_t at T_{lc} much lower than M_{TNZ} should be viewed as an indication for a strong

metabolic inhibition occurring during torpor and should invite for a careful consideration of

the estimated M_{TNZ} and T_{lc}.

While the performance of our method appears to be excellent with the selected data sets, its

main limitations correspond to those of the model on which it is based; the Scholander-Irving

model (Scholander et al., 1950) and its latter extensions (Hainsworth & Wolf, 1970; Humphries

et al., 2002; Speakman & Thomas, 2003; Geiser, 2011). Although extremely fruitful, this model

cannot be considered as more than a descriptive simplification of the standard energetics of

mammals and birds (McNab, 2002). It surely does not describe the thermal biology of all

heterothermic species. Causes for significant divergence from the hereby applied model (Fig.

1) include 1) a non-linear relationship between M_e and T_a (e.g. *Cercartetus nanus*: Song,

Körtner, & Geiser, 1997) that might be due to a variation of thermal conductance and/or T_b as

a function of T_a below thermal neutrality (McNab, 1980b); 2) a strong difference in the slopes

of the regressions of M vs T_a in regulated torpor and euthermic rest (Geiser, 2004); and 3) a

strong dependence of M_t on factors other than T_a, such as body mass (Kelm & von Helversen,

2007) and duration since the last activity period or since last meal (Grigg, Augee, & Beard,

1992; Morris, Curtin, & Thompson, 1994). This model also does not describe adequately

circadian (e.g. the shallow "rest-phase hypothermia" of many small birds (McKechnie &

Lovegrove, 2002) or ultradian (Heldmaier, Steinlechner, Ruf, Wiesinger, & Klingenspor, 1989)

variations in M. Finally, it must be recalled that our method constrains the minimal rate of

metabolism in torpor (*TMR*) between 0 and 80% of M_{TNZ} (Table 1). This constraint would

exclude heterotherms undergoing extremely shallow torpor with a *TMR* above 80% of M_{TNZ}.

Nevertheless, it is noteworthy that all heterotherms that were reviewed by Ruf & Geiser

(2015) had a lower (and usually much lower) relative TMR (4.4 – 66% in birds; 1 - 67% in

mammals). The above limitations should be kept in mind for a lucid application.

In the selected data sets, the metabolic values reported usually correspond to minimal and

stable rates for a given physiological state. Similarly, our method is designed for an analysis of

sets of stable M values. Transitions between activity and rest or between euthermic rest and

torpor (i.e. entrance into and arousal from torpor) as well as incomplete torpor bouts during

which M never reaches a stable, minimal level (e.g. "test drops" (Hudson & Scott, 1979); see

also (Lyman, 1982; Genoud, 1993)) should be excluded. It should also be recalled that the

duration of the runs can affect estimations of M (Cooper & Withers, 2009), and that long runs

may be necessary for some heterothermic animals to achieve a state of steady torpor (e.g.

Heldmaier, Ortmann, & Elvert, 2004). Our method is able to reveal the presence or the

absence of distinct groups of values (euthermic rest vs torpor) within a given data set. It is

clearly not intended to judge whether any of the data points entered correspond to stable

rates and/or to minimal values. Furthermore, throughout this paper, resting metabolic rate

within the TNZ is referred to as M_{TNZ}, rather than to the basal rate of metabolism (BMR).

Estimating the BMR requires several specific criteria: animals should be post-absorptive, adult,

non-reproductive, and resting during a major inactive phase of the daily cycle (McNab, 1997).

If these criteria are not met, the method still remains applicable, but the resting M estimated

within the TNZ (i.e. M_{TNZ}) will correspond to a minimal RMR (resting metabolic rate) rather

than to the BMR.

Then, individual variation is not explicitly considered in our method. Especially, body mass

variation within a studied population might affect the assignment of M to torpor or euthermia

in a complex manner. Indeed, *BMR* (thus also M_{TNZ}) and thermal conductance depend on body

mass in an allometric manner at an intraspecific level (Nespolo, Arim, & Bozinovic, 2003;

Glazier, 2005; Genoud, 2014; Sassi & Novillo, 2015), as they do at an interspecific level (e.g.

reviewed by Schleucher & Withers, 2001; Genoud, Isler, & Martin, 2018). Moreover, T_t and

TMR might also be affected by body mass within species, as they too are known to scale to

body mass at an interspecific level (Ruf & Geiser, 2015). However, there would be no particular

difficulty to apply our method to single individuals, other than having enough measurements

for each of them. The method could also be applied repeatedly on single individuals, in order

to account for intra-individual variation.

Despite its limitations, our method and the associated R-package (torpor) provide a way to

standardize the analysis of respirometry data in relation to T_a. Its major strength is that it uses

a probabilistic approach to assign metabolic values to torpor or euthermia. Rather than being

assigned on the basis of a particular threshold value, the method calculates for each data point

the probability of belonging to one state or another. Data points that might not be assignable

are clearly identified. Thus, the partly arbitrary nature of the assignment process is removed.

Furthermore, the method also estimates the parameters' functions which link M to T_a and

several parameters describing the standard energetics of the animal(s) under study, including

M_{TNZ}, *TMR*, T_{lc} and T_t. It can be applied to study intraspecific as well as interspecific variation.

Parameters of standard energetics might now be extracted from the literature for

comparative analysis, largely avoiding causes of variation due to the diversity of past

assignment techniques. Ultimately, this method and the associated R-package (torpor) will

ease intra- and inter-specific comparative analyses of endotherm energetics.

Acknowledgements

We are thankful to Philippe Christe and Nicolas Salamin for their support during the

elaboration of the model and to Martyn Plummer who gave us some ideas at the beginning of

this project. Finally, we thank two anonymous reviewers and Dr. Aaron Ellison (Editor) for their

constructive comments, which helped us to improve the manuscript. This study was financially

supported by the Department of Ecology and Evolution of the University of Lausanne

(Switzerland) and by the Swiss National Science Foundation (grant number: P2BEP3_168709

to NJF)

References

Barclay, R. M. R., Lausen, C. L., & Hollis, L. (2001). What's hot and what's not: defining torpor in free-ranging birds and mammals. *Canadian*

Journal of Zoology, *79*(10), 1885–1890. doi:10.1139/cjz-79-10-1885

Barnes, B. M. (1989). Freeze avoidance in a mammal: Body temperatures below 0°C in an arctic hibernator. *Science*, *244*(4912), 1593–1595.

doi:10.1126/science.2740905

Bartels, W., Law, B. S., & Geiser, F. (1998). Daily torpor and energetics in a tropical mammal, the northern blossom-bat *Macroglossus minimus*

(Megachiroptera). *Journal of Comparative Physiology - B Biochemical, Systemic, and Environmental Physiology*, *168*(3), 233–239.

doi:10.1007/s003600050141

Bonaccorso, F. J., & McNab, B. K. (1997). Plasticity of Energetics in Blossom Bats (Pteropodidae): Impact on Distribution. *Journal of*

Mammalogy, *78*(4), 1073–1088. doi:10.2307/1383050

Boyles, J. G., Johnson, J. S., Blomberg, A., & Lilley, T. M. (2020). Optimal hibernation theory. *Mammal Review*, *50*(1), 91–100.

doi:10.1111/mam.12181

Boyles, J. G., Smit, B., & McKechnie, A. E. (2011). Does use of the torpor cut-off method to analyze variation in body temperature cause more

problems than it solves? *Journal of Thermal Biology*, *36*(7), 373–375. doi:10.1016/j.jtherbio.2011.07.007

Boyles, J. G., Thompson, A. B., Mckechnie, A. E., Malan, E., Humphries, M. M., & Careau, V. (2013). A global heterothermic continuum in

mammals. *Global Ecology and Biogeography*, *22*(9), 1029–1039. doi:10.1111/geb.12077

Brice, P. H., Grigg, G. C., Beard, L. A., & Donovan, J. A. (2002). Patterns of activity and inactivity in echidnas (*Tachyglossus aculeatus*) free-

ranging in a hot dry climate: Correlates with ambient temperature, time of day and season. *Australian Journal of Zoology*, *50*(5), 461–475.

doi:10.1071/ZO01080

Brooks, S. P., & Gelman, A. (1998). General methods for monitoring convergence of iterative simulations. *Journal of Computational and*

Graphical Statistics, *7*(4), 434–455. doi:10.1080/10618600.1998.10474787

Coburn, D. K., & Geiser, F. (1998). Seasonal changes in energetics and torpor patterns in the subtropical blossom-bat *Syconycteris australis*

(Megachiroptera). *Oecologia*, *113*(4), 467–473. doi:10.1007/s004420050399

Cooper, C. E., & Withers, P. C. (2009). Effects of measurement duration on the determination of basal metabolic rate and evaporative water

loss of small marsupials: How long is long enough? *Physiological and Biochemical Zoology*, *82*(5), 438–446. doi:10.1086/603654

Daniels, H. L. (1984). Oxygen Consumption in *Lemur fulvus*: Deviation from the Ideal Model. *Journal of Mammalogy*, *65*(4), 584–592.

doi:10.2307/1380841

Geiser, F. (1988a). Daily torpor and thermoregulation in <i>Antechinus<i/> (Marsupialia): influence of body mass, season, development,

reproduction, and sex. *Oecologia*, *77*(3), 395–399. doi:10.1007/BF00378050

Geiser, F. (1988b). Reduction of metabolism during hibernation and daily torpor in mammals and birds: temperature effect or physiological

inhibition? *Journal of Comparative Physiology B*, *158*(1), 25–37. doi:10.1007/BF00692726

Geiser, F. (2004). Metabolic Rate and Body Temperature Reduction During Hibernation and Daily Torpor. *Annual Review of Physiology*, *66*(1),

239–274. doi:10.1146/annurev.physiol.66.032102.115105

Geiser, F. (2011). Hibernation: Endotherms. In *eLS*. Chichester, UK: John Wiley & Sons, Ltd. doi:10.1002/9780470015902.a0003215.pub2

Geiser, F., & Brigham, R. M. (2012). The other functions of torpor. In *Living in a Seasonal World* (pp. 109–121). Springer Berlin Heidelberg.

doi:10.1007/978-3-642-28678-0

Geiser, F., Coburn, D. K., Kourtner, G., & Law, B. S. (1996). Thermoregulation, energy metabolism, and torpor in blossom-bats, *Syconycteris*

australis (Megachiroptera). *Journal of Zoology*, *239*(3), 583–590. doi:10.1111/j.1469-7998.1996.tb05944.x

Geiser, F., & Kenagy, G. J. (1988). Torpor Duration in Relation to Temperature and Metabolism in Hibernating Ground Squirrels. *Physiological*

Zoology, *61*(5), 442–449. doi:10.1086/physzool.61.5.30161266

Geiser, F., & Masters, P. (1994). Torpor in relation to reproduction in the mulgara, *Dasycercus cristicauda* (Dasyuridae: Marsupialia). *Journal of*

Thermal Biology, *19*(1), 33–40. doi:10.1016/0306-4565(94)90007-8

Geiser, F., & Ruf, T. (1995). Hibernation versus Daily Torpor in Mammals and Birds: Physiological Variables and Classification of Torpor Patterns.

Physiological Zoology, *68*(6), 935–966. doi:10.1086/physzool.68.6.30163788

Genoud, M. (1993). Temperature regulation in subtropical tree bats. *Comparative Biochemistry and Physiology -- Part A: Physiology*.

doi:10.1016/0300-9629(93)90324-W

Genoud, M. (2014). Thermal energetics of the New-Guinean moss-forest rat (*Rattus niobe*) in comparison with other tropical murid rodents.

Journal of Thermal Biology, *41*(1), 95–103. doi:10.1016/j.jtherbio.2014.01.006

Genoud, M., Bonaccorso, F. J., & Anends, A. (1990). Rate of metabolism and temperature regulation in two small tropical insectivorous bats

(*peropteryx macrotis* and *Natalus tumidirostris*). *Comparative Biochemistry and Physiology -- Part A: Physiology*, *97*(2), 229–234.

doi:10.1016/0300-9629(90)90177-T

Genoud, M., Isler, K., & Martin, R. D. (2018). Comparative analyses of basal rate of metabolism in mammals: data selection does matter.

Biological Reviews, *93*(1), 404–438. doi:10.1111/brv.12350

Genoud, M., Martin, R. D., & Glaser, D. (1997). Rate of metabolism in the smallest simian primate, the pygmy marmoset (*Cebuella pygmaea*).

American Journal of Primatology, *41*(3), 229–245. doi:10.1002/(SICI)1098-2345(1997)41:3<229::AID-AJP5>3.0.CO;2-Z

Glazier, D. S. (2005). Beyond the '3/4-power law': Variation in the intra- and interspecific scaling of metabolic rate in animals. *Biological

Reviews of the Cambridge Philosophical Society*, *80*(4), 611–662. doi:10.1017/S1464793105006834

Grigg, G. C., Augee, M. L., & Beard, L. A. (1992). Thermal Relations Of Free-Living Echidnas During Activity And In Hibernation In A Cold Climate.

Platypus and Echidnas.

Grimpo, K., Legler, K., Heldmaier, G., & Exner, C. (2013). That's hot: Golden spiny mice display torpor even at high ambient temperatures.

Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology, *183*(4), 567–581. doi:10.1007/s00360-012-

0721-4

Hainsworth, F. R., & Wolf, L. L. (1970). Regulation of oxygen consumption and body temperature during torpor in a hummingbird, *Eulampis*

jugularis. *Science*, *168*(3929), 368–369. doi:10.1126/science.168.3929.368

Heldmaier, G., Ortmann, S., & Elvert, R. (2004). Natural hypometabolism during hibernation and daily torpor in mammals. *Respiratory*

Physiology and Neurobiology, *141*(3), 317–329. doi:10.1016/j.resp.2004.03.014

Heldmaier, G., Steinlechner, S., Ruf, T., Wiesinger, H., & Klingenspor, M. (1989). Photoperiod and Thermoregulation in Vertebrates: Body

Temperature Rhythms and Thermogenic Acclimation. *Journal of Biological Rhythms*, *4*(2), 139–153. doi:10.1177/074873048900400211

Hosken, D. J., & Withers, P. C. (1999). Metabolic Physiology of Euthermic and Torpid Lesser Long-Eared Bats, *Nyctophilus geoffroyi* (Chiroptera:

Vespertilionidae). *Journal of Mammalogy*, *80*(1), 42–52. doi:10.2307/1383206

Hudson, J. W., & Scott, I. M. (1979). Daily Torpor in the Laboratory Mouse, *Mus musculus* Var. Albino. *Physiological Zoology*, *52*(2), 205–218.

doi:10.1086/physzool.52.2.30152564

Humphries, M. M., Thomas, D. W., & Speakman, J. R. (2002). Climate-mediated energetic constraints on the distribution of hibernating

mammals. *Nature*, *418*(6895), 313–316. doi:10.1038/nature00828

Huwaldt, J. A., & Steinhorst, S. (2015). Plot Digitizer, version 2.6.8. Free Software Foundation. Retrieved from

http://plotdigitizer.sourceforge.net

IUPS Thermal Commission. (2003). Glossary of terms for thermal physiology. *Journal of Thermal Biology*, *28*(1), 75–106. doi:10.1016/S0306-

4565(02)00055-4

Kelm, D. H., & von Helversen, O. (2007). How to budget metabolic energy: Torpor in a small Neotropical mammal. *Journal of Comparative*

Physiology B: Biochemical, Systemic, and Environmental Physiology, *177*(6), 667–677. doi:10.1007/s00360-007-0164-5

Levesque, D. L. (2008). *Seasonal changes in behavioural and thermoregulatory responses to hypoxia in the eastern chipmunk (*Tamias striatus*)*.

Brock University, St. Catharines, Ontario, Canada. Retrieved from http://hdl.handle.net/10464/1604

Levesque, D. L., & Lovegrove, B. G. (2014). Increased homeothermy during reproduction in a basal placental mammal. *Journal of Experimental*

Biology, *217*(9), 1535–1542. doi:10.1242/jeb.098848

Lovegrove, B. G. (2012). The evolution of mammalian body temperature : the Cenozoic supraendothermic pulses, 579–589.

doi:10.1007/s00360-011-0642-7

Lovegrove, B. G., & Raman, J. (1998). Torpor patterns in the pouched mouse (*Saccostomus campestris*; Rodentia): A model animal for

unpredictable environments. *Journal of Comparative Physiology - B Biochemical, Systemic, and Environmental Physiology*, *168*(4), 303–

312. doi:10.1007/s003600050150

Lyman, C. P. (1982). Entering hibernation. In C. P. Lyman, J. S. Willis, A. Malan, & W. L. C. H. (Eds.), *Hibernation and Torpor in Mammals and*

Birds (pp. 37–53). Academic Press.

Lyman, C. P., Willis, J. S., Malan, A., & Wang, L. C. H. (1982). *Hibernation and Torpor in Mammals and Birds*. New York, NY: Academic Press.

McKechnie, A. E., Ashdown, R. A. M., Christian, M. B., & Brigham, R. M. (2007). Torpor in an African caprimulgid, the freckled nightjar

Caprimulgus tristigma. *Journal of Avian Biology*, *38*(3), 261–266. doi:10.1111/j.2007.0908-8857.04116.x

McKechnie, A. E., & Lovegrove, B. G. (2002). Avian Facultative Hypothermic Responses: a Review. *The Condor*, *104*(4), 705. doi:10.1650/0010-

5422(2002)104[0705:afhrar]2.0.co;2

McNab, B. K. (1980a). Energetics and the Limits to a Temperate Distribution in Armadillos. *Journal of Mammalogy*, *61*(4), 606–627.

doi:10.2307/1380307

McNab, B. K. (1980b). On Estimating Thermal Conductance in Endotherms. *Physiological Zoology*, *53*(2), 145–156.

doi:10.1086/physzool.53.2.30152577

McNab, B. K. (1988). Energy Conservation in a Tree-Kangaroo (*Dendrolagus matschiei*) and the Red Panda (*Ailurus fulgens*). *Physiological*

Zoology, *61*(3), 280–292. doi:10.1086/physzool.61.3.30161241

McNab, B. K. (1989). Temperature Regulation and Rate of Metabolism in Three Bornean Bats. *Journal of Mammalogy*, *70*(1), 153–161.

doi:10.2307/1381678

McNab, B. K. (1997). On the utility of uniformity in the definition of basal rate of metabolism. *Physiological Zoology*, *70*(6), 718–720.

doi:10.1086/515881

McNab, B. K. (2002). *The Physiological Ecology of Vertebrates. A View from Energetics*. Cumstock Publishing Associates, Cornell University

Press, Ithaca & London.

Mersmann, O., Trautmann, H., Steuer, D., & Bornkamp, B. (2018). truncnorm: Truncated Normal Distribution, package version 1.0-8. Retrieved

from https://cran.r-project.org/package=truncnorm

Morris, S., Curtin, A. L., & Thompson, M. B. (1994). Heterothermy, torpor, respiratory gas exchange, water balance and the effect of feeding in

Gould's long-eared bat Nyctophilus gouldi. *The Journal of Experimental Biology*, *197*(1), 309 LP – 335. Retrieved from

http://jeb.biologists.org/content/197/1/309.abstract

Mzilikazi, N., & Lovegrove, B. G. (2002). Reproductive activity influences thermoregulation and torpor in pouched mice, *Saccostomus*

campestris. *Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology*, *172*(1), 7–16.

doi:10.1007/s003600100221

Nespolo, R. F., Arim, M., & Bozinovic, F. (2003). Body size as a latent variable in a structural equation model: Thermal acclimation and

energetics of the leaf-eared mouse. *Journal of Experimental Biology*, *206*(13), 2145–2157. doi:10.1242/jeb.00396

Nowack, J., Levesque, D. L., Reher, S., & Dausmann, K. H. (2020). Variable Climates Lead to Varying Phenotypes : " Weird " Mammalian Torpor

and Lessons From Non-Holarctic Species, *8*(March). doi:10.3389/fevo.2020.00060

Nowack, J., Stawski, C., & Geiser, F. (2017). More functions of torpor and their roles in a changing world. *Journal of Comparative Physiology B:*

Biochemical, Systemic, and Environmental Physiology, *187*(5–6), 889–897. doi:10.1007/s00360-017-1100-y

R Development Core Team. (2012). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna,

Austria.

Reher, S., Ehlers, J., Rabarison, H., & Dausmann, K. H. (2018). Short and hyperthermic torpor responses in the Malagasy bat *Macronycteris*

commersoni reveal a broader hypometabolic scope in heterotherms. *Journal of Comparative Physiology B: Biochemical, Systemic, and*

Environmental Physiology, *188*(6), 1015–1027. doi:10.1007/s00360-018-1171-4

Ruf, T., & Geiser, F. (2015). Daily torpor and hibernation in birds and mammals. *Biological Reviews*, *90*(3), 891–926. doi:10.1111/brv.12137

Sassi, P. L., & Novillo, A. (2015). Acclimating to thermal changes: Intraspecific variation in a small mammal from the Andes Mountains.

Mammalian Biology, *80*(2), 81–86. doi:10.1016/j.mambio.2014.12.005

Schleucher, E., & Withers, P. C. (2001). Re-evaluation of the allometry of wet thermal conductance for birds. *Comparative Biochemistry and*

Physiology - A Molecular and Integrative Physiology, *129*(4), 821–827. doi:10.1016/S1095-6433(01)00356-7

Scholander, P. F., Hock, R., Walters, V., Johnson, F., & Irving, L. (1950). Heat regulation in some arctic and tropical mammals and birds. *The*

Biological Bulletin, *99*(2), 237–258. doi:10.2307/1538741

Song, X., Körtner, G., & Geiser, F. (1995). Reduction of metabolic rate and thermoregulation during daily torpor. *Journal of Comparative*

Physiology B, *165*(4), 291–297. doi:10.1007/BF00367312

Song, X., Körtner, G., & Geiser, F. (1997). Thermal relations of metabolic rate reduction in a hibernating marsupial. *American Journal of*

Physiology - Regulatory Integrative and Comparative Physiology, *273*(6 42-6), 2097–2104. doi:10.1152/ajpregu.1997.273.6.r2097

Sparti, A., & Genoud, M. (1989). Basal rate of metabolism and temperature regulation in Sorex coronatus and S. minutus (Soricidae:

Mammalia). *Comparative Biochemistry and Physiology -- Part A: Physiology*. doi:10.1016/0300-9629(89)90576-8

Speakman, J. R., & Thomas, D. W. (2003). Physiological ecology and energetics of bats. In T. H. Kunz & M. B. Fenton (Eds.), *Bat Ecology* (pp.

430–490). University of Chicago Press.

van Breukelen, F., & Martin, S. L. (2015). The Hibernation Continuum: Physiological and Molecular Aspects of Metabolic Plasticity in Mammals.

Physiology, *30*(4), 273–281. doi:10.1152/physiol.00010.2015

Willis, C. K. R. (2007). An Energy-Based Body Temperature Threshold between Torpor and Normothermia for Small Mammals. *Physiological and*

Biochemical Zoology, *80*(6), 643–651. doi:10.1086/521085

Willis, C. K. R., Lane, J. E., Liknes, E. T., Swanson, D. L., & Brigham, R. M. (2005). Thermal energetics of female big brown bats (*Eptesicus fuscus*

). *Canadian Journal of Zoology*, *83*(6), 871–879. doi:10.1139/z05-074

Withers, P. C., Cooper, C. E., Maloney, S. K., Bozinovic, F., & Cruz-Neto, A. P. (2016). *Ecological and Environmental Physiology of Mammals*.

Oxford University Press. doi:10.1093/acprof:oso/9780199642717.001.0001

Youngflesh, C. (2018). MCMCvis: Tools to Visualize, Manipulate, and Summarize MCMC Output. *Journal of Open Source Software*, *3*(24), 640.

doi:10.21105/joss.00640

Zeileis, A., & Hothorn, T. (2002). Diagnostic Checking in Regression Relationshipslmtest citation info. *R News*, *2*(3), 7--10. Retrieved from

http://cran.r-project.org/doc/Rnews/

Table 1: Model parameters and corroboration index for 28 data sets on small or medium-sized mammals described as either heterotherms or homeotherms.

Species	Type	m	Autors M_{TNZ}	M units	Authors T_{lc}	Authors T_t	Authors	N	T_{lc}	M_{TNZ}	TMR	T_t	β_t	α_e	β_c	α_c	α_r	MR inhibit	T_{be}	T_{bt}	PPO TMR	PPO M_r	Valid assignments	Corroboration	M assigned to torpor?
Unit	-	g	M units	-	°C	°C	-	-	°C	M units	M units	°C	M units/°C	M units	log(M unit)/°C	M units	M units	%	°C	°C	%	%	-	-	-
Callithrix pygmaea	Homeotherm	153	98	mlO$_2$ h^{-1}	27.5 [8]	–	Genoud et al. 1997	27	29.15	102.97	NI	NI	-9.78	388.45	NI	NI	NI	NI	39.77	NI	89.80	86.80	0.63	1.00	No
Cercartetus lepidus	Heterotherm	12.6	1.49	mlO$_2$ g^{-1} h^{-1}	28.8	5.3	Geiser 1987	57	29.01	1.54	0.11	4.16	-0.20	7.47	0.08	0.08	0.95	0.49	36.72	4.68	4.60	33.50	0.98	1.00	Yes
Cercartetus nanus	Heterotherm	36	0.66	mlO$_2$ g^{-1} h^{-1}	28.7	4.8	Song et al. 1997	171	26.15	0.64	0.06	2.86	-0.14	4.29	0.05	0.05	0.46	0.27	30.55	3.26	8.10	27.10	0.95	0.85	Yes
Cynopterus brachyotis	Heterotherm	37.4	1.27	mlO$_2$ g^{-1} h^{-1}	29.7 [6]	–	McNab 1989	51	29.08	1.25	0.74	25.23	-0.21	7.18	NI	NI	NI	NI	34.38	28.76	44.80	94.30	0.76	0.92	Yes
Dasyuroides byrnei [1]	Heterotherm	118.2	0.7	mlO$_2$ g^{-1} h^{-1}	28.3 [7]	15	Geiser & Baudinette 1987	75	26.13	0.64	0.38	12.97	-0.12	3.66	0.01	0.34	1.89	0.67	31.51	16.24	21.50	29.90	0.89	0.87	Yes
Eptesicus fuscus	Heterotherm	15	16.98	mlO$_2$ h^{-1}	26.7	0	Willis et al. 2005	29	27.46	6.69	0.61	NI	-2.81	83.39	NI	NI	NI	NI	29.66	NI	13.60	80.50	0.93	1.00	Yes
Lasiurus cinereus [2]	Heterotherm	23.5 [4]	9	mW g^{-1}	30	–	Cryan & Wolf 2003	48	26.87	8.95	1.67	8.68	-1.09	38.23	NI	NI	NI	NI	35.15	10.22	24.70	78.00	0.90	0.81	Yes
Lasiurus seminolus	Heterotherm	9.4 [5]	1.26 [5]	mlO$_2$ g^{-1} h^{-1}	32	5	Genoud 1993	204	30.69	1.43	0.12	6.27	-0.42	14.22	0.08	0.08	2.74	0.60	34.03	6.57	7.60	23.20	0.95	0.99	Yes
Macroglossus minimus	Heterotherm	16.3	1.29	mlO$_2$ g^{-1} h^{-1}	30.9	20.4	Bartels et al. 1998	46	30.54	0.90	0.47	20.38	-0.16	5.89	0.06	0.16	3.78	0.93	36.32	23.35	27.50	26.30	0.65	0.97	Yes
Macroglossus minimus [3]	Heterotherm	16.4	0.88	mlO$_2$ g^{-1} h^{-1}	31	21	Bonaccorso & McNab 1997	47	30.05	1.23	0.78	18.79	-0.22	7.75	NI	NI	NI	NI	35.78	22.39	30.50	79.50	0.91	1.00	Yes
Marmosa robinsoni	Heterotherm	122	0.8	mlO$_2$ g^{-1} h^{-1}	26.5	–	McNab 1978	38	28.73	0.77	NI	NI	-0.10	3.49	NI	NI	NI	NI	36.41	NI	77.70	92.10	0.68	1.00	No
Melonycteris melanops	Heterotherm	53.3	0.81	mlO$_2$ g^{-1} h^{-1}	28	–	Bonaccorso & McNab 1997	30	28.84	0.81	0.44	17.05	-0.13	4.47	NI	NI	NI	NI	35.00	20.53	64.60	87.30	0.83	1.00	Yes
Natalus tumidirostris	Heterotherm	5.4	1.54	mlO$_2$ g^{-1} h^{-1}	28	20	Genoud et al. 1990	30	27.22	1.46	0.82	19.00	-0.58	17.07	NI	NI	NI	NI	29.54	20.42	58.70	76.70	0.83	1.00	Yes
Nyctophilus geoffroyi	Heterotherm	8	1.42	mlO$_2$ g^{-1} h^{-1}	34	–	Hosken & Withers 1999	84	32.32	1.43	0.24	8.62	-0.27	10.12	0.05	0.18	2.56	0.52	37.60	9.53	23.70	65.60	1.00	1.00	Yes
Peropteryx macrotis	Heterotherm	5.1	2.31	mlO$_2$ g^{-1} h^{-1}	30.5	20	Genoud et al. 1990	49	35.01	2.50	1.80	20.36	-0.22	10.14	0.02	1.35	6.20	0.90	46.99	28.70	22.30	60.40	0.71	0.66	Yes
Phenacomys intermedius	Homeotherm	21.5	3.13	mlO$_2$ g^{-1} h^{-1}	25.1 [6]	–	McNab 1992	75	25.48	3.09	NI	NI	-0.25	9.36	NI	NI	NI	NI	37.78	NI	80.00	94.60	0.85	1.00	No
Pipistrellus pipistrellus	Heterotherm	4.9	7.6	mlO$_2$ h^{-1}	33.2	0	Genoud & Christe 2011	95	32.79	7.62	0.54	2.22	-3.01	106.17	0.08	0.46	7.20	0.80	35.32	2.40	6.40	19.60	1.00	1.00	Yes
Planigale gilesi	Heterotherm	8.3	1.43	mlO$_2$ g^{-1} h^{-1}	28.5 [8]	14	Geiser & Baudinette 1988	62	27.76	1.53	0.30	13.41	-0.37	11.65	0.07	0.13	5.19	0.55	31.93	14.22	19.90	34.40	0.89	1.00	Yes
Rattus niobe	Homeotherm	42.3	53.6	mlO$_2$ h^{-1}	26	–	Genoud 2014	62	27.56	56.18	NI	NI	-4.56	181.97	NI	NI	NI	NI	39.97	NI	84.90	89.00	0.76	0.98	No
Setifer setosus	Heterotherm	530	0.231	mlO$_2$ g^{-1} h^{-1}	28	–	McNab 1980a	56	29.37	0.23	0.14	NI	-0.03	1.18	0.01	0.14	NI	0.88	36.20	NI	34.70	56.60	0.77	0.95	Yes
Sminthopsis macroura	Heterotherm	24.8	0.89	mlO$_2$ g^{-1} h^{-1}	31.3	14	Song et al. 1995	155	30.80	0.84	0.18	16.39	-0.19	6.63	0.09	0.05	3.27	0.77	35.24	17.36	16.50	30.60	0.97	1.00	Yes
Sorex coronatus	Homeotherm	9.1	5.7	mlO$_2$ g^{-1} h^{-1}	27.5	–	Sparti & Genoud 1989	33	22.61	5.94	NI	NI	-0.24	11.37	NI	NI	NI	NI	47.56	NI	95.20	97.20	0.97	0.81	No
Sorex minutus	Homeotherm	3.3	8.6	mlO$_2$ g^{-1} h^{-1}	27.5	–	Sparti & Genoud 1989	28	25.36	8.67	NI	NI	-0.40	18.98	NI	NI	NI	NI	46.99	NI	95.60	95.00	0.89	1.00	No
Steatomys pratensis	Heterotherm	27.5	1.315	mlO$_2$ g^{-1} h^{-1}	28	20	Ellison 1995	39	26.73	1.24	0.26	16.38	-0.18	6.02	0.07	0.13	3.18	0.42	33.88	17.85	21.60	45.40	0.77	1.00	Yes
Syconycteris australis	Heterotherm	17.8	1.44	mlO$_2$ g^{-1} h^{-1}	29.5	18	Geiser et al. 1996	91	32.13	1.27	0.49	15.11	-0.14	5.78	0.05	0.24	2.61	0.90	41.31	18.62	21.60	28.60	0.88	0.95	Yes
Syconycteris australis [1]	Heterotherm	18	1.38	mlO$_2$ g^{-1} h^{-1}	30.2	18	Coburn & Geiser 1998	93	27.53	1.49	0.61	16.37	-0.17	6.00	0.07	0.23	3.33	0.86	36.22	20.07	32.00	43.10	0.85	0.97	Yes
Tolypeutes matacus	Homeotherm	1160	0.181	mlO$_2$ g^{-1} h^{-1}	28	–	McNab 1980a	66	29.57	0.19	NI	NI	-0.04	1.24	NI	NI	NI	NI	33.94	NI	76.80	95.30	0.85	1.00	No
Zaedyus pichiy	Heterotherm	1740	0.226	mlO$_2$ g^{-1} h^{-1}	28	–	McNab 1980a	57	31.82	0.23	0.15	22.49	-0.02	0.94	0.02	0.11	0.65	0.82	42.25	29.47	29.30	71.70	0.70	0.75	Yes

Subset of published data considered: [1]=summer group; [2]=males; [3]=lowland population.

m=average body mass; Autor's M_{TNZ}=author's estimation of metabolic rate within TNZ; M units=units for metabolic rate; Authors' T_{lc} and T_t=author's estimation of lower critical temperature and threshold ambient temperature for torpor, respectively. When m, M_{TNZ} or T_{lc} were not explicitly provided, they were: [4]=back-calculated from M_{TNZ} in % of expected from body mass; [5]=calculated as a weighted average of two seasonal groups; [6]=extracted from a graph; [7]=calculated as the intercept of the euthermic line and M_{TNZ} given by the authors; [8]=obtained as the

662   mean of a given range; N=number of digitized data points (sample size); Model parameters are defined in text. NI=Parameter considered as not

663   identifiable.

Figure 1: Relation between *M* during rest and torpor and T_a on which the present method is based. Torpor is indicated in blue, euthermic rest below the TNZ in red and rest within the TNZ in black. *TMR*: minimal metabolic rate in torpor, M_{TNZ}: resting metabolic rate within the TNZ, T_t: threshold T_a below which a minimum T_b in torpor is regulated by varying heat production (regulated torpor), T_{lc}: lower critical temperature of TNZ.

Figure 2: Four data sets of M measured at different ambient temperature (T_a). Values were assigned to torpor (blue), euthermia (red) and M_{tnz} (black) using the presented three-step method. Predicted values: median and 95% credible intervals are represented by continuous and segmented lines, respectively. Invalid assignments are highlighted with a cross and mismatches between authors and model assignments are surrounded with black circle. A/ Perfect corroboration between model and authors assignments in a heterotherm: *Nyctophilus geoffroyi* (Hosken & Withers, 1999). B/ Absence of mismatched assignment between model and authors assignments with some invalid assignments in a homeotherm: *Sorex minutus* (Sparti & Genoud, 1989). C/ Presence of mismatched or invalid assignments: *Peropteryx macrotis* (Genoud et al., 1990). D/ Absence of mismatched assignment between model and authors assignments with some invalid assignments, but insufficient number

of torpor values to identify some torpor function parameters: *Melonycteris melanops* (Bonaccorso & McNab, 1997).

Figure 3: Frequency distribution of all the measurements from the 28 considered studies in relation to the difference between the method's estimated T_{lc} and the experimental T_a (N=1898). Invalid assignments (i.e. assignment confidence < 0.8) are represented in yellow and assignments mismatches in orange and blue. Mismatches caused by the difference between the model's estimated T_{lc} and that of the authors are highlighted in orange and the remaining mismatches in blue.

670