Assigning metabolic rate measurements to torpor and

euthermia in heterothermic endotherms: "torpor" a new

package for R.

Nicolas J. Fasel[1,2], Colin Vullioud[2], Michel Genoud[1,3]

[1] Department of Ecology and Evolution, University of Lausanne, CH–1015 Lausanne,

Switzerland.

[2] Leibniz Institute for Zoo and Wildlife Research, D-10315 Berlin, Germany

[3] Division of Conservation Biology, Institute of Ecology and Evolution, Department of

Biology, University of Bern, CH–3012 Bern, Switzerland.

Corresponding author:

Nicolas J. Fasel

Department of Ecology and Evolution

University of Lausanne

CH–1015 Lausanne

Switzerland

fasel.nicolas@gmail.com

Abstract

1.   Torpor is a state of controlled reduction of metabolic rate (M) and body

temperature in endotherms. Assigning measurements of M to torpor or euthermia

can be challenging. All current techniques available to distinguish between those two

states have their limitations and are at least partly arbitrary.

2.   Here, we firstly propose a new R package (torpor) enabling distinction of torpid

versus euthermic M measured in stable environmental conditions. Functions are

based on the variation in M measured along varying ambient temperatures (T_a). This

model determines physiological state membership using a binary mixture model, and

further allows prediction of M along T_a. Secondly, we challenge this method by

applying it to previously published data (N=28) on the M of various mammals

covering a diversity of metabolic and thermal strategies.

3.   The fractions of values for which the original assignment matched that of the

model ranged between 0.78 and 1 (median=0.99). Most of the conflicts concerned M

measured at temperatures close to the thermoneutral zone. Parameters describing

torpid M could not always be identified accurately with some smaller data sets.

4.   The package "torpor" provides an objective method to assign measurements

of M to euthermia and/or torpor, and to predict M values at any given T_a below the

thermoneutral zone. Necessary considerations for a lucid application of this method

are further discussed. Ultimately, the use of this package should improve the

standardization of respirometry analyses in heterotherms.

Table 1: Abbreviations used in the present paper and descriptions of the functions of the package "torpor".	
M	metabolic rate
BMR	basal metabolic rate
TMR	minimal metabolic rate in torpor
T_a	ambient temperature
T_b	body temperature
T_{lc}	lower critical temperature (i.e., lower limit of the thermoneutral zone)
T_m	extrapolated modelled temperature where the conforming torpor function reaches BMR
T_t	threshold ambient temperature below which thermoregulation occurs in torpor
tor_fit ()	fits a binomial mixture model using Bayesian inference.
tor_plot ()	provides a plot of the M values over the respective T_a. Raw data and predictions are presented in different colors depending on the metabolic state. Predicted values are represented by continuous and stripped lines for the estimates' median and 95% credible interval bounds of the posterior distribution, respectively. The function is a wrapper function around the tor_fit () and tor_predict (). For more flexibility, output of tor_fit () can be directly included as argument.
tor_predict ()	provides the predicted M and 95% credible interval bounds at a given T_a, for the normothermic and torpid states.
tor_summary()	provides a comprehensive summary of the output returned by tor_fit (). Reported are the mean, 95% credible interval bounds, median and Brooks–Gelman–Rubin criterion (i.e., chain convergence

	check) of parameters posterior distributions. Additionally, prior posterior overlap values for different parameters are generated. Finally, the mean M value used to standardize the M variable is reported.
tor_classify ()	returns the raw data with the related predicted state values (between 1 and 2), which leads to the state classification (torpor or euthermy). Additionally, it also provides the predicted M at the given T_a.

Introduction

Torpor is a state of controlled reduction of M and T_b (Ruf & Geiser, 2015) that

underlies most of the heterothermia observed in endotherms (Geiser & Ruf, 1995;

Lovegrove, 2012). This physiological state has profound implications on energy use

and allocation (Lyman, Willis, Malan, & Wang, 1982), and potentially affects many

biological functions (Geiser & Brigham, 2012; Nowack, Stawski, & Geiser, 2017).

Hence, reports on its occurrence, frequency and depth are highly welcome.

Assigning steady-state measurements of M or T_b to torpor or euthermia can be

challenging, especially in species that are thermolabile (e.g., Geiser, Coburn,

Körtner, & Law, 1996) or that defend an only slightly lowered minimal T_b during torpor

(e.g., Mzilikazi & Lovegrove, 2002). Barclay, Lausen, and Hollis (2001) reviewed the

variety of criteria that have been chosen in the past to distinguish torpor from

euthermia. Apart from a minority of reports that identified torpor on the basis of

behavioural features (e.g., Brice, Grigg, Beard, & Donovan, 2002), the vast majority

of studies have used patterns of variation in body (or skin) temperature and/or rate of

metabolism to separate the two states in the field or in the laboratory. Animals were

usually deemed to be in torpor below a threshold T_b or M (Geiser et al., 1996; Coburn

& Geiser, 1998; Hosken & Withers; 1999, Mzilikazi & Lovegrove, 2002; Kelm & von

Helversen, 2007; Levesque, 2008), or below some threshold temperature differential

between the body and air (Levesque & Lovegrove, 2014), or below a threshold

percentage of the euthermic rate of metabolism (Hudson & Scott, 1979; Geiser,

1988). The diagnostic threshold value was sometimes calculated on the basis of the

parameter's variation in the euthermic state (e.g., Lovegrove & Raman, 1998;

McKechnie, Ashdown, Christian, & Brigham, 2007). It can also be predicted by an

equation on the basis of body mass and T_a (Willis, 2007). All available techniques

have their limitations and are at least partly arbitrary (Barclay et al., 2001), despite

the efforts made to render them more objective.

Here, we present a new package "torpor" running under R (R Development Core

Team, 2012) allowing an objective assignment of M values measured in stable

environmental conditions to torpor or euthermia. Furthermore, "torpor" can also be

useful to estimate T_t and any M values along a range of Ta, including TMR. The

package comprises several useful functions listed in table 1, that will improve

standardization of the analyses of steady-state metabolic measures.

This novel approach is based on the variation observed among measurements of M

made in steady-state conditions. We assume that M follows the "standard"

relationship with T_a modelled by Humphries, Thomas, and Speakman (2002) and

Speakman and Thomas (2003) for bats. We additionally consider the possibility for a

metabolic inhibition during torpor to occur (Geiser, 2004; Withers, Cooper, Maloney,

Bozinovic, & Cruz-Neto, 2016). This model describes fairly well the vast majority of

patterns observed among endotherms entering daily torpor or hibernation. We further

explore the advantages and limitations of this new method by applying it to previously

published data on the M of various mammals.

Material and Methods

The Model

The package "torpor" is based on a binary mixture model using Bayesian inference. It

determines the physiological state membership (euthermia vs torpor) of each M

measurement. The package is constructed around the standard model for bats

(Humphries, Thomas, & Speakman, 2002; Speakman & Thomas, 2003) with the

consideration of metabolic inhibition during torpor (Geiser, 2004; Withers, Cooper,

Maloney, Bozinovic, & Cruz-Neto, 2016; Fig. 1; more details in the package

documentation). In addition to measured values of M along T_a, the method only

requires two parameters from researchers (i.e., BMR and T_{lc}).

Data collection and model testing

We tested the model on 28 published sets of metabolic data obtained by

respirometry on 26 mammal species (supplementary table). In each set, M measured

at T_a's below the thermoneutral zone were assigned by the authors to either of two

states, hereafter referred to as euthermia and torpor (details in supplementary

material). Chosen studies reported BMR and T_{lc} estimates, although the latter had to

be extracted from graphs in two cases. We calculated a corroboration index as the

fraction of values for which the authors' assignment matched the model's one.

Results

Corroboration index ranged between 0.78 and 1 (median = 0.99, N = 28, supplementary material). It was equal to 1 in 12 data sets (e.g., Fig. 2A and 2D). Corroboration indexes lower than 0.95 were restricted to heterotherms for which a limited number of values ($N < 50$) were available (e.g., Fig. 2B). Most of the conflicts (27 out of 41 cases) concerned M measured less than 6°C below T_{lc}. In 12 data sets for which T_t was estimated both by the original authors and by the model, the difference between the two estimates amounted 1.47 ± 0.83°C (range 0.05°C – 2.9°C, supplementary material).

Figure 1: Representation of the relation between M and T_a following the standard model for bats (Humphries, Thomas, & Speakman, 2002; Speakman & Thomas, 2003) with the consideration of metabolic inhibition during torpor (Geiser, 2004; Withers, Cooper, Maloney, Bozinovic, & Cruz-Neto, 2016). Torpor is indicated in blue and euthermia in red. Lines are continuous within the temperature range considered by the model.

Figure 2: Four sets of *M* measured at different T_a below thermal neutrality. Values were assigned to torpor (blue) or euthermia (red) by the binary mixture model. Predicted values: median and 95% credible intervals are represented by continuous and dotted curves, respectively. Mismatches between authors and model assignments are highlighted with a cross. A/ Perfect corroboration between model and authors assignments: *Planigale gilesi* (Geiser & Baudinette, 1988). B/ Some mismatches in the assignments: *Lasiurus cinereus* (Cryan & Wolf, 2003). C/ Perfect corroboration between model and authors assignments, but insufficient number of torpor values to identify torpor function parameters: *Melonycteris melanops* (Bonaccorso & McNab, 1997). D/ Perfect corroboration between model and authors assignments in a strict homeotherm: *Rattus niobe* (Genoud, 2014).

123 Discussion

The new package "torpor" provides an objective method to assign measurements of

M to euthermia and/or torpor. Additionally, it enables prediction of M at any T_a

(including TMR), as well as of T_t. The estimation of T_t further enables the objective

assignment of M values to regulated versus conforming torpor.

The assignments made by the model and by the original authors agreed fairly well as

illustrated by a corroboration index that equalled or approached one. Most conflicting

assignments were observed close to the thermoneutral zone (e.g., Fig. 2B). At these

temperatures, the two states are associated with only slightly different M. Small

sample size and consequently, a poor description of the M in either torpor or

euthermia is surely one cause for these conflicts. Some of them also occurred in

species exhibiting a large intra-state variability in M or a minimum T_b in torpor that is

relatively close to the euthermic T_b (Geiser, Coburn, Körtner, & Law, 1996, Mzilikazi

& Lovegrove, 2002).

For a lucid application of torpor, some considerations should be kept in mind. Firstly,

the standard model on which our method is based (Humphries et al., 2002,

Speakman & Thomas 2003) may not describe the thermal biology of all

heterothermic species. Torpid M may sometimes depend more strongly on body

mass than on T_a (Kelm & von Helversen, 2007). In addition, T_b and/or thermal

conductance may sometimes vary with T_a below thermal neutrality (McNab, 1980) in

such a way that the relationship between euthermic M and T_a is no longer linear.

Secondly, the exclusive use of stable M meeting steady-state conditions during

respirometry runs is an important requirement of the proposed method. In particular,

transitions between euthermia and torpor (i.e., entrance into and arousal from torpor)

as well as incomplete torpor bouts during which M never reaches a stable, minimal

level (Lyman, 1982) should be excluded. Thirdly, the proposed method requires that

values of *BMR* and T_{lc} for the species under investigation are known. Because *M* in

euthermia is assumed to converge at a point of coordinates [T_{lc}, *BMR*], these two

parameters affect the assignment of the data points below the thermoneutral zone.

Ultimately and with consideration of the above-mentioned points, the new R-package

"torpor" should ease and standardize the analysis of respirometry data in

heterotherms.

Authors contribution

NJF designed methodology; MG collected data; CV wrapped the package; All

authors gave final approval for publication.

Acknowledgements

We are thankful to Martyn Plummer for discussions at the very beginning of the

project as well as to Philippe Christe and Nicolas Salamin for their support during the

elaboration of the model. This study was financially supported by the Department of

Ecology and Evolution of the University of Lausanne (Switzerland) and by the Swiss

National Science Foundation (grant number: P2BEP3_168709 to NJF).

REFERENCES

Barclay, R. M. R., Lausen, C. L., & Hollis, L. (2001). What's hot and what's not:

defining torpor in free-ranging birds and mammals. Canadian Journal of

Zoology, 79,1885–1890. doi: 10.1139/z01-138.

Bonaccorso, F. J., & McNab, B. K. (1997). Plasticity of energetics in blossom bats

(Pteropodidae): impact on distribution. Journal of Mammalogy, 78, 1073–

1088. doi: 10.2307/1383050.

Brigham, R.M., Willis, C.K.R., Geiser, F., & Mzilikazi, N. (2011). Baby in the

bathwater: Should we abandon the use of body temperature thresholds to

quantify expression of torpor? Journal of Thermal Biology, 36, 376–379. doi:

10.1016/j.jtherbio.2011.08.

Brice, P. H., Grigg, G. C., Beard, L. A., & Donovan, J. A. (2002). Patterns of activity

and inactivity in echidnas (*Tachyglossus aculeatus*) free-ranging in a hot dry

climate: correlates with ambient temperature, time of day and season.

Australian Journal of Zoology, 50, 461–475. doi: 10.1071/ZO01080.

Brooks, S. P., & Gelman, A. (1998). General methods for monitoring convergence of

iterative simulations. Journal of Computational and Graphical Statistics, 7,

434–455. doi: 10.1080/10618600.1998.10474787.

Coburn, D. K., & Geiser, F. (1998). Seasonal changes in energetics and torpor

patterns in the subtropical blossom-bat *Syconycteris australis*

(Megachiroptera). Oecologia, 113, 467–473. doi: 10.1007/s004420050399.

Cryan, P. M., & Wolf, B. O. (2003). Sex differences in the thermoregulation and

evaporative water loss of a heterothermic bat, *Lasiurus cinereus*, during its

spring migration. Journal of Experimental Biology, 206, 3381–3390. doi:

10.1242/jeb.00574.

Geiser, F. (1988). Daily torpor and thermoregulation in *Antechinus* (Marsupialia):

influence of body mass, season, development, reproduction, and sex.

Oecologia, 77, 395–399. doi: 10.1007/BF00378050.

Geiser, F. (2004). Metabolic rate and body temperature reduction during hibernation

and daily torpor. Annual Review of Physiology, 66, 239–274. doi:

10.1146/annurev.physiol.66.032102.115105.

Geiser, F., & Baudinette, R. V. (1988). Daily torpor and thermoregulation in the small

dasyurid marsupials *Planigale gilesi* and *Ningaui yvonneae*. Australian Journal

of Zoology, 36, 473–481. doi: 10.1071/ZO9880473.

Geiser, F., & Brigham, R. M. (2012). The Other Functions of Torpor. In Ruf, T.,

Bieber, C., Arnold, W., & Milesi, E. (Eds.), *Living in a Seasonal World* (109–

121). Heidelberg, Berlin, D: Springer-Verlag. doi: 10.1007/978-3-642-28678-

0_10.

Geiser, F., Coburn, D. K., Körtner, G., & Law, B. S. (1996). Thermoregulation, energy

metabolism, and torpor in blossom-bats, *Syconycteris australis*

(Megachiroptera). Journal of Zoology, 239, 583–590. doi: 10.1111/j.1469-

7998.1996.tb05944.x.

Geiser, F., & Ruf, T. (1995). Hibernation versus daily torpor in mammals and birds:

physiological variables and classification of torpor patterns. Physiological

Zoology, 68, 935–966. doi: 10.1086/physzool.68.6.30163788.

Genoud, M. (2014). Thermal energetics of the New-Guinean moss-forest rat (*Rattus*

niobe) in comparison with other tropical murid rodents. Journal of Thermal

Biology, 41, 95–103. doi: 10.1016/j.jtherbio.2014.01.006.

Hosken, D. J., & Withers, P. C. (1999). Metabolic physiology of euthermic and torpid

lesser long-eared bats, *Nyctophilus geoffroyi* (Chiroptera : Vespertilionidae).

Journal of Mammalogy, 80, 42–52. doi: 10.2307/1383206.

Hudson, J. W., & Scott, I. M. (1979). Daily torpor in the laboratory mouse, *Mus*

musculus var. albino. Physiological Zoology, 52, 205–218. doi:

10.1086/physzool.52.2.30152564.

Humphries, M. M., Thomas, D. W., & Speakman, J. R. (2002). Climate-mediated

energetic constraints on the distribution of hibernating mammals. Nature, 418,

313–316. doi: 10.1038/nature00828.

Kelm, D. H., & von Helversen, O. (2007). How to budget metabolic energy: torpor in a

small Neotropical mammal. Journal of Comparative Physiology, B, 177, 667–

677. doi: 10.1007/s00360-007-0164-5.

Levesque, D. L. (2008). Seasonal changes in behavioural and thermoregulatory

responses to hypoxia in the eastern chipmunk (*Tamias striatus*). M.Sc. Brock

University, St. Catharines, Ontario, Canada. URL:

http://hdl.handle.net/10464/1604.

Levesque, D. L., & Lovegrove, B. G. (2014). Increased homeothermy during

reproduction in a basal placental mammal. Journal of Experimental Biology,

217, 1535–1542. doi: 10.1242/jeb.098848.

Lovegrove, B. G. (2012). The evolution of endothermy in Cenozoic mammals: a

plesiomorphic-apomorphic continuum. Biological Reviews, 87, 128–162. doi:

10.1111/j.1469-185X.2011.00188.x.

Lovegrove, B. G., & Raman, J. (1998). Torpor patterns in the pouched mouse

(*Saccostomus campestris*; Rodentia): a model animal for unpredictable

environments. Journal of Comparative Physiology, B, 168, 303–312. doi:

10.1007/s003600050150.

Lyman, C. P. (1982). Entering hibernation. In C. P. Lyman, J. S. Willis, A. Malan, & L.

C. H. Wang (Eds.), *Hibernation and Torpor in Mammals and Birds* (pp. 37–

53). New York, NY: Academic Press.

Lyman, C. P., Willis, J. S., Malan, A., & Wang, L. C. H. (1982). Hibernation and

Torpor in Mammals and Birds. New York, NY: Academic Press.

McKechnie, A. E., Ashdown, R. A. M., Christian, M. B., & Brigham, R. M. (2007).

Torpor in an African caprimulgid, the freckled nightjar *Caprimulgus tristigma*.

Journal of Avian Biology, 38, 261–266. doi: 10.1111/j.2007.0908-

8857.04116.x.

McNab, B. K. (1980). On estimating thermal conductance in endotherms.

Physiological Zoology, 53, 145–156. doi: 10.1086/physzool.53.2.30152577.

Mzilikazi, N., & Lovegrove, B. G. (2002). Reproductive activity influences

thermoregulation and torpor in pouched mice, *Saccostomus campestris*.

Journal of Comparative Physiology, B, 172, 7–16. doi:

10.1007/s003600100221.

Nowack, J., Stawski, C., & Geiser, F. (2017). More functions of torpor and their roles

in a changing world. Journal of Comparative Physiology, B, 187, 889–897. doi:

10.1007/s00360-017-1100-y.

R Development Core Team. (2012). R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-

900051-07-0. URL: http://www.R-project.org.

Ruf, T., & Geiser, F. (2015). Daily torpor and hibernation in birds and mammals.

Biological Reviews, 90, 891–926. doi: 10.1111/brv.12137.

Speakman, J. R., & Thomas, D. W. (2003). Physiological ecology and energetics of

bats. In T. H. Kunz & M. B. Fenton (Eds.), *Bat Ecology* (pp. 430–490).

Chicago, IL: University of Chicago Press.

Willis, C. K. R., Lane, J. E., Liknes, E. T., Swanson, D. L., & Brigham, R. M. (2005).

Thermal energetics of female big brown bats (*Eptesicus fuscus*). Canadian

Journal of Zoology, 83, 871–879. doi: 10.1139/cjz-2016-0055.

Withers, P. C., Cooper, C. E., Maloney, S. K., Bozinovic, F., & Cruz-Neto, A. P.

(2016). Ecological and Environmental Physiology of Mammals. Oxford, UK:

Oxford University Press.