Assigning metabolic rate measurements to torpor and

euthermy in heterothermic endotherms: "torpor", a new

package for R.

Nicolas J. Fasel[1], Colin Vullioud[2], Michel Genoud[1]

1. Department of Ecology and Evolution, University of Lausanne, CH–1015 Lausanne,
Switzerland.
2. Department Evolutionary Genetics, Leibniz Institute for Zoo and Wildlife Research, D-10315
Berlin, Germany.

Corresponding author:
Nicolas J. Fasel
Department of Ecology and Evolution
University of Lausanne
CH–1015 Lausanne
Switzerland
fasel.nicolas@gmail.com

Running head: A R-package for thermal biology

Key words: Energetics, Metabolism, Bayesian, Mixture models, Energy saving

Summary statement: The presented method and its associated R-package (torpor) enable the
assignment of metabolic rate measurements to torpor or euthermy, ultimately improving the
standardization of respirometry analyses in heterotherms.

Abstract

Torpor is a state of controlled reduction of metabolic rate (M) in endotherms. Assigning

measurements of M to torpor or euthermy can be challenging, especially when the difference

between euthermic M and torpid M is small, in species defending a high minimal body

temperature in torpor, in thermolabile species, and slightly below the thermoneutral zone

(*TNZ*). Here, we propose a novel method for distinguishing torpor from euthermy. We use the

variation in M measured during euthermic rest and torpor at varying ambient temperatures

(T_a) to objectively estimate the lower critical temperature (T_{lc}) of the *TNZ* and to assign

measurements to torpor, euthermic rest or rest within TNZ. In addition, this method allows

the prediction of M during euthermic rest and torpor at varying T_a, including resting M within

the *TNZ*. The present method has shown highly satisfactory results using 28 published sets of

metabolic data obtained by respirometry on 26 species of mammals. Ultimately, this novel

method aims to facilitate analysis of respirometry data in heterothermic endotherms. Finally,

the development of the associated R-package (torpor) will enable widespread use of the

method amongst biologists.

List of main abbreviations:

BMR: Basal metabolic rate.

M: Resting rate of metabolism, expressed in W, $mlCO_2$ h^{-1} or mlO_2 h^{-1}. Often expressed as a

mass-specific rate (e.g. in W g^{-1}).

M_{TNZ}: Resting rate of metabolism within the *TNZ*.

M_e: Euthermic *M*.

M_r: predicted M_t at T_{lc}. M_r may be lower than M_{TNZ} if additional metabolic inhibition is involved,

other than the abolishment or reduction of thermogenesis necessary to maintain euthermy.

M_t: Torpid *M*.

PPO: prior and posterior distribution overlap.

T_a: Ambient temperature.

T_b: Body temperature.

T_{lc}: lower critical T_a or lower limit of the *TNZ*.

TMR: minimal metabolic rate in torpor, measured at T_t.

TNZ: Themoneutral zone. The range of T_a where *M* is independent of T_a.

T_t: Threshold T_a separating regulated from conforming torpor.

Introduction

Torpor is a state of controlled reduction of metabolic rate (M) and body temperature (T_b) observed in numerous mammals and birds (Ruf and Geiser, 2015). This physiological state can occur over short periods (i.e. < 24h), often referred to as 'daily torpor', or it can last up to many days, often referred to as 'hibernation', which typically involves multiday torpor bouts separated by short spontaneous arousals (McKechnie and Lovegrove, 2002; Ruf and Geiser, 2015; Schleucher and Withers, 2001). The drop in T_b may be considerable in hibernators, which often allow T_b to reach values close to 0°C, but it can also be much shallower, as is often the case in species entering daily torpor (Ruf and Geiser, 2015). It has been suggested that a continuum may exist between these types of torpor as well as between torpor and euthermic rest (Boyles et al., 2013; McKechnie and Lovegrove, 2002; van Breukelen and Martin, 2015). Torpor use has profound implications on energy expenditure and allocation (Lyman et al., 1982) and affects many biological functions (Geiser and Brigham, 2012; Nowack et al., 2017). Accordingly, a vast amount of literature describes the occurrence of torpor and its associated energy savings in a multitude of species (Geiser and Ruf, 1995; Lovegrove, 2012; Nowack et al., 2020; Ruf and Geiser, 2015).

Determining whether measurements of M can be assigned to torpor or euthermy can however be challenging. The distinction between these two states is usually straightforward in typical hibernators, due to the more than 90% reduction in M common in these species when torpid at their usually hibernation ambient temperature (T_a) (Geiser, 2004). Yet this distinction can become problematic when the difference between euthermic M (M_e) and torpid M (M_t) is small, which often occurs slightly below the thermoneutral zone (TNZ) (Geiser, 2011;

Hainsworth and Wolf, 1970; Humphries et al., 2002; Speakman and Selman, 2003). Additional

difficulties are encountered with species that either enter daily torpor, referred to as 'daily

heterotherms', whose minimum T_b in torpor often lies only slightly below the euthermic T_b

(Bartels et al., 1998; Bonaccorso and McNab, 1997; Genoud et al., 1990; McNab, 1980b), or

who exhibit a large variability in M and T_b at rest (Coburn and Geiser, 1998; Geiser et al., 1996).

Barclay, Lausen, & Hollis (2001) reviewed the variety of criteria that have been chosen in the

past to distinguish torpor from euthermy. Apart from a minority of reports that identified

torpor on the basis of behavioural features (Brice et al., 2002; Geiser and Kenagy, 1988; Geiser

and Masters, 1994), the vast majority of studies have used patterns of variation in T_b (or skin

temperature) and/or M to separate the two states in the field or in the laboratory. Animals

have been deemed to be in torpor below a threshold T_b or M (Coburn and Geiser, 1998; Geiser

et al., 1996; Hosken and Withers, 1999; Kelm and von Helversen, 2007; Levesque, 2008;

Mzilikazi and Lovegrove, 2002), below some threshold temperature differential between the

body and air (Levesque and Lovegrove, 2014), or below a threshold percentage of the

euthermic rate of metabolism (Geiser, 1988a; Hudson and Scott, 1979). This diagnostic

threshold value was sometimes calculated on the basis of the parameter's variation in the

euthermic state (e.g. Lovegrove & Raman, 1998; McKechnie, Ashdown, Christian, & Brigham,

2007), but has also been predicted by an equation based on body mass and T_a (Willis, 2007).

All available techniques have their limitations and are at least partly arbitrary (Barclay et al.,

2001; Boyles et al., 2011), despite the efforts made to render them more objective.

With this paper, we propose an automated method for distinguishing torpor from euthermic

rest in species that enter torpor, hereafter referred to as heterotherms following the

terminology of Geiser & Ruf (1995) or Lovegrove (2012). Here, assignment to an either

euthermic or torpid state is based on a probabilistic approach using the variation observed

among measurements of M at varying T_a. We assume that the relationship between M of

resting animals and T_a follows the classical "Scholander-Irving model" (Scholander, Hock,

Walters, Johnson, & Irving, 1950; for a discussion see McNab, 2002), which was later extended

to include torpor (Hainsworth & Wolf, 1970; Humphries, Thomas, & Speakman, 2002;

Speakman & Thomas, 2003; Geiser, 2011). This model (Fig. 1) predicts the vast majority of

patterns observed among endotherms entering daily torpor or hibernation. We explicitly

consider that additional metabolic inhibition involving mechanisms other than the

abolishment or reduction of thermogenesis necessary to maintain euthermy may occur during

torpor (Geiser, 1988b; Geiser, 2004; Geiser and Kenagy, 1988; Guppy and Withers, 1999;

Withers et al., 2016), by allowing the curve of torpor M to reach T_{lc} at a level equal to – or

lower than the resting metabolic rate within the TNZ. Thus, over all aforementioned

techniques, our novel method provides major improvements, as it assigns measurements

under an explicit framework (i.e. the extended Sholander-Irving model) and does so with a

probabilistic approach based on observed variation.

Our method is specifically intended to facilitate the discrimination between torpor and

euthermy in laboratory experiments using respirometry. Additionally, it can also be useful to

predict M at varying T_a, by modelling the different parameters of the thermoregulatory curves.

Further, we have assessed the performance of our method, by applying it to previously

published data on the M of various mammals, including both heterotherms and species not

undergoing torpor (hereafter referred to as homeotherms). Finally, we have provided a new

package running within R (R Development Core Team, 2012) to allow researchers to apply our

method to their own data. This package ("torpor") comprises several useful functions that will

improve standardization of the analyses of metabolic measurements for thermal biology.

Results

Over all datasets, the proportion of assignments, that could be validly assigned (i.e.

assignment confidence > 0.80, cf. "Assignment confidence" in section "Material and

Methods") ranged from 0.63 to 1.00 (median=0.87, Table 1). The corroboration index, which

assesses the similarity between valid assignments made by our method and by the authors of

the original datasets, ranged between 0.66 and 1.00 (median=1.00, Table 1). Complete

matches between the method valid assignments and the authors' descriptions were found for

15 out 28 datasets (e.g. *Nyctophilus geoffroyi* (Hosken and Withers, 1999), Fig. 2A).

Mismatches mostly occurred close to T_{lc} (e.g. *Peropteryx macrotis* (Genoud et al., 1990), Fig.

2C) but could also be found as far as 41.0°C below T_{lc} (Fig. 3). Among homeotherms, no M

measurement was assigned to torpor (e.g. *Sorex minutus* (Sparti & Genoud, 1989), Fig. 2B).

In studies of homeotherms, where no M was originally assigned to torpor, prior and posterior

distribution overlaps (*PPOs*) of *TMR* ranged from 76.80 to 95.60%. In identified heterotherms,

PPOs of *TMR* values range from 4.60 to 64.60% (Table 1).

Modelled values of T_{lc}, M_{TNZ} and T_t and extracted values from the original studies were

significantly correlated (T_{lc}: N=28, Pearson's coefficient: 0.65, p<0.001, M_{TNZ}: N=28, Pearson's

coefficient: 1.00, p<0.001, T_t: N=14, Pearson's coefficient: 0.97, p<0.001).

Discussion

Our method aims to facilitate the discrimination between torpor and euthermy using the variation of M measured at varying T_a. We tested it using 28 published sets of metabolic data obtained by respirometry on 26 species of small or medium-sized mammals. Selected species displayed a diversity of metabolic and thermal strategies ranging from permanent homeothermy to heterothermy including shallow, daily torpor and deep, long-term hibernation. The efficiency of our method proved satisfactory. Indeed, the corroboration index was generally high. Most conflicting assignments were a mere consequence of the difference between the estimated T_{lc} and that defined by the authors (Table 1, Figs. 2C & 3). In particular, all mismatches between the model's and the authors' assignments concerning M_e and M_{TNZ} measurements in homeotherm species were explained by differences in estimated T_{lc}. The remaining mismatches could generally be explained by intra-state frequency distributions of M that did not segregate clearly. This condition was often found close to TNZ, but also occurred in species where T_t was relatively close to T_{lc} (e.g. McNab, 1980a, 1989). Such discrepancies illustrate the difficulty of the assignment process when torpor and euthermy have to be distinguished solely on the basis of measurements of M. While the use of a statistical method enables an objective assignment, it is nevertheless worthwhile to recall that some authors also used patterns of T_b to assign their data (e.g. McNab, 1980a, 1989). These specific cases should deserve further attention as they can highlight mechanisms decoupling M from the control of T_b (e.g. Daniels, 1984; McNab, 1988; Heldmaier, Ortmann, & Elvert, 2004). One advantage of our method is to allow researchers to identify statistically data points that are difficult to assign. In this study, we considered assignments with a confidence lower than 0.8 as invalid. Obviously, an increase of this threshold value would lead to fewer assignment mismatches.

Our method also models the relationship between M and T_a specific to each state, as well as

several parameters describing the standard energetics of the animal(s) under study, including

M_{TNZ}, TMR, T_{lc} and T_t. Predicting M at any T_a is crucial to model energy costs (e.g. Boyles,

Johnson, Blomberg, & Lilley, 2020). It should however be recalled that our Bayesian inference-

based method models all parameters whatever the available data. It remains therefore crucial

to consider the parameters' identifiability. The prior and posterior distribution overlaps (PPO)

of T_{bt} and T_{lc} are not reliable for that purpose, because the prior distribution of those

parameters largely depends on the data provided. From the $PPOs$ of TMR obtained in studies

where no M was assigned to torpor (i.e. "homeotherms") and those obtained in identified

heterotherms, we define a PPO higher than 75% as indicative of a parameter that is not

identifiable.

The parameters estimated by our method correlated significantly with those extracted from

the original studies. The number of M_e values provided was probably not always sufficient to

model T_{lc} adequately. On one hand, a statistical estimation of T_{lc} (e.g. in Song, Körtner, &

Geiser, 1995, 1997; Bonaccorso & McNab, 1997; Genoud, Martin, & Glaser, 1997; Hosken &

Withers, 1999; Willis, Lane, Liknes, Swanson, & Brigham, 2005; Genoud, 2014) was often

lacking, leaving space for some potential misleading subjectivity by the authors of the original

studies. On the other hand, an overestimation of T_{lc} by our method remains possible if

appreciable additional metabolic inhibition occurs during torpor (see below). The estimated

T_t was always close to that of the original studies. This higher precision in the estimation of T_t

in comparison to that of T_{lc} is probably due to the lower variation of M exhibited during

conforming torpor in comparison to that measured during euthermic rest. It is also not

surprising that the M_{TNZ} obtained by our method and by the authors of the original studies

were close to identical, since both were calculated as the mean of nearly the same M values

(i.e. all values above T_{lc})

An important aspect incorporated in our method is that we allow function for M_t to be lower

than M_{TNZ} at T_{lc}. We thus explicitly allow for additional metabolic inhibition to occur in some

species during torpor, other than the cessation or reduction of the thermogenesis necessary

to maintain euthermy (Geiser, 2004; Guppy and Withers, 1999; Withers et al., 2016). Caution

should be kept in M_r interpretation as an insufficient number of M_t values corresponding to

conforming torpor leads to the unidentifiability of this parameter (e.g. *Melonycteris melanops*

(Bonaccorso & McNab, 1997), Fig. 2D). Consequently, the mentioned additional metabolic

inhibition should only be considered when sufficient M_t values corresponding to conforming

torpor are provided and M_r PPO indicates adequate parameter's identifiability. Considering

only identifiable parameters (i.e. Mr PPO < 75%), we quantified additional metabolic inhibition

as the fraction M_r / M_{TNZ}. Values ranged between 0.27 and 0.93 (N=18, median=0.77, Table 1).

By extrapolation, we should consider the possibility that a hypometabolic state might occur

within the *TNZ* as well (e.g. Grimpo, Legler, Heldmaier, & Exner, 2013; Reher, Ehlers,

Rabarison, & Dausmann, 2018). With our method, we did not implement an assignment to

such a hypometabolic state within the TNZ, hence M_{TNZ} corresponds to the mean of all values

from the *TNZ*. Our method might therefore underestimate the eumetabolic $M_{TNZ,}$ and might

consequently overestimate T_{lc}, if animals occasionally enter a hypometabolic state in the *TNZ,*.

A predicted M_t at T_{lc} much lower than M_{TNZ} should be viewed as an indication for a strong

additional metabolic inhibition occurring during torpor and should invite for a careful

consideration of the estimated M_{TNZ} and T_{lc}.

Throughout this paper, resting metabolic rate within the *TNZ* is referred to as M_{TNZ} rather than

to the basal rate of metabolism (*BMR*). Estimating the *BMR* requires several specific criteria:

animals should be post-absorptive, adult, non-reproductive, and resting during a major

inactive phase of the daily cycle (McNab, 1997). If these criteria are not met, the method still

remains applicable, but the resting *M* estimated within the *TNZ* (i.e. M_{TNZ}) will correspond to

a minimal resting metabolic rate rather than to the *BMR*.

While the performance of our method appeared to be excellent with the selected datasets,

its main limitations correspond to those of the model on which it is based. Indeed, although

extremely fruitful, the Scholander-Irving model (Scholander et al., 1950) and its latter

extensions (Geiser, 2011; Hainsworth and Wolf, 1970; Humphries et al., 2002; Speakman and

Thomas, 2003) does not describe the thermal biology of all heterothermic species. Causes for

significant divergence from the hereby model include a non-linear relationship between M_e

and T_a (e.g. *Cercartetus nanus*: Song, Körtner, & Geiser, 1997), a difference in the slopes of

the regressions of *M* versus T_a in regulated torpor and euthermic rest (Geiser, 2004) and a

strong dependence of M_t on factors other than T_a, such as body mass (Kelm and von

Helversen, 2007) and duration since the last activity period or since last meal (Grigg et al.,

1992; Morris et al., 1994). Furthermore, this model also does not describe adequately

circadian (e.g. the shallow "rest-phase hypothermy" of many small birds (McKechnie and

Lovegrove, 2002) or ultradian (Heldmaier et al., 1989) variations in *M*.

Our method is able to reveal the presence or the absence of distinct groups of values

(euthermic rest vs torpor), but is not intended to judge whether any of the data points entered

correspond to stable rates and/or to minimal values. Transitions between activity and rest or

between euthermic rest and torpor as well as incomplete torpor bouts during which M never

reaches a stable, minimal level (e.g. "test drops" (Hudson and Scott, 1979); see also (Genoud,

1993; Lyman, 1982)) should be excluded. It should also be recalled that long runs may be

necessary for some heterothermic animals to achieve a state of steady torpor (e.g. Heldmaier,

Ortmann, & Elvert, 2004). The above limitations should be kept in mind for a lucid application.

Then, individual variation is not explicitly considered in our method. Especially, body mass

variation within a studied population might affect the assignment of M to torpor or euthermy

in a complex manner (Genoud, 2014; Genoud et al., 2018; Glazier, 2005; Nespolo et al., 2003;

Sassi and Novillo, 2015; Schleucher and Withers, 2001). However, there would be no particular

difficulty to apply our method to single individuals, or repeatedly on single individuals.

Our method and the associated R-package (torpor) provide a way to standardize the analysis

of respirometry data in relation to T_a. Its major strength is that it uses a probabilistic approach

to assign metabolic values to torpor or euthermy, rather than assigning them on the basis of

a particular threshold value. Thus, the partly arbitrary nature of the assignment process is

removed. Further, it can be applied to study intraspecific as well as interspecific variation in

energetics. Parameters might now be extracted from the literature for comparative analysis,

largely avoiding causes of variation due to the diversity of past assignment techniques.

Ultimately, this method and the associated R-package (torpor) will ease intra- and inter-

specific comparative analyses of endotherm energetics.

Material and Methods

The three-steps Method

Our present method is based on three steps and requires measurements of M of resting or

torpid animals at T_a ranging from below – to within the TNZ. Animals are assumed to be in one

of three "states": euthermic rest below the TNZ (M_e), torpor below the TNZ (M_t) or rest within

the TNZ (M_{TNZ}). We do not consider here the relation between M and T_a above the TNZ.

Consequently, metabolic rates measured at T_a higher than the upper limit of the TNZ should

be excluded. During the different steps, a mixture model based on Bayesian inference is run

under varying conditions.

The model

The Scholander-Irving model and its extensions (Fig. 1) consider that resting M measured

within the TNZ is independent of T_a. This rate is hereafter referred to as M_{TNZ}. Below T_{lc}, the

M of euthermic animals (M_e) increases linearly with decreasing T_a :

$M_e \sim \alpha_e + \beta_e * T_a \mid T_a < T_{lc}$ (1)

The model assumes that M_e equals M_{TNZ} at T_{lc}, which enabled calculation of α_e, the intercept

of the line predicting M_e at varying T_a:

$\alpha_e = M_{TNZ} - \beta_e * T_{lc}$ (2)

and of the slope of that line (β_e):

$\beta_e = \frac{M_{TNZ}}{T_{lc} - T_{be}}$ (3)

T_{be} represents the hypothetical T_a where M_e equals 0, which correspond to the euthermic body

temperature, provided that thermal conductance and body temperature do not vary with T_a

below the *TNZ* (McNab, 1980a).

Below the threshold T_a separating regulated and conforming torpor (T_t), the metabolic rate in

torpor (M_t) increases linearly with decreasing T_a to defend the animal's setpoint T_b in torpor.

That state, corresponding to M_t measured at T_a lower than T_t, is usually referred to as

"regulated torpor":

$M_t \sim \alpha_r + \beta_r * T_a \mid T_a < T_t$ (4)

Where :

$T_t = \frac{(TMR - \alpha_r)}{\beta_r}$ (5)

The intercept of the line for regulated torpor (α_r) is obtained as:

$\alpha_r = -T_{bt} * \beta_r$ (6)

TMR represents the minimal M_t measured at T_t. T_{bt} is the hypothetical T_a at which M_t of the

regulated torpor would equal 0. T_{bt} would correspond to the minimal body temperature in

torpor (i.e. setpoint T_b in torpor) provided that thermal conductance and body temperature

during regulated torpor do not vary with T_a below T_t. Between T_t and T_{lc}, torpor is referred to

as "conforming torpor" and M_t follows an exponential curve:

$M_t \sim \alpha_c * \exp(\beta_c * T_a) \mid T_t \leq T_a \leq T_{lc}$ (7)

Where the coefficient of the exponent (β_c) is calculated as:

$\beta_c = \frac{\log(M_r) - \log(TMR)}{T_{lc} - T_t}$ (8)

and the intercept for the exponential curve (α_c) is calculated as:

$\alpha_c = \frac{M_r}{\exp(\beta_c * T_{lc})}$ (9)

M_r represents M_t measured at T_{lc}. In accordance with the general trend observed (Geiser,

2004), we assume that the slopes of the lines linking T_a to M_e (β_e) and to M_t of the regulated

torpor, (β_r) are similar:

$\beta_r = \beta_e$ (10)

The following parameters are modelled: the fractions of measurements belonging to each

state, which provides the value specific state membership probabilities for each measurement

(see Second step: Metabolic rate (M) measurements pre-assignment, below), T_{be}, T_{bt}, TMR,

and the standard deviations for M_{TNZ} and regulated/conforming metabolic rates (SD_{TNZ}, SD_r

and SD_c).

First step: Estimation of M_{TNZ} and T_{lc}

The method initially defines the highest possible T_a within the dataset that still underestimates

T_{lc} (T_{lc_low}). Above T_{lc_low}, a linear regression between M and T_a should neither result in a

significant negative slope nor be affected by heteroscedasticity. Beginning with the ten M

values measured at the highest T_a, linear regressions are performed on sets of M values

progressively including values at lower T_a. The inclusion of additional M values obtained below

T_{lc} eventually leads to a significantly negative slope in the case of animals not entering torpor

and/or to heteroscedasticity in the case of heterotherms. A significantly positive slope would

reveal an absence or insufficient number of values for euthermic rest below the TNZ, hence it

automatically stops the analysis as T_{lc_low} is undefinable. Heteroscedasticity is assessed with a

Breusch-Pagan test with the function "bptest" from package "lmtest" (Zeileis and Hothorn,

2002). Significance level for the Breusch-Pagan test is set at 0.05. The significance of the

negative regression is assessed with a linear regression, function "lm" and significance level is

set at 0.01.

Then, in order to get T_{lc}, the model described in the second step (see Second step: Metabolic

rate (M) measurements pre-assignment) is first run without data points measured above

T_{lc_low}. Moreover, within this step, the parameter T_{lc} is also modelled. M_{TNZ} provided in that

analysis is the mean of the M measured at T_a higher than T_{lc_low}. Once T_{lc} has been estimated,

M_{TNZ} is recalculated as the mean of the M measured at T_a higher than T_{lc}.

Second step: Metabolic rate (M) measurements pre-assignment

The assignment of M measurements to one of the three physiological states is based on a

probabilistic process. Below T_{lc}, a posterior categorical distribution is generated for each

coupled data (T_a and M), providing the probabilities to belong either to torpor or to euthermic

rest (i.e. value specific state membership probabilities). Independent of the pre-assignment,

M values higher than the predicted M_e or lower than the predicted M_t are automatically

assigned to M_e and M_t, respectively. This latter automatic procedure is, however, only

performed if at least one M measurement has been assigned to torpor during that second

step. As defined previously, M values located above T_{lc} are automatically assigned to M_{TNZ}.

Automatically assigned values get a state membership probability of one.

Third step: Metabolic rate (M) measurements final assignment and estimation

of the functions parameters

In the third step, only M values situated between the predicted M_e and M_t are assigned.

Measurements' states that are automatically assigned during the second step are provided

along with the coupled M and T_a values. During that final step, the parameters of the functions

relating M to T_a below the T_{lc}, as well as the standard deviation for the different physiological

states are modelled.

Bayesian parametrization

Prior distributions are defined either uninformatively or with biologically relevant limits.

Except when specified, all prior distributions are Gaussian with a mean of 0 and a precision

(i.e. $1/SD^2$) of 0.001 and are truncated based on the specified limits. Specifically, the prior

distribution of T_{lc} during the first step is constrained between T_{lc_low} and the maximal T_a

recorded within the dataset. The prior distribution of T_{be} is constrained between T_{lc} (T_{lc_low} for

the first step) and 50°C. The upper limit of the T_{bt} prior distribution satisfies two conditions.

First, T_t should be inferior to the lower limit of the 95% credible interval (CI) of T_{lc}. Second, the

ratio of conforming M_t values corresponding to body temperatures differing by 10°C (Q_{10})

should not exceed a value of 5 (Geiser, 1988b). These conditions are verified by fixing the

upper range of the prior distribution of T_{bt} as:

$\min \left(\dfrac{\text{lower 95CI } T_{lc} * \beta_t - TMR}{\beta_t} , T_{be} - \dfrac{M_{TNZ} * 2}{TMR} \right)$ (11)

The lower limit of the T_{bt} prior distribution is set at -5°C (Barnes, 1989). The upper limit of the

prior distribution of TMR is defined as 80% of the M_{TNZ} (Ruf and Geiser, 2015), while its lower

one is fixed at 0. For M_r, the prior distribution ranges from TMR to M_{TNZ}. An uninformative

Dirichlet distribution (i.e. two concentration values of 1) is used for the priors of the dataset

specific state membership fractions, which provide the value specific state membership

probabilities of each measurement (see Second step: Metabolic rate (M) measurements pre-

assignment, above). Finally, the prior distributions of the standard deviations for M_{TNZ} and

regulated/conforming metabolic rates (SD_{TNZ}, SD_r and SD_c) are uniform, with that of SD_r

constrained between 0 and 3, that of SD_c constrained between one fifth of the value of SD_r

and SD_r and that of SD_{TNZ} constrained between half the value of SD_r and SD_r.

Three different Markov chains are run during 50'000 iterations starting at initial values within

the range of parameter space. The initial convergence phase is excluded by dropping the first

30'000 iterations. Markov chains are thinned by a factor of 10 and the Brooks–Gelman–Rubin

criterion R̂ (Brooks and Gelman, 1998) is used to assess the convergence of chains (< 1.1). To

ease parameters' estimation, metabolic rate measurements are divided by their mean.

Identifiability of the parameters

The prior and posterior distribution overlap (PPO) enables the evaluation of the relevance of

some modelled parameters. Overlap values are obtained with the function "MCMCtrace"

from package "MCMCvis" (Youngflesh, 2018). Truncated Gaussian prior distributions are

provided with the function "rtruncnorm" from package "truncnorm" (Mersmann et al., 2018).

Assignment confidence

The M value is assigned to the physiological state having the highest specific state membership

probability. The assignment confidence represents the product of the highest specific state

membership probability and of the probability that T_a is above (for M_{TNZ}) or below (for M_t and

M_e) T_{lc}. That latter probability is calculated from the variation in the estimation of T_{lc} (cf. First

step: Estimation of M_{TNZ} and T_{lc}), as the value at T_a of the T_{lc} cumulative posterior distribution

function, or 1 minus that function respectively. A threshold proportion is then selected to

consider any assignment with a significantly lower assignment confidence as invalid. That

hypothesis is tested with a binomial test, function: "binom.test", with the alternative

hypothesis set as "greater". The significance level for the binomial test is set at 0.05. For the

present study, assignments with a confidence higher than 0.8 were considered as valid.

Data collection and method evaluation

We evaluated our method by applying it to 28 published sets of M data obtained by

respirometry. Mainly, we aimed at highlighting strengths and possible issues linked to the

application of our method to typical sets of measurements. Thus, we were specifically

interested in divergences between assignments made by our method and by the authors. The

26 species of small or medium-sized mammals investigated illustrate a diversity of metabolic

and thermal strategies ranging from permanent homeothermy to heterothermy including

shallow, daily torpor and deep, long-term hibernation. The selected datasets also differ in size

(Table 1). For each published set, we considered all metabolic data provided, except those

made above the upper critical temperature defined by the authors (see "the three-steps

Method" above). In 20 species (here considered "heterotherms", Table 1), rates of

metabolism measured at T_a's below the described thermoneutral zone were originally

assigned to either of two states, which were usually referred to as euthermy (or

normothermy) and torpor. In *Cynopterus brachyotis*, values corresponding to particularly low

T_b were identified but not explicitly referred to as torpor (McNab, 1989). Some of the values

obtained from torpid *Lasiurus seminolus* by Genoud (1993) were originally characterised by

"irregular fluctuations" in M (Genoud, 1993). For simplicity, we treated these values as if the

authors assigned them to torpor. The remaining six species are hereafter referred to as

homeotherms (Table 1), as none of the values in the corresponding datasets were assigned to

torpor in the original studies. Chosen studies reported M_{TNZ} (mostly referred to as BMR) and

T_{lc} estimates, although the latter had to be extracted from a graph in two cases, or calculated

as the intercept between the euthermic line below thermal neutrality and M_{TNZ} provided by

the authors in another case (Table 1). In all dataset the authors provided a sufficiently precise

graph of metabolic rate values as a function of T_a and several of them also estimated T_t.

Plotted data were digitized using the software Plot Digitizer (Huwaldt and Steinhorst, 2015)

except for the *Pipistrellus pipistrellus* dataset, which was provided by MG. Sample sizes of the

digitized data ranged between 27 and 204 (Table 1). We tested the correlation between the

M_{TNZ}, T_{lc} and T_t values modelled with the present method and those provided by the authors

with Pearson's paired-samples correlation test, function "cor.test". Cases in which the

estimation of T_t fell outside the range of measured T_a were excluded from this analysis as T_t

was considered unidentifiable. Then, we examined whether the assignment of the metabolic

values to torpor or euthermy made by the authors and provided by the method coincided,

and we calculated a corroboration index as the fraction of the matched assignments. Only

measures with valid assignments (i.e. assignment confidence > 0.80, cf. "Assignment

confidence" in previous section) were considered for the correlation tests and the

corroboration index calculations. Finally, in order to define a PPO range highlighting an

identifiable parameter, the $PPOs$ of TMR from studies where part of the M measurements

were assigned to torpor were compared with those from studies where no M measurement

was assigned to torpor. In those later studies TMR was considered unidentifiable.

Acknowledgements

We are thankful to Philippe Christe and Nicolas Salamin for their support during the

elaboration of the model. We also thank four anonymous reviewers who contributed to

improve the original manuscript.

Competing interests

No competing interests declared

Funding

This study was financially supported by the Department of Ecology and Evolution of the

University of Lausanne (Switzerland) and by the Swiss National Science Foundation (grant

number: P2BEP3_168709 to NJF)

References

Barclay, R. M. R., Lausen, C. L. and Hollis, L. (2001). What's hot and what's not: defining torpor in free-ranging birds and mammals. *Can. J. Zool.* **79**, 1885–1890.

Barnes, B. M. (1989). Freeze Avoidance in a Mammal: Body Temperatures Below 0°C in an Arctic Hibernator. *Science.* **244**, 1593–1595.

Bartels, W., Law, B. S. and Geiser, F. (1998). Daily torpor and energetics in a tropical mammal, the northern blossom-bat *Macroglossus minimus* (Megachiroptera). *J. Comp. Physiol. - B Biochem. Syst. Environ. Physiol.* **168**, 233–239.

Bonaccorso, F. J. and McNab, B. K. (1997). Plasticity of Energetics in Blossom Bats (Pteropodidae): Impact on Distribution. *J. Mammal.* **78**, 1073–1088.

Boyles, J. G., Smit, B. and McKechnie, A. E. (2011). Does use of the torpor cut-off method to analyze variation in body temperature cause more problems than it solves? *J. Therm. Biol.* **36**, 373–375.

Boyles, J. G., Thompson, A. B., Mckechnie, A. E., Malan, E., Humphries, M. M. and Careau, V. (2013). A global heterothermic continuum in mammals. *Glob. Ecol. Biogeogr.* **22**, 1029–1039.

Boyles, J. G., Johnson, J. S., Blomberg, A. and Lilley, T. M. (2020). Optimal hibernation theory. *Mamm. Rev.* **50**, 91–100.

Brice, P. H., Grigg, G. C., Beard, L. A. and Donovan, J. A. (2002). Patterns of activity and inactivity in echidnas (*Tachyglossus aculeatus*) free-ranging in a hot dry climate: Correlates with ambient temperature, time of day and season. *Aust. J. Zool.* **50**, 461–475.

Brooks, S. P. and Gelman, A. (1998). General methods for monitoring convergence of iterative simulations. *J. Comput. Graph. Stat.* **7**, 434–455.

Coburn, D. K. and Geiser, F. (1998). Seasonal changes in energetics and torpor patterns in

the subtropical blossom-bat *Syconycteris australis* (Megachiroptera). *Oecologia* **113**,

467–473.

Daniels, H. L. (1984). Oxygen Consumption in *Lemur fulvus*: Deviation from the Ideal Model.

J. Mammal. **65**, 584–592.

Geiser, F. (1988a). Daily torpor and thermoregulation in <i>Antechinus<i/> (Marsupialia):

influence of body mass, season, development, reproduction, and sex. *Oecologia* **77**,

395–399.

Geiser, F. (1988b). Reduction of metabolism during hibernation and daily torpor in mammals

and birds: temperature effect or physiological inhibition? *J. Comp. Physiol. B* **158**, 25–

37.

Geiser, F. (2004). Metabolic Rate and Body Temperature Reduction During Hibernation and

Daily Torpor. *Annu. Rev. Physiol.* **66**, 239–274.

Geiser, F. (2011). Hibernation: Endotherms. In *eLS*, p. Chichester, UK: John Wiley & Sons, Ltd.

Geiser, F. and Brigham, R. M. (2012). The other functions of torpor. In *Living in a Seasonal*

World, pp. 109–121. Springer Berlin Heidelberg.

Geiser, F. and Kenagy, G. J. (1988). Torpor Duration in Relation to Temperature and

Metabolism in Hibernating Ground Squirrels. *Physiol. Zool.* **61**, 442–449.

Geiser, F. and Masters, P. (1994). Torpor in relation to reproduction in the mulgara,

Dasycercus cristicauda (Dasyuridae: Marsupialia). *J. Therm. Biol.* **19**, 33–40.

Geiser, F. and Ruf, T. (1995). Hibernation versus Daily Torpor in Mammals and Birds:

Physiological Variables and Classification of Torpor Patterns. *Physiol. Zool.* **68**, 935–966.

Geiser, F., Coburn, D. K., Kourtner, G. and Law, B. S. (1996). Thermoregulation, energy

metabolism, and torpor in blossom-bats, *Syconycteris australis* (Megachiroptera). *J.*

Zool. **239**, 583–590.

Genoud, M. (1993). Temperature regulation in subtropical tree bats. *Comp. Biochem.*

Physiol. -- Part A Physiol.

Genoud, M. (2014). Thermal energetics of the New-Guinean moss-forest rat (*Rattus niobe*)

in comparison with other tropical murid rodents. *J. Therm. Biol.* **41**, 95–103.

Genoud, M., Bonaccorso, F. J. and Anends, A. (1990). Rate of metabolism and temperature

regulation in two small tropical insectivorous bats (*peropteryx macrotis* and *Natalus*

tumidirostris). *Comp. Biochem. Physiol. -- Part A Physiol.* **97**, 229–234.

Genoud, M., Martin, R. D. and Glaser, D. (1997). Rate of metabolism in the smallest simian

primate, the pygmy marmoset (*Cebuella pygmaea*). *Am. J. Primatol.* **41**, 229–245.

Genoud, M., Isler, K. and Martin, R. D. (2018). Comparative analyses of basal rate of

metabolism in mammals: data selection does matter. *Biol. Rev.* **93**, 404–438.

Glazier, D. S. (2005). Beyond the "3/4-power law": Variation in the intra- and interspecific

scaling of metabolic rate in animals. *Biol. Rev. Camb. Philos. Soc.* **80**, 611–662.

Grigg, G. C., Augee, M. L. and Beard, L. A. (1992). Thermal Relations Of Free-Living Echidnas

During Activity And In Hibernation In A Cold Climate. *Platypus and Echidnas* 160–173.

Grimpo, K., Legler, K., Heldmaier, G. and Exner, C. (2013). That's hot: Golden spiny mice

display torpor even at high ambient temperatures. *J. Comp. Physiol. B Biochem. Syst.*

Environ. Physiol. **183**, 567–581.

Guppy, M. and Withers, P. (1999). Metabolic depression in animals: Physiological

perspectives and biochemical generalizations. *Biol. Rev.* **74**, 1–40.

Hainsworth, F. R. and Wolf, L. L. (1970). Regulation of oxygen consumption and body

temperature during torpor in a hummingbird, *Eulampis jugularis*. *Science (80-.).* **168**,

368–369.

Heldmaier, G., Steinlechner, S., Ruf, T., Wiesinger, H. and Klingenspor, M. (1989).

Photoperiod and Thermoregulation in Vertebrates: Body Temperature Rhythms and

Thermogenic Acclimation. *J. Biol. Rhythms* **4**, 139–153.

Heldmaier, G., Ortmann, S. and Elvert, R. (2004). Natural hypometabolism during

hibernation and daily torpor in mammals. *Respir. Physiol. Neurobiol.* **141**, 317–329.

Hosken, D. J. and Withers, P. C. (1999). Metabolic Physiology of Euthermic and Torpid Lesser

Long-Eared Bats, *Nyctophilus geoffroyi* (Chiroptera: Vespertilionidae). *J. Mammal.* **80**,

42–52.

Hudson, J. W. and Scott, I. M. (1979). Daily Torpor in the Laboratory Mouse, *Mus musculus*

Var. Albino. *Physiol. Zool.* **52**, 205–218.

Humphries, M. M., Thomas, D. W. and Speakman, J. R. (2002). Climate-mediated energetic

constraints on the distribution of hibernating mammals. *Nature* **418**, 313–316.

Huwaldt, J. A. and Steinhorst, S. (2015). Plot Digitizer, version 2.6.8.

Kelm, D. H. and von Helversen, O. (2007). How to budget metabolic energy: Torpor in a

small Neotropical mammal. *J. Comp. Physiol. B Biochem. Syst. Environ. Physiol.* **177**,

667–677.

Levesque, D. L. (2008). Seasonal changes in behavioural and thermoregulatory responses to

hypoxia in the eastern chipmunk (*Tamias striatus*).

Levesque, D. L. and Lovegrove, B. G. (2014). Increased homeothermy during reproduction in

a basal placental mammal. *J. Exp. Biol.* **217**, 1535–1542.

Lovegrove, B. G. (2012). The evolution of mammalian body temperature : the Cenozoic

supraendothermic pulses. 579–589.

Lovegrove, B. G. and Raman, J. (1998). Torpor patterns in the pouched mouse (*Saccostomus*

campestris; Rodentia): A model animal for unpredictable environments. *J. Comp.*

Physiol. - B Biochem. Syst. Environ. Physiol. **168**, 303–312.

Lyman, C. P. (1982). Entering hibernation. In *Hibernation and Torpor in Mammals and Birds*

596   (ed. Lyman, C. P.), Willis, J. S.), Malan, A.), and H., W. L. C.), pp. 37–53. Academic Press.

Lyman, C. P., Willis, J. S., Malan, A. and Wang, L. C. H. (1982). *Hibernation and Torpor in*

598   *Mammals and Birds*. New York, NY: Academic Press.

McKechnie, A. E. and Lovegrove, B. G. (2002). Avian Facultative Hypothermic Responses: a

600   Review. *Condor* **104**, 705.

McKechnie, A. E., Ashdown, R. A. M., Christian, M. B. and Brigham, R. M. (2007). Torpor in

602   an African caprimulgid, the freckled nightjar Caprimulgus tristigma. *J. Avian Biol.* **38**,

603   261–266.

McNab, B. K. (1980a). On Estimating Thermal Conductance in Endotherms. *Physiol. Zool.* **53**,

605   145–156.

McNab, B. K. (1980b). Energetics and the Limits to a Temperate Distribution in Armadillos. *J.*

607   *Mammal.* **61**, 606–627.

McNab, B. K. (1988). Energy Conservation in a Tree-Kangaroo (*Dendrolagus matschiei*) and

609   the Red Panda (*Ailurus fulgens*). *Physiol. Zool.* **61**, 280–292.

McNab, B. K. (1989). Temperature Regulation and Rate of Metabolism in Three Bornean

611   Bats. *J. Mammal.* **70**, 153–161.

McNab, B. K. (1997). On the utility of uniformity in the definition of basal rate of

613   metabolism. *Physiol. Zool.* **70**, 718–720.

McNab, B. K. (2002). *The Physiological Ecology of Vertebrates. A View from Energetics*.

615   Cumstock Publishing Associates, Cornell University Press, Ithaca & London.

Mersmann, O., Trautmann, H., Steuer, D. and Bornkamp, B. (2018). truncnorm: Truncated

Normal Distribution, package version 1.0-8.

Morris, S., Curtin, A. L. and Thompson, M. B. (1994). Heterothermy, torpor, respiratory gas

exchange, water balance and the effect of feeding in Gould's long-eared bat

Nyctophilus gouldi. *J. Exp. Biol.* **197**, 309 LP – 335.

Mzilikazi, N. and Lovegrove, B. G. (2002). Reproductive activity influences thermoregulation

and torpor in pouched mice, *Saccostomus campestris*. *J. Comp. Physiol. B Biochem. Syst.*

Environ. Physiol. **172**, 7–16.

Nespolo, R. F., Arim, M. and Bozinovic, F. (2003). Body size as a latent variable in a

structural equation model: Thermal acclimation and energetics of the leaf-eared

mouse. *J. Exp. Biol.* **206**, 2145–2157.

Nowack, J., Stawski, C. and Geiser, F. (2017). More functions of torpor and their roles in a

changing world. *J. Comp. Physiol. B Biochem. Syst. Environ. Physiol.* **187**, 889–897.

Nowack, J., Levesque, D. L., Reher, S. and Dausmann, K. H. (2020). Variable Climates Lead to

Varying Phenotypes : " Weird " Mammalian Torpor and Lessons From Non-Holarctic

Species. **8**,.

R Development Core Team (2012). R: A language and environment for statistical computing.

Reher, S., Ehlers, J., Rabarison, H. and Dausmann, K. H. (2018). Short and hyperthermic

torpor responses in the Malagasy bat *Macronycteris commersoni* reveal a broader

hypometabolic scope in heterotherms. *J. Comp. Physiol. B Biochem. Syst. Environ.*

Physiol. **188**, 1015–1027.

Ruf, T. and Geiser, F. (2015). Daily torpor and hibernation in birds and mammals. *Biol. Rev.*

90, 891–926.

Sassi, P. L. and Novillo, A. (2015). Acclimating to thermal changes: Intraspecific variation in a

small mammal from the Andes Mountains. *Mamm. Biol.* **80**, 81–86.

Schleucher, E. and Withers, P. C. (2001). Re-evaluation of the allometry of wet thermal

conductance for birds. *Comp. Biochem. Physiol. - A Mol. Integr. Physiol.* **129**, 821–827.

Scholander, P. F., Hock, R., Walters, V., Johnson, F. and Irving, L. (1950). Heat regulation in

some arctic and tropical mammals and birds. *Biol. Bull.* **99**, 237–258.

Song, X., Körtner, G. and Geiser, F. (1995). Reduction of metabolic rate and

thermoregulation during daily torpor. *J. Comp. Physiol. B* **165**, 291–297.

Song, X., Körtner, G. and Geiser, F. (1997). Thermal relations of metabolic rate reduction in

a hibernating marsupial. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **273**, 2097–2104.

Sparti, A. and Genoud, M. (1989). Basal rate of metabolism and temperature regulation in

Sorex coronatus and S. minutus (Soricidae: Mammalia). *Comp. Biochem. Physiol. -- Part*

A Physiol.

Speakman, J. R. and Selman, C. (2003). Physical activity and resting metabolic rate. *Proc.*

Nutr. Soc. **62**, 621–634.

Speakman, J. R. and Thomas, D. W. (2003). Physiological ecology and energetics of bats. In

Bat Ecology (ed. Kunz, T. H.) and Fenton, M. B.), pp. 430–490. University of Chicago

Press.

van Breukelen, F. and Martin, S. L. (2015). The Hibernation Continuum: Physiological and

Molecular Aspects of Metabolic Plasticity in Mammals. *Physiology* **30**, 273–281.

Willis, C. K. R. (2007). An Energy-Based Body Temperature Threshold between Torpor and

Normothermia for Small Mammals. *Physiol. Biochem. Zool.* **80**, 643–651.

Willis, C. K. R., Lane, J. E., Liknes, E. T., Swanson, D. L. and Brigham, R. M. (2005). Thermal

energetics of female big brown bats (*Eptesicus fuscus*). *Can. J. Zool.* **83**, 871–879.

Withers, P. C., Cooper, C. E., Maloney, S. K., Bozinovic, F. and Cruz-Neto, A. P. (2016).

Ecological and Environmental Physiology of Mammals. Oxford University Press.

Youngflesh, C. (2018). MCMCvis: Tools to Visualize, Manipulate, and Summarize MCMC

Output. *J. Open Source Softw.* **3**, 640.

Zeileis, A. and Hothorn, T. (2002). Diagnostic Checking in Regression Relationshipslmtest

citation info. *R News* **2**, 7--10.

Table 1: Model parameters and corroboration index for 28 datasets on small or medium-sized mammals described as either heterotherms or homeotherms.

Species	Type	m	Authors M_{TNZ}	M units	Authors T_{lc}	Authors T_t	Authors	N	T_{lc}	M_{TNZ}	TMR	T_t	β_t	α_e	β_c	α_c	α_r	MR inhibit	T_{be}	T_{bt}	PPO TMR	PPO M_r	Valid assignments	Corroboration	M assigned to torpor?
Unit	-	g	M units	-	°C	°C	-	-	°C	M units	M units	°C	M units/°C	M units	log(M unit)/°C	M units	M units	%	°C	°C	%	%	-	-	-
Callithrix pygmaea	Homeotherm	153	98	mlO$_2$ h^{-1}	27.5 [8]	–	Genoud et al. 1997	27	29.15	102.97	NI	NI	-9.78	388.45	NI	NI	NI	NI	39.77	NI	89.80	86.80	0.63	1.00	No
Marmosa robinsoni	Heterotherm	122	0.8	mlO$_2$ g^{-1} h^{-1}	26.5	–	McNab 1978	38	28.73	0.77	NI	NI	-0.10	3.49	NI	NI	NI	NI	36.41	NI	77.70	92.10	0.68	1.00	No
Phenacomys intermedius	Homeotherm	21.5	3.13	mlO$_2$ g^{-1} h^{-1}	25.1 [6]	–	McNab 1992	75	25.48	3.09	NI	NI	-0.25	9.36	NI	NI	NI	NI	37.78	NI	80.00	94.60	0.85	1.00	No
Rattus niobe	Homeotherm	42.3	53.6	mlO$_2$ h^{-1}	26	–	Genoud 2014	62	27.56	56.18	NI	NI	-4.56	181.97	NI	NI	NI	NI	39.97	NI	84.90	89.00	0.76	0.98	No
Sorex coronatus	Homeotherm	9.1	5.7	mlO$_2$ g^{-1} h^{-1}	27.5	–	Sparti & Genoud 1989	33	22.61	5.94	NI	NI	-0.24	11.37	NI	NI	NI	NI	47.56	NI	95.20	97.20	0.97	0.81	No
Sorex minutus	Homeotherm	3.3	8.6	mlO$_2$ g^{-1} h^{-1}	27.5	–	Sparti & Genoud 1989	28	25.36	8.67	NI	NI	-0.40	18.98	NI	NI	NI	NI	46.99	NI	95.60	95.00	0.89	1.00	No
Tolypeutes matacus	Homeotherm	1160	0.181	mlO$_2$ g^{-1} h^{-1}	28	–	McNab 1980b	66	29.57	0.19	NI	NI	-0.04	1.24	NI	NI	NI	NI	33.94	NI	76.80	95.30	0.85	1.00	No
Cercartetus lepidus	Heterotherm	12.6	1.49	mlO$_2$ g^{-1} h^{-1}	28.8	5.3	Geiser 1987	57	29.01	1.54	0.11	4.16	-0.20	7.47	0.08	0.08	0.95	0.49	36.72	4.68	4.60	33.50	0.98	1.00	Yes
Cercartetus nanus	Heterotherm	36	0.66	mlO$_2$ g^{-1} h^{-1}	28.7	4.8	Song et al. 1997	171	26.15	0.64	0.06	2.86	-0.14	4.29	0.05	0.05	0.46	0.27	30.55	3.26	8.10	27.10	0.95	0.85	Yes
Cynopterus brachyotis	Heterotherm	37.4	1.27	mlO$_2$ g^{-1} h^{-1}	29.7 [6]	–	McNab 1989	51	29.08	1.25	0.74	25.23	-0.21	7.18	NI	NI	NI	NI	34.38	28.76	44.80	94.30	0.76	0.92	Yes
Dasyuroides byrnei [1]	Heterotherm	118.2	0.7	mlO$_2$ g^{-1} h^{-1}	28.3 [7]	15	Geiser & Baudinette 1987	75	26.13	0.64	0.38	12.97	-0.12	3.66	0.01	0.34	1.89	0.67	31.51	16.24	21.50	29.90	0.89	0.87	Yes
Eptesicus fuscus	Heterotherm	15	16.98	mlO$_2$ h^{-1}	26.7	0	Willis et al. 2005	29	27.46	6.69	0.61	NI	-2.81	83.39	NI	NI	NI	NI	29.66	NI	13.60	80.50	0.93	1.00	Yes
Lasiurus cinereus [2]	Heterotherm	23.5 [4]	9	mW g^{-1}	30	–	Cryan & Wolf 2003	48	26.87	8.95	1.67	8.68	-1.09	38.23	NI	NI	NI	NI	35.15	10.22	24.70	78.00	0.90	0.81	Yes
Lasiurus seminolus	Heterotherm	9.4 [5]	1.26 [5]	mlO$_2$ g^{-1} h^{-1}	32	5	Genoud 1993	204	30.69	1.43	0.12	6.42	-0.42	14.22	0.08	0.08	2.74	0.60	34.03	6.57	7.60	23.20	0.95	0.99	Yes
Macroglossus minimus	Heterotherm	16.3	1.29	mlO$_2$ g^{-1} h^{-1}	30.9	20.4	Bartels et al. 1998	46	30.54	0.90	0.47	20.38	-0.16	5.89	0.06	0.16	3.78	0.93	36.32	23.35	27.50	26.30	0.65	0.97	Yes
Macroglossus minimus [3]	Heterotherm	16.4	0.88	mlO$_2$ g^{-1} h^{-1}	31	21	Bonaccorso & McNab 1997	47	30.05	1.23	0.78	18.79	-0.22	7.75	NI	NI	NI	NI	35.78	22.39	30.50	79.50	0.91	1.00	Yes
Melonycteris melanops	Heterotherm	53.3	0.81	mlO$_2$ g^{-1} h^{-1}	28	–	Bonaccorso & McNab 1997	30	28.84	0.81	0.44	17.05	-0.13	4.47	NI	NI	NI	NI	35.00	20.53	64.60	87.30	0.83	1.00	Yes
Natalus tumidirostris	Heterotherm	5.4	1.54	mlO$_2$ g^{-1} h^{-1}	28	20	Genoud et al. 1990	30	27.22	1.46	0.82	19.00	-0.58	17.07	NI	NI	NI	NI	29.54	20.42	58.70	76.70	0.83	1.00	Yes
Nyctophilus geoffroyi	Heterotherm	8	1.42	mlO$_2$ g^{-1} h^{-1}	34	–	Hosken & Withers 1999	84	32.32	1.43	0.24	8.62	-0.27	10.12	0.05	0.18	2.56	0.52	37.60	9.53	23.70	65.60	1.00	1.00	Yes
Peropteryx macrotis	Heterotherm	5.1	2.31	mlO$_2$ g^{-1} h^{-1}	30.5	20	Genoud et al. 1990	49	35.01	2.50	1.80	20.36	-0.22	10.14	0.02	1.35	6.20	0.90	46.99	28.70	22.30	60.40	0.71	0.66	Yes
Pipistrellus pipistrellus	Heterotherm	4.9	7.6	mlO$_2$ h^{-1}	33.2	0	Genoud & Christe 2011	95	32.79	7.62	0.54	2.22	-3.01	106.17	0.08	0.46	7.20	0.80	35.32	2.40	6.40	19.60	1.00	1.00	Yes
Planigale gilesi	Heterotherm	8.3	1.43	mlO$_2$ g^{-1} h^{-1}	28.5 [8]	14	Geiser & Baudinette 1988	62	27.76	1.53	0.30	13.41	-0.37	11.65	0.07	0.13	5.19	0.55	31.93	14.22	19.90	34.40	0.89	1.00	Yes
Setifer setosus	Heterotherm	530	0.231	mlO$_2$ g^{-1} h^{-1}	28	–	McNab 1980b	56	29.37	0.23	0.14	NI	-0.03	1.18	0.01	0.14	NI	0.88	36.20	NI	34.70	56.60	0.77	0.95	Yes
Sminthopsis macroura	Heterotherm	24.8	0.89	mlO$_2$ g^{-1} h^{-1}	31.3	14	Song et al. 1995	155	30.80	0.84	0.18	16.39	-0.19	6.63	0.09	0.05	3.27	0.77	35.24	17.36	16.50	30.60	0.97	1.00	Yes
Steatomys pratensis	Heterotherm	27.5	1.315	mlO$_2$ g^{-1} h^{-1}	28	20	Ellison 1995	39	26.73	1.24	0.26	16.38	-0.18	6.02	0.07	0.13	3.18	0.42	33.88	17.85	21.60	45.40	0.77	1.00	Yes
Syconycteris australis	Heterotherm	17.8	1.44	mlO$_2$ g^{-1} h^{-1}	29.5	18	Geiser et al. 1996	91	32.13	1.27	0.49	15.11	-0.14	5.78	0.05	0.24	2.61	0.90	41.31	18.62	21.60	28.60	0.88	0.95	Yes
Syconycteris australis [1]	Heterotherm	18	1.38	mlO$_2$ g^{-1} h^{-1}	30.2	18	Coburn & Geiser 1998	93	27.53	1.49	0.61	16.37	-0.17	6.00	0.07	0.23	3.33	0.86	36.22	20.07	32.00	43.10	0.85	0.97	Yes
Zaedyus pichiy	Heterotherm	1740	0.226	mlO$_2$ g^{-1} h^{-1}	28	–	McNab 1980b	57	31.82	0.23	0.15	22.49	-0.02	0.94	0.02	0.11	0.65	0.82	42.25	29.47	29.30	71.70	0.70	0.75	Yes

Subset of published data considered: [1]=summer group; [2]=males; [3]=lowland population.

m=average body mass; Autor's M_{TNZ}=author's estimation of metabolic rate within *TNZ*; M units=units for metabolic rate; Authors' T_{lc} and T_t=author's estimation of lower critical temperature and threshold ambient temperature for torpor, respectively. When m, M_{TNZ} or T_{lc} were not explicitly provided, they were: [4]=back-calculated from M_{TNZ} in % of expected from body mass; [5]=calculated as a weighted average of two seasonal groups; [6]=extracted from a graph; [7]=calculated as the intercept of the euthermic line and M_{TNZ} given by the authors; [8]=obtained as the

679   mean of a given range; N=number of digitized data points (sample size); Model parameters are defined in text. NI=Parameter considered as not
680   identifiable.
681

Figure 1: Representation of the relation between M during rest and torpor and T_a on which the present method is based. Torpor is indicated in blue, euthermic rest below the *TNZ* in red and rest within the *TNZ* in black. The relation follows the classical Scholander-Irving model (Scholander, Hock, Walters, Johnson, & Irving, 1950; for a discussion see McNab, 2002), which was later extended to include torpor (Geiser, 2011; Hainsworth and Wolf, 1970; Humphries et al., 2002; Speakman and Thomas, 2003). We further consider the possible occurrence of additional metabolic inhibition (Geiser, 2004; Geiser and Kenagy, 1988; Withers et al., 2016), hence allow the torpor curve to reach T_{lc} at a level equal to – or below M_{TNZ}.

Figure 2: Four datasets of *M* measured at different ambient temperature (*T_a*). Values were assigned to torpor (blue), euthermy (red) and M_{tnz} (black) using the presented three-step method. Predicted values: median and 95% credible intervals are represented by continuous and segmented lines, respectively. Invalid assignments are highlighted with a cross and mismatches between authors and model assignments are surrounded with black circle. A/ Perfect corroboration between model and authors assignments in a heterotherm: *Nyctophilus geoffroyi* (Hosken and Withers, 1999). B/ Absence of mismatched assignment between model and authors assignments with some invalid assignments in a homeotherm: *Sorex minutus* (Sparti and Genoud, 1989). C/ Presence of

mismatched or invalid assignments: *Peropteryx macrotis* (Genoud et al., 1990). D/ Absence of mismatched assignment between model and authors assignments with some invalid assignments, but insufficient number of torpor values to identify some torpor function parameters: *Melonycteris melanops* (Bonaccorso & McNab, 1997).

Figure 3: Frequency distribution of all the measurements from the 28 considered studies in relation to the difference between the method's estimated T_{lc} and the experimental T_a (N=1898). Invalid assignments (i.e. assignment confidence < 0.8) are represented in yellow and assignments mismatches in orange and blue. Mismatches caused by the difference between the model's estimated T_{lc} and that of the authors are highlighted in orange and the remaining mismatches in blue.