Environmental specificity in *Drosophila*-bacteria symbiosis affects host developmental plasticity

Robin Guilhot[1], Antoine Rombaut[1], Anne Xuéreb[1], Kate Howell[2], Simon Fellous[1]

[1]INRA, Centre for Biology and Management of Populations (UMR CBGP), 34988 Montferrier-sur-Lez, France

[2]Faculty of Veterinary and Agricultural Sciences, University of Melbourne, Parkville, Vic 3010, Australia

Abstract

Environmentally acquired microbial symbionts could contribute to host adaptation to local conditions like vertically transmitted symbionts do. This scenario necessitates symbionts to have different effects in different environments. We investigated this idea in *Drosophila melanogaster*, a species which communities of bacterial symbionts vary greatly among environments. We isolated four bacterial strains isolated from the feces of a *D. melanogaster* laboratory strain and tested their effects in two conditions: the ancestral environment (i.e. the laboratory medium) and a new environment (i.e. fresh fruit with live yeast). All bacterial effects on larval and adult traits differed among environments, ranging from very beneficial to marginally deleterious. The joint analysis of larval development speed and adult size further shows bacteria affected developmental plasticity more than resource acquisition. This effect was largely driven by the opposite effects of the bacteria in each environment. Our study reveals that understanding *D. melanogaster* symbiotic interactions in the wild necessitates using ecologically realistic conditions. Besides, context-dependent effects of symbionts, and their underlying mechanisms, shed light on how environmentally acquired symbionts may contribute to host evolution.

Introduction

Symbiosis may contribute to host evolution through recruitment of beneficial microorganisms (Margulis & Fester 1991; Jaenike et al. 2010; Fellous et al. 2011). As the environment varies among localities, different symbionts may be most beneficial in different conditions, possibly explaining microbiota variation among populations of the same animal species (e.g. Chandler et al. 2011; McKenzie et al. 2017). Microbial symbionts may therefore contribute to local adaptation (Kawecki and Ebert 2004). Most studies exploring symbiont-mediated local adaptation have focused on vertically transmitted microorganisms (e.g. Moran et al. 2008). However, numerous animals form symbioses with bacteria that are in part acquired from the environment either by horizontal transmission between hosts or recruitment of free-living strains (Ebert 2013). In this context, little is known on the how microbial effects on host fitness change with environmental conditions (Schwab et al. 2016; Callens et al. 2016), a necessary condition for symbiont-mediated local adaptation (Kawecki and Ebert 2004). Here, we explore how the effects of extracellular symbiotic bacteria on *Drosophila melanogaster* traits change when host and bacteria are studied in an environment different from their ancestral one.

The *Drosophila melanogaster* fly, is a prevalent model organism for host-microbiota studies (Douglas 2018). In *Drosophila melanogaster*, bacterial symbionts contribute to a broad range of functions including resource acquisition, digestion, immunity and behavior (Broderick and Lemaitre 2012; Ankrah and Douglas 2018; Schretter et al. 2018). Several laboratory studies have established fly nutrition relies on interactions with gut bacteria (Shin et al. 2011; Storelli et al. 2011; Ridley et al. 2012; Wong et al. 2014; Huang et al. 2015; Leitão-Gonçalves et al. 2017; Téfit et al. 2017). In particular, bacterial genera frequently associated with laboratory flies, such as *Lactobacillus* and *Acetobacter*, can improve larval growth and development

when laboratory food is poor in proteins (Shin et al. 2011; Storelli et al. 2011; Téfit et al.

2017). Even though some bacterial taxa are frequent in laboratory colonies, the composition

of *Drosophila* bacterial gut communities largely varies among laboratories (Chandler et al.

2011; Staubach et al. 2013; Wong et al. 2013; Vacchini et al. 2017). Studies have shown that

bacterial microbiota composition is determined by laboratory conditions more than

Drosophila species (Chandler et al. 2011; Staubach et al. 2013), demonstrating these

symbionts are largely acquired from fly environment. Empirical studies have nonetheless

shown pseudo-vertical transmission of bacteria from mothers to offspring also occurs in the

laboratory (Bakula 1969; Ridley et al. 2012; Wong et al. 2015; Téfit et al. 2018). Microbiota

composition differences between laboratory and field flies have led authors to argue that

symbiotic phenomena as observed in the laboratory may not reflect those occurring in natural

conditions (Chandler et al. 2011; Winans et al. 2017). Numerous variables differ between

laboratory and natural environments of *D. melanogaster* flies. A substantial difference is the

composition of the nutritive substrate upon which the adults feed, copulate, oviposit and

within which larvae develop. Wild flies live on and in fresh or decaying fruit flesh, usually

colonized by yeast, whereas laboratory flies are reared on an artificial, jellified and

homogeneous diet that contains long-chained carbohydrates (e.g. starch), agar, preservatives

and dead yeast cells or yeast extract. To this date, very few studies have investigated

Drosophila-bacteria interactions in conditions comparable to those of the field. How much

Drosophila-bacteria interactions that occur in the laboratory are maintained in natural

substrate remains largely undescribed.

Here, we experimentally studied the symbiosis between a laboratory strain of *D.*

melanogaster and four bacterial symbionts (isolated from its feces) in the ancestral laboratory

medium and in a new environment (grape berry) where we reproduced natural egg and

bacterial deposition from mothers. After inoculating bacteria-free eggs with these four

bacterial isolates, we scored various phenotypic fly traits at the larval and adult stages. We investigated two questions. (1) We focused on the influence of environmental variation on bacterial effects analyzing each of the host's traits individually. Our aim was to unveil whether host-symbiont that occurred in the environment of origin (i.e. the laboratory) maintained in more realistic conditions. We further relate these observations to fly and microbe ecology. (2) We performed a new, simultaneous analysis of two traits in order to disentangle symbionts' effects on host developmental plasticity and resource acquisition, two non-excluding possibilities. Separating plasticity from resource acquisition is important for at least two reasons. First, long-term symbiotic associations would be more likely when symbionts provide new capabilities (i.e. resources) than when they affect quantitative traits (Fellous and Salvaudon 2009) or their plasticity (Chevin et al 2010). Second, recent literature shows that the evolution of symbiont transmission depends on which of host's traits it affects (Brown and Akçay 2019); importantly, this mathematical model is based on the plastic trade-off between survival and reproduction. Here, we based our analysis on another well-established trade-off in holometabolous insects, that between duration of larval development and adult size at emergence (Teder et al. 2014; Nunney 1996). In brief, we reasoned that bacterial effects on host developmental plasticity would move host phenotypes along the trade-off axis, while resource acquisition would allow faster development or larger size without detrimental effects on the other trait (see Materials and Methods for details).

Materials and Methods

Drosophila strain

Insects were from the Oregon-R *Drosophila melanogaster* strain that was founded in 1927 and has since been maintained in numerous laboratories. Our sub-strain was funded ±2 years earlier from a few dozen individuals provided by colleagues. They had been reared on a laboratory medium comprising banana, sugar, dead yeast, agar and a preservative (Table S1A). Before and during the experiment reported here, all insects were maintained at 21 °C (stocks) or 23 °C (experiment), with 70 % humidity and a 14 h photoperiod.

Microbial isolates

The starting point of this work was to isolate and cultivate symbiotic bacteria from the flies. These bacteria were chosen for their ease of cultivation and our ability to discriminate them morphologically on standard microbiological medium. Our aim was not to sample the whole community of bacteria associated with our flies stock but to carry out tractable experiments using a random subset of their symbionts. Note our isolation method favored aerobic bacteria and excluded the anaerobes *Lactobacillus* spp., some of the best known symbionts of *D. melanogaster*. However, all the bacterial strains we isolated had already been identified as associated to *Drosophila* flies (Chandler et al. 2011; Staubach et al. 2013). Available literature did point to a number of taxa which interactions with *Drosophila* flies are described, and that we could have sourced from other laboratories. However, working with strains we could readily isolate from our fly colony meant we were certain to investigate fly-bacteria associations in their environment of origin.

In order to isolate bacteria present in fly feces, several groups of twenty *Drosophila* *melanogaster* flies were placed in sterile glass vials for 1 h. After fly removal, vials were washed with sterile PBS (Phosphate-Buffered Saline) solution, which was then plated on Lysogeny Broth (LB) agar medium (Table S1B) and incubated at 24 °C. Four bacterial morphotypes of variable frequency were chosen based on visible and repeatable differences in size, color, general shape and transparency during repeated sub-culturing on fresh media (Figure S2). A single colony of each morphotype was amplified in liquid LB medium in aerobic conditions at 24 °C for 72 h, centrifuged and washed in PBS. Several sub-samples of equal concentration were stored at -80 °C in PBS with 15 % glycerol and further used for molecular identification and the main experiment (one per experimental block).

Molecular identification of each bacterium was carried out by Sanger sequencing. To this aim, a fresh colony of each bacterial type was picked with a sterile toothpick and dipped into sterile water, then boiled 10 min at 95 °C (Mastercycler, Eppendorf) and cooled in ice water. A sterile toothpick dipped into sterile water served as sterility control of the process. Fragments of the 16sRNA gene were amplified with bacterial primers Y2MOD (5-ACTYCTACGGRAGGCAGCAGTRGG-3') and 16SB1 (5'-TACGGYTACCTTGTTACGACTT-3') (Haynes et al. 2003; Carletto et al. 2008). PCRs were performed in a volume of 25 µl, containing each primer at 0.2 µM, 1x buffer (containing 2 mM MgCl$_2$), each dNTP at 0.2 mM, and 1 U of *DreamTaq* Taq (Thermo Scientific). PCRs cycles had an initial denaturation step at 95 °C for 15 min, followed by ten cycles at 94 °C / 40 s - 65 °C / 45 s – 72 °C / 45 s); followed by 30 cycles at 94 °C / 40 s – 55 °C / 45 s – 72 °C / 45 s; and finished with an extension step of 10 min at 72 °C. Negative PCR controls were included. PCR products were visualized under UV light in an agarose gel before sequencing. Consensus sequences were created with CodonCode Aligner 4.2.7. Online SINA alignment service (https://www.arb-silva.de/aligner/) (Pruesse et al. 2012) and NCBI GenBank blastn

service (https://blast.ncbi.nlm.nih.gov/Blast.cgi) were used to compare and assign the

sequences. The four bacteria were identified as a *Staphylococcus* (likely *S. xylosus*), an

Enterococcus (likely *E. faecalis*), an Enterobacteriaceae and an Actinobacteria (likely

Brevibacterium). Further in this article, theses bacteria are referred to as *Staphylococcus*,

Enterococcus, Enterobacteriaceae and Actinobacteria, respectively. All sequences were

deposited in the NCBI database under the accession numbers MK461976 (*Staphylococcus*),

MK461977 (*Enterococcus*), MK461978 (Enterobacteriaceae) and MK461979

(Actinobacteria).

A wild isolate of *Saccharomyces cerevisiae* yeast was used in experiments where larvae

developed in fresh grape berries. The yeast was isolated from a wild Drosophilid in a vineyard

in Southern France (*'Le Domaine de l'Hortus'*, Hérault, France) (see Hoang et al. (2015) for a

balanced discussion on *Drosophila-Saccharomyces* interactions). The isolate was grown in

YPD medium, washed, split into several samples, stored at -80 °C in sterile PBS with 15 %

glycerol, that were further used in the experiment (one per block).

158 **Experimental design**

Flies were associated with bacteria following a full-factorial design that resulted in twelve

different treatments. There were two types of fly environments: laboratory medium (the

ancestral environment, see Table S1A for composition) and grape berries (the new

environment, white grapes, unknown cultivar). We had six different symbiont treatments:

each of the four bacterial strains described above, a mix of the four bacteria and controls

without bacteria (Figure 1). Each treatment had 13 to 15 replicates organized in 15 blocks

launched over four days. Bacterial growth was also studied in fly-free grapes but is not

described here.

Figure 1: summary of the experimental design and the measured traits. T0: association of
Drosophila eggs with bacteria (β1, β2, β3, β4), bacterial mixture (M), or nothing (Ø), in the
two environments: laboratory medium (L) or grape berry (G) inoculated with live yeast (Y).
T+5: larval traits scoring after five days. T+n: adult size scoring on a randomly chosen subset
of adults from each replicate. T+x: analysis of the microbial content of the larval environment
two days after the end of pupal formation.

Grape berries were surface-sterilized in a 2 % bleach solution before use. Because *D.*
melanogaster females only oviposit in wounded fruit, we incised 5 mm of berry skin (Figure
S4) where we deposited twenty eggs free from culturable bacteria. These eggs were produced
by the oviposition of flies on laboratory medium supplemented with the antibiotic
streptomycin (1 mg / ml in 1 mM EDTA, Sigma-Aldrich ref. 85886). The efficacy of this
method for removing culturable bacteria from egg surface was confirmed by the lack of
bacterial growth after the deposition of such eggs onto LB agar plates (note however that
these conditions were not suitable for detection of anaerobic bacteria such as *Lactobacillus*).
Grape berries were inoculated with live yeast cells as it is a key component (Begg &

Robertson 1948; Becher et al. 2012) and was necessary for fly survival in our system (Figure

S3). For treatments with laboratory diet we deposited 20 eggs free from culturable bacteria on

incisions at the surface of 4 ml of medium placed in 2 cm * 2 cm plastic cubes. Berries and

laboratory media were all placed in 75 ml plastic vials closed by a foam plug.

Bacterial cells were inoculated to laboratory medium and grape berry immediately before egg

deposition. Single bacterial strain treatments received 10^4 live bacterial cells, and the mixed

treatment 2.5 x 10^3 cells of each bacterium (i.e. 10^4 cells in total), suspended in 10 µl of sterile

PBS. The number of inoculated bacterial cells, that is $< 10^4$ Colony Forming Units (CFUs),

was chosen based on the average number of bacteria previously reported in the guts of

second-instar larvae (Bakula 1969; Storelli et al. 2011). In control treatments, sterile PBS was

deposited instead of bacteria. On grape berries, 10^4 live cells of the yeast *Saccharomyces*

cerevisiae were inoculated. Note fruit substrate and live yeast presence are confounded factors

in our experiment because we did not intend to study the effect of live yeast onto larval

growth (Becher et al 2012) but to mimic field conditions where larvae develop in presence of

live yeast. Although the laboratory medium also contains yeast, cells are killed during

industrial production (Table S1A).

Fly phenotyping

We scored six different phenotypic traits in larvae and adults: larval size, mouthpart

movement rate, and visible number on medium surface after 5 days; survival rate to adult

emergence; time until adult emergence and a proxy of adult size. Larval mouthpart movement

speed was the number of back-and-forth movements of the mouthpart that could be observed

in 5 seconds. Newly formed pupae were transferred to empty sterile vials daily. We recorded

male and female emergences daily.

The size of adults, and their microbial content (see below), were estimated on a subset of

those that emerged from each vial. For each vial, one pupa was chosen randomly and all

adults that emerged on the same day as the focal pupa were collected, pooled by sex. These

pools were homogenized in 200 µl of sterile PBS using a sterile pestle, splat in two sub-

samples and stored at -80 °C with 15% sterile glycerol. One of the two sub-samples was used

to numerate live bacteria and yeast cells in newly emerged adults, the other one to estimate

adult size with the spectrophotometric method described in Fellous et al. (2018). We chose

this method as it allowed the simultaneous analysis of adult size and microbial content.

Briefly, we used log-transformed optical density at 202 nm of fly homogenate as a proxy of

adult size. This was measured several months after the experiment when samples were

thawed, crushed a second time using a Tissue Lyser II (Qiagen) for 30 s at 30 Hz with Ø3 mm

glass balls, centrifuged for 30 s at 2000 G. Optical density of 15 µL of supernatant was then

read on a Multiskan GO spectrometer (Thermo Scientific). This metrics correlates in both

males and females with wet weight and wing length (all $R^2 > 0.8$), two frequently used size

proxies in *Drosophila* studies. For figures and analyses of adult size we used the Log of

observed optical density divided by the number of individuals in the sample.

Analysis of bacterial presence and metabolism

We tested the presence of inoculated bacteria and yeast in substrates two days after the

appearance of the last pupa. Samples were analyzed by plating homogenates on LB agar

medium and incubated at 24 °C. In this manuscript we only report on the presence or absence

of inoculated bacteria in the larval substrate. Data of microorganism presence and numbers in

emerging adults will be reported separately.

The Enterobacteriaceae and the Actinobacteria were the main bacterial strains that affected fly

phenotypes. In order to shed light on the ecologies of these two strains and therefore on their

effects on hosts, we analyzed their metabolic capabilities with Eco Microplates (Biolog) (see

Text S5 for methodological details).

Bacteria and fungi morphologically different from those we had inoculated were observed in

samples from 17 % of the vials (either in adults on in the environment). Data from these vials

were excluded for all analyzes presented here. Both datasets are available in the open data

repository Zenodo (DOI: 10.5281/zenodo.2554194).

Statistical analyses

Individual traits

To study the response of each fly phenotypic trait to variation of larval substrate and bacterial

symbiont, we used linear mixed models (LMM) with Restricted Maximum Estimate

Likelihood (REML). 'Block identity' was defined as random factor. Fixed factors were the

'larval environment' (i.e. laboratory medium or fruit), 'bacterial treatment', 'fly sex' (for the

analyses of age at emergence and adult size only), and their full-factorial interactions. A

Backward, stepwise model selection was used to eliminate non-significant terms from initial,

full models. Homoscedasticity and residuals normality visually complied with model

assumptions. When the 'bacteria*environment' interaction was significant, and to investigate

hypotheses based on the visual observation of the data, we used independent contrasts to test

significant differences between bacterial treatments and controls from the same environment.

Joint effect of bacteria on adult age and size at emergence

The aim of this analysis was to study how bacteria affected simultaneously speed of larval

development and size. Importantly, we needed to discard the general effect of the nutritive

environment to single out the effects of the symbionts. Indeed, if one environment was

generally more favorable than the other, main environmental effects could create a positive

relationship between the two traits that would conceal how bacteria affect them. To this end,

all analyses were carried out after subtracting the mean trait value of the controls (i.e.

bacteria-free) in the relevant environment from the trait values of each combination of

bacteria and environment. In other words, data presented in Figures 5 and S6 represent the

incremental effects of the bacteria on host traits after removal of the transversal influence of

the nutritive substrate.

We carried out two types of analyses. (1) In order to unveil the overall pattern (Figures 5 and

S6) we worked with mean treatment effects (i.e. one single data point per treatment, two when

sex was taken into account) and univariate regressions. Because of the significant interaction

between sex, bacteria and environment for adult size, our initial analysis separated males from

females (Figure S6). However, the linear regression of size onto developmental speed was not

significantly different among sexes (Interaction Sex*Speed: $F_{1.16} = 2.93$, p = 0.11). Presented

results hence merge observations from males and females. (2) In order to explain the factors

behind the simultaneous effect of bacteria on developmental speed and adult size we carried

out a multivariate analysis of variance (MANOVA) using all data points (i.e. one data point

per experimental unit). MANOVA was chosen because it enables studying how factors affect

several variables jointly, in other words it considers effects onto the correlation between

several variables (Zar 2009, p.319). We used a "repeated measures" personality of MANOVA

and reported the tests based on the Sum response function (i.e. a M-matrix that is a single

vector of 1 s; between-subject report in JMP). Model contained the factors 'bacterial

treatment', 'environment' and their interaction. Homoscedasticity and residuals normality

visually complied with MANOVA assumptions. The dataset used for the MANOVA analysis

is available in the open data repository Zenodo (DOI: 10.5281/zenodo.3352230).

Analyzes were performed with JMP (SAS, 14.1).

Results

Effects of bacteria on individual traits reveal extensive environmental-dependence

Larval size after five days was influenced by an interaction between the environment and the bacterial treatment (Table 1, Figure 2A). In grapes, addition of the Actinobacteria decreased larval size relative to bacteria-free controls but had no effect in laboratory media. In laboratory media, addition of the Enterobacteriaceae alone or in mixture with the other bacterial strains produced larger larvae than bacteria-free controls (contrast 'Enterobacteriaceae treatment' vs 'Control treatment': $F_{1.90} = 28.92$, $p < 0.0001$), which did not happen when grown on a grape substrate (contrast 'Enterobacteriaceae treatment' vs 'Control treatment': $F_{1.86} = 0.92$, $p = 0.3405$) (Figure 2A).

The number of larvae visible on medium surface was influenced by an interaction between the environment and the bacterial treatment (Table 1, Figure 2B). In laboratory media, addition of the Enterobacteriaceae alone or in mixture with the other bacterial strains led to greater numbers of visible larvae compared to bacteria-free controls (contrast 'Enterobacteriaceae treatment' vs 'Control treatment': $F_{1.131} = 20.40$, $p < 0.0001$; contrast 'Mixture treatment' vs 'Control treatment': $F_{1.131} = 6.98$, $p = 0.0092$), which did not happen when grown on a grape substrate (contrast 'Enterobacteriaceae treatment' vs 'Control treatment': $F_{1.131} = 1.63$, $p = 0.2036$; contrast 'Mixture treatment' vs 'Control treatment': $F_{1.131} = 0.93$, $p = 0.3355$) (Figure 2B).

Mouthparts movement rate was influenced by an interaction between the environment and the bacterial treatment (Table 1, Figure 2C). Movements were generally faster in grapes than in laboratory media. However, addition of the Actinobacteria slowed down the movements of

mouthparts in grapes to a level comparable to the one of larvae reared on laboratory media

(contrast 'Actinobacteria treatment' vs 'Control treatment': $F_{1.99} = 4.54$, $p = 0.0355$) (Figure

2C).

The proportion of eggs surviving until the adult stage was only affected by the environment,

with a lower survival in grapes than in laboratory media (Table 1, Figure 2D).

Table 1: analysis of larval and adult phenotypes in response to bacterial treatment and larval environment. Linear mixed models (REML).

Response variables Factors	Median larval size	Number of larvae on the substrate surface	Larval foraging behavior	Developmental survival	Average age of emerging adults	Adult size proxy
Environment	$F_{1.18} = 137.51$ $p < 0.0001$	$F_{1.98} = 13.64$ $p = 0.0004$	$F_{1.25} = 28.43$ $p < 0.0001$	$F_{1.17} = 27.02$ $p < 0.0001$	$F_{1.18} = 77.86$ $p < 0.0001$	$F_{1.15} = 0.35$ $p = 0.5630$
Bacterial treatment	$F_{5.88} = 4.08$ $p = 0.0022$	$F_{5.131} = 2.02$ $p = 0.0806$	$F_{5.97} = 0.78$ $p = 0.5657$	$F_{5.115} = 0.78$ $p = 0.5688$	$F_{5.211} = 4.41$ $p = 0.0008$	$F_{5.183} = 0.79$ $p = 0.5609$
Environment*Bacterial treatment	$F_{5.88} = 4.64$ $p = 0.0008$	$F_{5.131} = 4.50$ $p = 0.0008$	$F_{5.97} = 2.80$ $p = 0.0211$	$F_{5.115} = 0.53$ $p = 0.7558$	$F_{5.211} = 7.85$ $p < 0.0001$	$F_{5.183} = 1.90$ $p = 0.0960$
Fly sex	-	-	-	-	$F_{1.199} = 1.68$ $p = 0.1961$	$F_{1.166} = 3.27$ $p = 0.0724$
Environment*Bacterial treatment*Fly sex	-	-	-	-	$F_{5.199} = 0.42$ $p = 0.8336$	$F_{5.166} = 2.75$ $p = 0.0204$

312 Linear mixed models, with block as random factor.

Figure 2: larval phenotypes in response to bacterial treatment and larval environment.

(A) Median larval size after 5 day; (B) Number of larvae on the medium surface after 5 days; (C) Number of larval mouthparts movements per 5 s observed after 5 days; (D) Survival from egg to adult. Symbols indicate means; error bars indicate standard errors around the mean. Stars (*) indicate treatments significantly different from controls in the same environment (post-hoc contrasts, $\alpha = 0.05$).

Age at adult emergence was not different among sexes but was influenced by an interaction between the environment and the bacterial treatment (Table 1, Figure 3). In laboratory media, flies reared with the Enterobacteriaceae, alone or in mixture, emerged nearly two days sooner

than bacteria-free flies in the same environment and almost four days earlier than bacteria-free

flies in grapes (contrast 'Enterobacteriaceae treatment' vs 'Control treatment': $F_{1.225} = 27.57$,

p < 0.0001; contrast 'Mixture treatment' vs 'Control treatment': $F_{1.225} = 24.94$, p < 0.0001)

(Figure 3). In grapes, flies reared with the bacterial mixture emerged one day later than

bacteria-free flies (contrast 'Mixture treatment' vs 'Control treatment': $F_{1.222} = 6.42$, p =

0.0120) (Figure 3).

Figure 3: age of _Drosophila_ adults at emergence in response to bacterial treatment and

larval environment. Symbols indicate means; error bars indicate standard errors around the

mean. Stars (*) indicate treatments significantly different from controls in the same

environment (post-hoc contrasts, α = 0.05).

Adult size was influenced by the triple interaction between sex, the environment and the

bacterial treatment (Table 1, Figure 4). Several bacterial treatments had sex-specific effects

that differed among the two environments. For example, inoculation of the mixture of the four

bacteria produced larger males than females in grapes (contrast 'Mixture treatment' vs

'Control treatment': $F_{1.166} = 5.30$, p = 0.0225), but smaller males than females in laboratory

media (contrast 'Mixture treatment' vs 'Control treatment': $F_{1.167} = 4.79$, p = 0.0300) (Figure

4). Similarly, inoculation of the *Staphylococcus* or *Enterococcus* led to larger males than

females in grape (contrast '*Staphylococcus* treatment' vs 'Control treatment': $F_{1.164} = 4.97$, p

= 0.0271; contrast '*Enterococcus* treatment' vs 'Control treatment': $F_{1.164} = 7.48$, p = 0.0069),

but no difference in laboratory medium (contrast '*Staphylococcus* treatment' vs 'Control

treatment': $F_{1.165} = 0.11$, p = 0.7367; contrast '*Enterococcus* treatment' vs 'Control

treatment': $F_{1.167} = 0.66$, p = 0.4182) (Figure 4).

Figure 4: *Drosophila* adult size proxy in response to bacterial treatment and larval

environment. Symbols indicate means; error bars indicate standard errors around the mean.

Stars (*) indicate significant differences between males and females in the same environment

(post-hoc contrasts, α = 0.05).

Joint analysis of adult age and size at emergence suggests bacteria affect host developmental plasticity along a trade-off

We expected three possible patterns when plotting average adult size in function of speed of

larval development (i.e. - age at emergence): a positive relationship indicative of a similar

effect of the bacteria on the two traits (i.e. bacteria modulate resource acquisition, mostly); a

negative relationship indicative of bacteria affecting host position along the trade-off (i.e.

bacteria modulate developmental plasticity, mostly); a lack of correlation that would have

been challenging to interpret on its own as several processes could produce this result (e.g.

bacterial effects on both host plasticity and resource acquisition).

The relationship between effects of bacteria on adult age and size at emergence was

marginally significant and negative (Linear model $F_{1.8}$ = 8.83, p = 0.018) (Figure 5). A

MANOVA shed light on the relative influence of the environment and the bacterial treatments

on the correlated effect of the treatments on the two traits (Table 2). It revealed the

environment was an important factor: in laboratory medium, addition of bacteria accelerated

development relative to controls at the cost of producing smaller adults; in grape addition of

bacteria slowed down development relative to controls but emerging adults were large (Figure

5). There was no significant main effect of the bacterial treatments but a significant

interaction with the environment, which confirms the bacterial treatments had different effects

on host phenotype in each environment.

Table 2: Multivariate Analysis of Variance of the joint effect of the bacteria on 'Age at emergence' and 'Adult size'. As in Figure 5, general effects of the environments were removed by subtracting trait values of controls (i.e. without bacterial addition) in each environment before carrying out the analysis.

Factor	*F*	*d.f.*	*p*
Environment	14.9	1.85	0.0002
Bacterial treatment	1.65	4.85	0.17
Environment*Bacterial treatment	3.86	4.85	0.006

384 **Figure 5: relationship between bacterial effects on age of emerging adults and bacterial**

385 **effects on adult size.** Effects of bacteria for each treatment were calculated by subtracting the

386 mean trait value of controls in the same environment to mean trait value of the treatment.

387 Error bars indicate standard errors around the means.

Bacterial occurrence in the environment and their metabolism

The Enterobacteriaceae isolate was the only bacterium to be consistently retrieved from the

laboratory medium in which larvae had developed (Figure 6). In one instance, the

Actinobacteria was found in a grape berry from which no live adult fly emerged (Figure 6).

The physiological profile of the Enterobacteriaceae revealed growth of the bacterium in a

broad panel of carbon sources (Figure S5A). The physiological profile of the Actinobacteria

revealed substantial growth of the bacterium on the carbon sources Pyruvic Acid Methyl Ester

and Tween 80 only (Figure S5B).

Figure 6: proportion of larval environments that contained the inoculated bacterium after fly pupation. Proportions were calculated over replicates (see after): Lab. (Laboratory medium) - Control (n = 12 replicates), Lab. - Staphyl. (n = 11), Lab. - Enteroc. (n = 7), Lab. - Enterob. (n = 10), Lab. - Actinob. (n = 10), Lab. - Mixture (n = 13), Grape - Control (n = 26), Grape - Staphyl. (n = 16), Grape - Enteroc. (n = 16), Grape - Enterob. (n = 13), Grape - Actinob. (n = 16), Grape - Mixture (n = 12).

Discussion

We studied the symbiotic interactions between a laboratory strain of *Drosophila melanogaster* and four bacterial strains isolated from its feces. Our results show different effects of bacterial symbionts on host phenotype in laboratory medium and in real fruit. All symbiont effects were environment-dependent, some of which may be explained by the ecology of laboratory-associated symbionts in artificial medium. The joint analysis of larval developmental speed and adult size further suggests bacteria influence host developmental plasticity along the well-known physiological trade-off between these two traits.

Different symbiont effects in different environments

The observation that all bacterial effects on host phenotype were different in laboratory medium and grape berry prompts the question of the reason behind this discrepancy. Focusing of the Enterobacteriaceae may shed light onto the ecologies of the symbiotic bacteria we isolated, and why they differed among environments.

In laboratory medium, inoculation of the Enterobacteriaceae induced greater larval size and accelerated larval development (Figures 2A and 3). Besides, adults produced by larvae associated with the Enterobacteriaceae in laboratory medium were not significantly smaller than adults produced by bacteria-free larvae (Figure 4). The bacterium hence accelerated larval growth. In its presence larvae could be observed in greater numbers at the surface of the medium than in the absence of the bacterium (Figure 2B), even though there were no mortality differences among Enterobacteriaceae-associated and bacteria-free larvae (Figure 2D). The Enterobacteriaceae was also the only bacterium to be retrieved from the medium after fly pupation (Figure 6). These elements may be explained by two mechanisms. (1) The

bacterium could serve as food and be grazed on medium surface by foraging larvae. The

phenomenon would be similar to that described by Yamada et al. (2015) where the yeast

Issatchenkia orientalis extracts amino acids from agar-based laboratory medium and

concentrates them on medium surface where adult flies harvest them. This hypothesis is

congruent with the visual observation that media inoculated with the Enterobacteriaceae

harbored white microbial growth on their surface (Figure S7). Along these lines, the wide

metabolic spectrum of this bacterium (Figure S5A) suggests the microorganism is a generalist

that would extract resources from the medium, possibly transform nutrients (Ankrah and

Douglas 2018; Sannino et al. 2018), and eventually concentrate them on medium surface. (2)

Microbial growth at the surface would interfere with larval development in such way that

larvae would remain at the surface. This behavior could also trigger accelerated development

if excessive microbial growth revealed detrimental. The two hypotheses above are non-

excluding; the joint-analysis of developmental speed and adult size sheds further light on this

question (see below). Why did the effect of the Enterobacteriaceae on host phenotype differ

among environments? The physical nature of laboratory medium is very different from that of

real fruit. In particular, the agar of laboratory medium permits the diffusion of simple

nutrients and their absorption by bacteria and yeast present on surface. Besides, in grape

nutrients are not free to diffuse but enclosed in cells. Surface growth is therefore more likely

in artificial medium than in grape berry, leading to different effects on larval development.

In addition to physical differences between laboratory medium and fresh fruit, the nature and

concentration of available nutrient are likely to differ. It is well known that lactic and acetic

acid bacteria, two taxa that were not investigated in our experiment, can promote larval

growth upon nutrient scarcity (Shin et al. 2011; Storelli et al. 2011, Téfit et al. 2017).

However, it is also well established that bacteria can affect *Drosophila* phenotype through

signaling (Storelli et al. 2011) as well as nutrient provisioning (Brownlie et al. 2009; Bing et

al. 2018; Sannino et al. 2018). In most cases, these effects which were described from laboratory flies and in laboratory medium, are condition specific (Douglas 2018). Indeed, bacteria are often only beneficial when laboratory food has a low concentration in dead yeast (i.e. amino acids) (Shin et al. 2011; Storelli et al. 2011). Along these lines, it may seem paradoxical the Enterobacteriaceae only accelerates larval growth in rich laboratory medium rather than in grape berry (unless the bacterium synthesized a rare nutrient). Metabolic profiling (Figure S5A) further shows the Enterobacteriaceae is a generalist bacterium able to grow on a variety of substrate. However, the Actinobacteria had a narrower metabolic spectrum (Figure S5B), suggesting it is a specialist which growth largely depends on the availability of specific nutrients. The bacterium slowed down larval growth in grape (Figure 2A) for an unknown reason - maybe because it exerted additional stress onto larvae in a relatively poor medium - but had no notable effect in laboratory medium. The environment-specific effect of the Actinobacteria compares to previous reports of *Drosophila* symbionts being beneficial in some environments only (e.g. *Lactobacillus plantarum* in rich medium), and further reveals that bacteria with little effect in an environment can become detrimental in new conditions.

Effects of bacteria on host developmental plasticity

In holometabolous insects, the duration of larval development and adult size are often negatively correlated due to a physiological trade-off: faster development reduces the duration of food intake and leads to smaller adult size (Teder et al. 2014; Nunney 1996). We propose to exploit this trade-off to separate symbionts' effects on host developmental plasticity and resource acquisition. As discussed above, symbionts of *Drosophila* flies can modify host's signaling (e.g. Shin et al. 2011; Storelli et al. 2011) as well as provide rare resources (e.g.

Brownlie et al. 2009; Sannino et al. 2018). These two mechanisms are expected to have

different effects on the trade-off between speed of development and size. Signaling (i.e.

plasticity) should move hosts along the trade-off, while the provisioning of greater resources

should enable faster growth and/or larger size without sacrificing the other trait. In order to

test this idea we extracted bacterial effects on host phenotype by subtracting control trait

values to those of each of the bacterial treatments in each environment. The resulting plot of

symbionts effects on developmental speed and adult size (Figures 5 and S6) reveals the

influence of the bacteria on the host independent of the general effects of the environment (i.e.

those not due to the bacteria).

Our original analysis of bacterial effects on larval development and adult size revealed a

negative relationship (Figures 5 and S6). Treatments that accelerated development produced

small adults and treatments that slowed down development produced large adults. The

negative relationship came as a surprise as we expected nutritional symbionts to affect

developmental speed and adult size either in an independent or similar fashion, which would

have led to an absence or a positive relationship between these two traits, respectively.

Instead, results suggest bacterial treatments influenced host development plastically along the

trade-off between speed of development and adult size. This result contrasts with previous

reports on other *Drosophila* symbionts that induce positive relationships between larval and

adult traits (Anagnostou et al. 2010; Bing et al. 2018). For example, the yeast species

Metschnikowia pulcherrima produces small adults that are also slow to develop (Anagnostou

et al. 2010). Different symbionts in different contexts can therefore affect host developmental

plasticity or its resource acquisition.

The visual examination of Figure 5 shows bacterial effects measured in laboratory medium

(blue points) group in the fast development-small size region of phenotypic space, while

effects in grape (red points) occur in the small speed-large size side of the trade-off. This

suggests that the environment could determine whether bacteria accelerate development (at

the cost of a smaller size) or favor size (at the cost of a slower development). A MANOVA

revealed a significant effect of the environment on the joint analysis of the two traits, hence

confirming that bacterial influence on host developmental plasticity is largely determined by

the environment. With only 5 bacterial treatments per environment it was not possible to test

if bacteria affect host development along the trade-off within a single environment.

Whether microbial symbionts influence hosts through effects on developmental plasticity or

resource availability (i.e. general vigor *sensus* Fry (1993)) may change the evolutionary fate

of the host-symbiont relationship. First, symbionts that plastically alter phenotypes would be

more dispensable that those providing functions host genomes are not capable of (Fellous and

Salvaudon 2009). Besides, it could be argued that the fitness effects of alternative plastic

strategies may depend on the environmental context more than general improvement of

resource availability (Chevin et al 2010). Therefore, symbionts that improve general

performance of the host through greater resource availability may be more likely to be fixed

among host individuals and populations than those that affect plasticity. By contrast, hosts

may dynamically acquire and lose symbionts which effects on fitness depend on the

environment, paving the way for the evolution of facultative symbiosis. Along these lines,

recent modelling of host-symbiont dynamics revealed that whether symbionts affect adult

survival or reproduction (another well-known trade-off between fitness components)

determines transmission mode evolution (Brown and Akçay 2019). Our experimental study

only considered one trade-off between two developmental traits, possibly overlooking effects

on other fitness components. Future analyses should increase in dimensionality and consider a

greater number of fitness components. Similarly, a precise description of the slopes and

shapes of considered trade-offs will be necessary to discriminate simultaneous effects of

symbionts on plasticity and resource acquisition. We are now pursuing further investigation to

determine if, and when, bacterial and yeast symbionts affect host developmental plasticity

rather than resource availability in *Drosophila* flies.

Context-dependent effects of bacteria enable symbiont-mediated adaptation

A consequence of *Drosophila* bacterial symbionts having different effects in different

environments is the possibility they contribute to the fine-tuning of host phenotype to local

conditions (Margulis & Fester 1991; Moran 2007; Sudakaran et al. 2017). The phenomenon is

well established in vertically transmitted symbionts of insects that protect their hosts from

parasites. For example, populations of aphids exposed to parasitoids harbor protective

Hamiltonella symbionts at greater frequency than parasitoid-free populations (Oliver et al.

2005). Similarly, in the fly *Drosophila neotestacea*, the spread of the bacterium *Spiroplasma*

allowed hosts to evolve greater resistance to parasitic nematodes (Jaenike et al. 2010).

Vertically transmitted bacterial symbionts of *Paramecium* ciliates can also improve host

resistance to stressful conditions (Hori & Fujishima 2003). Whether bacteria act as parasites

or mutualists depends partly on the genetic ability of the host to deal with stress in absence of

the symbiont (Duncan et al. 2010). However, the evolutionary role of symbionts that may be

acquired from the environment is less clear, in part because the mechanisms favoring the

association of hosts with locally beneficial symbionts are not as straightforward as for vertical

transmission (Ebert 2013). Nonetheless, several lines of evidence suggest environmentally

acquired microbial symbionts may contribute to local adaptation in *Drosophila*-microbe

symbiosis. First, symbionts can be transmitted across metamorphosis (i.e. transstadial

transmission from the larval to the adult stage) and pseudo-vertically during oviposition (i.e.

from mothers to offspring) (Bakula 1969; Starmer et al. 1988; Spencer 1992; Ridley et al.

2012; Wong et al. 2015; Téfit et al. 2018). Second, host immune system participates in the

destruction of harmful gut bacteria and the retention of beneficial ones (Lee et al. 2017; Lee et

al. 2018). Third, *Drosophila* larvae may preferentially associate with beneficial yeast species

ensuring they engage in symbiosis with locally adequate nutritional symbionts (Fogleman et

al. 1981; Fogleman et al. 1982). In addition to host genetic and preferential association with

beneficial microbes, *Drosophila* adaptation to local conditions thanks to microorganisms

further necessitates symbionts have different effects in different environments. Our results

show bacteria isolated from a fly population have different effects on host phenotype

depending on the substrate larvae were reared in (Figures 2, 3, 4 and 5). As a consequence,

extracellular bacterial symbionts may therefore contribute to *Drosophila* local adaptation

through variations in symbiont community composition.

Conclusion

In this study, we found that associations between laboratory *Drosophila* flies and their microbial symbionts result in different effects on host phenotype when the symbiosis is investigated under conditions comparable to nature. The context-dependency of bacterial effects, and the underlying mechanisms we unveiled (i.e. bacterial ecology and bacterial effects on host plasticity), shed light on the role of microorganisms in the evolution of their hosts. Our results point out that in order to understand the ecology and evolution of symbiotic interactions in the wild it is necessary to use ecologically realistic conditions, which is attainable in the *Drosophila* system.

Acknowledgements

We warmly thank L. Benoit, M.P. Chapuis, J. Collet, D. Duneau, R. Gallet, P. Gautier, N. Kremer, F. Leulier and N. Rode for methodological help and useful comments on an earlier version of this work. We thank Wolf Blanckenhorn, Pedro Simões and an anonymous referee for precious comments on an earlier version of the manuscript, as well as PCI (Peer Community In) Evol Biol managing board for their contribution to the advancement of science publishing.

Conflict of interest disclosure

The authors of this preprint declare that they have no financial conflict of interest with the content of this article. Simon Fellous is one of the PCI (Peer Community In) Evol Biol recommenders.

Funding

This project was supported by French National Research Agency through the 'SWING' project (ANR-16-CE02-0015) and by Agropolis Fondation under the reference ID 1505-002 through the 'Investissements d'avenir' programme (Labex Agro:ANR-10-LABX-0001-01).

References

Anagnostou, C., Dorsch, M., and Rohlfs, M. (2010). Influence of dietary yeasts on

Drosophila melanogaster life-history traits. *Entomologia Experimentalis et Applicata*, 136(1),

1-11.

Ankrah, N. Y., and Douglas, A. E. (2018). Nutrient factories: metabolic function of beneficial

microorganisms associated with insects. *Environmental microbiology*. 20(6), 2002-2011.

Bakula, M. (1969). The persistence of a microbial flora during postembryogenesis of

Drosophila melanogaster. *Journal of invertebrate pathology*, 14(3), 365-374.

Becher, P. G., Flick, G., Rozpędowska, E., Schmidt, A., Hagman, A., Lebreton, S., Larsson,

M. C., Hansson, B. S., Piskur, J., Witzgall, P., and Bengtsson, M. (2012). Yeast, not fruit

volatiles mediate *Drosophila melanogaster* attraction, oviposition and

development. *Functional Ecology*, 26(4), 822-828.

Begg, M., and Robertson, F. W. (1948). Nutritional requirements of *Drosophila*

melanogaster. *Nature*, 161(4098), 769.

Bing, X., Gerlach, J., Loeb, G., and Buchon, N. (2018). Nutrient-Dependent Impact of

Microbes on *Drosophila suzukii* Development. *mBio*, 9(2), e02199-17.

Bresson, J., Varoquaux, F., Bontpart, T., Touraine, B., and Vile, D. (2013). The PGPR strain

Phyllobacterium brassicacearum STM196 induces a reproductive delay and physiological

changes that result in improved drought tolerance in *Arabidopsis*. *New Phytologist*, 200(2), 558-569.

Broderick, N. A., and Lemaitre, B. (2012). Gut-associated microbes of *Drosophila melanogaster*. *Gut microbes*, 3(4), 307-321.

Brown, A., and Akçay, E. (2019). Evolution of transmission mode in conditional mutualisms with spatial variation in symbiont quality. *Evolution*, 73(2), 128-144.

Brownlie, J. C., Cass, B. N., Riegler, M., Witsenburg, J. J., Iturbe-Ormaetxe, I., McGraw, E. A., and O'Neill, S. L. (2009). Evidence for metabolic provisioning by a common invertebrate endosymbiont, *Wolbachia pipientis*, during periods of nutritional stress. *PLoS pathogens*, 5(4), e1000368.

Callens, M., Macke, E., Muylaert, K., Bossier, P., Lievens, B., Waud, M., and Decaestecker, E. (2016). Food availability affects the strength of mutualistic host–microbiota interactions in *Daphnia magna*. *The ISME journal*, 10(4), 911-920.

Carletto, J., Gueguen, G., Fleury, F., and Vanlerberghe-Masutti, F. (2008). Screening the bacterial endosymbiotic community of sap-feeding insects by terminal-restriction fragment length polymorphism analysis. *Entomologia experimentalis et applicata*, 129(2), 228-234.

Cass, B. N., Himler, A. G., Bondy, E. C., Bergen, J. E., Fung, S. K., Kelly, S. E., and Hunter, M. S. (2016). Conditional fitness benefits of the *Rickettsia* bacterial symbiont in an insect pest. *Oecologia*, 180(1), 169-179.

Chandler, J. A., Lang, J. M., Bhatnagar, S., Eisen, J. A., and Kopp, A. (2011). Bacterial communities of diverse *Drosophila* species: ecological context of a host–microbe model system. *PLoS genetics*, 7(9), e1002272.

Chevin, L. M., Lande, R., and Mace, G. M. (2010). Adaptation, plasticity, and extinction in a changing environment: towards a predictive theory. *PLoS biology*, 8(4), e1000357.

Daskin, J. H., and Alford, R. A. (2012). Context-dependent symbioses and their potential roles in wildlife diseases. *Proceedings of the Royal Society of London B: Biological Sciences*, 279(1733), 1457-1465.

De Vries, E. J., Jacobs, G., Sabelis, M. W., Menken, S. B., and Breeuwer, J. A. (2004). Diet–dependent effects of gut bacteria on their insect host: the symbiosis of *Erwinia* sp. and western flower thrips. *Proceedings of the Royal Society of London B: Biological Sciences*, 271(1553), 2171-2178.

Douglas, A. E. (2018). The *Drosophila* model for microbiome research. *Lab Animal*, 47(6), 157.

Duncan, A. B., Fellous, S., Accot, R., Alart, M., Chantung Sobandi, K., Cosiaux, A., and Kaltz, O. (2010). Parasite-mediated protection against osmotic stress for *Paramecium caudatum* infected by *Holospora undulata* is host genotype specific. *FEMS microbiology ecology*, 74(2), 353-360.

Ebert, D. (2013). The epidemiology and evolution of symbionts with mixed-mode
transmission. *Annual Review of Ecology, Evolution, and Systematics*, 44, 623-643.

Fellous, S., and Salvaudon, L. (2009). How can your parasites become your allies? *Trends in
parasitology*, 25(2), 62-66.

Fellous, S., Duron, O., and Rousset, F. (2011). Adaptation due to symbionts and conflicts
between heritable agents of biological information. *Nature Reviews Genetics*, 12(9), 663.

Fellous, S., Guilhot, R., Xuéreb, A., and Rombaut, A. (2018). A high-throughput
spectrophotometric assay of adult size in *Drosophila* that facilitates microbial and
biochemical content analysis. *Dros. Inf. Serv.*, 101, 69-74.

Fogleman, J. C., Starmer, W. T., and Heed, W. B. (1981). Larval selectivity for yeast species
by *Drosophila mojavensis* in natural substrates. *Proceedings of the National Academy of
Sciences*, 78(7), 4435-4439.

Fogleman, J. C., Starmer, W. T., and Heed, W. B. (1982). Comparisons of yeast florae from
natural substrates and larval guts of southwestern *Drosophila*. *Oecologia*, 52(2), 187-191.

Fry, J. D. (1993). The "general vigor" problem: can antagonistic pleiotropy be detected when
genetic covariances are positive?. *Evolution*, 47(1), 327-333.

Haynes, S., Darby, A. C., Daniell, T. J., Webster, G., Van Veen, F. J. F., Godfray, H. C. J.,

Prosser, J. I., and Douglas, A. E. (2003). Diversity of bacteria associated with natural aphid

populations. *Applied and Environmental Microbiology*, 69(12), 7216-7223.

Hoang, D., Kopp, A., and Chandler, J. A. (2015). Interactions between *Drosophila* and its

natural yeast symbionts—Is *Saccharomyces cerevisiae* a good model for studying the fly-

yeast relationship?. *PeerJ*, 3, e1116.

Huang, J. H., and Douglas, A. E. (2015). Consumption of dietary sugar by gut bacteria

determines *Drosophila* lipid content. *Biology letters*, 11(9), 20150469.

Jaenike, J., Unckless, R., Cockburn, S. N., Boelio, L. M., and Perlman, S. J. (2010).

Adaptation via symbiosis: recent spread of a *Drosophila* defensive

symbiont. *Science*, 329(5988), 212-215.

Kawecki, T. J., and Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology letters*,

7(12), 1225-1241.

Lee, J. H., Lee, K. A., and Lee, W. J. (2017). Microbiota, gut physiology, and insect

immunity. *Advances in Insect Physiology*, 52, 111-138.

Lee, K. A., Cho, K. C., Kim, B., Jang, I. H., Nam, K., Kwon, Y. E., Kim, M., Hyeon, D. Y.,

Hwang, D., Seol, J. H., and Lee, W. J. (2018). Inflammation-modulated metabolic

reprogramming is required for DUOX-dependent gut immunity in *Drosophila*. *Cell host and*

microbe, 23(3), 338-352.

Leitão-Gonçalves, R., Carvalho-Santos, Z., Francisco, A. P., Fioreze, G. T., Anjos, M.,

Baltazar, C., Elias, A. P., Itskov, P. M., Piper., M. D. W., and Ribeiro, C. (2017). Commensal

bacteria and essential amino acids control food choice behavior and reproduction. *PLoS*

biology, 15(4), e2000862.

Ma, D., and Leulier, F. (2018). The importance of being persistent: The first true resident gut

symbiont in *Drosophila*. *PLoS biology*, 16(8), e2006945.

Margulis, L., and Fester, R., eds (1991). *Symbiosis as a source of evolutionary innovation:*

speciation and morphogenesis. MIT Press.

McKenzie, V. J., Song, S. J., Delsuc, F., Prest, T. L., Oliverio, A. M., Korpita, T. M., Alexiev,

A., Amato, K. R., Metcalf, J. L., Kowalewski, M., Avenant, N. L., Link, A., Di Fiore, A.,

Seguin-Orlando, A., Feh, C., Orlando, L., Mendelson, J. R., Sanders, J., and Knight, R.

(2017). The effects of captivity on the mammalian gut microbiome. *Integrative and*

comparative biology, 57(4), 690-704.

Moran, N. A. (2007). Symbiosis as an adaptive process and source of phenotypic complexity.

Proceedings of the National Academy of Sciences, 104, 8627-8633.

Moran, N. A., McCutcheon, J. P., and Nakabachi, A. (2008). Genomics and evolution of

heritable bacterial symbionts. *Annual review of genetics*, 42, 165-190.

Nunney, L. (1996). The response to selection for fast larval development in *Drosophila*

melanogaster and its effect on adult weight: an example of a fitness trade-off. *Evolution*,

50(3), 1193-1204.

Oliver, K. M., Moran, N. A., and Hunter, M. S. (2005). Variation in resistance to parasitism in

aphids is due to symbionts not host genotype. *Proceedings of the National Academy of*

Sciences, 102(36), 12795-12800.

Pais, I. S., Valente, R. S., Sporniak, M., and Teixeira, L. (2018). *Drosophila melanogaster*

establishes a species-specific mutualistic interaction with stable gut-colonizing bacteria. *PLoS*

biology, 16(7), e2005710.

Pruesse, E., Peplies, J., and Glöckner, F. O. (2012). SINA: accurate high-throughput multiple

sequence alignment of ribosomal RNA genes. *Bioinformatics*, 28(14), 1823-1829.

Ridley, E. V., Wong, A. C., Westmiller, S., and Douglas, A. E. (2012). Impact of the resident

microbiota on the nutritional phenotype of *Drosophila melanogaster*. *PloS one*, 7(5), e36765.

Sannino, D. R., Dobson, A. J., Edwards, K., Angert, E. R., and Buchon, N. (2018). The

Drosophila melanogaster gut microbiota provisions thiamine to its host. *mBio*, 9(2), e00155-

18.

Schretter, C. E., Vielmetter, J., Bartos, I., Marka, Z., Marka, S., Argade, S., and Mazmanian,

S. K. (2018). A gut microbial factor modulates locomotor behaviour in *Drosophila*. *Nature*,

563(7731), 402.

Schwab, D. B., Riggs, H. E., Newton, I. L., and Moczek, A. P. (2016). Developmental and

ecological benefits of the maternally transmitted microbiota in a dung beetle. *The American*

Naturalist, 188(6), 679-692.

Shin, S. C., Kim, S. H., You, H., Kim, B., Kim, A. C., Lee, K. A., Yoon, J. H., Ryu, J. H., and

Lee, W. J. (2011). *Drosophila* microbiome modulates host developmental and metabolic

homeostasis via insulin signaling. *Science*, 334(6056), 670-674.

Spencer, D. M., Spencer, J. F. T., De Figueroa, L., and Heluane, H. (1992). Yeasts associated

with rotting citrus fruits in Tucumán, Argentina. *Mycological Research*, 96(10), 891-892.

Starmer, W. T., Peris, F., and Fontdevila, A. (1988). The transmission of yeasts by

Drosophila buzzatii during courtship and mating. *Animal behaviour*, 36(6), 1691-1695.

Staubach, F., Baines, J. F., Künzel, S., Bik, E. M., and Petrov, D. A. (2013). Host species and

environmental effects on bacterial communities associated with *Drosophila* in the laboratory

and in the natural environment. *PloS one*, 8(8), e70749.

Storelli, G., Defaye, A., Erkosar, B., Hols, P., Royet, J., and Leulier, F. (2011). *Lactobacillus*

plantarum promotes *Drosophila* systemic growth by modulating hormonal signals through

TOR-dependent nutrient sensing. *Cell metabolism*, 14(3), 403-414.

Sudakaran, S., Kost, C., and Kaltenpoth, M. (2017). Symbiont acquisition and replacement as

a source of ecological innovation. *Trends in Microbiology*, 25(5), 375-390.

Teder, T., Vellau, H., and Tammaru, T. (2014). Age and size at maturity: A quantitative

review of diet-induced reaction norms in insects. *Evolution*, 68(11), 3217-3228.

Téfit, M. A., and Leulier, F. (2017). *Lactobacillus plantarum* favors the early emergence of fit

and fertile adult *Drosophila* upon chronic undernutrition. *Journal of Experimental Biology*,

220, 900-907.

Téfit, M. A., Gillet, B., Joncour, P., Hughes, S., and Leulier, F. (2018). Stable association of a

Drosophila-derived microbiota with its animal partner and the nutritional environment

throughout a fly population's life cycle. *Journal of insect physiology*, 106(1), 2-12.

Vacchini, V., Gonella, E., Crotti, E., Prosdocimi, E. M., Mazzetto, F., Chouaia, B., ... and

Daffonchio, D. (2017). Bacterial diversity shift determined by different diets in the gut of the

spotted wing fly *Drosophila suzukii* is primarily reflected on acetic acid

bacteria. *Environmental microbiology reports*, 9(2), 91-103.

Winans, N. J., Walter, A., Chouaia, B., Chaston, J. M., Douglas, A. E., and Newell, P. D.

(2017). A genomic investigation of ecological differentiation between free-living and

Drosophila-associated bacteria. *Molecular ecology*, 26(17), 4536-4550.

Wong, A. C., Chaston, J. M., and Douglas, A. E. (2013). The inconstant gut microbiota of

Drosophila species revealed by 16S rRNA gene analysis. *The ISME journal*, 7(10), 1922-

1932.

Wong, A. C. N., Dobson, A. J., and Douglas, A. E. (2014). Gut microbiota dictates the

metabolic response of *Drosophila* to diet. *Journal of Experimental Biology*, 217(11), 1894-

1901.

Wong, A. C. N., Luo, Y., Jing, X., Franzenburg, S., Bost, A., and Douglas, A. E. (2015). The

host as the driver of the microbiota in the gut and external environment of *Drosophila*

melanogaster. *Applied and environmental microbiology*, 81(18), 6232-6240.

Yamada, R., Deshpande, S. A., Bruce, K. D., Mak, E. M., and William, W. J. (2015).

Microbes promote amino acid harvest to rescue undernutrition in *Drosophila*. *Cell*

reports, 10(6), 865-872.

Zar, J. H. (2009). *Biostatistical Analysis Fifth Edition*. Upper Saddle River, New Jersey:

Pearson.

Supplementary Material 1. Laboratory recipes.

Table S1A: laboratory medium recipe.

Component	*Amount for 1.5L*
Reverse osmosis water	1200 ml
Banana	280 g
Sugar	74 g
Dead yeast	74 g
Alcohol	30 ml
Agar	12 g
Nipagin	6 g

832   **Table S1B:** lysogeny broth (LB) recipes.

Component	Quantity / Volume for			
	Liquid LB	*Agar LB*	*Anti-bacteria Agar LB*	*Anti-yeast Agar LB*
Reverse osmosis water	1000 ml	1000 ml	1000 ml	1000 ml
Proteose peptone n°3 (Conda)	10 g	10 g	10 g	10 g
Yeast extract (Merck)	5 g	5 g	5 g	5 g
NaCl (Carlo Erba)	5 g	5 g	5 g	5 g
European Bacteriological Agar (Conda)		15 g	15 g	15 g
Ampicillin (Sigma) (pure)			100 mg	
Chloramphenicol (Sigma) (100 mg/ml in ethanol)			10 mg	
Cycloheximide (Sigma) (100 mg/ml in DMSO)				1 mg

Supplementary Material 2. Bacterial strains isolated from Oregon-R *Drosophila melanogaster* and used in the experiment.

Figure S2: bacterial strains isolated from Oregon-R *Drosophila melanogaster* and used in the experiment. (a) *Staphylococcus* sp.; (b) *Enterococcus* sp.; (c) Enterobacteriaceae.; (d) Actinobacteria.

Supplementary Material 3. Live yeast as a prerequisite to *D. melanogaster* larvae survival on pristine grape berry.

Figure S3: live yeast is necessary for the survival of *D. melanogaster* larvae on pristine grape berry. Prior to the experiment, we investigated survival of *D. melanogaster* larvae on fresh grape berries. Twenty bacteria-free *D. melanogaster* eggs were deposited next to an artificial wound with or without the bacterial isolates and *Saccharomyces cerevisiae*. In absence of yeast, larvae died quickly after hatching, with or without bacteria (Figure S1a). When live yeast was added to the system, numerous larvae developed up to the 3[thrd] instar (Figure S1b), when we stopped monitoring.

Supplementary Material 4. Experimental design for the grape

berry environment.

857 **Figure S4: experimental design for the grape berry environment.** (a) Experimental block

858 for grape berry treatments, (b) Experimental unit with grape berry, (c) Decaying grape berry

859 with live yeast, bacteria and larvae, (d) Egg cases visible near berry incision and active larvae

860 in fruit flesh.

Supplementary Material 5. Bacterial physiological profiles.

Text S5:

Eco Microplates (Biolog) were used to have an overview of the metabolic 'fingerprint' of the Enterobacteriaceae, the Actinobacteria isolate and the Actinobacteria variant. A fixed number of fresh bacteria cells suspended in sterile PBS were inoculated in well with one of 31 different carbon sources. Each combination Bacterial isolate*Carbon source was replicated three times. The plates were incubated at 25 °C and the absorbance at 595 nm was measured with a Multiskan GO spectrometer (Thermo Scientific) after 48 h and 120 h. A tetrazolium dye included with each carbon source entrained the production of red color when bacterial respiration occurred, i.e. when the carbon source was used. Variations of red color among carbon sources allowed establishing a physiological profile of each bacterial isolate.

Carbon sources

Figure S5A: physiological profile of the Enterobacteriaceae isolate after (A) 48 h- and (B) 120 h-long exposure to different carbon sources. Symbols indicate means; error bars indicate standard errors around the mean. X-axis labels correspond to abbreviations of tested carbon sources, with 2-Hyd for 2-Hydroxy Benzoic Acid; 4-Hyd for 4-Hydroxy Benzoic Acid; a-Cyc for α-Cyclodextrin; a-D-Lac for α-D-Lactose; a-Ket for α-Ketobutyric Acid; b-Met for β-Methyl-D-Glucoside; D-Cel for D-Cellobiose; D-Galacto for D-Galactonic Acid γ -Lactone; D-Galactu for D-Galacturonic Acid; D-Glu for D-Glucosaminic Acid; D-Mal for D-Malic Acid; D-Man for D-Mannitol; D-Xyl for D-Xylose; D L-a-Gly for D,L-α-Glycerol Phosphate; Glu for Glucose-1-Phosphate; Glyco for Glycogen; Glycy for Glycyl-L-Glutamic Acid; i-Ery for i-Erythritol; Ita for Itaconic Acid; L-Arg for L-Arginine; L-Asp for L-Asparagine; L-Phe for L-Phenylalanine; L-Ser for L-Serine; L-Thr for L-Threonine; N-Ace for N-Acetyl-D-Glucosamine; Phe for Phenylethylamine; Put for Putrescine; Pyr for Pyruvic Acid Methyl Ester; Tw40 for Tween 40, Tw80 for Tween 80, Wat for Water and y-Hyd for γ-Hydroxybutyric Acid.

Figure S5B: physiological profile of the Actinobacteria isolate after (A) 48 h- and (B) 120

h-long exposure to different carbon sources. Symbols indicate means; error bars indicate

standard errors around the mean. X-axis labels correspond to abbreviations of tested carbon sources, with

2-Hyd for 2-Hydroxy Benzoic Acid; 4-Hyd for 4-Hydroxy Benzoic Acid; a-Cyc for α-Cyclodextrin; a-D-Lac for

α-D-Lactose; a-Ket for α-Ketobutyric Acid; b-Met for β-Methyl-D-Glucoside; D-Cel for D-Cellobiose; D-

Galacto for D-Galactonic Acid γ -Lactone; D-Galactu for D-Galacturonic Acid; D-Glu for D-Glucosaminic

Acid; D-Mal for D-Malic Acid; D-Man for D-Mannitol; D-Xyl for D-Xylose; D L-a-Gly for D,L-α-Glycerol

Phosphate; Glu for Glucose-1-Phosphate; Glyco for Glycogen; Glycy for Glycyl-L-Glutamic Acid; i-Ery for i-

Erythritol; Ita for Itaconic Acid; L-Arg for L-Arginine; L-Asp for L-Asparagine; L-Phe for L-Phenylalanine; L-

Ser for L-Serine; L-Thr for L-Threonine; N-Ace for N-Acetyl-D-Glucosamine; Phe for Phenylethylamine; Put

for Putrescine; Pyr for Pyruvic Acid Methyl Ester; Tw40 for Tween 40, Tw80 for Tween 80, Wat for Water and

y-Hyd for γ-Hydroxybutyric Acid.

Supplementary Material 6. Joint analysis of bacterial effects on adult age and size for each sex.

Figure S6: relationship between bacterial effects on age of emerging adults and bacterial effects on adult size, in females and males. As the linear regressions were not significantly different (Interaction Sex*Speed: $F_{1.16} = 2.93$, p = 0.11), data was pooled for the analysis reported in the main text. Symbols indicate the phenotype mean of each combination of bacterium and environment. Error bars mark the SE of the mean for both axes.

914 **Supplementary Material 7. Laboratory medium inoculated with**

915 **the Enterobacteriaceae.**

918 **Figure S7: bacterial growth at the surface of laboratory medium five days after**

919 **Enterobacteriaceae inoculation.** This picture was taken in absence of larvae, but similar

920 growth could be observed in their presence.