A mutation in *Hnrnph1* that decreases methamphetamine-induced reinforcement, reward, and dopamine release and increases synaptosomal hnRNP H and mitochondrial proteins

Abbreviated title: *Hnrnph1* in methamphetamine addiction traits

Qiu T. Ruan[1,2,3,^], Neema Yazdani[1,2,3,^], Benjamin C. Blum[4], Jacob A. Beierle[1,2,3], Weiwei Lin[4], Michal A. Coelho[5], Elissa K. Fultz[5], Aidan F. Healy[5], John R. Shahin[5], Amarpreet K. Kandola[2], Kimberly P. Luttik[2], Karen Zheng[2], Nathaniel J. Smith[2], Justin Cheung[2], Farzad Mortazavi[6], Daniel J. Apicco[7], Durairaj Ragu Varman[8], Sammanda Ramamoorthy[8], Peter E. A. Ash[7], Douglas L. Rosene[6,] Andrew Emili[4], Benjamin Wolozin[7], Karen K. Szumlinski[5], Camron D. Bryant[1,2,3,9]*

1. Biomolecular Pharmacology Training Program, Department of Pharmacology and Experimental Therapeutics, Boston University School of Medicine, 02118
2. Laboratory of Addiction Genetics, Department of Pharmacology and Experimental Therapeutics, Boston University School of Medicine, 02118
3. Transformative Training Program in Addiction Science, Boston University, 02118
4. Center for Network Systems Biology, Boston University School of Medicine, 02118
5. Department of Psychological and Brain Sciences, University of California, Santa Barbara, 93106
6. Department of Anatomy and Neurobiology, Boston University School of Medicine, 02118
7. Laboratory of Neurodegeneration, Department of Pharmacology and Experimental Therapeutics and Neurology, Boston University School of Medicine, 02118
8. Department of Pharmacology & Toxicology, Virginia Commonwealth University, 23284
9. Department of Psychiatry, Boston University School of Medicine, 02118

^Equally contributing co-first authors
*Corresponding Author
Camron D. Bryant, Ph.D.
Department of Pharmacology and Experimental Therapeutics and Department of Psychiatry
72 E. Concord St. L-616
Boston, MA 02118 USA
E: camron@bu.edu   P: (617) 358-9581

Pages: 49
Figures: 10
Tables: 2
Extended data figures: 15
Abstract: 249 words
Introduction: 588 words
Discussion: 1497 words
Conflict of interest statement: The authors declare no competing interests.

Acknowledgments: This study was supported by National Institutes of Health/National Institute on Drug Abuse (NIH/NIDA) Grant Nos. R00DA029635 (C.D.B.), R01DA039168 (C.D.B.), F31DA40324 (N.Y.), and N01DA-14-7788; NIH/National Institute of General Medical Sciences (NIGMS) for Grant No. T32GM008541 (N.Y., Q.T.R. and J.A.B.); and the Burroughs Wellcome Fund Transformative Training Program in Addiction Science Grant No. 1011479 (N.Y., Q.T.R. J.A.B,). We would like to acknowledge Dr. Haley Melikian for providing guidance on the dopamine transporter immunoblot. We would also like to acknowledge Drs. David Moody, Wenfang B Fang, Olga Averin, and Jonathan Crites for providing the analytical service for measuring methamphetamine concentration in our biological samples.

ABSTRACT

Individual variation in the addiction liability of amphetamines has a heritable genetic component. We previously identified *Hnrnph1* (heterogeneous nuclear ribonucleoprotein H1) as a quantitative trait gene underlying decreased methamphetamine-induced locomotor activity in mice. Here, mice (both male and female) with a heterozygous mutation in the first coding exon of *Hnrnph1* (H1+/-) showed reduced methamphetamine reinforcement and intake and dose-dependent changes in methamphetamine reward as measured via conditioned place preference. Furthermore, H1+/- mice showed a robust decrease in methamphetamine-induced dopamine release in the nucleus accumbens with no change in baseline extracellular dopamine, striatal whole tissue dopamine, dopamine transporter protein, or dopamine uptake. Immunohistochemical and immunoblot staining of midbrain dopaminergic neurons and their forebrain projections for tyrosine hydroxylase did not reveal any major changes in staining intensity, cell number, or in the number of forebrain puncta. Surprisingly, there was a two-fold increase in hnRNP H protein in the striatal synaptosome of H1+/- mice with no change in whole tissue levels. To gain insight into the molecular mechanisms linking increased synaptic hnRNP H with decreased methamphetamine-induced dopamine release and behaviors, synaptosomal proteomic analysis identified an increased baseline abundance of several mitochondrial complex I and V proteins that rapidly decreased at 30 min post-methamphetamine administration in H1+/- mice. In contrast, the much lower level of basal synaptosomal mitochondrial proteins in wild-type mice showed a rapid increase in response to methamphetamine. We conclude that H1+/- decreases methamphetamine–induced dopamine release, reward, and reinforcement and induces dynamic changes in basal and methamphetamine-induced synaptic mitochondrial function.

SIGNIFICANCE STATEMENT

Methamphetamine dependence is a significant public health concern with no FDA-approved

treatment. We discovered a role for the RNA binding protein hnRNP H in methamphetamine

reward and reinforcement. *Hnrnph1* mutation also blunted methamphetamine-induced

dopamine release in the nucleus accumbens – a key neurochemical event contributing to

methamphetamine addiction liability. Finally, *Hnrnph1* mutants showed a marked increase in

basal level of synaptosomal hnRNP H and mitochondrial proteins that decreased in response to

methamphetamine whereas wild-type mice showed a methamphetamine-induced increase in

synaptosomal mitochondrial proteins. Thus, we identified a potential role for hnRNP H in basal

and dynamic mitochondrial function that informs methamphetamine-induced cellular adaptations

associated with reduced addiction liability.

INTRODUCTION

Addiction to psychostimulants including methamphetamine (MA) is a major public health

concern in the United States, with an estimated 535,000 individuals currently meeting the

criteria for MA dependence (Lipari et al., 2016). Despite the prevalence of MA addiction, there is

currently no FDA-approved treatment, in part because the neurobiological mechanisms

underlying MA addiction are still not clear. Variation in sensitivity to the locomotor stimulant

response to psychostimulants is a heritable trait and can sometimes predict differences in drug

self-administration in rodents (Hooks et al., 1991; Yamamoto et al., 2013) because shared

neurocircuits and neurochemical mechanisms underlie these behaviors. We recently used

quantitative trait locus (QTL) mapping, positional cloning and gene editing via Transcription

Activator-like Effector Nucleases (TALENs) to identify *Hnrnph1* as a quantitative trait gene for

MA sensitivity in mice (Yazdani et al., 2015). *Hnrnph1* (heterogenous nuclear ribonucleoprotein

H1) encodes an RNA-binding protein (RBP) that is expressed throughout the brain, and is a part

of a subfamily of hnRNPs that includes hnRNP H1, hnRNP H2 and hnRNP F that possess

structurally unique quasi-RNA recognition motifs (Honoré et al., 1995). hnRNP H1 regulates all

aspects of RNA metabolism, including pre-mRNA splicing through binding at specific intronic

sites, mRNA stability and translational regulation via binding to the 5'UTR and 3'UTR, and poly-

adenylation control (Chou et al., 1999; George K. Arhin, Monika Boots, Paramjeet S. Bagga,

2002; Katz et al., 2010; Witten and Ule, 2011; Wang et al., 2012; Song et al., 2016).

We previously demonstrated that *Hnrnph1* polymorphisms and heterozygous deletion in

the first coding exon of *Hnrnph1* affect behavioral sensitivity to acute MA-induced locomotor

stimulation; however, the effects on MA reward and reinforcement are unknown. Additionally,

the neurobiological mechanism(s) underlying the mutational effects of *Hnrnph1* on MA-induced

behaviors remain to be established. *Hnrnph1* mRNA is ubiquitously expressed throughout the

adult mouse brain (Lein et al., 2007). While the protein expression of hnRNP H1 appears to be

nuclear-restricted, studies assessing hnRNP H1 protein in the brain are limited (Kamma et al.,

1995; Honoré et al., 1999; Van Dusen et al., 2010). With regard to CNS function, hnRNP H

family proteins are described as master regulators of neuron and oligodendrocyte differentiation

via alternative splicing control (Wang et al., 2007; Grammatikakis et al., 2016). Whole-exome

sequencing identified coding variants in human *HNRNPH1* and *HNRNPH2* (located on the X

chromosome) associated with severe neurodevelopmental disorders (Bain et al., 2016; Pilch et

al. 2018), implicating a crucial role of the hnRNP H protein family in neurodevelopment.

The purpose of the present study was three-fold. First, in order to expand beyond MA

locomotor stimulant sensitivity, we examined the effect of the *Hnrnph1* mutation on oral MA

reinforcement and intake via operant-conditioning and MA reward via conditioned place-

preference (CPP). This mutation comprises a small, frameshift deletion in the first coding exon

of *Hnrnph1* (H1+/-) that causes reduced MA-induced locomotor activity (Yazdani et al., 2015).

To gain insight into the neurobiological mechanisms underlying behavioral differences in MA

sensitivity, we examined drug-induced DA release via *in vivo* microdialysis, dopamine (DA)

content of striatal tissue, and DA clearance from striatal tissue. Second, because we previously

implicated *Hnrnph1* polymorphisms in dopaminergic neuron development, we assessed the

effect of H1+/- on tyrosine hydroxylase (TH) levels in cell bodies and processes of the

mesolimbic pathway via immunoblotting and immunohistochemistry. Finally, to gain insight into

neural dysfunction in H1+/- mice at the protein level that could underlie behavioral and

neurochemical deficits, we examined the synaptosomal proteome of the striatum between the

H1+/- and wildtype (WT) mice at baseline and in response to MA.

MATERIALS AND METHODS

Mice

Hnrnph1 mutant mice (H1+/-) were generated using TALENs targeting the first coding exon of

Hnrnph1 (exon 4, UCSC Genome Browser). Deletion of a small region in the first coding (exon

4) in *Hnrnph1* leads to a premature stop codon and transcription of a truncated mRNA

message. Gene expression via qPCR with primers specific for exons 6 – 7 (not targeting the

deleted exon 4) detected a 50% increase of *Hnrnph1* transcript in the H1+/- (heterozygous for

deletion) mouse embryos relative to WT (Yazdani et al., 2015) and a 1.5-fold increase in the H1-

/- (homozygous for deletion) (**Figure 7-1A to B**) with no change in gene expression of *Hnrnph2*

(Yazdani et al., 2015; **Figure 7-1C**). In addition, gene expression via qPCR with primers specific

for exon 4 detected a 50% reduction of *Hnrnph*1 transcript in H1+/- adult brain tissues relative to

WT (Yazdani et al., 2015). In addition, no change in hnRNP H protein expression was detected

in H1+/- and H1-/- embryonic stage brain tissue homogenate (**Figure 7-1D to G**). Because

hnRNP H1 and H2 have 96% amino acid sequence homology, there is no commercially

available antibody that can differentiate the two proteins. Quantitative analysis of protein peptide

from a previously unpublished hnRNP H immunoprecipitation mass spectrometry study

identified a decrease in the peptide encoded by the deleted region in exon 4 of *Hnrnph1* in

H1+/- compared to the WT mice (**Figure 7-2 A to B**). No genotypic difference was observed in

unique peptides associated with hnRNP H2 (**Figure 7-2 C to D**).

175   Experimental mice were generated by mating H1+/- males with C57BL/6J females

purchased from The Jackson Laboratory in Bar Harbor, ME USA (for studies in Boston

University) or in Sacramento, CA USA (for studies in UC Santa Barbara). Offspring were

genotyped as previously described (Yazdani et al., 2015). Unless otherwise indicated, both

female and male littermates (56-100 days old at the start of the experiment), were used in the

studies. Mice were housed in same-sex groups of 3-5 in standard mouse cages and housed

within ventilated racks under standard housing conditions. All procedures conducted in mice

were approved by the Boston University, UC Santa Barbara and Virginia Commonwealth

University Animal Care and Use Committees. All experiments were conducted in strict

accordance with the Guide for the Care and Use of Laboratory Animals. Colony rooms were

maintained on a 12:12 h light–dark cycle.

Genotyping of H1+/- and H1-/- mice

The genotyping protocol for the H1+/- mice is published in Yazdani et al. (2015). For

genotyping of H1-/- embryo tissue, an *Hnrnph1* forward primer

(GATGATGCTGGGAGCAGAAG) and reverse primer (GGTCCAGAATGCACAGATTG) were

designed to target upstream and downstream of the deleted region in exon 4 of *Hnrnph1*.

Genomic DNA was used to amplify a 204-bp PCR product using DreamTaq Green PCR

Mastermix (ThermoScientific) followed by overnight restriction enzyme digest with BstNI (New

England Biolads).

Hnrnph1 and *Hnrnph2* RT-qPCR for mouse embryo tissue

Oligo-dT primers were used to synthesize cDNA from total RNA to examine mRNA expression

and qPCR for evaluating gene expression were performed using Taqman SYBR Green

(ThermoFisher Scientific Cat# 4309155). Each sample was run in triplicate and averaged.

Differential gene expression was reported as the fold-change in H1+/- and H1-/- relative to WT

littermates using the $2^{-(\Delta\Delta C_T)}$ method. The primer sequences used for evaluating expression of

Hnrnph1 (targeting exons 6 and 7 were ACGGCTTAGAGGACTCCCTTT and

CGTACTCCTCCCCTGGAAGT. The primer sequences used for quantifying expression of

Hnrnph2 (targeting exons 1 and 2) were TAGCCGTTTGAGGGAAGAAG and

CCCTGTTAGAGTTTCTTCCAGGTA. The house keeping gene used was *Hprt* (targeting exons

7 and 9) with following primer sequence: GCTGGTGAAAAGGACCTCT and

CACAGGACTAGAACACCTGC.

Oral MA self-administration

The procedures for MA operant-conditioning were similar to those recently described (Lominac

et al., 2016). Testing was conducted in operant chambers equipped with 2 nose-poke holes, 2

cue lights, a tone generator, and a liquid receptacle for fluid reinforcement (Med Associates

Inc.). Mice were not water-restricted at any point during oral MA self-administration procedures.

The vehicle for MA was filtered tap water. Under a fixed-ratio 1 (FR1) schedule of

reinforcement, mice were trained daily to self-administer MA during 1 h sessions where a single

active nose poke response resulted in delivery of 20 µl of liquid MA into the receptacle, with a

20 s illumination of the cue light, and the sounding of the tone. During the 20 s period, further

responding resulted in no programmed consequences. Inactive hole responses were recorded

but had no consequences, serving to gauge the selectively of responding in the MA-reinforced

hole. Mice were initially trained to nose-poke 80 mg/l MA, with the concentration of MA

progressively increased over weeks (120, 160, 200, 300 and 400 mg/l MA; five days per dose).

Upon the completion of each daily session, the volume of MA remaining in the receptacle was

determined by pipetting and was subtracted from the volume delivered to calculate MA intake

(Lominac et al., 2016).

Conditioned place preference (CPP)

Mice were trained for 1 h each day in Plexiglas activity boxes within sound-attenuating

chambers (40 cm length x 20 cm width x 45 cm tall; divided into two sides with different plastic

floor textures for CPP). Mice were recorded from above using infrared cameras (Swan) and

tracked with ANY-maze (RRID:SCR_014289). The CPP paradigm was described in Kirkpatrick

and Bryant (2015).

On training Days 2-5, mice were injected with either saline (Days 2 & 4) or MA (Days 3 & 5;

saline, 0.5 or 2 mg/kg, i.p.) and confined to either the saline- or MA-paired side for 1 h.

Stereotaxic surgery

The procedures to implant indwelling microdialysis guide cannulae bilaterally over the NAc of

mice were described previously (Lominac et al., 2014, 2016). Mice were anesthetized under

1.5–2% isoflurane with 4% oxygen as a carrier gas, mounted in a Kopf stereotaxic device with

tooth and ear bars adapted for mice. The mouse skull was exposed, leveled, and holes were

drilled based on coordinates from Bregma for the NAC (AP: +1.3 mm, ML: ±1 mm, DV: −2.3

mm), according to the mouse brain atlas of Paxinos and Franklin (2001). The guide cannulae

were lowered bilaterally such that the tips of the cannulae were 2 mm above the NAc. The skull

was then prepared for polymer resin application and the guide cannulae were secured to the

skull with dental resin. Post-surgery, mice were injected subcutaneously (s.c.) with warm saline

and 250 µl of 2.5 mg/ml Banamine (Henry Schein Animal Health) and allowed to recover on a

heating pad. Post-operative care was provided for four days, during which mice were injected

with 250 µl of 2.5 mg/mL Banamine s.c. daily for the first two days. Mice were allowed a

minimum 1-week recovery prior to *in vivo* microdialysis assessments.

***In vivo* microdialysis & HPLC analysis**

Conventional microdialysis was conducted using a within-subjects design to examine saline and

acute MA-induced DA release (0.5 or 2 mg/kg, i.p.), using procedures similar to those described

previously (Lominac et al., 2014, 2016). Microdialysis probes were inserted unilaterally and

perfused with artificial cerebrospinal fluid for 3 h (2 µl/min), allowing for neurotransmitter

equilibration. For DA no net-flux analysis, DA was infused at 0, 2.5 nM, 5 nM, and 10 nM, and

dialysate was collected in 20-min intervals for 1h/concentration. On a subsequent day, mice

were probed on the contralateral side, and following the 3 h equilibration period and 1 h of

baseline dialysate collection, mice were injected i.p. with either 0.5 or 2.0 mg/kg MA and

dialysate was collected in 20-min intervals for 3 h post-injection. HPLC analysis of DA was

conducted as described previously (Lominac et al., 2014). Cannulae placement was determined

on Nissl-stained coronal sections and only mice exhibiting correct placement within the NAc

were included in analyses.

Behavioral test battery

Prepulse inhibition of acoustic startle*.* This test was employed to assess sensorimotor

gating. The apparatus and procedures were identical to those previously described in

Szumlinski et al. (2005). Six trial types were conducted: startle pulse (st110, 110 dB/40 ms), low

prepulse stimulus alone (st74, 74 dB/20 ms), high prepulse stimulus alone (st90, 90 dB/20 ms),

low or high prepulse stimulus given 100 ms. before the onset of the startle pulse (pp74 and

pp90, respectively) and no acoustic stimulus (st0; only background noise). All trials were

presented in a randomized order; st0, st110, pp74, and pp90 trials were given 10 times,

whereas st74 and st90 were presented five times. Background noise in each chamber was 70

dB and the average inter-trial interval lasted 15 s.

Novel object test*.* To assess anxiety-like behavior, mice were placed into a rectangular box

(9.25 x 17.75 x 8" high) containing one small, inedible object for 2 min. During that time, animals

were allowed to explore and interact with the object. The number of contacts and time in contact

(sec) with the novel object were video-recorded and tracked with ANY-maze tracking software

(RRID:SCR_014289). The apparatus and procedures used were identical to those previously

described in (Szumlinski et al., 2005).

Marble burying*.* The marble burying test was used to measure anxiety-like defensive burying

(Njung'e and Handley, 1991). In our paradigm, 12 square glass pieces (2.5 cm^2 × 1.25 cm high)

were placed in the animals' home cage, 6 at each end. The latency to start burying the marbles

was determined by a blind observer using a stopwatch and the total number of marbles buried

following a 20 min trial was recorded.

Light/dark shuttle box*.* The light/dark shuttle box test was employed to assess exploratory and

anxiety-like behaviors. Mice were placed into a polycarbonate box (46 cm long × 24 cm high ×

22 cm wide) wide containing distinct open (light) and closed (dark) environments for a 15 min

trial. These two environments were separated by a central divider with an opening. Mice were

first placed on the dark side and the latency to enter the light side, number of light-side entries,

and total time spent in the light-side of the shuttle box were recorded using ANY-maze tracking

software (RRID:SCR_014289). An increase in latency to enter the light, uncovered, side was

interpreted as an index of anxiety-like behavior.

Porsolt swim test. To assess depressive-like behavior (Porsolt et al., 1977), mice were placed

into a pool (30 cm in diameter; 45 cm high) filled with room-temperature water up to 35 cm and

allowed to swim for a total of 6 min. Time immobile (s), immobile episodes, and immobile

latency (sec) were video-recorded and tracked by ANY-maze tracking software

(RRID:SCR_014289).

Accelerating rotarod. To assess motor coordination, mice were trained on the rotarod (IITC life

science ROTO-ROD series) for a total of 10 trials over 3 days: 3 trials the first two days and 4

trials on the final day. The rotarod started at 4 RPM and accelerated to 40 RPM in 60 s. The

time (s) it took a mouse to fall (physically falling or hanging off rotarod) was manually scored.

Time on the rotarod was averaged across the total ten trials for each mouse.

Bitter/Quinine Taste Sensitivity

H1+/- and WT mice were allowed continuous-access in the home cage to 4 sipper tubes

containing 0 (filtered tap water), 0.1, 0.3 and 0.6 mg/ml quinine (Sigma-Aldrich). The quinine

concentrations selected for study were based on Eastwood and Phillips (2014). The mice and

bottles were weighed prior to initial presentation and the bottle was weighed every 24 h

thereafter. The difference in bottle weight was used to determine the volume consumed from

each solution over each 24 h period, the average intake from each solution, and the average

total volume consumed.

Quantification of baseline monoamine neurotransmitters from whole striatal tissue

Drug-naïve striatum were harvested from H1+/- and WT littermates and flash-frozen on dry ice.

The dissected tissue was sent to Vanderbilt University Neurotransmitter Core for the

quantification of monoamine neurotransmitters using HPLC wth electrochemical detection.

DAT-mediated DA uptake

Saline or MA (2.0 mg/kg) was administered interperitoneally in a volume of 10 ml/kg. After 2 h

post-administration (2 h was chosen based on the microdialysis results), mice were decapitated,

and DAT-specific [^{3}H]DA uptake from synaptosome preparations was conducted as described

previously (Kivell et al., 2014). Mice were rapidly decapitated, and striatal regions were

dissected from the brain and collected in 10 volumes (wt/vol) of prechilled 0.32 M sucrose buffer

(0.32 M sucrose in 5 mM HEPES, pH 7.5). The striatal tissue was homogenized and centrifuged

at 1000 x g for 15 min at 4°C. The supernatant was centrifuged at 12,000 x g for 20 min, and the

pellet was suspended in 0.32 M sucrose buffer. Striatal synaptosomes (30 µg) were incubated

in a total volume of 0.3 ml of Krebs-Ringer-HEPES (KRH) buffer consisting of 120 mM NaCl, 4.7

mM KCl, 2.2 mM CaCl$_2$, 10 mM HEPES, 1.2 mM MgSO$_4$, 1.2 mM KH$_2$PO$_4$, 5 mM Tris, 10 mM

D- glucose, pH 7.4 containing 0.1 mM ascorbic acid, and 0.1 mM pargyline at 37°C for 10 min

with or without DAT- specific blocker GBR12909 (50 nM). Following incubation, 5 nM [^{3}H]DA

(63.2 Ci/mmol-dihydroxyphenylethylamine [2,5,6,7,8-3H]; PerkinElmer) and further incubated for

additional 5 min. Uptake of DA was terminated with the addition of 500 nM DAT blocker

GBR12909. The samples were filtrated over 0.3% polyethylenimine coated GF-B filters on a

Brandel Cell Harvester (Brandel Inc.), and washed rapidly with 5 ml cold PBS. Radioactivity

bound to the filter was counted using a liquid scintillation counter. DAT mediated [^{3}H]DA uptake

was determined by subtracting total accumulation of [^{3}H]DA (absence of GBR12909) and in the

presence of GBR12909. Uptake assays were performed in triplicates.

Immunohistochemistry

For immunohistochemistry (IHC), drug-naïve H1+/- and WT mice were anesthetized with

pentobarbital, and transcardially perfused with phosphate buffered saline (PBS) followed by 4%

paraformaldehyde in PBS at room temperature. Next, brains were dissected and processed for

tyrosine hydroxylase (TH) 3-3'-diaminobenzidine (DAB) IHC and analysis as previously

described (Burke et al., 1990; Hutson et al., 2011), or double immunofluorescent IHC for hnRNP

H and TH colocalization. For DAB IHC, coronal slices were blocked with 4% normal goat serum

and then incubated for 48 h at 4°C with anti-hnRNP H (1:50,000, Bethyl Cat# A300-511A, RRID:AB_203269) or tyrosine hydroxylase (TH) (1:500, Santa Cruz Cat# sc-14007, RRID:AB_671397), and processed for DAB staining and analyzed as previously described (Hutson et al., 2011). For co-staining studies with hnRNP H and TH, tissues were blocked with superblock (ThermoFisher Scientific Cat# 37515), and incubated with anti-hnRNP H (Santa Cruz Cat# sc-10042, RRID:AB_2295514) and TH (1:500, Santa Cruz Cat# sc-14007, RRID:AB_671397) for 48 hours at 4°C. Next, tissues were incubated with donkey anti-rabbit Alexa Fluor 488 (1:500, Molecular Probes Cat# A-21206, RRID:AB_141708) and donkey anti-goat Alexa Fluor 633 (1:500, Molecular Probes Cat# A-21082, RRID:AB_141493), washed, and then coated with ProLong Diamond Antifade Mountant (ThermoFisher Scientific Cat#P36961), mounted onto slides, and imaged on the Leica SPE Confocal microscope.

346 TH puncta quantification in the striatum

347   Entire coronal slices of rostral, medial, and caudal striatum were imaged at 40x magnification using a Nikon Eclipse 600 microscope. A 225,000 μm^2 grid was overlaid onto these images using Image J and every 3rd field of view within the striatum was graded. Number of puncta within a field of view was graded in ImageJ by subtracting out background signal, creating binary images from these files, and then counting puncta meeting roundness and diameter criteria (roundness <0.6, diameter 1-45 µm). Averages puncta densities for ventral, dorsal, and total striatum were calculated. Grading of the puncta was performed in Image J. The image was duplicated into a 1,000,000-pixel area followed by brightness/contrast adjustment and background subtraction. The threshold was set to 106. A binary image was then generated for puncta measurement. Roundness was set to 0.6-1 and size was set to 5 – 200 pixels.

357 Dissection of mouse brain regions: striatum and midbrain

Live, rapid decapitation was used to avoid the effects of anesthesia or CO_2 asphyxiation on gene expression. Immediately followed live-decapitation using large, sharpened shears with an incision just posterior from the ears, the mouse brain was removed quickly with forceps and

transfer to a cold surface. The striatum has a somewhat darker appearance than the

surrounding cortex. To dissect the dorsal and ventral striatum, fine-tip forceps were used to

separate the midline of the brain and then the cortex and hippocampus were removed to reveal

the striatum. To dissect the midbrain, a razor blade was used to make a rostral cut where the

cerebral aqueduct begins and another caudal cut just before the start of the cerebellum.

Methamphetamine-induced locomotor activity followed by tissue harvesting

On Days 1 and 2, all mice received a saline injection (10 ml/kg, i.p.) and were recorded for

locomotor activity in Plexiglas chambers (40cm length x 20 cm width x 45 cm height) for 1 h. On

Day 3, mice receive either saline or MA (2 mg/kg, i.p.; Sigma Aldrich) and were recorded for

locomotor activity for 30 min and the whole striatum was harvested as described above at 30

min post-injection. Whole striata (left and right sides) were flash frozen in ethanol/dry ice bath

and stored at -80°C for long-term storage. Four cohort of animals were run in this behavioral

paradigm for tissue collection: 1) hnRNP H immunoprecipitation followed by mass spec; 2)

synaptosome mass spec and hnRNP H immunoblot; 3) validation studies for mitochondrial

protein immunoblots; 4) measurement of MA concentration in MA-treated striatal tissues. The

mice that were used for hnRNP H immunoprecipitation were all MA-treated on Day 3.

Preparation of synaptosomes

Striatal tissue collection from saline- or MA-treated mice was performed as described above.

The tissues were subsequently processed to obtain synaptosomes using a Percoll (Sigma

Aldrich Cat# 1644) gradient fractionation method, which was adapted from Dunkley et al.

(Dunkley et al., 2008). Whole striata (left and right hemisphere) were placed in 1 ml sucrose

homogenization buffer (2 mM HEPES, pH7.4, 320 mM Sucrose, 50 mM EDTA, 20 mM DTT)

supplemented with protease and phosphatase inhibitor (ThermoFisher Scientific Cat# 78440).

The brain tissues were lightly homogenized using a handheld motorized pestle. The

homogenate for each sample was then centrifuged for 2 min at 3000 rcf and the supernatant

(S1) was collected. The pellet (P1) was then resuspended in 500 µl of sucrose homogenization

buffer and re-spun for 3 min at 3000 rcf. The supernatant (S1') was collected and combined with

S1 and then centrifuged for 15 min at 9200 rcf. The supernatant S2 was then removed and the

pellet (P2) was re-suspended in 500 µl of sucrose homogenization buffer and loaded onto a

Percoll density gradient consisting of 23%, 10%, and 3% Percoll (1 ml each) in Polycarbonate

centrifuge tubes (13 x 51 mm; Beckman Coulter Cat# 349622). The gradients were then

centrifuged for 15 min at 18,700 rcf. The distinct band between the 10% and 23% Percoll was

collected as the synaptosome. The synaptosome fraction was then washed to 5 mL with 1X

HBM buffer, pH 7.4 (140 mM NaCl, 5 mM KCl, 5 mM $NaHCO_3$, 1.2 mM NaH_2PO_4, 1 mM $MgCl_2$-

$6H_2O$, 10 mM glucose, 10 mM HEPES) to dilute out the Percoll by centrifuging for 12 min at

18700 rcf. The pellet was then re-suspended in 100 µl of HBM buffer to yield the final

synaptosme fraction and BCA assay was used to determine protein concentration. A total of 30

µg of synaptosome was loaded per sample for SDS-PAGE and Western blotting as described

below.

SDS-PAGE and Western Blot

Brain tissues were homogenized using hand-held homogenizer in RIPA buffer with

Halt^{TM} Proteatase & Phosphotase inhibitor cocktail (ThermoFisher Scientific Cat# 78840). For

each sample, 30 µg of protein was heated in a 70^{o}C water bath for 10 min prior to loading into

into a 4-15% Criterion TGX precast Midi protein gel (Bio-Rad) for SDS-PAGE followed by wet

transfer to nitrocellulose membrane. The membrane was then blocked with 5% milk for 1 h and

probed with primary antibodies. For evaluating TH expression in brain tissues, overnight

incubuation of the membrane at 4°C with anti-TH (1:50,000, Santa Cruz Cat# sc-14007,

RRID:AB_671397) was performed followed by 1 h incubation with donkey anti-rabbit HRP

(1:10,00, Jackson ImmunoResearch Labs Cat# 711-035-152, RRID:AB_10015282). For

evaluating hnRNP H protein expression in mouse embryo tissues and in striatal synaptosome,

the following antibodies were used: hnRNP H (C-term: 1:50,000, Bethyl Cat# A300-511A,

RRID:AB_203269; N-term: 1:50,000, Santa Cruz Cat# sc-10042, RRID:AB_2295514) followed

by 1 h incubation with the appropriate secondary antibodies. For vadilation of mitochondrial

protein expression following MA treatment, the following priamry antibodies were used: ATP5A1

(1:2000, Abcam Cat# ab14748, RRID:AB_301447); ATP5F1 (1:5000, Proteintech Cat# 15999-

1-AP, RRID:AB_2258817); and NDUFS2 (1:5000; abcam ab192022). We used the following

loading controls: anti-β-actin (1:20,000, Sigma-Aldrich Cat# A2228, RRID:AB_476697); anti-

GAPDH (1:20,000; Millipore Cat# MAB374, RRID:AB_2107445), and PSD95 (1:10,000, Cell

Signaling Technology Cat# 3450, RRID:AB_2292883).

Mouse brain tissue processing for SDS-PAGE and DAT immunblotting was modified

from Staal et al., 2007. Briefly, tissue was triturated using a 20-22 gauge needle in RIPA buffer

(10 mM Tris, pH 7.4, 150 mM, NaCl, 1 mM EDTA, 0.1% SDS, 1% Triton X-100) supplemented

with Halt^{TM} Proteatase & Phosphotase inhibitor cocktail (ThermoFisher Scientific Cat# 78840).

30 µg of protein of each sample was allowed to rotate at room temperature prior to SDS-PAGE

instead of heating the sample at high temperature. For evaluating DAT expression in whole

striatal tissue and striatal synaptosome, we conducted overnight incubation anti-DAT (1:2000;

Millipore Cat# MAB369, RRID:AB_2190413) followed by 1 h incubation with goat anti-rat (1:500,

Jackson ImmunoResearch Labs Cat# 112-035-003, RRID:AB_2338128).

All processed membranes were imaged via enhanced chemiluminescence photo-

detection. Densitometry analysis in Image J was used for quantification.

hnRNP H immunoprecipitation

Following the three-day locomotor paradigm assessing acute locomotor sensitivity in H1+/-

versus WT mice as described above, striata were dissected from the mice at 30 min post-

injection of MA or saline. Striatum was dissected from WT or H1+/- mice and frozen on dry ice

and stored at -80^{o}C. Striatal tissues were then homogenized using a microcentrifuge pestle in

ice-cold RIPA buffer (50 mM Tris-HCl, pH 6.8, 150 mM NaCl, 5 mM EDTA, 1% Triton X-100,

0.1% SDS, 0.5% sodium deoxycholate) supplemented with protease and phosphatase inhibitor

cocktails (ThermoFisher Scientific Cat# 78840) and incubated overnight at 4^{o}C with gentle

agitation. Lysates were centrifuged at 10,000 rpm for 15 min at 4^{o}C, and the supernatant

fraction was saved for protein quantification via BCA assay. 1 mg of striatal lysates was pre-

cleared for 1 h with 80 ul Protein G Sepharose coated beads (ThermoFisher Scientific Cat#

101243) and then centrifuged at 4^{o}C for 5 min at 1000 rpm. The pre-cleared lysates

(supernatant) were then incubated overnight with 10 µg of rabbit anti-hnRNP H (Bethyl Cat#

A300-511A, RRID:AB_203269) or control normal rabbit IgG antibody (Millipore Cat# NI01-

100UG, RRID:AB_490574). The next day, 80 µl of Protein G Sepharose coated beads were

added to the antibody-lysate mixture, and incubated for an additional 2 h. The beads were then

washed 4 times in 1 ml lysis buffer, resuspended and centrifuged for 1 min at 1000 rpm each

time. The beads were eluted by adding 60 µl of non-reducing SDS buffer and heating at 95^{o}C

for 10 min. hnRNP H immunoprecipitates were eluted in non-reducing SDS buffer as described

above. 50 µl of each sample was separated by SDS-PAGE at 100 V for 30 min (~2 cm) on a

Novex Bolt 4-12% Bis-Tris gel. The gel was then washed 3 times in deionized H$_2$O and stained

with Simply Blue Coomassie SafeStain (ThermoFisher Scientific Cat# LC6060). The gel was

then cut at ~160 kDa to exclude the prominent non-reduced IgG band. Individual gel lanes

were then separately excised and stored in pre-washed microcentrifuge tubes at 4^{o}C prior to

shipping to the UMass Worchester Proteomics and Mass Spectrometry facility for analysis by

OrbiTrap liquid chromatography tandem mass spectrometry.

TMT Labeling, High pH reverse phase HPLC fraction, followed by LC-MS/MS

Following the three-day locomotor paradigm assessing acute locomotor sensitivity in H1+/-

versus WT mice as described above, striata were dissected from the mice at 30 min post-

injection. Synaptosomes were isolated following the procedure outlined above for proteomic

characterization. Samples were resuspended in 8 M urea for 30 min, followed by the addition of

5 mM DTT for 1 h. Lodoacetamide was then add to the samples that were incubated in the dark

for 30 min. The urea concentration was diluted below 1 M with the addition of 50 mM of

ammonium bicarbonate. The samples were then digested with trypsin (50:1, protein to enzyme

ratio) overnight at 37°C and terminated with the addition of formic acid to 1%. The samples were

desalted with a C18 tip. Peptide was determined by Pierce Quantitative Colorimetric Assay

(ThermoFisher Scientific Cat# 23275), then 100 µg of peptide was resuspended in 0.1 M

triethylammonium bicarbonate (TEAB) and incubated with the TMT plex isobaric labeling for 1 h

at room temperature. To quench the reaction, 5% hydroxylamine was added to each sample

and incubated for 15 min. Each sample was combined at equal amount and cleaned with C18

tips. One mg of labeled peptides was fractioned using a Waters XBridge BEH C18 (3.5µm, 4.6

×250mm) column on an Agilent 1100 HPLC system. 48 fractions were collected and combined

to 12 fractions and then dried. A C18 Acclaim PepMap 100 pre-column (3µm, 100 Å, 75µm ×

2cm) hyphenated to a PepMap RSLC C18 analytical column (2µm, 100 Å, 75µm × 50cm) was

used to separate peptide mixture. LC-MS/MS analyses were completed using an EASY nLC

1200 system coupled to a Q Exactive HF-X mass spectrometer. Full MS spectra were collected

at a resolution of 120,000 with an AGC of 3×10^6 or maximum injection time of 50 ms and a

scan range of 350 to 1650 m/z. MS2 scans were performed at 45,000 resolution and using 32%

total normalized collision energy. Source ionization parameters were optimized with the spray

voltage at 2.1 kV, dynamic seclusion was set to 45 s.

Proteomics Data Analysis and Pathway Enrichment Results

The acquired data was searched by MaxQuant against the UniProt mouse proteome data base

with standard settings. (fragment ion mass tolerance of 20ppm, maximum missed cleavage of 2,

oxidation as variable modification, false discovery was 1%, only protein groups identified with at

least 2 or more peptides). The intensity data were filtered and normalized using R

(RRID:SCR_001905) and the LIMMA package (RRID:SCR_010943) was used for differential

analysis with Genotype and Treatment as factors. A ranked list was generated from the analysis

and the fgsea R package was used to perform pre-ranked analysis, with proteins filtered for

absolute $\log_2FC > 0.2$ and $p < 0.05$. Enrichment results for the comparisons were visualized

using Cytoscape (RRID:SCR_003032) and EnrichmentMap (RRID:SCR_016052), with nodes

representing pathways and edges representing overlap genes. Pathways were clustered and

annotated with themes using AutoAnnotate (Reimand et al., 2019).

Experimental design and statistical analyses

We used an effect size (Cohen's d = 0.9) based on genotypic differences in MA-induced

locomotor activity at 30 min post-MA in our previously published data from H1+/- mice (Yazdani

et al., 2015). We used G*Power 3 (Faul et al., 2007) to calculate a sample size of n = 16 per

genotype required for 80% power (p < 0.05). Statistical details of the experiments can be found

in the figure legends. Data are presented as means of replicates from each experiment ± SEM.

For experiments in which two conditions were compared, an unpaired, two-tailed Student's t-test

was used to analyze the data. For experiment in which multiple factors were evaluated, ANOVA

was used to calculate statistical significance as indicated in the figure legends. Statistical p

value threshold for Student's t test and ANOVA was set to 0.05. All statistical analyses were

performed in R (RRID:SCR_001905).

Data and Code Availability

All related data and materials are available upon request. The mass spectrometry proteomics

data are available in MassIVE under proteome exchange accession number PXD014813.

RESULTS

Oral MA self-administration

We previously demonstrated a robust reduction in sensitivity to the locomotor stimulant

response to MA in H1+/- mice (Yazdani et al., 2015). To more directly test MA-induced

motivated behaviors, we investigated MA reinforcement and intake in H1+/- mice using an oral

MA self-administration paradigm in C57BL/6J mice (Szumlinski et al., 2017). The study showed

that mice allocated the majority of their nose-poking behavior toward the MA-reinforced hole,

indicating drug reinforcement. H1+/- mice presented fewer active nose-pokes at the 160, 200,

300 and 400 mg/L MA doses compared to WT mice (**Figure 1A**), with no difference in non-

specific, inactive nose-pokes across a range of MA concentrations (80-400 mg/L) (**Figure 1B**).

Consistent with their lower level of MA reinforcement, H1+/- mice also consumed less MA

(intake; mg/kg/day) at the 200, 300 and 400 mg/L concentrations (**Figure 1C**). These results

indicate that the H1+/- mice are less motivated to work for the MA reinforcer. No sex differences

or interactions were observed for any measure during self-administration testing. The genotypic

differences in high-dose MA intake did not relate to bitter taste sensitivity as quinine intake in

the home cage was equivalent between H1+/- and WT mice (**Figure 1-1**).

The blunted escalation of oral MA intake observed at high MA concentrations (≥ 200

mg/L) in H1+/- mice initially trained to respond for 80 mg/L MA prompted us to determine if the

Hnrnph1 mutation would blunt the acquisition of oral MA self-administration when 200 mg/L MA

served as the reinforcer. However, we were unable to detect any genotypic difference in MA

intake and active nose-pokes when mice were trained at this higher MA concentration (**Figure**

1-2)**.** These results indicate that the *Hnrnph1* mutation interferes with the transition from low to

high-dose MA self-administration, without altering the reinforcing effects of MA during

acquisition.

MA-conditioned reward

To further investigate why H1+/- mice showed less oral self-administration of MA, we assessed

MA reward via conditioned place preference (CPP). When mice were tested in a MA-free state

(Day 8), H1+/- mice showed lower CPP to 0.5 mg/kg MA but higher CPP to 2 mg/kg MA

compared to the WT mice (**Figure 2A**). Similarly, during drug state-dependent CPP, H1+/- mice

also exhibited lower and higher CPP at 0.5 mg/kg and 2 mg/kg MA doses, respectively (**Figure**

2B). No sex differences or interactions were observed for any measure during MA-CPP testing.

The dose-dependent difference in CPP indicates that H1+/- showed a reduced sensitivity to the

rewarding effect of MA where a higher dose of CPP was required to elicit CPP in H1+/- mice. In

addition, although we did not detect significant genotypic differences in MA-induced locomotor

activity during the confined one-half of the CPP box during training, we observed a trend toward

a decrease in MA-induced locomotor activity in H1+/- mice on Day 9 during state-dependent

CPP (following the two previous MA exposures) when they had twice as much open access

space (**Figure** 2-1). This result was very similar to the previously published results in the same-

sized open arena minus the CPP divider and a single MA exposure of 2 mg/kg (Yazdani et al.,

2015). Furthermore, note that we replicated the reduced MA-induced locomotor activity

phenotype in H1+/- mice with the same previously published protocol (Yazdani et al., 2015)

prior to tissue harvesting for mass spectrometry analysis (**Figure 8-1**). There was no genotypic

difference in the striatal level of either MA or its metabolite amphetamine at 30 min i.p. post

injection of MA (**Figure 8-2**). Thus, differences in transport or metabolism of MA is an unlikely

explanation for the dramatic genotypic differences in MA-induced behavior at 30 min post-MA.

MA-induced DA release in H1+/- mice

Because H1+/- mice showed reduced MA self-administration and reward, we next

examined basal extracellular DA and MA-elicited DA release using *in vivo* microdialysis in the

NAc, a brain region and neurochemical event that are necessary for MA reinforcement and

reward (Prus et al., 2009; Keleta and Martinez, 2012; Bernheim et al., 2016). Linear regression

analysis of no net-flux *in vivo* microdialysis in the NAc indicated no significant change in either

DA release/reuptake (extraction fraction or slope, **Figure 3-1A**) or basal extracellular DA

content or (**Figure 3-1B**). Following administration of 0.5 mg/kg MA (i.p.), WT mice exhibited an

increase in extracellular DA whereas H1+/- mice showed a decrease in extracellular DA below

baseline (**Figure 3A**). Administration of 2 mg/kg MA (i.p.) induced an increase in DA within the

NAc of both genotypes, but H1+/- mice again showed markedly lower MA-induced extracellular

DA levels, in particular from 100 to 180 min post-MA (**Figure 3B**). Consistent with the genotypic

differences in extracellular DA levels, for both the 0.5 and 2 mg/kg MA doses, H1+/- mice also

showed a lower level of 3,4-Dihydroxyphenylacetic acid (DOPAC, metabolite of DA) compared

to WT mice (**Figure 3C-D**).

In examining glutamate levels, no net flux microdialysis study revealed no difference in

basal extracellular level of glutamate (**Figure 3-2A**). Acute administration of 0.5 mg/kg or 2

mg/kg MA did not induce any effect on extracellular glutamate levels, nor was there any

genotypic difference in glutamate levels (**Figures 3-2B and C**). To summarize, H1+/- mice

showed a significantly blunted MA-induced DA release in the NAc in the absence of any

significant difference in baseline DA levels and in the absence of any significant differences in

glutamate neurotransmission. No sex differences or interactions were observed. These results

point toward a MA-induced deficit in DA release as a plausible functional mechanism underlying

the deficit in the MA-induced behavioral responses in H1+/- mice.

To further investigate the possible causes of the decreased capacity of MA to induce DA

release in H1+/- mice, we quantified basal striatal tissue levels of biogenic amines in drug-naïve

H1+/- and WT mice. No genotypic difference was detected in the amount of DA and DOPAC, or

in the levels of other biogenic amines including HVA, 3-MT, 5-HT, etc. (**Figure 4A**). We also

tested whether H1+/- impacts the function of DAT by examining DAT-mediated DA uptake in

synaptosomes prepared from whole striatal tissue. However, no difference in basal DAT

function was detected (**Figure 4B**). In addition, there was no change in the protein level of DAT

in the total striatal brain lysate in H1+/- compared to WT mice (**Figure 4C-D**). There was also no

effect of MA on DAT protein level in the synaptosome of either H1+/- or WT mice (**Figure 4-1).**

Tyrosine hydroxylase expression in the mesolimbic and nigrostriatal brain regions of

H1+/- mice

In our previous transcriptome analysis of congenic mice that captured *Hnrnph1*

polymorphisms and decreased MA behavioral sensitivity, we identified a decrease in expression

of *Nurr1* (*Nr4a2*), a transcription factor critical for dopaminergic neuron development (Yazdani et

al., 2015). Thus, we hypothesized that dysfunctional mesolimbic and nigrostriatal DA pathways

at the neuroanatomical level could underlie decreased MA-induced behaviors and DA release in

H1+/- mice. We tested this hypothesis by examining changes in the number of neurons or the

number of projections of neurons in the mesolimbic and nigrostriatal pathway, which mediate

reward and motor activity, respectively (Adinoff 2004). Diagrams outlining the different brain

regions assessed are presented in **Figure 5-1**. We first examined co-expression of hnRNP H

(there are currently no commercially available antibodies for differentiating between hnRNP H1

and hnRNP H2) and tyrosine hydroxylase (TH; the rate-limiting enzyme for the synthesis of DA;

Daubner et al., 2011). Results showed that hnRNP H and TH were co-expressed in the same

midbrain TH neurons of the ventral tegmental area (VTA) and substantia nigra pars compacta

(SNc) at a similar level in both genotypes (**Figure 5A**). hnRNP H immunostaining was primarily

nuclear whereas TH immunostaining was cytoplasmic. We next examined expression of TH in

the dopaminergic neurons of the mesolimbic and nigrostriatal circuit (**Figures 5 and 6)**. TH DAB

immunohistostaining in the VTA and SNc of the midbrain (**Figure 5B**) indicated no significant

genotypic differences in TH optical density (OD) (**Figure 5C)**. There was also no genotypic

difference in the number (**Figure 5D**) or diameter (**Figure 5E**) of TH-positive neuron of the VTA

and SNc. TH DAB immunostaining in the striatum indicated no change in TH OD in the dorsal

striatum (**Figures 6A-C**) between H1+/- and WT mice. However, a small, but statistically

significant increase in TH OD was observed in the ventral striatum (which includes the NAc) of

the H1+/- mice (**Figures 6D-F**). To measure TH-positive puncta as an indirect estimate of the

number of DA terminals in the striatum, we performed stereology under higher magnification

(**Figure 6-1A**) and detected no difference in the number of TH-positive puncta between the

H1+/- and WT mice in the dorsal striatum or in the ventral striatum (**Figure 6-1B**). Immunoblot

analysis indicated no significant difference in TH protein level in either the midbrain (**Figures**

5F-G) or the whole striatum (**Figures 6G-H**). Taken together, analysis of the dopaminergic

mesolimbic and nigrostriatal DA circuit did not provide strong evidence for changes in the

expression of TH within cell bodies or terminals or changes in the number of TH-positive

terminals that could explain behavioral differences in H1+/- mice.

An important message gleaned from the above results is that the behavioral and

neurochemical deficits exhibited by H1+/- mice appear to manifest only under the influence of

MA. In further support of this notion, a screen of WT and H1+/- mice in a behavioral test battery

did not detect any genotypic differences in sensorimotor-gating, anxiety-like behaviors,

depressive-like behaviors, or in sensorimotor coordination (**Figure 3-3**). The null results from

this behavioral battery, combined with the lack of genotypic differences in saline-induced

locomotion/response to a novel environment, and in various indices of basal DA transmission in

the striatum argue for an active, MA-induced cell biological mechanism linking *Hnrnph1* function

to MA behavior.

Methamphetamine-induced changes in total and synaptic level of hnRNP H

The above neuroanatomical studies failed to support a neuroanatomical hypothesis of

reduced neurodevelopment of DA projection pathways that could underlie reduced MA-induced

DA release and behavior in H1+/- mice. Thus, to further explore the possibility of a synaptic,

MA-induced cell biological mechanism that could underlie H1+/- behavior, we next examined

changes in the synaptic localization of hnRNP H which is potentially relevant for understanding

how H1+/- could alter MA-induced DA release. Surprisingly, we identified a two-fold *increase* in

hnRNP H protein level in the striatal synaptosome of H1+/- mice, regardless of treatment

(**Figure 7 A-B**). In contrast, there was no significant genotypic difference in hnRNP H protein

level in bulk striatal tissue in response to saline or MA (**Figure 7 C-D**), indicating a change in

localization rather than total hnRNP H protein in H1+/- mice.

To identify changes in the levels of other synaptic proteins that could mechanistically link

the robust increase in synaptic hnRNP H with decreased MA-induced DA release and behavior,

we examined genotypic differences in the synaptosomal proteome in the striatum of H1+/- and

WT mice treated with MA versus saline using LC-MS/MS. At the behavioral level, H1+/- showed

reduced MA-induced locomotor activity only in response to MA but not to saline (**Figure 8-1**).

Overall, proteomic analysis of the main effect of Genotype identified a highly enriched

upregulation of mitochondrial proteins in the H1+/- striatal synaptosome regardless of treatment

(**Figure 8A**). Enrichment analysis for the set of top differentially expressed proteins (absolute

$log_2FC > 0.2$; $p < 0.05$) between H1+/- and WT mice revealed a highly significant enrichment for

mitochondrial respiratory chain complex I assembly (**Table 1**). In examining MA-induced

changes in synaptic proteins in H1+/- versus WT mice, we again identified an enrichment for

alterations in metabolic processes involving components of mitochondrial complex I and V

(**Figures 8B, 8-3, and Table 2**). Interestingly, in response to MA, there was a decrease in the

level of mitochondrial proteins in the H1+/- mice, but an increase in the WT mice (**Figure 9**). The

findings were independently validated with a separate cohort of mice, which also pointed to a

trending, though non-significant pattern of an increase in expression of all three mitochondrial

proteins that we tested in saline-treated H1+/- mice and a trending decrease in all three

mitochondrial proteins in response to MA (**Figure 9-1**).

DISCUSSION

Here, we extend a role for *Hnrnph1* in MA reinforcement and reward (**Figure 10**). A

heterozygous frameshift 16 bp deletion of the first coding exon of *Hnrnph1* reduced MA oral

operant self-administration and MA-induced CPP (**Figures 1 and 2**) and induced a robust

reduction in MA-induced DA release in the NAc (**Figure 3**). This DA anomaly occurred without

any differences in 1) basal extracellular DA (**Figure 3A-B**); 2) basal DA content within striatal

tissue (**Figure 4A**); 3) DA uptake (**Figure 4B**); 4) the number or staining of midbrain DA

neurons within the mesolimbic and nigrostriatal dopaminergic circuits (**Figure 5**) or 6) any

robust changes in the number of forebrain striatal puncta originating from these neurons (**Figure**

6). The combined results suggested an alternate mechanism underlying the decreases in MA-

induced DA release and behavior in H1+/- mice**.** In further support of a MA-induced cell

biological mechanism, there was no effect of H1+/- on spontaneous locomotion, anxiety- and

depressive-like behaviors, or sensorimotor function (**Figure 3-3**).

H1+/- mice showed less MA-CPP at 0.5 mg/kg but greater CPP at 2 mg/kg than WT

mice (**Figure 2**). One interpretation is that MA-CPP exhibits an inverted U-shaped dose-

response curve (Uhl et al., 2014) in WT and that H1+/- shifts this curve to the right, yielding

reduced sensitivity to positive and negative motivational valence of MA at lower versus higher

doses. Consistent with this interpretation, H1+/- mice exhibited blunted operant oral MA

reinforcement. The combined data are consistent with H1+/- mice self-administering less MA

because they are less sensitive to the physiological and interoceptive effects of MA due to a

reduced MA-induced DA release.

H1+/- mice exhibited a blunted DA response to 0.5 and 2 mg/kg MA (**Figure 3A-B**). This

effect on MA-induced DA release could not be explained by alterations in total DA levels

(**Figure 4A**), or in DAT levels or function at the presynaptic membrane (**Figures 4B-D and 4-1**).

An alternative explanation to DAT dysfunction is that the *Hnrnph1* mutation somehow decreases

MA binding to DAT, limiting its entry into presynaptic dopaminergic neuronal terminals and

decreasing DA release. Future studies are necessary to examine MA binding to DAT in H1+/-

mice.

Our findings support a dopaminergic mechanism underlying reduced MA reward and

reinforcement in H1+/- mice. Nevertheless, we acknowledge the potential involvement of

additional neurotransmitter systems and brain regions. We did not identify any difference in

basal or MA-induced changes in extracellular glutamate levels in the NAc of H1+/- mice. While

MA reward is generally attributed to an increase in DA release in the NAc (Segal and

Kuczenski, 1997; Adinoff, 2004), MA also increases release of norepinephrine and serotonin by

targeting their respective transporters that could modulate the locomotor stimulant and/or

rewarding response to psychostimulants such as MA (Haughey et al., 2002; Rothman et al.,

2001; Zaniewska et al., 2015). Future studies are warranted to address these other

neurotransmitters in MA-induced behavioral dysfunction in H1+/- mice as well as the possibility

that H1+/- has a pleiotropic influence on behavioral (e.g., cognitive, antidepressant) and

neurochemical responses to drugs targeting other membrane transporters such as NET and

SERT. For example, phosphorylation of the RBP hnRNP K increases expression of SERT

protein via changes in binding to the distal polyadenylation element of the transporter (Yoon et

al., 2013).

The two-fold increase in hnRNP H protein in the striatal synaptosome of H1+/- mice with

no change in total hnRNP H protein was surprising. This finding was observed in multiple

replication studies and suggests a redistribution of hnRNP H protein to the synapse in H1+/-

mice. We performed LC-MS/MS analyses on striatal synaptosomes following 2 mg/kg MA (i.p.)

to further understand the effect of increased synaptosomal hnRNP H on global changes in

protein expression and the underlying cell biological mechanisms. We identified a higher

abundance of several mitochondrial proteins, in particular, complex I of the mitochondrial

respiratory chain in H1+/- mice. The mammalian complex I consists of 38 nuclear DNA-encoded

subunits (Sharma et al., 2009) and our proteomic analysis identified 8 out of the 38 subunits that

showed higher expression in H1+/- mice (**Figures 8-9**). Proteomics has been widely used to

study the effects of MA on protein expression in the brain tissues of animals (Liao et al., 2005;

Faure et al., 2009; Bosch et al., 2015). Iwazaki et al. (2006) used two-dimensional gel

electrophoresis proteomics and found that a single low dose of MA (1 mg/kg) administration in

rats induced differential expression of proteins involved in mitochondria/oxidative metabolism.

Furthermore, chronic exposure of 1 mg/kg MA induced locomotor sensitization and neurotoxicity

along with a downregulation of numerous striatal proteins indicating mitochondrial dysfunction

and an oxidative response (Iwazaki et al., 2007; Chin et al., 2008).

Recent studies showed postsynaptic dendritic mitochondrial fission and fusion

processes mediate cellular and behavioral plasticity, spine and synapse formation, and synaptic

function (Li et al., 2004; Oettinghaus et al., 2016; Chandra et al., 2017; Divakaruni et al., 2018).

Dynamin-related protein (Drp1), a GTPase involved in mitochondria fission, has been shown to

regulate addiction-relevant behavior during early cocaine abstinence (Chandra et al., 2017),

with inhibition of mitochondrial fission blunting cocaine-seeking and locomotor sensitization.

While the results of our synaptosomal proteome dataset did not reveal any mitochondrial fission

and fusion mediators, it is still possible that changes in mitochondrial proteins in the post-

synaptic dendrites contribute to behavioral differences. Future studies will isolate potential pre-

versus postsynaptic mechanisms.

Most mitochondrial proteins are nuclear-encoded and must be transported to

mitochondria for organelle-coupled translation (Williams et al., 2014). RBPs play a critical role in

targeting mRNAs to membrane-bound organelles such as mitochondria. RBPs interact with

mRNAs and chaperone them toward mitochondrial outer membranes where they are translated

(Gerber et al., 2004; García-Rodríguez et al., 2007; Eliyahu et al., 2010; Gehrke et al., 2015).

RBPs recognize and bind mitochondria-targeting RNA elements to form higher-order units

called mRNA ribonucleoprotein (RNP) complexes consisting of mRNAs and associated RBPs

(Béthune et al., 2019; Rossoll and Bassell, 2019). The robust increase in hnRNP H protein in

the striatal synaptosome accompanied by the increase in several complex I subunits suggests a

novel function for hnRNP H in targeting mRNAs encoding for subunits of mitochondrial complex

I to the mitochondria, thus, regulating local translation (**Figure 10**). Once the mRNAs are

transported nearby the mitochondria, hnRNP H could coordinate with other RBPs to form an

RNP complex to stabilize mRNAs to the mitochondrial membrane where translational activators

initiate translation. An example of a RBP with such function is Puf3pm that binds to *ATP2*

mRNA encoding mitochondrial components of the F_1F_0 ATPase and localizes to the outer

mitochondrial membrane (García-Rodríguez et al., 2007). Future studies involving cross-linking

immunoprecipitation combined with RNA-seq (CLIP-seq) will identify target mRNAs bound to

hnRNP H and determine the degree of enrichment for mRNAs encoding mitochondrial complex

I subunits.

Mitochondria are abundant at the synapse where they generate ATP for Ca^{2+} buffering,

vesicle release, and recycling (Vos et al., 2010; Devine and Kittler, 2018). Mitochondria consist

of five oxidative phosphorylation complexes (I through V) (Mimaki et al., 2012; Sharma et al.,

2009). Complex I is the first enzyme of the respiratory chain and initiates electron transport

continuing to complex II through IV to generate redox energy for Complex V to produce ATP.

The increase in Complex I subunits in H1+/- mice could increase Complex I activity and

synaptic ATP production to support vesicle fusion, DA release, and DA transport back into the

cells via DAT and Na+/K+ ATPase to counteract MA-induced DA release. We found a Genotype

by Treatment effect on protein levels of F_0F_1 ATP synthase subunits (Atp5a1, ATP5f1 and

Atp5o) of Complex V (**Figure 9**) in the synaptosomal proteome whereby MA decreased these

ATP synthase subunits in H1+/- mice which could decrease ATP production in response to MA

in H1+/- mice and affect extracellular DA levels.

Besides binding and targeting RNAs to the mitochondria for translation, hnRNP H, like

other RBPs, could bind and target proteins via its glycine-rich domain in an activity-dependent

manner (Tiruchinapalli et al., 2008). In an animal model for frontotemporal dementia, ploy(GR)

aggregates (resulting from hexanucleotide repeats in *C9ORF72)* bind to mitochondrial complex

V protein ATP5A1 to increase its ubiquitination and degradation through the proteasome

pathway, thus disrupting mitochondrial function (Choi et al., 2019). The higher level of

synaptosomal hnRNP H protein in H1+/- mice could increase binding to mitochondrial complex I

and V proteins (e.g., via the glycine-rich domain) to prevent degradation, yielding higher protein

levels at baseline (**Figures 8 and 9**). MA administration could then decrease hnRNP H-protein

interactions in H1+/- mice, thus decreasing synaptic mitochondrial proteins and synaptic

function.

Taken together, the opposing effects of MA treatment on synaptic abundance of

mitochondrial complex I proteins between H1+/- and WT mice could represent a mechanism

underlying blunted MA-induced DA release in H1+/- mice. Future studies will focus on the

interaction between hnRNP H and mRNA encoding mitochondrial complex I and V subunits at

the protein-RNA and protein-protein level (pre- and postsynaptically) and determine whether

disruption of such interactions can alter ATP production and DA release.

REFERENCES

776 Adinoff B (2004) Neurobiologic processes in drug reward and addiction. Harv Rev Psychiatry

777   12:305–320.

778 Bain JM, Cho MT, Telegrafi A, Wilson A, Brooks S, Botti C, Gowans G, Autullo LA,

779   Krishnamurthy V, Willing MC, Toler TL, Ben-Zev B, Elpeleg O, Shen Y, Retterer K,

780   Monaghan KG, Chung WK (2016) Variants in HNRNPH2 on the X Chromosome Are

781   Associated with a Neurodevelopmental Disorder in Females. Am J Hum Genet 99:1–7.

782 Bernheim A, See RE, Reichel CM (2016) Chronic methamphetamine self-administration

783   disrupts cortical control of cognition. Neurosci Biobehav Rev 69:36–48.

784 Béthune J, Jansen RP, Feldbrügge M, Zarnack K (2019) Membrane-Associated RNA-Binding

785   Proteins Orchestrate Organelle-Coupled Translation. Trends Cell Biol 29:178–188.

786 Bosch PJ, Peng L, Kivell BM (2015) Proteomics analysis of dorsal striatum reveals changes in

787   synaptosomal proteins following methamphetamine self-administration in rats. PLoS One

788   10:1–17.

789 Burke RE, Lud Cadet J, Kent JD, Karanas AL, Jackson-Lewis V (1990) An assessment of the

790   validity of densitometric measures of striatal tyrosine hydroxylase-positive fibers:

791   relationship to apomorphine-induced rotations in 6-hydroxydopamine lesioned rats. J

792   Neurosci Methods 35:63–73.

793 Chandra R et al. (2017) Drp1 mitochondrial fission in D1 neurons mediates behavioral and

794   cellular plasticity during early cocaine abstinence. Neuron 96:1327–1341.

795 Chin MH, Qian WJ, Wang H, Petyuk VA, Bloom JS, Sforza DM, Laćan G, Liu D, Khan AH,

796   Cantor RM, Bigelow DJ, Melega WP, Camp DG, Smith RD, Smith DJ (2008) Mitochondrial

797   dysfunction, oxidative stress, and apoptosis revealed by proteomic and transcriptomic

798   analyses of the striata in two mouse models of Parkinson's disease. J Proteome Res

799   7:666–677.

800 Choi SY, Lopez-Gonzalez R, Krishnan G, Phillips HL, Li AN, Seeley WW, Yao WD, Almeida S,

801   Gao FB (2019) C9ORF72-ALS/FTD-associated poly(GR) binds Atp5a1 and compromises

mitochondrial function in vivo. Nat Neurosci 22.

Chou M-Y, Rooke N, Turck CW, Black DL (1999) hnRNP H Is a Component of a Splicing

Enhancer Complex That Activates a c-src Alternative Exon in Neuronal Cells. 19:69–77.

Daubner SC, Le T, Wang S (2011) Tyrosine hydroxylase and regulation of dopamine synthesis.

Arch Biochem Biophys 508:1–12.

Devine MJ, Kittler JT (2018) Mitochondria at the neuronal presynapse in health and disease.

Nat Rev Neurosci 19:63–80.

Divakaruni SS, Van Dyke AM, Chandra R, LeGates TA, Contreras M, Dharmasri PA, Higgs HN,

Lobo MK, Thompson SM, Blanpied TA (2018) Long-Term Potentiation Requires a Rapid

Burst of Dendritic Mitochondrial Fission during Induction. Neuron 100:860-875.e7.

Dunkley PR, Jarvie PE, Robinson PJ (2008) A rapid percoll gradient procedure for preparation

of synaptosomes. Nat Protoc 3:1718–1728.

Eastwood EC, Phillips TJ (2014) Morphine intake and the effects of naltrexone and

buprenorphine on the acquisition of methamphetamine intake. Genes, Brain Behav

13:226–235.

Eliyahu E, Pnueli L, Melamed D, Scherrer T, Gerber AP, Pines O, Rapaport D, Arava Y (2010)

Tom20 Mediates Localization of mRNAs to Mitochondria in a Translation-Dependent

Manner. Mol Cell Biol 30:284–294.

Faul F, Erdfelder E, Lang A-G, Buchner A (2007) G*Power 3: A flexible statistical power

analysis program for the social, behavioral, and biomedical sciences. Behav Res Methods

39:175–191.

Faure JJ, Hattingh SM, Stein DJ, Daniels WM (2009) Proteomic analysis reveals differentially

expressed proteins in the rat frontal cortex after methamphetamine treatment. Metab Brain

Dis 24:685–700.

García-Rodríguez LJ, Gay AC, Pon LA (2007) Puf3p, a Pumilio family RNA binding protein,

localizes to mitochondria and regulates mitochondrial biogenesis and motility in budding

yeast. J Cell Biol 176:197–207.

Gehrke S, Wu Z, Klinkenberg M, Sun Y, Auburger G, Guo S, Lu B (2015) PINK1 and Parkin

Control Localized Translation of Respiratory Chain Component mRNAs on Mitochondria

Outer Membrane Stephan. Cell Metab 21:95–108.

George K. Arhin, Monika Boots, Paramjeet S. Bagga CM and JW (2002) Downstream sequence

elements with different affinities for the hnRNP H/H' protein influence the processing

efficiency of mammalian polyadenylation signals. Nucleic Acids Res 30:1842–1850.

Gerber AP, Herschlag D, Brown PO (2004) Extensive association of functionally and

cytotopically related mRNAs with Puf family RNA-binding proteins in yeast. PLoS Biol

2:0342–0354.

Grammatikakis I, Zhang P, Panda AC, Kim J, Maudsley S, Abdelmohsen K, Yang X, Martindale

JL, Motino O, Hutchison ER, Mattson MP, Gorospe M (2016) Alternative Splicing of

Neuronal Differentiation Factor TRF2 Regulated by HNRNPH1/H2. Cell Rep 15:1–9.

Haughey HM, Fleckenstein AE, Metzger RR, Hanson GR (2002) The Effects of

Methamphetamine on Serotonin Transporter Activity. J Neurochem 75:1608–1617.

Honoré B, Rasmussen HH, Vorum H, Dejgaard K, Liu X, Gromov P, Madsen P, Gesser B,

Tommerup N, Celis JE (1995) Heterogeneous nuclear ribonucleoproteins H, H′, and F are

members of a ubiquitously expressed subfamily of related but distinct proteins encoded by

genes mapping to different chromosomes. J Biol Chem 270:28780–28789.

Honoré B, Vorum H, Baandrup U (1999) hnRNPs H, H' and F behave differently with respect to

posttranslational cleavage and subcellular localization. FEBS Lett 456:274–280.

Hooks MS, Jones GH, Smith AD, Neill DB, Justice JB (1991) Individual differences in locomotor

activity and sensitization. Pharmacol Biochem Behav 38:467–470.

Hutson CB, Lazo CR, Mortazavi F, Giza CC, Hovda D, Chesselet M-F (2011) Traumatic Brain

Injury in Adult Rats Causes Progressive Nigrostriatal Dopaminergic Cell Loss and

Enhanced Vulnerability to the Pesticide Paraquat. J Neurotrauma 28:1783–1801.

Iwazaki T, Matsumoto I, McGregor IS (2007) Protein expression profile in the striatum of rats

with methamphetamine-induced behavioral sensitization. Proteomics 7:1131–1139.

Iwazaki T, McGregor IS, Matsumoto I (2006) Protein expression profile in the striatum of acute

methamphetamine-treated rats. Brain Res 1097:19–25.

Kamma H, Portman DS, Dreyfuss G (1995) Cell Type-Specific Expression of hnRNP Proteins.

Exp Cell Res 221:187–196.

Katz Y, Wang ET, Airoldi EM, Burge CB (2010) Analysis and design of RNA sequencing

experiments for identifying isoform regulation. Nat Methods 7:1009–1015.

Keleta YB, Martinez JL (2012) Brain Circuits of Methamphetamine Place Reinforcement

Learning: The Role of the Hippocampus-VTA Loop. Brain Behav 2:128–141.

Kirkpatrick SL, Bryant CD (2015) Behavioral architecture of opioid reward and aversion in

C57BL / 6 substrains. Front Behav Neurosci 8:1–11.

Kivell B, Uzelac Z, Sundaramurthy S, Rajamanickam J, Ewald A, Chefer V, Jaligam V, Bolan E,

Simonson B, Annamalai B, Mannangatti P, Prisinzano TE, Gomes I, Devi LA, Jayanthi LD,

Sitte HH, Ramamoorthy S, Shippenberg TS (2014) Salvinorin A regulates dopamine

transporter function via a kappa opioid receptor and ERK1/2-dependent mechanism.

Neuropharmacology 86:228–240.

Lein ES et al. (2007) Genome-wide atlas of gene expression in the adult mouse brain. Nature

445:168–176.

Li Z, Okamoto KI, Hayashi Y, Sheng M (2004) The importance of dendritic mitochondria in the

morphogenesis and plasticity of spines and synapses. Cell 119:873–887.

Liao PC, Kuo YM, Hsu HC, Cherng CG, Yu L (2005) Local proteins associated with

methamphetamine-induced nigrostriatal dopaminergic neurotoxicity. J Neurochem 95:160–

168.

Lipari RN, Park-Lee E, Van Horn S (2016) America's Need for and Receipt of Substance Use

Treatment in 2015.

Lominac KD, McKenna CL, Schwartz LM, Ruiz PN, Wroten MG, Miller BW, Holloway JJ, Travis

KO, Rajasekar G, Maliniak D, Thompson AB, Urman LE, Phillips TJ, Szumlinski KK (2014)

Mesocorticolimbic monoamine correlates of methamphetamine sensitization and

motivation. Front Syst Neurosci 8:1–19.

Lominac KD, Quadir SG, Barrett HM, Mckenna CL, Schwartz LM, Ruiz PN, Wroten MG,

Campbell RR, Miller BW, Holloway JJ, Travis KO, Rajasekar G, Maliniak D, Thompson AB,

Urman LE, Kippin TE, Phillips TJ, Szumlinski KK (2016) Prefrontal glutamate correlates of

methamphetamine sensitization and preference. Eur J Neurosci 43:689–702.

Mimaki M, Wang X, McKenzie M, Thorburn DR, Ryan MT (2012) Understanding mitochondrial

complex I assembly in health and disease. Biochim Biophys Acta - Bioenerg 1817:851–

862.

Njung'e K, Handley SL (1991) Evaluation of marble-burying behavior as a model of anxiety.

Pharmacol Biochem Behav 38:63–67.

Oettinghaus B, Schulz JM, Restelli LM, Licci M, Savoia C, Schmidt A, Schmitt K, Grimm A,

Morè L, Hench J, Tolnay M, Eckert A, D'adamo P, Franken P, Ishihara N, Mihara K,

Bischofberger J, Scorrano L, Frank S (2016) Synaptic dysfunction, memory deficits and

hippocampal atrophy due to ablation of mitochondrial fission in adult forebrain neurons.

Cell Death Differ 23:18–28.

Paxinos G, Franklin KBJ (2001) Paxinos and Franklin's the Mouse Brain in Stereotaxic

Coordinates, 2nd ed. Academic Press.

Porsolt RD, Le Pichon M, Jalfre M (1977) Depression: a new animal model sensitive to

antidepressant treatments. Nature 266:730–732.

Prus AJ, James JR, Rosecrans JA (2009) Conditioned Place Preference. In: Methods of

Behavior Analysis in Neuroscience, 2nd ed. (Buccafusco J, ed). Boca Raton, FL: CRC

Press/Taylor & Francis.

Reimand J, Isserlin R, Voisin V, Kucera M, Tannus-Lopes C, Rostamianfar A, Wadi L, Meyer M,

Wong J, Xu C, Merico D, Bader GD (2019) Pathway enrichment analysis and visualization

of omics data using g:Profiler, GSEA, Cytoscape and EnrichmentMap. Nat Protoc 14:482–

517.

Rossoll W, Bassell GJ (2019) Crosstalk of Local Translation and Mitochondria: Powering

Plasticity in Axons and Dendrites. Neuron 101:204–206.

Rothman RB, Baumann MH, Dersch CM, Romero D V., Rice KC, Carroll FI, Partilla JS (2001)

Amphetamine-type central nervous system stimulants release norepinephrine more

potently than they release dopamine and serotonin. Synapse 39:32–41.

Segal DS, Kuczenski R (1997) Repeated binge exposures to amphetamine and

methamphetamine: behavioral and neurochemical characterization. J Pharmacol Exp Ther

282:561–573.

Sharma LK, Lu J, Bai Y (2009) Mitochondrial Respiratory Complex I: Structure, Function and

Implication in Human Diseases. Curr Med Chem 16:1266–1277.

Song KY, Choi HS, Law PY, Wei LN, Loh HH (2016) Post-Transcriptional Regulation of the

Human Mu-Opioid Receptor (MOR) by Morphine-Induced RNA Binding Proteins hnRNP K

and PCBP1. J Cell Physiol:1–9.

Staal RGW, Ray S, Sulzer D (2007) Terminals Amperometric Detection of Dopamine Exocytosis

from Synaptic Introduction to Dopamine Neurotransmission. In: Electrochemical Methods

for Neuroscience. (Michael A, Borland L, eds). Boca Raton, FL: CRC Press/Taylor &

Francis.

Szumlinski KK, Lominac KD, Campbell RR, Cohen M, Fultz EK, Brown CN, Miller BW, Quadir

SG, Martin D, Thompson AB, Jonquieres G von, Klugmann M, Phillips TJ, Kippin TE

(2017) Methamphetamine addiction vulnerability: The glutamate, the bad and the ugly. Biol

Psychiatry 81:959–970.

Szumlinski KK, Lominac KD, Kleschen MJ, Oleson EB, Dehoff MH, Schwartz MK, Seeberg PH,

Worley PF, Kalivas PW (2005) Behavioral and neurochemical phenotyping of Homer1

mutant mice: Possible relevance to schizophrenia. Genes, Brain Behav 4:273–288.

Tiruchinapalli DM, Ehlers MD, Keene JD (2008) Activity-dependent expression of RNA binding

protein HuD and its association with mRNAs in neurons. RNA Biol 5:157–168.

Uhl GR, Drgonova J, Hall FS (2014) Curious cases: Altered dose-response relationships in

addiction genetics. Pharmacol Ther 141:335–346.

Van Dusen CM, Yee L, McNally LM, McNally MT (2010) A glycine-rich domain of hnRNP H/F

promotes nucleocytoplasmic shuttling and nuclear import through an interaction with

transportin 1. Mol Cell Biol 30:2552–2562.

Vos M, Lauwers E, Verstreken P (2010) Synaptic mitochondria in synaptic transmission and

organization of vesicle pools in health and disease. Front Synaptic Neurosci 2:1–10.

Wang E, Aslanzadeh V, Papa F, Zhu H, de la Grange P, Cambi F (2012) Global Profiling of

Alternative Splicing Events and Gene Expression Regulated by hnRNPH/F. PLoS One

7:e52166.

Wang E, Dimova N, Cambi F (2007) PLP/DM20 ratio is regulated by hnRNPH and F and a

novel G-rich enhancer in oligodendrocytes. Nucleic Acids Res 35:4164–4178.

Williams CC, Jan CH, Weissman JS (2014) Targeting and plasticity of mitochondrial proteins

revealed by proximity-specific ribosome profiling. Science (80-) 346:748–751.

Witten JT, Ule J (2011) Understanding splicing regulation through RNA splicing maps. Trends

Genet 27:89–97.

Yamamoto DJ, Nelson AM, Mandt BH, Larson GA, Rorabaugh JM, Ng CMC, Barcomb KM,

Richards TL, Allen RM, Zahniser NR (2013) Rats classified as low or high cocaine

locomotor responders: A unique model involving striatal dopamine transporters that

predicts cocaine addiction-like behaviors. Neurosci Biobehav Rev 37:1738–1743.

Yazdani N, Parker CC, Shen Y, Reed ER, Guido MA, Kole LA, Kirkpatrick SL, Lim JE, Sokoloff

G, Cheng R, Johnson WE, Palmer AA, Bryant CD (2015) Hnrnph1 Is A Quantitative Trait

Gene for Methamphetamine Sensitivity. PLoS Genet 11:e1005713.

Yoon Y, McKenna MC, Rollins DA, Song M, Nuriel T, Gross SS, Xu G, Glatt CE (2013) Anxiety-

associated alternative polyadenylation of the serotonin transporter mRNA confers

translational regulation by hnRNPK. Proc Natl Acad Sci 110:11624–11629.

Zaniewska M, Filip M, Przegaliński E (2015) The involvement of norepinephrine in behaviors

related to psychostimulant addiction. Curr Neuropharmacol 13:407–418.

FIGURE/TABLE/EXTENDED DATA LEGENDS

Figure 1. Oral MA self-administration in H1+/- mice. H1+/- mice were less sensitive than WT

to the reinforcing effect of MA. H1+/- and WT mice were provided access to 80, 120, 160, 200,

300 and 400 mg/L of MA for a time period of five days per dose. **(A):** The average total active

nose-pokes emitted during five, 1 h, sessions varied as a function of MA Dose [$F_{(5,205)}$ =

23.93, $p < 0.0001$] and Genotype [$F_{(1,41)}$ = 6.33; $p = 0.02$], with no Genotype x Dose

interaction [$F_{(5,205)}$ = 1.582, $p = 0.166$] and no effect of Sex [$F_{(1,41)} < 1$] with H1+/- showing

less active nose pokes at MA doses 160, 200, 300, and 400 mg/l (main effect of Genotype at

each dose: *$p < 0.05$). **(B):** The average total inactive nose-pokes emitted during the five, 1 h,

sessions varied as a function of MA Dose [$F_{(5,205)}$ = 5.84, $p < 0.0001$], but not of Genotype

$[F(1,41) < 1]$, with no Genotype x Dose interaction $[F(5,205) = 1.27, p = 0.28]$ or effect Sex

$[F(1,41) < 1]$. **(C):** The average MA intake (mg/kg/day) exhibited by mice showed a significant

Genotype x Dose interaction $[F(5,205) = 4.47, p < 0.0001]$, with H1+/- mice consuming less MA

than WT mice at 200, 300, and 400mg/l doses of MA (main effect of Genotype at each dose: *p

< 0.05). n = 23 (10 females, 13 males) for H1+/- and n = 22 (9 females, 13 males) for WT.

Figure 2. MA-induced CPP in H1+/- mice. H1+/- mice were less sensitive than WT to the

rewarding effect of MA. **(A):** The genotypic difference in the time spent in the MA-paired side

between Day 8 and 1 (preference in s) varied with MA doses [Genotype x Dose interaction:

$F(2,121) = 3.92, p = 0.023$], with genotypic differences observed at 0.5 mg/kg [$t(45) = -2.13$, *p =

0.039] and 2 mg/kg MA doses [$t(45) = 2.18$, *p = 0.036]. No effect of Sex [$F(1,115) = 0.77, p =

0.381$] or Genotype x Sex interactions [$F(1,115) = 0.04, p = 0.852$] were observed. **(B):** In

examining state-dependent CPP following a challenge dose of MA that was the same as during

training, although a similar pattern of results was observed, there was no significant Genotype x

Dose interaction [$F(2,121) = 1.864, p = 0.16$]. Furthermore, there was no effect of Sex [$F(1,115)$

$= 0.09, p = 0.761$] or Genotype x Sex interactions [$F(1,115) = 0.02, p = 0.882$]. n = 24 (16

females, 8 males), 22 (14 females, 8 males) and 16 (9 female, 7 males) at 0, 0.5 and 2 mg/kg

MA for H1+/-; n = 23 (9 female, 14 males), 23 (12 female, 11 males) and 19 (7 female, 12

males) at 0, 0.5 and 2 mg/kg MA for WT.

Figure 3. *In vivo* microdialysis of MA-induced DA release in H1+/- mice. H1+/- mice

showed blunted MA-induced DA and DOPAC levels compared to WT. Mice were probed on the

contralateral side, perfused with artificial cerebrospinal fluid (aCSF), and administered a MA

challenge of either 0.5 mg/kg or 2 mg/kg (i.p., black arrow) after a 1 h baseline period. **(A-B):**

The capacity of MA to elevate DA in the NAc was blunted in the H1+/- mice, in a manner that

varied with the dose of MA administered [Genotype x Dose x Time: $F(11,572) = 2.31, p = 0.01$; no

Sex effects or interactions, $p > 0.25$]. **(A):** The 0.5 mg/kg MA dose elicited an increase in

extracellular DA in in WT but not in H1+/- mice [Genotype x Time: $F(11,275) = 4.71, p <

0.0001], with H1+/- mice exhibiting significantly lower extracellular DA levels than WT mice at all

time-points post-injection (main effect of Genotype: all *p's < 0.04). **(B):** The 2 mg/kg MA dose

elicited an increase in extracellular DA in both genotypes; however, the temporal patterning and

the magnitude of this rise was distinct between H1+/- and WT mice [Genotype x Time:

$F_{(11,341)} = 5.58$, $p < 0.0001$]. H1+/- mice exhibited lower DA levels compared to WT mice

during the first 20 min post-injection (main effect of Genotype: $p = 0.05$), as well as during the

second half of testing [main effect of Genotype, *p's < 0.025]. **(C-D)** MA dose-dependently

reduced NAc extracellular DOPAC levels [Dose X Time: $F_{(11,539)} = 4.23$, $p < 0.0001$];

however, irrespective of MA Dose or Sex, this reduction was, overall, greater in H1+/- versus

WT mice [main effect of Genotype: $F_{(1,49)}=7.62$, $p = 0.008$; Genotype X Time: $F_{(11,539)} =$

5.61, $p < 0.0001$]. **(C)** 0.5 mg/kg MA induced a lowering of extracellular DOPAC, relative to

baseline levels, with H1+/- mice exhibiting significantly lower DOPAC 80 min post-injection

[Genotype X Time: $F_{(11,308)} = 2.78$, $p = 0.002$; main effect of Genotype, *$p = 0.03$]. **(D):** 2

mg/kg MA induced a lowering of extracellular DOPAC in both genotypes; however, the effect

was amplified in H1+/- mice [Genotype x Time: $F_{(11,275)} = 10.42$, $p < 0.0001$]. Relative to WT

animals, H1+/- mice exhibited lower DA levels during the first 20 min post-injection (main effect

of Genotype: $p = 0.05$), as well as during the second half of testing [main effect of Genotype:

*p's < 0.025]. n=13 (7 females, 6 males) at 0.5 and n=17 (10 females, 7 males) at 2.0 mg/kg MA

for H1+/-; n=14 (8 females, 6 males) at 0.5 mg/kg MA and 17 (10 females, 7 males) at 2 mg/kg

MA for WT.

Figure 4. DA content and uptake and DAT levels in the striatum of H1+/- mice. No changes

in DA or in DAT were detected at baseline in the striatum of H1+/- mice compared to WT mice.

(A) Measurement of biogenic amine content in the striatum showed no difference in the level of

dopamine between H1+/- and WT mice [WT, n = 8; H1+/-, n = 8; $t_{(14)} = -0.30$, $p = 0.8$; unpaired

Student's T-test]. H1+/- also exhibited no change in the levels of other biogenic amines [$p > 0.1$;

unpaired Student's T-test]. **(B)** DAT-mediated DA uptake in the striatum of H1+/- mice. Striatum

was harvested 2 h post-saline or MA injection for DA uptake assay, based on the time course of

the microdialysis data in Figure 3B. There was no genotypic difference in the rate of DA uptake

in response to saline (SAL) or MA treatment [Genotype x Treatment: $F_{(1,14)} = 0.02$, p = 0.903].

(C-D): Immunoblot for DAT level in the striatum. Representative immunoblot for DAT shown in

(C) with quantification shown in **(D)**. There was no change in DAT protein level in the striatum of

H1+/- relative to WT mice [WT, n = 14; H1+/-, n = 11; t(23) = 0.32, p = 0.75; unpaired Student's

T-test]. There was a main effect of Sex where the female (regardless of Genotype) showed a

higher level of DAT compared to that of male mice [female, n = 13; male, n = 12; t(23) = 2.53; *p

= 0.02; unpaired Student's T-test].

Figure 5. TH levels in the midbrain of H1+/- mice. No difference in TH level was detected in

the VTA between H1+/- and WT mice. **(A):** Immunofluorescent staining of hnRNP H (*magenta*)

and TH (*green*) was conducted in coronal midbrain sections (Bregma: -3.28 mm to -3.64 mm)

containing the VTA dopaminergic neurons in adult H1+/- and WT mice. Higher magnification

images in panels (*i*) and (*ii*) demonstrate nuclear expression of hnRNP H across all TH-positive

dopaminergic neurons that we examined. Scale bars represent 200 µM (*top*) and 20 µM

(*bottom)*. **(B):** Representative image showing immunohistochemical DAB staining of TH in

coronal sections of the midbrain region (Bregma: -3.28 mm to -3.64 mm). Scale bars represent

represent 1 mm. **(C-E):** Immunohistochemical DAB staining of TH in the midbrain regions

revealed no genotypic difference in TH optical density **(C)**, in number of TH-positive cells **(D)**, or

in the diameter of TH-positive cells **(E)** between the H1+/- and WT mice [WT, n = 8; H1+/-, n =

8; t(14) < 1, all p's > 0.05; unpaired Student's t-test]. **(F-G):** Immunoblot for TH protein level in

the midbrain. Representative immunoblot for TH shown in **(F)** with quantification shown in **(G)**.

There was no change in TH protein level in the midbrain region of H1+/- relative to WT mice

[WT, n = 10; H1+/-, n = 10; t(18) = 0.67, p = 0.51; unpaired Student's T-test].

Figure 6. TH levels in striatum of H1+/- mice. No difference in TH level was detected in the

striatum between the H1+/- versus WT mice. **(A-F):** Representative images showing

immunohistochemical DAB staining on coronal sections of striatum (Bregma: 1.18 mm to 0.86

mm). **A-C:** dorsal striatum. **D-F:** ventral striatum which includes NAc. Scale bars represent

represent 1 mm. Optical density (OD) analysis revealed a nonsignificant increase in TH OD in

the dorsal striatum [**C:** WT, n = 7; H1+/-, n = 8; t(13) = 2.07, p = 0.10; unpaired Student's T-test]

and a small but significant increase in TH intensity in the ventral striatum of H1+/- compared to

WT mice [**F**: WT, n = 7; H1+/-, n = 8; t(13) = 2.30, *p = 0.04; unpaired Student's T-test]. **(G-H):**

Immunoblot for TH protein level in the striatum. Representative immunoblot for TH shown in **(G)**

with quantification shown in **(H)**. There was no change in TH protein expression in the striatum

of H1+/- compared to WT mice [WT, n = 14; H1+/-, n = 11; t(23) = 0.62, p = 0.54; unpaired

Student's T-test].

Figure 7. MA-induced changes in hnRNP H protein expression in striatal tissue and

striatal synaptosomes of H1+/- mice. An increase in hnRNP H protein level was detected in

the striatal synaptosome of H1+/- versus WT mice but no change in hnRNP H protein from total

striatal tissue. **(A-B):** Representative immunoblot for hnRNP H protein level in the striatal

synaptosome 30-min post saline (SAL) or MA treatment shown in **(A)** with quantification shown

in **(B)**. Genotypic difference in hnRNP H protein level was detected in the striatal synaptosome

of H1+/- and WT mice [main effect of Genotype: F(1,27) = 24.36, p < 0.001; main effect of

Treatment: F(1,27) = 0.23; Genotype x Treatment: F(1,27) = 0.41, p = 0.53]. Collapsing across

Treatment, an increase in hnRNP H protein was noted in the striatal synaptosome of H1+/-

versus WT mice regardless of Treatment [t(29) = -5.06, *p < 0.001, unpaired Student's T-test].

This finding was subsequently validated in multiple replications. **(C-D):** Representative

immunoblot for *total* hnRNP H protein level the striatum 30-min post-SAL or post-MA shown in

(C): with quantification shown in **(D).** There was no change in total striatal level of hnRNP H

relative to WT mice after SAL or MA treatment [Genotype x Treatment: interaction [F(1,20) =

0.37, p = 0.55; main effect of Genotype: F(1,20) = 0.60, main effect of Treatment: F(1,20) =

0.01, p = 0.95].

Figure 8. Proteomic analysis of the striatal synaptosome in H1+/- mice. On Days 1 and 2, mice were injected (i.p.) with saline (SAL) and placed into apparatus for 1 h. On Day 3, mice were injected (i.p.) with 2 mg/kg MA and placed into the apparatus for 30 min and then whole striatum was rapidly harvested via free form dissection. Locomotor activity for all three days is shown in Figure S12. The LIMMA package was used for differential analysis with Genotype and Treatment as factors. A ranked list was generated from the analysis and the fgsea R package was used to perform pre-ranked analysis, with proteins filtered for absolute $\log_2$FC > 0.2 and p < 0.05. **(A):** Proteomic analysis was performed to identify differences in protein abundance in the striatal synaptosome of H1+/- versus WT mice. The volcano plot shows the top differentially expressed proteins in the H1+/- versus WT striatal synaptosome. Proteins that are part of the mitochondrial respiratory complex I are circled in orange. **(B):** Proteomic analysis was performed to examine Genotype x Treatment interactions in protein abundance in the striatal synaptosome of H1+/- versus WT mice. This analysis accounted for baseline differences by examining the difference of the difference between the H1+/- and WT in response to MA: (H1+/-$_{MA}$ − H1+/-$_{SAL}$) − (WT$_{MA}$ − WT$_{SAL}$). Volcano plot showing the top differential expressed proteins between the H1+/- versus WT striatal synaptosome in response MA. Proteins that are part of the mitochondrial respiratory complex I are circled in orange.

Table 1. Differentially expressed proteins in H1+/- versus WT. Table showing the top three gene ontology (GO) terms for the top differential expressed protein (shown in Figure 8A) in the H1+/- versus WT striatal synaptosome.

Table 2. Differentially expressed proteins in [H1+/- $_{(MA)}$ − H1+/- $_{(SAL)}$] − [WT$_{(MA)}$ − WT$_{(SAL)}$]. Table showing the top 3 GO terms for the top differential expressed proteins (shown in Figure 8B) in the H1+/- versus WT striatal synaptosome in response to MA.

Figure 9. Protein expression profiles of select mitochondrial proteins in the striatal synaptosome of H1+/- and WT mice from the proteomic dataset. The protein abundance for the mitochondrial proteins are shown for the four groups: WT (SAL), H1+/- (SAL), WT (MA) and H1+/- (MA). Opposing Genotype x Treatment effects on protein levels are shown for the mitochondrial complex I components **(A-C)** and complex V ATPase subunits **(D-F)**. MA induced a decrease in protein expression of these subunits in H1+/- mice and an increase in WT mice.

Figure 10. Proposed model linking increased synaptic mitochondria with a decrease in MA-induced DA release and motivated behaviors. Compared to WT mice, H1+/- mice showed reduced MA-induced DA release in the NAc and reduced sensitivity to the locomotor stimulant, rewarding, and reinforcing effects of MA. In addition, H1+/- mice show an increase in hnRNP H protein in the striatal synaptosome, with no change in total hnRNP H. This increased localization of synaptosome hnRNP H is associated with an increase in complex I mitochondrial proteins. RBPs such as hnRNP H chaperone mRNAs to membrane-bound organelles such as the mitochondria for translation and subsequent assembly in the mitochondria. In the absence of MA, hnRNP H binds to mRNA transcripts encoding mitochondria proteins and transports them to the surface of the mitochondria. MA administration disrupts hnRNP H-RNA interactions which results in less mitochondrial mRNAs being transported to the mitochondria for translation.

Figure 1-1. Bitter/Quinine taste sensitivity in H1+/- mice. There were no genotypic differences between H1+/- and WT mice in average quinine intake (0.003-0.6 mg/ml) [Quinine effect: $F_{(5,100)} = 44.63$, $p < 0.0001$; Genotype effect: $F_{(1,20)} = 2.07$, $p = 0.17$; Genotype x Quinine interaction: $F_{(5,100)} = 0.50$, $p=0.78$].

Figure 1-2. High-dose MA oral self-administration in H1+/- mice. Acquisition of oral MA self-administration with initial training to respond for 200 mg/mL. Mice were trained to nose-poke under an FR1 schedule of reinforcement (20 s time-out) for 20 uL delivery of a 200 mg/L MA solution over the course of 2 weeks. **(A and B):** While both MA intake and the relative amount

of reinforced responding varied across days [for intake, Day effect: $F_{(13,273)}=8.39$, $p<0.0001$;

for % reinforced responding, $F_{(13,273)}=16.20$, $p<0.0001$], no genotypic differences were

detected for either variable during this phase of testing [no Genotype effects or interactions,

p's>0.50]. Furthermore, there was no difference between the number of WT (2 of 12) versus

H1+/- (3 of 11) mice that failed to meet our acquisition criteria [>10 responses with 70% of

responding directed at the MA-reinforced hole; $\lambda2=0.24$, $p=0.62$].

Figure 2-1. Locomotor activity in the CPP box on Day 1, 8, and 9 for 0 mg/kg (saline,

SAL), 0.5 mg/kg MA, and 2 mg/kg MA in H1+/- mice. For all data shown, mice were allowed

open access to both sides of the apparatus via an open entryway in the middle of the dividing

wall. On Day 1, mice were injected with SAL and assessed for initial preference for the MA-

paired side. On Day 8, mice were injected with SAL and tested for drug-free MA-CPP. On Day

9, mice were tested with either SAL or with the same dose of MA that they received during

training to test for state-dependent MA-CPP. In examining the main effect of Genotype and

interaction between Genotype and Time for each dose, a mixed-model two-way ANOVA with

Genotype (between-subjects) and Time (repeated measure) as factors was used. **(A-C):**

Locomotor activity for Day 1. No genotypic difference was detected for Day 1 locomotor activity

for 0.5 and 2 mg/kg MA [SAL: $F_{(1,45)} = 24.76$, $p = 3.81E-5$; 0.5 mg/kg MA: $F_{(1,43)} = 1.894$, $p =$

0.176; 2 mg/kg: $F_{(1, 33)} = 2.657$, $p = 0.113$]. **(D-F):** Locomotor activity for Day 8. Genotypic

difference was detected for Day 8 locomotor activity for 0.5 mg/kg MA but not for SAL and 2

mg/kg MA [SAL: $F_{(1,45)} = 2.22$, $p = 0.143$; 0.5 mg/kg MA: $F_{(1,43)} = 6.117$, $p = 0.017$; 2 mg/kg:

$F_{(1, 33)} = 2.897$, $p = 0.1$]. **(G-I):** Locomotor activity for Day 9. No genotypic difference was

detected for Day 9 locomotor activity for the SAL and 0.5 mg/kg MA dose [SAL: $F_{(1,44)} = 2.956$,

$p = 0.1$; 0.5 mg/kg MA: $F_{(1,42)} = 0.31$, $p = 0.58$]. In response to 2 mg/kg MA, H1+/- showed a

trend toward less total distance over the 1 h period compared to WT [main effect of Genotype:

$F_{(1,33)} = 3.032$, $p = 0.091$]. Sample sizes are indicated in the parentheses.

Figure 3-1. DA no net flux, baseline DA levels and probe placements in H1+/- mice. (A):

DA no net flux in H1+/- and WT mice revealed no genotypic differences at the point of no DA-

flux within the NAc [Genotype by Sex ANOVA, all p's > 0.35; y = 0: 5.00 ± 0.44 for WT (n = 16)

versus 4.68 ± 0.70 for H1+/- (n = 16)]. Although we failed to detect a genotypic difference in the

extraction fraction (slope; Genotype and interaction, p's>0.10], we detected a main effect of Sex

[F(1,31) = 12.09, p = 0.002] that reflected a greater extraction fraction in females (0.94 ± 0.02)

versus males (0.69 ± 0.04). These data do not support an effect of *Hnrnph1* deletion on basal

extracellular DA content (y = 0) or basal DA release/reuptake (extraction fraction) associated

with the reduced MA-induced DA release observed in H1+/- mice. **(B):** Summary of the average

extracellular DA content (nM) prior to an acute injection of MA in WT and H1+/- mice. When

assessed using conventional *in vivo* microdialysis procedures, we failed to detect any Genotype

or Sex differences in the average basal extracellular DA content, prior to MA injection (no

Genotype or Sex effects, nor any Genotype or Sex interactions, p's>0.40). Probe recovery was

lower in the 2 mg/kg MA study than it was in the 0.5 mg/kg study [Dose effect: F(1,33) = 47.83,

p < 0 .0001]. Sample sizes are indicated in parentheses. **(C):** Probe placements for WT and

H1+/- mice fall primarily within the NAc shell (Bregma: 1.54 mm to 1.18 mm).

Figure 3-2. *In vivo* microdialysis of MA-induced glutamate release in H1+/- mice. (A):

Linear regression analyses of the results of a glutamate no net-flux *in vivo* microdialysis study

(0, 2.5, 5 and 10 µM glutamate) revealed no genotypic differences in the x-intercept (an

estimate of basal extracellular glutamate content in µM; WT: 2.28 ± 0.56, n = 7 versus H1+/-

2.55 ± 0.18, n = 8) and no difference in the extraction fraction or slope of the regression (an

index of neurotransmitter release/clearance; WT: 0.73 ± 0.07 versus H1+/-: 0.80 ± 0.04). **(B-C):**

No genotypic or sex differences were detected in average baseline extracellular glutamate

levels, determined under conventional *in vivo* microanalysis procedures (Genotype x Sex x

Dose ANOVA, all p's>0.09; WT=3.27±0.39 µM, n=32 versus H1+/-=3.67±0.32 µM, n=30). Data

are expressed as the percent change from the average baseline value to better visualize the

time-course of the glutamate response to an acute injection of **(B)** 0.5 or **(C)** 2 mg/kg MA.

Figure 3-3. Behavioral test battery in H1+/- mice. (A): Timeline of behavioral battery testing in

drug-naïve H1+/- (n = 38) and wild-type (WT) mice. **(B):** Prepulse inhibition of acoustic startle

revealed no genotypic differences in the PPI response to 75Hz [*left*; t(69) = 1.26, p = 0.21] or

95Hz [*right*; t(69) < 1]. **(C):** Novel object test revealed no genotypic differences in novel object

contact number [*left*; t(69) < 1] or contact time [*right*; t(69) = 1.62, p = 0.11]. **(D):** Marble burying

test revealed no genotypic differences in the amount of time (s) spent burying [t(69) < 1]. **(E):**

Light/dark shuttle box test revealed no genotypic differences in the amount of time (s) spent in

the light zone [t(69) = 1.48, p = 0.15]. **(F):** Porsolt swim test revealed no genotypic differences in

the amount of time immobile [t(69) < 1]. **(G):** In the accelerating rotorod test, there were no

genotypic differences in the ten-trial average time spent on the rotorod [t(69) < 1].

Figure 4-1. MA-induced changes in DAT protein level in the striatal synaptosome of H1+/-

mice. There was no change in DAT protein level in the striatal synaptosome of H1+/- mice **(A):**

Immunoblot showing DAT protein level the striatal synaptosome at 30-min post saline (SAL) or

MA treatment. **(B):** The striatal synaptosome of H1+/- mice exhibited no change in DAT relative

to WT mice after SAL or MA treatment. A two-way ANOVA (Genotype x Treatment) indicated a

no effect of Genotype [F(1,20) = 0.031, p = 0.86], no effect of Treatment [F(1,20) = 0.037, p =

0.85], and no significant interaction between Genotype and Treatment [F(1,20) = 1.24, p = 0.28].

Figure 5-1. Schematics showing Bregma positions of the brain regions in IHC studies

and Western blots. (A): VTA and SNc of midbrain. Left: schematic showing the ventral

midbrain region dissected for Western blot analysis. Right: coronal brain diagrams showing

highlighted VTA (yellow) and SNc (teal) for IHC. **(B)**: Coronal brain diagrams showing

highlighted dorsal striatum (green) and NAc (yellow) for IHC.

Figure 6-1. Schematics and puncta count of TH IHC staining in H1+/- mice. (A): A 225,000

x 225,000 µm grid was overlaid onto the images in Image J, and every third field (indicated by

the black dots) was graded for total number of puncta. Regions of interest were graded by

subtracting the background, setting a threshold, creating a binary image, and conducting

particle analysis to count total puncta number based on roundness and total size (as indicated

by threshold black/white image on bottom right). **(B):** Stereological analysis on number of TH-

positive puncta revealed no genotypic difference on the number of dopaminergic terminals in

the dorsal and ventral striatum of H1+/- relative to WT mice [WT, n = 8; H1+/-, n = 8; t(14) < 1].

Figure 7-1. *Hnrnph1* **whole brain mRNA and hnRNP H whole body protein expression in**

WT, H1+/-, and H1-/- mice. H1-/- mice and WT littermates were generated by intercrossing

H1+/- and H1+/-. Embryos were harvested at E12 for genotyping using a restriction enzyme-

based assay. **(A):** A PCR amplicon capturing the deleted region was digested with BstNI. WT

mice had two copies of two BstNI restriction sites, and thus, restriction digest produced three

fragments (58 bp, 157 bp, and 153 bp) corresponding to two bands on the gel. H1-/- mice had

two copies of a single BstNI restriction sites, and thus, restriction digest produced two fragments

(153 bp and 198 bp). H1+/- mice possessed one copy of each of the two BstNI restriction sites

and one copy of a single BstNI restriction site, and thus, restriction digested produced 5

fragments (58 bp, 153 bp, 153 bp, 157 bp, and 198 bp) corresponding to three bands on the gel.

(B): There was a non-significant, gene dosage-dependent increase in the transcript level

Hnrnph1 in H1+/- and H1-/- mice [$F_{(2,9)} = 2.161$, p = 0.18; one-way ANOVA]. The 1.5-increase

in *Hnnph1* transcript level in H1+/- mice replicated our previously published data (Yazdani et al.

2015). The > 2-fold increase in *Hnrnph1* transcript level in H1-/- with two copies of the mutation

provides further functional support for increased expression of the mutant transcript [WT versus

H1-/-: t(6) = -2.51, **#**p = 0.05]. **(C):** There was no genotypic difference in *Hnrnph2* transcript level

[$F_{(2,9)} = 1.33$, p = 0.32; one-way ANOVA]. **(D-G):** Protein expression of hnRNP H in WT, H1+/-,

and H1-/- mice. There was no genotypic difference in hnRNP H protein expression using an

antibody targeting the C-terminus of hnRNP H [**D-E;** $F_{(2,9)} = 0.89$, p =0.44; one-way ANOVA],

nor was there a significant difference using an antibody specific for the N-term of hnRNP H [**F-**

G; $F_{(2,9)} = 1.64$, $p = 0.25$; one-way ANOVA]. n = 4 for each genotype.

Figure 7-2. Quantification of hnRNP H1 and hnRNP H2 protein expression using peptide

information from mass spec of hnRNP H immunoprecipitates from H1+/- and WT

striatum. A separate cohort of animals were used for this study. Co-immunoprecipitation was

formed to pull down hnRNP H and associated proteins. Using the "Similarity" function in the

Scaffold software, peptides exclusive to hnRNP H1 or hnRNP H2 were identified and quantified.

(A): A list is shown for the peptides that are unique to hnRNP H1. **(B):** Quantification of peptides

unique to hnRNP H1. There was a trend for a decrease in peptide GLPWSCSADEVQR in the

striatum of H1+/- versus WT mice [WT, n = 3; H1+/-, n = 3; $t_{(4)} = 1.938$, **#**$p = 0.06$; unpaired

Student's T-test]. Amino acids PWSCS within this peptide are encoded by the deleted region

(GCCCTGGTCCTGCTCC) in exon 4 of *Hnrnph1* in the H1+/- mice. **(C):** Table outlining the

peptides that are unique to hnRNP H2. **(D):** Quantification of hnRNP H2 unique peptides. No

change in unique peptides of hnRNP H2 was observed between WT and H1+/- mice [WT, n = 3;

H1+/-, n = 3; $t_{(4)} < 1$; unpaired Student's T-test].

Figure 8-1. MA-induced locomotor activity in H1+/- and WT mice that were used for

striatal synaptosome harvest for proteomic analysis. On Days 1 and 2, mice were injected

(i.p.) with saline (SAL) and placed into apparatus for 1 h. On Day 3, mice were injected (i.p.)

with 2 mg/kg MA and placed into apparatus for 30 min followed by subsequent removal of the

striatum. **(A-B):** Locomotor activity for Day 1 **(A)** and Day 2 **(B)** for 1 h in 5-min bin. No

difference in total distance traveled across [no interaction between Genotype x Treatment; Day

1: $F_{(1,12)} = 0.17$; Day 2: $F_{(1,12)} = 0.21$]. **(C):** Locomotor activity for Day 3 for 30 min in 5-min

bin. H1+/- and WT showed Genotype x Treatment difference in sensitivity to SAL and MA [main

effect of Genotype: $F_{(1,12)} = 8.53$, $p = 0.013$; main effect of Treatment: $F_{(1,12)} = 77.42$, $p <$

0.001; Genotype x Treatment: $F_{(1,12)} = 5.85$, $p = 0.03$]. In response to MA (2 mg/kg, i.p.), H1+/-

showed less distance traveled compared to WT [main effect of Genotype: $F_{(1,6)} = 7.84$, $p =$

0.03]. There was a significant MA-induced decrease in locomotor activity in the H1+/- mice [t(6)

= -2,46, -2.45, -3.02 for 20, 25 and 30-min time point respectively, *p < 0.05]. However, in

response to SAL, there was no genotypic difference in locomotor activity [main effect of

Genotype: F(1,6) = 0.70], all p's > 0.05].

Figure 8-2. Concentration of MA and amphetamine in the striatum at 30 min post-MA

injection in H1+/- mice. On Days 1 and 2, mice were injected (i.p.) with saline and placed into

apparatus for 1 h. On Day 3, mice were injected (i.p.) with 2 mg/kg MA and placed into

apparatus for 30 min followed by subsequent removal of the striatum. **(A):** No genotypic

difference was detected in the striatal concentration of MA [t(14) = 0.72, p = 0.48; unpaired

Student's T-test]. **(B):** Also, no genotypic difference was detected in the striatal concentration of

amphetamine [t(14) = 0.22, p = 0.83; unpaired Student's T-test].

Figure 8-3. Network and pathway analysis of synaptosomal proteome of H1+/- versus WT

mice treated with MA or SAL [(H1+/-$_{MA}$ − H1+/-$_{SAL}$) − (WT$_{MA}$ − WT$_{SAL}$)]. Enrichment results for

the comparisons were visualized using Cytoscape and EnrichmentMap. Pathways were

clustered and annotated with themes using AutoAnnotate. Nodes (grey circle) represent the

pathways and the size of each node represents the number of proteins. Each circle within each

node represents the individual protein (red is up and blue is down relative to WT). An

enrichment for metabolic processes is detected in the H1+/- mice and WT in response to MA in

which MA decreases the expression of proteins involved in metabolic processes in the H1+/-

mice. The cutoff for the differential analysis is set to absolute log$_2$FC = 0.2 and p < 0.05.

Figure 9-1. Immunoblots of select mitochondrial proteins in the synaptosome of H1+/-

and WT mice. Three mitochondrial proteins (NDUFS2, ATP5A, a d ATP5F1) were selected for

independent validation of mass spectrometry results in a separate sample cohort via

immunoblot. **(A):** Immunoblots for NDUFS2, ATP5A1 and ATP5F1. **(B-D):** Quantification of

protein expression of three select mitochondrial proteins. All three proteins show the same trend

as the mass spectrometry results depicted in Figure 9 where an increase in expression was

detected in the SAL-treated H1+/- mice. **(B):** NDUFS2: $t(6) = -1.39$ $p = 0.11$; **(C)** ATP5A1: $t(6) =$

-1.64, $p = 0.08$; **(D)** ATP5F1: $t(6) = -1.24$, $p = 0.13$]. n = 4 per genotype per dose.

1294

1295

A

Active Nose Pokes

* WT (n = 22)
○ H1+/- (n = 23)

MA dose (mg/l)

B

Inactive Nose Pokes

MA dose (mg/l)

C

Intake (mg/kg)

MA dose (mg/l)

A **Drug-free**

B **MA-state**

MA-induced DA release: 0.5 and 2 mg/kg (i.p.)

A 0.5 mg/kg

B 2 mg/kg

MA-induced DOPAC release: 0.5 and 2 mg/kg (i.p.)

C 0.5 mg/kg

D 2 mg/kg

A H1+/− vs WT

B [H1+/−$_{MA}$ − H1+/−$_{SAL}$] − [WT$_{MA}$ − WT$_{SAL}$]

Table 1. Differentially expressed proteins in H1+/- vs WT.

GO Term	p value	FDR	Proteins
mitochondrial respiratory chain complex I assembly	2.1E-12	1.4E-09	SAMM50, NDUFB10, NDUFAF6, NDUFB3, NDUFA1, NDUFAF2, NDUFV3, NDUFV1
Complex I biogenesis	1.7E-11	1.6E-09	NDUFB10, NDUFAF6, NDUFB3, NDUFA1, NDUFAF2, NDUFV3, NDUFV1
Respiratory electron transport	2.3E-11	1.6E-09	NDUFB10, NDUFAF6, NDUFB3, NDUFA1, NDUFAF2, NDUFV3, NDUFV1, COX5A

Table 2. Differentially expressed proteins in [H1+/- $_{(MA)}$ – H1+/- $_{(SAL)}$] – [WT$_{(MA)}$ – WT$_{(SAL)}$].

GO Term	p value	FDR	Proteins
Metabolic pathways	4.2E-10	1.7E-08	NDUFB10, NDUFB3, ATP5A1, MBOAT2, HSD17B4, ATP5F1, ATP5O, AGPAT1, UGT8, GANAB, MTHFD1L, ACOX1, AGPS, NDUFS1, ALDH18A1
Parkinson's disease	3.6E-09	7.0E-08	NDUFB10, NDUFB3, ATP5A1, NDUFS1, VDAC1, ATP5F1, ATP5O
mitochondrial ATP synthesis coupled proton transport	4.1E-06	1.2E-3	ATP5A1, SLC25A20, ATP5F1, ATP5O