Title

Human footprint differentially impacts genetic connectivity of four wide-ranging mammals in a

fragmented landscape

Authors

Prachi Thatte[1*], Anuradha Chandramouli[1], Abhinav Tyagi[1], Kaushal Patel[1], Phulmani Baro[1],

Himanshu Chhattani[1] and Uma Ramakrishnan[1]

Affiliations

1- National Center for Biological Sciences, Tata Institute of Fundamental Research

Corresponding author: prachi.thatte@gmail.com

National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bellary Road,
Bangalore 560065. Phone number: +91 80 23666030/31

Running title

Multi-species connectivity

Abstract

Aim

Maintaining connectivity is critical for long-term persistence of wild carnivores in landscapes fragmented due to anthropogenic activity. We examined spatial genetic structure and the impact of landscape features on connectivity in four wide-spread species- jungle cat (*Felis chaus*), leopard (*Panthera pardus*), sloth bear (*Melursus ursinus*) and tiger (*Panthera tigris*).

Location

Our study was carried out in the central Indian landscape, a stronghold in terms of distribution and abundance of large mammals. The landscape comprises fragmented forests embedded in a heterogeneous matrix of multiple land-use types.

Methods

Microsatellite data from non-invasively sampled individuals (90 jungle cats, 82 leopards, 104 sloth bears and 117 tigers) were used to investigate genetic differentiation. Impact of landscape features on gene flow was inferred using a multi-model landscape resistance optimization approach.

Results

All four study species revealed significant isolation by distance (IBD). The correlation between genetic and geographic distance was significant only over a short distance for jungle cat, followed by longer distances for sloth bear, leopard and tiger. Overall, human footprint had aa high negative impact on geneflow in tigers, followed by leopards, sloth bears and the least on

jungle cats. Individual landscape variables- land-use, human population density, density of linear features and roads- impacted the study species differently. Land-use was found to be the most important variable explaining genetic connectivity for all four species, although the scale at which it impacted them differed.

Main conclusions

As expected from theory, but rarely demonstrated using empirical data, the pattern of spatial autocorrelation of genetic variation scaled with dispersal ability of the study species. Landscape genetic analyses revealed species-specific impact of landscape features and provided insights into interactions between species biology and landscape structure. Our results emphasize the need for incorporating functional connectivity data from multiple species for landscape-level conservation planning.

Keywords: multi-species connectivity, human footprint, resistance surface, landscape genetics, microsatellites

Introduction

Habitat destruction and fragmentation due to human footprint is degrading habitat quality and increasing extinction risk of mammals globally (Crooks et al., 2017). Dispersal and genetic exchange between habitat fragments is critical for reducing extinction risk, maintaining genetic diversity and persistence of sub-divided populations (Moilanen & Nieminen, 2002; Thatte et al., 2018a). Maintaining connectivity is hence recognized as a key factor in conservation and management of endangered mammal species.

It is no surprise therefore that research on connectivity conservation has markedly increased over the last decade (Correa Ayram et al., 2016). However, connectivity studies in mammals have largely focused on a single focal species, usually a large predator (Beier et al., 2008; Segelbacher et al., 2010; McRae et al., 2012; Dudaniec et al., 2013; Joshi et al., 2013; execept in a few recent cases- Dudaniec et al., 2016; Wultsch et al., 2016; Marrotte et al., 2017). Conservation strategies based on information from a single species may not effectively capture varied ecological requirements for dispersal of other sympatric species (Brodie et al., 2015; Gangadharan et al., 2016).

Co-occurring species may respond differently to fragmentation based on the differences in body size, dispersal ability, generation time, area requirement and distribution (Gehring & Swihart, 2003; Davidson et al., 2009). Landscape structure or configuration of habitat patches in a landscape may also determine response to fragmentation. Habitat association and the ability to move through heterogeneous matrix between habitat patches can further affect species response to fragmentation (Kierepka et al., 2016). In summary, connectivity for different

species may vary based on their biology, landscape structure and interaction between the two, even in the context of a single landscape.

Conservation strategies for a landscape need to factor in the different requirements of species to prioritize areas of maximum diversity as well as areas where the most endangered species are likely to persist in the long term. For long-term persistence, areas that facilitate dispersal of multiple species need to be protected. While opinion articles emphasize the importance of multi-species landscape genetics studies when developing guidelines for landscape level conservation and management (Keller et al., 2015; Richardson et al., 2016), relatively few studies have compared connectivity for multiple species (eg. Engler et al., 2014; Dudaniec et al., 2016; Wultsch et al., 2016; Marrotte et al., 2017).

We studied genetic connectivity in four wide-spread species- jungle cat (*Felis chaus*), leopard (*Panthera pardus*), sloth bear (*Melursus ursinus*) and tiger (*Panthera tigris*) in the Central Indian landscape. Classified as a global priority tiger conservation landscape, it is a stronghold in terms of mammal distribution and abundance (Jhala, Y. V.;Qureshi, Q;Gopal 2015). Several studies have investigated tiger connectivity in the landscape (Joshi et al., 2013; Sharma et al., 2013; Yumnam et al., 2014; Reddy et al., 2017; Thatte et al., 2018a). However, there is dearth of information on habitat and population connectivity for other sympatric species (Dutta et al., 2013a, 2015).

Our study species vary in body size, dispersal ability, distribution and habitat association, traits that determine species sensitivity to habitat fragmentation (Davidson et al., 2009; Kierepka et al., 2016). Interestingly, considerations based on the above factors result in

contradictory predictions in terms of expected spatial genetic structure for our study species (discussed below and represented in Figure 1).

Empirical evidence across taxa suggests that organisms with specific habitat requirements (and hence habitat association) are usually characterized by low gene flow resulting in strong genetic differentiation (Stuart-Fox et al., 2001; Zayed et al., 2006; Khimoun et al., 2016; Kierepka et al., 2016; Harvey et al., 2017). Among our study species, tigers and sloth bears show stronger habitat associations (Kitchener & Dugmore, 2000; Ramesh et al., 2012; Joshi et al., 2013; Das et al., 2014; Yumnam et al., 2014; Puri et al., 2015) and hence we expect them to have higher genetic differentiation than jungle cats and leopards (Athreya et al., 2013; Gray et al., 2016; Stein et al., 2016).

Dispersal ability of a species can also impact genetic structure. High dispersal ability of a species is likely to homogenize genetic variation over space. Although species with low dispersal ability are likely to have low geneflow, their abundance in a given area is likely to be high (Jenkins et al., 2007; White et al., 2007), resulting in low impact of genetic drift and hence low genetic differentiation. Species with moderate dispersal ability may show higher genetic differentiation: geneflow may not be high enough to homogenize genetic differentiation and the abundance may not be high enough to minimize differentiation. Among our study species, median dispersal distance (calculated based on body size using allometric scaling equations (Sutherland et al., 2000; Bowman et al., 2002) is lowest for jungle cat (8 km), followed by sloth bear (13 km), leopard (35 km) and tiger (85 km) among the study species. Hence we expect sloth bear and leopard to have higher genetic differentiation than tiger and jungle cat.

Continuous and uniform population distribution has been shown to result in lower genetic differentiation (Cowled et al., 2009; Llorens et al., 2017). Spatial patchiness can restrict gene flow and hence increase genetic differentiation. Sloth bears are the most widely distributed large carnivores in central India, followed by leopards and tigers (Jhala et al., 2015). Hence, we expect tigers to have highest genetic differentiation followed by leopard and sloth bear. Jungle cats are also likely to be widely distributed in central India (Gray et al., 2016) although no research studies have been carried out. Hence we expect low genetic differentiation for jungle cats as well.

In summary, all three factors (habitat association, dispersal ability and distribution) predict jungle cats to have the lowest spatial genetic structure among the study species. Predictions for the other three study species are inconsistent. The above predictions ignore the potentially differential ability of the study species to navigate through human modified landscape. In this paper, we seek to address the genetic structure predictions based on dispersal ability, habitat association and distribution with empirical data on four species and also investigate how landscape features impact their movement.

Methods

Study area and sampling

With 34% forest cover, the central Indian landscape (CIL) is a stronghold in terms of mammal distribution and abundance (Yoganand et al., 2006; Jhala et al., 2011). 8.5 % of the forested area (~3 % of the total study area) is protected. Despite being a global hotspot of carnivore habitat, the central Indian landscape has been changing rapidly (Venter et al., 2016).

Similar to other tropical landscapes in the Americas, Asia and Africa, central India faces the threat of further habitat degradation due to anthropogenic activities (Crooks et al., 2011). Being a mineral-rich area, patches of forest have been cleared for mining and the landscape is dotted with mines. There are several railway lines and national and state highways crisscrossing through the landscape. There are five major urban centers (with a population of >1 million) and several big towns. Villages surrounded by agricultural fields cover ~64% of the land area.

We collected non-invasive scat samples (n=1,025), between November 2012 and April 2017, from forested areas inside and outside protected areas. Existing roads/trails were used to search for potential tiger, sloth bear, leopard and jungle cat scats and fresh scats were collected. Each road/trail was sampled only once so as to reduce recaptures and maximize the area covered.

We sampled nine protected areas: out of which (1) Kanha Tiger Reserve (KTR), (2) Pench Tiger Reserve (PTR), (3) Bandhavgarh Tiger Reserve (BTR), (4) Achanakmar Tiger Reserve (ATR), (5) Nagzira-Nawegaon Tiger Reserve(NN), (6) Satpura Tiger Reserve (STR), (7) Sitanadi-Udanti Tiger Reserve (S-U) and (8)Panna Tiger Reserve (PAN) are International Union for the Conservation of Nature (IUCN) category II protected areas and (9) Tipeshwar Wildlife Sanctuary (TIP) which is a IUCN category IV PA. Nine territorial forest divisions outside these protected areas that fall under IUCN categories IV and VI: (1) North and South Balaghat Forest Divisions (BAL), (2) Bramhapuri Forest Division (BPR), (3) North and South Gondia Forest Divisions (GON), (4) Bhandara Forest Division (BHA), (5) Kawardha Forest Division (KAW), (6) Khairagarh Forest Division (KHA) and (7) South Seoni Forest Division (SEO) were also sampled. Tiger samples collected and used in this study were also a part of Thatte et al. (2018).The scat samples collected

between 2012 and 2015 were stored in 30 ml wide mouth bottles containing absolute alcohol. During

2016-17, samples were collected using swabs (HiMedia Inc.) and stored in vials containing Longmire's

buffer (Longmire et al., 1997).

Genotyping and population genetic analysis

In order to quantify genetic connectivity, we first extracted DNA from scat samples using

QIAamp DNA Stool Mini Kit (QIAGEN Inc.) and HiPurA Stool DNA Purification Kit (HiMedia,

India). We then identified species using published primers for tiger, jungle cat and leopard and

a species specific primer that was designed for sloth bear (Farrell et al., 2000; Thatte et al.,

2018b). Jungle cat positive samples were genotyped using 14 felid microsatellite loci (FCA126,

FCA069, FCA672, FCA304, FCA628, FCA279, FCA232, msHDZ170, msFCA506, msFCA453, msF42,

msFCA441, msFCA391, msF41). For leopards the same set of loci were used, except that

msF115 was used instead of msF41. For sloth bears, a panel of 13 bear microsatellite loci was

used (G10H, G10X, G10J, G10C, G10O, UarMu59, UarMu05, UarMu09, UarMu23, UarMu51,

UarMu61, UarMu10 and UarMu64). G10O and UarMu10 were found to be homozygous for all

the samples and hence were not used for analysis. Genotyping details for tigers are reported in

Thatte et al., (2018a). To minimize the impact of genotyping error, the genotyping procedure

was repeated four times (4 independent PCRs) for each sample, for each locus (Mondol et al.,

2009). Genotyping error rates were calculated using GIMLET (Valière, 2002). Individual

identification was conducted using the identity analysis module in the program CERVUS

(Marshall et al., 1998). Further, heterozygosity based differentiation statistics were calculated

using PopGenReport (Adamack & Gruber, 2014), MMOD (Winter, 2012), and HIERFSTAT

(Goudet, 2005) in R (Ihaka & Gentleman, 1996). Inter-individual genetic distances was

calculated based on proportion of shared alleles (D_{PS}; (Bowcock et al., 1994)) using the package

'adegenet' (Jombart, 2008) in R.

Isolation by distance and spatial autocorrelation

Isolation by distance (IBD), the correlation between genetic (D_{PS}) and geographic

(Euclidean) distance, was calculated using 'ecodist' (Goslee & Urban, 2007) package in R. Spatial

autocorrelation analysis was carried out using GenAlEx version 6.5 (Peakall & Smouse, 2012).

Genetic and geographic pairwise distance matrices were correlated for a set of distance classes

(5 km-125 km for jungle cat and 20 km-500 km for the other three species). The correlation

coefficient (r) obtained for each distance class was plotted as a correlogram. The significance of

r was tested against a null hypothesis (r=0) by generating 95% confidence interval for each

distance class (999 permutations, 99 bootstrap repeats).

Landscape genetics analysis

The relationship between the observed genetic structure and the landscape variables

likely to affect the movement of the study species was systematically evaluated using a multi-

model inference and optimization approach (based on Shirk et al., 2010) described in the

following sections.

Landscape variables

We used four landscape variables to build resistance models for all the species- (1) land-use land-cover, (2) human population density, (3) roads and (4) density of linear features (Figure S1). We used BHUVAN land-cover data (http://bhuvan.nrsc.gov.in/bhuvan_links.php) and reclassified it into four broad land-cover types which were ranked in order of increasing resistance: (i) forest- this included deciduous and evergreen forest. Central India is dominated by dry and moist deciduous forests and hence the study area hardly has any pixels classified as evergreen, (ii) Scrub and degraded forest- this category also included orchards and plantations, (iii) agriculture-includes zaid, rabi, kharif croplands, those with two or three crops every year, also fallow and wasteland and (iv) built-up areas- areas of human habitation that is non-agricultural and has buildings, roads and other infrastructure. A layer of human population density, developed at the Foundation for Ecological Research, Advocacy and Learning (FERAL), based on the 2011 census of India data was used for this study. We used population density layer as a continuous variable. A vector layer of national highways, state highways and major public roads was reclassified into 6 categories based on the intensity of traffic on the road (based on Passenger Car Unit (PCU) data for 2006 from the Ministry of Road Transport and Highways). PCU is a measure of the impact of a vehicle on traffic and is calculated by converting vehicle units of different types of vehicles (e.g., trucks, buses, motorcycles) into passenger car units. For example, a motorcycle is equivalent to 0.5 passenger cars, whereas a bus is 3.5 PCUs. Road width (1, 2, 4 or 6 lanes) data was used wherever PCU data was not available. The following criteria was used for classification of roads (Indian Roads Congress, 2007, 2010): minor roads (PCU< 1500), very low traffic roads (2500- 5000 PCU- single lane roads), low traffic

roads (5000- 10000 PCU- intermediate lane roads), moderate traffic roads (10000- 25000 PCU- 2 lane roads), high traffic roads (25000- 55000 PCU- 4 lane roads), very high traffic roads (>55000 PCU- 6 lane roads). In order to calculate the density of linear features, road, railway lines and irrigation canal vector layers were combined into a single poly-lines layer. This layer was developed at FERAL. The density of lines in each cell was calculated to generate a raster layer that was used for analysis.

These four landscape variables were used at 6 different resolutions and the two continuous variables also at 7 different scales for landscape genetics modelling (described in the next section). Studies have shown that the spatial scale at which a landscape variable is measured can also affect the species-landscape relationship (Jackson & Fahrig, 2015). The scale at which a species responds to the surrounding is unknown. The common approach for dealing with the scale problem is analyzing the landscape variables at multiple scales (Krishnamurthy et al., 2016; Zeller et al., 2016). We analyzed all the landscape variables at 6 different spatial resolutions (0.25 km, 0.5 km, 1 km, 2.5 km, 5 km, 10 km). Additionally, the two continuous variables, density of linear features and human population density, were also analysed at multiple scales or ecological neighborhoods (0.25 km, 0.5 km, 1 km, 2.5 km, 5 km, 10 km, 25 km and 50 km) weighted by a Gaussian kernel using the 'kernel2dsmooth' function from the smoothie package in R. These ecological neighborhoods ranged from less than the radius of the home range to more than the median dispersal distance (as suggested by (Jackson & Fahrig, 2015)), except for the tiger for which we did not have a scale of larger than its median dispersal distance. Allometric scaling equations (Bowman et al., 2002) based on body size and trophic level predict the radius of home range and median dispersal distance in the study species to be

0.6 km and 8 km for jungle cat, 1.1 km and 13 km for sloth bear, 2.9 and 35 km for leopard and

6.9 km and 85 km for tiger respectively.

Multi-model optimization

We used a multi-model optimization approach to identify which landscape variables offer

resistance to species movement, the magnitude of resistance (Rmax) for each contributing

variable and the functional form or shape of the relationship between each variable and

resistance (x). Genetic data was used as a response variable to infer the landscape resistance

values and the functional form of the relationship between each variable and resistance. For

each landscape variable, we explored a range of parameter values for Rmax and x and

identified which combination of values best explained the genetic data. For each variable, six

levels of maximum resistance (Rmax = 2, 5, 10, 50, 100, 1000) and six levels of the shape

parameter (x= 0.01, 0.1, 1, 2, 5, 10, Figure S2 represents the transformations) were evaluated.

Shape parameter controlled the functional response between the landscape variable and

resistance to gene flow– 1 meant a linear relationship while values less than and greater than 1

were non-linear. For each combination of Rmax and x, cost-distance between pairs of

individuals was calculated. Cost-distance is the modified Euclidean distance between two points

that accounts for the resistance offered by the landscape. Two methods were used to relate

genetic distance and cost distance matrices and select the best fitting models- (1) partial

Mantel's test (Smouse et al., 1986), controlling for Euclidean distance and (2) linear mixed

effects model with a maximum likelihood population effects parameterization (Clarke et al.,

2002) fitted to the data using MLPE.lmm function from the ResistanceGA package (Peterman,

2018). The best fitting model was identified as the one with the highest correlation in the partial Mantel's test and the one with the lowest AIC score from the MLPE mixed effects model for each landscape variable.

For each landscape variable, we identified the scale and the combination of parameters (Rmax and x) of the best fitting model. The best fitting univariate models identified using partial Mantel's test for each landscape variable were combined (additively) and optimized again in a multivariate context (using partial Mantel's test) to account for interactions between different landscape variables for all the study species (Shirk et al., 2010). MLPE models were fitted and multivariate optimization was carried out based on AIC only for tigers. For multivariate optimization, we varied the model parameters (Rmax and x) of one landscape variable at a time, starting with the variable that had the highest correlation in the univariate optimization, while holding the parameter values of the other variables constant. If the optimal parameters changed for a landscape variable in the multivariate context, we held the new model parameters constant and varied the parameters of the next variable. We repeated this process until the parameterization of all the variables stabilized.

Results

Population genetic analysis

Out of the collected scat samples, based on genetic identification in the lab, 298 were tigers, 343 were sloth bears, 165 were jungle cats and 138 leopards. The remaining 81 samples either failed to amplify with any of the primers used or were from non-target species. Based on unique genotypes, we identified 117 tigers, 104 sloth bears, 92 jungle cats and 82 leopards

(Figure 2). 116 of the 117 tiger individuals were also a part of Thatte et al. (2018a). Genotyping

error rate, expected and observed heterozygosity (H_e and H_o respectively) and the probability

of identity (P(ID)) are presented in Table 1. The P(ID) (the probability of two different

individuals having the same genotype) and the more conservative measure Sib P(ID) (PID when

all individuals in the population are assumed to be siblings) was low for all the species,

indicating that even related individuals would have a very low probability of having identical

genotypes. Sloth bear had lower heterozygosity and higher P(ID) among the study species. The

genetic differentiation, estimated as F_{ST} and G_{ST} followed a pattern that was expected based on

the spatial distribution of the species (Figure 3).

Isolation by distance

        All four study species displayed significant correlation between genetic and geographic

distance (IBD for jungle cats r=0.055, p=0.059; for leopards r=0.213, p=0.001; for sloth bears

r=0.061, p=0.056 and for tigers r=0.461, p=0.001). To examine the spatial extent of genetic

structure, spatial autocorrelation analysis was carried out. IBD broke down at different

distances for different species. IBD relationship broke down at distances beyond ~15 km for

jungle cat, followed by ~40 km for sloth bear, ~60 km for leopard and ~140 km for tiger. Figure

4 presents the correlograms showing correlation between genetic and geographic distance (r)

as a function of geographic distance classes.

Landscape genetic analysis

        Overall among the study species, landscape variables were found to have the highest

impact on tigers, followed by leopards, sloth bears and the least on jungle cats. The optimum

parameters (Rmax, x and spatial resolutions) identified based on the two different model selection methods agreed with each other when the impact of the variables was high (high magnitude of correlation and good support based on AIC scores). Land-use land-cover (LULC) was found to be the most important variable explaining genetic connectivity of all the study species. However, the resistance offered by different land-use classes, strength of correlation and the scale at which it impacted the species varied.

For jungle cat, the effect of landscape variables on genetic connectivity was low based on partial Mantel's test and negligible based on AIC scores. None of the landscape models (Table 2) had AIC score that was lower than the IBD model (-8168). Based on Mantel's r, among the landscape variables tested, LULC had the highest magnitude of correlation followed by density of linear features, human population density and roads. The parameter estimates (maximum resistance offered R_{max} and shape parameter (x)) for each landscape variable are presented in Table 2 (univariate optimization) along with the optimum scale at which the variable showed strongest correlation. Shape parameter determines the shape of the relationship between the landscape variable and resistance.

Despite the low correlation, we combined the four variables (density of linear features, traffic intensity on roads, human population density and LULC) additively for multivariate optimization. On combining, the strength of correlation increased, the maximum resistance offered by the landscape variable (R_{max}) and the shape of the relationship between the landscape variable and the resistance offered (x) changed (Table 3). This suggested interaction between the landscape variables. Human population density, on combining the layers, was found to offer no resistance to jungle cat movement and only built-up areas offered resistance

among the land-use classes. Roads were found to have a high negative impact. Figure 5 presents the relationship between the landscape variables and the resistance they offer and Figure 6 presents the optimized resistance surface for jungle cats.

For leopards, when the landscape variables were tested separately, LULC was found to be the most important variable affecting connectivity, followed by roads, density of linear features and human population density based on AIC scores. Based on partial Mantel's tests, LULC was the only variable that was found to affect connectivity (Table 2). The shape parameter (x>1) suggested that the relationship was nonlinear. Non-linear transformations (x >1) indicate that low resistance is offered by smaller and middle values assigned to the variable and the resistance increases steeply with very high values. Both model selection methods identified 10 km as the optimum resolution for the LULC layer, suggesting that forest cover is important at a broad scale.

For sloth bears, LULC was the most important variable based on both the model selection methods. However, the optimum scale of LULC as identified by the two methods was different (2.5 km based on partial Mantel's test and 0.25 km based on LME). Human population density, roads and density of linear features also explained significant variation after controlling for Euclidean distance. However the magnitude of correlation and the support based on AIC scores was lower for these variables.

Combining the layers for multivariate optimization changed the optimum parameters of a few variables, suggesting an interaction between them. LULC, roads and density of linear features retained their high Rmax values and non-linear shape parameters (x< 1), suggesting

high negative impact. Human population density was also found to have an impact. However, its shape parameter (x= 2) suggested that the resistance offered increased gradually with increasing population density.

The effect of landscape on genetic connectivity of tigers was found to be higher than other species. LULC was the most important (highest magnitude of correlation) landscape variable explaining genetic differentiation in univariate optimization, based on both analyses methods. Both methods suggested that LULC is important at a fine scale (0.25 km resolution based on partial Mantel's test and 0.5 km based on LME), while human population density (50 km scale or ecological neighborhood based on both methods) and density of linear features (10 km scale based on Mantel's test and 25 km based on LME) are important at a larger scale for tigers. Density of linear features, human population density and traffic intensity on roads also explained significant variation after accounting for geographic distance (Table 2). These four variables were combined for multivariate optimization.

The resistance offered by the density of linear features became negligible based on both the model selection methods when all four variables were included. Roads retained their high impact, with resistance increasing highly non-linearly with increasing traffic (Rmax=1000; x=5 and x=10 with optimization based on partial Mantel's test and MLPE with AIC as model selection criteria, respectively). The two model selection methods identified different optimum parameters for the human population density variable. Table 3 presents the optimum values for the maximum resistance (R_{max}) and the shape parameter (x) on combining the variables. Figure 6 represents the optimized resistance surfaces for the study species.

Discussion

Understanding how anthropogenic features impact movement and connectivity of wild carnivores is important for their conservation and management. Conservation plans that incorporate data on multiple species are likely to be better than those based on a single umbrella species. Our objective was to understand how genetic variation is currently partitioned and how landscape features affect gene flow in jungle cat, leopard, sloth bear and tiger in the central Indian landscape. While the patterns of isolation by distance were in agreement with expectations based on dispersal ability and density, species' distribution explained global genetic differentiation. Human footprint negatively impacted genetic connectivity in all four study species. The nature and scale of impact of individual landscape features was species specific. Our results provide insights into how species life-history and ecology interact with the landscape features to shape spatial genetic structure, and are important for conservation planning.

Genetic diversity in central India was high for the studied felid species and moderate for the sloth bear. Ours is the first study to estimate genetic diversity based on microsatellites in jungle cats. An earlier study using mitochondrial data on jungle cats in India had also found them to have high genetic diversity (Mukherjee et al., 2010). The only other study on leopards in the landscape (Dutta et al., 2013b) found the observed (H_o) and expected (H_e) heterozygosity to be higher (0.74 and 0.84 respectively), possibly due to higher sample size in their study. Studies on tigers in the landscape have reported heterozygosity levels comparable to what this study finds (H_o=0.52, H_e=0.723- our study; H_o=0.54, H_e=0.81 (Joshi et al., 2013); H_o=0.71, H_e=0.75 (Yumnam et al., 2014); H_o=0.65, H_e=0.81 (Sharma et al., 2012)). The heterozygosity

based on microsatellites was also found to be low (H_e= 0.45) for spectacled bears that are endemic to the Andes mountain range in South America (Ruiz-García et al., 2005). The only other study on sloth bears in the landscape also found them to have moderate heterozygosity (H_o=0.54, H_e=0.66 (Dutta et al., 2015)).

Dispersal ability and density impact IBD patterns

All four study species revealed significant isolation by distance (IBD). IBD relationship broke down (correlation between genetic and geographic distance was not significant) at distances beyond ~15 km for Jungle cat, ~40 km for sloth bear, ~60 km for leopard and ~140 km for tiger. Allometric scaling equations (Bowman et al., 2002) based on body size and trophic level predict the median dispersal distance in the study species to be in the same order- jungle cat (8 km), sloth bear (13 km), leopard (35 km) and tiger (85 km). Based on literature, the density at which the study species occur is also likely to follow the same order, with jungle cats likely to occur at high densities (0.4-1.5/ km^2, Gray et al., 2016), followed by sloth bear (0.3-0.7/ km^2, Garshelis & Smith, 1999)), leopard (0.05-0.3/ km^2, Kalle et al., 2011; Athreya et al., 2013) and tiger (0.04-0.17/ km^2, (Karanth & Nichols, 1998; Karanth et al., 2004; Kalle et al., 2011). The observed pattern of IBD breakdown is in accordance with what one would expect based on Wright's genetic neighborhood size for the species (Wright, 1946). The neighborhood size depends on density of individuals and species dispersal ability and is non-linearly related to the spatial scale of autocorrelation. Although expected, such trend in breakdown of IBD has been rarely demonstrated with empirical data. It is interesting that we observe this trend despite the differential impact of landscape features on species' geneflow. Spatial autocorrelation of genetic variation accrues over time and can remain stable over long periods (Epperson, 2005).

The relatively recent landscape change may have altered the pattern of isolation by distance expected based on a more or less continuous historical distribution for all the study species. However, as observed in a recent analysis by Tucker et al (2018) for dispersal distances in terrestrial mammals, despite high impact of human footprint, body mass and dietary guild may also explain significant variation in spatial autocorrelation of genetic variation.

Distribution shapes spatial genetic structure

The partitioning of genetic variation, as measured using F_{ST}, G_{ST} and Mantel's r, is highest for the tiger, followed by leopard and then sloth bear and jungle cat. The observed spatial genetic structure can be expected based on the distribution of the species in the central Indian landscape. Sloth bears are the most widely distributed large carnivores in central India, followed by leopard and then tiger (Jhala et al., 2015). Jungle cats are also widely distributed in central India. Their habitat association suggests a more or less continuous distribution in the landscape, although no research studies have been carried out. Patchily distributed populations of species are expected to have low gene flow between patches and higher impact of drift (compared to semi-continuously distributed species) that could result in high genetic differentiation (Allendorf & Luikart, 2007). This has been demonstrated in a few recent studies on plants and birds. Two comparative studies on plants find that semi-continuously or continuously distributed populations have lower genetic differentiation than patchily distributed populations (Levy et al., 2016; Llorens et al., 2017). Robin et al. (2015) studied an entire community of montane birds of the Western Ghats in India and found, with a few exceptions, highest genetic differentiation in montane-restricted species while the widespread species showed no genetic differentiation. Although comparative studies in mammals are rare,

continuously distributed populations have been shown to have low genetic differentiation (Cowled et al., 2009; Côté et al., 2012; Brashear et al., 2015), unless the currently continuous population has a history of range shift or decline followed by spatial expansion (Haanes et al., 2011). Ours is the first comparative study in mammals that demonstrates a relationship between distribution and spatial genetic structure.

Human footprint impacts all four species

Landscape genetic analysis revealed that human footprint has an impact on connectivity of all the study species in the landscape. The two model selection methods used for univariate optimization in the study agreed with each other when the impact of the landscape variables was high (high magnitude of correlation and good support based on AIC scores). Figures S3 and S4 represent the relationship between the spatial resolution, model fit and the agreement between the two model selection methods. A recent simulation based study (Shirk et al., 2018) that compared regression based methods for model selection in individual based landscape genetics analyses found that none of the regression based methods is likely to identify the correct model in landscapes with low resistance that approach IBD. This could explain the lack of agreement in the optimum landscape resistance parameters identified for jungle cat and sloth bear by the two model selection methods. Landscape variables, when considered in isolation, had a low to negligible impact on both jungle cat and sloth bear. However, on combining the landscape variables, they explained significant variation in genetic distances in the two species, suggesting interaction between the different landscape features.

LULC was the most important variable that affected genetic connectivity in all four study species. While only built-up areas offered resistance to jungle cat movement, all non-forest areas offered high resistance to sloth bear movement. For tigers and leopards, the resistance increased gradually from degraded forests to agriculture to built-up areas. Density of linear features also had a strong negative impact on the sloth bear. Density of linear features (highway, rivers and canals) has been shown to influence genetic connectivity of roe deer (*Capreolus capreolus,* Coulon et al., 2006) and has also been shown to affect abundance (Beazley et al., 2004; Boulanger & Stenhouse, 2014) and occurrence (Wasserman et al., 2012) of mammals, which in-turn can impact genetic differentiation in a landscape (Weckworth et al., 2013; Llorens et al., 2017).

There has been a lot of focus on the effect of roads on wildlife populations (Van Der Ree et al., 2011; Laurance et al., 2014). Roads have been shown to reduce gene flow in bighorn sheep (*Ovis Canadensis*, (Epps et al., 2005)) and puma (*Puma concolor,* (Ernest et al., 2014)) among others. Most of these studies have been carried out on a single species (but see (Frantz et al., 2012)). Our results revealed differential impact of roads on our study species. All roads, irrespective of the traffic, were found to have a negative effect on sloth bear connectivity. This is unlike what was observed for tigers and jungle cats. Only roads with high intensity of traffic were found to offer high cost to tiger movement (results were consistent with individual based analysis used in this study and population based analysis carried out with a subset of tiger samples from this study in Thatte et al. (2018)). The resistance offered by roads increased gradually for jungle cat with increasing traffic on roads. There are several roads crisscrossing through the central Indian landscape and 70% of them have low to medium traffic intensity.

These low and medium traffic roads may not impact tiger connectivity but have a high negative impact on jungle cats and sloth bears. Our results emphasize the need to plan and install mitigation structures on not only national and state highways with high traffic but also on smaller roads.

Landscape variables impact species at different scales

Recent studies have suggested carrying out landscape genetic analysis at multiple spatial scales (Cushman & Landguth, 2010). Different landscape variables can impact the same species at different scales. For example, a study on western toads found that land-cover variables measured at a fine scale and topography variables at a coarse scale together explain the observed landscape connectivity (Murphy et al., 2010). At the same time, the scale at which a landscape variable affects connectivity can be different for different species. We found land-use to have the highest impact (strength of Mantel's r) on gene flow when it was measured at a coarse scale for leopard and jungle cat (10 km), but at a fine scale (0.25 km) for the tiger and sloth bear. This suggests that as long as there was forest cover at a coarse scale, land-use class at the fine scale mattered less for leopard and jungle cat gene flow. Forest cover distribution mattered for tiger and sloth bear gene flow at a fine scale, suggesting that maintaining well connected forested corridors is essential for connectivity. For sloth bears, LULC and density of linear features at a fine scale (0.25 km and 0.5 km, respectively) and human population density at a large scale (10 km) together explained the genetic structure.

Interactions between species life-history, ecology and landscape structure

We see interesting genetic patterns in our data that are influenced by both species morphology and ecology. We found dispersal ability impacted spatial autocorrelation between genetic and geographic distance in the study species, while distribution seems to explain the genetic differentiation pattern. Often considered in isolation in comparative studies, species life-history and ecological characteristics are likely to interact with each other and with landscape characteristics to shape spatial genetic structure. All three variables- habitat association, dispersal ability and distribution predicted that jungle cats would show the lowest genetic differentiation among the study species and our results corroborated these expectations. Both tigers and sloth bears were expected to show high genetic differentiation based on two of the three variables. While tiger had the highest genetic structure among the study species and a strong effect of landscape features, sloth bears had low genetic differentiation and milder (based on magnitude of partial Mantel's r) effect of landscape features. Sloth bears are likely to have a high effective population size owing to their high densities (Garshelis & Smith, 1999; Ratnayeke et al., 2007), ability to survive in small habitat patches (Bargali et al., 2012) and wide distribution in central India (Jhala et al., 2015). Hence, despite moderate dispersal ability, they are likely to experience low genetic drift resulting in low genetic differentiation. Additionally, sloth bear demographic decline has likely been gradual over the last few decades (Dharaiya et al., 2016). On the other hand, models of tiger demographic history suggest they went through a population bottleneck ~200 years ago and lost 90% of the population (Mondol et al., 2013). The high genetic structure observed in tigers, despite high dispersal ability, could also be due to a combination of demographic history of population decline and their current restricted distribution.

The observed patterns of connectivity in landscape genetic studies are likely to be a result of multiple interacting processes that influence gene flow and genetic drift. There is a dearth of conceptual and theoretical research investigating the interactions between the different variables including dispersal ability, habitat specialization, distribution and demographic history along with landscape characteristics (area and configuration) and their effect on gene flow and genetic drift (Balkenhol et al., 2016). Although outside the scope of the current study, using a combination of empirical data on multiple species and simulations can provide a better understanding of the interacting processes shaping spatial genetic structure.

Conservation implications

Our study finds differential impact of landscape features on gene flow in the four study species and highlights the importance of incorporating species-specific insights while making conservation decisions in a landscape. Conservation tends to focus on large mammals in protected areas. While this is important, it may not be an effective conservation strategy for a widely distributed species like the jungle cat, leopard and sloth bear, which are abundant outside protected areas. For such species, conservation value of non-protected areas and small fragments might be high despite being embedded in a human dominated landscape. The utility of non-protected forests may not be restricted to being 'stepping-stones' which species use to cross from one patch to another. Considering the rapid rate of deforestation and anthropogenic development in forest and non-forest areas, it is important to realize the conservation value of modified habitats (Pardini et al., 2009) and non-protected areas for conservation.

Landscapes of high conservation value, especially in the tropics, have been changing

rapidly due to increasing human population and exploitation of natural resource leading to

conversion and degradation of habitat (Crooks et al., 2011; Elmhagen et al., 2015; Newbold et

al., 2015). Understanding the impact of fragmentation on multiple species and identifying

strategies to maintain connectivity are critical for species persistence into the future. In order

to conserve areas important for dispersal of multiple species, insights gained from assessing

overlap between species-specific connectivity surfaces can be used. However, there are several

challenges that remain. Any planning approach that aims to conserve multiple species will be a

compromise between what is best for each species individually and what is optimum and

feasible when all species are considered together (Early & Thomas, 2007). Very few studies

have tried to incorporate data on multiple species that differ in their connectivity pattern and

dispersal ability into conservation planning exercises (but see Early & Thomas 2007; Magris *et*

al. 2016; Dudaniec *et al.* 2016). Species persistence being one of the key goals of conservation

planning (Pereira et al., 2010), a better conceptual understanding of how species interact with

their landscape in combination with empirical data on multiple species and models that

account for future landscape change would be critical for conservation planning exercises to

ensure survival of multiple species into the future.

Acknowledgements

We are thankful to the Maharashtra, Madhya Pradesh and Chhattisgarh Forest Departments for providing the necessary permissions (Letter numbers: Kramank/Takniki-I/1720, Kramank/Takniki-I/5100 and Kramank/Takniki-I/1468 for Madhya Pradesh; No./Prabandh-71/13/2866 and Kramank/Wa.Pra./Prabandh-71/16/348 for Chhattisgarh; D-22(8)/Research/1344(2012-13) and D-22(8)/WL/CR-97(15-16)/2972/2016-17 for Maharashtra) and logistical support for this research. We sincerely thank the forest department staff, field assistants and volunteers for their help and support during the fieldwork. Two landscape layers

used in this study, human population density and linear features density, were developed by Rajat Naik and Srinivas Vaidyanathan at Foundation for Ecological Research, Advocacy and Learning. We thank Shomita Mukherjee and Srinivas Vaidyanathan for helpful comments on the manuscript. We are grateful to the National Tiger Conservation Authority (NTCA) of India and the DeFries-Bajpai Foundation for partially funding this study. We thank the National Centre for Biological Sciences (NCBS) for institutional support. PT was supported by Council for Industrial and Scientific Research fellowship and NCBS. UR was supported by the DAE outstanding Scientist Award.

References

Adamack, A.T. & Gruber, B. (2014) PopGenReport: simplifying basic population genetic analyses in R. *Methods in Ecology and Evolution*, **5**, 384–387.

Allendorf, F.W. & Luikart, G. (2007) *Conservation and the genetics of populations.* Oxford: Blackwell Publishing,

Athreya, V., Odden, M., Linnell, J.D.C., Krishnaswamy, J., & Karanth, U. (2013) Big Cats in Our Backyards: Persistence of Large Carnivores in a Human Dominated Landscape in India. *PLoS ONE*, **8**, e57872.

Balkenhol, N., Cushman, S.A., Waits, L.P., & Storfer, A. (2016) Current status, future opportunities, and remaining challenges in landscape genetics. *Landscape Genetics: Concepts, Methods, Applications* (ed. by N. Balkenhol, S.A. Cushman, A.T. Storfer, and L.P. Waits), pp. 247–255. John Wiley & Sons, Ltd.,

Bargali, H.S., Bargali, H.S., Akhtar, N., & Chauhan, N.P.S. (2012) The Sloth Bear Activity and Movement in Highly Fragmented and Disturbed Habitat in Central India. *World Journal of Zoology*, **7**, 312–319.

Beazley, K.F., Snaith, T. V, Mackinnon, F., & Colville, D. (2004) Road density and potential impacts on wildlife species such as American moose in mainland Nova Scotia. *Proceedings of the Nova Scotian Institute of Science*, **42**, 339–357.

Beier, P., Majka, D.R., & Spencer, W.D. (2008) Forks in the Road: Choices in Procedures for Designing Wildland Linkages. *Conservation Biology*, **22**, 836–851.

Boulanger, J. & Stenhouse, G.B. (2014) The Impact of Roads on the Demography of Grizzly Bears in Alberta. *PLoS ONE*, **9**, e115535.

Bowcock, A.M., Ruiz-Linares, A., Tomfohrde, J., Minch, E., Kidd, J.R., & Cavalli-Sforza, L.L. (1994) High resolution of human evolutionary trees with polymorphic microsatellites. *Nature*, **368**, 455–457.

Bowman, J., Jaeger, J. a. G., & Fahrig, L. (2002) Dispersal Distance of Mammals Is Proportional To Home Range Size. *Ecology*, **83**, 2049–2055.

Brashear, W.A., Ammerman, L.K., & Dowler, R.C. (2015) Short-distance dispersal and lack of genetic structure in an urban striped skunk population. *Journal of Mammalogy*, **96**, 72–80.

Brodie, J.F., Giordano, A.J., Dickson, B., Hebblewhite, M., Bernard, H., Mohd-Azlan, J., Anderson, J., & Ambu, L. (2015) Evaluating multispecies landscape connectivity in a threatened tropical mammal community. *Conservation Biology*, **29**, 122–132.

Clarke, R.T., Rothery, P., & Raybould, A.F. (2002) Confidence limits for regression relationships between distance matrices: Estimating gene flow with distance. *Journal of Agricultural, Biological, and Environmental Statistics*, **7**, 361–372.

Correa Ayram, C.A., Mendoza, M.E., Etter, A., & Salicrup, D.R.P. (2016) Habitat connectivity in biodiversity conservation. *Progress in Physical Geography: Earth and Environment*, **40**, 7–37.

Côté, H., Garant, D., Robert, K., Mainguy, J., & Pelletier, F. (2012) Genetic structure and rabies spread potential in raccoons: the role of landscape barriers and sex-biased dispersal. *Evolutionary Applications*, **5**, 393–404.

Coulon, A., Guillot, G., Cosson, J.-F., Angibault, J.M.A., Aulagnier, S., Cargnelutti, B., Galan, M., & Hewison, A.J.M. (2006) Genetic structure is influenced by landscape features: empirical evidence from a roe deer population. *Molecular Ecology*, **15**, 1669–1679.

Cowled, B.D., Lapidge, S.J., Hampton, J.O., & SPENCER, P.B.S. (2009) Measuring the Demographic and Genetic Effects of Pest Control in a Highly Persecuted Feral Pig Population. *http://dx.doi.org/10.2193/0022-541X(2006)70[1690:MTDAGE]2.0.CO;2*, .

Crooks, K.R., Burdett, C.L., Theobald, D.M., King, S.R.B., Di Marco, M., Rondinini, C., & Boitani, L. (2017) Quantification of habitat fragmentation reveals extinction risk in terrestrial mammals. *Proceedings of the National Academy of Sciences of the United States of America*, **114**, 7635–7640.

Crooks, K.R., Burdett, C.L., Theobald, D.M., Rondinini, C., & Boitani, L. (2011) Global patterns of fragmentation and connectivity of mammalian carnivore habitat. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, **366**, 2642–2651.

Das, S., Dutta, S., Sen, S., S, J.A., Babu, S., Nagraj Kumara, H., & Singh, M. (2014) Identifying regions for conservation of sloth bears through occupancy modelling in north-eastern Karnataka, India. *Ursus*, **25**, 111–120.

Davidson, A.D., Hamilton, M.J., Boyer, A.G., Brown, J.H., & Ceballos, G. (2009) Multiple ecological pathways to extinction in mammals. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 10702–5.

Dharaiya, N., Bargali, H.S., & Sharp, T. (2016) Available at: http://www.iucnredlist.org/details/13143/0.

Dudaniec, R.Y., Rhodes, J.R., Worthington Wilmer, J., Lyons, M., Lee, K.E., McAlpine, C. a., & Carrick, F.N. (2013) Using multilevel models to identify drivers of landscape-genetic structure among management areas. *Molecular Ecology*, **22**, 3752–3765.

Dudaniec, R.Y., Worthington Wilmer, J., Hanson, J.O., Warren, M., Bell, S., & Rhodes, J.R. (2016) Dealing with uncertainty in landscape genetic resistance models: a case of three co-occurring marsupials. *Molecular Ecology*, **25**, 470–486.

Dutta, T., Sharma, S., Maldonado, J.E., Panwar, H.S., & Seidensticker, J. (2015) Genetic Variation, Structure, and Gene Flow in a Sloth Bear (Melursus ursinus) Meta-Population in the Satpura-Maikal Landscape of Central India. *PLOS ONE*, **10**, e0123384.

Dutta, T., Sharma, S., Maldonado, J.E., Wood, T.C., Panwar, H.S., & Seidensticker, J. (2013a) Fine-scale population genetic structure in a wide-ranging carnivore, the leopard (Panthera pardus fusca) in central India. *Diversity and Distributions*, **19**, 760–771.

Dutta, T., Sharma, S., Maldonado, J.E., Wood, T.C., Panwar, H.S., & Seidensticker, J. (2013b) Fine-scale population genetic structure in a wide-ranging carnivore, the leopard (Panthera

pardus fusca) in central India. *Diversity and Distributions*, **19**, 760–771.

Early, R. & Thomas, C.D. (2007) Multispecies conservation planning: identifying landscapes for the conservation of viable populations using local and continental species priorities. *Journal of Applied Ecology*, **44**, 253–262.

Elmhagen, B., Eriksson, O., & Lindborg, R. (2015) Implications of climate and land-use change for landscape processes, biodiversity, ecosystem services, and governance. *Ambio*, **44**, S1–S5.

Engler, J.O., Balkenhol, N., Filz, K.J., Habel, J.C., & Rödder, D. (2014) Comparative landscape genetics of three closely related sympatric Hesperid butterflies with diverging ecological traits. *PLoS ONE*, **9**, e106526.

Epperson, B.K. (2005) Estimating dispersal from short distance spatial autocorrelation. *Heredity*, **95**, 7–15.

Epps, C.W., Palsbøll, P.J., Wehausen, J.D., Roderick, G.K., Ramey, R.R., & McCullough, D.R. (2005) Highways block gene flow and cause a rapid decline in genetic diversity of desert bighorn sheep. *Ecology Letters*, **8**, 1029–1038.

Ernest, H.B., Vickers, T.W., Morrison, S.A., Buchalski, M.R., & Boyce, W.M. (2014) Fractured genetic connectivity threatens a southern california puma (Puma concolor) population. *PloS one*, **9**, e107985.

Farrell, L.E., Roman, J., & Sunquist, M.E. (2000) Dietary separation of sympatric carnivores identified by molecular analysis of scats. *Molecular Ecology*, **9**, 1583–1590.

Frantz, A.C., Bertouille, S., Eloy, M.C., Licoppe, A., Chaumont, F., & Flamand, M.C. (2012) Comparative landscape genetic analyses show a Belgian motorway to be a gene flow barrier for red deer (Cervus elaphus), but not wild boars (Sus scrofa). *Molecular Ecology*, **21**, 3445–3457.

Gangadharan, A., Vaidyanathan, S., & Clair, C.C.S. (2016) Categorizing species by niche characteristics can clarify conservation planning in rapidly-developing landscapes. *Animal Conservation*, **19**, 451–461.

Garshelis, D.L. & Smith, J.L.D. (1999) ESTIMATING DENSITY AND RELATIVE ABUNDANCE OF SLOTH BEARS. *Ursus*, **11**, 87–98.

Gehring, T.M. & Swihart, R.K. (2003) Body size, niche breadth, and ecologically scaled responses to habitat fragmentation: mammalian predators in an agricultural landscape. *Biological Conservation*, **109**, 283–295.

Goslee, S.C. & Urban, D.L. (2007) The ecodist Package for Dissimilarity-based Analysis of Ecological Data. *Journal of Statistical Software*, **22**, 1–19.

Goudet, J. (2005) hierfstat, a package for r to compute and test hierarchical F-statistics. *Molecular Ecology Notes*, **5**, 184–186.

Gray, T.N.E., Timmins, R.J., Jathana, D., Duckworth, J.W., Baral, H., & Mukherjee, S. (2016) Felis chaus. The IUCN red list of threatened species. .

Haanes, H., Røed, K.H., Solberg, E.J., Herfindal, I., & Sæther, B.-E. (2011) Genetic discontinuities in a continuously distributed and highly mobile ungulate, the Norwegian moose. *Conservation Genetics*, **12**, 1131–1143.

Harvey, M.G., Aleixo, A., Ribas, C.C., & Brumfield, R.T. (2017) Habitat Association Predicts Genetic Diversity and Population Divergence in Amazonian Birds. *The American naturalist*, **190**, 631–648.

Ihaka, R. & Gentleman, R. (1996) R: A Language for Data Analysis and Graphics. *Journal of Computational and Graphical Statistics*, **5**, 299–314.

Indian Roads Congress (2007) IRC SP 073: Manual of Standards and Specifications for Two Laning of State Highways. .

Indian Roads Congress (2010) Four-laning of highways through public private partnership: manual of specifications & standards. .

Jackson, H.B. & Fahrig, L. (2015) Are ecologists conducting research at the optimal scale? *Global Ecology and Biogeography*, **24**, 52–63.

Jenkins, D.G., Brescacin, C.R., Duxbury, C. V., Elliott, J. a., Evans, J. a., Grablow, K.R., Hillegass, M., Lyon, B.N., Metzger, G. a., Olandese, M.L., Pepe, D., Silvers, G. a., Suresch, H.N., Thompson, T.N., Trexler, C.M., Williams, G.E., Williams, N.C., & Williams, S.E. (2007) Does size matter for dispersal distance? *Global Ecology and Biogeography*, **16**, 415–425.

Jhala, Y. V., Qureshi, Q., & Gopal, R. (2015) Status of tigers in India, 2014. .

Jhala, Y. V., Qureshi, Q., Gopal, R., & Sinha, P.R. (2011) Status of tigers,co-predators and prey in India, 2010. .

Jombart, T. (2008) adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics (Oxford, England)*, **24**, 1403–1405.

Joshi, A., Vaidyanathan, S., Mondol, S., Edgaonkar, A., & Ramakrishnan, U. (2013) Connectivity of Tiger (Panthera tigris) Populations in the Human-Influenced Forest Mosaic of Central India. *PLoS ONE*, **8**, e77980.

Kalle, R., Ramesh, T., Qureshi, Q., & Sankar, K. (2011) Density of tiger and leopard in a tropical deciduous forest of Mudumalai Tiger Reserve, southern India, as estimated using photographic capture–recapture sampling. *Acta Theriologica*, **56**, 335–342.

Karanth, K.U., Chundawat, R.S., Nichols, J.D., & Kumar, N.S. (2004) Estimation of tiger densities in the tropical dry forests of Panna, Central India, using photographic capture–recapture sampling. *Animal Conservation*, **7**, 285–290.

Karanth, K.U. & Nichols, J.D. (1998) Estimation of tiger densities in India using photographic captures and recaptures. *Ecology*, **79**, 2852–2862.

Keller, D., Holderegger, R., van Strien, M.J., & Bolliger, J. (2015) How to make landscape genetics beneficial for conservation management? *Conservation Genetics*, **16**, 503–512.

Khimoun, A., Eraud, C., Ollivier, A., Arnoux, E., Rocheteau, V., Bely, M., Lefol, E., Delpuech, M., Carpentier, M.L., Leblond, G., Levesque, A., Charbonnel, A., Faivre, B., & Garnier, S. (2016) Habitat specialization predicts genetic response to fragmentation in tropical birds. *Molecular Ecology*, **25**, 3831–3844.

Kierepka, E.M., Anderson, S.J., Swihart, R.K., Rhodes, O.E., & Jr (2016) Evaluating the influence of life-history characteristics on genetic structure: a comparison of small mammals inhabiting complex agricultural landscapes. *Ecology and evolution*, **6**, 6376–96.

Kitchener, A.C. & Dugmore, A.J. (2000) Biogeographical change in the tiger, Panthera tigris. *Animal Conservation*, **3**, 113–124.

Laurance, W.F., Clements, G.R., Sloan, S., Connell, C.S.O., Mueller, N.D., Goosem, M., Venter, O., Edwards, D.P., Phalan, B., Balmford, A., Ree, R. Van Der, & Arrea, I.B. (2014) A global strategy for road building. *Nature*, .

Levy, E., Byrne, M., Coates, D.J., Macdonald, B.M., McArthur, S., & van Leeuwen, S. (2016) Contrasting Influences of Geographic Range and Distribution of Populations on Patterns of

Genetic Diversity in Two Sympatric Pilbara Acacias. *PLOS ONE*, **11**, e0163995.

Llorens, T.M., Tapper, S.-L., Coates, D.J., McArthur, S., Hankinson, M., & Byrne, M. (2017) Does population distribution matter? Influence of a patchy versus continuous distribution on genetic patterns in a wind-pollinated shrub. *Journal of Biogeography*, **44**, 361–374.

Magris, R.A., Treml, E.A., Pressey, R.L., & Weeks, R. (2016) Integrating multiple species connectivity and habitat quality into conservation planning for coral reefs. *Ecography*, **39**, 649–664.

Marrotte, R.R., Bowman, J., Brown, M.G.C., Cordes, C., Morris, K.Y., Prentice, M.B., & Wilson, P.J. (2017) Multi-species genetic connectivity in a terrestrial habitat network. *Movement Ecology*, **5**, 21.

Marshall, T.C., Slate, J., Kruuk, L.E., & Pemberton, J.M. (1998) Statistical confidence for likelihood-based paternity inference in natural populations. *Molecular ecology*, **7**, 639–655.

McRae, B.H., Hall, S.A., Beier, P., & Theobald, D.M. (2012) Where to restore ecological connectivity? Detecting barriers and quantifying restoration benefits. *PloS one*, **7**, e52604.

Moilanen, A. & Nieminen, M. (2002) Simple connrctivity measures in spatial ecology. *Ecology*, **83**, 1131–1145.

Mondol, S., Bruford, M.W., & Ramakrishnan, U. (2013) Demographic loss, genetic structure and the conservation implications for Indian tigers. *Proceedings. Biological sciences / The Royal Society*, **280**, 20130496.

Mondol, S., Ullas Karanth, K., Samba Kumar, N., Gopalaswamy, A.M., Andheria, A., & Ramakrishnan, U. (2009) Evaluation of non-invasive genetic sampling methods for estimating tiger population size. *Biological Conservation*, **142**, 2350–2360.

Mukherjee, S., Krishnan, A., Tamma, K., Home, C., Navya, R., Joseph, S., Das, A., & Ramakrishnan, U. (2010) Ecology driving genetic variation: a comparative phylogeography of jungle cat (Felis chaus) and leopard cat (Prionailurus bengalensis) in India. *PloS one*, **5**, e13724.

Newbold, T., Hudson, L.N., Hill, S.L.L., et al. (2015) Global effects of land use on local terrestrial biodiversity. *Nature*, **520**, 45–50.

Pardini, R., Faria, D., Accacio, G.M., Laps, R.R., Mariano-Neto, E., Paciencia, M.L.B., Dixo, M., & Baumgarten, J. (2009) The challenge of maintaining Atlantic forest biodiversity: A multi-taxa conservation assessment of specialist and generalist species in an agro-forestry mosaic in southern Bahia. *Biological Conservation*, **142**, 1178–1190.

Peakall, R. & Smouse, P.E. (2012) GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research--an update. *Bioinformatics*, **28**, 2537–2539.

Pereira, H.M., Leadley, P.W., Proença, V., et al. (2010) Scenarios for global biodiversity in the 21st century. *Science (New York, N.Y.)*, **330**, 1496–501.

Peterman, W.E. (2018) ResistanceGA: An R package for the optimization of resistance surfaces using genetic algorithms. *Methods in Ecology and Evolution*, **9**, 1638–1647.

Puri, M., Srivathsa, A., Karanth, K.K., Kumar, N.S., & Karanth, K.U. (2015) Multiscale distribution models for conserving widespread species: The case of sloth bear Melursus ursinus in India. *Diversity and Distributions*, **21**, 1087–1100.

Ramesh, T., Kalle, R., Sankar, K., & Qureshi, Q. (2012) Factors affecting habitat patch use by sloth bears in Mudumalai Tiger Reserve , Western Published by : International Association

for Bear Research and Management Stable URL⊡: http://www.jstor.org/stable/41818973 Linked references are available on JSTOR fo. *Ursus*, **23**, 78–85.

Ratnayeke, S., Manen, F.T. van, & Padmalal, U.K.G.K. (2007) Home Ranges and Habitat Use of Sloth Bears Melursus Ursinus Inornatus in Wasgomuwa National Park, Sri Lanka. *Wildlife Biology*, **13**, 272–284.

Reddy, P.A., Cushman, S.A., Srivastava, A., Sarkar, M.S., & Shivaji, S. (2017) Tiger abundance and gene flow in Central India are driven by disparate combinations of topography and land cover. *Diversity and Distributions*, **23**, 863–874.

Van Der Ree, R., Jaeger, J.A.G., Van Der Grift, E.A., & Clevenger, A.P. (2011) Effects of Roads and Traffic on Wildlife Populations and Landscape Function Effects of Roads and Traffic on Wildlife Populations and Landscape Function: Road Ecology is Moving toward Larger Scales. *Ecology and Society*, **16**, 48.

Richardson, J.L., Brady, S.P., Wang, I.J., & Spear, S.F. (2016) Navigating the pitfalls and promise of landscape genetics. *Molecular Ecology*, **25**, 849–863.

Robin, V. V, Vishnudas, C.K., Gupta, P., & Ramakrishnan, U. (2015) Deep and wide valleys drive nested phylogeographic patterns across a montane bird community. *Proceedings. Biological sciences*, **282**, 20150861.

Ruiz-García, M., Orozco-terWengel, P., Castellanos, A., & Arias, L. (2005) Microsatellite analysis of the spectacled bear (Tremarctos ornatus) across its range distribution. *Genes & genetic systems*, **80**, 57–69.

Segelbacher, G., Cushman, S.A., Epperson, B.K., Fortin, M.-J., Francois, O., Hardy, O.J., Holderegger, R., Taberlet, P., Waits, L.P., & Manel, S. (2010) Applications of landscape genetics in conservation biology: concepts and challenges. *Conservation Genetics*, **11**, 375–385.

Sharma, S., Dutta, T., Maldonado, J.E., Wood, T.C., Panwar, H.S., & Seidensticker, J. (2012) Spatial genetic analysis reveals high connectivity of tiger (Panthera tigris) populations in the Satpura-Maikal landscape of Central India. *Ecology and evolution*, **3**, 48–60.

Sharma, S., Dutta, T., Maldonado, J.E., Wood, T.C., Panwar, H.S., Seidensticker, J., Maldonado, J.E., & Wood, T.C. (2013) Forest corridors maintain historical gene flow in a tiger metapopulation in the highlands of central India. *Proceedings of the Royal Society B: Biological Sciences*, **280**, 20131506.

Shirk, A.J., Landguth, E.L., & Cushman, S.A. (2018) A comparison of regression methods for model selection in individual-based landscape genetic analysis. *Molecular Ecology Resources*, **18**, 55–67.

Shirk, A.J., Wallin, D.O., Cushman, S.A., Rice, C.G., & Warheit, K.I. (2010) Inferring landscape effects on gene flow: a new model selection framework. *Molecular Ecology*, **19**, 3603–3619.

Smouse, P.E., Long, J.C., & Sokal, R.R. (1986) Multiple Regression and Correlation Extensions of the Mantel Test of Matrix Correspondence. *Systematic Zoology*, **35**, 627–632.

Stein, A.B., Athreya, V., Gerngross, P., Balme, G., Henschel, P., Karanth, U., Miquelle, D., Rostro-Garcia, S., Kamler, J.F., Laguardia, A., Khorozyan, I., & Ghoddousi, A. (2016) Panthera pardus. The IUCN red list of threatened species. .

Stuart-Fox, D.M., Schneider, C.J., Moritz, C., & Couper, P.J. (2001) Comparative phylogeography of three rainforest-restricted lizards from mid-east Queensland. *Australian Journal of*

Zoology, **49**, 119.

Sutherland, G.D., Harestad, A.S., Price, K., & Lertzman, K.P. (2000) Scaling of Natal Dispersal Distances in Terrestrial Birds and Mammals. *Conservation Ecology*, **4**, 16.

Thatte, P., Joshi, A., Vaidyanathan, S., Landguth, E., & Ramakrishnan, U. (2018a) Maintaining tiger connectivity and minimizing extinction into the next century: Insights from landscape genetics and spatially-explicit simulations. *Biological Conservation*, **218**, 181–191.

Thatte, P., Patel, K., & Ramakrishnan, U. (2018b) Rapid species identification of sloth bears from non-invasive samples: a PCR based assay. *Ursus*, **29**, .

Thompson, C.M. & McGarigal, K. (2002) The influence of research scale on bald eagle habitat selection along the lower Hudson River, New York (USA). *Landscape Ecology*, **17**, 569–586.

Tucker, M.A., Böhning-Gaese, K., Fagan, W.F., et al. (2018) Moving in the Anthropocene: Global reductions in terrestrial mammalian movements. *Science (New York, N.Y.)*, **359**, 466–469.

Valière, N. (2002) gimlet: a computer program for analysing genetic individual identification data. *Molecular Ecology Notes*, **2**, 377–379.

Venter, O., Sanderson, E.W., Magrach, A., Allan, J.R., Beher, J., Jones, K.R., Possingham, H.P., Laurance, W.F., Wood, P., Fekete, B.M., Levy, M.A., & Watson, J.E.M. (2016) Sixteen years of change in the global terrestrial human footprint and implications for biodiversity conservation. *Nature Communications*, **7**, 12558.

Wasserman, T.N., Cushman, S.A., Wallin, D.O., & Hayden, J. (2012) Multi scale habitat relationships of Martes americana in northern Idaho, U.S.A. .

Weckworth, B. V., Musiani, M., DeCesare, N.J., McDevitt, A.D., Hebblewhite, M., & Mariani, S. (2013) Preferred habitat and effective population size drive landscape genetic patterns in an endangered species. *Proceedings of the Royal Society of London B: Biological Sciences*, **280**, .

White, E.P., Ernest, S.K.M., Kerkhoff, A.J., & Enquist, B.J. (2007) Relationships between body size and abundance in ecology. *Trends in Ecology & Evolution*, **22**, 323–330.

Winter, D.J. (2012) MMOD: an R library for the calculation of population differentiation statistics. *Molecular ecology resources*, **12**, 1158–1160.

Wright, S. (1946) Isolation by distance under diverse systems of mating. *Genetics*, **31**, 39–59.

Wultsch, C., Waits, L.P., Kelly, M.J., Lizama, D., Correa, E., & Cruz, F. (2016) A Comparative Analysis of Genetic Diversity and Structure in Jaguars (Panthera onca), Pumas (Puma concolor), and Ocelots (Leopardus pardalis) in Fragmented Landscapes of a Critical Mesoamerican Linkage Zone. *PLOS ONE*, **11**, e0151043.

Yoganand, K., Rice, C.G., Johnsingh, A.J.T., & Seidensticker, J. (2006) Is the sloth bear in India secure? A priliminary report on distribution, threats and conservation requirements. *Journal of the Bombay Natural History Society*, **103**, 172–181.

Yumnam, B., Jhala, Y. V, Qureshi, Q., Maldonado, J.E., Gopal, R., Saini, S., Srinivas, Y., & Fleischer, R.C. (2014) Prioritizing tiger conservation through landscape genetics and habitat linkages. *PloS one*, **9**, e111207.

Zayed, A., Packer, L., Grixti, J.C., Ruz, L., Owen, R.E., & Toro, H. (2006) Increased genetic differentiation in a specialist versus a generalist bee: implications for conservation. *Conservation Genetics*, **6**, 1017–1026.

Zeller, K.A., McGarigal, K., & Whiteley, A.R. (2012) Estimating landscape resistance to movement: a review. *Landscape Ecology*, **27**, 777–797.

Data accessibility statement

Genotypes of final set of individuals used for analysis will be uploaded on Dryad repository.

Tables

Table 1. Genotyping error and allele frequency based statistics. He and Ho represent the expected and observed heterozygosities, respectively. P(ID) represents the probability of two different individuals having the same genotype and Sib P(ID) is the PID when all individuals in the population are assumed to be siblings.

Species	Allelic Dropout	False alleles	He	Ho	P(ID)	Sib P(ID)
Jungle cat (N= 90)	0.045	0.045	0.794	0.688	5.12×10^{-18}	0.92×10^{-6}
Leopard (N= 82)	0.049	0.018	0.778	0.607	4×10^{-18}	4.1×10^{-7}
Sloth bear (N= 104)	0.060	0.030	0.51	0.392	1.6×10^{-7}	1.4×10^{-3}
Tiger (N= 117)	0.062	0.019	0.723	0.524	1.48×10^{-11}	5.1×10^{-5}

Table2. Univariate optimization results. The table reports optimum parameters (Rmax and x) for each landscape variable based on the two different model selection methods and the respective estimates of model fit (partial Mantel's r and AIC), along with the optimum spatial resolution for LULC and ecological neighborhood (scale) for density of linear features and human population density at which it explained the most amount of variation in genetic differentiation.

Species	Variable	Rmax	x	Partial Mantel's r	Scale	Rmax	x	AIC	Scale
JC	Human density	1000	2	0.013	25 km	1000	1	-8168.29	5 km
	LULC	100	10	0.05	10 km	1000	10	-8168.97	10 km
	Roads	1000	0.01	0.031		1000	0.01	-8168.34	
	Linear density	1000	1	0.044	0.5 km	1000	2	-8169	25 km
LPD	Human density	1000	1	-0.003	10 km	1000	1	-5595.08	10 km
	LULC	1000	2	0.11	10 km	1000	5	-5655.09	10 km
	Roads	1000	0.1	-0.02		1000	0.1	-5586.22	
	Linear density	1000	10	-0.02	15 km	10	0.1	-5578.62	0.5 km
SB	Human density	1000	1	0.050	2.5 km	1000	2	-8791.33	10 km
	LULC	1000	1	0.099	2.5 km	50	2	-8794.12	0.25 km
	Roads	1000	0.1	0.042		5	0.01	-8789.5	

	Linear density	1000	0.1	0.056	0.5 km	1000	2	-8791.17	0.5 km
TIG	Human density	10	1	0.303	50 km	50	2	-9724.41	50 km
	LULC	50	1	0.44	0.25 km	100	2	-10189.8	0.5 km
	Roads	1000	5	0.2963		1000	2	-9522.27	
	Linear density	1000	1	0.402	10 km	1000	2	-9773.94	25 km and 5 km

Table3. Multivariate optimization. The table reports maximum resistance offered by each variable (Rmax) and the shape of the relationship between the value of a variable and the resistance it offers (x=1 implies a linear relationship, x> or < 1 implies a non-linear relationship) on combining with other layers. The correlation coefficient between genetic distance and cost-distance controlling for Euclidean distance (partial Mantel's r), significance of the partial Mantel's test are reported for the combined layers (optimized resistance surfaces).

Species	Variable	Rmax	X	Partial Mantel's r	Significance
Jungle cat	Human density	0	0.01	0.073	0.05
	LULC	50	10		
	Roads	1000	2		
	Linear density	50	2		
Sloth Bear	Human density	100	2	0.119	0.001
	LULC	1000	0.01		
	Roads	500	0.01		
	Linear density	500	0.1		
Tiger*	Human density	500 (10)	10 (0.01)	0.445 (AIC= -10210.3)	0.001
	LULC	500 (1000)	2 (2)		
	Roads	1000 (1000)	5 (10)		
	Linear Density	10 (0)	0.01 (0.01)		

* Optimum parameter values based on model selection using MPLE and AIC value are

represented in parenthesis

Figure Legends

Figure 1. Spatial genetic structure expectations based on habitat association, dispersal ability and distributions of the study species.

Figure 2. Study area and sampling. The figure contains a map of India with the study landscape highlighted and enlarged. In the enlarged study area, protected areas are marked by black outline and genetically identified individuals of study species as coloured dots.

Figure 3. Observed genetic differentiation matched the genetic differentiation pattern expected based on distribution

Figure 4. Spatial autocorrelation. x-axis represents distance classes in km. y-axis is the correlation coefficient (Mantel's r) for the correlation between genetic and geographic distance for samples falling within each distance class. Jungle cats and sloth bears show low correlation (r) and the relationship breaks down at ~20-25 km and ~30-60 km respectively. Leopards and Tigers show higher correlation and the relationship breaks down ~60 km and 140-180 km respectively.

Figure 5. Relationship of landscape features with resistance. x-axis represents each land-use variable used. The categories for roads represent 1- no roads, 2- minor roads, 3- very low traffic roads, 4- low traffic roads, 5- moderate traffic roads, 6- high traffic roads and 7- very high traffic roads. Land-use land-cover categories: 1- forest, 2- degraded forest and scrub, 3- agriculture and 4- built-up. Density of linear features and human population density, being continuous variables, had different maximum values at different resolutions (scales). The optimum scales for different species are different. y-axis is the resistance offered by the different categories/values of the landscape variable.

Figure 6. Optimized resistance surfaces. Dark colours represent high resistance and light colours represent low resistance. Purple dots in each map are the individual locations and green polygons are protected areas. RAT- Ratapani wildlife sanctuary, NOR- Noradehi wildlife sanctuary, BTR- Bandhavgarh tiger reserve, SAN- Sanjay tiger reserve, PAN- Panna tiger reserve, STR- Satpura tiger reserve, ATR- Achanakmar tiger reserve, KTR- Kanha tiger reserve, PTR- Pench tiger reserve, NN- Nagzira Navegaon tiger reserve, BOR- Bor tiger reserve, UK- Umred Karhandla wildlife sanctuary, TATR- Tadoba Andhari tiger reserve, TIP- Tipeshwar tiger reserve, CHH- Chhaprala wildlife sanctuary and SU- Sitanadi Udanti tiger reserve

a Expected based on distribution

Continuous   Restricted

Spatial genetic structure

JC   SB   LPD   TIG

Species

b Observed

Spatial genetic structure

JC   SB   LPD   TIG

Species

(a) Jungle cat

(b) Leopard

(c) Sloth bear

(d) Tiger