Social interactions across groups of colobus monkeys (*Colobus vellerosus*) explain similarities in their gut

microbiomes

Running title: Social networks predict gut microbiome similarity across groups

Eva Wikberg[1#], Diana Christie[2], Pascale Sicotte[3], Nelson Ting[2*]

[1]Department of Anthropology, University of Texas at San Antonio

[2]Department of Anthropology and Institute of Ecology and Evolution, University of Oregon

[3]Department of Anthropology and Archaeology, University of Calgary

[#*]Corresponding authors

[#]Department of Anthropology, University of Texas at San Antonio, One UTSA Circle, San Antonio, TX

78249, USA; eva.wikberg@utsa.edu; 210-458-5722

[*]Department of Anthropology, 1218 University of Oregon, Eugene, OR 97403; nting@uoregon.edu; 541-

346-8878

Abstract

The gut microbiome is structured by social groups in a variety of host taxa. Whether this pattern is

driven by relatedness, similar diets, or shared social environments is under debate because few studies

have had access to the data necessary to disentangle these factors. We investigated whether diet,

relatedness, or 1-meter proximity networks best explains differences in the gut microbiome among 45

female colobus monkeys in 8 groups residing at Boabeng-Fiema, Ghana. We combined demographic and

behavioural data collected May – August 2007 and October 2008 – April 2009 with 16S rRNA sequencing

of faecal samples collected during the latter part of each observation period. Group identity explained a

large percentage of the variation in gut microbiome beta-diversity. When comparing the predictive

power of dietary dissimilarity, relatedness, and distance in the 1-meter proximity network, the models

with social distance received the strongest support. Thus, social proximity plays a larger role than diet

and relatedness in generating group-level gut microbiome signatures, corroborating previous findings of

the importance of social transmission of gut microbes within groups. Our novel findings also show how a

social network metric predicts gut microbiome similarity of non-group members, indicating that

microbes may be transmitted during intergroup encounters. Lastly, some of the socially transmitted gut

microbial taxa are associated with digestion of plant material, but further research is needed to

investigate whether social transmission of gut microbes yields health benefits, which could provide an

incentive for the formation and maintenance of social bonds within and between groups.

Key words: 16S rRNA gene, between-group encounters, colobines, diet, gut microbiome, microbe

transmission, relatedness, social networks, social transmission, host-microbe

Introduction

The gut microbiome consists of thousands of species that affects its host's nutritional status, immune

function, and behavior (McFall-Ngai et al., 2013), and can have important health and fitness

consequences for hosts in wild (Koch & Schmid-Hempel, 2011; Vlčková et al., 2018) and captive settings

(Turnbaugh et al., 2006). Because of these potential fitness effects, it is important to investigate the

acquisition and maintenance of the gut microbiome (Amato, 2016; Archie & Tung, 2015).

The gut microbiome of individuals or groups become more distinct the further apart they live

(Barelli et al., 2015; Grieneisen et al., 2019; Hansen et al., 2019; Hird, Carstens, Cardiff, Dittmann, &

Brumfield, 2014; Lankau, Hong, & Mackie, 2012; Phillips et al., 2012), and the microbiome is structured

by group or family co-residency in a variety of host taxa, such as humans (Lax et al., 2014; Song et al.,

2013; Yatsunenko et al., 2012), non-human primates (Amato et al., 2017; Degnan et al., 2012;

Goodfellow et al., 2019; Orkin, Webb, & Melin, 2019; Springer et al., 2017; Tung et al., 2015), carnivores

(Leclaire, Nielsen, & Drea, 2014; Theis et al., 2013), birds (White et al., 2010), and insects (Anderson et

al., 2012; Koch & Schmid-Hempel, 2011). For example, we previously reported how the gut microbiomes

of colobus monkeys diverged over the course of nine months as a social group fissioned into two

daughter groups (Goodfellow et al., 2019). Changes in the kin composition of groups did not explain this

pattern, while we were unable to discern whether this divergence was due to the daughter groups

having separate home ranges with different food resources or due to the social network changes that

resulted from the group fission (Goodfellow et al., 2019). Indeed, few studies have had access to the

necessary detailed behavioural, ecological, and genetic data to disentangle whether the divergence in

gut microbiomes with home range separation is due to dietary differences, lower degrees of

relatedness, or lack of shared social environments (Archie & Tung, 2015; Björk, Dasari, Grieneisen, &

Archie, 2019).

Diet is suggested to be one of the most important factors affecting the gut microbiome

(Voreades, Kozil, & Weir, 2014). The gut microbial composition fluctuates within hosts with seasonal or

experimental dietary changes (Davenport et al., 2014; David et al., 2013; Hicks et al., 2018; Leamy et al.,

2014; Mallott, Amato, Garber, & Malhi, 2018; Maurice et al., 2015; Michl et al., 2019; Orkin et al., 2019),

and dietary similarities may explain whether groups have distinct gut microbiomes (Orkin et al., 2019).

However, this pattern could also be due to closer genetic relatedness within than between groups as the

host's genetic makeup can affect microbe colonization (Opstal & Bordenstein, 2015; Spor, Koren, & Ley,

2011), and a number of genomic regions are associated with gut microbial composition in rodents

(Bonder et al., 2016; Leamy et al., 2014). This may explain why closely related individuals have more

similar gut microbiomes than unrelated individuals in some studies of humans and captive rodents

(Faith et al., 2013; Kovacs et al., 2011; Ley et al., 2005). In contrast, genetic differentiation between

populations was a poor predictor of their gut microbiome (Grieneisen et al., 2019), and relatedness did

not have a significant effect on the gut microbiome in some studies of humans (Rothschild et al., 2018),

non-human primates (Moeller et al., 2016) and carnivores (Leclaire et al., 2014). Moeller and colleagues

(2016) suggest that this may be due to an overriding effect of transmission among unrelated social

partners. Indirect social contact via shared environments, such as touching common surfaces, may

facilitate microbiome transmission within households (Lax et al., 2014). Direct social contact such as

grooming or sitting in body contact further increases microbiome transmission between close social

partners within groups of monkeys (*Alouatta pigra*: Amato et al., 2017; *Papio cynocephalus*: Tung et al.,

2015; Grieneisen et al., 2017) and lemurs (Eulemur rubriventer: Raulo et al., 2014). Gut microbiomes are

also more similar among socially connected than disconnected siblings and married couples (Dill-

McFarland et al., 2019). In contrast, social distance between non-group members did not predict gut

microbiome similarity in Verreaux's sifaka (*Propithecus verreauxi*). Taken together, these studies

indicate that social interactions transmit gut microbes within groups, while it is unclear whether social

interactions between groups have a similar effect.

To investigate whether the pattern of increasing between-individual differences in the gut

microbiome (i.e., beta-diversity) with home range separation is best explained by lower dietary overlap,

relatedness, or social connectedness, we focus on the black-and-white colobus monkeys (*Colobus*

vellerosus) at Boabeng-Fiema, Ghana. This is one of several rare species of arboreal leaf-eating monkeys

distributed across the forested regions of the African tropics, and it is closely related to guerezas

(*Colobus guereza*) and western black-and-white colobus (*Colobus polykomos*) (Ting, 2008). To break

down hard-to-digest items in their primarily folivorous diet (Saj & Sicotte, 2007; Teichroeb & Sicotte,

2009), they rely on behavioural traits, physiological traits, and their gut microbiome (Amato et al., 2016;

Lambert, 1998). At Boabeng-Fiema, all colobus groups utilize a highly folivorous diet, but the most

important food species differ between groups (Saj & Sicotte, 2007; Teichroeb & Sicotte, 2009). More

seeds and fruits are available during the dry season, during which they eat up to 43% of these food

items (Teichroeb & Sicotte, 2017). Possibly due to constraints imposed by their highly folivorous diet,

colobus monkeys spend a low percentage of their time engaging in direct social activities such as

grooming (Teichroeb, Saj, Paterson, & Sicotte, 2003). Female colobus spend on average 3% of their time

within 1 meter and 0.1% of their time grooming each female group member (Wikberg, Ting, & Sicotte,

2014b). However, females still form preferred friendships, which are only occassionally based on kinship

and never based on their relatively weakly expressed dominance hierarchies (Wikberg, Teichroeb,

Bădescu, & Sicotte, 2013; Wikberg, Ting, & Sicotte, 2014a; Wikberg et al., 2014b; Wikberg, Ting, &

Sicotte, 2015). Instead, females prefer social partners with similar immigration status (Wikberg et al.,

2014b, 2014a) in this population where all males and half of the females disperse (Sicotte et al., 2017;

Teichroeb, Wikberg, & Sicotte, 2009, 2011; Wikberg, Sicotte, Campos, & Ting, 2012). This flexible female

dispersal pattern results in groups with different female kin composition and some close maternal

female kin residing in different groups (Wikberg et al., 2012). Neighbouring groups encounter each other

in the large zones of home range overlap on an almost daily basis. During these encounters, groups

sometimes chase each other away from food trees while they at other times engage in affiliative or

sexual between-group interactions (Sicotte & MacIntosh, 2004; Teichroeb & Sicotte, 2017).

The frequent between-group interactions coupled with variation in diet and relatedness within

and between groups makes this a good study population to investigate whether the pattern of

increasing female gut microbial beta-diversity with home range separation is best explained by lower

degrees of dietary similarity, relatedness, or social connectedness. We take a cross-sectional approach

using observational and genetic data from eight groups to first test whether the gut microbiome was

structured by social groups. We predicted gut microbiome beta-diversity to be structured by groups and

to increase with home range separation. We then evaluated which factors explained gut microbiome

beta-diversity between females across different groups. We expected gut microbiome beta-diversity to

be negatively correlated with dietary similarity, relatedness, and connectedness in the 1-meter

proximity network. Finally, the significant predictor from the analyses above (social distance) was used

in a subsequent population-level analysis of Operational Taxonomic Unit (OTU) abundance to determine

which microbial taxa were transmitted this way.

Materials and Methods

Behavioral data collection

Demographic data have been collected since 2000 from the black-and-white colobus monkeys (*Colobus*

vellerosus) at Boabeng-Fiema, Ghana. In this study, we also use behavioral and ecological data as well as

DNA samples from eight groups (Fig. A1) collected during two study periods: the rainy season May-

August 2007 and the pre-dry and dry seasons October 2008 - April 2009 (Table A2). During this time

period, the study groups contained 4-9 adult (i.e., parous) females (Table A2), 1-4 adult males, and 8-24

immatures. Data collection was approved by the Boabeng-Fiema Monkey Sanctuary's management

committee, Ghana Wildlife Division, and the University of Calgary's Animal Care Committee (BI 2006-28,

BI 2009-25).

We recoded the group's location every hour using a map with trails, roads, villages, and large

trees (>40 cm DBH) in order to determine home ranges (Fig. A1). During 10-minute focal samples

(Altmann, 1974) of adult females females, we continuously recorded all social behaviors (including the

identity of the interactant and the duration of the behavior) and plant species and part (i.e., mature leaf,

young leaf, flower, fruit, seed, or other) for each ingested food item. Females fed on a total of 210 food

item-plant species combinations, and to assess dietary differences, we calculated Sørensen dissimilarity

indices using ingested plant parts and plant species during focal samples. We choose this diversity index

because it only takes the presence or absence of an ingested food item into account, which we have a

robust estimate of using the focal data. The Sørensen dissimilarity indices in our data set had a high

median value of 0.83 and it was lower within than between groups (Fig. A2).

We observed over 300 between-group encounters (i.e., two groups located within 50 meters of

each other), of which 53% of the encounters lacked female aggression and 35% of the encounters lacked

male aggression. Because close proximity between individuals of different groups are rare and unlikely

to be recorded during focal sampling, we recorded approaches to 1 meter collected *ad libitum* (Altmann,

1974). Some of these approaches only led to brief close proximity while others led to prolonged contact

like copulations, grooming, and play. We created an undirected proximity network based on the

presence and absence of approaches to 1 meter between all individuals (N = 177) present in the eight

study groups. We used the software UCINET (Borgatti, Everett, & Freeman, 2002) to compute the dyadic

Geodesic distance in the 1-meter proximity network (hereafter referred to as social distance), which is

the number of steps (i.e., recorded interaction ties) in the shortest path from one individual to another.

Group members were in 1-meter proximity with each other (i.e., a geodesic distance of 1) or separated

by two to three partners (i.e., a geodesic distance of 2-4) (Fig. A2). Individuals belonging to different

groups had a geodesic distance of 1 to 9 while some were not connected at all and assigned a distance

of 10 for the purpose of our data analysis (Fig. 1; Fig. A2). Note that the seemingly unconnected

individuals only occurred in the 2007 data set, and they were unconnected because we only had access

to data collected from a 3-month period. These individuals are connected and separated with up to

eight steps in the 2008-2009 network, which is based on six months of data.

Genetic data collection

We collected faecal samples June-August 2007 and January-April 2009. Immediately after a female

defecated, we collected approximatey 1g of feces and dissolved it in 6ml RNAlater. The samples were

stored in a fridge at the field site until the end of the field season when they were transported to the

Ting lab and stored in a -20-degree C freezer. Note that we lack information on soil type, which was

driving between-site differences in the gut microbiomes in a large-scale study of terrestrial baboons

(Grieneisen et al., 2019). However, our samples were collected from arboreal primates within a small

study area, and sampling site does not have a significant effect on beta-diversity in our study population

(Goodfellow et al., 2019).

(a) 2007

(a) 2009

(b) 2007

(b) 2009

Focal Group

BS NP SP
DA RT WW

Unweighted Uni−Frac
0.10 0.15 0.20 0.25 0.30

Focal Group

BO OD WW

Unweighted Uni−Frac
0.20 0.25

Figure 1. Social networks for: **a)** the entire population. Each group member is depicted as a node in

yellow or blue depending if they resided in a group that was or was not used for behavioural analyses.

Between-group dyads with a Geodesic distance of one (i.e., observed in 1-meter proximity) are

connected with lines. **b)** females included in the behavioural analyses with lines connecting non-group

members (i.e., nodes of different color) where color represents gut microbiome beta-diversity (i.e.,

unweighted Unifrac distances) ranging from similar (dark) to dissimilar (light) and thickness indicates

social distances ranging from short (thick) to long distances (thin).

We extracted DNA from the samples and genotyped the extracts at 17 short tandem repeat loci

(STR) as previously described (Wikberg et al., 2012). To make sure that the samples used in the

relatedness and gut microbiome analyses were collected from the correct individual, we compared the

short tandem repeat STR genotypes obtained from these samples with a second sample collected from

the same individual at a different time. We calculated dyadic estimated relatedness values (R) in

MLRelate (Kalinowski, Wagner, & Taper, 2006) because this method provided the most accurate

relatedness estimates in our study population (Wikberg et al., 2012). We used R-values calculated from

STR loci rather than theoretical relatedness (r) calculated from pedigrees, because R-values predict

kinship relatively accurately in our study population (Wikberg et al., 2014a) and they are more accurate

than r in studies such as ours with limited access to pedigrees (Forstmeier, Schielzeth, Mueller, Ellegren,

& Kempenaers, 2012; Robinson, Simmons, & Kennington, 2013). The median female relatedness was

low both within and between groups, but there were at least some closely related females residing in

the same groups (Fig. A2).

198   For generating the gut microbial data, we conducted fresh DNA extracts from 61 previously

199 genotyped samples from 45 females (Table A2) using the QIAamp DNA Stool Mini Kit with a modified

200 protocol. More details regarding the extraction protocol are presented in the Appendix and in

201 Goodfellow et al. (2019). The V4 hypervariable region of the bacterial 16S ribosomal RNA gene was

202 amplified and libraries were prepared using the 515F and 806R primers containing 5' Illumina adapter

203 tails and dual indexing barcodes, and libraries were sequenced as part of a 150bp paired-end sequencing

204 run on the Illumina NextSeq platform following Goodfellow et al. (2019). We used a custom pipeline

205 (https://github.com/kstagaman/Process_16S) for quality filtering and assembly (see Appendix). We

206 performed *de novo* OTU picking in UCLUST (Edgar, 2010), and sequences with 97% overlap were defined

207 as belonging to the same bacterial Operational Taxonomic Unit (OTU). We obtained a total of 2,597

208 OTUs. To guard against sequencing errors, we filtered out OTU's with a frequency lower than 0.00005 as

209 recommended (Bokulich et al., 2012). After filtering, the 2007 data set contained 450 OTUs and had a

210 mean read depth of 91,833 (range: 61,469 – 118,633) and the final 2009 data set contained 396 OTUs

211 with a mean read depth of 91,833 (range: 61,469 – 101,081). We did not rarefy the data set to an even

212 read depth, because it is recommended against (McMurdie & Holmes, 2014).

213   We initially calculated four different measures of gut microbiome beta-diversity (Sørensen

214 dissimilarity index, Bray-Curtis dissimilarity index, unweighted UniFrac distances, and weighted UniFrac

215 distances) in the R package vegan (Oksanen et al., 2017). Because the two presence/absence indices

216 were strongly correlated with each other (Sorenson dissimilarity indices and unweighted UniFrac

217 distances: Mantel r = 0.93, p = 0.001) as were the two abundance indices (Bray-Curtis dissimilarity

indices and weighted UniFrac distances: Mantel r = 0.77, p = 0.001), in our analyses, we only used the

one presence/absence index (unweighted UniFrac distances) and the one abundance index (weighted

UniFrac distances) that take phylogenetic relationships of OTUs into account.

Data analyses

We combined the 2007 and 2009 data sets and included study year (aka season) and individual ID as

predictor variables whenever possible (i.e., permutational multivariate analysis of variance and linear

mixed models) while we had to create squared interaction matrices for each study year separately when

using matrix correlations (i.e., Mantel tests and Moran's test for autospatial correlations). We only used

the full data set (N = 61 samples from 45 females) for the initial analysis regarding the effect of group

identitiy. All subsequent analyses examined the effects of behavioural variables on beta-diversity in a

subset of the full dataset (N = 48 samples from 42 females), as we removed duplicate samples from the

same year and female, one adult female with incomplete dietary information, and groups in which the

majority of females were not sampled to make sure we had a representative sample of social distance

from each group (Table A2).

The initial analysis investigated the effects of season, social group, individual identity, and read

depth in the full data set (N = 61 samples from 45 females) using permutational multivariate analysis of

variance (PERMANOVA) with 10,000 permutations using the adonis function in the R package vegan

(Oksanen et al., 2017). The terms were added sequentially in the order listed above.

We used non-parametric Mantel correlations implemented in the R package vegan (Oksanen et

al., 2017) to investigate whether the two measures of gut microbiome beta-diversity were correlated

with home range separation (0 = same group and home range; 1 = different groups but adjacent home

ranges, 2 = different groups and non-adjacent home ranges).

To investigate which combination of dyadic traits predicted gut microbiome beta-diversity

between females, we created linear mixed models (LMMs) with the outcome variable gut microbiome

beta-diversity and the random effects animal identities. We created a null model that did not contain

any fixed effects, alternative models with one fixed effect that represented one of the hypotheses

outlined in the introduction (dietary dissimilarities, R-values, or social distances), and a full model with

all three predictor variables. We included data collection year as a fixed effect in all alternative models

because the two sampling years occurred in different seasons and several other studies show strong

seasonal shifts in gut microbiome composition (Amato et al., 2015; Hicks et al., 2018; Orkin et al., 2019;

Smits et al., 2017; Springer et al., 2017). All numerical predictor variables were centered and scaled

(Schielzeth, 2010). We included group and focal identity as random effects in all models. We did not

have any issues with collinearity based on low Variance Inflation Factors for the full models (all VIF <

1.43). We evaluated the support for each model using Akaike Information Criterion (AIC) (Akaike, 1974),

and this approach allowed us to determine which hypotheses (diet, relatedness, or social distance) was

best supported by our data (Burnham & Anderson, 2002). In the first set of analyses, we included dyads

that resided in the same group and dyads that resided in different groups. To make sure that the effect

of social distance was not driven by the close social bonds within groups, we repeated the analyses with

between-group dyads only. These analyses were done using the packages lme4 (Bates et al., 2014) and

MuMIn (Barton, 2013) in R (R Core Team, 2018).

To infer which of the gut microbial taxa may be transmitted via close proximity, which was a

better predictor of beta-diversity than diet and relatedness (see Results), we investigated whether the

abundance of each OTU was correlated with Geodesic distance in the 1-meter approach network using

Moran's test for autospatial correlations implemented in the package ape (Paradis, Claude, & Strimmer,

2004). We included within-group and between-group dyads in this analysis (N = 342 dyads). We counted

the number of OTUs in each phylum (or family) that were socially structured based on the autospatial

correlation results. We conducted hypergeometric tests to investigate whether this number was higher

than expected by chance based on the total number of OTUs in the phylum (or family) using the phyper

function implemented in R. In all analyses of taxonomic differences, we used the 10% false discovery

rate to correct p-values for multiple testing (sensu Tung et al., 2015). The gut microbial taxa we

expected to be shaped by sociality are listed in the Table A1 (Amato et al., 2017; Goodfellow et al., 2019;

Tung et al., 2015).

Results

Factors predicting gut microbiome beta-diversity

We investigated the relative effects of season, group, individual ID, and read depth in the full data set

(PERMANOVA, N = 61 samples collected 2007-2009). Of the observed variation in the taxonomic

composition of the gut microbiome (i.e., beta-diversity), individual identity explained the largest

percentage (54-55% depending on which beta-diversity index was used as outcome variable), group

identity explained a more moderate percentage (19-28%), while year explained much smaller

percentage (8-12%) (Table 1). Read depth did not have a significant effect on beta-diversity (Table 1).

Beta- diversity index	Factor	Df	Sums of squares	Mean squares	F	R^2	P
Unweighted UniFrac	Season	1	0.103	0.103	12.325	0.084	<0.001
	Group	7	0.347	0.050	5.943	0.282	<0.001
	ID	39	0.663	0.017	2.035	0.538	<0.001
	Read depth	1	0.010	0.010	1.185	0.008	0.249
Weighted UniFrac	Season	1	0.136	0.136	12.277	0.118	<0.001
	Group	7	0.219	0.031	2.812	0.189	<0.001
	ID	39	0.639	0.016	1.477	0.553	<0.001
	Read depth	1	0.018	0.018	1.637	0.016	0.104

Table 1. Results from the PERMANOVA with factors added sequentially in the ordered listed in the table.

Gut microbiome beta-diversity and home range separation were correlated in the 2007 data set

(N = 870 dyads in 6 groups, Mantel tests, unweighted UniFrac distance: r = 0.22, p = 0.002; weighted

UniFrac distance: r = 0.10, p = 0.049) and in the 2009 data set (N = 342 dyads in 3 groups, Mantel tests;

unweighted UniFrac distance: r = 0.36, p = 0.005; weighted UniFrac distance: r = 0.20, p = 0.024),

meaning that females residing farther from each other had less similar gut microbiomes. This pattern

can potentially be explained by group members having more similar diets, higher relatedness, or

stronger social ties than non-group members (Fig. A2).

We created several competing generalized linear mixed models to investigate which of the three

hypotheses best explained increasing beta-diversity with home range separation: dietary dissimilarity,

relatedness, or social distance, controlling for data collection year. In our data set with both within-

group and between-group dyads (N = 1,212 dyads in 2007-2009), the models with social distance

received the greatest support (Table 2). Social distance predicted gut microbiome beta-diversity, and

females located further apart in the social network had less similar gut microbiomes (Figs. 1-3). Year also

predicted gut microbiome beta-diversity (Fig. 3), and females had more similar gut microbiomes during

the rainy season of 2007 than the dry season of 2009. Without taking the random effects into account,

the model with social distance and year explained 9% to 24% of the observed variation depending on

which beta-diversity index was used (Table 2). The alternative models with dietary dissimilarity and

relatedness did not receive strong support based on their high AIC values (Table 2).

Figure 2. Gut microbiome beta-diversity (i.e., unweighted UniFrac distances) for dyads located close

(geodesic distance 1-2), intermediate (geodesic distance 3-4), and far away from each other (geodesic

distance 5 or greater) in the population's 1-meter proximity network.

Outcome variable	Fixed effect	AIC	Delta	Weight	R^2 marginal	R^2 conditional
a) Within-group and between-group dyads						
Unweighted UniFrac	Season + Proximity	-4768.97	0.00	0.84	0.24	0.48
	Season + Diet + Relatedness + Proximity	-4765.69	3.29	0.16		
	Season + Diet	-4714.71	54.26	0.00		
	Season + Relatedness	-4670.38	98.60	0.00		
	-	-4561.86	207.11	0.00		
Weighted UniFrac	Season + Proximity	-4616.00	0.00	1.00	0.09	0.22
	Season + Diet	-4593.90	22.10	0.00		
	Season + Relatedness	-4593.47	22.53	0.00		
	Season + Diet + Relatedness + Proximity	-4588.86	27.13	0.00		
	-	-4572.74	43.26	0.00		
b) Between-group dyads						
Unweighted UniFrac	Season + Proximity	-4230.98	0.00	0.97	0.32	0.56
	Season + Diet	-4223.20	7.79	0.02		
	Season + Relatedness	-4222.57	8.42	0.01		
	Season + Diet + Relatedness + Proximity	-4203.70	27.29	0.00		
	-	-4061.37	169.62	0.00		
Weighted UniFrac	Season + Proximity	-3712.02	0.00	0.94	0.10	0.25
	Season + Diet	-3705.37	6.65	0.03		
	Season + Relatedness	-3704.60	7.42	0.02		
	Season + Diet + Relatedness + Proximity	-3688.41	23.61	0.00		
	-	-3677.10	34.92	0.00		

Table 2. The competing LMMs' fixed effects, Akaike Information Criterion, delta (i.e., difference in AIC

between the current model and the best-fit model), and Akaike weights (i.e., relative likelihood of the

model), and marginal and conditional R^2 for the best fitting model (i.e., without versus with random

effects) when including **a)** within-group and between-group dyads and **b)** only between-group dyads.

To assess whether the effect of social distance on gut microbiome beta-diversity was driven by

closely connected within-group dyads having very similar gut microbiomes, we repeated the analyses

with between-group dyads only (N = 966 dyads). The social distance models were again the strongest

supported models and explained 10% to 32% of the observed variation depending on which beta-

diversity index was used as outcome variable (Table 2). Again, both year and social distance predicted

beta-diversity (Fig. 3). The alternative models with diet or relatedness were not well supported because

of their high AIC values were much higher than those for the proximity models (Table 2).

Figure 3. Coefficient estimates and their 95% confidence intervals for the best fitting model for

unweighted versus weighted UniFrac distances in the data set including within-group and between-

group dyads and in the data set with between-group dyads only. Social distance is a continuous variable.

Of the two study periods, year 2007 was used as the baseline level against which we depict the effect of

year 2009.

Socially structured OTUs

Our data set contained OTUs from 14 phyla, of which the most well-represented was Firmicutes,

followed by Bacteroidetes, Spirochetes, and Verrucomicrobia (Supplementary Material Fig. 1). In each

social group, at least 70% of the OTUs belonged to the phylum Firmicutes (Supplementary Material Fig.

1) and at least 50% of the OTUs belonged to the families Lachnospiraceae and Ruminococcaceae in the

phylum Firmicutes (Supplementary Material Fig. 2).

Social distance predicted differences in abundances for 73 of the 396 OTUs in the 2009 data set

(Moran's I range: -0.27 – -0.14, all $p < 0.05$, Supplementary Material Table 1). The number of OTUs with

a significant relationship to social distance was greater than expected in the phylum Firmicutes (N = 64,

Hypergeometric test, $p < 0.001$). The numbers of socially structured OTUs in the phyla Bacteroidetes

(N=6), Planctomycetes (N = 1), Proteobacteria (N = 1) and Tenericutes (N = 1) were not greater than

expected based on the total number of OTUs in these phyla (Hypergeometric tests, $p > 0.050$). The other

phyla did not contain any socially structured OTUs. Four families had a higher than expected number of

socially structured OTUs: Bacteroidaceae (N = 4), Lachnospiraceae (N = 20), Peptococcaceae 2 (N = 1),

and Ruminococcaceae (N = 31). There was also a greater than expected number of socially structured

OTUs in 14 of 34 genera (Fig. 4).

Figure 4. The proportion of Operational Taxonomic Units (OTUs) whose abundance was correlated with

distance in the proximity network for genera that contained a higher than expected number of socially

structured OTUs.

Social distance predicted differences in abundances for 1 of the 450 OTUs in the 2007 data set

(Moran's I range: -0.27 – -0.14, all p < 0.05), which belonged to the phylum Firmicutes, the family

Lachnospiraceae, and the genus *Roseburia*. As a result, these taxa had a greater than expected number

of socially structured OTUs (Hypergeometric tests, all p > 0.001).

Discussion

The aim of this study was to investigate if the increase in gut microbiome beta-diversity with home

range separation in female colobus monkeys was best explained by diet, relatedness, or sociality.

Distance in the proximity network was a better predictor than diet and relatedness, similar to findings in

more social primates (Amato et al., 2017; Perofsky, Lewis, Abondano, Di Fiore, & Meyers, 2017; Raulo et

al., 2014; Tung et al., 2015). Although these previous studies suggest that strong social bonds within

groups drive between-group differences in the gut microbiome after ruling out the effects of relatedness

and diet, this is the first report of a relationship between gut microbiome beta-diversity and social

connectedness between individuals in different groups. These mostly brief instances of close proximity

during between-group encounters explained up to 32% of the variation between non-group members.

In contrast, gut microbiome dissimilarity between individuals residing in different groups did not

increase with grooming network distance in sifakas (Perofsky et al., 2017). These contrasting results may

be due to the nature of the host population's between-group interactions, particularly the presence or

absence of friendly behaviours. Colobus monkeys sometimes engage in affiliative, sexual, and playful

behaviours with non-group members (Supplemental Information; Sicotte & MacIntosh, 2004; Teichroeb,

Marteinson, & Sicotte, 2005; Teichroeb et al., 2011), which differ from the almost exclusively aggressive

nature of between-group encounters in some other taxa. Similar to these colobus monkeys, mountain

gorillas (*Gorilla beringei beringei*) occasionally affiliate with members from other groups (Forcina et al.,

2019) and human foraging societies form extended social networks to optimize resource flow (Hamilton,

Milne, Walker, Burger, & Brown, 2007). These extended networks could possibly affect their gut

microbiome in similar ways as documented here in colobus monkeys.

To determine the consequences of such social transmission, the first step is to determine which

type of microbes are transmitted this way. The socially transmitted OTUs in this study included all taxa

(family Porphyromonadaceae and genera *Parabacteroides* and *Coprococcus)* that diverged after a group

fission at our site (Goodfellow et al., 2019) and genera (*Bacteroides, Clostridium,* and *Roseburia*) that

were transmitted via grooming and close proximity in howlers (Amato et al., 2017). In contrast, the

socially transmitted OTUs in our study did not overlap with those transmitted via grooming within

groups of baboons (Tung et al., 2015), despite the host species relatively close phylogenetic relationship.

This discrepancy with the baboon study is not particularly surprising given that the colobus lifestyle is

more similar to howlers than baboons, including factors that influence the gut microbiome like diet and

degree of terrestriality (Perofsky, Lewis, & Meyers, 2019). This is despite recent findings that host

phylogeny has a stronger effect than diet on gut microbiome composition (Amato et al., 2019). It is thus

possible that while phylogeny has the strongest overall effect on the gut microbiome, the same gut

microbial taxa are structured by sociality in primates with similar lifestyle.

We found that the majority of socially transmitted OTUs belonged to the most dominant

families in our host population and other folivorous primates (Barelli et al., 2015; Perofsky et al., 2017),

the families Lachnospiraceae and Ruminococcaceae in the phylum Firmicutes. These taxa are well-suited

for breaking down hard-to-digest plant material (Biddle, Stewart, Blanchard, & Leschine, 2013), and it is

therefore possible that socially transmitted gut microbes benefit hosts in terms of improved digestion of

mature leaves, which make up the majority of the colobus diet (Saj & Sicotte, 2007). Several studies

imply that social gut microbe transmission benefits the host. For example, Tung and colleagues (2015)

suggest that the positive health and fitness effects that baboons accrue from forming close social bonds

with group members are mediated by the gut microbiome. Our results support the notion that this may

be the case in a wide range of gregarious species, including those with relatively low frequencies of

social interactions. If social transmission sustains a healthy gut microbiome (as documented in Koch &

Schmid-Hempel, 2011; Vlčková et al., 2018), it could provide an incentive for the formation and

maintenance of social bonds within groups (Lombardo, 2008). Our findings leave open the as-of-yet

unexplored possibility that social transmission of microbes may even explain the occurrence of friendly

between-group encounters, especially in the absence of limiting resouces such as fertile females and

important food sources.

The results of this paper ultimately lead us to an important outstanding question, which is how

gut microbes are transmitted among animals that spend considerably less time grooming or in direct

contact than other primates with socially mediated gut microbe transmission (Amato et al., 2017; Raulo

et al., 2014; Tung et al., 2015). It might be that microbes are transmitted directly during the occasions

we observed non-group members copulating, grooming, and playing. However, it could also be that the

microbes are transmitted indirectly between hosts when they are touching shared surfaces within a

certain time period (Münger, Montiel-Castro, Langhans, & Pacheco-López, 2018). This reasoning is

consistent with spatial proximity predicting the gut microbiome in other gregarious species with low

frequencies of social behaviours like the Welsh Mountain ponies (*Equus ferus caballus*) (Antwis, Lea,

Unwin, & Shultz, 2018) and in more solitary species such as North American red squirrels (*Tamiasciurus*

hudsonicus) (Ren et al., 2017) and gopher tortoise (*Gopherus polyphemus*) (M. L. Yuan et al., 2015). The

occurrences of direct and indirect social transmission are difficult to tease apart when the two are

correlated and when brief physical contact between extra-group members often go unnoticed, but

carefully designed studies in the future may be able to address this question.

418   Finally, relatedness and dietary differences within a season were not good predictors of beta-

diversity in comparison to social distance. In contrast, seasonal changes in diet may be associated with

changes in the colobus gut microbiome, because beta-diversity was higher during the 2009 dry season

when their diet was more diverse than during the 2007 rainy season when they ate mostly mature

leaves. We will continue to investigate whether this seasonal dietary switch is linked to changes in the

gut microbiome, as previously reported from other species inhabiting seasonal environments (Amato et

al., 2015; Hicks et al., 2018; Orkin et al., 2019; Smits et al., 2017; Springer et al., 2017). These authors

concluded that gut microbiome dynamics determine nutrient uptake and is key for dietary flexibility

(Amato et al., 2015; Hicks et al., 2018; Orkin et al., 2019; Smits et al., 2017; Springer et al., 2017), while

the potential three-way interaction between social, dietary, and gut microbial dynamics is still poorly

understood. An interesting venue for further research is therefore to investigate whether the gut

microbiomes of socially well-connected individuals map more quickly onto ecological changes, which

could help them deal with the rapidly changing environments that many wild animals inhabit today.

Acknowledgements

We would like to thank Keaton Stagaman for use of his custom pipeline; Fernando Campos, Teresa

Holmes, and Robert Koranteng for field data collection; BFMS management committee, Ghana Wildlife

Division, and the University of Calgary's Life and Environmental Sciences Animal Care Committee for

permission to conduct this study; Alberta Innovates Technology Futures, American Society of

Primatologists, Animal Behaviour Society, International Primatological Society, Leakey Foundation,

Natural Sciences and Engineering Research Council of Canada, Sweden-America Foundation, Research

Services at the University of Calgary, the National Institute of General Medical Sciences (P50GM098911)

via the META Center for Systems Biology, and Wenner-Gren Foundation for funding.

References

Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic*

Control, *19*(6), 716–723. doi: 10.1109/TAC.1974.1100705

Altmann, J. (1974). Observational study of behavior: sampling methods. *Behaviour*, *49*(3), 227–267.

Amato, K. R. (2016). Incorporating the gut microbiota into models of human and non-human primate

ecology and evolution: Gut microbiota influences host nutrition, health, and behavior. *American*

Journal of Physical Anthropology, *159*, 196–215. doi: 10.1002/ajpa.22908

Amato, K. R., G Sanders, J., Song, S. J., Nute, M., Metcalf, J. L., Thompson, L. R., … R Leigh, S. (2019).

Evolutionary trends in host physiology outweigh dietary niche in structuring primate gut

microbiomes. *The ISME Journal*, *13*(3), 576–587. doi: 10.1038/s41396-018-0175-0

Amato, K. R., Leigh, S. R., Kent, A., Mackie, R. I., Yeoman, C. J., Stumpf, R. M., … Garber, P. A. (2015). The

gut microbiota appears to compensate for seasonal diet variation in the wild black howler

monkey (*Alouatta pigra*). *Microbial Ecology*, *69*(2), 434–443. doi: 10.1007/s00248-014-0554-7

Amato, K. R., Metcalf, J. L., Song, S. J., Hale, V. L., Clayton, J., Ackermann, G., … Braun, J. (2016). Using

the gut microbiota as a novel tool for examining colobine primate GI health. *Global Ecology and*

Conservation, *7*, 225–237. doi: 10.1016/j.gecco.2016.06.004

Amato, K. R., Van Belle, S., Di Fiore, A., Estrada, A., Stumpf, R., White, B., … Leigh, S. R. (2017). Patterns

in gut microbiota similarity associated with degree of sociality among sex classes of a

Neotropical primate. *Microbial Ecology*. doi: 10.1007/s00248-017-0938-6

Anderson, K. E., Russell, J. A., Moreau, C. S., Kautz, S., Sullam, K. E., Hu, Y., … Wheeler, D. E. (2012).

Highly similar microbial communities are shared among related and trophically similar ant

species. *Molecular Ecology*, *21*(9), 2282–2296. doi: 10.1111/j.1365-294X.2011.05464.x

Antwis, R. E., Lea, J. M. D., Unwin, B., & Shultz, S. (2018). Gut microbiome composition is associated with

spatial structuring and social interactions in semi-feral Welsh Mountain ponies. *Microbiome*,

6(1), 207. doi: 10.1186/s40168-018-0593-2

Archie, E. A., & Tung, J. (2015). Social behavior and the microbiome. *Current Opinion in Behavioral*

Sciences, *6*, 28–34. doi: 10.1016/j.cobeha.2015.07.008

Barelli, C., Albanese, D., Donati, C., Pindo, M., Dallago, C., Rovero, F., … Filippo, C. D. (2015). Habitat

fragmentation is associated to gut microbiota diversity of an endangered primate: implications

for conservation. *Scientific Reports*, *5*, 14862. doi: 10.1038/srep14862

Barton, K. (2013). *MuMIn: Multi-model inference. R package version 1.9.5. http://CRAN.R-*

project.org/package=MuMIn. Accessed 1 May 2013.

Bates, D., Maechler, M., Bolker, B., Walker, S., Bojesen Christensen, R. H., & Singman, H. (2014). *lme4:*

Linear mixed-effects models using Eigen and S4. http://lme4.r-forge.r-project.org/ Accessed 1

Jun 2014.

Biddle, A., Stewart, L., Blanchard, J., & Leschine, S. (2013). Untangling the genetic basis of fibrolytic

specialization by Lachnospiraceae and Ruminococcaceae in diverse gut communities. *Diversity*,

5(3), 627–640. doi: 10.3390/d5030627

Björk, J. R., Dasari, M., Grieneisen, L., & Archie, E. A. (2019). Primate microbiomes over time:

Longitudinal answers to standing questions in microbiome research. *American Journal of*

Primatology, *0*(0), e22970. doi: 10.1002/ajp.22970

Bokulich, N. A., Subramanian, S., Faith, J. J., Gevers, D., Gordon, J. I., Knight, R., … Caporaso, J. G. (2012).

Quality-filtering vastly improves diversity estimates from Illumina amplicon sequencing. *Nature*

Methods, *10*(1), 57–59. doi: 10.1038/nmeth.2276

Bonder, M. J., Kurilshikov, A., Tigchelaar, E. F., Mujagic, Z., Imhann, F., Vila, A. V., … Zhernakova, A.

(2016). The effect of host genetics on the gut microbiome. *Nature Genetics*, *48*(11), 1407–1412.

doi: 10.1038/ng.3663

Borgatti, S., Everett, M., & Freeman, L. (2002). *Ucinet for Windows: software for social network analysis*.

Harvard, MA: Analytic Technologies.

Burnham, K., & Anderson, D. (2002). *Model selection and multimodel inference: a practical information-*

theoretic approach. New York: Springer.

Davenport, E. R., Mizrahi-Man, O., Michelini, K., Barreiro, L. B., Ober, C., & Gilad, Y. (2014). Seasonal

Variation in Human Gut Microbiome Composition. *PLoS ONE*, *9*(3), e90731. doi:

10.1371/journal.pone.0090731

David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe, B. E., … Turnbaugh, P.

J. (2013). Diet rapidly and reproducibly alters the human gut microbiome. *Nature*, *505*(7484),

559–563. doi: 10.1038/nature12820

Degnan, P. H., Pusey, A. E., Lonsdorf, E. V., Goodall, J., Wroblewski, E. E., Wilson, M. L., … Ochman, H.

(2012). Factors associated with the diversification of the gut microbial communities within

chimpanzees from Gombe National Park. *Proceedings of the National Academy of Sciences*,

109(32), 13034–13039. doi: 10.1073/pnas.1110994109

Dill-McFarland, K. A., Tang, Z.-Z., Kemis, J. H., Kerby, R. L., Chen, G., Palloni, A., … Herd, P. (2019). Close

social relationships correlate with human gut microbiota composition. *Scientific Reports*, *9*(1),

703. doi: 10.1038/s41598-018-37298-9

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*, *26*(19),

2460–2461. doi: 10.1093/bioinformatics/btq461

Faith, J. J., Guruge, J. L., Charbonneau, M., Subramanian, S., Seedorf, H., Goodman, A. L., … Gordon, J. I.

(2013). The Long-Term Stability of the Human Gut Microbiota. *Science*, *341*(6141), 1237439–

1237439. doi: 10.1126/science.1237439

Forcina, G., Vallet, D., Le Gouar, P. J., Bernardo-Madrid, R., Illera, G., Molina-Vacas, G., … Vila, C. (2019).

From groups to communities in western lowland gorillas. *Proceedings of the Royal Society B-*

Biological Sciences, *286*(1896), 20182019. doi: 10.1098/rspb.2018.2019

Forstmeier, W., Schielzeth, H., Mueller, J. C., Ellegren, H., & Kempenaers, B. (2012). Heterozygosity-

fitness correlations in zebra finches: microsatellite markers can be better than their reputation.

Molecular Ecology, *21*(13), 3237–3249. doi: 10.1111/j.1365-294X.2012.05593.x

Goodfellow, C. K., Whitney, T., Christie, D. M., Sicotte, P., Wikberg, E. C., & Ting, N. (2019). Divergence in

gut microbial communities mirrors a social group fission event in a black-and-white colobus

monkey (Colobus vellerosus). *American Journal of Primatology*, *0*(0), e22966. doi:

10.1002/ajp.22966

Grieneisen, L. E., Charpentier, M. J. E., Alberts, S. C., Blekhman, R., Bradburd, G., Tung, J., & Archie, E. A.

(2019). Genes, geology and germs: gut microbiota across a primate hybrid zone are explained by

site soil properties, not host species. *Proceedings of the Royal Society B: Biological Sciences*,

286(1901), 20190431. doi: 10.1098/rspb.2019.0431

Hamilton, M. J., Milne, B. T., Walker, R. S., Burger, O., & Brown, J. H. (2007). The complex structure of

hunter-gatherer social networks. *Proceedings of the Royal Society B: Biological Sciences*,

274(1622), 2195–2203. doi: 10.1098/rspb.2007.0564

Hannon Lab. (2010). *FASTX Toolkit. Available from http:// hannonlab.cshl.edu/fastx_toolkit/index.html.*

Hansen, M. E. B., Rubel, M. A., Bailey, A. G., Ranciaro, A., Thompson, S. R., Campbell, M. C., … Tishkoff, S.

A. (2019). Population structure of human gut bacteria in a diverse cohort from rural Tanzania

and Botswana. *Genome Biology*, *20*(1), 16. doi: 10.1186/s13059-018-1616-9

Hicks, A. L., Lee, K. J., Couto-Rodriguez, M., Patel, J., Sinha, R., Guo, C., … Williams, B. L. (2018). Gut

microbiomes of wild great apes fluctuate seasonally in response to diet. *Nature*

Communications, *9*(1), 1786. doi: 10.1038/s41467-018-04204-w

Hird, S. M., Carstens, B. C., Cardiff, S. W., Dittmann, D. L., & Brumfield, R. T. (2014). Sampling locality is

more detectable than taxonomy or ecology in the gut microbiota of the brood-parasitic Brown-

headed Cowbird (*Molothrus ater*). *PeerJ*, *2*, e321. doi: 10.7717/peerj.321

Kalinowski, S. T., Wagner, A. P., & Taper, M. L. (2006). ML-RELATE: a computer program for maximum

likelihood estimation of relatedness and relationship. *Molecular Ecology Notes*, *6*(2), 576–579.

doi: 10.1111/j.1471-8286.2006.01256.x

Koch, H., & Schmid-Hempel, P. (2011). Socially transmitted gut microbiota protect bumble bees against

an intestinal parasite. *Proceedings of the National Academy of Sciences*, *108*(48), 19288–19292.

doi: 10.1073/pnas.1110474108

Kovacs, A., Ben-Jacob, N., Tayem, H., Halperin, E., Iraqi, F. A., & Gophna, U. (2011). Genotype Is a

Stronger Determinant than Sex of the Mouse Gut Microbiota. *Microbial Ecology*, *61*(2), 423–428.

doi: 10.1007/s00248-010-9787-2

Lambert, J. E. (1998). Primate digestion: Interactions among anatomy, physiology, and feeding ecology.

Evolutionary Anthropology: Issues, News, and Reviews, *7*(1), 8–20. doi: 10.1002/(SICI)1520-

6505(1998)7:1<8::AID-EVAN3>3.0.CO;2-C

Lankau, E. W., Hong, P.-Y., & Mackie, R. I. (2012). Ecological drift and local exposures drive enteric

bacterial community differences within species of Galápagos iguanas: GALÁPAGOS IGUANA

ENTERIC COMMUNITY DRIVERS. *Molecular Ecology*, *21*(7), 1779–1788. doi: 10.1111/j.1365-

294X.2012.05502.x

Lax, S., Smith, D. P., Hampton-Marcell, J., Owens, S. M., Handley, K. M., Scott, N. M., … Gilbert, J. A.

(2014). Longitudinal analysis of microbial interaction between humans and the indoor

environment. *Science*, *345*(6200), 1048–1052. doi: 10.1126/science.1254529

Leamy, L. J., Kelly, S. A., Nietfeldt, J., Legge, R. M., Ma, F., Hua, K., … Pomp, D. (2014). Host genetics and

diet, but not immunoglobulin A expression, converge to shape compositional features of the gut

microbiome in an advanced intercross population of mice. *Genome Biology*, *15*(12). doi:

10.1186/s13059-014-0552-6

Leclaire, S., Nielsen, J. F., & Drea, C. M. (2014). Bacterial communities in meerkat anal scent secretions

vary with host sex, age, and group membership. *Behavioral Ecology*, *25*(4), 996–1004. doi:

10.1093/beheco/aru074

Ley, R. E., Backhed, F., Turnbaugh, P., Lozupone, C. A., Knight, R. D., & Gordon, J. I. (2005). Obesity alters

gut microbial ecology. *Proceedings of the National Academy of Sciences*, *102*(31), 11070–11075.

doi: 10.1073/pnas.0504978102

Lombardo, M. P. (2008). Access to mutualistic endosymbiotic microbes: an underappreciated benefit of

group living. *Behavioral Ecology and Sociobiology*, *62*(4), 479–497. doi: 10.1007/s00265-007-

0428-9

Magoc, T., & Salzberg, S. L. (2011). FLASH: fast length adjustment of short reads to improve genome

assemblies. *Bioinformatics*, *27*(21), 2957–2963. doi: 10.1093/bioinformatics/btr507

Mallott, E. K., Amato, K. R., Garber, P. A., & Malhi, R. S. (2018). Influence of fruit and invertebrate

consumption on the gut microbiota of wild white-faced capuchins (*Cebus capucinus*). *American*

Journal of Physical Anthropology, *165*(3), 576–588. doi: 10.1002/ajpa.23395

Maurice, C. F., Knowles, S. C. L., Ladau, J., Pollard, K. S., Fenton, A., Pedersen, A. B., & Turnbaugh, P. J.

(2015). Marked seasonal variation in the wild mouse gut microbiota. *The ISME Journal*, *9*, 2423–

2434. doi: 10.1038/ismej.2015.53

McFall-Ngai, M., Hadfield, M. G., Bosch, T. C. G., Carey, H. V., Domazet-Lošo, T., Douglas, A. E., …

Wernegreen, J. J. (2013). Animals in a bacterial world, a new imperative for the life sciences.

Proceedings of the National Academy of Sciences, *110*(9), 3229–3236. doi:

10.1073/pnas.1218525110

McMurdie, P. J., & Holmes, S. (2014). Waste Not, Want Not: Why Rarefying Microbiome Data Is

Inadmissible. *PLoS Computational Biology*, *10*(4), e1003531. doi: 10.1371/journal.pcbi.1003531

Michl, S. C., Beyer, M., Ratten, J.-M., Hasler, M., LaRoche, J., & Schulz, C. (2019). A diet-change

modulates the previously established bacterial gut community in juvenile brown trout (Salmo

trutta). *Scientific Reports*, *9*(1), 2339. doi: 10.1038/s41598-019-38800-7

Moeller, A. H., Foerster, S., Wilson, M. L., Pusey, A. E., Hahn, B. H., & Ochman, H. (2016). Social behavior

shapes the chimpanzee pan-microbiome. *Science Advances*, *2*(1), e1500997–e1500997. doi:

10.1126/sciadv.1500997

Münger, E., Montiel-Castro, A. J., Langhans, W., & Pacheco-López, G. (2018). Reciprocal interactions

between gut microbiota and host social behavior. *Frontiers in Integrative Neuroscience*, *12*, 21.

doi: 10.3389/fnint.2018.00021

Oksanen, J., Guillaume Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., … Szoecs, E. (2017).

vegan: Community Ecology Package (http://CRAN.R-project.org/ package=vegan accessed 1

December 2017).

Opstal, E. J. v., & Bordenstein, S. R. (2015). Rethinking heritability of the microbiome. *Science*, *349*(6253),

1172–1173. doi: 10.1126/science.aab3958

Orkin, J. D., Campos, F. A., Myers, M. S., Cheves Hernandez, S. E., Guadamuz, A., & Melin, A. D. (2019).

Seasonality of the gut microbiota of free-ranging white-faced capuchins in a tropical dry forest.

The ISME Journal, *13*(1), 183–196. doi: 10.1038/s41396-018-0256-0

Orkin, J. D., Webb, S. E., & Melin, A. D. (2019). Small to modest impact of social group on the gut

microbiome of wild Costa Rican capuchins in a seasonal forest. *American Journal of Primatology*,

0(0), e22985. doi: 10.1002/ajp.22985

Paradis, E., Claude, J., & Strimmer, K. (2004). APE: analyses of phylogenetics and evolution in R language.

Bioinformatics, *20*, 289–290.

Perofsky, A. C., Lewis, R. J., Abondano, L. A., Di Fiore, A., & Meyers, L. A. (2017). Hierarchical social

607   networks shape gut microbial composition in wild Verreaux's sifaka. *Proceedings of the Royal*

608   *Society B: Biological Sciences*, *284*(1868), 20172274. doi: 10.1098/rspb.2017.2274

609   Perofsky, A. C., Lewis, R. J., & Meyers, L. A. (2019). Terrestriality and bacterial transfer: a comparative

610   study of gut microbiomes in sympatric Malagasy mammals. *The ISME Journal*, *13*(1), 50–63. doi:

611   10.1038/s41396-018-0251-5

612   Phillips, C. D., Phelan, G., Dowd, S. E., McDONOUGH, M. M., Ferguson, A. W., Delton Hanson, J., … Baker,

613   R. J. (2012). Microbiome analysis among bats describes influences of host phylogeny, life history,

614   physiology and geography: MICROBIOME ANALYSIS AMONG BATS. *Molecular Ecology*, *21*(11),

615   2617–2627. doi: 10.1111/j.1365-294X.2012.05568.x

616   Pollock, J., Glendinning, L., Wisedchanwet, T., & Watson, M. (2018). The Madness of Microbiome:

617   Attempting To Find Consensus "Best Practice" for 16S Microbiome Studies. *Applied and*

618   *Environmental Microbiology*, *84*(7). doi: 10.1128/AEM.02627-17

619   R Core Team. (2018). *R: A language and environment for statistical computing (R Foundation for*

620   *Statistical Computing. http://www.R-project.org. Accessed 1 November 2018).*

621   Raulo, A., Ruokolainen, L., Lane, A., Amato, K., Knight, R., Leigh, S., … Tecot, S. R. (2014). Social behaviour

622   and gut microbiota in red-bellied lemurs (*Eulemur rubriventer*): In search of the role of immunity

623   in the evolution of sociality. *Journal of Animal Ecology*. doi: 10.1111/1365-2656.12781

624   Ren, T., Boutin, S., Humphries, M. M., Dantzer, B., Gorrell, J. C., Coltman, D. W., … Wu, M. (2017).

625   Seasonal, spatial, and maternal effects on gut microbiome in wild red squirrels. *Microbiome*,

626   *5*(1), 163. doi: 10.1186/s40168-017-0382-3

627   Robinson, S. P., Simmons, L. W., & Kennington, W. J. (2013). Estimating relatedness and inbreeding using

628   molecular markers and pedigrees: the effect of demographic history. *Molecular Ecology*, *22*(23),

629   5779–5792. doi: 10.1111/mec.12529

630   Rothschild, D., Weissbrod, O., Barkan, E., Kurilshikov, A., Korem, T., Zeevi, D., … Segal, E. (2018).

631   Environment dominates over host genetics in shaping human gut microbiota. *Nature*, *555*(7695),

632   210–215. doi: 10.1038/nature25973

633   Saj, T. L., & Sicotte, P. (2007). Predicting the Competitive Regime of Female *Colobus vellerosus* from the

634   Distribution of Food Resources. *International Journal of Primatology*, *28*(2), 315–336. doi:

635   10.1007/s10764-007-9124-x

636   Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients. *Methods

637   in Ecology and Evolution*, *1*(2), 103–113. doi: 10.1111/j.2041-210X.2010.00012.x

638   Sicotte, P., & MacIntosh, A. J. (2004). Inter-group encounters and male incursions in *Colobus vellerosus*

639   in central Ghana. *Behaviour*, *141*, 533–553.

640   Sicotte, P., Teichroeb, J. A., Vayro, J. V., Fox, S. A., Bădescu, I., & Wikberg, E. C. (2017). The influence of

641   male takeovers on female dispersal in *Colobus vellerosus*: Takeovers and Female Dispersal in

642   Colobus. *American Journal of Primatology*, *79*(7), e22436. doi: 10.1002/ajp.22436

Smits, S. A., Leach, J., Sonnenburg, E. D., Gonzalez, C. G., Lichtman, J. S., Reid, G., … Sonnenburg, J. L.

(2017). Seasonal cycling in the gut microbiome of the Hadza hunter-gatherers of Tanzania.

Science, *357*(6353), 802–806. doi: 10.1126/science.aan4834

Song, S. J., Lauber, C., Costello, E. K., Lozupone, C. A., Humphrey, G., Berg-Lyons, D., … Knight, R. (2013).

Cohabiting family members share microbiota with one another and with their dogs. *ELife*, *2*. doi:

10.7554/eLife.00458

Spor, A., Koren, O., & Ley, R. (2011). Unravelling the effects of the environment and host genotype on

the gut microbiome. *Nature Reviews Microbiology*, *9*(4), 279–290. doi: 10.1038/nrmicro2540

Springer, A., Fichtel, C., Al-Ghalith, G. A., Koch, F., Amato, K. R., Clayton, J. B., … Kappeler, P. M. (2017).

Patterns of seasonality and group membership characterize the gut microbiota in a longitudinal

study of wild Verreaux's sifakas (*Propithecus verreauxi*). *Ecology and Evolution*, *7*(15), 5732–

5745. doi: 10.1002/ece3.3148

Teichroeb, J. A., Marteinson, S., & Sicotte, P. (2005). Individuals' behaviors following dye-marking in wild

black-and-white colobus (*Colobus vellerosus*). *American Journal of Primatology*, *65*(2), 197–203.

doi: 10.1002/ajp.20108

Teichroeb, J. A., Saj, T. L., Paterson, J. D., & Sicotte, P. (2003). Effect of group size on activity budgets of

Colobus vellerosus in Ghana. *International Journal of Primatology*, *24*(4), 743–758. doi:

10.1023/A:1024672604524

Teichroeb, J. A., & Sicotte, P. (2009). Test of the ecological-constraints model on ursine colobus monkeys

(*Colobus vellerosus*) in Ghana. *American Journal of Primatology*, *71*(1), 49–59. doi:

10.1002/ajp.20617

Teichroeb, J. A., & Sicotte, P. (2017). Cascading competition: the seasonal strength of scramble

influences between-group contest in a folivorous primate. *Behavioral Ecology and Sociobiology*,

72(1), 6. doi: 10.1007/s00265-017-2418-x

Teichroeb, J. A., Wikberg, E. C., & Sicotte, P. (2009). Female dispersal patterns in six groups of ursine

colobus (*Colobus vellerosus*): infanticide avoidance is important. *Behaviour*, *146*(4), 551–582.

doi: 10.1163/156853909X426363

Teichroeb, J. A., Wikberg, E. C., & Sicotte, P. (2011). Dispersal in male ursine colobus monkeys (*Colobus*

vellerosus): influence of age, rank and contact with other groups on dispersal decisions.

Behaviour, *148*(7), 765–793. doi: 10.1163/000579511X577157

Theis, K. R., Venkataraman, A., Dycus, J. A., Koonter, K. D., Schmitt-Matzen, E. N., Wagner, A. P., …

Schmidt, T. M. (2013). Symbiotic bacteria appear to mediate hyena social odors. *Proceedings of*

the National Academy of Sciences, *110*(49), 19832–19837. doi: 10.1073/pnas.1306477110

Ting, N. (2008). Mitochondrial relationships and divergence dates of the African colobines: evidence of

Miocene origins for the living colobus monkeys. *Journal of Human Evolution*, *55*(2), 312–325.

doi: 10.1016/j.jhevol.2008.02.011

Tung, J., Barreiro, L. B., Burns, M. B., Grenier, J.-C., Lynch, J., Grieneisen, L. E., … Archie, E. A. (2015).

Social networks predict gut microbiome composition in wild baboons. *ELife*, *4*, e05224. doi:

10.7554/eLife.05224

Turnbaugh, P. J., Ley, R. E., Mahowald, M. A., Magrini, V., Mardis, E. R., & Gordon, J. I. (2006). An

obesity-associated gut microbiome with increased capacity for energy harvest. *Nature*,

444(7122), 1027–1131. doi: 10.1038/nature05414

Vlčková, K., Pafčo, B., Petrželková, K. J., Modrý, D., Todd, A., Yeoman, C. J., … Gomez, A. (2018).

Relationships between gastrointestinal parasite infections and the fecal microbiome in free-

ranging western lowland gorillas. *Frontiers in Microbiology*, *9*(JUN), 1202. doi:

10.3389/fmicb.2018.01202

Voreades, N., Kozil, A., & Weir, T. L. (2014). Diet and the development of the human intestinal

microbiome. *Frontiers in Microbiology*, *5*. doi: 10.3389/fmicb.2014.00494

White, J., Mirleau, P., Danchin, E., Mulard, H., Hatch, S. A., Heeb, P., & Wagner, R. H. (2010). Sexually

transmitted bacteria affect female cloacal assemblages in a wild bird: Sexually transmitted

bacteria in kittiwakes. *Ecology Letters*, *13*(12), 1515–1524. doi: 10.1111/j.1461-

0248.2010.01542.x

Wikberg, E. C., Sicotte, P., Campos, F. A., & Ting, N. (2012). Between-group variation in female dispersal,

kin composition of groups, and proximity patterns in a black-and-white colobus monkey

(*Colobus vellerosus*). *PLoS ONE*, *7*(11), e48740. doi: 10.1371/journal.pone.0048740

Wikberg, E. C., Teichroeb, J. A., Bădescu, I., & Sicotte, P. (2013). Individualistic female dominance

hierarchies with varying strength in a highly folivorous population of black-and-white colobus.

Behaviour, 1–26. doi: 10.1163/1568539X-00003050

Wikberg, E. C., Ting, N., & Sicotte, P. (2014a). Familiarity is more important than phenotypic similarity in

shaping social relationships in a facultative female dispersed primate, *Colobus vellerosus*.

Behavioural Processes, *106*, 27–35. doi: 10.1016/j.beproc.2014.04.002

Wikberg, E. C., Ting, N., & Sicotte, P. (2014b). Kinship and similarity in residency status structure female

social networks in black-and-white colobus monkeys (*Colobus vellerosus*). *American Journal of

Physical Anthropology*, *153*(3), 365–376. doi: 10.1002/ajpa.22435

Wikberg, E. C., Ting, N., & Sicotte, P. (2015). Demographic factors are associated with intergroup

variation in the grooming networks of female colobus (*Colobus vellerosus*). *International Journal

of Primatology*, *36*(1), 124–142. doi: 10.1007/s10764-015-9816-6

Yatsunenko, T., Rey, F. E., Manary, M. J., Trehan, I., Dominguez-Bello, M. G., Contreras, M., … Gordon, J.

I. (2012). Human gut microbiome viewed across age and geography. *Nature*. doi:

10.1038/nature11053

Yuan, M. L., Dean, S. H., Longo, A. V., Rothermel, B. B., Tuberville, T. D., & Zamudio, K. R. (2015). Kinship,

inbreeding and fine-scale spatial structure influence gut microbiota in a hindgut-fermenting

tortoise. *Molecular Ecology*, *24*(10), 2521–2536. doi: 10.1111/mec.13169

Yuan, S., Cohen, D. B., Ravel, J., Abdo, Z., & Forney, L. J. (2012). Evaluation of Methods for the Extraction

and Purification of DNA from the Human Microbiome. *PLOS ONE*, *7*(3), e33865. doi:

10.1371/journal.pone.0033865

Data Accessibility Statement

All raw data are stored in the PaceLab database hosted by the University of Calgary. The 16S sequencing

data will be uploaded to NCBI's Short Read Archive. The data used for the analyses presented here will

be uploaded to Dryad.

Author contribution

E.C.W., P.S., and N.T. designed and funded the study. E.C.W. and D.C. conducted the DNA extractions

and analysed the data. All authors contributed to writing the manuscript.

Appendix

Microbial taxa predicted to be structured by social distance

The last aim of this study was to investigate whether social distance was correlated with Operational

Taxonomic Unit (OTU) abundances in certain taxa (Table A1) previously reported as structured by social

relationships (Amato et al., 2017; Tung et al., 2015). We also expected social distance to be correlated

with the abundances of three gut microbial taxa that diverged between the two daughter groups after a

group fission (DA and NP in Fig. A1), because we suspect that this pattern was driven by social network

changes (Goodfellow et al., 2019).

737
738   **Figure A1.** Home ranges and core areas for groups in the main forest fragment at Boabeng-Fiema,

739   Ghana. Solid lines indicate home ranges of groups from which we collected behavioural data. Dashed

740   lines indicate partial home ranges from other groups present in this fores

Taxon	Socially structured in 2007	Socially structured in 2009	Reference
Actinobacteria:	X	X	Tung et al. 2015
Bifidobacteriaceae:	NA	X	Tung et al. 2015
Bifidobacterium	NA	X	Tung et al. 2015
Coriobacteriaceae	X	X	Tung et al. 2015
Bacteroidetes:	-	-	-
Bacteroidaceae:	X	(✓)	-
Bacteroides	X	✓	Amato et al. 2017
Porphyromonadaceae:	NA	X	Goodfellow et al. 2019
Parabacteroides	NA	✓	Goodfellow et al. 2019
Firmicutes:	✓	(✓)	-
Clostridiaceae	-	-	-
Clostridium	X	✓	Amato et al. 2017
Eubacteriaceae	-	-	-
Eubacterium	X	(✓)	-
Lachnospiraceae:	✓	(✓)	-
Coprococcus	X	✓	Goodfellow et al. 2019
Lachnospiracea incertae sedis	X	(✓)	-
Roseburia	✓	X	Amato et al. 2017
Peptococcaceae:	X	(✓)	-
Desulfurispora	X	(✓)	-
Ruminococcaceae	X	(✓)	-
Flavonifractor	X	(✓)	-
Subdoligranulum	NA	(✓)	-
Streptococcaceae:	NA	-	-
Streptococcus	NA	NA	Amato et al. 2017
Veillonellaceae:	X	X	Tung et al. 2015
Propionispira	X	(✓)	-
Centipeda	X	(✓)	-
Fuscobacteria	NA	X	Tung et al. 2015
Fusobacteriaceae	NA	X	Tung et al. 2015
Fusobacterium	NA	X	Tung et al. 2015
Proteobacteria:	-	-	-
Desulfovibrionaceae	-	-	-
Desulfovibrio	X	(✓)	-
Enterobacteriaceae	X	X	Tung et al. 2015
Tenericutes:	X	X	Tung et al. 2015
Anaeroplasmataceae	-	-	-
Asteroleplasma	X	(✓)	-
Mycoplasmataceae	NA	X	Tung et al. 2015
Mycoplasma	NA	X	Tung et al. 2015

Table A1. OTUs in these phyla, families, and genera are expected to be structured by sociality based on

previous studies. Predictions were supported ✓; not supported X; no prediction made but structured in

our data set (✓); or no prediction made and not structured in our data set (-). Grey text indicates rare

taxa (N < 3 OTUs). NA denotes taxa not present in our data set.

DNA extraction, amplification, and sequencing protocols for the gut microbiome analysis

We extracted DNA from 200 μl of sample using QIAmp DNA stool extraction protocol the following

modifications. Step 2: Added 50 μl Proteinase K. Step 4: Pipetted all of the supernatant. Step 5: Used

half of the InhibitEX tablet. Step 6: Centrifuged for 5 minutes. Step 9: Added 4ul RNAse and vortexed for

15 seconds. Step 19: Used 50 μl Buffer AE and incubated at 10 minutes. Step 20: Pipetted the same 50 μl

of buffer AE back onto filter and incubated at room temperature for 15 minutes. Centrifuged at full

speed for 2 minutes. Our DNA extraction protocol did not include a bead-beating step, which could bias

against lysis-resistant taxa such as Gram-positive and spore-forming bacteria that are less likely to be

dependent on direct social contact for transmission between hosts because they can survive for

prolonged periods outside the host (Pollock, Glendinning, Wisedchanwet, & Watson, 2018; Yuan, Cohen,

Ravel, Abdo, & Forney, 2012).

We determined the concentration of the extracts using Qubit dsDNA BR Assay Kit (Invitrogen)

and diluted products to 2nM for downstream reactions. We amplified the bacterial v4 region of the 16S

ribosomal RNA gene using the following 515F and 806R primers containing 5' Illumina adapter tails and

dual indexing barcodes:

515F 5' AATGATACGGCGACCACCGAGATCTACACTAGATCGCTATGGTAATTGTGTGCCAGCMGCCGCGGTAA

806R 5' CAAGCAGAAGACGGCATACGAGATTCACCTAGAGTCAGTCAGCCGGACTACHVGGGTWTCTAAT.

We set the PCRs with 12.5 μl NEB Q5 Hot start 2x Master mix, 1.25 μl 10uM Primer mix, 1 μl template

DNA, and 10.25 μl MoBio certified DNA free water and used the following cycling protocol: 98 degrees

for 30 seconds (1x) followed by 98 degrees for 10 seconds, 61 degrees for 20 seconds, and 72 degrees

for 20 seconds (20x), followed by 72 degrees for 2 minutes and 4 degrees. The amplification products

were cleaned up using Ampure XP beads and normalized into a final pool with an Eppendorf liquid

handling robot. Libraries were sequenced as part of a 150bp paired-end sequencing run on the Illumina

NextSeq platform following the manufacturer's protocol.

We used a custom pipeline that contained the following steps: joining pair-end reads; removing

low-quality and chimeric reads; dereplication and dropping unique reads with low abundance; clustering

OTUs; making OTU table; alignment; building a reference tree; and taxon assignment using FLASH

(Magoc & Salzberg, 2011), the FASTX Toolkit (Hannon Lab, 2010), and the USEARCH pipeline (Edgar,

2010). See https://github.com/kstagaman/Process_16S and Goodfellow et al. (2019) for further details.

We performed *de novo* OTU picking in UCLUST (Edgar, 2010), and sequences with 97% overlap were

defined as belonging to the same bacterial Operational Taxonomic Unit (OTU). After these steps, we had

a total of 2,597 OTUs and a mean read depth of 91,833 (range: 68,252 – 120,119). To guard against

sequencing errors, we filtered out OTU's with a frequency lower than 0.00005 as recommended

(Bokulich et al., 2012).

783
Variation in predictor and outcome variables

Of the females included in the analyses with behavioural predictor variables (Table A2), dietary

dissimilarity (i.e., Sørensen diversity index) varied from 0 to 1, dissimilarity in relatedness calculated as

their *R*-value subtracted from 1 ranged from 0.31 to 1, and social distance (i.e., Geodesic distance in the

1-meter proximity network) varied from 1 to 10 where 10 represents unconnected dyads (Fig. A2).

In our full data set, mean unweighted Unifrac distances within the same season and year was

0.052 ± 0.004 for samples collected from the same individual (N = 4 samples) and 0.205 ± 0.042 for

samples collected from different individuals within the same season and year (N = 82 samples). The low

amount of within-individual variation in comparison to the between-individual variation suggests that

one sample per individual is representative of its gut microbiome during that season and sufficient for

analysis of beta-diversity. Furthermore, the beta-diversity of matched samples from the same adult

female in the wet season 2007 and the dry season 2009 (N = 22 samples from 11 females) was lower

(0.152 ± 0.031) than the female's mean beta-diversity with samples from a different female and year

(0.183 ± 0.021) for all but one female, and there was a significant difference in beta-diversity between

samples from the same versus different females in this sample (N = 11 females, Wilcoxon signed rank

test, p < 0.001).

Year	Group	AF group size	AF sampled	AF omitted	Reason for omitting samples
2007	BS	4	4	0	
2007	DA	5	5	1	*Only omitted one female with incomplete dietary information
2007	NP	4	4	0	
2007	RT	6	5	0	
2007	SP	4	4	0	
2007	WW	9	9	0	
2009	BO	8	8	0	
2009	BS	6	1	1	Lacked samples from majority of AF
2009	DA	7	2	2	Lacked samples from majority of AF
2009	NP	5	3	3	Lacked samples from majority of AF
2009	OD	6	6	0	
2009	RT	7	2	2	Lacked samples from majority of AF
2009	SP	3	0	-	
2009	WW	7	5	0	

801
802   **Table A2.** Number of adult females (AF) present, sampled, and omitted from data analyses with

803   behavioural predictor variables (i.e., Mantel tests and LMMs).

Figure A2. Variation in dietary dissimilarity, relatedness dissimilarity, and social distance for within-

807 group and between-group dyads.