Interactions between social groups of colobus monkeys (*Colobus vellerosus*) explain similarities in their

gut microbiomes

Abstract

The gut microbiome is structured by social groups in a variety of host taxa. Whether this pattern is

driven by relatedness, similar diets, or shared social environments is under debate because few studies

have had access to the data necessary to disentangle these factors in wild populations. We investigated

whether diet, relatedness, or the 1-meter proximity network best explains differences in the gut

microbiome among 45 female colobus monkeys in 8 social groups residing at Boabeng-Fiema, Ghana.

We combined demographic and behavioural data collected May-August 2007 and October 2008-April

2009 with 16S rRNA sequencing of faecal samples collected during the latter part of each observation

period. Social group identity explained a large percentage of the variation in gut microbiome beta-

diversity. When comparing the predictive power of dietary dissimilarity, relatedness, and connectedness

in the 1-meter proximity network, the models with social connectedness received the strongest support,

even in our analyses that excluded within-group dyads. This novel finding indicates that microbes may

be transmitted during intergroup encounters, which could occur either indirectly via shared

environments or directly via social contact. Lastly, some of the gut microbial taxa that appear to be

transmitted via 1-meter proximity are associated with digestion of plant material, but further research is

needed to investigate whether this type of gut microbe transmission yields health benefits, which could

provide an incentive for the formation and maintenance of social bonds within and between social

groups.

Key words: 16S rRNA gene, between-group encounters, colobines, diet, gut microbiome, microbe

transmission, relatedness, social networks, social transmission, host-microbe

Introduction

The gut microbiome consists of thousands of species that affects its host's nutritional status, immune

function, and behavior (McFall-Ngai et al., 2013). It is associated with parasite resistance and stress

response of hosts in the wild (Koch & Schmid-Hempel, 2011; Vlčková et al., 2018) and with obesity in

captive settings (Turnbaugh et al., 2006). Because of these potential health consequences, it is

important to investigate the acquisition and maintenance of the gut microbiome (Amato, 2016; Archie &

Tung, 2015).

The gut microbiome of individuals or social groups become more distinct with geographic

distance (Barelli et al., 2015; Grieneisen et al., 2019; Hansen et al., 2019; Hird, Carstens, Cardiff,

Dittmann, & Brumfield, 2014; Lankau, Hong, & Mackie, 2012; Phillips et al., 2012), and the microbiome

is structured by social group or family co-residency in a variety of host taxa, such as humans (Lax et al.,

2014; Song et al., 2013; Yatsunenko et al., 2012), non-human primates (Amato et al., 2017; Degnan et

al., 2012; Goodfellow et al., 2019; Orkin, Webb, & Melin, 2019; Springer et al., 2017; Tung et al., 2015),

carnivores (Leclaire, Nielsen, & Drea, 2014; Theis et al., 2013), birds (White et al., 2010), and insects

(Anderson et al., 2012; Koch & Schmid-Hempel, 2011). For example, the gut microbiome of immigrant

male baboons *(Papio cynocephalus)* converge over time with that of their new group members

(Grieneisen, Livermore, Alberts, Tung, & Archie, 2017), and the gut microbiomes of colobus monkeys

(*Colobus vellerosus*) diverged over the course of nine months after a social group fissioned into two

daughter groups (Goodfellow et al., 2019).

The divergence in gut microbiomes with home range separation could potentially be due to

dietary differences, lower degrees of relatedness, or lack of shared social environments (Archie & Tung,

2015; Björk, Dasari, Grieneisen, & Archie, 2019). Diet is suggested to be one of the most important

factors affecting the gut microbiome (Voreades, Kozil, & Weir, 2014). Gut microbial composition

fluctuates within hosts with seasonal or experimental dietary changes (Davenport et al., 2014; David et

al., 2013; Hicks et al., 2018; Leamy et al., 2014; Mallott, Amato, Garber, & Malhi, 2018; Maurice et al.,

2015; Michl et al., 2019; Orkin, Campos, et al., 2019), and dietary similarities may explain whether social

groups have distinct gut microbiomes (Orkin, Webb, et al., 2019). However, this pattern could also

reflect the genetic similarity of hosts in societies where at least some closely related individuals remain

together in their natal group. When this is the case, closely related group members are expected to have

more similar gut microbiomes than non-group members with lower degree of relatedness, because the

host's genetic makeup affects microbe colonization (Opstal & Bordenstein, 2015; Spor, Koren, & Ley,

2011) and a number of genomic regions are associated with gut microbial composition in rodents

(Bonder et al., 2016; Leamy et al., 2014). This may explain why closely related individuals have more

similar gut microbiomes than unrelated individuals in some studies of humans and captive rodents

(Faith et al., 2013; Kovacs et al., 2011; Ley et al., 2005). In contrast, genetic differentiation between

baboon populations was a poor predictor of their gut microbiome (Grieneisen et al., 2019), and

relatedness did not have a significant effect on the gut microbiome in some studies of humans

(Rothschild et al., 2018), non-human primates (Moeller et al., 2016) and carnivores (Leclaire et al.,

2014). Moeller and colleagues (2016) suggest that this may be due to an overriding effect of

transmission among unrelated social partners. Indirect social contact via shared environments, such as

touching common surfaces, may facilitate microbiome transmission within households (i.e., indirect

social transmission) (Lax et al., 2014). Direct social contact such as grooming or sitting in body contact

further increases microbiome transmission (i.e., direct social transmission) between close social partners

within social groups of monkeys (*Alouatta pigra*: Amato et al., 2017; *Papio cynocephalus*: Tung et al.,

2015; Grieneisen et al., 2017) and lemurs *(Eulemur rubriventer:* Raulo et al., 2017*)*. Gut microbiomes are

also more similar among socially connected than disconnected siblings and married couples (Dill-

McFarland et al., 2019). In contrast, social connectedness between non-group members did not predict

gut microbiome similarity in sifakas (*Propithecus verreauxi*) (Perofsky et al., 2017). Even if intergroup

encoutners promote the transmission of microbes, there may not be an association between intergroup

interactiowns and gut microbiome similarity if groups rarely interact at close distances. Taken together,

these studies indicate that gut microbes are transmitted via social interactions within social groups,

while it is unclear whether this is the case for social interactions between social groups.

To investigate whether the pattern of increasing between-individual differences in the gut

microbiome (i.e., beta-diversity) with home range separation is best explained by lower dietary overlap,

relatedness, or social connectedness, we focus on the black-and-white colobus monkeys (*Colobus*

vellerosus) at Boabeng-Fiema, Ghana. This is one of several rare species of arboreal leaf-eating monkeys

distributed across the forested regions of the African tropics, and it is closely related to guerezas

(*Colobus guereza*) and western black-and-white colobus (*Colobus polykomos*) (Ting, 2008). At Boabeng-

Fiema, all colobus social groups utilize a highly folivorous diet, but the most important food species

differ between social groups (Saj & Sicotte, 2007; Teichroeb & Sicotte, 2009). More seeds and fruits are

available during the dry season, during which they eat up to 43% of these food items (Teichroeb &

Sicotte, 2017). To break down hard-to-digest items in their primarily folivorous diet (Saj & Sicotte, 2007;

Teichroeb & Sicotte, 2009), they rely on behavioural traits, physiological traits, and their gut microbiome

(Amato et al., 2016; Lambert, 1998). Possibly due to constraints imposed by their highly folivorous diet,

colobus monkeys spend a low percentage of their time engaging in direct social activities such as

grooming (Teichroeb, Saj, Paterson, & Sicotte, 2003). Female colobus spend on average 3% of their time

within 1 meter and 0.1% of their time grooming each female group member (Wikberg, Ting, & Sicotte,

2014b). However, females still form preferred friendships, which are only occassionally based on kinship

and never based on their relatively weakly expressed dominance hierarchies (Wikberg, Teichroeb,

Bădescu, & Sicotte, 2013; Wikberg, Ting, & Sicotte, 2014a; Wikberg et al., 2014b; Wikberg, Ting, &

Sicotte, 2015). Instead, females prefer to affiliate with females with similar immigration status (Wikberg

et al., 2014b, 2014a) in this population where all males and half of the females disperse (Sicotte et al.,

2017; Teichroeb, Wikberg, & Sicotte, 2009, 2011; Wikberg, Sicotte, Campos, & Ting, 2012). This flexible

female dispersal pattern results in social groups with different female kin composition and some close

100 maternal female kin residing in different social groups (Wikberg et al., 2012). Neighbouring social groups

101 encounter each other in the large zones of home range overlap on an almost daily basis. During these

102 encounters, social groups sometimes chase each other away from food trees, while at other times, they

103 engage in affiliative or sexual between-group interactions (Sicotte & MacIntosh, 2004; Teichroeb &

104 Sicotte, 2017).

105   The frequent between-group interactions coupled with variation in diet and relatedness within

106 and between social groups makes this a good study population to investigate whether the pattern of

107 increasing gut microbial beta-diversity with home range separation is best explained by lower degrees of

108 dietary similarity, relatedness, or social connectedness. We take a cross-sectional approach using

109 observational and genetic data from eight social groups to first test whether the gut microbiome was

110 structured by social groups. We predicted gut microbiome beta-diversity to be structured by social

111 groups and to increase with home range separation. We then evaluated which factors explained gut

112 microbiome beta-diversity between females across different social groups. We expected gut

113 microbiome beta-diversity to decrease with dietary similarity and relatedness and increase with distance

114 in the 1-meter proximity network. Finally, the significant predictor from the analyses above (social

115 connectedness) was used in a subsequent population-level analysis of Operational Taxonomic Unit

116 (OTU) abundance to determine which microbial taxa may be socially transmitted. Our definition of social

117 transmission includes both direct social transmission via physical contact and indirect social transmission

118 via shared substrates (e.g., Perofsky et al., 2017), and we will not attempt here to tease these two social

119 transmission routes. Males and females of all age-classes were used to create social networks, but the

gut microbiome data are only available for adult females. Therefore, our analyses of beta-diversity focus

on adult females.

Methods

Behavioral data collection

Demographic data have been collected since 2000 from the black-and-white colobus monkeys (*Colobus*

vellerosus) at Boabeng-Fiema, Ghana. In this study, we also use behavioral and ecological data as well as

DNA samples from eight social groups (Fig. A1) collected during two study periods: the rainy season

May-August 2007 and the pre-dry and dry seasons October 2008 - April 2009 (Table A2). During this

time period, the study groups contained 3-9 adult (i.e., parous) females (Table A2), 1-4 adult males, and

8-24 immatures. Our research adheres to ASAB/ABS Guidelines for the Use of Animals in Research, the

laws of Ghana, and data collection was approved by the Boabeng-Fiema Monkey Sanctuary's

management committee, Ghana Wildlife Division, and the University of Calgary's Animal Care

Committee (BI 2006-28, BI 2009-25).

We recorded the social group's location every hour using a map with trails, roads, villages, and

large trees (>40 cm DBH) in order to determine home ranges (Fig. A1). During 10-minute focal samples

(Altmann, 1974) of adult females, we continuously recorded all social behaviors (including the identity of

the interactant and the duration of the behavior) and plant species and part (i.e., mature leaf, young

leaf, flower, fruit, seed, or other) for each ingested food item. Females fed on a total of 210 food item-

plant species combinations, and to assess dietary differences, we calculated Sørensen dissimilarity

indices using ingested plant parts and plant species during focal samples. We choose this diversity index

because it only takes the presence or absence of an ingested food item into account, which we have a

robust estimate of using the focal data. The Sørensen dissimilarity indices in our data set had a high

median value of 0.83 and it was lower within than between social groups (Fig. A2).

We observed 61 and 285 between-group encounters (i.e., two social groups located within 50

meters of each other) during the first and second data collection period respectively. Of these

encounters, 53% lacked female aggression and 35% lacked male aggression. Because close proximity

between individuals of different social groups are rare and unlikely to be recorded during focal sampling,

we recorded approaches to 1 meter *ad libitum* (Altmann, 1974). Some of these approaches only led to

brief close proximity while others led to prolonged contact like copulations, grooming, and play. We

created an undirected proximity network based on the presence and absence of approaches to 1 meter

between all individuals (N = 177 adult females, adult males, and immatures) present in the eight study

groups. We used the software UCINET (Borgatti, Everett, & Freeman, 2002) to compute inverse shortest

path length (i.e., Geodesic distance) in the 1-meter proximity network (hereafter referred to as social

connectedness): 1/[the number of steps (i.e., recorded interaction ties) in the shortest path from one

individual to another]. Social group members were in 1-meter proximity with each other (i.e., an inverse

path length of 1) or separated by two to three partners (i.e., an inverse path length of 0.5 and 0.33) (Fig.

A2). The inverse path length for males and females belonging to different social groups ranged from 0 to

1 (Fig. 1; Fig. A2). The seemingly unconnected individuals in the 2007 data set were most likely

unconnected because we only had access to data collected from a 3-month period. These individuals

were connected and separated with up to eight steps in the 2008-2009 network, which was based on six

months of data.

Genetic data collection

We collected faecal samples June-August 2007 and January-April 2009. Immediately after a female

defecated, we collected approximatey 1g of feces and dissolved it in 6ml RNAlater. The samples were

stored in a fridge at the field site until the end of the field season when they were transported to the

Ting lab and stored in a -20-degree C freezer. Note that we lack information on soil type, which was

driving between-site differences in the gut microbiomes in a large-scale study of terrestrial baboons

(Grieneisen et al., 2019). However, our samples were collected from arboreal primates within a small

study area, and sampling site does not have a significant effect on beta-diversity in our study population

(Goodfellow et al., 2019).

(a) 2007

(a) 2009

(b) 2007

(b) 2009

Focal Group   Unweighted Uni−Frac

BS   NP   SP

DA   RT   WW   0.10 0.15 0.20 0.25 0.30

Focal Group   Unweighted Uni−Frac

BO   OD   WW

0.20   0.25

Figure 1. Social networks for: **a)** the entire population where each group member is depicted as a node

in yellow (group used for behavioural analyses) or blue (group not used for behavioural analyses) and

between-group dyads observed in 1-meter proximity are connected with lines; **b)** females included in

the behavioural analyses with lines connecting between-group dyads (i.e., nodes of different color)

where color represents gut microbiome beta-diversity (i.e., unweighted Unifrac distances) ranging from

similar (dark) to dissimilar (light) and thickness indicates social connectedness ranging from strongly

connected (thick) to more disconnected (thin). The black lines connect group members and are not

weighted based on beta-diversity or social connectedness.

We extracted DNA from the samples and genotyped the extracts at 17 short tandem repeat loci

(STR) as previously described (Wikberg et al., 2012). To make sure that the samples used in the

relatedness and gut microbiome analyses were collected from the correct individual, we compared the

STR genotypes obtained from these samples with a second sample collected from the same individual at

a different time. We calculated dyadic estimated relatedness values (*R*) in MLRelate (Kalinowski,

Wagner, & Taper, 2006) because this method provided the most accurate relatedness estimates in our

study population (Wikberg et al., 2012). We used *R*-values calculated from STR loci rather than

theoretical relatedness (*r*) calculated from pedigrees, because *R*-values predict kinship relatively

accurately in our study population (Wikberg et al., 2014a) and they are more accurate than *r* in studies

such as ours with limited access to pedigrees (Forstmeier, Schielzeth, Mueller, Ellegren, & Kempenaers,

2012; Robinson, Simmons, & Kennington, 2013). The median female relatedness was low both within

and between social groups, but there were at least some closely related females residing in the same

social groups (Fig. A2).

For generating the gut microbial data, we conducted fresh DNA extracts from 61 previously

genotyped samples from 45 females (Table A2) using the QIAamp DNA Stool Mini Kit with a modified

protocol. More details regarding the extraction protocol are presented in the Appendix and in

Goodfellow et al. (2019). The V4 hypervariable region of the bacterial 16S ribosomal RNA gene was

amplified and libraries were prepared using the 515F and 806R primers containing 5' Illumina adapter

tails and dual indexing barcodes, and libraries were sequenced as part of a 150bp paired-end sequencing

run on the Illumina NextSeq platform following Goodfellow et al. (2019). We obtained a mean read

depth of 127,628 per sample (range: 86,924-166,438). Then, we used a custom pipeline

(https://github.com/kstagaman/Process_16S) for quality filtering and assembly (see Appendix). We

performed *de novo* OTU picking in UCLUST (Edgar, 2010), and sequences with 97% overlap were defined

as belonging to the same bacterial Operational Taxonomic Unit (OTU). After this processing, we had a

total of 2,597 OTUs and an average of 89,483 reads per sample (range: 59,817-120,119). To further

guard against sequencing errors, we filtered out OTU's with a frequency lower than 0.00005 as

recommended (Bokulich et al., 2012). After filtering, the 2007 data set contained 450 OTUs and the

2009 data set contained 396 OTUs. The mean read depth was 88,346 (range: 59,005 – 118,633). We did

not rarefy the data set to an even read depth, because it is recommended against (McMurdie & Holmes,

2014). First, rarefying leads to increased false positives and decreased true positives, especially in data

sets with read depths comparable to ours (Pereira et al., 2018). Second, unrarefied counts are

particularly accurate when using our measure of beta-diversity—weighted UniFrac distances (McMurdie

& Holmes, 2014).

We initially calculated four different measures of gut microbiome beta-diversity (Sørensen

dissimilarity index, Bray-Curtis dissimilarity index, unweighted UniFrac distances, and weighted UniFrac

distances) in the R package vegan (Oksanen et al., 2017). Because the two presence/absence indices

were strongly correlated with each other (Sorenson dissimilarity indices and unweighted UniFrac

distances: Mantel r = 0.93, p = 0.001) as were the two abundance indices (Bray-Curtis dissimilarity

indices and weighted UniFrac distances: Mantel r = 0.77, p = 0.001), in our analyses, we only used the

one presence/absence index (unweighted UniFrac distances) and the one abundance index (weighted

UniFrac distances) that take phylogenetic relationships of OTUs into account.

Data analyses

We combined the 2007 and 2009 data sets and included study year (aka season) and individual ID as

predictor variables whenever possible (i.e., permutational multivariate analysis of variance and linear

mixed models) while we had to create squared interaction matrices for each study year separately when

using matrix correlations (i.e., Mantel tests and Moran's test for autospatial correlations). We only used

the full data set (N = 61 samples from 45 females) for the initial analysis regarding the effect of social

group identity. All subsequent analyses examined the effects of behavioural variables on beta-diversity

in a subset (N = 49 samples from 42 unique females) from which we removed: 1) duplicate samples from

the same year and same female; 2) one adult female with incomplete dietary information, and 3) social

groups from which the majority of females remained unsampled to make sure we had a representative

sample of social connectedness from each social group.

The initial analysis investigated the effects of season, social group, individual identity, and read

depth on beta-diversity of all dyads in the full data set (N = 61 samples from 45 females in 2007 and

2009) using permutational multivariate analysis of variance (PERMANOVA) with 10,000 permutations

using the adonis function in the R package vegan (Oksanen et al., 2017). The terms were added

sequentially in the order listed above.

We used non-parametric Mantel correlations implemented in the R package vegan (Oksanen et

al., 2017) to investigate whether the two measures of gut microbiome beta-diversity were correlated

with home range separation (0 = same social group and home range; 1 = different social groups but

adjacent home ranges, 2 = different social groups and non-adjacent home ranges) using beta-diversity

indices from 30 samples from unique females in 6 social groups in 2007 and 19 samples from unique

females in 3 social groups in 2009. We used beta-diversity indices of all dyads, but analysed the two

years separately.

To investigate which combination of dyadic traits predicted gut microbiome beta-diversity

between females, we created generalized linear mixed models (GLMMs) with the outcome variable gut

microbiome beta-diversity using the beta family function in the package glmmTMB (Magnusson et al.,

2019) in R (R Core Team, 2018). Again, we used 30 samples from unique females in 6 social groups in

2007 and 19 samples from unique females in 3 social groups in 2009. We created a null model that did

not contain any fixed effects, alternative models with one fixed effect that represented one of the

hypotheses outlined in the introduction (dietary dissimilarities, *R*-values, or social connectedness), and a

full model with all three predictor variables. We included data collection year as a fixed effect in all

alternative models because the two sampling years occurred in different seasons and several other

studies show strong seasonal shifts in gut microbiome composition (Amato et al., 2015; Hicks et al.,

2018; Orkin, Campos, et al., 2019; Smits et al., 2017; Springer et al., 2017). All numerical predictor

variables were centered and scaled (Schielzeth, 2010). We included social group and focal identities as

random effects in all GLMMs, including the null models. We did not have any issues with collinearity

based on low Variance Inflation Factors for the full models (all VIF < 1.43). We evaluated the support for

each model using Akaike Information Criterion (AIC) (Akaike, 1974), and this approach allowed us to

determine which hypotheses (diet, relatedness, or social connectedness) was best supported by our

data (Burnham & Anderson, 2002). Because several models received similar support, we took model

selection uncertainly into account by averaging coefficients across models (Burnham & Anderson, 2002)

using the R package MUMIN (Barton, 2013). In the first set of analyses, we included dyads that resided

in the same social group and dyads that resided in different social groups. To make sure that the effect

of social connectedness was not driven by the close social bonds within social groups, we repeated the

analyses with between-group dyads only.

To infer which of the gut microbial taxa may be transmitted via close proximity, which was a

better predictor of beta-diversity than diet and relatedness (see Results), we investigated whether the

abundance of each OTU was correlated with Geodesic distance in the 1-meter approach network using

Moran's test for autospatial correlations implemented in the package ape (Paradis, Claude, & Strimmer,

2004). We included within-group and between-group dyads in this analysis (N = 342 dyads). We counted

the number of OTUs in each phylum (or family) that were socially structured based on the autospatial

correlation results. We conducted hypergeometric tests to investigate whether this number was higher

than expected by chance based on the total number of OTUs in the phylum (or family) using the phyper

function implemented in R. In all analyses of taxonomic differences, we used the 10% false discovery

rate to correct p-values for multiple testing (sensu Tung et al., 2015). The gut microbial taxa we

expected to be shaped by sociality are listed in the Table A1 (Amato et al., 2017; Goodfellow et al., 2019;

Tung et al., 2015).

Results

Factors predicting gut microbiome beta-diversity

We investigated the relative effects of season, social group, individual ID, and read depth in the full data

set (PERMANOVA: N = 61 samples collected 2007-2009). Of the observed variation in the taxonomic

composition of the gut microbiome (i.e., beta-diversity), individual identity explained the largest

percentage (54-55% depending on which beta-diversity index was used as outcome variable), social

group identity explained a more moderate percentage (19-28%), while year explained much smaller

percentage (8-12%) (Table 1). Read depth did not have a significant effect on beta-diversity (Table 1).

Beta-diversity index	Factor	Df	Sums of squares	Mean squares	F	R^2	P
Unweighted UniFrac	Season	1	0.103	0.103	12.325	0.084	<0.001
	Group	7	0.347	0.050	5.943	0.282	<0.001
	ID	39	0.663	0.017	2.035	0.538	<0.001
	Read depth	1	0.010	0.010	1.185	0.008	0.249

Weighted UniFrac	Season	1	0.136	0.136	12.277	0.118	<0.001
	Group	7	0.219	0.031	2.812	0.189	<0.001
	ID	39	0.639	0.016	1.477	0.553	<0.001
	Read depth	1	0.018	0.018	1.637	0.016	0.104

Table 1. Results from the PERMANOVA with factors added sequentially in the ordered listed in the table.

Gut microbiome beta-diversity and home range separation were correlated in the 2007 data set

(N = 870 dyads in 6 social groups, Mantel tests: unweighted UniFrac distance: r = 0.22, P = 0.002;

weighted UniFrac distance: r = 0.10, P = 0.049) and in the 2009 data set (N = 342 dyads in 3 social

groups, Mantel tests: unweighted UniFrac distance: r = 0.36, P = 0.005; weighted UniFrac distance: r =

0.20, P = 0.024), meaning that females residing farther from each other had less similar gut

microbiomes. This pattern can potentially be explained by group members having more similar diets,

higher relatedness, or stronger social connectedness than non-group members (Fig. A2).

We created several competing generalized linear mixed models to investigate which of the three

hypotheses best explained increasing beta-diversity with home range separation: dietary dissimilarity,

relatedness, or social connectedness, controlling for data collection year. In our data set with both

within-group and between-group dyads (N = 1,212 dyads in 2007-2009), the full models and the models

with social connectedness received the greatest support (Table 2). Social connectedness predicted gut

microbiome beta-diversity, and females located further apart in the social network had less similar gut

microbiomes (Figs. 1-2). Year also predicted gut microbiome beta-diversity (Fig. 2), and females had

more similar gut microbiomes during the rainy season of 2007 than the dry season of 2009. In contrast,

diet and relatedness did not have significant effects on gut microbiome beta-diversity (Fig. 2).

Outcome variable	Fixed effect	AIC	Delta	Weight
Unweighted UniFrac	Season + Proximity	-5247.30	0.00	0.73
	Season + Diet + Relatedness + Proximity	-5245.28	2.02	0.27
	Season + Diet	-5103.63	143.67	0.00
	Season + Relatedness	-5060.36	186.94	0.00
	-	-4940.73	306.58	0.00
Weighted UniFrac	Season + Proximity	-4684.20	0.00	0.56
	Season + Relatedness	-4683.69	0.51	0.44
	Season + Diet	-4658.94	25.25	0.00
	Season + Diet + Relatedness + Proximity	-4657.85	26.35	0.00
	-	-4614.36	69.84	0.00
Unweighted UniFrac	Season + Proximity	-4307.62	0.00	0.76
	Season + Diet	-4303.92	3.71	0.12
	Season + Relatedness	-4302.67	4.95	0.06
	Season + Diet + Relatedness + Proximity	-4302.44	5.19	0.06
	-	-4119.55	188.07	0.00
Weighted UniFrac	Season + Proximity	-3770.21	0.00	0.64
	Season + Diet	-3768.91	1.30	0.33
	Season + Relatedness	-3762.64	7.58	0.01
	Season + Diet + Relatedness + Proximity	-3761.96	8.25	0.01
	-	-3713.73	56.48	0.00

Table 2. The competing GLMMs' fixed effects, Akaike Information Criterion, delta (i.e., difference in AIC between the current model and the best-fit model), and Akaike weights (i.e., relative likelihood of the model), and marginal and conditional R^2 for the best fitting model (i.e., without versus with random effects) when including **a)** within-group and between-group dyads and **b)** only between-group dyads.

To assess whether the effect of social connectedness on gut microbiome beta-diversity was driven by closely connected within-group dyads having very similar gut microbiomes, we repeated the analyses with between-group dyads only (N = 966 dyads). The full models and the social connectedness

models were again the strongest supported models (Table 2). Beta-diversity was predicted by year and

social connectedness, but not by diet and relatedness (Fig. 2).

Figure 2. Coefficient estimates and their 95% confidence intervals for the best fitting model for

unweighted versus weighted UniFrac distances in the data set including within-group and between-

group dyads and in the data set with between-group dyads only. Social connectedness, dietary

dissimilarity, and relatedness ranges from 0 to 1. Of the two study periods, year 2007 was used as the

baseline level against which we depict the effect of year 2009.

Socially structured OTUs

Our data set contained OTUs from 14 phyla, of which the most well-represented was Firmicutes,

followed by Bacteroidetes, Spirochetes, and Verrucomicrobia (Supplementary Material Fig. 1). In each

social group, at least 70% of the OTUs belonged to the phylum Firmicutes (Supplementary Material Fig.

1) and at least 50% of the OTUs belonged to the families Lachnospiraceae and Ruminococcaceae in the

phylum Firmicutes (Supplementary Material Fig. 2).

Social connectedness predicted differences in abundances for 73 of the 396 OTUs in the 2009

data set (Moran's I range: -0.27 – -0.14, all $P < 0.05$, Supplementary Material Table 1). The number of

OTUs with a significant relationship to social connectedness was greater than expected in the phylum

Firmicutes (N = 64, Hypergeometric test: $P < 0.001$). The numbers of socially structured OTUs in the

phyla Bacteroidetes (N=6), Planctomycetes (N = 1), Proteobacteria (N = 1) and Tenericutes (N = 1) were

not greater than expected based on the total number of OTUs in these phyla (Hypergeometric tests, $p >$

0.050). The other phyla did not contain any socially structured OTUs. Four families had a higher than

expected number of socially structured OTUs: Bacteroidaceae (N = 4), Lachnospiraceae (N = 20),

Peptococcaceae 2 (N = 1), and Ruminococcaceae (N = 31). There was also a greater than expected

number of socially structured OTUs in 14 of 34 genera (Fig. 4).

Figure 4. The proportion of Operational Taxonomic Units (OTUs) whose abundance was correlated with

distance in the proximity network for genera that contained a higher than expected number of socially

structured OTUs.

Social connectedness predicted differences in abundances for 1 of the 450 OTUs in the 2007

data set (Moran's I range: -0.27 – -0.14, all $p < 0.05$), which belonged to the phylum Firmicutes, the

family Lachnospiraceae, and the genus *Roseburia*. As a result, these taxa had a greater than expected

number of socially structured OTUs (Hypergeometric tests, all $p > 0.001$).

Discussion

The aim of this study was to investigate whether the increase in gut microbiome beta-diversity with

home range separation in female colobus monkeys was best explained by diet, relatedness, or sociality.

Distance in the proximity network was a better predictor than diet and relatedness, similar to findings in

more social primates (Amato et al., 2017; Perofsky et al., 2017; Raulo et al., 2017; Tung et al., 2015).

Although these previous studies suggest that strong social bonds within social groups drive between-

group differences in the gut microbiome after ruling out the effects of relatedness and diet, this is the

first report of a relationship between gut microbiome beta-diversity and social connectedness between

individuals in different social groups. In contrast, gut microbiome dissimilarity between individuals

residing in different social groups did not increase with grooming network distance in sifakas (Perofsky

et al., 2017). These contrasting results may be due to the nature or frequency of the host population's

between-group interactions. Colobus monkeys sometimes engage in affiliative, sexual, and playful

behaviours with non-group members (Supplemental Information; Sicotte & MacIntosh, 2004; Teichroeb,

Marteinson, & Sicotte, 2005; Teichroeb et al., 2011), which differ from the almost exclusively aggressive

nature of between-group encounters in some other taxa. Similar to these colobus monkeys, mountain

gorillas (*Gorilla beringei beringei*) occasionally affiliate with members from other social groups (Forcina

et al., 2019) and human foraging societies form extended social networks to optimize resource flow

(Hamilton, Milne, Walker, Burger, & Brown, 2007). These extended networks could possibly affect their

gut microbiome in similar ways as documented here in colobus monkeys.

To determine the consequences of such socially-mediated transmission, the first step is to

determine which types of microbes are transmitted this way. The socially transmitted OTUs in this study

included all taxa (family Porphyromonadaceae and genera *Parabacteroides* and *Coprococcus)* that

diverged after a social group fission at our site (Goodfellow et al., 2019) and genera (*Bacteroides,*

Clostridium, and *Roseburia*) that were transmitted via grooming and close proximity in howlers (Amato

et al., 2017). The close match in socially transmitted taxa in howlers and colobus is not particularly

surprising given both have a folivorous diet and low degree of terrestriality, which are factors that

influence the gut microbiome (Perofsky et al., 2019). In contrast, the socially transmitted OTUs in our

study did not overlap with those transmitted via grooming within social groups of baboons (Tung et al.,

2015), despite the host species relatively close phylogenetic relationship. Recent findings show that host

phylogeny has a stronger effect than diet on gut microbiome composition (Amato et al., 2019), and it is

thus possible that while phylogeny has the strongest overall effect on the gut microbiome, the same gut

microbial taxa are structured by sociality in primates with similar lifestyle.

We found that the majority of socially transmitted OTUs belonged to the most dominant

families in our host population and other folivorous primates (Barelli et al., 2015; Perofsky et al., 2017),

the families Lachnospiraceae and Ruminococcaceae in the phylum Firmicutes. These taxa are well-suited

for breaking down hard-to-digest plant material (Biddle, Stewart, Blanchard, & Leschine, 2013), and it is

therefore possible that socially transmitted gut microbes benefit hosts in terms of improved digestion of

mature leaves, which make up the majority of the colobus diet (Saj & Sicotte, 2007). Several studies

imply that socially-mediated transmission benefits the host. For example, Tung and colleagues (2015)

suggest that the positive health and fitness effects that baboons accrue from forming close social bonds

with group members are mediated by the gut microbiome. Our results support the notion that this may

be the case in a wide range of gregarious species, including those with relatively low frequencies of

social interactions. If social transmission sustains a healthy gut microbiome (as documented in Koch &

Schmid-Hempel, 2011), it could provide an incentive for the formation and maintenance of social bonds

within social groups (Lombardo, 2008). Our findings leave open the as-of-yet unexplored possibility that

social transmission of microbes may even explain the occurrence of friendly between-group encounters,

especially in the absence of limiting resources such as fertile females and important food sources.

The results of this paper ultimately lead us to an important outstanding question, which is how

gut microbes are transmitted among animals that spend considerably less time grooming or in direct

contact than other primates with socially mediated gut microbe transmission (Amato et al., 2017; Raulo

et al., 2017; Tung et al., 2015). It might be that microbes are transmitted directly during the occasions

we observed non-group members copulating, grooming, and playing. However, it could also be that the

microbes are transmitted indirectly between hosts when they are touching shared surfaces within a

certain time period (Münger et al., 2018). This reasoning is consistent with spatial proximity predicting

the gut microbiome in other gregarious species with low frequencies of social behaviours like the Welsh

Mountain ponies (*Equus ferus caballus*) (Antwis et al. 2018) and in more solitary species such as North

American red squirrels (*Tamiasciurus hudsonicus*) (Ren et al., 2017) and gopher tortoise (*Gopherus*

polyphemus) (Yuan et al., 2015). The occurrences of direct and indirect social transmission are difficult

to tease apart when the two are correlated and when brief physical contact between extra-group

members often go unnoticed, but carefully designed studies in the future may be able to address this

question.

Finally, relatedness and dietary differences within a season were not good predictors of beta-

diversity in comparison to social connectedness. In contrast, seasonal changes in diet may be associated

with changes in the colobus gut microbiome, because beta-diversity was higher during the 2009 dry

season when their diet was more diverse than during the 2007 rainy season when they ate mostly

mature leaves. We will continue to investigate whether this seasonal dietary switch is linked to changes

in the gut microbiome, as previously reported from other species inhabiting seasonal environments

(Amato et al., 2015; Hicks et al., 2018; Orkin, Campos, et al., 2019; Smits et al., 2017; Springer et al.,

2017). These authors concluded that gut microbiome dynamics determine nutrient uptake and is key for

dietary flexibility (Amato et al., 2015; Hicks et al., 2018; Orkin, Campos, et al., 2019; Smits et al., 2017;

Springer et al., 2017), while the potential three-way interaction between social, dietary, and gut

microbial dynamics is still poorly understood. An interesting venue for further research is therefore to

investigate whether the gut microbiomes of socially well-connected individuals map more quickly onto

ecological changes, which could help them adjust to the rapidly changing environments that many wild

animals inhabit today.

Acknowledgements

We would like to thank two anonymous reviewers for their help with improving this manuscript; Keaton

Stagaman for use of his custom pipeline; Fernando Campos, Teresa Holmes, and Robert Koranteng for

field data collection; BFMS management committee, Ghana Wildlife Division, and the University of

Calgary's Life and Environmental Sciences Animal Care Committee for permission to conduct this study;

Alberta Innovates Technology Futuresn (200600343), American Society of Primatologists, Animal

Behaviour Society, International Primatological Society, Leakey Foundation, Natural Sciences and

Engineering Research Council of Canada, Sweden-America Foundation, Research Services at the

University of Calgary, the National Institute of General Medical Sciences (P50GM098911) via the META

Center for Systems Biology, and Wenner-Gren Foundation (8172) for funding.

References

Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic*

Control, *19*(6), 716–723. doi: 10.1109/TAC.1974.1100705

Altmann, J. (1974). Observational study of behavior: sampling methods. *Behaviour*, *49*(3), 227–267.

Amato, K. R. (2016). Incorporating the gut microbiota into models of human and non-human primate

ecology and evolution: Gut microbiota influences host nutrition, health, and behavior. *American*

Journal of Physical Anthropology, *159*, 196–215. doi: 10.1002/ajpa.22908

Amato, K. R., G Sanders, J., Song, S. J., Nute, M., Metcalf, J. L., Thompson, L. R., … R Leigh, S.

(2019). Evolutionary trends in host physiology outweigh dietary niche in structuring primate gut

microbiomes. *The ISME Journal*, *13*(3), 576–587. doi: 10.1038/s41396-018-0175-0

Amato, K. R., Leigh, S. R., Kent, A., Mackie, R. I., Yeoman, C. J., Stumpf, R. M., … Garber, P. A.

(2015). The gut microbiota appears to compensate for seasonal diet variation in the wild black

howler monkey (*Alouatta pigra*). *Microbial Ecology*, *69*(2), 434–443. doi: 10.1007/s00248-014-

0554-7

Amato, K. R., Metcalf, J. L., Song, S. J., Hale, V. L., Clayton, J., Ackermann, G., … Braun, J. (2016).

Using the gut microbiota as a novel tool for examining colobine primate GI health. *Global*

Ecology and Conservation, *7*, 225–237. doi: 10.1016/j.gecco.2016.06.004

Amato, K. R., Van Belle, S., Di Fiore, A., Estrada, A., Stumpf, R., White, B., … Leigh, S. R. (2017).

Patterns in gut microbiota similarity associated with degree of sociality among sex classes of a

Neotropical primate. *Microbial Ecology*. doi: 10.1007/s00248-017-0938-6

Anderson, K. E., Russell, J. A., Moreau, C. S., Kautz, S., Sullam, K. E., Hu, Y., … Wheeler, D. E.

(2012). Highly similar microbial communities are shared among related and trophically similar

ant species. *Molecular Ecology*, *21*(9), 2282–2296. doi: 10.1111/j.1365-294X.2011.05464.x

Antwis, R. E., Lea, J. M. D., Unwin, B., & Shultz, S. (2018). Gut microbiome composition is associated

with spatial structuring and social interactions in semi-feral Welsh Mountain ponies. *Microbiome*,

6(1), 207. doi: 10.1186/s40168-018-0593-2

Archie, E. A., & Tung, J. (2015). Social behavior and the microbiome. *Current Opinion in Behavioral*

Sciences, *6*, 28–34. doi: 10.1016/j.cobeha.2015.07.008

Barelli, C., Albanese, D., Donati, C., Pindo, M., Dallago, C., Rovero, F., … Filippo, C. D. (2015). Habitat

fragmentation is associated to gut microbiota diversity of an endangered primate: implications for

conservation. *Scientific Reports*, *5*, 14862. doi: 10.1038/srep14862

Barton, K. (2013). *MuMIn: Multi-model inference. R package version 1.9.5. http://CRAN.R-*

project.org/package=MuMIn. Accessed 1 May 2013.

Biddle, A., Stewart, L., Blanchard, J., & Leschine, S. (2013). Untangling the genetic basis of fibrolytic

specialization by Lachnospiraceae and Ruminococcaceae in diverse gut communities. *Diversity*,

5(3), 627–640. doi: 10.3390/d5030627

Björk, J. R., Dasari, M., Grieneisen, L., & Archie, E. A. (2019). Primate microbiomes over time:

Longitudinal answers to standing questions in microbiome research. *American Journal of*

Primatology, *0*(0), e22970. doi: 10.1002/ajp.22970

Bokulich, N. A., Subramanian, S., Faith, J. J., Gevers, D., Gordon, J. I., Knight, R., … Caporaso, J. G.

(2012). Quality-filtering vastly improves diversity estimates from Illumina amplicon sequencing.

Nature Methods, *10*(1), 57–59. doi: 10.1038/nmeth.2276

Bonder, M. J., Kurilshikov, A., Tigchelaar, E. F., Mujagic, Z., Imhann, F., Vila, A. V., … Zhernakova, A.

(2016). The effect of host genetics on the gut microbiome. *Nature Genetics*, *48*(11), 1407–1412.

doi: 10.1038/ng.3663

Borgatti, S., Everett, M., & Freeman, L. (2002). *Ucinet for Windows: software for social network*

analysis. Harvard, MA: Analytic Technologies.

Burnham, K., & Anderson, D. (2002). *Model selection and multimodel inference: a practical*

information-theoretic approach. New York: Springer.

Davenport, E. R., Mizrahi-Man, O., Michelini, K., Barreiro, L. B., Ober, C., & Gilad, Y. (2014). Seasonal

Variation in Human Gut Microbiome Composition. *PLoS ONE*, *9*(3), e90731. doi:

10.1371/journal.pone.0090731

David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe, B. E., …

Turnbaugh, P. J. (2013). Diet rapidly and reproducibly alters the human gut microbiome. *Nature*,

505(7484), 559–563. doi: 10.1038/nature12820

Degnan, P. H., Pusey, A. E., Lonsdorf, E. V., Goodall, J., Wroblewski, E. E., Wilson, M. L., … Ochman,

H. (2012). Factors associated with the diversification of the gut microbial communities within

chimpanzees from Gombe National Park. *Proceedings of the National Academy of Sciences*,

109(32), 13034–13039. doi: 10.1073/pnas.1110994109

Dill-McFarland, K. A., Tang, Z.-Z., Kemis, J. H., Kerby, R. L., Chen, G., Palloni, A., … Herd, P. (2019).

Close social relationships correlate with human gut microbiota composition. *Scientific Reports*,

9(1), 703. doi: 10.1038/s41598-018-37298-9

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*,

26(19), 2460–2461. doi: 10.1093/bioinformatics/btq461

Faith, J. J., Guruge, J. L., Charbonneau, M., Subramanian, S., Seedorf, H., Goodman, A. L., … Gordon, J.

I. (2013). The Long-Term Stability of the Human Gut Microbiota. *Science*, *341*(6141), 1237439–

1237439. doi: 10.1126/science.1237439

Forcina, G., Vallet, D., Le Gouar, P. J., Bernardo-Madrid, R., Illera, G., Molina-Vacas, G., … Vila, C.

(2019). From groups to communities in western lowland gorillas. *Proceedings of the Royal*

Society B-Biological Sciences, *286*(1896), 20182019. doi: 10.1098/rspb.2018.2019

Forstmeier, W., Schielzeth, H., Mueller, J. C., Ellegren, H., & Kempenaers, B. (2012). Heterozygosity-

fitness correlations in zebra finches: microsatellite markers can be better than their reputation.

Molecular Ecology, *21*(13), 3237–3249. doi: 10.1111/j.1365-294X.2012.05593.x

Goodfellow, C. K., Whitney, T., Christie, D. M., Sicotte, P., Wikberg, E. C., & Ting, N. (2019).

Divergence in gut microbial communities mirrors a social group fission event in a black-and-

white colobus monkey (Colobus vellerosus). *American Journal of Primatology*, *0*(0), e22966.

doi: 10.1002/ajp.22966

Grieneisen, L. E., Charpentier, M. J. E., Alberts, S. C., Blekhman, R., Bradburd, G., Tung, J., & Archie,

E. A. (2019). Genes, geology and germs: gut microbiota across a primate hybrid zone are

explained by site soil properties, not host species. *Proceedings of the Royal Society B: Biological*

Sciences, *286*(1901), 20190431. doi: 10.1098/rspb.2019.0431

Grieneisen, L. E., Livermore, J., Alberts, S., Tung, J., & Archie, E. A. (2017). Group Living and Male

Dispersal Predict the Core Gut Microbiome in Wild Baboons. *Integrative and Comparative*

Biology, *57*(4), 770–785. doi: 10.1093/icb/icx046

Hamilton, M. J., Milne, B. T., Walker, R. S., Burger, O., & Brown, J. H. (2007). The complex structure of

hunter-gatherer social networks. *Proceedings of the Royal Society B: Biological Sciences*,

274(1622), 2195–2203. doi: 10.1098/rspb.2007.0564

Hannon Lab. (2010). *FASTX Toolkit. Available from http:// hannonlab.cshl.edu/fastx_toolkit/index.html.*

Hansen, M. E. B., Rubel, M. A., Bailey, A. G., Ranciaro, A., Thompson, S. R., Campbell, M. C., …

Tishkoff, S. A. (2019). Population structure of human gut bacteria in a diverse cohort from rural

Tanzania and Botswana. *Genome Biology*, *20*(1), 16. doi: 10.1186/s13059-018-1616-9

Hicks, A. L., Lee, K. J., Couto-Rodriguez, M., Patel, J., Sinha, R., Guo, C., … Williams, B. L. (2018).

Gut microbiomes of wild great apes fluctuate seasonally in response to diet. *Nature*

Communications, *9*(1), 1786. doi: 10.1038/s41467-018-04204-w

Hird, S. M., Carstens, B. C., Cardiff, S. W., Dittmann, D. L., & Brumfield, R. T. (2014). Sampling

locality is more detectable than taxonomy or ecology in the gut microbiota of the brood-parasitic

Brown-headed Cowbird (*Molothrus ater*). *PeerJ*, *2*, e321. doi: 10.7717/peerj.321

Kalinowski, S. T., Wagner, A. P., & Taper, M. L. (2006). ML-RELATE: a computer program for

maximum likelihood estimation of relatedness and relationship. *Molecular Ecology Notes*, *6*(2),

576–579. doi: 10.1111/j.1471-8286.2006.01256.x

Koch, H., & Schmid-Hempel, P. (2011). Socially transmitted gut microbiota protect bumble bees against

an intestinal parasite. *Proceedings of the National Academy of Sciences*, *108*(48), 19288–19292.

doi: 10.1073/pnas.1110474108

Kovacs, A., Ben-Jacob, N., Tayem, H., Halperin, E., Iraqi, F. A., & Gophna, U. (2011). Genotype Is a

Stronger Determinant than Sex of the Mouse Gut Microbiota. *Microbial Ecology*, *61*(2), 423–

428. doi: 10.1007/s00248-010-9787-2

Lambert, J. E. (1998). Primate digestion: Interactions among anatomy, physiology, and feeding ecology.

Evolutionary Anthropology: Issues, News, and Reviews, *7*(1), 8–20. doi: 10.1002/(SICI)1520-

6505(1998)7:1<8::AID-EVAN3>3.0.CO;2-C

Lankau, E. W., Hong, P.-Y., & Mackie, R. I. (2012). Ecological drift and local exposures drive enteric

bacterial community differences within species of Galápagos iguanas: GALÁPAGOS IGUANA

ENTERIC COMMUNITY DRIVERS. *Molecular Ecology*, *21*(7), 1779–1788. doi:

10.1111/j.1365-294X.2012.05502.x

Lax, S., Smith, D. P., Hampton-Marcell, J., Owens, S. M., Handley, K. M., Scott, N. M., … Gilbert, J. A.

(2014). Longitudinal analysis of microbial interaction between humans and the indoor

environment. *Science*, *345*(6200), 1048–1052. doi: 10.1126/science.1254529

Leamy, L. J., Kelly, S. A., Nietfeldt, J., Legge, R. M., Ma, F., Hua, K., … Pomp, D. (2014). Host genetics

and diet, but not immunoglobulin A expression, converge to shape compositional features of the

gut microbiome in an advanced intercross population of mice. *Genome Biology*, *15*(12). doi:

10.1186/s13059-014-0552-6

Leclaire, S., Nielsen, J. F., & Drea, C. M. (2014). Bacterial communities in meerkat anal scent secretions

vary with host sex, age, and group membership. *Behavioral Ecology*, *25*(4), 996–1004. doi:

10.1093/beheco/aru074

Ley, R. E., Backhed, F., Turnbaugh, P., Lozupone, C. A., Knight, R. D., & Gordon, J. I. (2005). Obesity

alters gut microbial ecology. *Proceedings of the National Academy of Sciences*, *102*(31), 11070–

11075. doi: 10.1073/pnas.0504978102

Lombardo, M. P. (2008). Access to mutualistic endosymbiotic microbes: an underappreciated benefit of

group living. *Behavioral Ecology and Sociobiology*, *62*(4), 479–497. doi: 10.1007/s00265-007-

0428-9

Magnusson, A., Skaug, H., Nielsen, A., Berg, C., Kristensen, K., Maechler, M., … Brooks, M. (2019).

Generalized Linear Mixed Models using Template Model Builder.

Magoc, T., & Salzberg, S. L. (2011). FLASH: fast length adjustment of short reads to improve genome

assemblies. *Bioinformatics*, *27*(21), 2957–2963. doi: 10.1093/bioinformatics/btr507

Mallott, E. K., Amato, K. R., Garber, P. A., & Malhi, R. S. (2018). Influence of fruit and invertebrate

consumption on the gut microbiota of wild white-faced capuchins (*Cebus capucinus*). *American*

Journal of Physical Anthropology, *165*(3), 576–588. doi: 10.1002/ajpa.23395

Maurice, C. F., Knowles, S. C. L., Ladau, J., Pollard, K. S., Fenton, A., Pedersen, A. B., & Turnbaugh, P.

J. (2015). Marked seasonal variation in the wild mouse gut microbiota. *The ISME Journal*, *9*,

2423–2434. doi: 10.1038/ismej.2015.53

McFall-Ngai, M., Hadfield, M. G., Bosch, T. C. G., Carey, H. V., Domazet-Lošo, T., Douglas, A. E., …

Wernegreen, J. J. (2013). Animals in a bacterial world, a new imperative for the life sciences.

Proceedings of the National Academy of Sciences, *110*(9), 3229–3236. doi:

10.1073/pnas.1218525110

McMurdie, P. J., & Holmes, S. (2014). Waste Not, Want Not: Why Rarefying Microbiome Data Is

Inadmissible. *PLoS Computational Biology*, *10*(4), e1003531. doi: 10.1371/journal.pcbi.1003531

Michl, S. C., Beyer, M., Ratten, J.-M., Hasler, M., LaRoche, J., & Schulz, C. (2019). A diet-change

modulates the previously established bacterial gut community in juvenile brown trout (Salmo

trutta). *Scientific Reports*, *9*(1), 2339. doi: 10.1038/s41598-019-38800-7

Moeller, A. H., Foerster, S., Wilson, M. L., Pusey, A. E., Hahn, B. H., & Ochman, H. (2016). Social

behavior shapes the chimpanzee pan-microbiome. *Science Advances*, *2*(1), e1500997–e1500997.

doi: 10.1126/sciadv.1500997

Münger, E., Montiel-Castro, A. J., Langhans, W., & Pacheco-López, G. (2018). Reciprocal interactions

between gut microbiota and host social behavior. *Frontiers in Integrative Neuroscience*, *12*, 21.

doi: 10.3389/fnint.2018.00021

Oksanen, J., Guillaume Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., … Szoecs, E.

(2017). *vegan: Community Ecology Package (http://CRAN.R-project.org/ package=vegan*

accessed 1 December 2017).

Opstal, E. J. v., & Bordenstein, S. R. (2015). Rethinking heritability of the microbiome. *Science*,

349(6253), 1172–1173. doi: 10.1126/science.aab3958

Orkin, J. D., Campos, F. A., Myers, M. S., Cheves Hernandez, S. E., Guadamuz, A., & Melin, A. D.

(2019). Seasonality of the gut microbiota of free-ranging white-faced capuchins in a tropical dry

forest. *The ISME Journal*, *13*(1), 183–196. doi: 10.1038/s41396-018-0256-0

Orkin, J. D., Webb, S. E., & Melin, A. D. (2019). Small to modest impact of social group on the gut

microbiome of wild Costa Rican capuchins in a seasonal forest. *American Journal of*

Primatology, *0*(0), e22985. doi: 10.1002/ajp.22985

Paradis, E., Claude, J., & Strimmer, K. (2004). APE: analyses of phylogenetics and evolution in R

language. *Bioinformatics*, *20*, 289–290.

Pereira, M. B., Wallroth, M., Jonsson, V., & Kristiansson, E. (2018). Comparison of normalization

methods for the analysis of metagenomic gene abundance data. *BMC Genomics*, *19*(1), 274. doi:

10.1186/s12864-018-4637-6

Perofsky, A. C., Lewis, R. J., Abondano, L. A., Di Fiore, A., & Meyers, L. A. (2017). Hierarchical social

networks shape gut microbial composition in wild Verreaux's sifaka. *Proceedings of the Royal*

Society B: Biological Sciences, *284*(1868), 20172274. doi: 10.1098/rspb.2017.2274

Perofsky, A. C., Lewis, R. J., & Meyers, L. A. (2019). Terrestriality and bacterial transfer: a comparative

study of gut microbiomes in sympatric Malagasy mammals. *The ISME Journal*, *13*(1), 50–63.

doi: 10.1038/s41396-018-0251-5

Phillips, C. D., Phelan, G., Dowd, S. E., McDONOUGH, M. M., Ferguson, A. W., Delton Hanson, J., …

Baker, R. J. (2012). Microbiome analysis among bats describes influences of host phylogeny, life

history, physiology and geography: MICROBIOME ANALYSIS AMONG BATS. *Molecular*

Ecology, *21*(11), 2617–2627. doi: 10.1111/j.1365-294X.2012.05568.x

Pollock, J., Glendinning, L., Wisedchanwet, T., & Watson, M. (2018). The Madness of Microbiome:

Attempting To Find Consensus "Best Practice" for 16S Microbiome Studies. *Applied and*

Environmental Microbiology, *84*(7). doi: 10.1128/AEM.02627-17

R Core Team. (2018). *R: A language and environment for statistical computing (R Foundation for*

Statistical Computing. http://www.R-project.org. Accessed 1 November 2018).

Raulo, A., Ruokolainen, L., Lane, A., Amato, K., Knight, R., Leigh, S., … Tecot, S. R. (2017). Social

behaviour and gut microbiota in red-bellied lemurs (*Eulemur rubriventer*): In search of the role of

immunity in the evolution of sociality. *Journal of Animal Ecology*. doi: 10.1111/1365-

2656.12781

Ren, T., Boutin, S., Humphries, M. M., Dantzer, B., Gorrell, J. C., Coltman, D. W., … Wu, M. (2017).

Seasonal, spatial, and maternal effects on gut microbiome in wild red squirrels. *Microbiome*, *5*(1),

163. doi: 10.1186/s40168-017-0382-3

Robinson, S. P., Simmons, L. W., & Kennington, W. J. (2013). Estimating relatedness and inbreeding

using molecular markers and pedigrees: the effect of demographic history. *Molecular Ecology*,

22(23), 5779–5792. doi: 10.1111/mec.12529

Rothschild, D., Weissbrod, O., Barkan, E., Kurilshikov, A., Korem, T., Zeevi, D., … Segal, E. (2018).

Environment dominates over host genetics in shaping human gut microbiota. *Nature*, *555*(7695),

210–215. doi: 10.1038/nature25973

Saj, T. L., & Sicotte, P. (2007). Predicting the Competitive Regime of Female *Colobus vellerosus* from

the Distribution of Food Resources. *International Journal of Primatology*, *28*(2), 315–336. doi:

10.1007/s10764-007-9124-x

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients. *Methods in*

Ecology and Evolution, *1*(2), 103–113. doi: 10.1111/j.2041-210X.2010.00012.x

Sicotte, P., & MacIntosh, A. J. (2004). Inter-group encounters and male incursions in *Colobus vellerosus*

in central Ghana. *Behaviour*, *141*, 533–553.

Sicotte, P., Teichroeb, J. A., Vayro, J. V., Fox, S. A., Bădescu, I., & Wikberg, E. C. (2017). The influence

of male takeovers on female dispersal in *Colobus vellerosus*: Takeovers and Female Dispersal in

Colobus. *American Journal of Primatology*, *79*(7), e22436. doi: 10.1002/ajp.22436

Smits, S. A., Leach, J., Sonnenburg, E. D., Gonzalez, C. G., Lichtman, J. S., Reid, G., … Sonnenburg, J.

L. (2017). Seasonal cycling in the gut microbiome of the Hadza hunter-gatherers of Tanzania.

Science, *357*(6353), 802–806. doi: 10.1126/science.aan4834

Song, S. J., Lauber, C., Costello, E. K., Lozupone, C. A., Humphrey, G., Berg-Lyons, D., … Knight, R.

(2013). Cohabiting family members share microbiota with one another and with their dogs. *ELife*,

2. doi: 10.7554/eLife.00458

Spor, A., Koren, O., & Ley, R. (2011). Unravelling the effects of the environment and host genotype on

the gut microbiome. *Nature Reviews Microbiology*, *9*(4), 279–290. doi: 10.1038/nrmicro2540

Springer, A., Fichtel, C., Al-Ghalith, G. A., Koch, F., Amato, K. R., Clayton, J. B., … Kappeler, P. M.

(2017). Patterns of seasonality and group membership characterize the gut microbiota in a

longitudinal study of wild Verreaux's sifakas (*Propithecus verreauxi*). *Ecology and Evolution*,

7(15), 5732–5745. doi: 10.1002/ece3.3148

Teichroeb, J. A., Marteinson, S., & Sicotte, P. (2005). Individuals' behaviors following dye-marking in

wild black-and-white colobus (*Colobus vellerosus*). *American Journal of Primatology*, *65*(2),

197–203. doi: 10.1002/ajp.20108

Teichroeb, J. A., Saj, T. L., Paterson, J. D., & Sicotte, P. (2003). Effect of group size on activity budgets

of *Colobus vellerosus* in Ghana. *International Journal of Primatology*, *24*(4), 743–758. doi:

10.1023/A:1024672604524

Teichroeb, J. A., & Sicotte, P. (2009). Test of the ecological-constraints model on ursine colobus

monkeys (*Colobus vellerosus*) in Ghana. *American Journal of Primatology*, *71*(1), 49–59. doi:

10.1002/ajp.20617

Teichroeb, J. A., & Sicotte, P. (2017). Cascading competition: the seasonal strength of scramble

influences between-group contest in a folivorous primate. *Behavioral Ecology and Sociobiology*,

72(1), 6. doi: 10.1007/s00265-017-2418-x

Teichroeb, J. A., Wikberg, E. C., & Sicotte, P. (2009). Female dispersal patterns in six groups of ursine

colobus (*Colobus vellerosus*): infanticide avoidance is important. *Behaviour*, *146*(4), 551–582.

doi: 10.1163/156853909X426363

Teichroeb, J. A., Wikberg, E. C., & Sicotte, P. (2011). Dispersal in male ursine colobus monkeys

(*Colobus vellerosus*): influence of age, rank and contact with other groups on dispersal decisions.

Behaviour, *148*(7), 765–793. doi: 10.1163/000579511X577157

Theis, K. R., Venkataraman, A., Dycus, J. A., Koonter, K. D., Schmitt-Matzen, E. N., Wagner, A. P., …

Schmidt, T. M. (2013). Symbiotic bacteria appear to mediate hyena social odors. *Proceedings of*

the National Academy of Sciences, *110*(49), 19832–19837. doi: 10.1073/pnas.1306477110

Ting, N. (2008). Mitochondrial relationships and divergence dates of the African colobines: evidence of

Miocene origins for the living colobus monkeys. *Journal of Human Evolution*, *55*(2), 312–325.

doi: 10.1016/j.jhevol.2008.02.011

Tung, J., Barreiro, L. B., Burns, M. B., Grenier, J.-C., Lynch, J., Grieneisen, L. E., … Archie, E. A.

(2015). Social networks predict gut microbiome composition in wild baboons. *ELife*, *4*, e05224.

doi: 10.7554/eLife.05224

Turnbaugh, P. J., Ley, R. E., Mahowald, M. A., Magrini, V., Mardis, E. R., & Gordon, J. I. (2006). An

obesity-associated gut microbiome with increased capacity for energy harvest. *Nature*,

444(7122), 1027–1131. doi: 10.1038/nature05414

Vlčková, K., Pafčo, B., Petrželková, K. J., Modrý, D., Todd, A., Yeoman, C. J., … Gomez, A. (2018).

Relationships between gastrointestinal parasite infections and the fecal microbiome in free-

ranging western lowland gorillas. *Frontiers in Microbiology*, *9*(JUN), 1202. doi:

10.3389/fmicb.2018.01202

Voreades, N., Kozil, A., & Weir, T. L. (2014). Diet and the development of the human intestinal

microbiome. *Frontiers in Microbiology*, *5*. doi: 10.3389/fmicb.2014.00494

White, J., Mirleau, P., Danchin, E., Mulard, H., Hatch, S. A., Heeb, P., & Wagner, R. H. (2010). Sexually

transmitted bacteria affect female cloacal assemblages in a wild bird: Sexually transmitted

bacteria in kittiwakes. *Ecology Letters*, *13*(12), 1515–1524. doi: 10.1111/j.1461-

0248.2010.01542.x

Wikberg, E. C., Sicotte, P., Campos, F. A., & Ting, N. (2012). Between-group variation in female

dispersal, kin composition of groups, and proximity patterns in a black-and-white colobus

monkey (*Colobus vellerosus*). *PLoS ONE*, *7*(11), e48740. doi: 10.1371/journal.pone.0048740

Wikberg, E. C., Teichroeb, J. A., Bădescu, I., & Sicotte, P. (2013). Individualistic female dominance

hierarchies with varying strength in a highly folivorous population of black-and-white colobus.

Behaviour, 1–26. doi: 10.1163/1568539X-00003050

Wikberg, E. C., Ting, N., & Sicotte, P. (2014a). Familiarity is more important than phenotypic similarity

in shaping social relationships in a facultative female dispersed primate, *Colobus vellerosus*.

Behavioural Processes, *106*, 27–35. doi: 10.1016/j.beproc.2014.04.002

Wikberg, E. C., Ting, N., & Sicotte, P. (2014b). Kinship and similarity in residency status structure

female social networks in black-and-white colobus monkeys (*Colobus vellerosus*). *American

Journal of Physical Anthropology*, *153*(3), 365–376. doi: 10.1002/ajpa.22435

Wikberg, E. C., Ting, N., & Sicotte, P. (2015). Demographic factors are associated with intergroup

variation in the grooming networks of female colobus (*Colobus vellerosus*). *International Journal

of Primatology*, *36*(1), 124–142. doi: 10.1007/s10764-015-9816-6

Yatsunenko, T., Rey, F. E., Manary, M. J., Trehan, I., Dominguez-Bello, M. G., Contreras, M., …

Gordon, J. I. (2012). Human gut microbiome viewed across age and geography. *Nature*. doi:

10.1038/nature11053

Yuan, M. L., Dean, S. H., Longo, A. V., Rothermel, B. B., Tuberville, T. D., & Zamudio, K. R. (2015).

Kinship, inbreeding and fine-scale spatial structure influence gut microbiota in a hindgut-

fermenting tortoise. *Molecular Ecology*, *24*(10), 2521–2536. doi: 10.1111/mec.13169

Yuan, S., Cohen, D. B., Ravel, J., Abdo, Z., & Forney, L. J. (2012). Evaluation of Methods for the

Extraction and Purification of DNA from the Human Microbiome. *PLOS ONE*, *7*(3), e33865. doi:

10.1371/journal.pone.0033865

Data Accessibility Statement

All raw data are stored in the PaceLab database hosted by the University of Calgary. The 16S sequencing

data will be uploaded to NCBI's Short Read Archive. The data used for the analyses presented here will

be uploaded to Dryad.

Appendix

Microbial taxa predicted to be structured by social connectedness

The last aim of this study was to investigate whether social connectedness was correlated with

Operational Taxonomic Unit (OTU) abundances in certain taxa (Table A1) previously reported as

structured by social relationships (Amato et al., 2017; Tung et al., 2015). We also expected social

connectedness to be correlated with the abundances of three gut microbial taxa that diverged between

two daughter groups after a group fission (DA and NP in Fig. A1), because we suspect that this pattern

was driven by social network changes (Goodfellow et al., 2019).

747
748 **Figure A1.** Home ranges and core areas for groups in the main forest fragment at Boabeng-Fiema,

749 Ghana. Solid lines indicate home ranges of groups from which we collected behavioural data. Dashed

750 lines indicate partial home ranges from other groups present in this forest.

Taxon	Socially structured in 2007	Socially structured in 2009	Reference
Actinobacteria:	X	X	Tung et al. 2015
Bifidobacteriaceae:	NA	X	Tung et al. 2015
Bifidobacterium	NA	X	Tung et al. 2015
Coriobacteriaceae	X	X	Tung et al. 2015
Bacteroidetes:	-	-	-
Bacteroidaceae:	X	(✓)	-
Bacteroides	X	✓	Amato et al. 2017
Porphyromonadaceae:	NA	X	Goodfellow et al. 2019
Parabacteroides	NA	✓	Goodfellow et al. 2019
Firmicutes:	✓	(✓)	-
Clostridiaceae	-	-	-
Clostridium	X	✓	Amato et al. 2017
Eubacteriaceae	-	-	-
Eubacterium	X	(✓)	-
Lachnospiraceae:	✓	(✓)	-
Coprococcus	X	✓	Goodfellow et al. 2019
Lachnospiracea incertae sedis	X	(✓)	-
Roseburia	✓	X	Amato et al. 2017
Peptococcaceae:	X	(✓)	-
Desulfurispora	X	(✓)	-
Ruminococcaceae	X	(✓)	-
Flavonifractor	X	(✓)	-
Subdoligranulum	NA	(✓)	-
Streptococcaceae:	NA	-	-
Streptococcus	NA	NA	Amato et al. 2017
Veillonellaceae:	X	X	Tung et al. 2015
Propionispira	X	(✓)	-
Centipeda	X	(✓)	-
Fuscobacteria	NA	X	Tung et al. 2015
Fusobacteriaceae	NA	X	Tung et al. 2015
Fusobacterium	NA	X	Tung et al. 2015
Proteobacteria:	-	-	-
Desulfovibrionaceae	-	-	-
Desulfovibrio	X	(✓)	-
Enterobacteriaceae	X	X	Tung et al. 2015
Tenericutes:	X	X	Tung et al. 2015
Anaeroplasmataceae	-	-	-
Asteroleplasma	X	(✓)	-
Mycoplasmataceae	NA	X	Tung et al. 2015
Mycoplasma	NA	X	Tung et al. 2015

Table A1. OTUs in these phyla, families, and genera are expected to be structured by sociality based on

previous studies. Predictions were supported ✓; not supported X; no prediction made but structured in

our data set (✓); or no prediction made and not structured in our data set (-). Grey text indicates rare

taxa (N < 3 OTUs). NA denotes taxa not present in our data set.

DNA extraction, amplification, and sequencing protocols for the gut microbiome analysis

We extracted DNA from 200 μl of sample using QIAmp DNA stool extraction protocol the following

modifications. Step 2: Added 50 μl Proteinase K with overnight lysis before proceeding to step 3. Step 4:

Pipetted all of the supernatant. Step 5: Used half of the InhibitEX tablet. Step 6: Centrifuged for 5

minutes. Step 9: Added 4ul RNAse and vortexed for 15 seconds. Step 19: Used 50 μl Buffer AE and

incubated at 10 minutes. Step 20: Pipetted the same 50 μl of buffer AE back onto filter and incubated at

room temperature for 15 minutes. Centrifuged at full speed for 2 minutes. Our DNA extraction protocol

did not include a bead-beating step, which could bias against lysis-resistant taxa such as Gram-positive

and spore-forming bacteria that are less likely to be dependent on direct social contact for transmission

between hosts because they can survive for prolonged periods outside the host (Pollock, Glendinning,

Wisedchanwet, & Watson, 2018; Yuan, Cohen, Ravel, Abdo, & Forney, 2012).

We determined the concentration of the extracts using Qubit dsDNA BR Assay Kit (Invitrogen)

and diluted products to 2nM for downstream reactions. We amplified the bacterial v4 region of the 16S

ribosomal RNA gene using the following 515F and 806R primers containing 5' Illumina adapter tails and

dual indexing barcodes:

515F 5' AATGATACGGCGACCACCGAGATCTACACTAGATCGCTATGGTAATTGTGTGCCAGCMGCCGCGGTAA

806R 5' CAAGCAGAAGACGGCATACGAGATTCACCTAGAGTCAGTCAGCCGGACTACHVGGGTWTCTAAT.

We set the PCRs with 12.5 μl NEB Q5 Hot start 2x Master mix, 1.25 μl 10uM Primer mix, 1 μl template

DNA, and 10.25 μl MoBio certified DNA free water and used the following cycling protocol: 98 degrees

for 30 seconds (1x) followed by 98 degrees for 10 seconds, 61 degrees for 20 seconds, and 72 degrees

for 20 seconds (20x), followed by 72 degrees for 2 minutes and 4 degrees. The amplification products

were cleaned up using Ampure XP beads and normalized into a final pool with an Eppendorf liquid

handling robot. Libraries were sequenced as part of a 150bp paired-end sequencing run on the Illumina

NextSeq platform following the manufacturer's protocol.

We used a custom pipeline that contained the following steps: joining pair-end reads; removing

low-quality and chimeric reads; dereplication and dropping unique reads with low abundance; clustering

OTUs; making OTU table; alignment; building a reference tree; and taxon assignment using FLASH

(Magoc & Salzberg, 2011), the FASTX Toolkit (Hannon Lab, 2010), and the USEARCH pipeline (Edgar,

2010). See https://github.com/kstagaman/Process_16S and Goodfellow et al. (2019) for further details.

We performed *de novo* OTU picking in UCLUST (Edgar, 2010), and sequences with 97% overlap were

defined as belonging to the same bacterial Operational Taxonomic Unit (OTU). To guard against

sequencing errors, we filtered out OTU's with a frequency lower than 0.00005 as recommended

(Bokulich et al., 2012).

792
Variation in predictor and outcome variables

Of the females included in the analyses with behavioural predictor variables (Table A2), dietary

dissimilarity (i.e., Sørensen diversity index) varied from 0 to 1, dissimilarity in relatedness calculated as

their *R*-value subtracted from 1 ranged from 0.31 to 1, and social connectedness (i.e., inversed path

length or Geodesic distance in the 1-meter proximity network) varied from 0 to 1 where 0 represents

unconnected dyads (Fig. A2).

In our full data set, mean unweighted Unifrac distances within the same season and year was

0.052 ± 0.004 for samples collected from the same individual (N = 4 samples) and 0.205 ± 0.042 for

samples collected from different individuals within the same season and year (N = 61 samples). The low

amount of within-individual variation in comparison to the between-individual variation suggests that

one sample per individual is representative of its gut microbiome during that season and sufficient for

analysis of beta-diversity. Furthermore, the beta-diversity of matched samples from the same adult

female in the wet season 2007 and the dry season 2009 (N = 22 samples from 11 females) was lower

(0.152 ± 0.031) than the female's mean beta-diversity with samples from a different female and year

(0.183 ± 0.021) for all but one female, and there was a significant difference in beta-diversity between

samples from the same versus different females in this sample (N = 11 females, Wilcoxon signed rank

test, p < 0.001).

Year	Group	AF group size	AF sampled	AF omitted	Reason for omitting samples
2007	BS	4	4	0	
	DA	5	5	1	Incomplete dietary information
	NP	4	4	0	
	RT	6	5	0	
	SP	4	4	0	
	WW	9	9	0	
2009	BO	8	8	0	
	BS	6	1	1	Lacked samples from majority of AF
	DA	7	2	2	Lacked samples from majority of AF
	NP	5	3	3	Lacked samples from majority of AF
	OD	6	6	0	
	RT	7	2	2	Lacked samples from majority of AF
	SP	3	0	-	
	WW	7	5	0	

811
812   **Table A2.** Number of adult females (AF) present, sampled, and omitted from data analyses with

813   behavioural predictor variables.

816   **Figure A2.** Dietary dissimilarity, relatedness dissimilarity, and social connectedness were lower for

817   within-group than between-group female-female dyads (Wilcoxon signed rank tests, N = 88 samples, all

818   p < 0.001).