A modification to the life cycle of the parasite

Trypanosoma brucei

Sarah Schuster[#], Ines Subota[#], Jaime Lisack, Henriette Zimmermann, Christian

Reuter, Brooke Morriswood and Markus Engstler*

* these authors contributed equally

[#] corresponding author: markus.engstler@biozentrum.uni-wuerzburg.de

Abstract

African trypanosomes cause sleeping sickness in humans and nagana in cattle. These

unicellular parasites are transmitted by the blood-sucking tsetse fly[1]. In the mammalian

host's circulation, tissues, and interstitium, at least two main life cycle stages exist:

slender and stumpy bloodstream forms[2,3,4,5,6]. Proliferating slender forms differentiate

into cell cycle-arrested stumpy forms at high levels of parasitaemia. This

developmental stage transition occurs in response to the quorum sensing factor SIF

(stumpy induction factor)[7], and is thought to fulfil two main functions. First, it auto-

regulates the parasite load in the host. Second, the stumpy stage is regarded as pre-

adapted for tsetse fly infection and the only form capable of successful vector

transmission[8]. Differentiation to the stumpy form is accompanied by fundamental

morphological and metabolic changes, including expression of the stumpy marker,

protein associated with differentiation 1 (PAD1)[9]. Here, we show that proliferating

slender stage trypanosomes are equally able to infect the tsetse fly, and that a single

parasite is sufficient. The slender parasites complete the complex life cycle in the fly

with comparable overall success rates and kinetics as stumpy forms. We further show

that in the tsetse midgut, the slender parasites activate the canonical PAD1 pathway,

without undergoing either cell cycle arrest or a morphological transition to the stumpy

form. Instead, with the onset of PAD1 expression, the parasites directly differentiate

into the procyclic (insect) stage. Our findings not only propose a revision to the

traditional view of the trypanosome life cycle, but also suggest a solution to a long-

acknowledged paradox in the transmission event: parasitaemia in chronic infections is

typically quite low, and so the probability of a tsetse ingesting a stumpy cell during a

bloodmeal is also low[10,11,12,13].

Results and Discussion

Slender and stumpy bloodstream form trypanosomes can be distinguished based on

cell cycle, morphological, and metabolic criteria. The genome of the single

mitochondrion (kinetoplast, K) and the cell nucleus (N) can be readily visualised using

DNA stains, and their prescribed sequence of replication (1K1N, 2K1N, 2K2N) allows

cell cycle stage to be inferred[14]. Slender cells are found in all three K/N ratios, while

stumpy cells, which are cell cycle-arrested, are found only as 1K1N cells (Fig. 1A).

Expression of _protein associated with differentiation 1_ (PAD1) is accepted as a marker

for development to the stumpy stage[9]. Cells expressing an NLS-GFP reporter fused

to the 3' UTR of the PAD1 gene (GFP:PAD1$_{UTR}$) will have GFP-positive nuclei when

the PAD1 gene is active. Hence, slender cells are GFP-negative; stumpy cells are

GFP-positive (Fig. 1A). We have previously shown that stumpy cells can be formed

independently of high cell population density by ectopic expression of a second variant

surface glycoprotein (VSG) isoform, a process that mimics one of the pathways

involved in trypanosome antigenic variation[15,16,17,18]. These so-called expression site

(ES)-attenuated stumpy cells can complete the developmental cycle in the tsetse fly[18].

It remained an open question whether this occurred with the same efficiency as with

SIF-produced stumpy cells. To address this question, we quantitatively compared the

transmission efficiency of stumpy populations generated by either SIF-treatment or

through ES-attenuation. Tsetse flies (*Glossina morsitans morsitans*) were infected

during membrane feeding (Fig. 1B; Supplementary Video 1) with defined numbers of

pleomorphic trypanosomes, which are capable of completing the entire developmental

cycle. Two transgenic trypanosome cell lines, both of which contained the

GFP:PAD1$_{UTR}$ reporter construct, were used. One was subjected to ectopic VSG

expression to drive ES attenuation (line ES;[18]). The other was treated with stumpy

induction factor (line SIF). Both treatments resulted in expression of the

GFP:PAD1$_{UTR}$ reporter and synchronous differentiation to the stumpy stage. The

resulting stumpy populations were fed to tsetse flies at concentrations ranging from

120,000 to 10 cells/ml. A feeding tsetse typically ingests 20 µl of blood[19], meaning that

between 2,400 and 0.2 trypanosomes were ingested per bloodmeal on average (Fig.

1C, rows i-vi, column 2, Total). The trypanosomes had been previously scored for

expression of the GFP:PAD1$_{UTR}$ reporter to confirm their identity as stumpy forms (Fig.

1C, columns 3-4). To analyse the infections, we carried out microscopic analyses of

dissected tsetse digestive tracts (Fig. 1D, E). The presence of mammal-infective,

metacyclic trypanosomes in explanted tsetse salivary glands indicated completion of

the tsetse transmission cycle (Fig. 1F). The uptake, on average, of two stumpy

parasites of either cell line produced robust infections of tsetse midgut (MG),

proventriculus (PV), and salivary glands (SG) (Fig. 1C, columns 5-7). Ingestion, on

average, of even a single stumpy cell was sufficient to produce salivary gland

infections in almost 5% of all tsetse (Fig. 1C, row v). When the stumpy parasite number

was further reduced to 0.2 cells on average per bloodmeal, 0.9% of flies still acquired

salivary gland infections (Fig. 1C, row vi). As a measure of the incidence of life cycle

completion in the tsetse fly, we calculated the transmission index (TI; ratio of salivary

gland to midgut infections) for each condition[20]. We found that for flies infected with 2

trypanosomes on average, the TI was comparable between SIF-induced (TI = 0.29)

and ES-induced (TI = 0.31) stumpy trypanosomes (Fig. 1C, rows iii-iv). A similar TI of

0.23 was observed in flies ingesting on average 1 trypanosome (Fig. 1C, row v). Thus,

our data clearly show that a single stumpy cell is sufficient to infect a tsetse fly. As a

control, infections were also carried out using a monomorphic trypanosome strain

(Supplementary Table 1A). Monomorphic trypanosomes are able to infect the tsetse

midgut, but are incapable of completing the developmental cycle in the fly[21,22]. As

expected, no salivary gland infections were seen using these cells. Next, we did an

experiment that was originally intended as an additional *negative* control: we infected

tsetse flies with proliferating PAD1-negative slender trypanosomes from the two

pleomorphic cell lines used (Fig. 1C, rows vii-xi). We expected these cells not to

passage through the fly. Remarkably, however, there was almost no difference in the

infection efficiency when the flies were fed with either 20 stumpy trypanosomes or 20

pleomorphic slender trypanosomes (Fig. 1C, compare TI for rows ii and vii). When flies

were fed with on average 2 slender parasites each, the TI was actually higher for

slender cells (0.60) than for stumpy cells (0.31) (Fig. 1C, compare TI for rows viii and

iii). This TI of 0.60 was identical for both populations of slender cells (Fig. 1C, rows viii-

ix). Next, when given on average just one PAD1-negative slender cell per bloodmeal,

parasite infections still made it through the midgut, proventriculus, and salivary glands

with incidences of 4.9%, 4.3%, and 2.1% respectively, at a TI of 0.44 (Fig. 1C, row x).

In order to be absolutely sure that slender trypanosomes can passage through the

tsetse, we repeated the experiment with naïve slender parasites that had been freshly

differentiated from insect-derived metacyclic trypanosomes, i.e. cells that had just

restarted the mammalian stage life cycle (Supplementary Table 1, row iii). Infections

with on average two freshly-differentiated slender trypanosomes per bloodmeal

revealed 6.9% midgut and 2.5% salivary gland infections. The transmission index was

0.36.

It is important to note however that while ES-attenuated cells showed similar midgut,

proventriculus, and salivary gland infection incidence as either stumpy or slender

forms (Fig. 1C, rows ii-iii and vii-viii), the SIF-induced stumpy cells were better at

establishing infections than their slender counterparts (Fig. 1C, rows iv-vi and ix-xi).

Infections with 1-2 slender cells, however, produced higher TI values than those with

the same numbers of stumpy cells (Fig. 1G). This suggests that the proliferative

slender cells are more capable of progressing from a midgut infection to a salivary

gland one, and thus have at least comparable developmental competence to the

stumpy forms. In summary, our experiments not only establish that single

trypanosomes (either slender or stumpy) can infect the tsetse fly, but also strongly

suggest that slender cells can efficiently complete the passage through the tsetse fly.

To determine how slender trypanosomes manage to establish infections, we observed

the early events following trypanosome ingestion by tsetse flies (Supplementary Video

2). The canonical version of events is that ingested stumpy (i.e. PAD1-positive) cells

reactivate the cell cycle, begin to express the EP procyclin protein on their cell surface,

and differentiate to the procyclic life cycle stage. We infected tsetse with pleomorphic

trypanosomes, which not only contained the stumpy-specific GFP:PAD1$_{UTR}$ marker,

but also encoded an EP1:YFP fusion[23]. In this way, the onset of stumpy development

was observable as GFP fluorescence in the nucleus, and further differentiation to the

procyclic life cycle stage as YFP fluorescence on the parasite cell surface. In addition,

the cell cycle status (K/N counts, see Fig. 1A), morphology, and the characteristic

motile behaviour of the trypanosomes were also assessed as criteria of developmental

progress. In total, 114 tsetse flies (57 male and 57 female) were dissected after at

least six independent infections with either 12,000 slender or stumpy parasites each

(Fig. 2). These high initial parasite numbers allowed the microscopic analysis of

individual living slender (n = 1845) and stumpy trypanosomes (n = 1237) within the

convoluted microenvironment of midgut explants[24]. As early as 2-4 h post-infection

with slender trypanosomes, a few (0.8%) dividing trypanosomes with a nuclear PAD1

signal could be observed (Fig. 2A). After 8-10 hours however, half (50.4%) of all

trypanosomes in the explants were PAD1-positive. After 24 hours, 84.3% of the

parasites expressed PAD1, and 9.8% had already initiated developmental progression

to the procyclic insect stage, as evidenced by EP1:YFP fluorescence on their cell

surface. At 48-50 h post-infection with slender trypanosomes, virtually the entire

trypanosome population (91.8%) expressed PAD1, and almost one fifth (19.1%) was

EP1-positive. To examine cell cycle progression, we counted the number of 1K1N,

2K1N, and 2K2N cells in the PAD1-positive and PAD1-negative slender cell

populations (Fig. 2B, slender). Remarkably, 8-10 h post-infection, replicating (i.e.

2K1N, 2K2N) cells that were PAD1-positive could be readily observed. Over the

duration of the experiment, PAD1-negative cells gradually decreased in numbers,

while PAD1-positive slender cells at all cell cycle stages were increasingly observed

(Supplementary Video 2C). Thus, the PAD1 pathway is triggered in the fly, but

apparently without pushing the slender parasites towards cell cycle arrest. Of note,

EP1 expression did not exactly correlate with acquisition of procyclic morphology. At

24-26h, only 9.8% of slender cells are EP1-positive (Fig. 2A), but the EP1-negative

cells frequently exhibited procyclic morphology (Fig. 2C, upper panels). An example

of a dividing (2K2N), PAD1-positive, EP1-positive cell is also shown (Fig. 2C, lower

panels; Supplementary Video 2D). Thus, it appears that a seamless developmental

stage transition from slender bloodstream forms to the procyclic insect forms takes

place, which is accompanied by the typical reorganisation of the cytoskeleton and the

concomitant switch of swimming styles[25,24].

In order to directly compare the kinetics of slender-to-procyclic development with that

of stumpy stage trypanosomes, we fed flies with SIF-induced, PAD1-positive stumpy

trypanosomes (Fig. 2B, stumpy). These cells remained in cell cycle arrest for the first

day, and re-entered the cell cycle as procyclic parasites after day 2-3. Stumpy

trypanosomes showed a higher incidence of EP1:YFP expression than slender cells

at all timepoints (Fig. 2A, grey bars). The procyclic marker EP1:YFP was already

visible on the cell surface of 16.2 % of trypanosomes after 10 hours, showing that EP

expression was initiated before release of cell cycle arrest. Uncoupling of EP surface

expression from the commitment to differentiation has been reported before[23].

We further investigated the developmental potential of pleomorphic slender

bloodstream forms *in vitro* using the same cell lines and analysis as above.

Differentiation to the procyclic insect stage was induced by the addition of *cis*-aconitate

and a temperature drop from 37°C to 27°C[26,27,28,23] (Fig. 3). Slender trypanosomes

activated the PAD1 pathway immediately, with 9.8% of all parasites being PAD1-

positive within 2-4 hours, and 83.2% after 10 hours. PAD1 expression peaked after

one day (98.3%), and declined thereafter (Fig. 3A). Simultaneously with PAD1 reporter

expression, EP1 appeared on the cell surface of 19.6% of all parasites within 8-10

hours, increasing to 98.3% after 3 days (Fig. 3A). Throughout the timecourse, PAD1-

positive 2K1N and 2K2N cells were consistently observed, demonstrating that the

PAD1-positive slender parasites did not arrest in the cell cycle, and continued dividing

throughout *in vitro* differentiation to the procyclic stage (Fig. 3B). After 3 days of *cis*-

aconitate treatment *in vitro*, slender trypanosomes had established a proliferating

procyclic parasite population.

By comparison, stumpy parasites (Fig. 3A, grey bars) responded to *in vitro cis-*

aconitate treatment with rapid expression of the EP1:YFP marker, with 28.6% of all

cells being positive within 2-4 hours. After one day, EP1 was present on almost all

(96.7%) stumpy trypanosomes. The cell cycle analysis revealed that the parasites

were not dividing, however (Fig. 3B). The first cells re-entered the cell cycle only after

15-17 hours, and a normal procyclic cell cycle profile was not reached until 3 days.

Thus, the *in vitro* differentiation supported the *in vivo* observations, demonstrating that

pleomorphic slender trypanosomes are able to directly differentiate to the procyclic

stage without becoming cell cycle-arrested stumpy cells. Furthermore, the overall

developmental capacity and differentiation kinetics of both life cycle stages are

comparable, *in vitro* and *in vivo*.

In conclusion, our observations suggest a revised view on the life cycle of African

trypanosomes (Fig. 4). One trypanosome suffices to produce robust infections of the

vector, and the stumpy stage is not essential for tsetse transmission. Slender parasites

can complete the complex life cycle in the fly with comparable overall success rates

and kinetics as the stumpy forms. The stumpy stage appears more able to establish

initial infections in the fly midgut, however (Fig. 1C, column 5, MG). This may be

related to a greater resistance against the digestive environment in the fly's gut, as

has been discussed[29,30]. On the other hand, stumpy trypanosomes are not replicative,

and their lifetime is limited to roughly 3 days[31]. In the fly, re-entry into the cell cycle is

by no means immediate, but takes at least two days. Thus, stumpy cells may run into

an age-related problem. Conversely, slender-derived parasites appear to travel more

efficiently to their final destination within the tsetse, the salivary glands.   The

significantly higher mobility of slender forms when compared to stumpy

trypanosomes[32], and the seamless differentiation to the motile procyclic stage could

be involved in this migratory success. Along similar lines, it is worth noting that

Trypanosoma congolense, the principal causative agent of the cattle plague nagana,

infects tsetse flies without manifesting a cell cycle-arrested stumpy life cycle stage[33].

This raises the question of what the true biological function of the stumpy life cycle

stage actually is.

Methods

Trypanosome culture

Pleomorphic *Trypanosoma brucei brucei* strain EATRO 1125 (serodome AnTat1.1)[34]

bloodstream forms were grown in HMI-9 medium[35], supplemented with 10% (v/v) fetal

bovine serum and 1.1% (w/v) methylcellulose (Sigma 94378)[36] at 37°C and 5% CO_2.

Slender stage parasites were maintained at a maximum cell density of $5x10^5$ cells/ml.

For cell density-triggered differentiation to the stumpy stage, cultures seeded at $5x10^5$

cells/ml were cultivated for 48 hours without dilution. Pleomorphic parasites were

harvested from the viscous medium by 1:4 dilution with trypanosome dilution buffer

(TDB; 5 mM KCl, 80 mM NaCl, 1 mM $MgSO_4$, 20 mM Na_2HPO_4, 2 mM NaH_2PO_4, 20

mM glucose, pH 7.6), followed by filtration (MN 615 ¼, Macherey-Nagel, Germany)

and centrifugation (1,400x*g*, 10 min, 37°C)[18]. Monomorphic *T. brucei* 427 MITat 1.2

13-90 bloodstream forms[37] were grown in HMI-9 medium[35], supplemented with 10%

(v/v) fetal bovine serum at 37°C and 5% CO_2.

For *in vitro* differentiation to the procyclic insect stage, bloodstream stage

trypanosomes were pooled to a cell density of $2x10^6$ cells/ml in DTM medium

immediately before use[38]. *Cis*-aconitate was added to a final concentration of 6

mM[26,38] and temperature was adjusted to 27°C. Procyclic parasites were grown in

SDM79 medium[39], supplemented with 10% (v/v) fetal bovine serum[35] and 20 mM

glycerol[40,24].

Genetic manipulation of trypanosomes

Transfection of pleomorphic trypanosomes was done as previously described[18], using

an AMAXA Nucleofector II (Lonza, Switzerland). Transgenic trypanosome clones were

selected by limiting dilution in the presence of the appropriate antibiotic. The

GFP:PAD1$_{UTR}$ reporter construct[18] was used to transfect AnTat1.1 trypanosomes to

yield the cell line 'SIF'. The trypanosome 'ES' line was described previously[18]. It

contains the reporter GFP:PAD1$_{UTR}$ construct and an ectopic copy of VSG gene MITat

1.6 under the control of a tetracycline-inducible T7-expression system. The EP1:YFP

construct was integrated into the EP1-procyclin locus as described previously[23].

Tsetse maintenance

238 The tsetse fly colony (*Glossina morsitans morsitans*) was maintained at 27°C and 70%

239 humidity. Flies were kept in Roubaud cages and fed 3-times a week through a silicon

240 membrane, with pre-warmed, defibrinated, sterile sheep blood (Acila, Germany).

Fly infection and dissection

Teneral flies were infected 1-3 days post-eclosion during their first meal. Depending

on the experiment, trypanosomes were diluted in either TDB or sheep blood. The

infective meals were supplemented with 60 mM N-acetylglucosamine[41]. For infection

with monomorphic parasites, cells were additionally treated for 48 hours with 12.5 mM

glutathione (GSH)[42] and 100 µM 8-pCPT-cAMP (cAMP)[7].

Tsetse infection status was analysed between 35 and 40 days post-infection. Flies

were euthanized with chloroform and dissected in PBS. Intact tsetse alimentary tracts

were explanted and analysed microscopically, as described previously (Schuster,

2017). For the analysis of early trypanosome differentiation *in vivo*, slender or stumpy

trypanosomes at a concentration of $6x10^5$ cells/ml were resuspended in TDB to the

required final concentration and fed to flies. The numbers of flies used and the number

of independent experiments carried out are indicated in the figure legends. Results

are presented as sample means.

Fluorescence microscopy and video acquisition

Live trypanosome imaging was performed with a fully automated DMI6000B wide field

fluorescence microscope (Leica microsystems, Germany), equipped with a

DFC365FX camera (pixel size 6.45 µm) and a 100x oil objective (NA 1.4). For

high- speed imaging, the microscope was additionally equipped with a pco.edge

sCMOS camera (PCO, Germany; pixel size 6.5 µm). Fluorescence video acquisition

was performed at frame rates of 250 fps. For visualisation of parasite cell cycle and

morphology, slender and stumpy trypanosomes were harvested and incubated with 1

mM AMCA-sulfo-NHS (Thermo Fisher Scientific, Germany) for 10 minutes on ice.

Cells were chemically fixed in 4% (w/v) formaldehyde and 0.05% (v/v) glutaraldehyde

overnight at 4°C. DNA was visualised with 1 µg/ml DAPI immediately before analysis.

3D-Imaging was done with a fully automated iMIC wide field fluorescence microscope

(FEI-TILL Photonics, Germany), equipped with a Sensicam qe CCD camera (PCO,

Germany; pixel size 6.45 µm) and a 100x oil objective (NA 1.4). Deconvolution of

image stacks was performed with the Huygens Essential software (Scientific Volume

Imaging B.V., Netherlands). Fluorescence images are shown as maximum intensity

projections of 3D-stacks in false colours with green fluorescence in green and blue

fluorescence in grey.

Scanning electron microscopy

Explanted tsetse alimentary tracts were fixed in Karnovsky solution (2% formaldehyde,

2.5% glutaraldehyde in 0.1M cacodylate buffer, pH 7.4) and incubated overnight at

4°C. Samples were washed 3-times for 5 minutes at 4°C with 0.1M cacodylate buffer,

pH 7.4, followed by incubation for 1 hour at 4°C in post-fixation solution (2.5%

glutaraldehyde in 0.1M cacodylate buffer, pH 7.4). After additional washing, the

samples were incubated for 1 hour at 4°C in 2% tannic acid in cacodylate buffer, pH

7.4, 4.2% sucrose, and washed again in water (3x for 5 minutes, 4°C). Finally, serial

dehydration in acetone was performed, followed by critical point drying and platinum

coating. Scanning electron microscopy was done using the JEOL JSM-7500F field

emission scanning electron microscope.

Data Availability

All datasets generated during this project are provided as online source data. The cell

lines used are available from the corresponding author on request.

Acknowledgements

We thank Nicola Jones, Susanne Kramer, Manfred Alsheimer, Christian Janzen and

Ricardo Benavente for discussion and critical reading of the manuscript. ME is

supported by DFG grants EN305, SPP1726 (Microswimmers – From Single Particle

Motion to Collective Behaviour), and GRK2157 (3D Tissue Models to Study Microbial

Infections by Obligate Human Pathogens). ME is member of the Wilhelm Conrad

Roentgen Center for Complex Material Systems (RCCM).

Author contributions

S.S. designed the experiments, performed the experiments, analysed the data, interpreted the results and wrote the manuscript. I.S. designed the experiments, performed the experiments, analysed the data and interpreted the results. J.L. designed the experiments, performed the experiments, analysed the data, interpreted the results and wrote the manuscript. H.Z., designed the experiments, performed the experiments, analysed the data and interpreted the results. C.R. designed the experiments, performed the experiments, analysed the data and interpreted the results. B.M. interpreted the results and wrote the manuscript. M.E. conceived the study, designed the experiments, analysed the data, interpreted the results and wrote the manuscript.

Competing interests

The authors declare no competing interests.

References

1. Bruce, D. Preliminary Report on Tsetse Fly Disease or Nagana in Zululand, Bennet & Davis, Durban (1895).

2. Robertson, M. Notes on the Polymorphism of *Trypanosoma gambiense* in the Blood and Its Relation to the Exogenous Cycle in *Glossina palpalis*. *Proceedings of the Royal Society B: Biological Sciences* **85**, 527–539 (1912).

3. Goodwin, L. G. The pathology of African Trypanosomiasis. *Trans R Soc Trop Med Hyg* **64**, 797–812 (1970).

4. Trindade, S. *et al. Trypanosoma brucei* Parasites Occupy and Functionally

Adapt to the Adipose Tissue in Mice. *Cell Host & Microbe* **19**, 837–848 (2016).

5. Capewell, P. *et al.* The skin is a significant but overlooked anatomical reservoir

for vector-borne African trypanosomes. *eLife* **5**, e17716 (2016).

6. Krüger, T., Schuster, S. & Engstler, M. Beyond Blood: African Trypanosomes

on the Move. *Trends in Parasitology* **34**, 1056–1067 (2018).

7. Vassella, E., Reuner, B., Yutzy, B. & Boshart, M. Differentiation of African

trypanosomes is controlled by a density sensing mechanism which signals cell

cycle arrest via the cAMP pathway. *J. Cell. Sci.* **110**, 2661–2671 (1997).

8. Rico, E. *et al.* Bloodstream form pre-adaptation to the tsetse fly in *Trypanosoma*

brucei. *Front. Cell. Infect. Microbiol.* **3:78** (2013).

9. Dean, S., Marchetti, R., Kirk, K. & Matthews, K. R. A surface transporter family

conveys the trypanosome differentiation signal. *Nature* **459**, 213–217 (2009).

10. Koch, R., Beck, M., Kleine, F. K. *Bericht über die Tätigkeit der zur Erforschung*

der Schlafkrankheit im Jahre 1906/07 nach Ostafrika entsandten Kommission.

Arbeiten aus dem Kaiserlichen Gesundheitsamt, Band **31**. Berlin : J. Springer

(1909).

11. Frezil, J. L. Application of xenodiagnosis in the detection of *T. gambiense*

trypanosomiasis in immunologically suspect patients. *Bull Soc Pathol Exot*

Filiales **64**, 871–878 (1971).

12. Wombou Toukam, C. M., Solano, P., Bengaly, Z., Jamonneau, V. & Bucheton,

B. Experimental evaluation of xenodiagnosis to detect trypanosomes at low

parasitaemia levels in infected hosts. *Parasite* **18**, 295–302 (2011).

13. Capewell, P. *et al.* Resolving the apparent transmission paradox of African

sleeping sickness. *PLOS Biology* **17**, e3000105 (2019).

14. Sherwin, T. & Gull, K. The cell division cycle of *Trypanosoma brucei brucei*:

timing of event markers and cytoskeletal modulations. *Philos. Trans. R. Soc.*

Lond., B, Biol. Sci. **323**, 573–588 (1989).

15. Cross, G. A. Identification, purification and properties of clone-specific

glycoprotein antigens constituting the surface coat of *Trypanosoma brucei*.

Parasitology **71**, 393–417 (1975).

16. Hertz-Fowler, C. *et al.* Telomeric Expression Sites Are Highly Conserved in

Trypanosoma brucei. *PLoS ONE* **3**, e3527 (2008).

17. Batram, C., Jones, N. G., Janzen, C. J., Markert, S. M. & Engstler, M.

Expression site attenuation mechanistically links antigenic variation and

development in *Trypanosoma brucei*. *Elife* **3**, e02324 (2014).

18. Zimmermann, H. *et al.* A quorum sensing-independent path to stumpy

development in *Trypanosoma brucei*. *PLOS Pathogens* **13**, e1006324 (2017).

19. Gibson, W. & Bailey, M. The development of *Trypanosoma brucei* within the

tsetse fly midgut observed using green fluorescent trypanosomes. *Kinetoplastid*

Biol Dis **2**, 1 (2003).

20. Peacock, L., Ferris, V., Bailey, M. & Gibson, W. The influence of sex and fly

species on the development of trypanosomes in tsetse flies. *PLoS Negl Trop*

Dis **6**, e1515 (2012).

21. Herder, S. *et al. Trypanosoma brucei* 29-13 strain is inducible in but not

permissive for the tsetse fly vector. *Exp. Parasitol.* **117**, 111–114 (2007).

22. Peacock, L., Ferris, V., Bailey, M. & Gibson, W. Fly transmission and mating

of *Trypanosoma brucei brucei* strain 427. *Mol. Biochem. Parasitol.* **160**, 100–

106 (2008).

23. Engstler, M. & Boshart, M. Cold shock and regulation of surface protein

trafficking convey sensitization to inducers of stage differentiation in

Trypanosoma brucei. *Genes Dev* **18**, 2798–2811 (2004).

24. Schuster, S. *et al.* Developmental adaptations of trypanosome motility to the

tsetse fly host environments unravel a multifaceted *in vivo* microswimmer

system. *eLife* **6**, e27656 (2017).

25. Heddergott, N. *et al.* Trypanosome Motion Represents an Adaptation to the

Crowded Environment of the Vertebrate Bloodstream. *PLoS Pathogens* **8**,

e1003023 (2012).

26. Brun, R., Jenni, L., Schönenberger, M. & Schell, K. F. *In vitro* cultivation of

bloodstream forms of *Trypanosoma brucei*, *T. rhodesiense*, and *T. gambiense*.

J. Protozool. **28**, 470–479 (1981).

27. Czichos, J., Nonnengaesser, C. & Overath, P. *Trypanosoma brucei: cis-*

Aconitate and temperature reduction as triggers of synchronous transformation

of bloodstream to procyclic trypomastigotes *in vitro*. *Experimental Parasitology*

62, 283–291 (1986).

28. Ziegelbauer, K., Quinten, M., Schwarz, H., Pearson, T. W. & Overath, P.

Synchronous differentiation of *Trypanosoma brucei* from bloodstream to

procyclic forms *in vitro*. *Eur. J. Biochem.* **192**, 373–378 (1990).

29. Nolan, D. P., Rolin, S., Rodriguez, J. R., Van Den Abbeele, J. & Pays, E.

Slender and stumpy bloodstream forms of *Trypanosoma brucei* display a

differential response to extracellular acidic and proteolytic stress. *Eur. J.*

Biochem. **267**, 18–27 (2000).

30. Matetovici, I., De Vooght, L. & Van Den Abbeele, J. Innate immunity in the tsetse fly (*Glossina*), vector of African trypanosomes. *Dev. Comp. Immunol.* **98**, 181–188 (2019).

31. Turner, C. M., Aslam, N. & Dye, C. Replication, differentiation, growth and the virulence of *Trypanosoma brucei* infections. *Parasitology* **111**, 289–300 (1995).

32. Bargul, J. L. *et al.* Species-Specific Adaptations of Trypanosome Morphology and Motility to the Mammalian Host. *PLoS Pathogens* **12**, e1005448 (2016).

33. Rotureau, B. & Van Den Abbeele, J. Through the dark continent: African trypanosome development in the tsetse fly. *Front Cell Infect Microbiol* **3**, 53 (2013).

34. Le Ray, D., Barry, J. D., Easton, C. & Vickerman, K. First tsetse fly transmission of the 'AnTat' serodeme of *Trypanosoma brucei*. *Ann Soc Belg Med Trop* **57**, 369–381 (1977).

35. Hirumi, H. & Hirumi, K. Continuous cultivation of *Trypanosoma brucei* blood stream forms in a medium containing a low concentration of serum protein without feeder cell layers. *J. Parasitol.* **75**, 985–989 (1989).

36. Vassella, E. *et al.* Deletion of a novel protein kinase with PX and FYVE-related domains increases the rate of differentiation of *Trypanosoma brucei*. *Mol. Microbiol.* **41**, 33–46 (2001).

37. Wirtz, E., Leal, S., Ochatt, C. & Cross, G. A. A tightly regulated inducible expression system for conditional gene knock-outs and dominant-negative genetics in *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* **99**, 89–101 (1999).

38. Overath, P., Czichos, J. & Haas, C. The effect of citrate/*cis*-aconitate on oxidative metabolism during transformation of *Trypanosoma brucei*. *Eur. J. Biochem.* **160**, 175–182 (1986).

39. Brun, R. & Schönenberger, M. Cultivation and *in vitro* cloning or procyclic

culture forms of *Trypanosoma brucei* in a semi-defined medium. *Acta Trop.* **36**,

289–292 (1979).

40. Vassella, E. *et al.* A major surface glycoprotein of *Trypanosoma brucei* is

expressed transiently during development and can be regulated post-

transcriptionally by glycerol or hypoxia. *Genes Dev.* **14**, 615–626 (2000).

41. Peacock, L., Ferris, V., Bailey, M. & Gibson, W. Multiple effects of the lectin-

inhibitory sugars D-glucosamine and N-acetyl-glucosamine on tsetse-

trypanosome interactions. *Parasitology* **132**, 651–658 (2006).

42. MacLeod, E. T., Maudlin, I., Darby, A. C. & Welburn, S. C. Antioxidants

promote establishment of trypanosome infections in tsetse. *Parasitology* **134**,

827–831 (2007).

Figure 1

C

		Number of trypanosomes per blood meal			Tsetse fly infection (%)							
		1	2	3	4	5	6	7	8	9	10	11	
			Total	PAD1-negative	PAD1-positive	MG	PV	SG	TI	Tsetse infected	Tsetse dissected	Sex ratio (♀/♂)	n
i	Stumpy ES		2400	105	2295	19.3	18.1	12	0.63	107	83	1.04	3
ii			20	1	19	25.3	22.9	9.6	0.38	115	83	0.39	4
iii			2	0	2	14.6	14.6	4.5	0.31	110	89	0.96	4
iv	Stumpy SIF		2	0	2	38.8	29.3	11.2	0.29	120	116	1.70	2
v			1	0	1	20.2	18.3	4.6	0.23	122	109	0.60	2
vi			0.2	0	0.2	7.5	4.7	0.9	0.13	114	107	1.28	2
vii	Slender ES		20	19.16	0.84	22.5	22.5	5.0	0.22	104	80	0.67	3
viii			2	1.93	0.07	13.2	13.2	7.9	0.60	100	76	0.96	3
ix	Slender SIF		2	1.93	0.07	7.8	7.8	4.7	0.60	156	129	1.44	3
x			1	0.99	0.01	4.9	4.3	2.1	0.44	383	329	1.01	6
xi			0.2	198	0,002	0.9	0.9	0.0	0.00	130	114	1.35	2

Legend to Figure 1

Slender trypanosomes can complete the entire life cycle in the tsetse fly vector.

(A) Cell cycle (G1/S/post-mitotic), morphology, and differentiation of bloodstream form (mammalian-infective stage) trypanosomes. Proliferation of slender trypanosomes is detectable by duplication and segregation of the mitochondrial genome (kinetoplast, K) and nuclear DNA (N) over time. Quorum sensing causes cell cycle arrest (G0) and expression of the stumpy marker PAD1. Images are false-coloured, maximum intensity projections of deconvolved 3D stacks. The green colour indicates the nuclear GFP:PAD1$_{UTR}$ fluorescence, while the DAPI-stained kinetoplast and nucleus, and the AMCA-sulfo-NHS-labelled parasite cell surface, are shown in grey. Scale bar: 5 μm.

(B) Trypanosome infections of tsetse flies were achieved via bloodmeal, which consists typically of 20 μl, through a silicon membrane. The corresponding video is available in the Supplementary information (Supplementary Video 1).

(C) Slender trypanosomes can complete the entire tsetse infection cycle, and a single parasite is sufficient for tsetse passage. The flies were infected with either stumpy or slender trypanosomes. Stumpy trypanosomes were generated by induction of expression site attenuation (ES), or SIF-treatment (SIF). MG, midgut infection; PV, proventriculus infection; SG, salivary gland infection; TI, transmission index (SG/MG); n, number of independent fly infection experiments.

(D) Dissection of an infected tsetse fly for explantation of the alimentary tract. Scale bar: 5 mm.

(E) Explanted alimentary tract of the tsetse, with the different subcompartments labelled. Scale bar: 5 mm.

(F) Scanning electron micrograph of a typical trypanosome infection of the tsetse salivary glands, with epimastigote and mammal-infective metacyclic trypanosomes. Parasites are pseudocoloured yellow. Scale bar: 1 μm.

(G) Graphical representation of the transmission index TI (SG/MG) of slender (striped, SL) and stumpy (solid, ST) trypanosomes at different numbers per bloodmeal (data reproduced from Figure 1C). A high TI indicates successful completion of the life cycle in the tsetse vector. At low infective doses, slender trypanosomes had a higher TI compared to stumpy parasites. There was no difference between stumpy parasites generated by SIF-treatment (SIF) or expression site attenuation (ES).

Figure 2

A

	Time after infective meal (h)					
	2-4	**8-10**	**15-17**	**24-26**	**48-50**	**72-74**

PAD1 slender
EP1 slender
EP1 stumpy

B

Percentage of PAD1

			2-4	8-10	15-17	24-26	48-50	72-74
Slender	positive	1K1N	5.5	38.3	48.2	56.3	59.5	51.1
		2K1N	0.8	6.8	13.3	15.0	19.0	16.1
		2K2N	0	5.3	8.8	13.0	13.4	10.9
	negative	1K1N	67.5	39.1	21.9	11.0	5.9	15.3
		2K1N	16.7	3.8	6.4	3.5	1.1	3.6
		2K2N	9.4	6.8	1.5	1.2	1.1	2.9
Stumpy	positive	1K1N	99.7	100	95.2	97.1	82.6	75.6
		2K1N	0	0	3.3	2.3	11.4	16.5
		2K2N	0.3	0	1.4	0.6	6.0	7.1
	negative	1K1N	0	0	0	0	0	0.8
		2K1N	0	0	0	0	0	0
		2K2N	0	0	0	0	0	0

C

Legend to Figure 2

Slender trypanosomes differentiate to the procyclic life cycle stage in the tsetse fly without undergoing cell cycle arrest. Tsetse flies were infected with either slender (3.6% PAD1-positive) or stumpy (100% PAD1-positive) trypanosomes. 72 (slender) or 42 (stumpy) flies were dissected (equal sex ratios) at different timepoints after infection. Experiments were done at least three times; data are presented as sample means.

(A) Living trypanosomes (>100 cells per time point) were microscopically analysed in the explants and scored for the expression of the fluorescent stumpy reporter GFP:PAD1$_{UTR}$ in the nucleus (green bar), and the procyclic insect stage reporter EP1:YFP on the cell surface (yellow bar). The grey bars show EP1:YFP expression in infections using stumpy cells. GFP:PAD1$_{UTR}$ expression in stumpy trypanosomes is not shown, because all cells were positive.

(B) Slender (n=1845) and stumpy (n=1237) trypanosomes scored as PAD1-positive or -negative in (A), were stained with DAPI, and the cell cycle position determined based on the configuration of kinetoplast (K) to nucleus (N) at the timepoints indicated.

(C) Exemplary images of procyclic trypanosomes in the tsetse explants 24h post-infection with slender cells. Morphology (DIC panels, left), cell cycle status (DAPI label, middle panels) and expression of fluorescent reporters (right) were scored. Note that the upper panels show a cell with procyclic morphology that is nonetheless EP1:YFP-negative, indicating that the EP1 signal underestimates the total numbers of procyclic cells in the population. Scale bar: 5 μm.

Figure 3

A

Time after *cis*-aconitate addition (h)						
0	2-4	8-10	15-17	24-26	48-50	72-74

Percentage of trypanosomes

- PAD1 slender (green)
- EP1 slender (yellow)
- EP1 stumpy (grey)

PAD1 slender: 2.6, 9.8, 83.2, 94.9, 98.3, 68.2, 25.6
EP1 slender: 0, 0.8, 19.6, 14.4, 62.1, 89.9, 98.3
EP1 stumpy: 0, 28.6, 72, 79.8, 96.7, 93.4, 92.4

B

Percentage of PAD1

			0	2-4	8-10	15-17	24-26	48-50	72-74
Slender	positive	1K1N	2.6	7.3	54.2	70.3	64.1	49.6	23.1
		2K1N	0	1	13.1	6.8	17.6	10.9	0.9
		2K2N	0	1.5	15.9	17.8	16.6	7.8	1.7
	negative	1K1N	64.6	56	10.7	4.2	1.7	24	53.8
		2K1N	22.2	20.6	6.1	0	0	3.9	12.8
		2K2N	10.6	13.6	0	0.8	0	3.9	7.7
Stumpy	positive	1K1N	96.1	96.7	100	93.5	92.9	60.6	26.7
		2K1N	0	0	0	4.0	4.3	4.4	5.1
		2K2N	3.9	3	0	2.4	2.4	6.6	0
	negative	1K1N	0	0.2	0	0	0.5	22.6	50.8
		2K1N	0	0	0	0	0	2.2	4.2
		2K2N	0	0	0	0	0	3.6	10.2

Legend to Figure 3

Slender trypanosomes differentiate to the procyclic life cycle stage *in vitro* without cell cycle arrest. Cultured slender or stumpy trypanosomes were differentiated *in vitro* by the addition of *cis*-aconitate and temperature reduction to 27°C.

(A) At the times indicated, trypanosomes were analysed for the expression of the fluorescent reporters GFP:PAD1$_{UTR}$ and EP1:YFP, as in Fig. 2.

(B) Slender (n=1653) and stumpy (n=1798) trypanosomes were stained with DAPI and the configuration of the nucleus (N) and kinetoplast (K) was microscopically determined to identify the cell cycle stage. Data are compiled from five independent experiments, with each timepoint being analysed in at least two separate experiments.

Figure 4

Figure 4

A revised life cycle for the parasite *Trypanosoma brucei*. Cell-cycle-arrested metacyclic trypanosomes are injected by the tsetse fly into the mammalian host's skin. There, the parasites re-enter the cell cycle, and proliferate as slender forms in the blood, while disseminating into the interstitium and various tissues, including fat, and brain. At least two triggers (SIF or ES) launch the PAD1-dependent differentiation pathway to the cell cycle-arrested stumpy bloodstream stage. Stumpy trypanosomes can establish a fly infection when taken up with the bloodmeal of a tsetse. This work reveals that proliferating slender stage trypanosomes are equally effective for tsetse transmission, that a single parasite suffices, and that no cell cycle arrest is required for differentiation to the procyclic insect stage.

Supplementary Table 1

		Number of trypanosomes per blood meal			Tsetse fly infection (%)							
		1	2	3	4	5	6	7	8	9	10	11	
		Total	PAD1-negative	PAD1-positive	MG	PV	SG	TI	Tsetse infected	Tsetse dissected	Sex ratio (♀/♂)	n
i	Monomorph	2400	2395	5	11.0	0.6	0.0	0.0	186	155	0.64	3
ii	Monomorph + GSH +cAMP	2400	495	1905	25.5	0.0	0.0	0.0	52	47	0.82	1
iii	Slender naïve	2	750	11	6.9	6.3	2.5	0.36	165	159	1.04	3

(i, ii) Monomorphic trypanosomes do colonise the tsetse midgut, but cannot pass the proventriculus to infect the salivary glands. For infection with monomorphic parasites, cells were additionally treated for 48 hours with 12.5 mM glutathione (GSH) and 100 µM 8-pCPT-cAMP (cAMP).
(iii) Pleomorphic, naïve slender trypanosomes, freshly differentiated from metacyclic trypanosomes, readily passage through the tsetse fly.

Supplementary Video 1

Video of a tsetse fly taking a bloodmeal through a silicon membrane.

Supplementary Video 2

A

Dividing (2K2N) long slender trypanosome in the tsetse fly, 2-4 hours post infection (h.p.i). No GFP:PAD1$_{UTR}$ signal is detectable

B

Cell cycle arrested (1K1N) short stumpy trypanosome in the tsetse fly, 2-4 h p.i. The GFP:PAD1$_{UTR}$ reporter is expressed

C

Dividing (2K1N) long slender trypanosome in the tsetse fly, 15-17 h.p.i. The GFP:PAD1$_{UTR}$ signal is clearly visible

D

10 μm

Dividing (2K2N) procyclic trypanosome in the tsetse fly, 48-50 h p.i. The cell expresses both, GFP:PAD1$_{UTR}$ and EP1:YFP.

After uptake by the tsetse fly, slender trypanosomes promptly launch the PAD1 pathway, without arresting in the cell cycle. All videos were recorded at 250 fps, and the cell cycle position is indicated by DAPI staining.