1

Spontaneous polyploids and antimutators compete during the evolution of mutator cells

Maxwell A. Tracy, Mitchell B. Lee, Brady L. Hearn, Ian T. Dowsett, Luke C. Thurber, Jason Loo, Anisha M. Loeb, Kent Preston, Miles I. Tuncel, Niloufar Ghodsian, Anna Bode, Thao T. Tang, Andy R. Chia, and Alan J. Herr*

Department of Pathology, University of Washington, Seattle, Washington 98195

Running Title: Mutator evolution

Key Words: DNA replication, spontaneous polyploidization, antimutator, mutation spectra

*Correspondence: Alan J. Herr
                   Department of Pathology, Box 357705
                   University of Washington
                   1959 NE Pacific St.
                   Seattle, WA 98195-7705
                   alanherr@uw.edu
                   206-616-5063 (tel)
                   206-543-3967 (fax)

Abstract

Heterozygous mutations affecting DNA polymerase (Pol) exonuclease domains and homozygous inactivation of mismatch repair (MMR) each generate "mutator" phenotypes capable of driving tumorigenesis. Cancers with both defects exhibit an explosive increase in mutation burden that appears to reach a threshold, consistent with selection acting against further mutation accumulation. In haploid yeast, simultaneous defects in polymerase proofreading and MMR select for "antimutator" mutants that suppress the mutator phenotype. We report here that spontaneous polyploids also escape this "error-induced extinction" and routinely out-compete antimutators in evolved haploid cultures. We performed similar experiments to explore how diploid yeast adapt to the mutator phenotype. We first evolved cells with homozygous mutations affecting proofreading and MMR, which we anticipated would favor tetraploid emergence. While tetraploids arose with a low frequency, in most cultures, a single antimutator clone rose to prominence carrying biallelic mutations affecting the polymerase mutator alleles. Variation in mutation rate between subclones from the same culture suggests there exists continued selection pressure for additional antimutator alleles. We then evolved diploid yeast modeling MMR-deficient cancers with the most common heterozygous exonuclease domain mutation (*POLE-P286R*). Although these cells grew robustly, within 120 generations, all subclones carried truncating or nonsynonymous mutations in the *POLE-P286R* homologous allele (*pol2-P301R*) that suppressed the mutator phenotype as much as 100-fold. Independent adaptive events in the same culture were common. Our findings suggest that analogous tumor cell populations may adapt to the threat of extinction by polyclonal mutations that neutralize the *POLE* mutator allele and preserve intra-tumoral genetic diversity for future adaptation.

Introduction

The high fidelity of DNA replication and repair prevents mutations that might otherwise lead to cancer. As first proposed more than 40 years ago, cells with defects in these pathways exhibit an elevated mutation rate ("mutator phenotype") and an accelerated path to malignancy (Loeb *et al.* 1974; Loeb 2016). The mutator hypothesis of cancer first gained support from the discovery that Lynch syndrome was caused by defects in mismatch repair (MMR), which normally corrects DNA replication errors (LYNCH *et al.* 2009). Mouse studies later demonstrated that loss of polymerase proofreading by the major leading and lagging strand DNA polymerases (Polε and Polδ, respectively) also promoted tumorigenesis (Goldsby *et al.* 2001; Albertson *et al.* 2009). More recently, cancer genome sequencing revealed exonuclease-domain mutations in the catalytic subunit genes for Polε (*POLE*) and Polδ (*POLD1*) in a subset of highly mutated colorectal and endometrial human cancers (Yoshida *et al.* 2011; Network 2012; Palles *et al.* 2012; Church *et al.* 2013; Kandoth *et al.* 2013). Heterozygous mutations in *POLE* predominate in these "ultra-hypermutated" cancers, with the most common *POLE* variant encoding a P286R substitution (Barbari *et al.* 2018). *POLE-P286R* confers strong mutator and cancer phenotypes when engineered into mice in the heterozygous state (Li *et al.* 2018). The corresponding allele in budding yeast (*pol2-P301R*) was found to confer a strong dominant mutator phenotype that far exceeds that of an exonuclease-null allele (*pol2-4*) (Kane and Shcherbakova 2014). Recent biochemical and crystallography experiments indicate that the yeast Polε P301R variant retains residual exonuclease activity but encodes a hyperactive polymerase that excels at mispair extension (Xing *et al.* 2019) due to occlusion of the DNA binding site of the exonuclease domain by the arginine substitution (Parkash *et al.* 2019). Many of the other *POLE* exonuclease domain alleles, which confer weaker mutator phenotypes than *pol2-*

P301R when modeled in yeast, may also promote mispair extension rather than inhibit proof-

reading (Barbari *et al.* 2018).

Tandem defects in proofreading and MMR pathways produce greater than additive in-

creases in mutation rate in bacteria and yeast (Morrison *et al.* 1993; Schaaper 1993; Morrison

and Sugino 1994) indicating that polymerase fidelity and mismatch repair act redundantly to

limit replication errors. Correspondingly, some human tumors with heterozygous *POLE* or

POLD1 mutations also carry mutations affecting MMR (Network 2012; Kandoth *et al.* 2013).

Far from incidental, the MMR defects appear to synergize with the polymerase variants to ac-

celerate mutagenesis. In so doing, they create mutational signatures consistent with the inter-

action between these two pathways (Campbell *et al.* 2017; Haradhvala *et al.* 2018; Hodel *et al.*

2018). Likewise, patients with inherited biallelic MMR deficiency frequently develop rapidly mu-

tating tumors with heterozygous mutations affecting the fidelity of Polε or Polδ (Shlien *et al.*

2015). Thus, powerful mutator phenotypes based on the synergistic interactions between poly-

merase fidelity and MMR defects represent a reoccurring event in cancer evolution.

As an evolutionary strategy, cells that employ a high mutation rate run the risk of error-

induced extinction (*EEX*), where every one of their descendants eventually acquires a lethal

mutation (Morrison *et al.* 1993; Fijalkowska and Schaaper 1996; Herr *et al.* 2011a). As an ex-

treme manifestation of *EEX*, double mutant haploid yeast spores deficient in Polδ proofreading

and MMR cease dividing within 10 cellular generations (Morrison *et al.* 1993). Strong mutator

phenotypes can also drive *EEX* of diploid yeast, although the maximum tolerated mutation rate

(error threshold) is higher than in haploids, consistent with genetic buffering by the diploid ge-

nome (Herr *et al.* 2014). Colony formation slows and becomes irregular once mutation rates

are within an order of magnitude of the diploid error threshold. High mutation loads may

inherently restrict cellular longevity. Recent lifespan experiments in yeast demonstrate that

strong diploid mutators exhibit a form of genetic anticipation: the longer a lineage exists, the

higher the mutation burden, and the shorter the cellular lifespan (Lee *et al.* 2019). Evidence

suggests that error thresholds also exist in mammals. Mouse embryos completely deficient in

both proofreading and MMR initiate development but fail in the second week of gestation

(Albertson *et al.* 2009). Moreover, tumors with combined deficiencies in polymerase fidelity and

MMR appear to reach an upper limit of ~ 250 mutations/Mb, which has been interpreted as evi-

dence for a mutation threshold in cancer (Shlien *et al.* 2015).

It is unclear whether strong mutator-driven cancers adapt to the threat of extinction and,

if so, how. In yeast, one third of haploid *"eex"* mutants with combined defects in proofreading

and MMR acquire intragenic "antimutator" mutations that suppress the mutator polymerase

(Herr *et al.* 2011a; Williams *et al.* 2013). Polymerase antimutators also arise in bacteria and

bacteriophage expressing strong mutator polymerases, suggesting that these adaptive mecha-

nisms are highly conserved (Reha-Krantz 1988; Fijalkowska and Schaaper 1995). The yeast

antimutator mutations encode amino acid substitutions in a wide variety of positions within the

polymerase including the polymerase active site, DNA binding domains, polymerase structural

elements, or even the exonuclease domain itself (Herr *et al.* 2011a; Herr *et al.* 2011b; Williams

et al. 2013; Dennis *et al.* 2017). Since Polδ and Polε both play essential roles in DNA replica-

tion, these antimutator alleles found in haploid cells likely produce functional polymerases.

Some substitutions may increase nucleotide selectivity while others may promote polymerase

dissociation from mispaired primer termini, allowing extrinsic proofreading (Herr *et al.* 2011b).

Evidence for extrinsic proofreading in yeast between Polδ and Polε exists in the literature

(Morrison and Sugino 1994; Flood *et al.* 2015; Bulock *et al.* 2020) and processing of mispaired

primer termini by other DNA repair pathways may also be possible (Herr *et al.* 2011b).

The majority of *eex* mutants in the above haploid yeast genetic screens, which utilized

plasmid shuffling, carried genetic changes that were extragenic to the plasmid-borne mutator

polymerase gene and were never mapped (Herr *et al.* 2011a; Williams *et al.* 2013). Here, we

performed new genetic screens that would allow us to more easily identify these loci using

Mendelian segregation of mutator and antimutator phenotypes. In the process, we discovered

that many *eex* mutants were, in fact, not antimutators, but spontaneous polyploids, which are

buffered from mutation accumulation as described above. This finding led us to explore the

competition between antimutators and spontaneous polyploids during the evolution of diploid

mutators. These diploid mutator strains did not display an initial growth defect but became sub-

ject to selection during propagation. This scenario models a population of mutator tumor cells

having to adapt to the threat of extinction during cancer progression.

Results

Discovery of spontaneous polyploid *eex* mutants

The genetic screens that led to the discovery of spontaneous polyploid *eex* mutants be-

gan with diploid parent strains that were heterozygous for deletion of *MSH6 (msh6Δ)* and *pol3-*

01 (Polδ proofreading defective), or *msh2Δ* and *pol2-4* (Polε proofreading defective) (Figure

S1A,B). After inducing the diploids to undergo sporulation (meiosis) we isolated *eex* mutants

that suppressed the *EEX* phenotypes of freshly isolated double-mutant haploid cells. Both pa-

rental strains were heterozygous for two distinct transgene insertions at the *CAN1* locus: each

eex mutant carried either *CAN1::natMX*, which confers sensitivity to canavanine and re-

sistance to nourseothricin (NTC), or *can1Δ::HIS3*, which confers canavanine resistance and

histidine prototrophy. We crossed all *eex* mutants to WT haploids with the opposite *CAN1* al-

lele and mating type and then isolated diploid mapping strains by plating for His⁺ and NTC-

resistant colonies. We sporulated the resulting mapping strains and although they routinely

formed tetrads, the majority of mapping strains from both genetic screens failed to produce any

viable spores (Figure S1B). Surprisingly, several mapping strains with intermediate spore via-

bility produced progeny that carried both *CAN1* alleles. Flow cytometry revealed that these un-

usual mapping strains were in fact tetraploids (4n) (Figure S1*B*), while the mapping strains with

no viable spores were triploids (3n). We inferred that the original *eex* mutants for the 3n map-

ping strains were *MAT**a**/MAT**a*** or *MATα/MATα* diploids and that the 4n strains were derived

from *MAT**a**/MAT**a**/MAT**a*** or *MATα/MATα/MATα* triploid *eex* mutants. Since polyploid *eex* mu-

tants arose in both genetic screens, this escape mechanism is a general one, likely related to

the ability of polyploids to better withstand mutation accumulation.

Evolution of haploid mutator cells

A tendency to pick larger candidate *eex* colonies could explain why polyploids were

more common than antimutators in our genetic screens. To determine the relative success of

these two adaptations in an unbiased manner, we isolated 89 independent *pol2-4 msh2Δ*

spore clones by dissecting tetrads from *POL2/pol2-4 MSH2/msh2Δ* diploids on growth media

that selected for the transgenes linked to the mutator alleles (Figure 1A). Freshly dissected

pol2-4 msh2Δ haploids form colonies with an uneven perimeter, consistent with the onset of

EEX and the emergence of *eex* mutant subclones with improved survival. After two days, we

isolated four cells from the perimeter of each colony and moved them elsewhere on the plate

to assess their colony forming capacity, mutation rate and ploidy (Figure S2A). Two-thirds of

the cells isolated from the perimeters of the initial colonies failed to form a second colony — a

manifestation of *EEX*. The 120 colonies from viable cells contained 12 that were uniformly pol-

yploid (including one triploid (A4.1)) (Figure S2B, Dataset S1). To test for antimutators, we

measured mutation rates of the haploid subclones that carried the *CAN1::natMX* allele. This

involved subcloning these cells a second time to obtain replica colonies, which we assessed

for ploidy. A high frequency of polyploid replica colonies prevented measurements for 12 sub-

clones. Thirteen subclones exhibited the expected mutation rate for *pol2-4 msh2Δ* cells, while

twelve subclones displayed an antimutator phenotype (Figure S2C). These results indicate that

antimutators and polyploids emerge with similar frequencies during the initial growth of *pol2-4*

msh2Δ haploids and that polyploids continue to arise.

To investigate which *eex* strategy prevails during long-term growth, after removing indi-

vidual cells for the above analysis, we suspended the original colonies in liquid growth media

and evolved them in 96-well format with interspersed empty wells to serve as controls for con-

tamination. We repeatedly sub-cultured the cells by diluting them 1:500 (~10^5 cells/inoculum)

and growing them to saturation (~10^8 cells), yielding an estimated 10-12 generations per trans-

fer. We assessed ploidy at the initial time point (T1) and then after the 4th (T4) and the 7th

transfer (T7). The ploidies of the initial colony suspensions were overwhelming haploid, alt-

hough 15 out of 89 cultures at this early stage showed evidence of a minor diploid subclone

(Figure 1B, Dataset S1). The colony suspension from the above triploid subclone (A4) con-

tained both haploid and triploid cells in equal proportions. Prolonged growth of the original cul-

tures revealed profound fitness differences between haploid antimutators and spontaneous

polyploids (Figure 1B). At T7 only 2 cultures remained haploid; 10 were mixtures of 1n and 2n

cells, and 73 cultures were uniformly 2n or 3n. Thus, polyploids routinely win out over antimu-

tators during the adaptation of haploid cells to the mutator phenotype.

Mechanism of polyploidization in haploid mutator cultures

In the haploid evolution experiment, cells could have become polyploid by either mating

or failed cytokinesis (Ganem *et al.* 2007). Although deletion of *Homing Endonuclease* (*HO*)

stabilizes mating type in our studies, mating-type switching may still occur through random

double-stranded DNA breaks at the *MAT* locus, which are then repaired using the transcrip-

tionally silent *HML* (MATα) or *HMR* (MAT**a**) loci (Haber 2012). Conceivably, the rate of these

events could be elevated in mutator cells. To determine the relative contributions of mating and

spontaneous polyploidization, we isolated a single colony from each evolved culture and per-

formed flow cytometry. Of the 87 isolates, there were 83 diploids, 1 haploid, 1 triploid, and 2

mixed cultures with both 1n and 2n cells (Figure 1C). Mating-type PCR assays revealed 41

*MAT**a**/MAT**a***, 36 *MAT*α/*MAT*α and 6 *MAT**a**/MAT*α diploids (Figure 1D). Three of the

*MAT**a**/MAT*α isolates were heterozygous at the *CAN1* locus (*CAN1::natMX/can1Δ::HIS3*) (Da-

taset S2), providing evidence for rare mating events between cells from neighboring wells. The

three remaining *MAT**a**/MAT*α cells were homozygous for *CAN1::natMX* or *can1Δ::HIS3* and

could have arisen through contamination or mating-type switching. We specifically investigated

the frequency of mating-type switching with both *MAT**a*** and *MAT*α cells. Diploids formed in

both same-sex pairings with similar frequencies. But while *MAT**a** x MAT**a*** crosses nearly al-

ways gave rise to *MAT**a**/MAT*α diploids, *MAT*α *x MAT*α crosses produced *MAT*α*/MAT*α dip-

loids (Figures S3, S4). Same-sex mating has previously been observed (Strathern *et al.* 1981).

Given the similarity in frequencies, a dsDNA break at the *MAT* locus likely initiates both mat-

ing-type switching of *MAT**a*** cells and *MAT*α same-sex mating (see Methods). The 3

*MAT**a**/MAT*α clones that were homozygous for the *CAN1::natMX* or *can1Δ::HIS3* allele, which

arose by mating type switching, suggests that only a small percentage of the 36 *MAT*α*/MAT*α

cultures in the evolution experiment arose by same-sex mating. Together these observations

argue that the majority of *MAT**a**/MAT**a*** or *MATα/MATα* isolates in our evolving populations

arose by spontaneous polyploidization due to failed cytokinesis, which recent studies indicate

occurs with a frequency of 7×10^{-5} in haploid yeast (Harari *et al.* 2018). Since tetraploids have

been observed in various cancers, we reasoned that spontaneous polyploidization could con-

ceivably represent a conserved mechanism of escape from *EEX* for mutator-driven tumors.

Tetraploidization further protects spontaneous diploids.

As a first test of whether tetraploidy confers an advantage in a diploid mutator popula-

tion we utilized the above spontaneous *pol2-4/pol2-4 msh2Δ/msh2Δ* diploids. We isolated

*MAT**a**/MAT**a*** and *MATα/MATα* cells from different evolved cultures by microdissection and

mated their daughter cells. We moved the parental diploid cells as well as the resulting tetra-

ploid zygotes to distinct locations on the agar plates to form colonies (Figure 2A). We then

mixed each parent diploid strain separately with their tetraploid progeny at different ratios

(100:1, 10:1, 1:1, 1:10, 1:100) and grew them in competition. The tetraploids overtook the pa-

rental mutator diploids in 7 out of 8 competitions. These experiments suggest that tetraploid

mutators would potentially have a sizeable fitness advantage over the diploid mutator popula-

tion from which they arise. An important caveat to this experiment, however, is that both diploid

parents likely carry homozygous mutations that first arose in the original haploid mutators.

These genetic liabilities would became heterozygous in their tetraploid offspring. In diploid cells

that spontaneously acquire a mutator phenotype, mutations would accumulate from the begin-

ning in a buffered, heterozygous state.

Adaptation of diploid yeast to a strong sublethal mutator phenotype

To explore how diploid cells with a near WT mutation burden adapt to a strong mutator

phenotype, we began with *pol3-01/pol3-01 msh6Δ/msh6Δ* cells, which form robust colonies

despite having a mutation rate (1 x 10^{-3} Can^R mutants/division) just below the diploid error

threshold (Figure 3A)(Herr *et al.* 2014). Using a mathematical model in which the decrease in

fitness was driven solely by lethal homozygous inactivation of essential genes (Herr *et al.*

2014), we estimated that a tetraploid or antimutator *eex* mutant would overtake a *pol3-01/pol3-*

01 msh6Δ/msh6Δ diploid culture in 200 generations (Figure S5, Dataset 1). With two sets of

homozygous mutator alleles driving the mutator phenotype, we anticipated tetraploids would

be at their greatest advantage relative to antimutators.

To set up the evolution experiment, we first isolated numerous independent *pol3-*

01/pol3-01 msh6Δ/msh6Δ zygotes by mating *pol3-01 msh6Δ* haploids that were freshly dis-

sected from the *pol3-01/POL3 msh6Δ/MSH6* parental strain (Figure 3A). We evolved 89 zy-

gote clones through ~300 generations, monitoring ploidy at the T1, T10, T19, and T27 trans-

fers (Figure 3B). A single pure tetraploid culture emerged at the end of the experiment. The re-

maining cultures were overwhelmingly diploid, suggesting that they adapted through antimuta-

tor mutations. To assay for antimutator phenotypes, we isolated independent subclones from

T25 *CAN1::natMX/can1Δ::HIS3* cultures, assessed their ploidy, and measured mutation rates.

Almost all subclones were diploids (Dataset 1) and exhibited a clear antimutator phenotype

(Figure 4A). Antimutators also dominated the T13 cultures and were evident as early as T1

(Figure 4A). T1 cells from every culture formed variably sized colonies, which were notably

smaller than T13 colonies, suggesting that strong selection for *eex* mutants exists by ~30 cellu-

lar generations (20 generations/colony and 10 generations for T1) (Figure S6).

Independent T25 isolates from the same *pol3-01/pol3-01 msh6Δ/msh6Δ* culture some-

times displayed different mutation rates (Figure 4B). We wondered whether they represented

independent adaptations, and therefore sequenced the genomes of eight sets of subclones. In

each set, the subclones could be arranged into a phylogenetic tree with ~1300 to 4500 shared

single nucleotide variants (SNVs) in the trunk and ~1400 to 3000 unique SNVs in the branches

(Figure 4C,D). This indicates that a single adapted clone rose to prominence in each culture

and then diverged. Thus, differences in mutation rates between subclones likely reflect the

presence of additional antimutator mutations in some isolates but not others.

The mutations that delineate the phylogenetic trees bear the imprint of the mutator and

antimutator phenotypes involved in the evolution of these cultures. We classified the shared

and unique SNVs according to the 96 trinucleotide contexts defined by the six different muta-

tion subtypes (C:G→A:T, C:G→G:C, C:G→T:A, T:A→A:T, T:A→C:G, T:A→G:C), flanked by

all possible 5' and 3' nucleotides (Figure 4D). We then compared mutation spectra using the

hypergeometric and cosine similarity tests. By both tests, the spectra of the trunk mutations

were largely the same between different cultures, consistent with each culture beginning with a

common mutator phenotype (Figures 4D, S7, Dataset S2). The spectrum of unique mutations

in each branch of a given tree differed substantially from that of the trunk mutations ($p <$

0.0001), although generally not from each other, consistent with the antimutator phenotype

arising prior to their divergence (Figure 4D, S7). The antimutator alleles caused particularly

sharp decreases in the relative abundance of C:G→A:T mutations in the branches of all trees.

Despite these similarities, the spectra of unique mutations from unrelated evolved cultures of-

ten differed statistically from each other, suggesting that the antimutator alleles altered replica-

tion fidelity in distinct ways (Figure 4D, Dataset S2). Thus, acquisition of an antimutator pheno-

type that fundamentally changes the mutation spectrum coincides with the key adaptive event

in each culture.

To understand how the cells attenuated the mutator phenotype, we annotated the muta-

tions (Dataset S2) and compared the affected genes. All 8 strain sets acquired multiple *pol3*

mutations prior to their divergence (Figure 4C,D, see trunks). Polδ plays an essential role in

DNA replication. Both *pol3-01* alleles contribute to the onset of error-induced extinction, but

only one allele needs to remain functional for cells to survive. Thus, the premature nonsense

codons observed in three trees (B1, C2, D2) contribute to mutator suppression by inactivating

one of the two *pol3-01* alleles, which allows the other *pol3-01* allele with an antimutator muta-

tion to serve as the sole source for Polδ. Many missense mutations affect the same amino acid

residue as (or are identical to) known antimutator mutations (*A704V(2x), A786V*, *A677T,*

D831A, R475S, R815C) (Herr *et al.* 2011a; Herr *et al.* 2014; Dennis *et al.* 2017). The other

missense mutations affecting *pol3-01* may inactivate polymerase function or may be *bona fide*

antimutator alleles that have not previously been identified. Candidate antimutator mutations

include *P942L* and *M578I*, which were each observed in two independent lineages, as well as

mutations affecting P796, which were isolated three independent times (*P796L*(2x) and

P796S) — once in combination with a stop codon. In some lineages, additional *pol3* alleles

were observed in the branches. In two cases (G5-3, A6-2), these may account for the variation

in mutation rates between subclones from the same culture (Figure 4B), but in other cases

they clearly do not (B2-4, C2-3, D2-1, B6-2, B6-3, C7-3), implicating other loci in the attenua-

tion of mutation rates.

We looked for genes under positive selection by comparing the ratio of nonsynonymous

and synonymous mutations (dN/dS) across the genome using *dNdscv*, which normalizes the

observed dN/dS ratio by the 192-trinucleotide mutation spectra of each gene in order to assess

whether the observed ratio departs from neutrality (dN/dS = 1) (Martincorena *et al.* 2017). We

analyzed the mutations from the trunks of the 8 lineages and found that only *pol3* mutations

are under positive selection (q = 0.004, corrected *p*-value using Benjamini and Hochberg's

false discovery rate)(Dataset S2). Mutations in other genes, within the resolution of this da-

taset, accumulate in a neutral fashion. Repeating this analysis for the 23 unique sets of muta-

tions from the branches failed to reveal any other genes under strong selection despite evi-

dence for their existence, suggesting that more observations are needed for sufficient statisti-

cal power, especially if there are multiple antimutator genes. Interestingly, when branch and

trunk mutations were analyzed together, *pol3* mutations do not emerge from the multiple test-

ing correction as significant, underscoring the importance of using phylogenetic information for

the identification of candidate antimutators by dN/dScv.

Being able to classify mutations according to the phylogenetic tree also provided evi-

dence for negative selection acting against the mutator phenotype. For subclasses of trunk

mutations involving missense or nonsense mutations, which arose early during mutator evolu-

tion, the genome-wide (or global) dN/dS ratio was 1.0, indicative of neutrality (Figure 4E)(Da-

taset S2). However, the genome-wide dN/dS ratio in the branches for nonsense mutations was

0.8, indicating that negative selection begins to act against the most deleterious class of muta-

tions during extended propagation of these lines.

Modeling adaptation to the *POLE-P286R* mutator phenotype in cancer

Having established that diploid mutators rapidly evolve to avoid extinction in long-term

culture, we sought to understand how cells would adapt to the most common *POLE* mutator

allele in cancer, *P286R*. Most *POLE-P286R* tumors appear microsatellite-stable and bear mu-

tational signatures that suggest that they are MMR proficient (Haradhvala *et al.* 2018). How-

ever, one *POLE-P286R* patient curated on cBioPortal (Cerami *et al.* 2012) carried two distinct

null alleles of *MSH6* (TCGA-IB-7651)(Dataset 1) and a clear mutation signature consistent with

synergy between *POLE-P286R* and MMR deficiency (Haradhvala *et al.* 2018). To model this

extreme scenario, we took a similar approach as before and isolated numerous independent

pol2-P301R/POL2 msh2Δ/msh2Δ zygotes by crossing freshly dissected *pol2-P301R msh2Δ*

and *POL2 msh2Δ* haploid spores (Figure 5A). Whereas concurrently isolated double homozy-

gous mutants formed small colonies of around 1 x 10^5 cells, the *pol2-P301R/POL2*

msh2Δ/msh2Δ zygotes grew robustly (Figure 5B, Figure S8) and we evolved 81 independent

clones through 10 transfers (~120 generations). By T10, one culture had evolved a prominent

tetraploid subclone (Figure 5C). To test for antimutator phenotypes in the others, we isolated

three subclones from 44 *CAN1::natMX/can1Δ::HIS3* cultures. Genotyping revealed 1 out of the

132 isolates had lost the heterozygous *pol2-P301R* allele through mitotic recombination (H9-

1). Most of the other isolates had mutation rates more than 10-fold lower than the original zy-

gote clones, and many as low as the mitotic recombinant, which serves as an internal

msh2Δ/msh2Δ control (Figure 5D) (Dataset S1). Thus, as with *pol3-01/pol3-01 msh6Δ/msh6Δ*

diploids, there is strong selection pressure for *pol2-P301R/POL2 msh2Δ/msh2Δ* cells to sup-

press their mutator phenotype, even though they do not show an obvious initial growth defect.

To understand the basis for the antimutator phenotypes, we sequenced the genomes of

119 subclones from 40 evolved cultures. The subclones averaged 665±178(SD) SNVs and

225±35(SD) Indels, corresponding to 60 SNVs/Mb and 20 Indels/Mb (Dataset S2). We anno-

tated all variants (Dataset S2) and constructed phylogenetic trees using the SNVs (Figure 5E).

Secondary mutations in *pol2* were found in all subclones, suggesting that they are adaptive

(Figure 5E, Dataset S2). In 6 out of 40 evolved cultures, the *pol2* mutation preceded the diver-

gence of the subclones (B1, B11, D9, F6, F7, H11). As they diverged, the subclones

accumulated few additional mutations, providing strong evidence for a direct role of the *pol2*

mutations in mutator suppression. Altogether, we observed 88 candidate adaptations in *pol2*,

including 45 frameshifts, 23 nonsense mutations, and 36 missense (or combinations of mis-

sense) mutations. Two mutant alleles, *I279fs* (H2-2) and *D306Y* (H9-2,3), were close enough

to *P301R* to confirm they reside on the same chromosome. The abundance of truncation muta-

tions suggests that inactivation of the *pol2-P301R* allele allows cells to use WT *POL2* as the

sole source for Polε, thereby dramatically lowering their mutation rate. This mirrors how the

combination of nonsense and antimutator mutations suppresses the mutator phenotype of

some *pol3-01/pol3-01 msh6Δ/msh6Δ* cultures. The abundance of truncating *pol2* mutations in

these strains also indicates that the observed missense mutations may either inactivate the

mutator polymerase or function as antimutators.

The central role of the *pol2* mutations in the evolution of these cultures helps to interpret

the structure of the phylogenetic trees. Since most *pol2* mutations occurred after divergence of

the subclones, most of the 300±126(SD) shared mutations in the trunks likely arose prior to zy-

gote formation in the *pol2-P301R msh2Δ* haploids. The magnitude of the mutator phenotype of

these haploid cells is likely similar to *pol2-P301R/pol2-P301R msh2Δ/msh2Δ* diploid cells,

which formed small colonies (Figures 5B, S8). In the mathematical model of diploid *EEX*, ho-

mozygous inactivation of essential genes limits growth (Herr *et al.* 2014). A terminal colony

size of 1×10^5 cells corresponds to a mutation rate of ~5×10^{-3} Can^R mutants per division,

which is 10 times greater than the observed *pol2-P301R/POL2 msh2Δ/msh2Δ* mutation rate

(5.43×10^{-4} Can^R mutants/division). We can deduce the genome-wide mutation rate of *pol2-*

P301R/POL2 msh2Δ/msh2Δ cells from the minimal internodal distance between branch points

in trees that rapidly diverged and acquired independent *pol2* mutations (Dataset S2). The

lowest number of shared SNVs between any two subclones was 24. Pairs of subclones from 8

cultures (A5, A7, A8, C11, F10, G6, G8, and H6), with similar low increments of shared SNVs,

averaged 30±7 (SD) SNVs/diploid genome. This would place the mutation rate of *pol2-*

P301R/pol2-P301R msh2Δ/msh2Δ diploids at ~300 SNVs/division and the *pol2-P301R msh2Δ*

haploids cells at 150 SNVs/division, easily accounting for the observed number of mutations in

the trunks of the phylogenetic trees.

If a mutation rate of 30 SNVs/division continued unabated in *pol2-P301R/POL2*

msh2Δ/msh2Δ cells, the subclones would have accumulated ~3600 mutations over 120 gener-

ations. However, the average branch length was 418(±188 SD). A truncating mutation in *pol2-*

P301R that nullifies the mutator phenotype should lower the mutation rate to that observed in

msh2Δ/msh2Δ cells (5 x 10^{-6} Can^R mutants/division)(see H9-1, Figure 5C, Dataset S2) , which

we estimate would be ~0.3 mutations/diploid genome/division. A hundred cellular generations

at this suppressed mutation rate would produce 30 SNVs, equal to the number that occurs in

just one division of the original mutator strain. Subtracting this value from the average branch

length, suggests that the average adaptative event occurred by ~ 13 cellular generations, be-

fore the colony reached 10^4 cells (2^{13}). Thus, adaptive mutations in *pol2-P301R* occur early

during the evolution of these strains, substantially moderating the mutation burden.

Since the bulk of mutations in branches with strong *pol2* null alleles come from these

early divisions, we pooled them to determine the mutation spectrum produced by the *pol2-*

P301R/POL2 msh2Δ/msh2Δ mutator phenotype (Figure 5F)*.* The spectrum from the branches

matches the spectrum of the pooled mutations from the trunks (cosine similarity of 0.994) de-

spite likely differences in the magnitude of their mutator phenotypes (Figure 5F). We also com-

pared the branch mutation spectrum to all mutational signatures from the Catalogue of

Somatic Mutations in Cancer (Cosmic) database (Alexandrov *et al.* 2020) to test how well

these yeast cells recapitulate the mutational processes occurring in human disease. We found

the highest similarity with Cosmic mutational signature 14 (0.793), which results from the syn-

ergy between *POLE* and MMR mutator alleles (Haradhvala *et al.* 2018; Alexandrov *et al.* 2020)

(Table S1). Thus, our analysis of the phylogenic trees of evolved *pol2-P301R/POL2*

msh2Δ/msh2Δ strains reveal a severe mutator phenotype that resembles the mutagenesis in

cancers with combined *POLE* and MMR deficiencies. In this model, we show that adaptation

readily occurs through suppression of the *pol2-P301R* allele. In nearly every culture, the adap-

tation is polyclonal, resulting in a genetically diverse population of cells that survive extinction.

Discussion

Cell lineages with very high mutation rates face a progressive loss of fitness and even

extinction unless they adapt by becoming polyploid or acquiring antimutator mutations. The

competition between polyploids and antimutators during the evolution of mutators is influenced

by the rate at which they arise and the extent to which they preserve fitness.

In our study of *pol2-4 msh2Δ* haploids, we found that two-thirds of cells sampled from

the perimeter of the initial colonies (<20 cellular generations) were inviable, illustrating the

strong selection pressure for *eex* mutants. At this early stage, polyploids and antimutators

were present with similar frequencies, which is in keeping with the comparable rates of sponta-

neous polyploidization of haploid yeast (7×10^{-5} per division) (Harari *et al.* 2018) and gene in-

activation of *pol2-4 msh2Δ* cells (1×10^{-4} Can^R mutants/division). Diploids frequently emerge in

WT haploid evolution studies, with a fitness advantage of around 3.5% per generation

(Gerstein *et al.* 2006; Venkataram *et al.* 2016; Fisher *et al.* 2018). The fitness advantage of

polyploids during the evolution of haploid mutators is likely much larger and constantly

increasing as mutations accumulate, as suggested by studies with chemically mutagenized

haploid yeast (Mable and Otto 2001). As the mutator parent strain loses fitness and polyploids

and antimutators compete for supremacy, continued mutation accumulation in antimutators

may place them at a competitive disadvantage. Even more consequential, the antimutator mu-

tations may inherently compromise fitness by diminishing polymerase processivity. In some

cases, antimutators may even cause DNA replication stress and a need for homologous re-

combination, which would favor the emergence of polyploid variants within the antimutator

population.

While spontaneous polyploids and antimutators also arise during the evolution of strong

diploid mutators, antimutators dominate nearly every culture. This may reflect stark differences

in how frequently the two types of mutants arise. The rate of tetraploidization of diploid yeast,

to our knowledge, has not been reported, but our data provide an estimate based on the fol-

lowing logic. Sequencing shows that a single adapted clone rose to prominence in each *pol3-*

01/pol3-01 msh6Δ/msh6Δ culture. We observed that 1% of these cultures became tetraploid

during their evolution. Mutator suppression of *pol3-01/pol3-01 msh6Δ/msh6Δ* cells requires

two independent *pol3* mutations, whereas polyploidization is a singular event. Biallelic attenua-

tion of the *pol3-01* mutator phenotype will occur with a rate no greater than the square of the

per gene mutation rate, which for *pol3-01/pol3-01 msh6Δ/msh6Δ* cells is $\sim(1 \times 10^{-3})^2$ or 10^{-6}. If

we assume that tetraploids and antimutators have a similar stabilizing effect on the loss of fit-

ness, the rate of tetraploid formation may be no greater than 10^{-8}. Of course, the rate of tetra-

ploid formation could potentially be higher if tetraploids are inherently less fit than diploid anti-

mutator strains. Growth competition studies between WT tetraploids and diploids suggests this

may be true (Gerstein *et al.* 2006). Tetraploids suffer higher rates of genomic instability and

depend more heavily on factors involved in homologous recombination (Storchova *et al.* 2006).

In the context of a mutator phenotype, a higher rate of homologous recombination in tetra-

ploids may limit buffering of deleterious mutations by promoting loss of heterozygosity. Diploid

antimutators are already buffered from mutations and may retain that protection better than tet-

raploids if they indeed have lower recombination rates.

The ease with which *pol3-01/pol3-01 msh6Δ/msh6Δ* cells adapted to *EEX* despite being

homozygous for the mutator alleles, suggests that similar events may be common in cancers

with heterozygous *POLE* mutations. We modeled this scenario using *pol2-P301R/POL2*

msh2Δ/msh2Δ cells, which incur 2.7 SNVs/Mb/division (30 SNVs/11Mb). This level of muta-

genesis imposes a strong selection for suppressor mutations, and we found a wide variety of

suppressor mutations that reduced mutation rate as much as a 100-fold. The average mutation

burden of these adapted subclones (60 SNVs/Mb) is less than the clonal mutation burden of

many ultra-mutated tumors (>100 SNVs/Mb). Half of *POLE-P286R* tumors curated on cBioPor-

tal (Cerami *et al.* 2012) carry clonal, secondary mutations in *POLE* (Figure 5G, Dataset S1)*.*

Intriguingly, four of these carry truncating *POLE* mutations including TCGA-IB-7651

(X1335_splice), whose mutation spectrum reflects synergy between *POLE* and MMR defects

(Haradhvala *et al.* 2018). The short-read sequencing used for these samples does not reveal

whether the mutations are co-linear with *P286R*. But there are only two possibilities: the muta-

tions either suppress the mutator or WT allele. Yeast cells that only express *pol2-P301R* have

a 27-fold higher mutation rate (1.4×10^{-4} Can^R mutants/division) than cells that are heterozy-

gous for the allele (Kane and Shcherbakova 2014). Thus, truncating *POLE* mutations in the

WT allele may enhance the *POLE-P286R* mutator phenotype. The TCGA-IB-7651 tumor car-

ries four other mutations in *POLE* besides the truncating mutation, making it a reasonable

candidate for mutator suppression regardless of which *POLE* allele carries the truncating mu-

tation. The mutational signatures of the other three tumors with truncating *POLE* mutations

(TCGA-AX-A05Z, P0010967, P0005824) have high mutation loads but lack a mutational signa-

ture of combined *POLE* and MMR defects (Haradhvala *et al.* 2018). Thus, these truncating mu-

tations may enhance the *P286R* mutator phenotype. Given their high mutation loads, we pre-

dict that these tumors may also experience selection pressure for *P286R* suppressor muta-

tions. However, since *POLE* is essential, adaptations in these types of tumors may be limited

to missense mutations that restore Polε fidelity.

The adaptive missense mutations in our study encode amino acid changes that may in-

activate the polymerase or attenuate the mutator phenotype. They group together in different

locations within the Pol2p structure (Swan *et al.* 2009; Parkash *et al.* 2019), suggesting dis-

crete mechanisms of suppression (see Figure S9 for details). The most striking antimutator

candidate is D306Y, which is adjacent to P301R in the exonuclease structure (Figure S9B) and

may even be allele specific. Together, the truncating and missense mutations argue that

POLE cancer cells have numerous ways to escape error-induced extinction. The frequent

emergence of independent *eex* mutants in the same evolved culture indicates that polyclonal

adaptation may be common in *POLE* tumors, potentially obscuring the critical role of suppres-

sor mutations in preventing error-induced extinction of the cancer and preserving genetic diver-

sity for tumor evolution.

Other changes that attenuate the mutator phenotype may also be possible. In yeast, de-

letion of *DUN1* suppresses Polδ (Datta *et al.* 2000; Mertz *et al.* 2015) and Polε mutator pheno-

types (Williams *et al.* 2015) in haploid MMR-proficient cells. The *dun1Δ* antimutator effect in-

volves a reduction in dNTP pools, since restoration of dNTPs by deletion of *SML1*, a Dun1

target, restores the mutator phenotype (Williams *et al.* 2015). Like intragenic mutations in poly-

merase genes predicted to compromise DNA binding, mutations that lower dNTP pools may

enhance polymerase dissociation from the primer•template, permitting extrinsic proofreading

(Williams *et al.* 2015). Although heterozygous variants in *DUN1* are observed in some *pol3-*

01/pol3-01 msh6Δ/msh6Δ lineages, they do not explain the observed variation in mutation

rates between subclones. While MMR-proficient *pol3-01* cells appear to activate the check-

point (Datta *et al.* 2000), this activation may depend, in part, on processing of mismatches by

MMR (Reha-Krantz *et al.* 2011). Therefore, in the absence of MMR, the contribution of Dun1p

to the *pol3-01* mutator phenotype may be substantially reduced. Recent evolution studies in

diploid yeast pointed to unconventional mutator phenotypes caused by heterozygous muta-

tions in house-keeping genes (Coelho *et al.* 2019). The vast phenotypic space created by het-

erozygous mutations and their epistatic interactions remains largely unexplored. If mutation

rate can be enhanced by such mutations, perhaps it may be attenuated as well.

Materials and Methods

Culture conditions. Yeast propagation and tetrad dissection followed standard proce-

dures (Sherman 2002). Unless otherwise noted, reagents were purchased from Fisher Scien-

tific or Sigma. For rich liquid media, we used YPD (1% wt/vol yeast extract, 2% wt/vol peptone,

2% wt/vol dextrose). For rich solid media we used YPD or synthetic complete (SC) [6.7 g Difco

yeast nitrogen base without amino acids, 2% wt/vol dextrose, 2 g/L SC amino acid Mix (SCM)

(Bufferad)] supplemented with 2% wt/vol agar. For drop-out media, SCM powders lacking

specified amino acids were either purchased from Bufferad or made to the same specifications

(Kaiser 1994). For haploid mutation rates, we used SC plates lacking arginine (SC-Arg) with 60

μg/mL canavanine (Can). For mutation rates with diploid strains we used SC_MSG-Arg [1.7 g

Difco yeast nitrogen base without amino acids or ammonium sulfate, 1 g/L monosodium gluta-

mate (for a nitrogen source), 2% wt/vol dextrose, 2 g/L SCM-arginine)] with 60 µg/mL Can and

100 µg/mL nourseothricin (NTC) (Herr *et al.* 2014). We used similar plates lacking histidine

and containing NTC (SC_MSG-His+NTC) to select for mating between *CAN1::natMX* and

can1Δ::HIS3 cells (see below).

Yeast Strains. The diploid parent strain used for these experiments, AH0401, is a

BY4743 derivative engineered to be heterozygous at the *CAN1* locus (Herr *et al.* 2014). One

allele carries the *natMX* transgene (encoding resistance to NTC) inserted just downstream of

the *CAN1* coding sequence (*CAN1::natMX*), while the other *CAN1* allele is deleted and re-

placed with *HIS3* (*can1Δ::HIS3*). Selection for *can1* mutants in the presence of canavanine

and NTC allows us to score *bona fide can1* mutants without the high background of

can1Δ::HIS3/can1Δ::HIS3 mitotic recombinants (Herr *et al.* 2014). The four WT strains used

for generating mapping strains (Figure S1) and for the mating-type switching experiments (Fig-

ures S3, S4) were dissected from the same AH0401 tetrad: AH4002 (*MATα CAN1::natMX)*,

AH4003 (*MAT**a** CAN1::natMX)*, AH4010 (*MATα can1Δ::HIS3*), and AH4015 (*MAT**a***

can1Δ::HIS3). AH12721 is a *POL2/URA3::pol2-4 MSH2/msh2Δ::LEU2* strain isolated by mat-

ing freshly dissected spores from AH2801 (*POL2/URA3::pol2-4 MSH6/msh6Δ::LEU2*)

(Kennedy *et al.* 2015) and AH5610 (*MSH2/msh2Δ::LEU2*). Zygotes were isolated by microdis-

section after 8 hours and allowed to form colonies, which were then genotyped. AH5610 was

constructed for this study by transforming AH0401 with a *LEU2* PCR product amplified from

pRS416 with Msh2U and Msh2D as described (Williams *et al.* 2013). AH2601 is a previously

described *POL3/URA3::pol3-01 MSH6/msh6Δ::LEU2* strain (Lee *et al.* 2019) constructed in the

AH0401 background. All strains derived from mating between *pol3-01 msh6Δ* spores used in

the evolution experiment were designated AH164_NN, where "NN" refers to their coordinate in

the 96-well plate. AH11304 is a *URA3::pol2-P301R* transformant of AH5610 in which a dupli-

cated *POL2* promoter flanks the *URA3* transgene upstream of the *POL2* coding sequence. We

engineered the strain using a chimeric URA3::*pol2P301R* PCR product amplified in two frag-

ments from pRS416-POL2 (Williams *et al.* 2013) with Phusion polymerase (New England Bi-

olabs) and the following conditions: 98°C, 1min; 25 x (98°C, 10 sec; 51°C, 30 sec; 72°C, 2

min); 72°C 2 min. One fragment was amplified with pol2U (ATGATGAAAGAGCACAT-

TCTATCAAGATAACACTCTCAGGGGACAAGTATAGATTGTACTGAGAGTGCAC) and

POL2P301R-r (CATTATTTGATCTACGGCGGAATCCcGGAATTTTAAAGGCGGCTTCGTGG;

pol2-P301R mutation, lower case). A second *POL2* fragment was amplified with POL2P301R

(CCACGAAGCCGCCTTTAAAATTCCgGGATTCCGCCGTAGATCAAATAATG) and pol2S7

(ATGTGGATAACTTGGTCTGCG). The two PCR fragments were gel purified, equimolar ratios

were combined without primers in a second PCR reaction for ten cycles (98°C, 10 sec; 55°C,

30 sec; 72°C, 5 min), outside primers pol2U and pol2S7 were added, and the reaction was

continued for another 10 cycles. The entire *pol2-P301R* gene was initially confirmed by Sanger

sequencing and then by whole genome sequencing of the evolved clones.

Isolation of eex mutants. For *pol3-01 msh6Δ eex* mutants (Figure S1), we plated large

numbers of random haploid spores onto media that selected for the *URA3* and *LEU2*

transgenes tightly linked to each mutator allele. The *pol3-01 msh6Δ* cells do not form visible

colonies unless they acquire an *eex* mutation. The isolation of *pol2-4 msh2Δ eex* mutants re-

quired a slightly different approach. During our initial tetrad analysis of the *pol2-4/POL2*

msh2Δ/MSH2 strain, we found that *pol2-4 msh2Δ* cells were not synthetically lethal, as they

were in our earlier plasmid shuffling system, but grew slowly (Williams *et al.* 2013). To isolate

eex mutants, we simply re-plated the slow-growing double mutant spore clones for faster-

growing colonies. Haploid yeast cells grow vegetatively as one of two mating-types, *MAT***a** and

*MAT*α, which can mate to form *MAT***a**/*MAT*α diploids. To isolate mapping strains, we mated

each *eex* mutant in parallel with AH4002, AH4003, AH4010, and AH4015 and plated the mu-

tants on SC_MSG-His+NTC. Candidates that only mated with one of the four strains were car-

ried forward for genetic analyses.

Assay for mating-type switching or same-sex mating. We mixed ~5 x 10^6 cells of two

strains with the same mating type but opposite *CAN1* alleles (*can1Δ::HIS3* and *CAN1::natMX*)

in a microtiter plate with YPD and incubated the cells at 30°C for 8 hours without shaking.

CAN1::natMX/can1Δ::HIS3 diploids were then selected on SC_MSG-His+NTC media. Wild

type and *pol2-4 msh6Δ* mutator cells from the T1 transfer were used for Figure S3. Wild-type

cells picked from individual colonies were used for Figure S4. Mating type PCRs were per-

formed as described (Harari *et al.* 2018).

Explaining the similar rates of mating-type switching and same-sex mating. Our finding

that diploid colonies occur with similar frequencies indicates that a dsDNA break likely initiates

both mating-type switching and same sex mating. In *MAT***a** cells the dsDNA break is repaired

by gene conversion from the HML locus, converting the cell to a *MAT*α cell, which then mates

with an adjacent *MAT***a** cell to form a *MAT***a**/*MAT*α diploid. In *MAT*α cells, the dsDNA break

likely disrupts expression of the MATα2 repressor, which normally suppresses the expression

of genes required for the *MAT***a** program (Strathern *et al.* 1981). These phenotypic "*MAT***a**"

cells then mate with adjacent *MAT*α cells and repair the break through intra- or inter-chromoso-

mal recombination.

Evolution Experiments. To obtain *URA3::pol2-4 msh2Δ::LEU2* double mutant spores we

dissected tetrads on SC-Ura-Leu. After two days of growth, we isolated single cells from the

perimeter of each colony by microdissection and moved them below the colony to form new

colonies (Figure S2A). We then suspended the main colonies in 100 μls of H_2O in 96-well for-

mat (Figure 1). Random empty wells were interspersed among the samples to monitor for con-

tamination. We made glycerol stocks from part of the cell suspension. From the remaining cell

suspensions, we diluted the cells 1:50 in YPD, and grew 3 hours before removing aliquots for

the first flow cytometry measurement after which the cultures were grown to saturation and

then subcloned as described below.

To isolate mutator diploids for evolution experiments, AH2601 or AH11304 tetrads were

dissected on SC-Ura-Leu plates and spores were separated from each other by ~100 μm (the

width of two diameters of the dissecting needle). After 2 to 3 divisions, dividing cells from differ-

ent tetrads were placed next to each other to allow mating. Half of the pairings produced zy-

gotes after 2 to 3 additional divisions. Zygotes were moved to a defined coordinate on the plate

and allowed to form a colony. In our initial experiment with AH2601, we noted that Ura⁻ Leu⁺

spores germinated and divided once on this media and that we could isolate *pol3-01/POL3*

msh6Δ/msh6Δ heterozygotes by crossing them with more rapidly dividing Ura⁺ Leu⁺ cells. We

used this approach to enrich for *pol2-P301R/POL2 msh2Δ/msh2Δ* strains in the AH11304

crosses. We also obtained numerous *pol2-P301R/pol2-P301R msh2Δ/msh2Δ* zygotes in the

process, which allowed us to compare the relative colony forming capacity of the two geno-

types. Colonies were picked from the plate and suspended in 100 μls of H_2O. We used 30 μls

of the cell suspensions to inoculate 500 μls cultures, 40 μls for glycerol stocks, and the remain-

ing cell suspension to measure mutation rates (see below). For AH11304 derived zygotes, we

reserved 10 uls for *pol2P301R* genotyping. The *pol2P301R* allele creates a sequence

(…AATTCCgGG…) that is one bp from a DraI site (CCCGGG). To genotype *pol2-P301R* we

amplified *POL2* with pol2-P301R-dctF (TGTGGTAATGGCATTTGATATAGAAACCACGAA-

GCCGCCTTTAAAATcCC), which has a mismatch (lower case) that creates a DraI site, and

pol2-P301R-dctF (GGATACTCCGGTTTCGGTGTATACTCAAAGTCTTCAATATCC-

TCAGAGA). We digested the amplified products with DraI in the NEB Cutsmart buffer and re-

solved the fragments on a 3% agarose gel. The 180bp product is cleaved into fragments of 50

and 130bp if the *pol2-P301R* allele is present.

In both evolution experiments, the strains were propagated at 30°C with constant shak-

ing in sterile Nunc 96 deep-well storage blocks (Thermo Scientific, 260252), covered with an

AeraSeal (Excel Scientific) gas-permeable disposable plate sealer. Each well had 500 µls of

YPD and a single 3 mm glass bead to maintain cells in suspension. Sub-culturing steps were

performed by transferring 1 ul volumes from saturated cultures to 500 µls of fresh YPD using

an 8-channel pipettor. Frozen -80°C glycerol stocks of the cultures were made every three

transfers in 96-well format. To recover cultures from storage, the plates were first thawed at

room temperature and then 5 µls were transferred into 250 µl of fresh media.

Mutation rates. Mutation rates by fluctuation analysis were performed essentially as de-

scribed (Herr *et al.* 2014). To determine the mutation rate of the zygote colonies, 10 µls of the

original suspension (see above) were spot-plated directly on SC_MSG-Arg+CAN+NTC plates.

Another 10 µls were used for a series of 10-fold serial dilutions. With each dilution, 10 µls were

plated on the mutation rate plate and 10 µls on SC media to estimate the total number cells in

each colony (Nt). Only half of the zygote colonies had the *CAN1::natMX/can1Δ::HIS3* geno-

type suitable for mutation rate measurements. Thirty colonies with similar Nt values were used

as replica colonies to estimate the average number of mutational events (*m*) by maximum like-

lihood using the newton.LD.plating function in the R package rSalvador (Zheng 2015). The

confint.LD.plating function was used to calculate confidence intervals. For mutation rates of all

subclonal isolates (Figure 4A, S2), replica colonies were obtained by plating serial dilutions

onto SC media. For mutation rate calculations, we typically used mutation counts from 8 rep-

lica colonies. In some assays of T1 subclones in Figure 4A, 1 or 2 replica colonies had very

low mutation counts (0 or 1 mutants vs 100's). These clear outliers were censored under the

assumption that they represented an antimutator or polyploid subclone. Cohorts with more

than 2 such outliers were censored completely.

Flow Cytometry. Overnight cultures were diluted 1:50 in fresh YPD and grown for 3-5

hours. 100 µls of cells were recovered by centrifugation and washed twice in an equal volume

of water by resuspending them and centrifuging again. We then resuspended the cells in 70%

ethanol and incubated them at least 1 hour at room temperature or overnight at 4°C. We re-

covered the cells again by centrifugation, washed once with water, and then resuspended

them in 50 mM sodium citrate with RNaseA (10 µg/mL). The cells were heated to 95°C for 15

minutes and then incubated for 3 to 4 hours at 37°C. The cells were recovered by centrifuga-

tion and resuspended at a concentration of 10^6 cells/mL in 50 mM sodium citrate with 2 µM

Sytox Green (Life Technologies). After 1 hour at room temperature, the cells were sonicated

for 30 seconds with 1 second pulses in a Misonix cup horn sonicator and then analyzed on a

BD Accuri c6 flow cytometer using the FL1 detector. Plots in Dataset S1 are organized in the

96 well-format used for each experiment. Histogram plots of the data were generated using the

c6 software, which allows multiple plots to be overlaid. Plots used as controls are indicated at

the top of each experiment.

Genome Sequencing and Analysis. We performed whole genome sequencing of sub-

clones from the AH164 evolved cultures as described (Herr *et al.* 2014; Lee *et al.* 2019). We

used a custom sequencing analysis pipeline (eex_yeast_pileline.sh) that operates in a Unix

command-line. It aligns the sequencing reads against a repeat-masked version of the R64-1-1

assembly of the S288C *S. cerevisiae* genome using the Burrows-Wheeler Aligner (0.7.17).

Discordant and split reads were removed using Samblaster (0.1.24). Picard tools (2.21.9) Ad-

dorReplaceReadGroups was used to add information to the header needed for later steps. Af-

ter indexing the BAM files with Samtools (1.8), we sequentially processed them with the Ge-

nome Analysis Toolkit (GATK3) RealignerTargetCreator, IndelRealigner, LeftAlignIndels, Bas-

eRecalibrator, and PrintReads to minimize false variant calls. We used SamTools to make a

pileup file and then VarScan (v2.3.9) mpileup2snp and mpileup2indel to call SNPs and indels,

respectively. We used a variant frequency cutoff of 0.1 for SNPs to demonstrate that the se-

quenced genomes were indeed from 2n and not 4n cells (Dataset 2). We used a variant fre-

quency cutoff of 0.3 for indels to minimize false positives at repetitive sequences due to

frameshift errors during library preparation and filtered out SNPs and indels that were present

in the parental strains. For AH11304 evolved cultures we prepared DNA sequencing libraries

with the NEBNext Ultra II FS DNA library kit, which utilizes enzymatic fragmentation instead of

sonication. The same sequencing analysis pipeline was used except that during the variant

calls, we used a variant frequency cut-off of 0.22 for SNPs and 0.3 for indels.

We wrote a python script to compare variants found in subclones from the same culture

and create lists of shared and unique mutations found in each strain. The program only as-

sesses positions in the genome with 18-fold depth coverage in all subclones (Dataset S2). We

annotated the SNP lists using Annotate 0.1 (Pashkova *et al.* 2013) and the Indel lists with a

modified version of the program that reports the 3' peptide produced by the frameshift (anno-

tatefs2.py). We used the SNP lists to determine the 96-trinucleoide mutation spectra using

snv-spectrum (https://github.com/aroth85/snv-spectrum). We wrote a python script which cre-

ates a 2 x 96 table of the output of snv-spectrum for two different variant lists and then esti-

mates p-values of the hypergeometric test (Adams and Skopek 1987) and the cosine similarity

test, under the null hypothesis that they came from the same population.

Cosine similarity, commonly used in mutation signature analyses, is a measure of the

angle between two vectors projected into an n-dimensional space, where n is the number of

categories used for the comparison. In this case, the categories are the 96 trinucleotide con-

texts. The number of mutations at each context together define the direction of the vector. The

closer two spectra, the smaller the angle, and the closer the cosine similarity is to 1. The statis-

tical significance of such comparisons is typically not reported. However, the hypergeometric

test, which is an expansion of Fisher's exact test, has previously been used to test the null hy-

pothesis that two spectra are the same. Formally, the hypergeometric probability of each pos-

sible contingency table is calculated and the number of tables with a higher probability than the

observed table provides a measure of significance. With a 2 x 96 contingency table, an exact

calculation of significance becomes computationally impractical but can be estimated using a

Monte Carlo approach. Implementation of the test involves construction of 10,000 random 2 x

96 tables that maintain the same row and column totals as the original table (Agresti *et al.*

1979) and then calculation of the hypergeometric probability or cosine similarity of each table

(Dataset S2). The fraction of random tables with a hypergeometric probability higher than the

original table is an estimate for the probability that two spectra are the same. Likewise, the co-

sine similarity p-value represents the number of random tables whose cosine similarity was

higher than the observed cosine similarity. We used the conservative Bonferroni multiple test-

ing correction to estimate the *p*-value cut-off for significance for each set of comparisons (Da-

taset S2). Finally, we used the SNP lists described above to determine the dN/dS ratios of indi-

vidual genes and of the entire genome using the R-package *dNdscv* (Martincorena *et al.* 2017)

https://github.com/im3sanger/dndscv.

Acknowledgements

We thank Dr. Maitreya Dunham for use of her flow cytometer and horn sonicator. We

thank Dr. Iñigo Martincorena for help implementing *dNdscv*. We thank Drs. Dunham and Scott

Kennedy for critically reading the manuscript. We acknowledge funding (R01GM118854) from

the National Institute of General Medical Sciences (NIGMS). MBL was supported by the How-

ard Hughes Medical Institute (HHMI) Gilliam Fellowship for Advanced Study, the National Insti-

tutes of Health (NIH) Cellular and Molecular Biology training grant (NIH T32GM7270-39), and

the University of Washington Graduate Opportunities and Minority Achievement Program (UW

GO-MAP) Bank of America Fellowship. ITD was supported by the UW Molecular Medicine

Training Grant (NIH/NIGMS T32GM095421) and Genetic Approaches to Aging Training Grant

(NIH/NIA T32AG000057). The content is solely the responsibility of the authors and does not

necessarily represent the official views of the NIH, NIA, NIGMS or HHMI.

Competing Interests

716 The authors declare that there are no competing financial or non-financial interests.

Data Availability

Strains and plasmids are available upon request. All whole genome sequencing data is

available from the Sequence Read Archive (BioProject ID: PRJNA629499). Supplemental files

are available at FigShare. Dataset 1 contains flow cytometry results, mutation rates, and mod-

eling of diploid mutator evolution. Dataset 2 describes whole genome sequencing results of

diploid mutator strains. Datasets 1 and 2 can also be found at github

(https://github.com/mutatorUW/MutatorEvolution) along with all computer scripts used in this

work.

References

Adams, W. T., and T. R. Skopek, 1987 Statistical test for the comparison of samples from
727   mutational spectra. J Mol Biol **194:** 391-396.
728   Agresti, A., D. Wackerly and J. M. Boyett, 1979 Exact conditional tests for cross-classifications:
729   approximation of attained significance levels. Psychometrika **44:** 75-83.
730   Albertson, T. M., M. Ogawa, J. M. Bugni, L. E. Hays, Y. Chen *et al.,* 2009 DNA polymerase ε
731   and δ proofreading suppress discrete mutator and cancer phenotypes in mice. Proc Natl
732   Acad Sci U S A **106:** 17101-17104.
733   Alexandrov, L. B., J. Kim, N. J. Haradhvala, M. N. Huang, A. W. Tian Ng *et al.,* 2020 The
734   repertoire of mutational signatures in human cancer. Nature **578:** 94-101.
735   Barbari, S. R., D. P. Kane, E. A. Moore and P. V. Shcherbakova, 2018 Functional Analysis of
736   Cancer-Associated DNA Polymerase epsilon Variants in Saccharomyces cerevisiae. G3
737   (Bethesda) **8:** 1019-1029.
738   Bulock, C. R., X. Xing and P. V. Shcherbakova, 2020 DNA polymerase δ proofreads errors
739   made by DNA polymerase ε. Proceedings of the National Academy of Sciences **117:**
740   6035-6041.
741   Campbell, B. B., N. Light, D. Fabrizio, M. Zatzman, F. Fuligni *et al.,* 2017 Comprehensive
742   Analysis of Hypermutation in Human Cancer. Cell **171:** 1042-1056.e1010.
743   Cerami, E., J. Gao, U. Dogrusoz, B. E. Gross, S. O. Sumer *et al.,* 2012 The cBio Cancer
744   Genomics Portal: An Open Platform for Exploring Multidimensional Cancer Genomics
745   Data. Cancer Discovery **2:** 401-404.
746   Church, D. N., S. E. W. Briggs, C. Palles, E. Domingo, S. J. Kearsey *et al.,* 2013 DNA
747   polymerase ε and δ exonuclease domain mutations in endometrial cancer. Human
748   Molecular Genetics **22:** 2820-2828.
749   Coelho, M. C., R. M. Pinto and A. W. Murray, 2019 Heterozygous mutations cause genetic
750   instability in a yeast model of cancer evolution. Nature **566:** 275-278.
751   Datta, A., J. L. Schmeits, N. S. Amin, P. J. Lau, K. Myung *et al.,* 2000 Checkpoint-dependent
752   activation of mutagenic repair in *Saccharomyces cerevisiae pol3-01* mutants. Mol Cell
753   **6:** 593-603.
754   Dennis, D. G., J. McKay-Fleisch, K. Eitzen, I. Dowsett, S. R. Kennedy *et al.,* 2017 Normally
755   lethal amino acid substitutions suppress an ultramutator DNA Polymerase δ variant.
756   Scientific Reports **7:** 46535.
757   Fijalkowska, I. J., and R. M. Schaaper, 1995 Effects of *Escherichia coli dnaE* antimutator
758   alleles in a proofreading-deficient *mutD5* strain. J Bacteriol **177:** 5979-5986.

Fijalkowska, I. J., and R. M. Schaaper, 1996 Mutants in the Exo I motif of *Escherichia coli*
dnaQ: defective proofreading and inviability due to error catastrophe. Proc Natl Acad
Sci U S A **93:** 2856-2861.
Fisher, K. J., S. W. Buskirk, R. C. Vignogna, D. A. Marad and G. I. Lang, 2018 Adaptive
genome duplication affects patterns of molecular evolution in Saccharomyces
cerevisiae. PLoS Genetics **14:** e1007396.
Flood, C. L., G. P. Rodriguez, G. Bao, A. H. Shockley, Y. W. Kow *et al.*, 2015 Replicative DNA
Polymerase δ but Not ε Proofreads Errors in *Cis* and in *Trans*. PLoS Genet **11:**
e1005049.
Ganem, N. J., Z. Storchova and D. Pellman, 2007 Tetraploidy, aneuploidy and cancer. Current
Opinion in Genetics and Development **17:** 157-162.
Gerstein, A. C., H. J. Chun, A. Grant and S. P. Otto, 2006 Genomic convergence toward
diploidy in Saccharomyces cerevisiae. PLoS Genet **2:** e145.
Goldsby, R. E., N. A. Lawrence, L. E. Hays, E. A. Olmsted, X. Chen *et al.*, 2001 Defective DNA
polymerase-δ proofreading causes cancer susceptibility in mice. Nat Med **7:** 638-639.
Haber, J. E., 2012 Mating-type genes and MAT switching in Saccharomyces cerevisiae.
Genetics **191:** 33-64.
Haradhvala, N. J., J. Kim, Y. E. Maruvka, P. Polak, D. Rosebrock *et al.*, 2018 Distinct
mutational signatures characterize concurrent loss of polymerase proofreading and
mismatch repair. Nature Communications **9:** 1746.
Harari, Y., Y. Ram, N. Rappoport, L. Hadany and M. Kupiec, 2018 Spontaneous Changes in
Ploidy Are Common in Yeast. Current Biology **28:** 825-835 e824.
Herr, A. J., S. R. Kennedy, G. M. Knowels, E. M. Schultz and B. D. Preston, 2014 DNA
replication error-induced extinction of diploid yeast. Genetics **196:** 677-691.
Herr, A. J., M. Ogawa, N. A. Lawrence, L. N. Williams, J. M. Eggington *et al.*, 2011a Mutator
suppression and escape from replication error–induced extinction in yeast. PLoS Genet
7: e1002282.
Herr, A. J., L. N. Williams and B. D. Preston, 2011b Antimutator variants of DNA polymerases.
Critical Reviews in Biochemistry and Molecular Biology **46:** 548-570.
Hodel, K. P., R. de Borja, E. E. Henninger, B. B. Campbell, N. Ungerleider *et al.*, 2018
Explosive mutation accumulation triggered by heterozygous human Pol ε proofreading-
deficiency is driven by suppression of mismatch repair. eLife **7:** e32692.
Kaiser, C., Michaelis, S., Mitchell, A., 1994 *Methods in Yeast Genetics: A Laboratory Course*
Manual. Cold Spring Harbor Laboratory Press.
Kandoth, C., N. Schultz, A. D. Cherniack, R. Akbani, Y. Liu *et al.*, 2013 Integrated genomic
characterization of endometrial carcinoma. Nature **497:** 67-73.
Kane, D. P., and P. V. Shcherbakova, 2014 A common cancer-associated DNA polymerase ε
mutation causes an exceptionally strong mutator phenotype, indicating fidelity defects
distinct from loss of proofreading. Cancer Research.
Kennedy, S. R., E. M. Schultz, T. M. Chappell, B. Kohrn, G. M. Knowels *et al.*, 2015 Volatility
of Mutator Phenotypes at Single Cell Resolution. PLoS Genet **11:** e1005151.
Lee, M. B., I. T. Dowsett, D. T. Carr, B. M. Wasko, S. G. Stanton *et al.*, 2019 Defining the
impact of mutation accumulation on replicative lifespan in yeast using cancer-
associated mutator phenotypes. Proc Natl Acad Sci U S A **116:** 3062-3071.

Li, H.-D., I. Cuevas, M. Zhang, C. Lu, M. M. Alam *et al.*, 2018 Polymerase-mediated
ultramutagenesis in mice produces diverse cancers with high mutational load. The
Journal of Clinical Investigation **128:** 4179-4191.
Loeb, L. A., 2016 Human Cancers Express a Mutator Phenotype: Hypothesis, Origin, and
Consequences. Cancer Research **76:** 2057-2059.
Loeb, L. A., C. F. Springgate and N. Battula, 1974 Errors in DNA replication as a basis of
malignant changes. Cancer Res **34:** 2311-2321.
Lynch, H. T., P. M. Lynch, S. J. Lanspa, C. L. Snyder, J. F. Lynch *et al.*, 2009 Review of the
Lynch syndrome: history, molecular genetics, screening, differential diagnosis, and
medicolegal ramifications. Clinical Genetics **76:** 1-18.
Mable, B. K., and S. P. Otto, 2001 Masking and purging mutations following EMS treatment in
haploid, diploid and tetraploid yeast (Saccharomyces cerevisiae). Genetical Research
77: 9-26.
Martincorena, I., K. M. Raine, M. Gerstung, K. J. Dawson, K. Haase *et al.*, 2017 Universal
Patterns of Selection in Cancer and Somatic Tissues. Cell **171:** 1029-1041.e1021.
Mertz, T. M., S. Sharma, A. Chabes and P. V. Shcherbakova, 2015 Colon cancer-associated
mutator DNA polymerase δ variant causes expansion of dNTP pools increasing its own
infidelity. Proceedings of the National Academy of Sciences of the United States of
America.
Morrison, A., A. L. Johnson, L. H. Johnston and A. Sugino, 1993 Pathway correcting DNA
replication errors in *Saccharomyces cerevisiae*. EMBO J **12:** 1467-1473.
Morrison, A., and A. Sugino, 1994 The 3´→5´ exonucleases of both DNA polymerases δ and ε
participate in correcting errors of DNA replication in *Saccharomyces cerevisiae*. Mol
Gen Genet **242:** 289-296.
Network, C. G. A., 2012 Comprehensive molecular characterization of human colon and rectal
cancer. Nature **487:** 330-337.
Palles, C., J. B. Cazier, K. M. Howarth, E. Domingo, A. M. Jones *et al.*, 2012 Germline
mutations affecting the proofreading domains of POLE and POLD1 predispose to
colorectal adenomas and carcinomas. Nature Genetics **45:** 136-144.
Parkash, V., Y. Kulkarni, J. ter Beek, P. V. Shcherbakova, S. C. L. Kamerlin *et al.*, 2019
Structural consequence of the most frequently recurring cancer-associated substitution
in DNA polymerase ε. Nature Communications **10:** 373.
Pashkova, N., L. Gakhar, S. C. Winistorfer, A. B. Sunshine, M. Rich *et al.*, 2013 The yeast Alix
homolog Bro1 functions as a ubiquitin receptor for protein sorting into multivesicular
endosomes. Dev Cell **25:** 520-533.
Reha-Krantz, L. J., 1988 Amino acid changes coded by bacteriophage T4 DNA polymerase
mutator mutants. Relating structure to function. J Mol Biol **202:** 711-724.
Reha-Krantz, L. J., M. S. P. Siddique, K. Murphy, A. Tam, M. O'Carroll *et al.*, 2011 Drug
sensitive DNA polymerase δ reveals a role for mismatch repair in checkpoint activation
in yeast. Genetics **189:** 1211-1224.
Schaaper, R. M., 1993 Base selection, proofreading, and mismatch repair during DNA
replication in *Escherichia coli*. J Biol Chem **268:** 23762-23765.
Sherman, F., 2002 Getting started with yeast, pp. 3-41 in *Part B: Guide to Yeast Genetics and*
Molecular and Cell Biology, edited by C. Guthrie and G. R. Fink. Academic Press, San
Diego.

Shlien, A., B. B. Campbell, R. de Borja, L. B. Alexandrov, D. Merico *et al.*, 2015 Combined
hereditary and somatic mutations of replication error repair genes result in rapid onset
of ultra-hypermutated cancers. Nature Genetics **47:** 257-262.
Storchova, Z., A. Breneman, J. Cande, J. Dunn, K. Burbank *et al.*, 2006 Genome-wide genetic
analysis of polyploidy in yeast. Nature **443:** 541-547.
Strathern, J., J. Hicks and I. Herskowitz, 1981 Control of cell type in yeast by the mating type
locus: The α1-α2 hypothesis. Journal of Molecular Biology **147:** 357-372.
Swan, M. K., R. E. Johnson, L. Prakash, S. Prakash and A. K. Aggarwal, 2009 Structural basis
of high-fidelity DNA synthesis by yeast DNA polymerase δ. Nat Struct Mol Biol **16:** 979-
986.
Venkataram, S., B. Dunn, Y. Li, A. Agarwala, J. Chang *et al.*, 2016 Development of a
Comprehensive Genotype-to-Fitness Map of Adaptation-Driving Mutations in Yeast. Cell
166: 1585-1596.e1522.
Williams, L. N., A. J. Herr and B. D. Preston, 2013 Emergence of DNA polymerase ε
antimutators that escape error-Induced extinction in yeast. Genetics **193:** 751-770.
Williams, L. N., L. Marjavaara, G. M. Knowels, E. M. Schultz, E. J. Fox *et al.*, 2015 dNTP pool
levels modulate mutator phenotypes of error-prone DNA polymerase ε variants.
Proceedings of the National Academy of Sciences of the United States of America.
Xing, X., D. P. Kane, C. R. Bulock, E. A. Moore, S. Sharma *et al.*, 2019 A recurrent cancer-
associated substitution in DNA polymerase ε produces a hyperactive enzyme. Nature
Communications **10:** 374.
Yoshida, R., K. Miyashita, M. Inoue, A. Shimamoto, Z. Yan *et al.*, 2011 Concurrent genetic
alterations in DNA polymerase proofreading and mismatch repair in human colorectal
cancer. European Journal of Human Genetics **19:** 320-325.
Zheng, Q., 2015 A new practical guide to the Luria–Delbrück protocol. Mutation
Research/Fundamental and Molecular Mechanisms of Mutagenesis **781:** 7-13.

875
876

Figure 1. Spontaneous polyploids routinely emerge during the evolution of *pol2-4*

***msh2Δ* haploids.** A) Isolation of *pol2-4 msh2Δ* haploids from sporulated parental diploid cells

(2n). Diploid genotype indicated to the right. Arrows, from left to right, indicate manipulations of

cells: sporulation, tetrad dissection, and inoculation into 96-well culture blocks. Following tetrad

dissection, double mutant *pol2-4 msh2Δ* spores (blue cells) selectively form colonies on media

lacking uracil and leucine (green box). B) Evolution of *pol2-4 msh2Δ* mutators. Independent

isolates were cultured in 96-well format by diluting 1:500 with each transfer (T). Ploidy was as-

sessed by flow cytometry at T1, T4, and T7 (see Dataset S1 for histograms). Color of boxes

indicate the ploidy or mixture of ploidies of each culture (see top for key). C) Ploidy of single

cell subclones isolated from T7. D) Mating-type of single cell subclones. *Top*, Schematic of

*MAT***a** and *MAT*α mating type loci, PCR primers (arrows), and expected DNA sizes. *Bottom*,

PCR products run on 2% agarose gels, labeled by their coordinates in the 96-well block; *-ve*,

negative (no cells) controls; *cn,* culture censored during evolution. The 500bp fragment of the

Invitrogen 1kb plus ladder is indicated to the left.

Figure 2. Competition between diploid mutator parents and tetraploid offspring. A) Experimental design. *MATα/MATα* and *MATa/MATa* subclones from the *pol2-4 msh2Δ* evolved cultures (see Figure 1B) were mated. Cell suspensions of the resulting tetraploid and parental diploid colonies were counted, mixed and then competed through 8 serial transfers. B) Ploidy measurements after the first (T1) and eighth (T8) serial transfers (see Dataset 1 for histograms). Letter and numbers to the left indicate location in the 96 well culture block described in Figure.1. Color of boxes indicate the ploidy or mixture of ploidies of each culture (see top).

Figure 3. Strong diploid mutators maintain diploidy during evolution. A) Isolation of *pol3-01/pol3-01 msh6Δ/msh6Δ* zygotes. *Top,* Double mutant *pol3-01 msh6Δ* haploids, identified by dissecting tetrads on selective media (Sc-Ura-Leu), were mated to form zygotes. Zygotes were then moved to new locations on the plate to form colonies. *Bottom,* Growth phenotypes of the initial zygote colonies. Colonies were picked from left to right. Those used for evolution are indicated by either their 96-well coordinate or a dash (-). Colonies without a label or dash were not evolved. B) Ploidy of evolved cultures after one (T1), ten (T10), nineteen (T19), and twenty-seven (T27) serial transfers (see Dataset S1 for histograms). Color of boxes indicate the ploidy or mixture of ploidies of each culture as depicted in the key at the top.

Figure 4. Strong diploid mutators adapt through antimutator alleles. A) Mutation rates of

pol3-01/pol3-01 msh6Δ/msh6Δ subclones. Single cell subclones were isolated after the first

(T1), thirteenth (T13), and twenty-fifth (T25) transfers. Mutation rates were determined by a ca-

navanine resistance (Can^R) fluctuation assay (see Dataset S1 for mutation rates and 95% con-

fidence intervals). Solid circles, mutation rates; transparent circles, 95% confidence intervals.

B) Variation in mutation rates depicted in (A) of T25 subclones from the same culture. Alternat-

ing shades of red indicate groups of subclones. C) Phylogenetic trees of subclones depicted in

(B) based on whole genome sequencing. Observed *pol3* mutations are in blue lettering using

single letter amino acid codes: *, nonsense mutation; bold type, known or likely antimutators).

D) Representative mutation spectra of shared (trunk) and unique mutations (see Figure S7 for

mutation spectra of all trees in (C)). Each mutation site (represented here by the pyrimidine

base T or C) was categorized into one of the 96 mutation types created by subdividing the 6

general mutation types (C→A, C→G, C→T, T→A, T→C, T→G) into the 16 possible tri-nucleo-

tide contexts (see boxes to the lower right for contexts; blue, black, and red bars correspond to

mutation sites involving C; gray, green and pink bars correspond to mutation sites involving T).

Bar height corresponds to the fraction each mutation type represents of the total mutations

(given in the right-hand corner of each plot). Inset table: *p*-values of the hypergeometric test of

similarity between mutation spectra. E) Genome-wide (global) dN/dS ratios of trunk and branch

mutations.

Figure 5. Modeling adaptation of POLE-P286R MMR-deficient cancers to *EEX*. A) Isola-

tion of *pol2-P301R/POL2 msh2Δ/msh2Δ* zygotes. *POL2 msh2Δ* spores slowly divide once on

SC-Ura-Leu media, allowing mating with more rapidly dividing *pol2-P301R msh2Δ* cells. B)

Representative growth phenotypes of the initial zygote colonies (see Figure S8 for all colo-

nies). C) Ploidy of evolved cultures after ten (T10) serial transfers. Box color indicates the

ploidy or mixture of ploidies of each culture as depicted in the key to the right (see Dataset S1

for histograms). D) Mutation rates as determined by a canavanine resistance (Can^R) fluctua-

tion assay (see Dataset S1 for mutation rates and 95% confidence intervals). Solid circles, mu-

tation rates; transparent circles, 95% confidence intervals (for Zygote Clones only). E) Phylo-

genetic trees based on whole genome sequencing of the subclones. Line length, the number

of shared or unique SNVs; lettering at the bottom of each tree, culture position in 96 well plate;

lettering at the end of each branch, subclone ID number; *pol2* mutations are in blue lettering:

fs, frameshift; *, nonsense mutation. F) Spectra of grouped mutations from trunk and branches.

Format is the same as in Figure 4. Trunk spectrum determined from all trunk mutations com-

bined. Branch spectrum determined from combined mutations from branches with truncating

mutations at E945 or before (A1-3, A5-1, A5-2, A5-3, A8-1, A12-2, B3-3, C11-1, F1-2, F1-3,

F12-2, G6-2, G7-1, H2-2, H2-3, and H3-1/H3-2). G) Oncoprint of POLE-P286R tumors on cBi-

oPortal showing presence of additional *POLE* alleles and mutations affecting MMR compo-

nents (see Dataset 1).

 Supplementary Information

Table S1: Similarity of the *pol2/POL2 msh2Δ/msh2Δ* mutation spectra to mutation signatures in cancer.

Signature	Cosine Similarity
SBS1	0.138
SBS2	0.178
SBS3	0.485
SBS4	0.483
SBS5	0.537
SBS6	0.239
SBS7a	0.191
SBS7b	0.209
SBS7c	0.175
SBS7d	0.218
SBS8	0.519
SBS9	0.456
SBS10a	0.634
SBS10b	0.149
SBS11	0.344
SBS12	0.233
SBS13	0.070
SBS14	0.793
SBS15	0.217
SBS16	0.276
SBS17a	0.053
SBS17b	0.092
SBS18	0.626
SBS19	0.301
SBS20	0.536
SBS21	0.238
SBS22	0.046
SBS23	0.299
SBS24	0.333
SBS25	0.447
SBS26	0.236
SBS27	0.140
SBS28	0.159
SBS29	0.472
SBS30	0.340
SBS31	0.258
SBS32	0.501
SBS33	0.149
SBS34	0.075
SBS35	0.424
SBS36	0.701
SBS37	0.276
SBS38	0.317
SBS39	0.241
SBS40	0.623
SBS41	0.304
SBS42	0.453
SBS43	0.052
SBS44	0.561
SBS45	0.396
SBS46	0.266
SBS47	0.201
SBS48	0.026
SBS49	0.071
SBS50	0.282
SBS51	0.212
SBS52	0.180
SBS53	0.128
SBS54	0.144
SBS55	0.082
SBS56	0.653
SBS57	0.370
SBS58	0.375
SBS59	0.322
SBS60	0.025

Figure S1. Discovery of spontaneous polyploid *eex* mutants. A) Synthetic growth pheno-
types of freshly dissected *pol3-01 msh6Δ* and *pol2-4 msh2Δ* spore clones. *Top,* Schematic of
strain genotypes and tetrad dissections. *Bottom,* Images of tetrad dissections after 48 hours of
growth. Colored circles indicate genotypes. B) Isolation of *eex* mutants and evidence for poly-
ploidization. *Top,* Scheme for isolating *eex* mutants and the generation of mapping strains.
Bottom, Flow cytometry traces of two representative polyploid strains (AH13214 and AH13238;
see Dataset 1) and 1n, 2n, and 4n controls as well as tally of the ploidies of the mapping
strains.

Figure S2. Polyploids and antimutator subclones emerge with similar frequencies during the evolution of *pol2-4 msh2Δ* cultures. A) Representative images of colonies. *Clockwise from the top left:* initial colonies after 24 hours, A4 (a mixture of 1n and 3n cells) is circled in yellow; close-up of A4 just after removing single cells from the perimeter (holes were punched into the agar to mark origin of cells 1 through 4); colony forming capacity of single cells (A4.1 is circled in yellow). B) Ploidy of colonies derived from single-cell clones (see Dataset S1 for histograms), named for the coordinate of the parent strain in the evolved 96-well plate and the isolate number. C) Mutation rate measurements of haploid cultures as determined by fluctuation analysis of canavanine resistance (Can^R). Error bars represent 95% confidence intervals. (See Dataset S1 for Table of values.)

Figure S3. Test for mating-type switching and same-sex mating between mutator cells.
A) Experimental design. Same sex haploid cultures with opposite *CAN1* alleles (*CAN1::natMX*, *can1Δ::HIS3)* were mixed and plated for *CAN1::natMX/can1Δ::HIS3* diploids. B) Summary of flow cytometry and mating-type PCR assays of colonies resulting from the crosses indicated in the white boxes. The mutator cultures used for each cross are indicated by their 96-well coordinate (see Figure 1). The color of the box indicates ploidy. Blue 1n boxes are haploids included as controls (see Dataset S1 for individual histograms). Results of Mating type PCR: **α**, *MATα*; **a**, *MATa* C) Mating-type PCR gels.

989
990

Figure S4. Frequency of mating-type switching and same-sex mating between wild-type
cells. A) Experimental design. Same sex haploid colonies with opposite *CAN1* alleles
(*CAN1::natMX*, *can1Δ::HIS3)* were mixed, incubated in media, and plated for
CAN1::natMX/can1Δ::HIS3 diploids. B) Frequencies of His$^+$ NTC^R colonies per 10^7 cells. C)
Summary of flow cytometry and mating-type PCR assays of colonies resulting from MATα x
MATα matings. The color of the box indicates ploidy (see Dataset S1 for individual histo-
grams). D) Summary of flow cytometry and mating-type PCR assays of colonies resulting from
MAT**a** x MAT**a** matings. AH4002, AH4003, AH4010 represent haploid strains; 2 x 15 and 3 x
10 are MAT**a**/α control diploids derived from matings between AH4002 x AH4015 and AH4003
x AH4010.

1002

1004

Figure S5. Simulation of *eex* adaption during diploid mutator evolution. Exponential growth of *pol3-01/pol3-01 msh6Δ/msh6Δ* cells (Mutator, red line) and an *eex* mutant (blue line) with a 10-fold lower mutation rate were modeled assuming the only limiting factor for growth was homozygous inactivation of essential genes (see Dataset S1 for calculations). Each peak models a 1:500 dilution of the culture from ~10^8 cells to ~10^5 cells.

1011

1012 **Figure S6. Colony forming capacity of evolved *pol3-01/pol3-01 msh6Δ/msh6Δ* cultures.**
1013 Ten-fold serial dilutions from overnight cultures were plated on SC plates and incubated for
1014 two days at 30°C. T1 and T13 refer to first and thirteenth sub-culturing steps after the initial in-
1015 oculations. The culture designation on the left refers to the position of the culture in the 96 well
1016 plate (see Figure 3).

1017

Figure S7. Mutation spectra and phylogenetic relationships of evolved *pol3-01/pol3-01*

***msh6Δ/msh6Δ* T25 subclones.** Mutational variants identified by whole genome sequencing of

subclones isolated from different evolved cultures were classified as shared (trunk) and

unique. The nucleotide context 5' and 3' of each mutational site (represented as the pyrimidine

base T or C) were used to classify the 6 mutation types (C:G→A:T, C:G→G:C, C:G→T:A,

T:A→A:T, T:A→C:G, T:A→G:C) into 96 trinucleotide contexts. Each mutation type family is

color-coded (see Key for color and context order). The bar height for mutation type represents

the fraction of the total mutations (to the right of each plot). Phylogenetic trees: line length cor-

responds to the number of mutations, line color corresponds to mutation spectra, observed

pol3 mutations are in blue lettering (bold represents known or likely antimutators). Inset tables:

p-values determined by the hypergeometric test.

1031

1032

Figure S8. Growth phenotypes of *pol2-P301R/POL2 msh2Δ/msh2Δ* and *pol2-P301R/pol2-*

***P301R msh2Δ/msh2Δ* zygotes.** Zygotes colonies were isolated, as described in Figure 5, on

synthetic complete media lacking uracil and leucine (SC-Ura-Leu) by mating cells dissected

from tetrads of AH11304 (*URA3::pol2-P301R/POL2 msh2Δ::LEU2/MSH2*). Each colony used

for the evolution experiment in Figure 5 is labeled to the left by the 96-well coordinate position.

Small, unlabeled colonies are derived from *pol2-P301R/pol2-P301R msh2Δ/msh2Δ* zygotes

and were not evolved.

1040

Figure S9. Locations of amino acid substitutions encoded by *pol2-P301R* mutator suppressors. A) Two orientations of the wild-type *S. cerevisiae* Polε structure (Protein database accession code 4M80 (Swan *et al.* 2009)) are shown as ribbon diagrams of the α-carbon backbone (rendered in PyMol). Structural domains are color coded as follows: Amino (gray), Exo (red), processivity (P) (chartreuse), Palm (purple), Fingers (blue) and Thumb (green). Yellow spheres correspond to amino acid substitutions encoded by mutator suppressors. The incoming dNTP is denoted by green CPK sticks. The primer DNA is represented by tan sticks and the template DNA, by brown sticks. Active-site carboxylate side chains are gray CPK sticks coming out of the purple (palm) or red (exo) ribbons. L160R, S162F, and A202V affect interacting residues that play a structural role in the amino terminal domain in a region not previously implicated in fidelity. B) Locations of *eex* substitutions within the Exo domain of the Pol2p-P301R mutant polymerase (Protein database accession code 6i8a (Parkash *et al.* 2019)), which may be candidates for enhancing proofreading function. C) Substitutions positioned to influence dNTP binding and the KKRYA Motif. D640 is 3.21 Å and 3.46 Å from the γ and α-phosphates of the incoming dNTP, respectively. K966, K967(K), and R968(R) from the KKRYA motif, which monitors the minor groove for non-Watson-Crick base-pairing, are depicted as

1058   purple CPK space-filling spheres. We previously isolated an antimutator allele affecting K966

1059   (*pol2-K966Q*) (Williams *et al.* 2013). P956S is 3.23 Å from the primary amino group of R968.

1060

1061

Dataset S1. Flow cytometry, mutation rates, and modeling of diploid evolution.	
Page	*Description*
1	Ploidy measurements of *eex* mapping strains described in S1B Fig.
2	Ploidy measurements of *pol2-4 msh6Δ* evolved cultures described in Fig 1B.
3	Ploidy measurements of *pol2-4 msh6Δ* T8 subclones described in Fig 1C.
4	Ploidy measurements of *pol2-4 msh6Δ* subclones described in S2A Fig.
5	Mutation rate measurements of *pol2-4 msh6Δ* subclones described in S2B Fig.
6	Ploidy measurements of strains isolated by same sex mating between T1 *pol2-4 msh6Δ* cultures described in S3 Fig.
7	Ploidy measurements of competition experiment between *pol2-4 msh6Δ* derived diploid and tetraploid strains described in Fig 2B.
8	Simulation of evolution of *pol3-01/pol3-01 msh6Δ/msh6Δ* evolved cultures described in S4 Fig.
9	Ploidy measurements of *pol3-01/pol3-01 msh6Δ/msh6Δ* evolved cultures described in Fig 3B.
10	Ploidy measurments of T25 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones analyzed in Fig 4
11	Mutation rate measurements of T25 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones described in Fig 4.
12	Ploidy measurements of pol2-P301R/POL2 msh2Δ/msh2Δ evolved cultures described in Fig.5C.
13	Mutation rate measurements of T10 pol2-P301R/POL2 msh2Δ/msh2Δ subclones described in Fig. 5D and pol2 changes in Fig.5E.
14	POLE-P286R cancers on cBioPortal depicted in Fig. 5G.

1062

Dataset S2. Whole genome sequencing.	
Page	*Description*
1	AH164_B1_SNPs: Sequencing coverage, evolutionary relationships, distribution of variant allele frequencies (VAF), and annotation of SNPs for AH164_B1 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
2	AH164_B1_Indels: Sequencing coverage, evolutionary relationships, and annotation of Indels for AH164_B1 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
3	AH164_B2_SNPs: Sequencing coverage, evolutionary relationships, distribution of VAFs, and annotation of SNPs for AH164_B2 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
4	AH164_B2_Indels: Sequencing coverage, evolutionary relationships, and annotation of Indels for AH164_B2 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.

5	AH164_C2_SNPs: Sequencing coverage, evolutionary relationships, distribution of VAFs, and annotation of SNPs for AH164_C2 p*ol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
6	AH164_C2_Indels: Sequencing coverage, evolutionary relationships, and annotation of Indels for AH164_C2 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
7	AH164_D2_SNPs: Sequencing coverage, evolutionary relationships, distribution of VAFs, and annotation of SNPs for AH164_D2 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
8	AH164_D2_Indels: Sequencing coverage, evolutionary relationships, and annotation of Indels for AH164_D2 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
9	AH164_G5_SNPs: Sequencing coverage, evolutionary relationships, distribution of VAFs, and annotation of SNPs for AH164_G5 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
10	AH164_G5_Indels: Sequencing coverage, evolutionary relationships, and annotation of Indels for AH164_G5 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
11	AH164_A6_SNPs: Sequencing coverage, evolutionary relationships, distribution of VAFs, and annotation of SNPs for AH164_A6 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
12	AH164_A6_Indels: Sequencing coverage, evolutionary relationships, and annotation of Indels for AH164_A6 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
13	AH164_B6_SNPs: Sequencing coverage, evolutionary relationships, distribution of VAFs, and annotation of SNPs for AH164_B6 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
14	AH164_B6_Indels: Sequencing coverage, evolutionary relationships, and annotation of Indels for AH164_B6 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
15	AH164_C7_SNPs: Sequencing coverage, evolutionary relationships, distribution of VAFs, and annotation of SNPs for AH164_C7 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
16	AH164_C7_Indels: Sequencing coverage, evolutionary relationships, and annotation of Indels for AH164_C7 *pol3-01/pol3-01 msh6Δ/msh6Δ* subclones.
17	Table of cosine similarity and hypergeometric tests of similarity between *pol3-01/pol3-01 msh6Δ/msh6Δ* mutation spectra.
18	Output of dNdScv analysis of AH164 evolved cultures reported in Figure 4E.
19	Evolutionary relationships, variant counts, and annotated *pol2* variants for *pol2-P301R/POL2 msh2Δ/msh2Δ* evolved strains used for Figure 5E.
20	Annotations of all variants in *pol2-P301R/POL2 msh2Δ/msh2Δ* evolved strains described in Figure 5.
21	Table of cosine similarity and hypergeometric tests of similarity between *pol2-P301R/POL2 msh2Δ/msh2Δ* mutation spectra.

1063