Developmental Genome-Wide DNA Methylation Asymmetry Between

Mouse Placenta and Embryo

Legault LM[1,2], Doiron K[1], Lemieux A[1,2], Caron M[1,3], Chan D[3], Lopes FL[4], Bourque G[5,6,7],
Sinnett D[1,8], McGraw S[1,2,9,#].

[1]Research Center of the CHU Sainte-Justine, Montreal, Canada.
[2]Department of Biochemistry and Molecular Medicine, Université de Montréal, Montreal,
Canada.
[3]Research Institute of the McGill University Health Centre, Montreal, Quebec, Canada
[4]São Paulo State University (Unesp), School of Veterinary Medicine, Aracatuba, Brazil
[5]Department of Human Genetics, McGill University, Montreal, Quebec, Canada
[6]McGill University and Genome Quebec Innovation Centre, Montreal, Quebec, Canada
[7]Canadian Center for Computational Genomics, Montreal, Quebec, Canada
[8]Department of Pediatrics, Université de Montréal, Montreal, Canada.
[9]Department of Obstetrics and Gynecology, Université de Montréal, Montreal, Canada.
17
18 [#]Corresponding author: serge.mcgraw@umontreal.ca

19
20
21 **Running title:** Embryonic & Placental DNAme Asymmetry

23 **Summary statement:** The kinetics of asymmetric DNA methylation acquisition of embryo-
24 placenta differentially methylated regions is not a stepwise process occurring in embryonic and
25 extraembryonic lineages, but a prompt progression in the early post-implanted conceptus.
26

27   **ABSTRACT**
28
In early embryos, DNA methylation is remodelled to initiate the developmental program but for
mostly unknown reasons, methylation marks are acquired unequally between embryonic and
placental cells. To better understand this, we generated high-resolution DNA methylation maps
of mouse mid-gestation (E10.5) embryo and placenta. We uncovered specific subtypes of
differentially methylated regions (DMRs) that contribute directly to the developmental
asymmetry existing between mid-gestation embryonic and placental DNA methylation patterns.
We show that the asymmetry occurs rapidly during the acquisition of marks in the post-
implanted conceptus (E3.5-E6.5), and that these patterns are long-lasting across subtypes of
DMRs throughout prenatal development and in somatic tissues. We reveal that at the peri-
implantation stages, the *de novo* methyltransferase activity of DNMT3B is the main driver of
methylation marks on asymmetric DMRs, and that DNMT3B can largely compensate for lack of
DNMT3A in the epiblast and extraembryonic ectoderm, whereas DNMT3A can only partially
palliate in the absence of DNMT3B. However, as development progresses and as DNMT3A
becomes the principal *de novo* methyltransferase, the compensatory DNA methylation
mechanism of DNMT3B on DMRs becomes less effective.

INTRODUCTION

Throughout the eutherian mammalian gestation, the placenta plays an essential role in mediating maternal–embryonic exchanges of gas, nutrients and waste, and also provides the developing embryo with a protective layer against adverse environmental exposures and the maternal immune system (Rossant and Cross, 2001). These unique placental functions are orchestrated by several distinct trophoblast cell subtypes organized in separate layers (Cross, 2000). The initial steps of lineage specialization of both placental and embryonic cells occur promptly following fertilization during the first few embryonic cleavages as DNA methylation marks are being reprogrammed (Morgan et al., 2005).

DNA methylation is an epigenetic mechanism that is critical in the determination of lineage-specific differentiation and development, and is mainly recognized for its involvement in processes such as transcriptional repression, genomic imprinting and X-inactivation (Bestor, 2000). DNA methylation marks are mediated by the action of DNA methyltransferases (DNMTs). Establishment of new or *de novo* DNA methylation patterns required for cell lineage determination during development is mediated by DNMT3A and DNMT3B, with cofactor DNMT3L, (Okano et al., 1999, Li, 2002), whereas DNMT1 maintains heritable DNA methylation patterns during cellular divisions (Leonhardt et al., 1992, Lei et al., 1996). These enzymes are critical, as deletion of *Dnmt3b* or *Dnmt1* is embryonic lethal, while *Dnmt3a*-deficient offspring die shortly after birth (Li et al., 1992, Okano et al., 1999). During gametogenesis, the acquisition of genome-wide and allele-specific methylation patterns (i.e. genomic imprinting) in both oocytes and sperm is essentially due to the activity of DNMT3A (Kaneda et al., 2004, Kato et al., 2007). Following fertilization, a reprogramming wave removes most methylation signatures across the genome, except for imprinted regions, some types of repeat sequences, as well as imprinted-like sequences, to trigger the developmental program (Hirasawa et al., 2008, Howell et al., 2001, McGraw et al., 2015). Then, during the peri-implantation process, DNA methylation profiles are re-acquired in a sex-, cell- and tissue-specific manner across most parts of the genome by the combined action of DNMT3A and DNMT3B. In the early stages of the *de novo* methylation wave (E4.5-E7.5), the expression of *Dnmt3b* is more robust than *Dnmt3a* in the epiblast and embryonic-derived cells (Watanabe et al., 2002, Auclair et al., 2014, Smith et al., 2017), with the relative expression of *Dnmt3b* and *Dnmt3a* being considerably reduced in the extraembryonic ectoderm (ExE) and trophoblast

lineages (Smith et al., 2017, Senner et al., 2012). This discrepancy in *Dnmt3a* and *Dnmt3b*
expression levels coincides with the initiation of divergent DNA methylation acquisition
between the trophoblast and the inner cell mass of the blastocyst (Monk, 1987, Nakanishi et al.,
2012, Oda et al., 2013, Santos et al., 2002, Fulka et al., 2004, Guo et al., 2014, Smith et al.,
2014), a difference that becomes extremely apparent by E6.5, as the epiblast has acquired most
of its global DNA methylation compared to the lower-methylation state of the ExE (Auclair et
al., 2014, Smith et al., 2017). This divergence is a common feature across mammalian placenta,
as a heterogeneous and lower-methylation state compared to somatic tissues and other cell types
is constantly observed (Smith et al., 2017, Decato et al., 2017, Schroeder et al., 2013, Chatterjee
et al., 2016).

Although the functional role of reduced methylation levels observed across the placental
genome is still not fully understood, studies suggest that it may activate transposable elements
that are typically silenced in other tissues (Chuong, 2013). DNA methylation plays an important
role in suppressing retrotransposons in mammalian cells, for which the activity has been
associated with genomic instability and disease development (Church et al., 2009, Slotkin and
Martienssen, 2007). Following the *de novo* methylation wave, in embryonic-derived cells from
the inner cell mass, transposable elements acquire higher levels of DNA methylation causing
transcriptional silencing, whereas in the trophectoderm-derived cells that will form the placenta,
these transposable elements are maintained in a relaxed methylation state and preferentially
expressed (Price et al., 2012, Okahara et al., 2004, Warren et al., 2015). The low-methylation
levels on these elements contributed to the evolution and diversification of the placenta function
through the regulation of gene expression by providing placenta-specific enhancers, cryptic
promoters and other cis-regulatory elements (Haig, 2012, Mi et al., 2000, Cohen et al., 2011, Xie
et al., 2013, Macaulay et al., 2011, Emera and Wagner, 2012).

Despite the evident distinction between embryonic and placental DNA methylation
levels, it remains unclear how, when and where methylation levels are acquired unequally across
these genomes. To better understand the developmental dynamics of epigenetic asymmetry that
exists between embryo and placenta, we first generated high-resolution maps of DNA
methylation marks using Reduced Representation Bisulfite Sequencing (RRBS) at mid-gestation,
when the mouse placenta is first considered mature (E10.5), to identify embryo-placenta
differentially methylated regions (DMRs). Then, using publicly available DNA methylation data

sets and computational analyses, we defined how various categories of DMRs are established in
early stages and maintained throughout development. In addition, we outlined the contribution of
Dnmt3a and *Dnmt3b* in the acquisition and maintenance of embryonic and extraembryonic
specific DMR patterns.
110

METHODS
Animals and Sample Collection
Female C57BL/6 (8-10 week-old) were purchased from Harlan Sprague-Dawley
Laboratories (Indianapolis, IN) and were mated with male C57BL/6 (4 months of age).
Following natural mating, embryos and placental tissues were collected at E10.5 (presence of
vaginal plug at E0.5). Samples were frozen immediately in liquid nitrogen and stored at -80 °C
until analyzed.
118

DNA Methylation Analyses
RRBS libraries were generated as published protocols (Boyle et al., 2012, Gu et al., 2011)
with our specifications (Magnus et al., 2014, McGraw et al., 2015, Legault et al., 2019). 500 ng
of extracted DNA (Qiagen) from placenta (male n=2, female n=2) and embryo (male n=2,
female n=2) samples was *MspI* digested, adaptor ligated and PCR amplified (multiplex).
Multiplexed samples were pooled and 100 bp paired-end sequenced (HiSeq-2000, Illumina). The
data analyses were done according to the pipeline established at the McGill Epigenomics
Mapping and Data Coordinating Centers (Magnus et al., 2014, McGraw et al., 2015) that include
BSMAP and methylKit. Specific parameters were chosen including 100 bp step-wise tiling
windows, containing a minimum of 2 CpGs per tile and a minimum $15\times$ CpG coverage of each
tile per sample. The methylation level of a 100-bp tile was the result of all CpG C/T read counts
within the tile after coverage normalization between samples, and the methylation level reported
for a sample on autosomal chromosomes was the average methylation level across replicates.
Significant DNA methylation changes were designated as $\pm\geq20\%$ average differences between
groups of replicates and a q-value < 0.01 using the logistic regression function of methylKit
(Akalin et al., 2012). Direct comparisons between DNA methylation averages were done using
Wilcoxon-Mann-Whitney test in R. Gene ontology (GO) terms and pathway analyses for

136 RefSeqs associated to promoter-TSS were conducted using Metascape gene annotation and

137 analysis resource (Zhou et al., 2019).

139 *Sequencing Data*

140  Publicly available DNA methylation data sets (Auclair et al., 2014, Decato et al., 2017,

141 Smith et al., 2014, Smith et al., 2017, Whidden et al., 2016, Hon et al., 2013) were analyzed

142 using a custom script to intersect single CpG site methylation calls from these data sets within

143 defined 100bp tiles associated to the embryo-placenta DMRs categories

145 **RESULTS**

146 *Increased Fluctuation in Genome-Wide DNA Methylation Levels in Placental Cells*

147  To identify the overall epigenetic asymmetry that exists between placental and embryonic

148 genomes during mouse *in utero* development, we first established genome-wide DNA

149 methylation profiles using RRBS (McGraw et al., 2015, Magnus et al., 2014, Legault et al.,

150 2019) of embryos and their corresponding placentas at mid-gestation (E10.5), the developmental

151 stage at which the mouse placenta is considered mature (Cross et al., 1994). Using this approach,

152 we quantified the DNA methylation profiles of ~1.8 million CpG sites in each sample (embryo

153 n=4, placenta n=4). We found that the accumulation of CpG methylation was very distinct

154 between the placenta and the embryo. In the placenta, we detected a greater proportion of CpGs

155 in the 0-50% methylation range and a lower proportion of CpGs in the 80-100% methylation

156 range (Figure 1A). Although we observed that a large proportion of CpGs within the examined

157 regions had no methylation marks in both the embryo and the placenta, interestingly, most CpGs

158 with partial methylation (20-50%) in placenta showed high methylation levels (>80%) in

159 embryos (Figure 1B). The divergence in global DNA methylation profiles conferred a high

160 degree of clustering between both tissue types (Figure S1). These results are consistent with

161 previous studies indicating lower levels of overall DNA methylation in extraembryonic tissues

162 (Price et al., 2012, Schroeder et al., 2013, Smith et al., 2017), reviewed in (Robinson and Price,

163 2015).

164  To further investigate the dynamics of DNA methylation between placental and

165 embryonic genomes and enable direct comparison of precise regions, we segmented the genome

166 of autosomal chromosomes into 100bp non-overlapping genomic windows (*tiles*; see methods

section). After removal of sex chromosomes, we identified 245 048 unique sequenced tiles

(referred to as All-tiles) containing 896 820 common CpGs between all placenta and embryo

samples and with a minimum of 15x sequencing depth. We observed a strong reduction in the

fraction of highly methylated tiles (80-100%) in the placenta compared to the embryo (Figure

1C), which correlated with a sharp increase in the number of placenta tiles in the 0-20%, 20-40%

and 40-60% methylation range. Globally, we found that the average DNA methylation level

across all placental tiles was significantly lower compared to all embryo tiles (27% vs 45%,

p<0.0001) (Figure 1C). This overall epigenetic disparity in embryo and placenta DNA

methylation levels was especially noticeable when we mapped the methylation mean of All-tiles

with respect to regions surrounding the transcription start sites (TSS) (Figure 1D). When we

focused on a specific chromosome section (e.g., chr 7, 35Mb) (Figure 1E), we observed that

genomic segments with high DNA methylation levels in embryos were predominantly

hypomethylated in placenta. We also observed that gene or CpG island (CGI) poor regions had

consistently high methylation levels in embryos and lower methylation levels in the placenta.

Together, these results indicate that the mid-gestation placenta has very distinctive global DNA

methylation profiles compared to the embryo, and that this lower level of global DNA

methylation (i.e., DNA hypomethylation) across the placental genome is due to a significant

lower number of highly methylated (≥80%) genomic regions. We can also conclude that despite

the cellular heterogeneity in E10.5 placental and embryonic tissues, the vast majority of the

conceptus possesses specific genomic regions with either low (0-20%) or high (80-100%) levels

of methylation. However, the placenta genome presents an increased number of regions having a

broader distribution of DNA methylation (20-80%), revealing a greater diversity in methylation

levels across placental cell types compared to embryonic cell types.

Embryonic and Placental DNA Methylation Divergences Across Genomic Features

To explain the developmentally divergent methylated states between the mouse embryo

and the placenta, we next sought to precisely determine the genomic features revealing DNA

methylation differences. We defined differentially methylated regions (DMRs) as 100bp

genomic segments showing a significant difference of methylation levels between embryonic

and placental samples with an absolute mean methylation difference of 20% or higher (McGraw

et al., 2015, Shaffer et al., 2015, Piche et al., 2019). Using these conditions, we screened the 245

048 unique tiles common between all samples and identified 110 240 DMRs (~45% of All-tiles;

Supplementary Table 1) with tissue- and/or developmental-specific DNA methylation variations

between the embryo and the placenta (Figure2A; random subset of 20,000 DMRs shown).

Consistent with our findings (Figure 1), the majority (96.8%; n=106 712) of DMRs had lower

DNA methylation in the placenta (referred to as Hypo-DMRs) and only a small proportion

(3.2%; n=3 528) showed increased methylation levels (referred to as Hyper-DMRs) compared to

the embryo (Figure S2). For Hypo-DMRs, tiles mainly overlapped (96%; n=102 982) with

intergenic, intron, exon and promoter-TSS regions (Figure S2). For each of the genomic feature

categories of Hypo-DMRs, the average DNA methylation levels in the placenta were essentially

half of those present in the embryo (Figure 2B). As for DMRs with higher methylation levels in

the placenta (Hyper-DMRs), we noticed that the vast majority of these tiles (83%; n=2 929) had

very low methylation levels in the embryo (<20%) (Figure 2A-B). Most of Hyper-DMRs

overlapped intergenic, intron, exon and promoter-TSS regions (93%; n=3 272, Figure S2). We

then investigated the relationship between gene expression and DMR DNA methylation levels.

Highly expressed genes in placental tissue such as *Syna* (*Syncytin A*) (Mi et al., 2000) showed

overall lower methylation levels in the placenta when compared to the embryo (Figure 2D,

smoothed representation) (Hansen et al., 2012). Similar observations were made in the gene body

of *Atf6b* (*Activating Transcription Factor 6 Beta*), a gene implicated in the transcriptional

downregulation of *Pgf* (*Placental Growth Factor*) in response to endoplasmic reticulum stress in

pathological placentas (Mizuuchi et al., 2016). As for genes predominantly expressed in the

embryo, *Mir219a-2/Mir219b*, a brain-specific non-coding microRNA, showed higher

methylation in the placenta (Figure 2D). It was recently shown that *Mir219* was abnormally

expressed during pregnancy in amnion retrieved specifically from obese women (Nardelli et al.,

2014) and, interestingly, obesity has been suggested to cause epigenetic changes (Lillycrop and

Burdge, 2011). Another example of placenta Hyper-DMRs is *Sox6* (*SRY-Box 6*), which is

implicated in the terminal differentiation of muscle (Figure 2D) (Kamachi and Kondoh, 2013).

These observations point to an association between tissue-specific DNA methylation levels of

DMRs and biological functions. Gene ontology enrichment analyses further support that Hypo-

DMRs were strongly associated to germline functions and reproduction (*e.g.*, male and female

gamete generation, reproduction, piRNA metabolic process, meiotic cell cycle, germ cell

development) (Figure 2C). As for Hyper-DMRs, top biological processes were mostly associated

with developmental and differentiation processes (*e.g*., regionalization, embryo development,
pattern specification process, skeletal system, head development) (Figure 2C). Fittingly, the
biological functions were completely divergent between Hypo- and Hyper-DMRs. Altogether,
these results denote that Hypo- and Hyper-DMRs are present across genomic features between
mid-gestation embryo and placenta, and that these DNA methylation divergences are implicated
in promoting/repressing specific processes during embryonic and placental development.

Presence of Distinctive DMR Categories between Mid-Gestation Embryo and Placenta

Amongst DMRs, our analyses also suggest the presence of particular DMR categories
based on their level of DNA methylation in the embryo and placenta (Figure 2A). By defining
subsets of DMRs and establishing their dynamic properties between tissues, we might better
understand the genome-wide asymmetry in DNA methylation levels observed between the
embryo and the placenta. To do so, we first clustered DMR-associated tiles in 6 different
categories based on their range of low, mid and high methylation level (Low; <20%, Mid; ≥20 to
<80%, High; ≥80%) in the embryo and the placenta, and followed the DMR category transitions
between both tissues. We observed that DMRs with High-levels of methylation in the embryo
overlapped with a large proportion of DMRs that showed Mid-levels of methylation in the
placenta (Figure 3A,B; High-Mid n=72 715), whereas only a fraction corresponded to DMRs
with Low-levels of methylation in the placenta (Figure 3A,B; High-Low n=1 889). As for DMRs
with Mid-levels of methylation in embryo, the largest part remained in that same Mid-levels
category in the placenta (Figure 3A; Mid-Mid n=23 640). Nonetheless, a portion of these
embryonic Mid-levels DMRs were directed to DMRs with either Low- (Figure 3A; Mid-Low
n=9 055) or High- (Figure 3A; Mid-High n= 21) levels of methylation in the placenta. Finally,
DMRs with Low-levels of methylation in the embryo all showed Mid-levels of methylation in
the placenta (Figure 3A; Low-Mid n=2 920). Thus, when we subdivide our DMRs into
distinctive categories, we uncover that embryonic cells possess a large proportion of DMRs of
High-level (≥80%) of DNA methylation, which remain potentially static in embryonic tissues,
whereas the vast majority of these DMRs have inferior and wide-range DNA methylation levels
within placental tissue.

Specific Genomic Features Associated with DMR Categories

We next aimed to determine if the genomic distribution of embryo-placenta DMR
categories was associated with distinct genomic features. First, by classifying by genomic
annotations, we observed that DMRs with reduced methylation levels in the placenta (High-Mid,
High-Low, Mid-Low) were prevailingly found in intergenic regions (>50% of tiles) (Figure 3C),
whereas DMR categories with equivalent (Mid-Mid) or superior methylation level (Mid-High,
Low-Mid) in the placenta were more frequent in genic associated regions (>50% of tiles).
However, divergence between DMR categories was observed when we performed ontology
analyses on promoter regions (Figure S3), as each DMR subtype clearly showed distinct
biological functions. For example, High-Mid DMRs top functions were related to genes (e.g.,
Aszl, Tnp1, *Piwil2*) associated with male gamete generation, piRNA metabolic process and male
meiotic nuclear division, whereas those of the Low-Mid category were mostly associated with
developmental aspects (e.g., skeletal system development, regionalization, nervous system
development). Since we know that in placenta, activation of retrotransposon-derived genes is
interrelated with low DNA methylation levels (Macaulay et al., 2011, Reiss et al., 2007, Cohen
et al., 2011), we next assessed how DMR categories overlapped with major types of
retrotransposons (LINE; long interspersed nuclear elements, SINE; short interspersed nuclear
element, and LTR; long terminal repeats). Out of the DMR categories, those with High-levels of
methylation in the embryo and either Mid- or Low-levels in the placenta (High-Mid and High-
Low) showed the most enriched overlap with retrotransposons, with 92% and 96% respectively
(Figure 3D). DMR categories associated with gain of methylation in the placenta (Mid-High and
Low-Mid) showed the least overlap with retrotransposons, especially the Low-Mid subtype. This
highlights that during the *de novo* acquisition of DNA methylation patterns, DMRs with High-
levels of methylation in the embryo and lower levels in the placenta are almost exclusively
within retrotransposons-associated sequences, whereas DMRs with Low-levels of methylation in
the embryo and higher DNA methylation in the placenta are preferentially outside
retrotransposons-associated sequences. Finally, we investigated the proximity of the DMR
categories in regards to CpG rich (CpG islands; CGI), neighbouring (shore; < 2kb away from
CGIs, shelf; 2-4kb away from CGIs) and distant (open sea; > 4kb away from CGIs) regions
(Figure 3E). We observed that most DMR categories are depleted from CGIs and are mostly
found in open sea regions. In contrast, ~60% of tiles in Low-Mid DMRs overlapped with CGI,
shores and shelves, revealing that the acquisition of *de novo* methylation for these genomic

fragments in the extraembryonic lineage preferentially targets sequences inside or surrounding
CGIs.

Altogether, these results indicate that the asymmetry within DMR categories, based on
their methylation levels in the embryo and the placenta, can be associated to specific biological
functions and genomic-derived features (e.g., CpG, retrotransposon contents). This is particularly
apparent for DMRs with High-levels in the embryo (High-Mid, High-Low) and those with higher
methylation levels in the placenta (Low-Mid).

DMR Categories are Established During the De Novo Methylation Wave and Maintained
Throughout Development

To gain insights into the kinetics of lineage-specific DMR establishment between mid-
gestation embryo and placenta, as well as their status during development, we assessed the levels
of methylation associated with tiles for each DMR category as a function of their developmental
stage. Publicly available sequencing data (Smith et al., 2012, Whidden et al., 2016) were
analyzed using our custom script to generate 100bp tiles, and the DNA methylation levels for
each tile were calculated. In E3.5 blastocysts, when the mouse genome is mostly depleted from
DNA methylation marks, we observed low global DNA methylation levels (average <20%) for
all DMR subtypes, with similar median levels between the committed cells of the inner cell mass
(ICM) and trophectoderm lineages (Figure 4). DNA methylation levels tended to be higher in the
trophectoderm for regions falling in the Mid-High DMRs, although measurements are based on
very few DMRs for this specific category (n=21, Figure 3B, Mid-High). Since global DNA
methylation is re-acquired in the next few subsequent developmental stages, we then asked
whether the contrast in DNA methylation levels associated with the various DMR categories at
mid-gestation would already be present between E6.5 epiblast and extraembryonic ectoderm
(ExE) cell lineages, layers that are mostly composed of homogeneous and undifferentiated cell
populations. For all DMR categories, DNA methylation levels in the E6.5 epiblast and ExE
already showed similar pattern trends to those observed in the E10.5 embryo and placenta
(Figure 4, Table S2). Interestingly, for all DMR categories we observed slightly higher DNA
methylation levels in E6.5 ExE compared to E10.5 placenta (Figure 4, Table S2). However,
when measured in the subsequent developmental stages (E10.5, E11.5, E15 and E18), levels
associated with each category stabilized and remained in the same methylation range. This is true

for all categories, except for the Mid-High DMRs where methylation levels are reduced by ~20%

at E18. As for the E6.5 epiblast DNA methylation profiles, they closely matched those observed

in the E10.5 embryo, except again for the Mid-High DMRs. For each of the DMR subtypes, the

DNA methylation levels were largely stable and persisted across other time points (E6.5, E10.5,

E11.5) in embryonic cells. Although no public DNA methylation data were available for whole

embryos at later stages, when we overlapped tiles associated with E10.5 DMR categories with

data from differentiated tissues of adult mice, the global DNA methylation profiles closely

matched those for the majority of DMR categories (High-Mid, High-Low, Mid-Mid and Mid-

Low) (Figure 4, Figure S4). We conclude that the various DMR categories observed at E10.5 are

established during the embryonic and extraembryonic lineage-specification processes occurring

during the peri-implantation wave of *de novo* methylation, and that these DNA methylation

landscapes are widely retained throughout embryo and placenta development. Furthermore, for

most DMR categories, the DNA methylation levels observed in developing embryos are long-

lasting and conserved throughout somatic cell differentiation.

Dnmt3a- or Dnmt3b-Deficiency Alters Proper Establishment of DMR-Associated Patterns

Since the combined activity of DNMT3A and DNMT3B is essential for proper

establishment of DNA methylation profiles and normal development, we next sought to define

the contribution of each enzyme in the *de novo* establishment of DNA methylation in subtypes of

DMRs in embryonic and extraembryonic cell lineages. To do so, we used DNA methylation data

from publicly available datasets (whole genome bisulfite sequencing and RRBS) at E6.5

(epiblast and ExE) and E8.5 (embryo) with various inactive forms of *Dnmts*. First, when we

overlapped our All-tiles subset, we observed a substantial reduction (p□0.0001) in average DNA

methylation in absence of DNMT3A (37.81%) or DNMT3B (31.48%) in the E6.5 epiblast

compared to wild-type (44.81%), whereas in the E6.5 ExE such a comparable loss was only

associated with a *Dnmt3b*-deficiency (wt: 34.38% vs *Dnmt3b*$^{-/-}$: 19.33%, p□0.0001) (Figure S5).

Similarly, in the E8.5 embryo, a significant reduction in average DNA methylation level was

measured with lack of *Dnmt3a* or *Dnmt3b* expression compared to wild-type (wt: 47.5%;

Dnmt3a$^{-/-}$: 43.6%; *Dnmt3b*$^{-/-}$: 34%, p□0.0001) (Figure S5). For most DMR subtypes, absence of

DNMT3A caused modest or no reduction on overall DNA methylation levels in E6.5 epiblast

and ExE (Figure 5A and S6, Table S3). Intriguingly, *Dnmt3a*-deficiency caused a slight gain of

methylation (wt: 76.9% vs *Dnmt3a*$^{-/-}$: 81.5%, p□0.0001) level in the E6.5 epiblast for High-Mid

DMRs, suggesting a potential overcompensation effect by DNMT3B in the epiblast on this

specific DMR subtype. In comparison, *Dnmt3b*-depletion led to a substantial loss of average

methylation levels in all DMR categories in the E6.5 epiblast and ExE, except for epiblast High-

Mid DMRs (Figure 5A, Table S3) for which a compensatory mechanism provided high levels of

methylation (wt: 76.9% vs *Dnmt3b*$^{-/-}$: 71.6%). The compensatory mechanism on High-Mid

DMRs in response to *Dnmt3b*-deficiency was ineffective in the E6.5 ExE. When we focused on

promoter-TSS for each DMR categories, we observed again that loss of methylation for these

regulatory regions was principally associated with *Dnmt3b*-deficiency (Figure 5B). The

promoter-TSS regions associated with High-Mid DMRs retained relatively high methylation

levels for either *Dnmt3a*$^{-/-}$ or *Dnmt3b*$^{-/-}$ epiblast samples, demonstrating a robust and

compensatory *de novo* methylation mechanism. To further underline the impact of DNMT3A or

DNMT3B on the *de novo* methylation of DMRs, we measured methylation levels for DMRs

selected from our gene enrichment analyses (*Fgb*, *P2rx7*, *Pcyt2*, *Etnppl*, *Ralgds*, *Lrp5*) and other

genomic segments covered by multiple tiles (*Cxxc1*, *Fcgrt*, *Lamp5*, *Mbd1*, *Lphn1*, *Pick1*, *Irf1*)

(Figure S7 & S8) in *Dnmt3a*- or *Dnmt3b*-deficient epiblast and ExE. In line with our global

observations, for most DMR-associated tiles, a *Dnmt3b*-deficiency in E6.5 epiblast or ExE

caused a more severe loss of methylation compared to lack of *Dnmt3a*. However, for some

regions in the E6.5 epiblast (e.g., *Pick1*, *Irf1*), *Dnmt3a*$^{-/-}$ methylation levels were inferior to those

of *Dnmt3b*$^{-/-}$. Noteworthy, for most DMRs, lack of co-factor *Dnmt3l* affected DNA methylation

levels to greater extent than *Dnmt3a*-deficiency (Figure S6). Overall, we show that of DNMT3A

and DNMT3B participate in the establishment of the asymmetric methylation patterns associated

with the various DMR categories in both embryonic and extraembryonic cells, with DNMT3B

being the principal contributor in both cell lineages during the *de novo* reprogramming wave as it

can compensate almost entirely for the absence of DNMT3A.

Decline in Compensatory DNA Methylation Mechanisms in Response to Dnmt3a- or Dnmt3b-

Deficiency.

381   As embryonic development progresses from E6.5 to E8.5, our data suggest that lack of

DNMT3A or DNMT3B further reduces global methylation levels of DMRs (High-Mid, High-

Low, Mid-Low and Mid-Mid), evoking that absence of either enzymatic activity causes an

additive and extended effect (Figure 5A-B). To further define the robustness in compensatory

mechanism between DNMT3A or DNMT3B in the establishment and maintenance of DNA

methylation on DMRs categories, we focused on High-Mid promoter-TSS- associated DMRs as

they have the highest DNA methylation levels and require the most *de novo* methyltransferase

activity. Methylation differences of ≥20% between wild-type and *Dnmt3a-* or *Dnmt3b*-deficient

samples were considered as regions showing substantial lack of compensation. In agreement

with our results (Figure 5A-B), we observed that at E6.5, DNMT3B can compensate almost

entirely for DNMT3 Aloss by maintaining methylation levels on most promoters-TSS associated

tiles in the epiblast (410/437 = 93.8%) and ExE (558/602 = 92.7%), whereas DNMT3A can only

partially alleviate the lack of DNMT3B in the epiblast (315/410 = 78.6%) and ExE (244/524 =

46.6%) (Figure 6A-B). As embryonic cell lineages development evolves between E6.5 and E8.5,

global methylation levels increase on promoter-TSS of High-Mid DMRs. However, we detected

a sharp decline in the compensatory mechanism to rescue the *de novo* and/or maintenance of

methylation levels of promoter-TSS DMRs in the *Dnmt3a-* (524/627 = 80.9%) or *Dnmt3b-*

(256/622 = 41.2%) deficient E8.5 embryos. This is further highlighted when we focus on a

subgroup of 314 promoter-TSS associated tiles overlapping all of 9 data sets (Figure 6C),

including gene promoters (e.g., *Asz1, Catsper1, Ccdc42, Dmrtb1, Piwil2, Rpl10l, Sox30, Sycp1,*

Tnp1, Ttll1, Zfn42) related to our top enriched biological functions (i.e., piRNA, gamete

generation and germ cells, meiotic nuclear division). Although E8.5 *Dnmt3a-* or *Dnmt3b-*

deficient ExE data was not available, our data suggest that greater compensation failure would

also be observed in this tissue. Thus, embryonic-placental DMRs need the combined action of

DNMT3Aand DNMT3Bto both establish and maintain proper asymmetric levels during early

development as compensatory DNA methylation mechanisms during the *de novo* wave of

methylation decline overtime and fail to overcome methyltransferase shortage in later

development stages.

DISCUSSION

With recent breakthroughs in high-throughput sequencing, we now have a better

understanding of the dynamic of DNA methylation erasure occurring in early cleavage stage

embryos following fertilization. However, the discrepancies in acquisition of genome-wide DNA

methylation patterns in early post-implantation embryonic and extraembryonic cell lineages

remain to be methodically delineated. To address this issue and further our understanding of the DNA methylation asymmetry that guides the developmental trajectory of embryonic and extraembryonic cell lineages, we established genome-wide DNA methylation profiles of mouse embryo and placenta at mid-gestation, and analyzed various publicly available developmental stage specific embryo and placenta DNA methylation datasets. Using this strategy, we uncovered that 45% of the genomic regions analyzed differ in DNA methylation status (≥20%) between mid-gestation embryo and placenta, and that these DMRs can be further divided into categories based on their levels of DNA methylation (Low; <20%, Mid; ≥20 to <80%, High; ≥80%) in the embryo and placenta. We show that the embryo and placenta acquire specific DMR categories during the early stage of the *de novo* DNA methylation wave, and that these DMRs persist throughout prenatal development, as well as into somatic adult tissues. Furthermore, we show that *Dnmt3b* primarily drives the divergence in DNA methylation levels associated with these specific DMRs and that *de novo* methyltransferase activity of *Dnmt3b* can almost entirely compensate for lack of *Dnmt3a* in the asymmetric establishment of embryonic and placental DMRs, but that *Dnmt3a* can only partially palliate the absence of *Dnmt3b*. However, with developmental progression, this compensatory DNA methylation mechanism becomes less effective.

Our results indicate that the kinetics of DNA methylation acquisition leading to specific embryo-placenta DMR categories is not a stepwise process occurring throughout cell fate decisions and patterning of embryonic and extraembryonic lineages, but a prompt progression in the early post-implanted conceptus. These results are in line with studies reflecting that the initiation of asymmetric DNA methylation levels begins within the trophoblast and the inner cell mass of the blastocyst (Santos et al., 2002), and becomes highly evident by the time the epiblast acquires its initial global DNA methylation patterns at E6.5 during the *de novo* reprogramming wave (Auclair et al. 2014, Smith et al. 2017). Our results show that the acquisition period between E4.5 and E6.5 is particularly key to establishing asymmetric DNA methylation patterns associated with mid-gestation embryo-placenta DMR classes, and that these DMR associated-patterns are long-lasting across stages of prenatal development. Since we studied cell populations derived from embryonic and placental tissues, we cannot dismiss the prevalence of cell-to-cell heterogeneity in the acquisition kinetics of specific DMR patterns. Despite this concern, our results indicate that embryonic cells have a large body of DMRs with High-levels (≥80%) and

Low-levels (<20%) of methylation that remain static across development, as well as in somatic
cell types (High-Mid & High-Low DMRs), whereas compared to the embryo, the placental
DMRs have overall lower methylation levels and are more broadly distributed amongst
methylation levels, which has also been shown in other studies where genome-wide DNA
methylation profiles of placental cells were compared to other tissues and specific cell types
(Schroeder et al., 2013; Chatterjee et al., 2016; Smith et al., 2017). Lower methylation levels in
the placenta have been associated with reduced *de novo* methyltransferase activity in the ExE
during the *de novo* reprogramming wave (Fulka et al., 2004, Santos et al., 2002). Nevertheless,
we observe methylation level peaks for all DMR categories within the ExE at E6.5 before levels
stabilize at E10.5. As of now, the implication of these methylation level peaks on future
regulation mechanisms and methylation profiles is unknown. Moreover, it remains to be
determined whether global reduction in DNA methylation marks on DMRs between the E6.5 and
E10.5 extraembryonic cell lineages is stochastic or targeted, and whether it occurs through
passive or active mechanisms.

We also observed an enrichment of retrotransposons (i.e., LINE, SINE, LTR) in DMRs
especially in those with High-levels of methylation in the embryo and lower-levels in the
placenta, whereas DMRs with Low-levels of methylation in the embryo and higher levels in the
placenta are preferentially outside retrotransposons-associated sequences. Earlier findings of
Chapman et *al*. (Chapman et al., 1984) revealed that repeat regions in the placenta appear to lack
tight control of their methylation patterns, perhaps indicating that maintaining methylation, and
therefore repression of these elements for genome stability and integrity, is not critical given the
relatively short lifespan of this organ. However, we do observe Mid-range methylation levels
(20-80%) in the placenta for a substantial portion of DMRs that are associated with
retrotransposons, revealing that specific genomic regions associated with repetitive elements do
need tight regulation in extraembryonic cell lineages for proper development. These results are in
line with findings exposing that the deletion of genome-defense gene *Tex19.1* leads to the de-
repression of LINE1 and compromises placental development, suggesting that disparities
between retrotransposon suppression and genome-defense mechanisms might contribute to
placenta dysfunction and disease (Reichmann et al., 2013).

DNMT3A and DNMT3B are required to establish proper methylation profiles on the
embryonic genome during the *de novo* reprogramming wave, as both methyltransferase enzymes

have redundant, but also specific functions. However, the activity of DNMT3B is the main

contributor in the acquisition of profiles in epiblast cells, and especially commands methylation

on CGIs associated with developmental genes (Auclair et al., 2014). Auclair et *al*. highlighted

that in the absence of DNMT3B, DNMT3A was not able to counterbalance, leading to the loss of

promoter-CGI methylation and gain of expression of germline genes (e.g., *Sycp1*, *Sycp2*, *Mael*,

Rpl10l, *Dmrtb1*) in somatic cells of the embryo. Here, we show that the promoter of these

germline genes, associated with meiotic and piRNA processes as well as other genes with similar

biological functions, are highly enriched in High-Mid DMRs. We also reveal that for this

particular set of High-Mid DMRs, mainly deprived of CGIs, DNMT3A is able to substantially

counterbalance the lack of DNMT3B in the E6.5 epiblast and establish global DNA methylation

levels, whereas in the ExE, this compensatory mechanism is rather inefficient. Globally, we

show that DNMT3B is much more potent at compensating than DNMT3A in both the epiblast

and ExE for all DMR categories, indicating that DNMT3B is the main *de novo* enzyme driving

asymmetric DNA methylation patterns between the embryo and placenta. Although

compensatory mechanisms have been observed in *Dnmt3a* or *Dnmt3b*-depleted conceptuses, we

still do not fully understand the process as *Dnmt3a* and *Dnmt3b* have cell lineage specific

expression during the peri-implantation developmental period. As development progresses, we

observed that the compensation mechanism in *Dnmt3a* or *Dnmt3b*-deficient embryos remains

apparent at E8.5 on most DMR categories, but is less effective as the methylation gaps increase

compared to wild-type. This correlates with a developmental period where DNMT3A is now the

main *de novo* methyltransferase enzyme in both the embryo and placenta (Okano et al., 1999,

Watanabe et al., 2002), and with a compensatory activity of DNMT3A being less effective. This

suggests that DNMT3B activity is critical to ensure proper establishment of DNA methylation

asymmetry between the embryonic and extraembryonic cell lineages during the *de novo*

reprogramming wave, but that DNMT3A is required during the developmental progression to

safeguard methylation levels on DMR categories.

CONCLUSION

505   We demonstrate that asymmetry between embryo and placenta DNA methylation patterns

occurs rapidly during *de novo* acquisition of methylation marks in the early post-implanted

conceptus, and that these patterns are long-lasting across subtypes of DMRs. We also reveal that

at the peri-implantation stages, *de novo* methyltransferase activity of DNMT3B is the main
provider of asymmetric methylation marks on DMRs, and that it largely palliates the lack of
DNMT3A in the epiblast and ExE. However, as development progresses, DNMT3A becomes the
principal *de novo* methyltransferase by mid-gestation, and DNMT3B methyltransferase activity
is less effective at promoting compensation. These results further underline why embryos
developing without DNMT3B have severe DNA methylation defects and die at mid-gestation,
whereas those without DNMT3A only die postnatally. Further investigation is required to
determine the molecular mechanisms controlling the precise *de novo* acquisition of long-lasting
methylation marks on specific DMR subtypes in the embryonic and extraembryonic cell
lineages, and how errors in this process could lead to abnormal development and diseases.

ACKNOWLEDGMENTS

We thank the McGraw lab for critical comments and suggestions, as well as Elizabeth Maurice-
Elder for editing.

COMPETING INTERESTS
No competing interests declared

FUNDING

This work was supported by a research grant from the Natural Sciences and Engineering
Research Council of Canada. We acknowledge the Fonds de Recherche du Québec en Santé
(S.M, L.M.L) and the Réseau Québécois en Reproduction and Fonds de recherche du Québec –
Nature et technologies (K.D) for salary awards.

DATA AVAILABILITY
The data from this study have been submitted to the Gene Expression Omnibus (#GSE95610).
Reviewer link:

Table S1. Annotation and DNA methylation values associated with DMRs.

Table S2. DNA methylation levels for DMR categorises across development.

Table S3. DNA methylation levels for DMR categories in various *Dnmt* knockout mouse
models.

Table S4. Number of 100bp tiles in each category of public data that overlap with Embryo-
Placenta DMRs (related to Figure 4; 5A; S4; S5; S6)

REFERENCES

AKALIN, A., KORMAKSSON, M., LI, S., GARRETT-BAKELMAN, F. E., FIGUEROA, M.
E., MELNICK, A. & MASON, C. E. 2012. methylKit: a comprehensive R package for
the analysis of genome-wide DNA methylation profiles. *Genome Biol,* 13**,** R87.
AUCLAIR, G., GUIBERT, S., BENDER, A. & WEBER, M. 2014. Ontogeny of CpG island
methylation and specificity of DNMT3 methyltransferases during embryonic
development in the mouse. *Genome Biol,* 15**,** 545.
BESTOR, T. H. 2000. The DNA methyltransferases of mammals. *Hum Mol Genet,* 9**,** 2395-402.
BOYLE, P., CLEMENT, K., GU, H., SMITH, Z. D., ZILLER, M., FOSTEL, J. L., HOLMES,
L., MELDRIM, J., KELLEY, F., GNIRKE, A. & MEISSNER, A. 2012. Gel-free
multiplexed reduced representation bisulfite sequencing for large-scale DNA methylation
profiling. *Genome Biol,* 13**,** R92.
CHAPMAN, V., FORRESTER, L., SANFORD, J., HASTIE, N. & ROSSANT, J. 1984. Cell
lineage-specific undermethylation of mouse repetitive DNA. *Nature,* 307**,** 284-6.
CHATTERJEE, A., MACAULAY, E. C., RODGER, E. J., STOCKWELL, P. A., PARRY, M.
F., ROBERTS, H. E., SLATTER, T. L., HUNG, N. A., DEVENISH, C. J. & MORISON,
I. M. 2016. Placental Hypomethylation Is More Pronounced in Genomic Loci Devoid of
Retroelements. *G3 (Bethesda),* 6**,** 1911-21.
CHUONG, E. B. 2013. Retroviruses facilitate the rapid evolution of the mammalian placenta.
Bioessays, 35**,** 853-61.
CHURCH, D. M., GOODSTADT, L., HILLIER, L. W., ZODY, M. C., GOLDSTEIN, S., SHE,
X., BULT, C. J., AGARWALA, R., CHERRY, J. L., DICUCCIO, M., HLAVINA, W.,
KAPUSTIN, Y., MERIC, P., MAGLOTT, D., BIRTLE, Z., MARQUES, A. C.,
GRAVES, T., ZHOU, S., TEAGUE, B., POTAMOUSIS, K., CHURAS, C., PLACE, M.,
HERSCHLEB, J., RUNNHEIM, R., FORREST, D., AMOS-LANDGRAF, J.,
SCHWARTZ, D. C., CHENG, Z., LINDBLAD-TOH, K., EICHLER, E. E., PONTING,
C. P. & MOUSE GENOME SEQUENCING, C. 2009. Lineage-specific biology revealed
by a finished genome assembly of the mouse. *PLoS Biol,* 7**,** e1000112.
COHEN, C. J., REBOLLO, R., BABOVIC, S., DAI, E. L., ROBINSON, W. P. & MAGER, D.
L. 2011. Placenta-specific expression of the interleukin-2 (IL-2) receptor beta subunit
from an endogenous retroviral promoter. *J Biol Chem,* 286**,** 35543-52.

CROSS, J. C. 2000. Genetic insights into trophoblast differentiation and placental
morphogenesis. *Semin Cell Dev Biol,* 11**,** 105-13.
CROSS, J. C., WERB, Z. & FISHER, S. J. 1994. Implantation and the placenta: key pieces of
the development puzzle. *Science,* 266**,** 1508-18.
DECATO, B. E., LOPEZ-TELLO, J., SFERRUZZI-PERRI, A. N., SMITH, A. D. & DEAN, M.
D. 2017. DNA Methylation Divergence and Tissue Specialization in the Developing
Mouse Placenta. *Mol Biol Evol,* 34**,** 1702-1712.
EMERA, D. & WAGNER, G. P. 2012. Transposable element recruitments in the mammalian
placenta: impacts and mechanisms. *Brief Funct Genomics,* 11**,** 267-76.
FULKA, H., MRAZEK, M., TEPLA, O. & FULKA, J., JR. 2004. DNA methylation pattern in
human zygotes and developing embryos. *Reproduction,* 128**,** 703-8.
GU, H., SMITH, Z. D., BOCK, C., BOYLE, P., GNIRKE, A. & MEISSNER, A. 2011.
Preparation of reduced representation bisulfite sequencing libraries for genome-scale
DNA methylation profiling. *Nat Protoc,* 6**,** 468-81.
GUO, H., ZHU, P., YAN, L., LI, R., HU, B., LIAN, Y., YAN, J., REN, X., LIN, S., LI, J., JIN,
X., SHI, X., LIU, P., WANG, X., WANG, W., WEI, Y., LI, X., GUO, F., WU, X., FAN,
X., YONG, J., WEN, L., XIE, S. X., TANG, F. & QIAO, J. 2014. The DNA methylation
landscape of human early embryos. *Nature,* 511**,** 606-10.
HAIG, D. 2012. Retroviruses and the placenta. *Curr Biol,* 22**,** R609-13.
HANSEN, K. D., LANGMEAD, B. & IRIZARRY, R. A. 2012. BSmooth: from whole genome
bisulfite sequencing reads to differentially methylated regions. *Genome Biol,* 13**,** R83.
HIRASAWA, R., CHIBA, H., KANEDA, M., TAJIMA, S., LI, E., JAENISCH, R. & SASAKI,
H. 2008. Maternal and zygotic Dnmt1 are necessary and sufficient for the maintenance of
DNA methylation imprints during preimplantation development. *Genes Dev,* 22**,** 1607-
16.
HON, G. C., RAJAGOPAL, N., SHEN, Y., MCCLEARY, D. F., YUE, F., DANG, M. D. &
REN, B. 2013. Epigenetic memory at embryonic enhancers identified in DNA
methylation maps from adult mouse tissues. *Nat Genet,* 45**,** 1198-206.
HOWELL, C. Y., BESTOR, T. H., DING, F., LATHAM, K. E., MERTINEIT, C., TRASLER, J.
M. & CHAILLET, J. R. 2001. Genomic imprinting disrupted by a maternal effect
mutation in the Dnmt1 gene. *Cell,* 104**,** 829-38.
KAMACHI, Y. & KONDOH, H. 2013. Sox proteins: regulators of cell fate specification and
differentiation. *Development,* 140**,** 4129-44.
KANEDA, M., OKANO, M., HATA, K., SADO, T., TSUJIMOTO, N., LI, E. & SASAKI, H.
2004. Essential role for de novo DNA methyltransferase Dnmt3a in paternal and maternal
imprinting. *Nature,* 429**,** 900-3.
KATO, Y., KANEDA, M., HATA, K., KUMAKI, K., HISANO, M., KOHARA, Y., OKANO,
M., LI, E., NOZAKI, M. & SASAKI, H. 2007. Role of the Dnmt3 family in de novo
methylation of imprinted and repetitive sequences during male germ cell development in
the mouse. *Hum Mol Genet,* 16**,** 2272-80.
LEGAULT, L.-M., CHAN, D. & MCGRAW, S. 2019. Rapid Multiplexed Reduced
Representation Bisulfite Sequencing Library Prep (rRRBS). *BIO-PROTOCOL,* 9.
LEI, H., OH, S. P., OKANO, M., JUTTERMANN, R., GOSS, K. A., JAENISCH, R. & LI, E.
1996. De novo DNA cytosine methyltransferase activities in mouse embryonic stem
cells. *Development,* 122**,** 3195-205.

LEONHARDT, H., PAGE, A. W., WEIER, H. U. & BESTOR, T. H. 1992. A targeting sequence
622   directs DNA methyltransferase to sites of DNA replication in mammalian nuclei. *Cell,*
623   71**,** 865-73.
624   LI, E. 2002. Chromatin modification and epigenetic reprogramming in mammalian development.
625   *Nat Rev Genet,* 3**,** 662-73.
626   LI, E., BESTOR, T. H. & JAENISCH, R. 1992. Targeted mutation of the DNA
627   methyltransferase gene results in embryonic lethality. *Cell,* 69**,** 915-26.
628   LILLYCROP, K. A. & BURDGE, G. C. 2011. The effect of nutrition during early life on the
629   epigenetic regulation of transcription and implications for human diseases. *J Nutrigenet*
630   *Nutrigenomics,* 4**,** 248-60.
631   MACAULAY, E. C., WEEKS, R. J., ANDREWS, S. & MORISON, I. M. 2011.
632   Hypomethylation of functional retrotransposon-derived genes in the human placenta.
633   *Mamm Genome,* 22**,** 722-35.
634   MAGNUS, N., GARNIER, D., MEEHAN, B., MCGRAW, S., LEE, T. H., CARON, M.,
635   BOURQUE, G., MILSOM, C., JABADO, N., TRASLER, J., PAWLINSKI, R.,
636   MACKMAN, N. & RAK, J. 2014. Tissue factor expression provokes escape from tumor
637   dormancy and leads to genomic alterations. *Proc Natl Acad Sci U S A,* 111**,** 3544-9.
638   MCGRAW, S., ZHANG, J. X., FARAG, M., CHAN, D., CARON, M., KONERMANN, C.,
639   OAKES, C. C., MOHAN, K. N., PLASS, C., PASTINEN, T., BOURQUE, G.,
640   CHAILLET, J. R. & TRASLER, J. M. 2015. Transient DNMT1 suppression reveals
641   hidden heritable marks in the genome. *Nucleic Acids Res*.
642   MI, S., LEE, X., LI, X., VELDMAN, G. M., FINNERTY, H., RACIE, L., LAVALLIE, E.,
643   TANG, X. Y., EDOUARD, P., HOWES, S., KEITH, J. C., JR. & MCCOY, J. M. 2000.
644   Syncytin is a captive retroviral envelope protein involved in human placental
645   morphogenesis. *Nature,* 403**,** 785-9.
646   MIZUUCHI, M., CINDROVA-DAVIES, T., OLOVSSON, M., CHARNOCK-JONES, D. S.,
647   BURTON, G. J. & YUNG, H. W. 2016. Placental endoplasmic reticulum stress
648   negatively regulates transcription of placental growth factor via ATF4 and ATF6beta:
649   implications for the pathophysiology of human pregnancy complications. *J Pathol,* 238**,**
650   550-61.
651   MONK, M. 1987. Genomic imprinting. Memories of mother and father. *Nature,* 328**,** 203-4.
652   MORGAN, H. D., SANTOS, F., GREEN, K., DEAN, W. & REIK, W. 2005. Epigenetic
653   reprogramming in mammals. *Hum Mol Genet,* 14 Spec No 1**,** R47-58.
654   NAKANISHI, M. O., HAYAKAWA, K., NAKABAYASHI, K., HATA, K., SHIOTA, K. &
655   TANAKA, S. 2012. Trophoblast-specific DNA methylation occurs after the segregation
656   of the trophectoderm and inner cell mass in the mouse periimplantation embryo.
657   *Epigenetics,* 7**,** 173-82.
658   NARDELLI, C., IAFFALDANO, L., FERRIGNO, M., LABRUNA, G., MARUOTTI, G. M.,
659   QUAGLIA, F., CAPOBIANCO, V., DI NOTO, R., DEL VECCHIO, L., MARTINELLI,
660   P., PASTORE, L. & SACCHETTI, L. 2014. Characterization and predicted role of the
661   microRNA expression profile in amnion from obese pregnant women. *Int J Obes (Lond),*
662   38**,** 466-9.
663   ODA, M., OXLEY, D., DEAN, W. & REIK, W. 2013. Regulation of lineage specific DNA
664   hypomethylation in mouse trophectoderm. *PLoS One,* 8**,** e68846.

OKAHARA, G., MATSUBARA, S., ODA, T., SUGIMOTO, J., JINNO, Y. & KANAYA, F.
2004. Expression analyses of human endogenous retroviruses (HERVs): tissue-specific
and developmental stage-dependent expression of HERVs. *Genomics,* 84**,** 982-90.
OKANO, M., BELL, D. W., HABER, D. A. & LI, E. 1999. DNA methyltransferases Dnmt3a
and Dnmt3b are essential for de novo methylation and mammalian development. *Cell,*
99**,** 247-57.
PICHE, J., GOSSET, N., LEGAULT, L. M., PACIS, A., ONEGLIA, A., CARON, M.,
CHETAILLE, P., BARREIRO, L., LIU, D., QI, X., NATTEL, S., LECLERC, S.,
BRETON-LARRIVEE, M., MCGRAW, S. & ANDELFINGER, G. 2019. Molecular
Signature of CAID Syndrome: Noncanonical Roles of SGO1 in Regulation of TGF-beta
Signaling and Epigenomics. *Cell Mol Gastroenterol Hepatol,* 7**,** 411-431.
PRICE, E. M., COTTON, A. M., PENAHERRERA, M. S., MCFADDEN, D. E., KOBOR, M. S.
& ROBINSON, W. 2012. Different measures of "genome-wide" DNA methylation
exhibit unique properties in placental and somatic tissues. *Epigenetics,* 7**,** 652-63.
REICHMANN, J., REDDINGTON, J. P., BEST, D., READ, D., OLLINGER, R., MEEHAN, R.
R. & ADAMS, I. R. 2013. The genome-defence gene Tex19.1 suppresses LINE-1
retrotransposons in the placenta and prevents intra-uterine growth retardation in mice.
Hum Mol Genet, 22**,** 1791-806.
REISS, D., ZHANG, Y. & MAGER, D. L. 2007. Widely variable endogenous retroviral
methylation levels in human placenta. *Nucleic Acids Res,* 35**,** 4743-54.
ROBINSON, W. P. & PRICE, E. M. 2015. The human placental methylome. *Cold Spring Harb*
Perspect Med, 5**,** a023044.
ROSSANT, J. & CROSS, J. C. 2001. Placental development: lessons from mouse mutants. *Nat*
Rev Genet, 2**,** 538-48.
SANTOS, F., HENDRICH, B., REIK, W. & DEAN, W. 2002. Dynamic reprogramming of DNA
methylation in the early mouse embryo. *Dev Biol,* 241**,** 172-82.
SCHROEDER, D. I., BLAIR, J. D., LOTT, P., YU, H. O., HONG, D., CRARY, F.,
ASHWOOD, P., WALKER, C., KORF, I., ROBINSON, W. P. & LASALLE, J. M. 2013.
The human placenta methylome. *Proc Natl Acad Sci U S A,* 110**,** 6037-42.
SENNER, C. E., KRUEGER, F., OXLEY, D., ANDREWS, S. & HEMBERGER, M. 2012.
DNA methylation profiles define stem cell identity and reveal a tight embryonic-
extraembryonic lineage boundary. *Stem Cells,* 30**,** 2732-45.
SHAFFER, B., MCGRAW, S., XIAO, S. C., CHAN, D., TRASLER, J. & CHAILLET, J. R.
2015. The dnmt1 intrinsically disordered domain regulates genomic methylation during
development. *Genetics,* 199**,** 533-41.
SLOTKIN, R. K. & MARTIENSSEN, R. 2007. Transposable elements and the epigenetic
regulation of the genome. *Nat Rev Genet,* 8**,** 272-85.
SMITH, Z. D., CHAN, M. M., HUMM, K. C., KARNIK, R., MEKHOUBAD, S., REGEV, A.,
EGGAN, K. & MEISSNER, A. 2014. DNA methylation dynamics of the human
preimplantation embryo. *Nature,* 511**,** 611-5.
SMITH, Z. D., CHAN, M. M., MIKKELSEN, T. S., GU, H., GNIRKE, A., REGEV, A. &
MEISSNER, A. 2012. A unique regulatory phase of DNA methylation in the early
mammalian embryo. *Nature,* 484**,** 339-44.
SMITH, Z. D., SHI, J., GU, H., DONAGHEY, J., CLEMENT, K., CACCHIARELLI, D.,
GNIRKE, A., MICHOR, F. & MEISSNER, A. 2017. Epigenetic restriction of
extraembryonic lineages mirrors the somatic transition to cancer. *Nature,* 549**,** 543-547.

WARREN, I. A., NAVILLE, M., CHALOPIN, D., LEVIN, P., BERGER, C. S., GALIANA, D.
& VOLFF, J. N. 2015. Evolutionary impact of transposable elements on genomic
diversity and lineage-specific innovation in vertebrates. *Chromosome Res,* 23**,** 505-31.
WATANABE, D., SUETAKE, I., TADA, T. & TAJIMA, S. 2002. Stage- and cell-specific
expression of Dnmt3a and Dnmt3b during embryogenesis. *Mech Dev,* 118**,** 187-90.
WHIDDEN, L., MARTEL, J., RAHIMI, S., RICHARD CHAILLET, J., CHAN, D. &
TRASLER, J. M. 2016. Compromised oocyte quality and assisted reproduction
contribute to sex-specific effects on offspring outcomes and epigenetic patterning. *Hum*
Mol Genet.
XIE, M., HONG, C., ZHANG, B., LOWDON, R. F., XING, X., LI, D., ZHOU, X., LEE, H. J.,
MAIRE, C. L., LIGON, K. L., GASCARD, P., SIGAROUDINIA, M., TLSTY, T. D.,
KADLECEK, T., WEISS, A., O'GEEN, H., FARNHAM, P. J., MADDEN, P. A.,
MUNGALL, A. J., TAM, A., KAMOH, B., CHO, S., MOORE, R., HIRST, M.,
MARRA, M. A., COSTELLO, J. F. & WANG, T. 2013. DNA hypomethylation within
specific transposable element families associates with tissue-specific enhancer landscape.
Nat Genet, 45**,** 836-41.
ZHOU, Y., ZHOU, B., PACHE, L., CHANG, M., KHODABAKHSHI, A. H., TANASEICHUK,
O., BENNER, C. & CHANDA, S. K. 2019. Metascape provides a biologist-oriented
resource for the analysis of systems-level datasets. *Nat Commun,* 10**,** 1523.

730
731
Figure 1. Distinctive patterns of genome-wide DNA methylation accumulation between
mid-gestation embryo and placenta. Analyses of genome-wide DNA methylation sequencing
results for embryo (n=4) and placenta (n=4). **A)** Density histograms showing the distribution of
CpG methylation levels for embryo (*purple*) and placenta (*green*). **B)** Pairwise comparison of
CpG methylation between embryo and placenta. Density increases from blue to red. **C)** CpG
fraction of 100 bp tiles within 0-20%, 20-40%, 40-60%, 60-80% and 80-100% ranges in embryo

and placenta. Shown above bars, is the average CpG methylation for each unique tile represented
in graph. **D)** DNA methylation means surrounding the transcription start site (TSS) for All-tiles
in each experimental group. **E)** Circle plot showing methylation average of embryo (*purple*) and
placenta (*green*) across a portion of chromosome 7.

750
751
752 **Figure 2. Genome-wide methylation asymmetry across genomic features between embryo**
753 **and placenta**. Analyses of genome-wide DNA methylation sequencing results for embryo (n=4)

and placenta (n=4) samples. **A)** Heatmap representation of DNA methylation levels for the 20 000 tiles with the most variable levels between embryo and placenta (DNA methylation variance >20% and *P*<0.05). **B)** Box-plots representing DNA methylation distribution in embryo and placenta for the different genomic annotation regions (intergenic, intron, exon, promoter-TSS, 3′UTR, TTS, 5'UTR and non-coding). **C)** Summary of biological functions associated with promoter regions in Hypo- and Hyper-DMRs. **D)** Examples of smoothed methylation profiles (*BSmooth* tool) in regions with lower methylation profiles (*Atf6b,* and *Syna*) or higher methylation profiles (*Mir219a-2/Mir219b,* and *Sox6*) in placenta.

Figure 3. Distinct genomic features in DMRs based on their methylation status in embryo and placenta. DMR analysis by methylation levels in embryo and placenta. **A)** Sankey diagram dividing DMRs by methylation levels in embryo (*purple*) and placenta (*green*) for the associated tiles. High : ≥80% methylation, Mid (intermediate) : ≥20% - <80% methylation , Low : <20% methylation. **B)** Venn diagram showing the proportion of tiles in the different DMR categories based on DNA methylation levels between embryo and placenta. **C), D)** and **E)** Analysis of all tiles, all the DMRs, as well as the 6 different DMR categories based on levels of DNA methylation in embryo and placenta for : **C)** Genomic annotations, **D)** Main retrotransposons and **E)** Proximity of CpG rich regions. Neighboring CpG dense regions were defined as shore; up to 2kb away from CGIs, shelf; 2-4kb away from CGIs, and open sea; >4kb away from CGIs.

Figure 4. Dynamics of DNA methylation profiles associated with DMR categories and their evolution across embryo and placenta development. Box plots representing the DNA methylation distribution and median values for each DMR category in various developmental stages. Tiles associated with DMR categories at E10.5 were overlapped with previously published and publicly available data, and methylation levels were determined at each developmental stage (Smith et *al* 2012, Smith et *al*. 2014, Whidden et *al*. 2016) or in adult somatic tissues (Hon et *al*. 2013). ICM: inner cell mass, TE: trophectoderm, Epi: epiblast, ExE: extraembryonic ectoderm, E: embryo, Pla: placenta. See Table S2 for median and mean methylation values, and Table S4 for number of overlapping tiles analysed.

804
805
Figure 5. *Dnmt3a*- or *Dnmt3b*-deficiency alter proper establishment of DMR-associated
patterns. A) Violin plot representing DNA methylation distribution and median values of tiles
associated with the different DMR categories at E10.5, and their methylation levels in the
different tissues and genotypes at E6.5 and E8.5. **B)** Promoter-TSS density graphs for E10.5
embryo and E6.5 epiblast (wt, *Dnmt3a*[-/-] and *Dnmt3b*[-/-]) (left panel) and for E10.5 placenta and
E6.5 extraembryonic ectoderm (wt, *Dnmt3a*[-/-] and *Dnmt3b*[-/-]) (right panel). E: embryo, Epi:
epiblast, ExE: extraembryonic ectoderm. See Table S3 for median and mean methylation values,
and Table S4 for number of overlapping tiles analysed.
814
815

816
Figure 6. Developmental decline in compensatory DNA methylation mechanism in response
to *Dnmt3a*- or *Dnmt3b*-deficiency at promoter-TSS associated with High-Mid DMRs. A)
Heatmaps comparing the DNA methylation profiles of wt vs *Dnmt3a*$^{-/-}$ or wt vs *Dnmt3b*$^{-/-}$ for
E6.5 epiblast, E6.5 extraembryonic ectoderm or E8.5 embryo in 100bp tiles overlapping
promoter-associated DMRs that are highly methylated in the embryo (≥80%) and mildly
methylated in the placenta (≥20 and <80%). Upper panel represents regions with a difference of
methylation of at least 20% between wt and *Dnmt3a*$^{-/-}$ or wt and *Dnmt3b*$^{-/-}$. Lower panel

represents stable regions (lower than 20% difference in methylation). **B)** Pie charts representing
DMR numbers for each comparison. **C)** Heatmap of the methylation levels of tiles commonly
represented in all 9 public data samples that overlap with promoters-associated High-Mid DMRs
(n = 314 tiles) (left panel). Heatmap of the methylation level of gene associated with piRNA,
gamete generation and germ cells or meiotic nuclear division that are covered in all 9 public
datasets (right panel).
830