Developmental Genome-Wide DNA Methylation Asymmetry Between

Mouse Placenta and Embryo

Legault LM[1,2], Doiron K[1], Lemieux A[1,2], Caron M[1,3], Chan D[3], Lopes FL[4], Bourque G[5,6,7],
Sinnett D[1,8], McGraw S[1,2,9, #].

[1]Research Center of the CHU Sainte-Justine, Montreal, Canada.
[2]Department of Biochemistry and Molecular Medicine, Université de Montréal, Montreal,
Canada.
[3]Research Institute of the McGill University Health Centre, Montreal, Quebec, Canada
[4]São Paulo State University (Unesp), School of Veterinary Medicine, Aracatuba, Brazil
[5]Department of Human Genetics, McGill University, Montreal, Quebec, Canada
[6]McGill University and Genome Quebec Innovation Centre, Montreal, Quebec, Canada
[7]Canadian Center for Computational Genomics, Montreal, Quebec, Canada
[8]Department of Pediatrics, Université de Montréal, Montreal, Canada.
[9]Department of Obstetrics and Gynecology, Université de Montréal, Montreal, Canada.

[#]Corresponding author: serge.mcgraw@umontreal.ca

Running title: Embryonic & Placental DNAme Asymmetry

Summary statement: The kinetics of asymmetric DNA methylation acquisition of embryo-
placenta differentially methylated regions is not a stepwise process occurring in embryonic and
extraembryonic lineages, but a prompt progression in the early post-implanted conceptus.

ABSTRACT
28
When early embryo DNA methylation is remodelled to initiate the developmental program,
methylation marks are acquired unequally between embryonic and placental cells. To better
understand this process, we generated high-resolution DNA methylation maps of mouse mid-
gestation (E10.5) embryo and placenta. We uncovered specific subtypes of differentially
methylated regions (DMRs) contributing to this developmental asymmetry, which occurs rapidly
during the acquisition of marks in the post-implanted conceptus (E3.5-E6.5). These patterns are
long-lasting across subtypes of DMRs throughout prenatal development and in somatic tissues.
During peri-implantation stages, the *de novo* methyltransferase activity of DNMT3B is the main
driver of methylation marks, and DNMT3B can largely compensate for lack of DNMT3A in the
epiblast and extraembryonic ectoderm, whereas DNMT3A can only partially palliate in the
absence of DNMT3B. However, as development progresses and as DNMT3A becomes the
principal *de novo* methyltransferase, the compensatory DNA methylation mechanism of DNMT3B
on DMRs becomes less effective.

INTRODUCTION

Throughout the eutherian mammalian gestation, the placenta plays an essential role in mediating maternal–embryonic exchanges of gas, nutrients and waste, and also provides the developing embryo with a protective layer against adverse environmental exposures and the maternal immune system (Rossant & Cross, 2001). These unique placental functions are orchestrated by several distinct trophoblast cell subtypes organized in separate layers (Cross, 2000). The initial steps of lineage specialization of both placental and embryonic cells occur promptly following fertilization during the first few embryonic cleavages as DNA methylation marks are being reprogrammed (Morgan, Santos, Green, Dean, & Reik, 2005).

DNA methylation is an epigenetic mechanism that is critical in the determination of lineage-specific differentiation and development, and is mainly recognized for its involvement in processes such as transcriptional repression, genomic imprinting and X-inactivation (Bestor, 2000). DNA methylation marks are mediated by the action of DNA methyltransferases (DNMTs). Establishment of new or *de novo* DNA methylation patterns required for cell lineage determination during development is mediated by DNMT3A and DNMT3B, with cofactor DNMT3L, (Li, 2002; Okano, Bell, Haber, & Li, 1999), whereas DNMT1 maintains heritable DNA methylation patterns during cellular divisions (Lei et al., 1996; Leonhardt, Page, Weier, & Bestor, 1992). These enzymes are critical, as deletion of *Dnmt3b* or *Dnmt1* is embryonic lethal, while *Dnmt3a*-deficient offspring die shortly after birth (Li, Bestor, & Jaenisch, 1992; Okano et al., 1999). During gametogenesis, the acquisition of genome-wide and allele-specific methylation patterns (i.e. genomic imprinting) in both oocytes and sperm is essentially due to the activity of DNMT3A (Kaneda et al., 2004; Kato et al., 2007). Following fertilization, a reprogramming wave removes most methylation signatures across the genome, except for imprinted regions, some types of repeat sequences, as well as imprinted-like sequences, to trigger the developmental program (Hirasawa et al., 2008; Howell et al., 2001; McGraw et al., 2015). Then, during the peri-implantation process, DNA methylation profiles are re-acquired in a sex-, cell- and tissue-specific manner across most parts of the genome by the combined action of DNMT3A and DNMT3B. In the early stages of the *de novo* methylation wave (E4.5-E7.5), the expression of *Dnmt3b* is more robust than *Dnmt3a* in the epiblast and embryonic-derived cells (Auclair, Guibert, Bender, & Weber, 2014; Smith et al., 2017; Watanabe, Suetake, Tada, & Tajima, 2002), with the relative expression of *Dnmt3b* and *Dnmt3a* being considerably reduced in the extraembryonic ectoderm (ExE) and trophoblast

lineages (Senner, Krueger, Oxley, Andrews, & Hemberger, 2012; Smith et al., 2017). This
discrepancy in *Dnmt3a* and *Dnmt3b* expression levels coincides with the initiation of divergent
DNA methylation acquisition between the trophoblast and the inner cell mass of the blastocyst
(Fulka, Mrazek, Tepla, & Fulka, 2004; Guo et al., 2014; Monk, 1987; Nakanishi et al., 2012; Oda,
Oxley, Dean, & Reik, 2013; Santos, Hendrich, Reik, & Dean, 2002; Smith et al., 2014), a
difference that becomes extremely apparent by E6.5, as the epiblast has acquired most of its global
DNA methylation compared to the lower-methylation state of the ExE (Auclair et al., 2014; Smith
et al., 2017). This divergence is a common feature across mammalian placenta, as a heterogeneous
and lower-methylation state compared to somatic tissues and other cell types is constantly
observed (Chatterjee et al., 2016; Decato, Lopez-Tello, Sferruzzi-Perri, Smith, & Dean, 2017;
Schroeder et al., 2013; Smith et al., 2017).

Although the functional role of reduced methylation levels observed across the placental
genome is still not fully understood, studies suggest that it may activate transposable elements that
are typically silenced in other tissues (Chuong, 2013). DNA methylation plays an important role
in suppressing retrotransposons in mammalian cells, for which the activity has been associated
with genomic instability and disease development (Church et al., 2009; Slotkin & Martienssen,
2007). Following the *de novo* methylation wave, in embryonic-derived cells from the inner cell
mass, transposable elements acquire higher levels of DNA methylation causing transcriptional
silencing, whereas in the trophectoderm-derived cells that will form the placenta, these
transposable elements are maintained in a relaxed methylation state and preferentially expressed
(Okahara et al., 2004; Price et al., 2012; Warren et al., 2015). The low-methylation levels on these
elements contributed to the evolution and diversification of the placenta function through the
regulation of gene expression by providing placenta-specific enhancers, cryptic promoters and
other cis-regulatory elements (Cohen et al., 2011; Emera & Wagner, 2012; Haig, 2012; Macaulay,
Weeks, Andrews, & Morison, 2011; Mi et al., 2000; Xie et al., 2013).

Despite the evident distinction between embryonic and placental DNA methylation levels,
it remains unclear how, when and where methylation levels are acquired unequally across these
genomes. To better understand the developmental dynamics of epigenetic asymmetry that exists
between embryo and placenta, we first generated high-resolution maps of DNA methylation marks
using Reduced Representation Bisulfite Sequencing (RRBS) at mid-gestation, when the mouse
placenta is first considered mature (E10.5), to identify embryo-placenta differentially methylated

regions (DMRs). Then, using publicly available DNA methylation data sets and computational

analyses, we defined how various categories of DMRs are established in early stages and

maintained throughout development. In addition, we outlined the contribution of *Dnmt3a* and

Dnmt3b in the acquisition and maintenance of embryonic and extraembryonic specific DMR

patterns.

RESULTS

Increased Fluctuation in Genome-Wide DNA Methylation Levels in Placental Cells

To identify the overall epigenetic asymmetry that exists between placental and embryonic

genomes during mouse *in utero* development, we first established genome-wide DNA methylation

profiles using RRBS (Legault et al., 2019; Magnus et al., 2014; McGraw et al., 2015) of embryos

and their corresponding placentas at mid-gestation (E10.5), the developmental stage at which the

mouse placenta is considered mature (Cross, Werb, & Fisher, 1994). Using this approach, we

quantified the DNA methylation profiles of ~1.8 million CpG sites in each sample (embryo n=4,

placenta n=4). We found that the accumulation of CpG methylation was very distinct between the

placenta and the embryo. In the placenta, we detected a greater proportion of CpGs in the 0-50%

methylation range and a lower proportion of CpGs in the 80-100% methylation range (Figure 1A).

Although we observed that a large proportion of CpGs within the examined regions had no

methylation marks in both the embryo and the placenta, interestingly, most CpGs with partial

methylation (20-50%) in placenta showed high methylation levels (>80%) in embryos (Figure 1B).

The divergence in global DNA methylation profiles conferred a high degree of clustering between

both tissue types (Figure S1). These results are consistent with previous studies indicating lower

levels of overall DNA methylation in extraembryonic tissues (Price et al., 2012; Schroeder et al.,

2013; Smith et al., 2017), reviewed in (Robinson & Price, 2015).

To further investigate the dynamics of DNA methylation between placental and embryonic

genomes and enable direct comparison of precise regions, we segmented the genome of autosomal

chromosomes into 100bp non-overlapping genomic windows (*tiles*; see methods section). After

removal of sex chromosomes, we identified 245 048 unique sequenced tiles (referred to as All-

tiles) containing 896 820 common CpGs between all placenta and embryo samples and with a

minimum of 15x sequencing depth. We observed a strong reduction in the fraction of highly

methylated tiles (80-100%) in the placenta compared to the embryo (Figure 1C), which correlated

with a sharp increase in the number of placenta tiles in the 0-20%, 20-40% and 40-60%
methylation range. Globally, we found that the average DNA methylation level across all placental
tiles was significantly lower compared to all embryo tiles (27% vs 45%, p<0.0001) (Figure 1C).
This overall epigenetic disparity in embryo and placenta DNA methylation levels was especially
noticeable when we mapped the methylation mean of All-tiles with respect to regions surrounding
the transcription start sites (TSS) (Figure 1D). When we focused on a specific chromosome section
(e.g., chr 7, 35Mb) (Figure 1E), we observed that genomic segments with high DNA methylation
levels in embryos were predominantly hypomethylated in placenta. We also observed that gene or
CpG island (CGI) poor regions had consistently high methylation levels in embryos and lower
methylation levels in the placenta. Together, these results indicate that the mid-gestation placenta
has very distinctive global DNA methylation profiles compared to the embryo, and that this lower
level of global DNA methylation (i.e., DNA hypomethylation) across the placental genome is due
to a significant lower number of highly methylated (≥80%) genomic regions. We can also conclude
that despite the cellular heterogeneity in E10.5 placental and embryonic tissues, the vast majority
of the conceptus possesses specific genomic regions with either low (0-20%) or high (80-100%)
levels of methylation. However, the placenta genome presents an increased number of regions
having a broader distribution of DNA methylation (20-80%), revealing a greater diversity in
methylation levels across placental cell types compared to embryonic cell types.

Embryonic and Placental DNA Methylation Divergences Across Genomic Features

To explain the developmentally divergent methylated states between the mouse embryo
and the placenta, we next sought to precisely determine the genomic features revealing DNA
methylation differences. We defined differentially methylated regions (DMRs) as 100bp genomic
segments showing a significant difference of methylation levels between embryonic and placental
samples with an absolute mean methylation difference of 20% or higher (McGraw et al., 2015;
Piche et al., 2019; Shaffer et al., 2015). Using these conditions, we screened the 245 048 unique
tiles common between all samples and identified 110 240 DMRs (~45% of All-tiles;
Supplementary Table 1) with tissue- and/or developmental-specific DNA methylation variations
between the embryo and the placenta (Figure2A; random subset of 20,000 DMRs shown).
Consistent with our findings (Figure 1), the majority (96.8%; n=106 712) of DMRs had lower
DNA methylation in the placenta (referred to as Hypo-DMRs) and only a small proportion (3.2%;

n=3 528) showed increased methylation levels (referred to as Hyper-DMRs) compared to the
embryo (Figure S2). For Hypo-DMRs, tiles mainly overlapped (96%; n=102 982) with intergenic,
intron, exon and promoter-TSS regions (Figure S2). For each of the genomic feature categories of
Hypo-DMRs, the average DNA methylation levels in the placenta were essentially half of those
present in the embryo (Figure 2B). As for DMRs with higher methylation levels in the placenta
(Hyper-DMRs), we noticed that the vast majority of these tiles (83%; n=2 929) had very low
methylation levels in the embryo (<20%) (Figure 2A-B). Most of Hyper-DMRs overlapped
intergenic, intron, exon and promoter-TSS regions (93%; n=3 272, Figure S2). We then
investigated the relationship between gene expression and DMR DNA methylation levels. Highly
expressed genes in placental tissue such as *Syna* (*Syncytin A*) (Mi et al., 2000) showed overall
lower methylation levels in the placenta when compared to the embryo (Figure 2D, smoothed
representation) (Hansen, Langmead, & Irizarry, 2012). Similar observations were made in the gene
body of *Atf6b* (*Activating Transcription Factor 6 Beta*), a gene implicated in the transcriptional
downregulation of *Pgf* (*Placental Growth Factor*) in response to endoplasmic reticulum stress in
pathological placentas (Mizuuchi et al., 2016). As for genes predominantly expressed in the
embryo, *Mir219a-2/Mir219b*, a brain-specific non-coding microRNA, showed higher methylation
in the placenta (Figure 2D). It was recently shown that *Mir219* was abnormally expressed during
pregnancy in amnion retrieved specifically from obese women (Nardelli et al., 2014) and,
interestingly, obesity has been suggested to cause epigenetic changes (Lillycrop & Burdge, 2011).
Another example of placenta Hyper-DMRs is *Sox6* (*SRY-Box 6*), which is implicated in the
terminal differentiation of muscle (Figure 2D) (Kamachi & Kondoh, 2013). These observations
point to an association between tissue-specific DNA methylation levels of DMRs and biological
functions. Gene ontology enrichment analyses further support that Hypo-DMRs were strongly
associated to germline functions and reproduction (*e.g.*, male and female gamete generation,
reproduction, piRNA metabolic process, meiotic cell cycle, germ cell development) (Figure 2C).
As for Hyper-DMRs, top biological processes were mostly associated with developmental and
differentiation processes (*e.g.*, regionalization, embryo development, pattern specification
process, skeletal system, head development) (Figure 2C). Fittingly, the biological functions were
completely divergent between Hypo- and Hyper-DMRs. Altogether, these results denote that
Hypo- and Hyper-DMRs are present across genomic features between mid-gestation embryo and

placenta, and that these DNA methylation divergences are implicated in promoting/repressing
specific processes during embryonic and placental development.

Presence of Distinctive DMR Categories between Mid-Gestation Embryo and Placenta

Amongst DMRs, our analyses also suggest the presence of particular DMR categories
based on their level of DNA methylation in the embryo and placenta (Figure 2A). By defining
subsets of DMRs and establishing their dynamic properties between tissues, we might better
understand the genome-wide asymmetry in DNA methylation levels observed between the embryo
and the placenta. To do so, we first clustered DMR-associated tiles in 6 different categories based
on their range of low, mid and high methylation level (Low; <20%, Mid; ≥20 to <80%, High;
≥80%) in the embryo and the placenta, and followed the DMR category transitions between both
tissues. We observed that DMRs with High-levels of methylation in the embryo overlapped with
a large proportion of DMRs that showed Mid-levels of methylation in the placenta (Figure 3A,B;
High-Mid n=72 715), whereas only a fraction corresponded to DMRs with Low-levels of
methylation in the placenta (Figure 3A,B; High-Low n=1 889). As for DMRs with Mid-levels of
methylation in embryo, the largest part remained in that same Mid-levels category in the placenta
(Figure 3A; Mid-Mid n=23 640). Nonetheless, a portion of these embryonic Mid-levels DMRs
were directed to DMRs with either Low- (Figure 3A; Mid-Low n=9 055) or High- (Figure 3A;
Mid-High n= 21) levels of methylation in the placenta. Finally, DMRs with Low-levels of
methylation in the embryo all showed Mid-levels of methylation in the placenta (Figure 3A; Low-
Mid n=2 920). Thus, when we subdivide our DMRs into distinctive categories, we uncover that
embryonic cells possess a large proportion of DMRs of High-level (≥80%) of DNA methylation,
which remain potentially static in embryonic tissues, whereas the vast majority of these DMRs
have inferior and wide-range DNA methylation levels within placental tissue.

Specific Genomic Features Associated with DMR Categories

We next aimed to determine if the genomic distribution of embryo-placenta DMR
categories was associated with distinct genomic features. First, by classifying by genomic
annotations, we observed that DMRs with reduced methylation levels in the placenta (High-Mid,
High-Low, Mid-Low) were prevailingly found in intergenic regions (>50% of tiles) (Figure 3C),
whereas DMR categories with equivalent (Mid-Mid) or superior methylation level (Mid-High,

Low-Mid) in the placenta were more frequent in genic associated regions (>50% of tiles).
However, divergence between DMR categories was observed when we performed ontology
analyses on promoter regions (Figure S3), as each DMR subtype clearly showed distinct biological
functions. For example, High-Mid DMRs top functions were related to genes (e.g., *Aszl, Tnp1,*
Piwil2) associated with male gamete generation, piRNA metabolic process and male meiotic
nuclear division, whereas those of the Low-Mid category were mostly associated with
developmental aspects (e.g., skeletal system development, regionalization, nervous system
development). Since we know that in placenta, activation of retrotransposon-derived genes is
interrelated with low DNA methylation levels (Cohen et al., 2011; Macaulay et al., 2011; Reiss,
Zhang, & Mager, 2007), we next assessed how DMR categories overlapped with major types of
retrotransposons (LINE; long interspersed nuclear elements, SINE; short interspersed nuclear
element, and LTR; long terminal repeats). Out of the DMR categories, those with High-levels of
methylation in the embryo and either Mid- or Low-levels in the placenta (High-Mid and High-
Low) showed the most enriched overlap with retrotransposons, with 92% and 96% respectively
(Figure 3D). DMR categories associated with gain of methylation in the placenta (Mid-High and
Low-Mid) showed the least overlap with retrotransposons, especially the Low-Mid subtype. This
highlights that during the *de novo* acquisition of DNA methylation patterns, DMRs with High-
levels of methylation in the embryo and lower levels in the placenta are almost exclusively within
retrotransposons-associated sequences, whereas DMRs with Low-levels of methylation in the
embryo and higher DNA methylation in the placenta are preferentially outside retrotransposons-
associated sequences. Finally, we investigated the proximity of the DMR categories in regards to
CpG rich (CpG islands; CGI), neighbouring (shore; < 2kb away from CGIs, shelf; 2-4kb away
from CGIs) and distant (open sea; > 4kb away from CGIs) regions (Figure 3E). We observed that
most DMR categories are depleted from CGIs and are mostly found in open sea regions. In
contrast, ~60% of tiles in Low-Mid DMRs overlapped with CGI, shores and shelves, revealing
that the acquisition of *de novo* methylation for these genomic fragments in the extraembryonic
lineage preferentially targets sequences inside or surrounding CGIs.
Altogether, these results indicate that the asymmetry within DMR categories, based on their
methylation levels in the embryo and the placenta, can be associated to specific biological
functions and genomic-derived features (e.g., CpG, retrotransposon contents). This is particularly

apparent for DMRs with High-levels in the embryo (High-Mid, High-Low) and those with higher
methylation levels in the placenta (Low-Mid).

DMR Categories are Established During the De Novo Methylation Wave and Maintained
Throughout Development

To gain insights into the kinetics of lineage-specific DMR establishment between mid-
gestation embryo and placenta, as well as their status during development, we assessed the levels
of methylation associated with tiles for each DMR category as a function of their developmental
stage. Publicly available sequencing data (Smith et al., 2012; Whidden et al., 2016) were analyzed
using our custom script to generate 100bp tiles, and the DNA methylation levels for each tile were
calculated. In E3.5 blastocysts, when the mouse genome is mostly depleted from DNA methylation
marks, we observed low global DNA methylation levels (average <20%) for all DMR subtypes,
with similar median levels between the committed cells of the inner cell mass (ICM) and
trophectoderm lineages (Figure 4). DNA methylation levels tended to be higher in the
trophectoderm for regions falling in the Mid-High DMRs, although measurements are based on
very few DMRs for this specific category (n=21, Figure 3B, Mid-High). Since global DNA
methylation is re-acquired in the next few subsequent developmental stages, we then asked
whether the contrast in DNA methylation levels associated with the various DMR categories at
mid-gestation would already be present between E6.5 epiblast and extraembryonic ectoderm (ExE)
cell lineages, layers that are mostly composed of homogeneous and undifferentiated cell
populations. For all DMR categories, DNA methylation levels in the E6.5 epiblast and ExE already
showed similar pattern trends to those observed in the E10.5 embryo and placenta (Figure 4, Table
S2). Interestingly, for all DMR categories we observed slightly higher DNA methylation levels in
E6.5 ExE compared to E10.5 placenta (Figure 4, Table S2). However, when measured in the
subsequent developmental stages (E10.5, E11.5, E15 and E18), levels associated with each
category stabilized and remained in the same methylation range. This is true for all categories,
except for the Mid-High DMRs where methylation levels are reduced by ~20% at E18. As for the
E6.5 epiblast DNA methylation profiles, they closely matched those observed in the E10.5 embryo,
except again for the Mid-High DMRs. For each of the DMR subtypes, the DNA methylation levels
were largely stable and persisted across other time points (E6.5, E10.5, E11.5) in embryonic cells.
Although no public DNA methylation data were available for whole embryos at later stages, when

we overlapped tiles associated with E10.5 DMR categories with data from differentiated tissues of

adult mice, the global DNA methylation profiles closely matched those for the majority of DMR

categories (High-Mid, High-Low, Mid-Mid and Mid-Low) (Figure 4, Figure S4). We conclude

that the various DMR categories observed at E10.5 are established during the embryonic and

extraembryonic lineage-specification processes occurring during the peri-implantation wave of *de*

novo methylation, and that these DNA methylation landscapes are widely retained throughout

embryo and placenta development. Furthermore, for most DMR categories, the DNA methylation

levels observed in developing embryos are long-lasting and conserved throughout somatic cell

differentiation.

Dnmt3a- or Dnmt3b-Deficiency Alters Proper Establishment of DMR-Associated Patterns

Since the combined activity of DNMT3A and DNMT3B is essential for proper

establishment of DNA methylation profiles and normal development, we next sought to define the

contribution of each enzyme in the *de novo* establishment of DNA methylation in subtypes of

DMRs in embryonic and extraembryonic cell lineages. To do so, we used DNA methylation data

from publicly available datasets (whole genome bisulfite sequencing and RRBS) at E6.5 (epiblast

and ExE) and E8.5 (embryo) with various inactive forms of *Dnmts*. First, when we overlapped our

All-tiles subset, we observed a substantial reduction (p<0.0001) in average DNA methylation in

absence of DNMT3A (37.81%) or DNMT3B (31.48%) in the E6.5 epiblast compared to wild-type

(44.81%), whereas in the E6.5 ExE such a comparable loss was only associated with a *Dnmt3b*-

deficiency (wt: 34.38% vs *Dnmt3b*$^{-/-}$: 19.33%, p<0.0001) (Figure S5). Similarly, in the E8.5

embryo, a significant reduction in average DNA methylation level was measured with lack of

Dnmt3a or *Dnmt3b* expression compared to wild-type (wt: 47.5%; *Dnmt3a*$^{-/-}$: 43.6%; *Dnmt3b*$^{-/-}$:

34%, p<0.0001) (Figure S5). For most DMR subtypes, absence of DNMT3A caused modest or no

reduction on overall DNA methylation levels in E6.5 epiblast and ExE (Figure 5A and S6, Table

S3). Intriguingly, *Dnmt3a*-deficiency caused a slight gain of methylation (wt: 76.9% vs *Dnmt3a*$^{-/-}$

: 81.5%, p<0.0001) level in the E6.5 epiblast for High-Mid DMRs, suggesting a potential

overcompensation effect by DNMT3B in the epiblast on this specific DMR subtype. In

comparison, *Dnmt3b*-depletion led to a substantial loss of average methylation levels in all DMR

categories in the E6.5 epiblast and ExE, except for epiblast High-Mid DMRs (Figure 5A, Table

S3) for which a compensatory mechanism provided high levels of methylation (wt: 76.9% vs

Dnmt3b^{-/-}: 71.6%). The compensatory mechanism on High-Mid DMRs in response to *Dnmt3b-*

deficiency was ineffective in the E6.5 ExE. When we focused on promoter-TSS for each DMR

categories, we observed again that loss of methylation for these regulatory regions was principally

associated with *Dnmt3b*-deficiency (Figure 5B). The promoter-TSS regions associated with High-

Mid DMRs retained relatively high methylation levels for either *Dnmt3a*^{-/-} or *Dnmt3b*^{-/-} epiblast

samples, demonstrating a robust and compensatory *de novo* methylation mechanism. To further

underline the impact of DNMT3A or DNMT3B on the *de novo* methylation of DMRs, we

measured methylation levels for DMRs selected from our gene enrichment analyses (*Fgb*, *P2rx7*,

Pcyt2, *Etnppl*, *Ralgds*, *Lrp5*) and other genomic segments covered by multiple tiles (*Cxxc1*, *Fcgrt*,

Lamp5, *Mbd1*, *Lphn1*, *Pick1*, *Irf1*) (Figure S7 & S8) in *Dnmt3a-* or *Dnmt3b*-deficient epiblast and

ExE. In line with our global observations, for most DMR-associated tiles, a *Dnmt3b*-deficiency in

E6.5 epiblast or ExE caused a more severe loss of methylation compared to lack of *Dnmt3a.*

However, for some regions in the E6.5 epiblast (e.g., *Pick1*, *Irf1*), *Dnmt3a*^{-/-} methylation levels

were inferior to those of *Dnmt3b*^{-/-}. Noteworthy, for most DMRs, lack of co-factor *Dnmt3l* affected

DNA methylation levels to greater extent than *Dnmt3a*-deficiency (Figure S6). Overall, we show

that of DNMT3A and DNMT3B participate in the establishment of the asymmetric methylation

patterns associated with the various DMR categories in both embryonic and extraembryonic cells,

with DNMT3B being the principal contributor in both cell lineages during the *de novo*

reprogramming wave as it can compensate almost entirely for the absence of DNMT3A.

Decline in Compensatory DNA Methylation Mechanisms in Response to Dnmt3a- or Dnmt3b-

Deficiency.

As embryonic development progresses from E6.5 to E8.5, our data suggest that lack of

DNMT3A or DNMT3B further reduces global methylation levels of DMRs (High-Mid, High-

Low, Mid-Low and Mid-Mid), evoking that absence of either enzymatic activity causes an additive

and extended effect (Figure 5A-B). To further define the robustness in compensatory mechanism

between DNMT3A or DNMT3B in the establishment and maintenance of DNA methylation on

DMRs categories, we focused on High-Mid promoter-TSS- associated DMRs as they have the

highest DNA methylation levels and require the most *de novo* methyltransferase activity.

Methylation differences of $\geq 20\%$ between wild-type and *Dnmt3a-* or *Dnmt3b*-deficient samples

were considered as regions showing substantial lack of compensation. In agreement with our

results (Figure 5A-B), we observed that at E6.5, DNMT3B can compensate almost entirely for

DNMT3 Aloss by maintaining methylation levels on most promoters-TSS associated tiles in the

epiblast (410/437 = 93.8%) and ExE (558/602 = 92.7%), whereas DNMT3A can only partially

alleviate the lack of DNMT3B in the epiblast (315/410 = 78.6%) and ExE (244/524 = 46.6%)

(Figure 6A-B). As embryonic cell lineages development evolves between E6.5 and E8.5, global

methylation levels increase on promoter-TSS of High-Mid DMRs. However, we detected a sharp

decline in the compensatory mechanism to rescue the *de novo* and/or maintenance of methylation

levels of promoter-TSS DMRs in the *Dnmt3a-* (524/627 = 80.9%) or *Dnmt3b-* (256/622 = 41.2%)

deficient E8.5 embryos. This is further highlighted when we focus on a subgroup of 314 promoter-

TSS associated tiles overlapping all of 9 data sets (Figure 6C), including gene promoters (e.g.,

Asz1, Catsper1, Ccdc42, Dmrtb1, Piwil2, Rpl10l, Sox30, Sycp1, Tnp1, Ttll1, Zfn42) related to our

top enriched biological functions (i.e., piRNA, gamete generation and germ cells, meiotic nuclear

division). Although E8.5 *Dnmt3a-* or *Dnmt3b*-deficient ExE data was not available, our data

suggest that greater compensation failure would also be observed in this tissue. Thus, embryonic-

placental DMRs need the combined action of DNMT3Aand DNMT3Bto both establish and

maintain proper asymmetric levels during early development as compensatory DNA methylation

mechanisms during the *de novo* wave of methylation decline overtime and fail to overcome

methyltransferase shortage in later development stages.

DISCUSSION

With recent breakthroughs in high-throughput sequencing, we now have a better

understanding of the dynamic of DNA methylation erasure occurring in early cleavage stage

embryos following fertilization. However, the discrepancies in acquisition of genome-wide DNA

methylation patterns in early post-implantation embryonic and extraembryonic cell lineages

remain to be methodically delineated. To address this issue and further our understanding of the

DNA methylation asymmetry that guides the developmental trajectory of embryonic and

extraembryonic cell lineages, we established genome-wide DNA methylation profiles of mouse

embryo and placenta at mid-gestation, and analyzed various publicly available developmental

stage specific embryo and placenta DNA methylation datasets. Using this strategy, we uncovered

that 45% of the genomic regions analyzed differ in DNA methylation status ($\geq$20%) between mid-

gestation embryo and placenta, and that these DMRs can be further divided into categories based

on their levels of DNA methylation (Low; <20%, Mid; ≥20 to <80%, High; ≥80%) in the embryo

and placenta. We show that the embryo and placenta acquire specific DMR categories during the

early stage of the *de novo* DNA methylation wave, and that these DMRs persist throughout prenatal

development, as well as into somatic adult tissues. Furthermore, we show that *Dnmt3b* primarily

drives the divergence in DNA methylation levels associated with these specific DMRs and that *de*

novo methyltransferase activity of *Dnmt3b* can almost entirely compensate for lack of *Dnmt3a* in

the asymmetric establishment of embryonic and placental DMRs, but that *Dnmt3a* can only

partially palliate the absence of *Dnmt3b*. However, with developmental progression, this

compensatory DNA methylation mechanism becomes less effective.

Our results indicate that the kinetics of DNA methylation acquisition leading to specific

embryo-placenta DMR categories is not a stepwise process occurring throughout cell fate

decisions and patterning of embryonic and extraembryonic lineages, but a prompt progression in

the early post-implanted conceptus. These results are in line with studies reflecting that the

initiation of asymmetric DNA methylation levels begins within the trophoblast and the inner cell

mass of the blastocyst (Santos et al., 2002), and becomes highly evident by the time the epiblast

acquires its initial global DNA methylation patterns at E6.5 during the *de novo* reprogramming

wave (Auclair et al. 2014, Smith et al. 2017). Our results show that the acquisition period between

E4.5 and E6.5 is particularly key to establishing asymmetric DNA methylation patterns associated

with mid-gestation embryo-placenta DMR classes, and that these DMR associated-patterns are

long-lasting across stages of prenatal development. Since we studied cell populations derived from

embryonic and placental tissues, we cannot dismiss the prevalence of cell-to-cell heterogeneity in

the acquisition kinetics of specific DMR patterns. Despite this concern, our results indicate that

embryonic cells have a large body of DMRs with High-levels (≥80%) and Low-levels (<20%) of

methylation that remain static across development, as well as in somatic cell types (High-Mid &

High-Low DMRs), whereas compared to the embryo, the placental DMRs have overall lower

methylation levels and are more broadly distributed amongst methylation levels, which has also

been shown in other studies where genome-wide DNA methylation profiles of placental cells were

compared to other tissues and specific cell types (Schroeder et al., 2013; Chatterjee et al., 2016;

Smith et al., 2017). Lower methylation levels in the placenta have been associated with reduced

de novo methyltransferase activity in the ExE during the *de novo* reprogramming wave (Fulka et

al., 2004; Santos et al., 2002). Nevertheless, we observe methylation level peaks for all DMR

categories within the ExE at E6.5 before levels stabilize at E10.5. As of now, the implication of
these methylation level peaks on future regulation mechanisms and methylation profiles is
unknown. Moreover, it remains to be determined whether global reduction in DNA methylation
marks on DMRs between the E6.5 and E10.5 extraembryonic cell lineages is stochastic or targeted,
and whether it occurs through passive or active mechanisms.

We also observed an enrichment of retrotransposons (i.e., LINE, SINE, LTR) in DMRs
especially in those with High-levels of methylation in the embryo and lower-levels in the placenta,
whereas DMRs with Low-levels of methylation in the embryo and higher levels in the placenta
are preferentially outside retrotransposons-associated sequences. Earlier findings of Chapman et
al. (Chapman, Forrester, Sanford, Hastie, & Rossant, 1984) revealed that repeat regions in the
placenta appear to lack tight control of their methylation patterns, perhaps indicating that
maintaining methylation, and therefore repression of these elements for genome stability and
integrity, is not critical given the relatively short lifespan of this organ. However, we do observe
Mid-range methylation levels (20-80%) in the placenta for a substantial portion of DMRs that are
associated with retrotransposons, revealing that specific genomic regions associated with
repetitive elements do need tight regulation in extraembryonic cell lineages for proper
development. These results are in line with findings exposing that the deletion of genome-defense
gene *Tex19.1* leads to the de-repression of LINE1 and compromises placental development,
suggesting that disparities between retrotransposon suppression and genome-defense mechanisms
might contribute to placenta dysfunction and disease (Reichmann et al., 2013).

DNMT3A and DNMT3B are required to establish proper methylation profiles on the
embryonic genome during the *de novo* reprogramming wave, as both methyltransferase enzymes
have redundant, but also specific functions. However, the activity of DNMT3B is the main
contributor in the acquisition of profiles in epiblast cells, and especially commands methylation
on CGIs associated with developmental genes (Auclair et al., 2014). Auclair et *al*. highlighted that
in the absence of DNMT3B, DNMT3A was not able to counterbalance, leading to the loss of
promoter-CGI methylation and gain of expression of germline genes (e.g., *Sycp1*, *Sycp2*, *Mael*,
Rpl10l, *Dmrtb1*) in somatic cells of the embryo. Here, we show that the promoter of these germline
genes, associated with meiotic and piRNA processes as well as other genes with similar biological
functions, are highly enriched in High-Mid DMRs. We also reveal that for this particular set of
High-Mid DMRs, mainly deprived of CGIs, DNMT3A is able to substantially counterbalance the

lack of DNMT3B in the E6.5 epiblast and establish global DNA methylation levels, whereas in

the ExE, this compensatory mechanism is rather inefficient. Globally, we show that DNMT3B is

much more potent at compensating than DNMT3A in both the epiblast and ExE for all DMR

categories, indicating that DNMT3B is the main *de novo* enzyme driving asymmetric DNA

methylation patterns between the embryo and placenta. Although compensatory mechanisms have

been observed in *Dnmt3a* or *Dnmt3b*-depleted conceptuses, we still do not fully understand the

process as *Dnmt3a* and *Dnmt3b* have cell lineage specific expression during the peri-implantation

developmental period. As development progresses, we observed that the compensation mechanism

in *Dnmt3a* or *Dnmt3b*-deficient embryos remains apparent at E8.5 on most DMR categories, but

is less effective as the methylation gaps increase compared to wild-type. This correlates with a

developmental period where DNMT3A is now the main *de novo* methyltransferase enzyme in both

the embryo and placenta (Okano et al., 1999; Watanabe et al., 2002), and with a compensatory

activity of DNMT3A being less effective. This suggests that DNMT3B activity is critical to ensure

proper establishment of DNA methylation asymmetry between the embryonic and extraembryonic

cell lineages during the *de novo* reprogramming wave, but that DNMT3A is required during the

developmental progression to safeguard methylation levels on DMR categories.

CONCLUSION

We demonstrate that asymmetry between embryo and placenta DNA methylation patterns

occurs rapidly during *de novo* acquisition of methylation marks in the early post-implanted

conceptus, and that these patterns are long-lasting across subtypes of DMRs. We also reveal that

at the peri-implantation stages, *de novo* methyltransferase activity of DNMT3B is the main

provider of asymmetric methylation marks on DMRs, and that it largely palliates the lack of

DNMT3A in the epiblast and ExE. However, as development progresses, DNMT3A becomes the

principal *de novo* methyltransferase by mid-gestation, and DNMT3B methyltransferase activity is

less effective at promoting compensation. These results further underline why embryos developing

without DNMT3B have severe DNA methylation defects and die at mid-gestation, whereas those

without DNMT3A only die postnatally. Further investigation is required to determine the

molecular mechanisms controlling the precise *de novo* acquisition of long-lasting methylation

marks on specific DMR subtypes in the embryonic and extraembryonic cell lineages, and how

errors in this process could lead to abnormal development and diseases.

METHODS

Animals and Sample Collection

Female C57BL/6 (8-10 week-old) were purchased from Harlan Sprague-Dawley Laboratories (Indianapolis, IN) and were mated with male C57BL/6 (4 months of age). Following natural mating, embryos and placental tissues were collected at E10.5 (presence of vaginal plug at E0.5). Samples were frozen immediately in liquid nitrogen and stored at -80 °C until analyzed.

DNA Methylation Analyses

RRBS libraries were generated as published protocols (Boyle et al., 2012; Gu et al., 2011) with our specifications (Legault, Chan, & McGraw, 2019; Magnus et al., 2014; McGraw et al., 2015). 500 ng of extracted DNA (Qiagen) from placenta (male n=2, female n=2) and embryo (male n=2, female n=2) samples was *MspI* digested, adaptor ligated and PCR amplified (multiplex). Multiplexed samples were pooled and 100 bp paired-end sequenced (HiSeq-2000, Illumina). The data analyses were done according to the pipeline established at the McGill Epigenomics Mapping and Data Coordinating Centers (Magnus et al., 2014; McGraw et al., 2015) that include BSMAP and methylKit. Specific parameters were chosen including 100 bp step-wise tiling windows, containing a minimum of 2 CpGs per tile and a minimum $15\times$ CpG coverage of each tile per sample. The methylation level of a 100-bp tile was the result of all CpG C/T read counts within the tile after coverage normalization between samples, and the methylation level reported for a sample on autosomal chromosomes was the average methylation level across replicates. Significant DNA methylation changes were designated as $\pm\geq20\%$ average differences between groups of replicates and a q-value < 0.01 using the logistic regression function of methylKit (Akalin et al., 2012). Direct comparisons between DNA methylation averages were done using Wilcoxon-Mann-Whitney test in R. Gene ontology (GO) terms and pathway analyses for RefSeqs associated to promoter-TSS were conducted using Metascape gene annotation and analysis resource (Zhou et al., 2019).

Sequencing Data

Publicly available DNA methylation data sets (Auclair et al., 2014; Decato et al., 2017; Hon et al., 2013; Smith et al., 2014; Smith et al., 2017; Whidden et al., 2016) were analyzed using

a custom script to intersect single CpG site methylation calls from these data sets within defined

100bp tiles associated to the embryo-placenta DMRs categories

ACKNOWLEDGMENTS

We thank the McGraw lab for critical comments and suggestions, as well as Elizabeth Maurice-

Elder for editing.

COMPETING INTERESTS

No competing interests declared

FUNDING

This work was supported by a research grant from the Natural Sciences and Engineering Research

Council of Canada. We acknowledge the Fonds de Recherche du Québec en Santé (S.M, L.M.L)

and the Réseau Québécois en Reproduction and Fonds de recherche du Québec – Nature et

technologies (K.D) for salary awards.

DATA AVAILABILITY

The data from this study have been submitted to the Gene Expression Omnibus (#GSE95610).

Reviewer link:

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?token=ghsnmqoctlgjlcj&acc=GSE95610

Table S1. Annotation and DNA methylation values associated with DMRs.

Table S2. DNA methylation levels for DMR categorises across development.

Table S3. DNA methylation levels for DMR categories in various *Dnmt* knockout mouse

models.

Table S4. Number of 100bp tiles in each category of public data that overlap with Embryo-

Placenta DMRs (related to Figure 4; 5A; S4; S5; S6)

REFERENCES

Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F. E., Figueroa, M. E., Melnick, A., &
Mason, C. E. (2012). methylKit: a comprehensive R package for the analysis of genome-
wide DNA methylation profiles. *Genome Biol, 13*(10), R87. doi:10.1186/gb-2012-13-10-
r87

Auclair, G., Guibert, S., Bender, A., & Weber, M. (2014). Ontogeny of CpG island methylation
and specificity of DNMT3 methyltransferases during embryonic development in the
mouse. *Genome Biol, 15*(12), 545. doi:10.1186/s13059-014-0545-5

Bestor, T. H. (2000). The DNA methyltransferases of mammals. *Hum Mol Genet, 9*(16), 2395-
2402.

Boyle, P., Clement, K., Gu, H., Smith, Z. D., Ziller, M., Fostel, J. L., . . . Meissner, A. (2012).
Gel-free multiplexed reduced representation bisulfite sequencing for large-scale DNA
methylation profiling. *Genome Biol, 13*(10), R92. doi:10.1186/gb-2012-13-10-r92

Chapman, V., Forrester, L., Sanford, J., Hastie, N., & Rossant, J. (1984). Cell lineage-specific
undermethylation of mouse repetitive DNA. *Nature, 307*(5948), 284-286.

Chatterjee, A., Macaulay, E. C., Rodger, E. J., Stockwell, P. A., Parry, M. F., Roberts, H. E., . . .
Morison, I. M. (2016). Placental Hypomethylation Is More Pronounced in Genomic Loci
Devoid of Retroelements. *G3 (Bethesda), 6*(7), 1911-1921. doi:10.1534/g3.116.030379

Chuong, E. B. (2013). Retroviruses facilitate the rapid evolution of the mammalian placenta.
Bioessays, 35(10), 853-861. doi:10.1002/bies.201300059

Church, D. M., Goodstadt, L., Hillier, L. W., Zody, M. C., Goldstein, S., She, X., . . . Mouse
Genome Sequencing, C. (2009). Lineage-specific biology revealed by a finished genome
assembly of the mouse. *PLoS Biol, 7*(5), e1000112. doi:10.1371/journal.pbio.1000112

Cohen, C. J., Rebollo, R., Babovic, S., Dai, E. L., Robinson, W. P., & Mager, D. L. (2011).
Placenta-specific expression of the interleukin-2 (IL-2) receptor beta subunit from an
endogenous retroviral promoter. *J Biol Chem, 286*(41), 35543-35552.
doi:10.1074/jbc.M111.227637

Cross, J. C. (2000). Genetic insights into trophoblast differentiation and placental
morphogenesis. *Semin Cell Dev Biol, 11*(2), 105-113. doi:10.1006/scdb.2000.0156

Cross, J. C., Werb, Z., & Fisher, S. J. (1994). Implantation and the placenta: key pieces of the
development puzzle. *Science, 266*(5190), 1508-1518. doi:10.1126/science.7985020

Decato, B. E., Lopez-Tello, J., Sferruzzi-Perri, A. N., Smith, A. D., & Dean, M. D. (2017). DNA
Methylation Divergence and Tissue Specialization in the Developing Mouse Placenta.
Mol Biol Evol, 34(7), 1702-1712. doi:10.1093/molbev/msx112

Emera, D., & Wagner, G. P. (2012). Transposable element recruitments in the mammalian
placenta: impacts and mechanisms. *Brief Funct Genomics, 11*(4), 267-276.
doi:10.1093/bfgp/els013

Fulka, H., Mrazek, M., Tepla, O., & Fulka, J., Jr. (2004). DNA methylation pattern in human
zygotes and developing embryos. *Reproduction, 128*(6), 703-708.
doi:10.1530/rep.1.00217

Gu, H., Smith, Z. D., Bock, C., Boyle, P., Gnirke, A., & Meissner, A. (2011). Preparation of
reduced representation bisulfite sequencing libraries for genome-scale DNA methylation
profiling. *Nat Protoc, 6*(4), 468-481. doi:10.1038/nprot.2010.190

Guo, H., Zhu, P., Yan, L., Li, R., Hu, B., Lian, Y., . . . Qiao, J. (2014). The DNA methylation
landscape of human early embryos. *Nature, 511*(7511), 606-610.
doi:10.1038/nature13544

Haig, D. (2012). Retroviruses and the placenta. *Curr Biol, 22*(15), R609-613.
doi:10.1016/j.cub.2012.06.002
Hansen, K. D., Langmead, B., & Irizarry, R. A. (2012). BSmooth: from whole genome bisulfite
sequencing reads to differentially methylated regions. *Genome Biol, 13*(10), R83.
doi:10.1186/gb-2012-13-10-r83
Hirasawa, R., Chiba, H., Kaneda, M., Tajima, S., Li, E., Jaenisch, R., & Sasaki, H. (2008).
Maternal and zygotic Dnmt1 are necessary and sufficient for the maintenance of DNA
methylation imprints during preimplantation development. *Genes Dev, 22*(12), 1607-
1616. doi:10.1101/gad.1667008
Hon, G. C., Rajagopal, N., Shen, Y., McCleary, D. F., Yue, F., Dang, M. D., & Ren, B. (2013).
Epigenetic memory at embryonic enhancers identified in DNA methylation maps from
adult mouse tissues. *Nat Genet, 45*(10), 1198-1206. doi:10.1038/ng.2746
Howell, C. Y., Bestor, T. H., Ding, F., Latham, K. E., Mertineit, C., Trasler, J. M., & Chaillet, J.
R. (2001). Genomic imprinting disrupted by a maternal effect mutation in the Dnmt1
gene. *Cell, 104*(6), 829-838.
Kamachi, Y., & Kondoh, H. (2013). Sox proteins: regulators of cell fate specification and
differentiation. *Development, 140*(20), 4129-4144. doi:10.1242/dev.091793
Kaneda, M., Okano, M., Hata, K., Sado, T., Tsujimoto, N., Li, E., & Sasaki, H. (2004). Essential
role for de novo DNA methyltransferase Dnmt3a in paternal and maternal imprinting.
Nature, 429(6994), 900-903. doi:10.1038/nature02633
Kato, Y., Kaneda, M., Hata, K., Kumaki, K., Hisano, M., Kohara, Y., . . . Sasaki, H. (2007). Role
of the Dnmt3 family in de novo methylation of imprinted and repetitive sequences during
male germ cell development in the mouse. *Hum Mol Genet, 16*(19), 2272-2280.
doi:10.1093/hmg/ddm179
Legault, L.-M., Chan, D., & McGraw, S. (2019). Rapid Multiplexed Reduced Representation
Bisulfite Sequencing Library Prep (rRRBS). *BIO-PROTOCOL, 9*(4).
Lei, H., Oh, S. P., Okano, M., Juttermann, R., Goss, K. A., Jaenisch, R., & Li, E. (1996). De
novo DNA cytosine methyltransferase activities in mouse embryonic stem cells.
Development, 122(10), 3195-3205.
Leonhardt, H., Page, A. W., Weier, H. U., & Bestor, T. H. (1992). A targeting sequence directs
DNA methyltransferase to sites of DNA replication in mammalian nuclei. *Cell, 71*(5),
865-873. doi:10.1016/0092-8674(92)90561-p
Li, E. (2002). Chromatin modification and epigenetic reprogramming in mammalian
development. *Nat Rev Genet, 3*(9), 662-673. doi:10.1038/nrg887
Li, E., Bestor, T. H., & Jaenisch, R. (1992). Targeted mutation of the DNA methyltransferase
gene results in embryonic lethality. *Cell, 69*(6), 915-926.
Lillycrop, K. A., & Burdge, G. C. (2011). The effect of nutrition during early life on the
epigenetic regulation of transcription and implications for human diseases. *J Nutrigenet
Nutrigenomics, 4*(5), 248-260. doi:10.1159/000334857
Macaulay, E. C., Weeks, R. J., Andrews, S., & Morison, I. M. (2011). Hypomethylation of
functional retrotransposon-derived genes in the human placenta. *Mamm Genome, 22*(11-
12), 722-735. doi:10.1007/s00335-011-9355-1
Magnus, N., Garnier, D., Meehan, B., McGraw, S., Lee, T. H., Caron, M., . . . Rak, J. (2014).
Tissue factor expression provokes escape from tumor dormancy and leads to genomic
alterations. *Proc Natl Acad Sci U S A, 111*(9), 3544-3549. doi:10.1073/pnas.1314118111

McGraw, S., Zhang, J. X., Farag, M., Chan, D., Caron, M., Konermann, C., . . . Trasler, J. M.
(2015). Transient DNMT1 suppression reveals hidden heritable marks in the genome.
Nucleic Acids Res. doi:10.1093/nar/gku1386

Mi, S., Lee, X., Li, X., Veldman, G. M., Finnerty, H., Racie, L., . . . McCoy, J. M. (2000).
Syncytin is a captive retroviral envelope protein involved in human placental
morphogenesis. *Nature, 403*(6771), 785-789. doi:10.1038/35001608

Mizuuchi, M., Cindrova-Davies, T., Olovsson, M., Charnock-Jones, D. S., Burton, G. J., &
Yung, H. W. (2016). Placental endoplasmic reticulum stress negatively regulates
transcription of placental growth factor via ATF4 and ATF6beta: implications for the
pathophysiology of human pregnancy complications. *J Pathol, 238*(4), 550-561.
doi:10.1002/path.4678

Monk, M. (1987). Genomic imprinting. Memories of mother and father. *Nature, 328*(6127), 203-
204. doi:10.1038/328203a0

Morgan, H. D., Santos, F., Green, K., Dean, W., & Reik, W. (2005). Epigenetic reprogramming
in mammals. *Hum Mol Genet, 14 Spec No 1*, R47-58. doi:10.1093/hmg/ddi114

Nakanishi, M. O., Hayakawa, K., Nakabayashi, K., Hata, K., Shiota, K., & Tanaka, S. (2012).
Trophoblast-specific DNA methylation occurs after the segregation of the trophectoderm
and inner cell mass in the mouse periimplantation embryo. *Epigenetics, 7*(2), 173-182.
doi:10.4161/epi.7.2.18962

Nardelli, C., Iaffaldano, L., Ferrigno, M., Labruna, G., Maruotti, G. M., Quaglia, F., . . .
Sacchetti, L. (2014). Characterization and predicted role of the microRNA expression
profile in amnion from obese pregnant women. *Int J Obes (Lond), 38*(3), 466-469.
doi:10.1038/ijo.2013.121

Oda, M., Oxley, D., Dean, W., & Reik, W. (2013). Regulation of lineage specific DNA
hypomethylation in mouse trophectoderm. *PLoS One, 8*(6), e68846.
doi:10.1371/journal.pone.0068846

Okahara, G., Matsubara, S., Oda, T., Sugimoto, J., Jinno, Y., & Kanaya, F. (2004). Expression
analyses of human endogenous retroviruses (HERVs): tissue-specific and developmental
stage-dependent expression of HERVs. *Genomics, 84*(6), 982-990.
doi:10.1016/j.ygeno.2004.09.004

Okano, M., Bell, D. W., Haber, D. A., & Li, E. (1999). DNA methyltransferases Dnmt3a and
Dnmt3b are essential for de novo methylation and mammalian development. *Cell, 99*(3),
247-257.

Piche, J., Gosset, N., Legault, L. M., Pacis, A., Oneglia, A., Caron, M., . . . Andelfinger, G.
(2019). Molecular Signature of CAID Syndrome: Noncanonical Roles of SGO1 in
Regulation of TGF-beta Signaling and Epigenomics. *Cell Mol Gastroenterol Hepatol,
7*(2), 411-431. doi:10.1016/j.jcmgh.2018.10.011

Price, E. M., Cotton, A. M., Penaherrera, M. S., McFadden, D. E., Kobor, M. S., & Robinson,
W. (2012). Different measures of "genome-wide" DNA methylation exhibit unique
properties in placental and somatic tissues. *Epigenetics, 7*(6), 652-663.
doi:10.4161/epi.20221

Reichmann, J., Reddington, J. P., Best, D., Read, D., Ollinger, R., Meehan, R. R., & Adams, I. R.
(2013). The genome-defence gene Tex19.1 suppresses LINE-1 retrotransposons in the
placenta and prevents intra-uterine growth retardation in mice. *Hum Mol Genet, 22*(9),
1791-1806. doi:10.1093/hmg/ddt029

Reiss, D., Zhang, Y., & Mager, D. L. (2007). Widely variable endogenous retroviral methylation
levels in human placenta. *Nucleic Acids Res, 35*(14), 4743-4754.
doi:10.1093/nar/gkm455
Robinson, W. P., & Price, E. M. (2015). The human placental methylome. *Cold Spring Harb
Perspect Med, 5*(5), a023044. doi:10.1101/cshperspect.a023044
Rossant, J., & Cross, J. C. (2001). Placental development: lessons from mouse mutants. *Nat Rev
Genet, 2*(7), 538-548. doi:10.1038/35080570
Santos, F., Hendrich, B., Reik, W., & Dean, W. (2002). Dynamic reprogramming of DNA
methylation in the early mouse embryo. *Dev Biol, 241*(1), 172-182.
doi:10.1006/dbio.2001.0501
Schroeder, D. I., Blair, J. D., Lott, P., Yu, H. O., Hong, D., Crary, F., . . . LaSalle, J. M. (2013).
The human placenta methylome. *Proc Natl Acad Sci U S A, 110*(15), 6037-6042.
doi:10.1073/pnas.1215145110
Senner, C. E., Krueger, F., Oxley, D., Andrews, S., & Hemberger, M. (2012). DNA methylation
profiles define stem cell identity and reveal a tight embryonic-extraembryonic lineage
boundary. *Stem Cells, 30*(12), 2732-2745. doi:10.1002/stem.1249
Shaffer, B., McGraw, S., Xiao, S. C., Chan, D., Trasler, J., & Chaillet, J. R. (2015). The dnmt1
intrinsically disordered domain regulates genomic methylation during development.
Genetics, 199(2), 533-541. doi:10.1534/genetics.114.173609
Slotkin, R. K., & Martienssen, R. (2007). Transposable elements and the epigenetic regulation of
the genome. *Nat Rev Genet, 8*(4), 272-285. doi:10.1038/nrg2072
Smith, Z. D., Chan, M. M., Humm, K. C., Karnik, R., Mekhoubad, S., Regev, A., . . . Meissner,
A. (2014). DNA methylation dynamics of the human preimplantation embryo. *Nature,
511*(7511), 611-615. doi:10.1038/nature13581
Smith, Z. D., Chan, M. M., Mikkelsen, T. S., Gu, H., Gnirke, A., Regev, A., & Meissner, A.
(2012). A unique regulatory phase of DNA methylation in the early mammalian embryo.
Nature, 484(7394), 339-344. doi:10.1038/nature10960
Smith, Z. D., Shi, J., Gu, H., Donaghey, J., Clement, K., Cacchiarelli, D., . . . Meissner, A.
(2017). Epigenetic restriction of extraembryonic lineages mirrors the somatic transition to
cancer. *Nature, 549*(7673), 543-547. doi:10.1038/nature23891
Warren, I. A., Naville, M., Chalopin, D., Levin, P., Berger, C. S., Galiana, D., & Volff, J. N.
(2015). Evolutionary impact of transposable elements on genomic diversity and lineage-
specific innovation in vertebrates. *Chromosome Res, 23*(3), 505-531.
doi:10.1007/s10577-015-9493-5
Watanabe, D., Suetake, I., Tada, T., & Tajima, S. (2002). Stage- and cell-specific expression of
Dnmt3a and Dnmt3b during embryogenesis. *Mech Dev, 118*(1-2), 187-190.
Whidden, L., Martel, J., Rahimi, S., Richard Chaillet, J., Chan, D., & Trasler, J. M. (2016).
Compromised oocyte quality and assisted reproduction contribute to sex-specific effects
on offspring outcomes and epigenetic patterning. *Hum Mol Genet*.
doi:10.1093/hmg/ddw293
Xie, M., Hong, C., Zhang, B., Lowdon, R. F., Xing, X., Li, D., . . . Wang, T. (2013). DNA
hypomethylation within specific transposable element families associates with tissue-
specific enhancer landscape. *Nat Genet, 45*(7), 836-841. doi:10.1038/ng.2649
Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., . . . Chanda, S.
K. (2019). Metascape provides a biologist-oriented resource for the analysis of systems-
level datasets. *Nat Commun, 10*(1), 1523. doi:10.1038/s41467-019-09234-6

714
715
Figure 1. Distinctive patterns of genome-wide DNA methylation accumulation between mid-
gestation embryo and placenta. Analyses of genome-wide DNA methylation sequencing results
for embryo (n=4) and placenta (n=4). **A)** Density histograms showing the distribution of CpG
methylation levels for embryo (*purple*) and placenta (*green*). **B)** Pairwise comparison of CpG
methylation between embryo and placenta. Density increases from blue to red. **C)** CpG fraction of
100 bp tiles within 0-20%, 20-40%, 40-60%, 60-80% and 80-100% ranges in embryo and placenta.
Shown above bars, is the average CpG methylation for each unique tile represented in graph. **D)**
DNA methylation means surrounding the transcription start site (TSS) for All-tiles in each
experimental group. **E)** Circle plot showing methylation average of embryo (*purple*) and placenta
(*green*) across a portion of chromosome 7.

734
735
Figure 2. Genome-wide methylation asymmetry across genomic features between embryo
and placenta. Analyses of genome-wide DNA methylation sequencing results for embryo (n=4)
and placenta (n=4) samples. **A)** Heatmap representation of DNA methylation levels for the 20 000
tiles with the most variable levels between embryo and placenta (DNA methylation variance >20%
and *P*<0.05). **B)** Box-plots representing DNA methylation distribution in embryo and placenta for
the different genomic annotation regions (intergenic, intron, exon, promoter-TSS, 3′UTR, TTS,
5'UTR and non-coding). **C)** Summary of biological functions associated with promoter regions in
Hypo- and Hyper-DMRs. **D)** Examples of smoothed methylation profiles (*BSmooth* tool) in
regions with lower methylation profiles (*Atf6b,* and *Syna*) or higher methylation profiles (*Mir219a-*
2/Mir219b, and *Sox6*) in placenta.
746

747
748
Figure 3. Distinct genomic features in DMRs based on their methylation status in embryo
and placenta. DMR analysis by methylation levels in embryo and placenta. **A)** Sankey diagram
dividing DMRs by methylation levels in embryo (*purple*) and placenta (*green*) for the associated
tiles. High : ≥80% methylation, Mid (intermediate) : ≥20% - <80% methylation , Low : <20%
methylation. **B)** Venn diagram showing the proportion of tiles in the different DMR categories
based on DNA methylation levels between embryo and placenta. **C), D)** and **E)** Analysis of all
tiles, all the DMRs, as well as the 6 different DMR categories based on levels of DNA methylation
in embryo and placenta for : **C)** Genomic annotations, **D)** Main retrotransposons and **E)** Proximity
of CpG rich regions. Neighboring CpG dense regions were defined as shore; up to 2kb away from
CGIs, shelf; 2-4kb away from CGIs, and open sea; >4kb away from CGIs.

766
767
Figure 4. Dynamics of DNA methylation profiles associated with DMR categories and their
evolution across embryo and placenta development. Box plots representing the DNA
methylation distribution and median values for each DMR category in various developmental
stages. Tiles associated with DMR categories at E10.5 were overlapped with previously published
and publicly available data, and methylation levels were determined at each developmental stage
(Smith et *al* 2012, Smith et *al*. 2014, Whidden et *al*. 2016) or in adult somatic tissues (Hon et *al*.
2013). ICM: inner cell mass, TE: trophectoderm, Epi: epiblast, ExE: extraembryonic ectoderm, E:
embryo, Pla: placenta. See Table S2 for median and mean methylation values, and Table S4 for
number of overlapping tiles analysed.

788
789
Figure 5. *Dnmt3a-* or *Dnmt3b*-**deficiency alter proper establishment of DMR-associated**
patterns. A) Violin plot representing DNA methylation distribution and median values of tiles
associated with the different DMR categories at E10.5, and their methylation levels in the different
tissues and genotypes at E6.5 and E8.5. **B)** Promoter-TSS density graphs for E10.5 embryo and
E6.5 epiblast (wt, *Dnmt3a^-/-^* and *Dnmt3b^-/-^*) (left panel) and for E10.5 placenta and E6.5
extraembryonic ectoderm (wt, *Dnmt3a^-/-^* and *Dnmt3b^-/-^*) (right panel). E: embryo, Epi: epiblast,
ExE: extraembryonic ectoderm. See Table S3 for median and mean methylation values, and Table
S4 for number of overlapping tiles analysed.
798
799

**Figure 6. Developmental decline in compensatory DNA methylation mechanism in response
to *Dnmt3a*- or *Dnmt3b*-deficiency at promoter-TSS associated with High-Mid DMRs. A)**
Heatmaps comparing the DNA methylation profiles of wt vs *Dnmt3a⁻/⁻* or wt vs *Dnmt3b⁻/⁻* for E6.5
epiblast, E6.5 extraembryonic ectoderm or E8.5 embryo in 100bp tiles overlapping promoter-
associated DMRs that are highly methylated in the embryo (≥80%) and mildly methylated in the
placenta (≥20 and <80%). Upper panel represents regions with a difference of methylation of at
least 20% between wt and *Dnmt3a⁻/⁻* or wt and *Dnmt3b⁻/⁻*. Lower panel represents stable regions
(lower than 20% difference in methylation). **B)** Pie charts representing DMR numbers for each
comparison. **C)** Heatmap of the methylation levels of tiles commonly represented in all 9 public
data samples that overlap with promoters-associated High-Mid DMRs (n = 314 tiles) (left panel).

811   Heatmap of the methylation level of gene associated with piRNA, gamete generation and germ
812   cells or meiotic nuclear division that are covered in all 9 public datasets (right panel).
813