Title: **Genome wide analysis in *Drosophila* reveals diet by gene interactions and uncovers diet-**

responsive genes

Author list: Deanne Francis[1,2], Shila Ghazanfar[1,4,6], Essi Havula[1,2], James R. Krycer[1,2], Alistair Senior[1,2],

Annabel Y. Minard[1,2,5], Thomas Geddes[1,2], Fiona Weiss[1,2], Jacqueline Stöckli[1,2], Jean Y.H. Yang[1,4] and David

E. James[1,2,3*]

1 Charles Perkins Centre, The University of Sydney, Camperdown, 2006, Australia

2 School of Life and Environmental Sciences, The University of Sydney, Camperdown, 2006, Australia

3 Sydney Medical School, The University of Sydney, Camperdown, 2006, Australia

4 School of Mathematics and Statistics, University of Sydney, Sydney, NSW 2006, Australia.

5 Current Address: Department of Molecular Physiology and Biophysics, Carver College of Medicine, The University of Iowa,

Iowa City, IA, 52242

6 Current Address: Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre,

Robinson Way, Cambridge, CB2 0RE, United Kingdom

* Corresponding author: David E. James, The University of Sydney, Camperdown, 2006, New South Wales, Australia. Email:

david.james@sydney.edu.au

Author contributions: D.F. and D.E.J. conceived the study. D.F., J.R.K and D.E.J participated in experimental

design. D.F., F.W., and A.M. performed the experiments and D.F., A.S., E.H. , T.G. and S.G. analysed data

for the experiments. J.S. and J.Y. provided intellectual input. D.F., J.S. and D.E.J. wrote the manuscript, and

all the authors edited the manuscript. D.E.J. supervised the study.

Acknowledgements

The authors would like to thank Elise Needham for edits, Roel Bevers and Bart Deplancke for useful

suggestions and members of the James Lab for helpful discussions. The authors acknowledge the facilities,

and the scientific and technical assistance, of the Australian Microscopy & Microanalysis Research Facility

at the Charles Perkins Centre, The University of Sydney.

Financial disclosure

The work is supported by National Health and Medical Research (NHMRC) project grants (APP1086851 for

D.E.J.) and Diabetes Australia Research Program grants (J.R.K. & D.F.). J.R.K. is recipient of the Australian

Diabetes Society Skip Martin Fellowship. D.E.J. is an NHMRC Senior Principal Research Fellow

(APP1117078). The contents of the published material are solely the responsibility of the authors and do not

reflect the views of the NHMRC. The authors have no financial conflicts of interest to declare.

Abstract: Genetic and environmental factors play a major role in metabolic health. However, they do not

act in isolation, as a change in an environmental factor such as diet may exert different effects based on an

individual's genotype. Here, we sought to understand how such gene-diet interactions influenced nutrient storage and utilisation, a major determinant of metabolic disease. We subjected the Drosophila Genetic Reference Panel (DGRP), comprising 200 genetically divergent inbred fly strains, to diets varying in sugar, fat and protein. We assessed starvation resistance, a holistic phenotype of nutrient storage and utilisation that can be robustly measured. Diet influenced the starvation resistance of each strain, but this effect varied markedly between strains. This demonstrates that genetics plays a major role in the response to diet. Furthermore, heritability analysis revealed that the greatest variability arose from diets either high in sugar or high in protein. To uncover the genetic underpinnings of this variation, we mapped 1,239 diet-responsive SNPs in 534 genes, 325 of which have human orthologues. Using whole-body knockdown, we confirmed that 30 candidate genes were required for glucose tolerance, storage and utilization. In particular, we characterised CG4607, a GLUT6/GLUT8 homolog, as a key protein involved in sugar tolerance. Overall, this provides strong evidence that genetics is a major contributor to how individuals respond to diets of varying nutrient composition. It is likely that a similar principle may be applied to metabolic disease in higher organisms thus supporting the case for nutrigenomics as an important health strategy.

Introduction:

Although it is widely appreciated that gene and environment interactions play a major role in determining various phenotypes in different animals, the application of this paradigm to health outcomes in humans remains an unmet need. For instance, a well-accepted culprit of metabolic disease is the widespread introduction of the high-fat, high-sugar 'Western' diet. Interestingly, indigenous populations only experienced a considerable burden of metabolic disease upon exposure to Western diets, after subsisting on different diets, whether rich in fats (Greenlandic Inuits) or carbohydrates (American Pima Indians), for thousands of years (Schulz & Chaudhari, 2015; Andersen & Hansen, 2018). However, studies in mice and humans have demonstrated that individuals display heterogeneous metabolic responses to the same diets (Zeevi *et al*, 2015; Parks *et al*, 2015). While genome-wide association studies (GWAS) of body mass index, insulin resistance and other metabolic traits have identified several causative loci, these loci do not account for the majority of phenotypic variation (Phillips, 2013). In fact, studies in mono- and di-zygotic twins revealed that a combination of genetic and environmental factors contributed to variance in body weight (Stunkard *et al.*;

Dubois *et al.* 2012), with obesity being linked to metabolic disease risk. Overall, this emphasises the need to

consider not only genetic predisposition or diet alone, but specifically gene-diet interactions (Heianza *et al.*

2017). Furthermore, one's genotype can affect metabolic outcomes in response to a particular diet, and this

'diet-responsiveness' can vary markedly between individuals.

The notion that individuals display discrete sensitivity to certain macronutrients, with dietary

interventions being based on one's genotype, is the key premise to the field of personalised nutrition, or

nutrigenomics (Sales *et al.* 2014) and is supported by recent preliminary clinical trials (Qi, 2014; Tan *et al*,

2019). However, we do not have a thorough understanding of how genes and diet interact to influence the risk

of metabolic disease (Drabsch & Holzapfel, 2019). Here, we aimed to set up a model system for recapitulating

gene-diet interactions. Studying this in humans remains a challenge because the environmental variables are

difficult to control at a sufficient scale to facilitate genetic mapping. In contrast, the *Drosophila* fruit fly model

system overcomes many of these logistical issues. Importantly, >70% of known human disease genes have

fly orthologs (Reiter *et al.* 2001), and genetic tools such as the *Drosophila* genetics reference panel (DGRP)

with 200 inbred and fully sequenced lines are available, thereby allowing identification of causal genetic

variants (Mackay *et al.* 2012).

We capitalised on the high-throughput nature of the DGRP to study the effect of gene-diet interaction

on nutrient storage and utilisation. A phenotype that encapsulates these outcomes is survival during starvation,

which measures an organism's ability to metabolise and store dietary nutrients during feeding and utilise them

efficiently during fasting. For instance, adult flies fed a high carbohydrate diet showed greater resistance to

starvation, as well as higher triacylglycerides (TAGs), suggesting a correlation between diets and starvation

resistance via energy storage (Lee and Jang 2014) – however, does this apply to all individuals? Indeed, there

was significant variation in starvation resistance between strains across the DGRP when fed a single diet

(Mackay *et al.* 2012). Thus, we utilised the genetic diversity of the DGRP to identify SNPs that alter starvation

resistance after exposure to diets varying in fat, sugar and protein content. Our study provided strong evidence

establishing starvation resistance in *Drosophila* as a prime example of a phenotype governed by gene-diet

interactions, motivating the notion that a similar principle may be applied to metabolic disease in higher

organisms. Here, we combine the high-throughput nature of the *Drosophila* model with the genetic diversity

of the DGRP to dissect diet-gene interactions on a population level. In this study, we aimed to identify "diet-

responsive" genes and determine the mechanism by which genes in combination with diet control metabolic

phenotypes. To do this, we used the DGRP to perform a GWAS to identify SNPs that contribute to variation

in response to diets that differ in fat, sugar and protein contents. Our study uncovers a previously under-

appreciated influence of diet on the heritability of starvation resistance. Finally, we provide a rich resource of

diet specific genes for further study.

Results

Starvation Resistance of DGRP across 4 different diets

We first sought to identify novel diet-responsive genes that affect nutrient storage and utilisation in

Drosophila. We used survival during starvation as a surrogate for an obesogenic phenotype to screen for

dietary effects. This is a powerful and sensitive assay as starvation resistant flies are often replete with fat

stores immediately prior to starvation and feeding flies a high sugar diet increases fat stores and prolongs

starvation resistance (Djawdan *et al.* 1998; Hoffmann and Harshman 1999; Bjedov *et al.* 2010). The diets we

selected, (normal food (NF), high carbohydrate diet (HCD), high fat diet (HFD) and high protein diet

(HPD),Table S1), were based on previous studies that explored the effect of different sugar and protein

concentrations on starvation resistance in a single strain (Skorupa *et al.* 2008; Lee and Jang 2014; Chandegra

et al. 2017). The dietary composition is indicated in Table S1: the carbohydrate is sugar, the protein is yeast

and the fat is coconut oil. We exposed 3 to 5-day old adult males from 178 DGRP strains to the 4 diets for 10

days and then measured starvation resistance by removing food and assessing survival (Fig. 1A, Fig. S7). The

DGRP has previously been used to examine starvation resistance in flies fed NF (Mackay *et al.* 2012) and

there was a strong correlation in starvation resistance across the 178 strains between the two studies (males,

NF, Pearson's R= 0.58, Fig. 1B), demonstrating the robustness of the starvation phenotype and the DGRP

resource. Interestingly, we found that previously published food intake data (Garlapow *et al.* 2015) was

negatively correlated with starvation resistance across genotypes (males, NF, Pearson's R= -0.32, Fig. 1B) in

both our study and previously published starvation data (Mackay *et al.* 2012). This is intriguing as it suggests

that strains that ate the most were the least resistant to starvation. This could be due to differences in metabolic

rate, an increase in hunger cues, nutrient storage capacity or differences in hormonal responses.

Irrespective of strain, flies fed NF, HFD and HCD displayed the greatest variance in starvation

resistance, while HPD fed flies had the least amount of variation across the strains (Fig 1C, 1D). We

normalised the starvation resistance to NF for each strain and performed hierarchical clustering, indicating

clusters of strains that were starvation sensitive to either HCD, HPD or HFD (Fig S1). Surprisingly these data

showed a disparity between the responses to different diets, likely driven by genetic background, indicating

that there is no single optimal diet where all strains display similar starvation resistance responses (Fig 1D,

Fig S1).

We quantified the differential contributions of diet and gene by diet interactions to the variation in starvation

resistance among DGRP strains. We determined the broad sense heritability (H_2) of the starvation response

using linear-mixed models (LMMs). We found that across the whole data set, after accounting for differences

among diets, the H_2 of starvation resistance was 20%. Dietary effects accounted for 8% of the variance in

starvation-resistance, leaving 72% residual (unexplained) variance. A model including a term to estimate

genetic variance in dietary effects (random-slopes LMM) had a significantly better fit than a model without

(likelihood ratio test, L = 11611, d.f. = 14, p<0.001) indicating the presence of gene by environment

interactions. Within-diet, the H_2 of starvation resistance on NF, HCD, HFD and HPD was 19%, 50%, 20%

and 65%, respectively (Table 1 and S2). Thus, across populations certain diets expose genetic diversity more

strongly than others and we estimate that genetics plays a particularly strong role in determining starvation

resistance in flies fed either HPD or HCD.

Mapping 'diet-responsive genes'

We aimed to identify gene-diet interactions by uncovering SNPs associated with diet-responsive

starvation resistance (Fig.2A). We first identified diet-associated SNPs across all diets by performing

multivariate ANOVA testing with a significance threshold of unadjusted P-value < $1x10_{-4}$. This is higher than

the conventional threshold used (P value $1x 10_{-5}$, (Mackay *et al.* 2012)) to increase the number of true positive

SNPs as further filtering was applied. Such filtering reduced the number of SNPs to exclusively contain SNPs

that exhibited a significant and large difference in starvation survival compared to NF in at least one diet

(unadjusted p-value < 0.01, absolute log fold change > 0.3), by performing a univariate rank-based Wilcoxon

Rank Sum Test. Using these stringent filters, we identified 1,239 SNPs that were associated with diet-

responsive starvation resistance. These SNPs were located within 534 genes (Table S7), 325 of which had

human orthologs with a high proportion of the SNPs (>80%) found in non-coding regions (Table S3 and S4).

We observed that the largest number of SNPs were associated with HFD (Fig. 2B) and only 7 SNPs affected

starvation resistance across all three diets (Fig 2. B). In order to assess the SNP data, we looked at the genomic

distribution of all SNPs, including those with predicted diet interactions (Fig. 2C, black dots). To observe the

genomic distribution of SNPs on each diet we generated Manhattan plots and observed a higher density of

diet-responsive SNPs around chromosome 2L and 3R in flies fed HFD, while HCD and HPD had fewer

associated SNPs (Fig 2, D-F black dots). Next, Quantile-Quantile (Q-Q) plots showed an enrichment of

significant SNPs compared to the expected p-value distribution, taking into account the genomic population

structure of the DGRP, and the correlation structure of diet survival phenotypes (Fig. 2G). We observed no

significant associations between the phenotype and covariates (Wolbachia status and large genomic

inversions) (data not shown). Thus, we conclude that neither population structure nor additional covariates

such as Wolbachia status affected the outcome of our SNP analysis. Overall using conventional SNP analysis

resulted in an enrichment of diet-responsive SNPs.

We next undertook a pathway analysis of genes containing significant diet-responsive SNPs using

gene ontology (GO) analysis on the human orthologs of the 'diet-responsive genes' (Table 2). This revealed

stark differences in enriched biochemical processes between the different diets. For instance, genes associated

with enhanced starvation resistance ('up') on HFD were enriched for peptidases in proteolytic pathways. In

contrast, genes associated with reduced starvation resistance ('down') after HFD and HPD were enriched for

signal transduction and for HFD, the MAPK pathway in particular (Table 2). Meanwhile, extracellular matrix

pathways and pathways linked to sulfotransferases were respectively linked to enhanced and reduced

starvation resistance after HCD feeding (Table 2). This suggests that individual diet responsiveness is

controlled at the pathway level rather than at the level of individual genes *per se*. These data provide an

abundant and valuable resource of diet-responsive genes and pathways.

Candidate gene validation

The following three main criteria were used to select genes for further validation: whether the SNPs

passed the more stringent fold change cut-off with respect to diet response (>0.3 log2 fold change); the

presence of a human ortholog; and an annotated gene function (Table 3). Using these criteria, we selected 30

genes for further validation using the GAL4-UAS system (Brand and Perrimon 1993). We generated whole-

body knockdown flies for each gene in the *W1118* background strain, pre-fed them the four diets and used a

more precise (low-throughput) method of monitoring starvation with the *Drosophila* activity monitoring

system (DAMs)(Pfeiffenberger *et al.* 2010). Candidate genes were considered validated when knocking down

the gene yielded a starvation resistance phenotype (Fig. S2). Strikingly, of the 30 genes tested, whole body

depletion of 8 candidate genes led to lethality, while 16 of the remaining 22 candidates displayed a significant

diet interaction – 62.5% of these were significant according to a cox hazard multivariate analysis (p<0.05,

Table 3, Fig. S2). This validated the use of RNAi knockdown as a useful tool and is consistent with the notion

that for many but not all of these genes the corresponding SNP is associated with reduced gene expression or

loss of function.

Of these genes, two were previously associated with starvation resistance: *spaghetti-squash activator*

(*sqa*) a myosin light chain kinase that is activated by ATG1 to form autophagosomes during starvation (Tang

et al. 2011) and *methuselah-like 3* (*mthl3*) known to enhance starvation resistance in the red flour beetle, T.

casteneum (Li *et al.* 2014). We also identified a lipid transport protein, *Lipophorin receptor 2 (LpR2)*, required

for the transport and binding of lipid molecules from the plasma membrane (Parra-Peralbo and Culi 2011;

Rodríguez-Vázquez *et al.* 2015); *Cdc42-interacting protein 4* (*Cip4*), which is implicated in phospholipid

binding in *Drosophila* and GLUT4 trafficking in mice (Feng *et al.* 2010; Zobel *et al.* 2015). Furthermore, we

identified *CCHamide-2 receptor* (*CCHa-2R*), a G-protein coupled receptor and *Drosophila* ortholog of the

bombesin receptor 3 (BRS3), which regulates secretion of *Drosophila* insulin-like peptides in response to

nutrient availability (Sano *et al.* 2015). Knockdown of all three genes (*LpR2, Cip4, CCHa-2R*) were associated

with increased starvation sensitivity after HFD feeding. Another interesting candidate, *happyhour* (*hppy*),

Drosophila ortholog of MAP4K3, a serine/threonine kinase that regulates triglyceride homoeostasis and

mTOR signalling (Bryk *et al*, 2010), modulated the response to HCD. Loss of *CG3339*, the *Drosophila*

ortholog of dynein heavy chain (DNAH9), a protein that regulates minus end directed microtubule trafficking,

enhanced starvation resistance on HCD. The most striking starvation sensitivity in response to diet was

observed for *CG9674* (glutamate synthase) and *CG4607* (GLUT6/GLUT8 orthologue). Knockdown of these

genes in flies modulated the response to HCD and HPD, respectively (Table 3).

Analysis of CG4607, an example of a diet responsive gene

CG4607 was of particular interest because it was starvation resistant on NF, sensitive to HPD, the diet

with the lowest amount of sugar, and is homologous to the human glucose transporters SLC2A6 (GLUT6)

and SLC2A8(GLUT8). Very little is known about the function of these proteins in mammals and so this raised

the possibility that they might play a unique role in the response to sugar. GLUT6 (formerly Glut9) is over

expressed in endometrial cancer and is highly expressed in brain, spleen and leukocytes in mice (DOEGE *et*

al. 2000). GLUT8 is highly expressed in the testis, heart, brain, liver, fat, and kidneys in mouse and has been

shown to respond to insulin and transports trehalose in hepatocytes (Carayannopoulos *et al.* 2000; Mueckler

and Thorens 2013; Mayer *et al.* 2016). In flies, *CG4607* is highly expressed in the adult midgut but is modestly

expressed in other tissues such as the heart, fat body, salivary gland and fly heads(Robinson *et al.* 2013).

However, depletion of *CG4607* with both midgut and fat body drivers did not phenocopy the whole body

starvation phenotype (data not shown). Whole body expression of the second hairpin *CG4607 GD3268* was

lethal, however, expression of both hairpins using *mef2*-GAL4 was also lethal (data not shown) and suggests

that the two hairpins overlap. Remarkably, groups of flies expressing UAS-*CG4607* KK104152 RNAi in the

whole body (*CG4607* KD) died after only three days of eating HCD (Fig. 3A), while surviving on NF. The

DGRP strain 45, which contained the *CG4607* SNP, was also not viable when fed HCD (Table S6), suggesting

that the lethality of strain 45 may be due to the presence of the SNP in *CG4607*. To further assess this

phenotype, we sought to monitor the behaviour of in single housed control and *CG4607* KD flies while they

were eating NF or HCD. We found a striking hyperactivity on HCD that abruptly stopped after 12 h when

flies died, as indicated by a total lack of activity (Fig. 3 B, C). The hyperactivity was reminiscent of flies

during starvation (Keene *et al.* 2010) and we speculated that the lethality was due to an inability of the flies

to consume HCD, thereby experiencing starvation. However, upon measuring caloric intake in control and

CG4607 KD flies we observed that knockdown flies consumed 38% more calories of HCD than controls (Fig.

3D). Thus, the observed hyperactivity of *CG4607*KD flies on HCD may be due to a perception of hunger

brought about by the loss of glucose sensitivity, or a defect in the ability to utilise stored nutrients that

ultimately lead to lethality.

Next, we investigated the mechanism by which *CG4607* controlled nutrient storage or utilisation in

response to HPD. NF pre-fed *CG4607* KD flies were starvation resistant but became starvation sensitive after

becoming low carbohydrate stressed on HPD (Table 3). We posited that the starvation phenotype reflected the

levels of energy stores. Fed nutrient levels were similar between *CG4607* KD and control flies on both NF and

HPD (Figure 4 A and C), with the exception of TAGs, which were higher in *CG4607* KD animals (Fig. 4 E).

However major differences in nutrient storage levels were observed upon starvation on both diets. Fasting of

HPD-fed flies resulted in a marked depletion of nutrients in both genotypes with undetectable fasted nutrient

levels in control flies. Yet, fasted levels of glycogen, glucose and TAGs were still higher in *CG4607* KD flies

than in control flies (Fig. 4 B, D, G). Overall, our data shows that relative to control flies, *CG4607* KD flies

have increased caloric intake and excessive energy stores on NF, which are reduced upon HPD feeding. This

suggests that although *CG4607* KD flies have an excess of energy stores, these are still mobilised during

starvation. However, given that HPD *CG4607* KD flies still die faster than HPD control flies during the

starvation resistance assay, this suggests an inefficient mobilisation of energy stores.

To further probe the role that *CG4607* plays in mobilising energy stores during starvation, we wanted

to explore if CG4607 functions like a glucose transporter. To address this, we measured glucose utilisation by

feeding flies an NF diet containing 14C-radiolabelled glucose and monitoring glucose incorporation into CO_2

and lipids (Francis *et al.* 2019; Krycer *et al.* 2019). We observed that *CG4607* KD flies exhibited lower levels

of glucose oxidation and incorporation into lipids compared to controls (Fig. 5 A, B), indicating reduced

glucose utilisation. This was not due to lower food intake in the *CG4607* KD flies (Fig. 5C). Given the effect

of depleting *CG4607* on glucose utilisation, we wanted to see where it was localised and posited that its

localisation would be similar to its mammalian homologs. The mammalian GLUT6 and GLUT8 both localise

to lysosomes (Diril *et al.* 2009; Maedera *et al.* 2019). To determine if the localisation of CG4607 was

conserved, we created and expressed CG4607-mRUBY3 in HeLa cells. We used immunofluorescence

microscopy to observe that, like GLUT6 and GLUT8, CG4607 partially co-localised (Avg. Pearson's r = 0.57,

n=13 cells) with lysosomal markers (Fig.5 D, E). Taken together, our data show that CG4607, a GLUT6/8

orthologue, is an example of a diet-responsive gene and that responds to dietary sugar by regulating lysosomal

glucose metabolism.

Discussion

255
256
The DGRP is a powerful tool for understanding the genetics driving variation of metabolic phenotypes

(Mackay *et al.* 2012; Garlapow *et al.* 2015; Unckless *et al.* 2015; Nelson *et al.* 2016). Our approach measured

starvation resistance in the adult male population after exposure to diets that vary in sugar, protein and fat

content. Using this method, the DGRP supported the notion that nutrition responses vary considerably between

individuals of different genetic background. Although, diet had a major influence on starvation resistance, the

diets that enhanced or reduced starvation resistance differed markedly between *Drosophila* strains. In

particular, exposure to diets high in sugar (HCD) or protein (HPD) elicited a greater genetic contribution to

phenotypic variation. These data highlight a heterogeneity in the response to diet that underlies the

fundamental principles of personalised nutrition. The SNPs we identified provide a resource dataset for further

study. Out of 39 candidate diet responsive genes that were selected for more detailed analysis, we validated

21 genes as bona fide diet responsive genes. Knock down of one such gene, *CG4607*, was linked to increased

caloric intake and nutrient storage combined with reduced glucose utilisation.

In relation to our study, since the natural food sources of *Drosophila* are rotting fruits that contain

sugar and yeast (Markow 2015), we postulate that genetic variants involved in the processing, storage and

utilization of sugar or protein (from yeast) contribute to differences in starvation resistance. Indeed, changes

in sugar and lipid handling enzymes have previously been linked to starvation resistance (Harshman *et al.*

1999). Alternatively, sugar and protein (from yeast) are dominant drivers of food intake with major sensory

systems to control intake of these macronutrients and hence these are more likely to be under genetic control

(Chng *et al.* 2017; May *et al.* 2019). Overall, this demonstrates that the mechanisms that underpin diet

sensitivity are complex and involve differences in food preference and thus food intake as well as differences

in nutrient metabolism (energy storage and utilisation), all of which are regulated at least in part by one's

genotype.

This study provides a rich resource of diet responsive genes and pathways. The majority (>80%) of

the SNPs in candidate genes were non-coding, a finding that is consistent with previous DGRP (Mackay *et al.*

2012) and human GWAS (Gallagher and Chen-Plotkin 2018) and the functional consequences of such SNPs

remain to be identified. As a proof of principle, we were able to validate candidate genes with intronic SNPs

using RNAi knockdown indicating that these types of mutations may control expression of these genes.

However, further studies examining transcriptional control are required to validate this conclusion.

We focused on *CG4607*, a validated diet-responsive gene. The SNP in *CG4607* was non-coding and

within the 5'-UTR, suggesting that it could affect splicing or post-transcriptional processing. Additionally,

the SNP contains a regulatory transcription factor binding site for the *invective* transcriptional repressor. While

it may be of interest to determine if the SNP alters the expression of CG4607, our validation using RNAi

knock down clearly shows that CG4607 is a diet-responsive gene. SNPs in *CG4607* were identified in a

GWAS of sleep/activity (Harbison *et al.* 2013). Our study provides several lines of evidence that CG4607

mediates its effects via regulating glucose metabolism. First, whole body depletion of *CG4607* resulted in

lethality after 3 days on a high sugar diet (HCD) (Fig. 3A) and the lethality was accompanied by hyperactivity

and an increase in caloric intake on HCD compared to control flies. This is symptomatic of a starvation

phenotype (Yu *et al*, 2016; Yang *et al*, 2015). Second, *CG4607 KD* animals have higher energy stores on NF

compared to control flies, likely due to impaired glucose utilisation. Third, our metabolic labelling

experiments show that *CG4607 KD* animals exhibit reduced glucose utilisation.

Last, the closest mammalian orthologues of *CG4607* are the human glucose transporters GLUT6 and

GLUT8. Recently, GLUT8 has been shown to transport trehalose in mammalian cells (Mayer *et al.* 2016)

suggesting that CG4607 may play a similar role in flies. However, loss of *CG4607* does not phenocopy

elevated or reduced trehalose levels and may not play a major role in trehalose transport(Matsuda *et al.* 2015;

Yasugi *et al.* 2017). Furthermore, GLUT8 regulates AMPK phosphorylation and signalling via trehalose

transport (Mayer *et al.* 2016; Narita *et al.* 2019) and AMPK deficient flies are starvation sensitive (Johnson *et*

al. 2010). It would be of interest to determine if CG4607 regulates AMPK in a similar manner as its

mammalian ortholog. Given that AMPK is an energy sensor perhaps energy sensing pathways play a part in

diet-dependent differences in starvation resistance.

Like these GLUT transporters, CG4607 is also targeted to lysosomes instead of the plasma membrane,

unlike other facilitative sugar transporters (Lisinski *et al.*; Maedera *et al.* 2019). This is intriguing as this co-

localises with mTORC1 (Lee *et al.* 2009; Efeyan and Sabatini 2013), which regulates numerous metabolic

processes including glycogen breakdown in autophagic vesicles (Mony *et al.* 2016; Zhao *et al.* 2018). Given

that compared to control flies, *CG4607 KD* flies exhibited higher glycogen levels after starvation, and broke

down more TAGs during starvation, the metabolic phenotype of *CG4607 KD* flies resembles a lysosomal

glycogen storage disease, where lysosomal glycogen cannot be accessed for energy utilisation in the cytosol.

Together, this provides strong evidence that CG4607 mediates its effects on starvation resistance by regulating

glucose metabolism. Hence this highlights the importance of glucose metabolism as a potential diet responsive

pathway.

Overall, our findings demonstrate that gene-diet interactions are an impact factor to consider in

metabolic homeostasis. This has substantial ramifications for human health because it means that the concept

of a 'healthy' diet varies between individuals, thus questioning population-wide nutritional recommendations.

While our study provides the basis for a nutrigenomics initiative such an endeavour is likely to require a

substantial future investment at the clinical level.

Materials and Methods

Materials and Methods

Drosophila Stocks and procedures

Stocks: DGRP (Bloomington *Drosophila* Stock Center, Indiana, USA), RNAi reagents (CG4607 v107219 and

v5450, VDRC, Vienna, Austria), *ubiquitous*-GAL4 (Bloomington # 32551) and *CG*-Gal4 (Bloomington #

7011). Flies were maintained at standard temperature and 12h light/dark cycle. DGRP flies were expanded in

bottles before collecting adult males for experiments. 5 replicates of ten 3-5 day old adult males from each

strain were collected and passaged onto each diet. Food was changed every other day, and the mortality rate

was monitored for the ten days of diet treatment. The diets were well tolerated with a similar lethality on HFD

during pre-feeding compared to other diets and did not affect the starvation assay. Afterwards, males were

placed into starvation vials with kimwipes and 1 mL of water and monitored every 12 hours for death. We

initially tested 5 diets (NF, HCD, HFD, HPD and CR). However, we discovered that the CR diet (6.5% sugar

and yeast) was not very different compared to NF. Thus, we chose to focus only on NF, HCD, HPD and HFD.

The median and mean starvation resistance was analysed using Prism and the R (CRAN, survival and

survminer packages).

Heritability estimates using LMMs

336 We estimated heritability from two LMMs of log survival. Model 1 included diet as a fixed effect and a

337 random intercept for genotype. Model 2 also included a random slope for diet effects among genotype.

338 Heritabilities were calculated as $\sigma_{2Genotype} / (\sigma_{2Genotype} + \sigma_{2Residual}) \times 100$. For model 2, diet specific $\sigma_{2Genotype}$

339 was calculated as $\sigma_{2Intercept} + \sigma_{2Diet} + 2 \times \rho_{Intercept} \times \sigma_{Intercept} \times \sigma_{Diet}$. A likelihood ratio test suggested model 2

340 gives significantly better fit to the data than model 1.

Statistical SNP analysis

In order to determine the diet-responsive SNPs we performed bioinformatic analysis on mean starvation

resistance data of each DGRP strain. To use the effect size of the diets, the mean starvation resistance of

DGRP strains on HCD, HFD and HPD was divided by the mean starvation resistance on NF. This normalised

starvation resistance was fed into the analysis pipeline. DGRP SNP genotypes were downloaded from the

DGRP Freeze 2 online resource http://dgrp2.gnets.ncsu.edu/data.html for all lines considered in this

manuscript. SNPs were filtered so at least five lines contained one of each of the reference and alternate alleles,

resulting in testing across 2,455,135 SNPs. Lines with missing allele information for a given variant were not

considered within each test. Statistical testing included multivariate analysis of variance (MANOVA) testing,

with Wolbachia status as a covariate per variant, with an unadjusted P-value < 1e-4 as significant, as well as

Wilcoxon Rank Sum Tests per diet. The overall enrichment of the multivariate response was assessed by

comparison to the null p-value distribution under random permutation of line labels, thereby retaining the

overall phenotype and SNP-SNP correlation structure (Fig 2G). To assess the effect size, we calculated the

median difference in phenotype between 'reference' and 'alternate' allele groups per diet, phenotype being

the log2 ratio of survival to NF per line. We selected diet-SNP pairs for further consideration if they were

significant for both the MANOVA test and Wilcoxon Rank Sum Test, and had an absolute difference in

median phenotypes of at least 0.3 (log2 fold change).

Gene ontology analysis:

Gene ontology and pathway analyses were performed using Fisher's Exact Test in the 'goseq' R package. We

used human orthologs for the analysis since GO analysis of fly genes led to very broad non-specific categories

(results not shown). Furthermore, using only the human orthologs would limit us to conserved genes. A total

of 6,056 pathways, based on human genes, with at least 10 and at most 500 genes were kept, with a 'gene

universe' of 7,677 human genes that are homologues of the fly genes tested. Pathways were considered

significant if FDR-corrected values were below 0.05, per diet and direction of phenotype change.

Validation/Automated starvation resistance (DAMS assay):

We used the GAL4-UAS gene expression system to validate the 30-candidate diet-responsive genes. RNAi

knockdown fly lines (see table S5 for reagent ID) were mated to 20 *ubi*-Gal4 females. Sixteen 3-5 day old

males were placed on five different diets for ten days. The food was changed every other day for ten days until

males were placed into the DAMS apparatus (Trikinetics, inc., USA). The flies were loaded into DAMS tubes

containing 2% agar and monitored every 5 minutes for starvation resistance. Candidate genes were considered

validated when the fold change in survival on each diet (relative to NF) corroborated with the SNP analysis

(Table 3, Table S5). The gene by diet p-values were generated using the Cox proportional hazards regression

model (survival package, R).

Measuring Activity

To measure fly activity while eating NF or HCD we placed 3-5 day old male flies into DAMs tubes with either

NF or HCD and monitored for activity for ten days. Since *CG4607 KD* flies died within the first 24 hours after

being placed on HCD, we analysed the activity data from the beginning of the monitoring. DAMS data were

analysed using the R survminer and rethomics (Geissmann *et al.* 2019) package.

Blue Dye Extraction and measurements:

To determine the amount of food ingested during the gas-trap assay, we extracted and measured blue dye that

was mixed with the radiolabelled food. In short term experiments, the blue dye serves as a proxy for food

intake (Krycer *et al.* 2019). 4 replicates of ten flies were collected and homogenized (Reche MM400) in 100uL

of water. Samples were briefly spun down and the supernatant was dried down in the Genevac personal

evaporator. Dried samples were reconstituted in 50uL of water, vortexed and placed into a 96 well plate for

measurement at 628nm in a spectrophotometer. Dilutions of blue dye (Queenie Brand, Coles Supermarket,

Australia) were used as a reference. Data was analysed in Excel (Microsoft) and plotted in Prism (Graphpad).

Statistical significance was calculated between genotypes using Student's t-test.

Capillary feeding assay:

We used the Café assay as previously described (Lee *et al.* 2008) to determine the amount of the

experimental diet eaten. 10 replicates of five 3-5 old adult males were placed into vials with water soaked

kimwipes and sealed with a rubber stopper with two holes. 5uL capillary tubes with food was placed

through one hole and into the vials to allow flies to feed for 24 hours(Lee *et al.* 2008). The diets were

composed of yeast (MP Biomedicals cat # 2232731), and sucrose (Table Sugar, Coles Supermarket,

Australia).

Experimental Diets:

The experimental diets that were used throughout this study were made up of agar (Sigma) and torula yeast

(H.J Langdon & Co, Victoria, Australia), Sucrose (Table sugar, Coles, Victoria, Australia) and extra virgin

coconut oil (Absolute Organics, NSW, Australia).

Gas Trap Assay to measure CO2 and Triglycerides:

We used a Gas Trap Assay to assess the capability of *CG4607 KD* flies to utilise glucose. The gas trap protocol

has been previously described(Francis *et al.* 2019). Briefly, 4 replicates of ten male adult 3-5-day old flies

were starved overnight with a Kim wipe and 1mL of water. Flies were placed into 12 well plates containing

glucose radiolabelled food and blue dye. We measured glucose oxidation and processed the flies for TAGs

and blue dye content as described.

Triglycerides Assay:

To measure triglyceride content we used a triglyceride extraction method performed as previously described

(FOLCH *et al.* 1957). 6 to 10 replicates of six 3-5 day old flies were collected and washed in 4 dilutions of

isopropanol to remove excess food. The lipids were collected after extraction, evaporated under N_2 gas and

reconstituted with 95% ethanol. Scintillant was added to samples with radioactive tracer instead of ethanol

and read on a beta counter (Beckman Coulter). For non-radioactive samples: samples were spun and placed

into 96 well plates (Sigma-Aldrich, # CLS9018BC) and incubated with triglyceride reagent (200uL; Thermo

Fischer Cat #TR22421) for at 37c for 30min. Precimat glycerol reagent (Thermo Fischer # NC0091901) was

used as a reference. Total absorbance at 500 nm was measured in a plate reader (Beckman) and subtracted

from a blank before determining the amount of triglyceride using the reference standard curve. All calculations

were performed in Excel (Microsoft) and graphed in Prism.

Glycogen Assay:

To determine the amount of glycogen we collected 6 replicates of six male flies and washed them in several

dilutions of isopropanol to remove food. Fasted flies were collected after 24 hours of starvation. Flies were

homogenised in 1M KOH for 30 seconds using steel balls and a tissue lyser (Resche MM400). Samples were

heated for 30 min at 70 degrees C. Saturated Na_2SO_4 was added following 95% Ethanol for precipitation. The

pellet was spun down and then reconstituted in water, heated at 70 degrees and 95% Ethanol was added again.

The pellet was spun down, and amyloglucosidase (Merck # A7420) were added overnight at 37 degrees.

Samples were spun and placed into 96 well plates (Sigma-Aldrich, # CLS9018BC) and incubated with glucose

oxidase reagent (200uL; Thermo Fischer, TR15221) for at 37C for 30min. 1mg/ml glucose was used as a

reference. Total absorbance at 500nm was measured in a plate reader (Beckman) and subtracted from a blank

before determining the amount of glucose using the reference standard curve. All calculations were performed

in Excel (Microsoft) and graphed in Prism.

Glucose assay:

Glucose was measured from the aqueous phase of the triglyceride extraction (see above): the aqueous mixture

was evaporated in a Genevac E2-3 evaporator until a dried pellet was visible. The pellet was reconstituted

with water and glucose was measured as described for glycogen.

Generation of CG4607-mRuby3:

In order to determine whether the localisation of CG4607 was conserved we created a plasmid to express

CG4607 in mammalian cells. CG4607-mRuby3 construct was created through Gibson cloning (Gibson *et al.*

2009). The CG4607 cDNA (clone RH58543 #11058, Drosophila Genome Resource Center, Indiana, USA)

was PCR amplified using the following primers:

dCG4607GibF1: GGACTCAGATCTCGAGACAAGATGAAGGGCCAGCAGGAGGAG

dCG4607GibR1: CATGCTGCCttCAGCTGAGGACAATTTCTTTAGGAACACTT

The backbone GLUT4-mRuby3 was PCR amplified to include overhangs using the following primers:

mRuby3GibF1: TCCTCAGCTGAAGGCAGCATG

mRuby3GibR1: AGCTGAGGATCCCTTGTCTCGAGATCTGAGTCC

PCR products were placed together with Gibson master mix, and the resulting plasmid was sequenced before

cell transfection.

Cell Culture and Immunostaining:

In order to observe the localisation of CG4607 we expressed an ectopic construct, encoding CG4607 fused to

mRuby3, in mammalian HeLa cells. HeLa cells were kept in DMEM with 1% glutamax and 10% FCS at 37

degrees and 5% CO2. Cells were transfected with lipofectamine 2000 and split onto coverslips at two x10$_5$

cells/mL. To determine the localisation of CG4607 with lysosomes we used immunostaining with anti-

LAMP1 antibodies and then looked at colocalization between the lysosomal marker and CG4607-mRuby3.

Coverslips were fixed in 4% paraformaldehyde, washed with PBS, and blocked for 30 minutes with 0.02%

saponin (Sigma,) and 2% BSA in PBS. Primary antibodies: ms anti-LAMP1(1:100, Developmental Studies

Hybridoma Bank 4C CR). Secondary antibodies: (Gt anti-mouse 488 (1:200, Invitrogen)). Coverslips were

mounted in mowiol and imaged using a 60X water objective on the A1R confocal (Nikon). Colocalisation

analysis was performed using the coloc2 plugin in Fiji (Schindelin *et al.* 2012) ImageJ, NIH, Bethesda, MD).

qPCR of knockdown:

We wanted to determine the degree of whole body depletion of CG4607. We used qPCR to look at the number

of transcripts in control and CG4607 ᴋᴅ animals. Three replicates of 10 flies were homogenized in TRIzol™

Reagent (Invitrogen, 15596026) and RNA was precipitated out. cDNA was created (superscript II, Invitrogen

18064014). Tubulin was used as a housekeeping gene and the following primers were used to amplify

CG4607:

        CG4607 F3   ACTCCCACGCGAAGGAGAA
        CG4607 R3   GCTGATTGAGAGTAACTGCCG

463 The samples were run using the ROCHE Lightcycler 480 II (Roche). The knockdown efficiency was

464 calculated using the delta-delta Ct method (Excel and Graphpad, Prism) and the Ct values were graphed.

465 Significance between the control and knockdown transcript was calculated using a student's t- test (p<0.0001

466 ****).

467 Figure Legends:
468
Figure 1: Inter-strain variation in response to diet. (A) DGRP screen schematic outlines the work flow. Each

individually coloured fly represents an individual DGRP line and survival curves represent the measured starvation resistance

after exposure of flies to each diet. (B) Pairs plot showing a positive correlation with starvation resistance data from Mackay et al,

2012 and negative correlation with food intake to starvation resistance data on NF from this study. (C) Histograms showing the

distribution of the mean starvation resistance of each fly independent of strain on each diet. (D) Heatmap and hierarchical

clustering of the mean starvation resistance of each DGRP strain.

Figure 2: Mapping the SNPs in diet-responsive genes (A) Schematic of SNP analysis used in this study. (B) An

upset plot indicating the degree of overlap of the numbers of highly significant SNPs between each diet and the number of

SNPs per diet. (C) Manhattan plot of selected SNPs based on a multi-variate significance p-value of $< 1 \times 10^{-4}$, filtered by a

univariate significance of $p < 0.01$ and a fold change of >0.3. Highly significant SNPs with a univariate p-value <0.01 and a

fold change >0.3 are in black. The y-axis in the Manhattan plot is -log (P-value) for the multivariate test. (D-F) Manhattan

plots of selected SNPs based on a multi-variate significance p-value of $< 1 \times 10^{-4}$, filtered by a univariate significance of $p<$

0.01 and a fold change of >0.3 for each diet. Highly significant SNPs with a univariate p-value <0.01 and a fold change

>0.3 are in black. G) Quantile-Quantile plots of the distribution of significant MANOVA p-values for all SNPs.

Figure 3: Whole body knockdown of _CG4607_ is lethal upon HCD feeding. Survival of _ubi_-GAL4>_CG4607_

KK104152 flies on NF and HCD. (A) Adult male _ubi_-GAL4>_CG4607_ KK104152 flies (n=60) were placed on either NF or HCD and

manually monitored for lethality every 24 hours. We observed that all _ubi_-GAL4>_CG4607_ KK104152 flies were dead within 60

hours after HCD feeding. (B) Actogram of activity and sleep (n=16 flies) wrapped over 24 hours of single housed individual

control and _ubi_-GAL4>_CG4607_ KK104152 flies in DAMS monitors and beam crosses were monitored every 5 minutes while they

ate NF or HCD (C) The same actogram as in B, but only the activity over the first 9 hours is shown. Activity data was analysed

using the rethomics package in r (Geissmann _et al._ 2019). (D) Caloric intake of adult male flies (n=50/genotype) using the

CAFE assay. _ubi_-GAL4>_CG4607_ KK104152 flies ingested significantly more calories from HCD (p<0.02) than control. Data is

mean +/- SD and significance between genotypes was calculated using a t-test , ** p<0.01 (Prism).

Figure 4: _CG4607_ interacts with HPD to regulate starvation resistance. (A) Glycogen content was

significantly reduced between control flies fed NF and HPD. However, glycogen content was the same between NF and HPD fed

ubi-GAL4>_CG4607_ KK104152 flies (B) Fasted glycogen was reduced in NF fed control flies compared to NF fed _ubi_-

GAL4>_CG4607_ KK104152 flies, but pre-feeding with HPD reduced glycogen levels of _ubi_-GAL4>_CG4607_ KK104152 flies, while

starvation after control flies ate HPD was lethal. (C) Glucose content was not significantly different across genotypes. (D) Fasted

glucose levels of NF pre-fed flies had similar glucose levels, however, glucose levels were significantly reduced in HPD pre-fed

ubi-GAL4>*CG4607* KK104152. (E)Triglyceride levels of *ubi*-GAL4>*CG4607* KK104152 animals were increased on NF feeding, but

reduced upon eating HPD. (F) Fasted TAG stores were reduced in control flies pre-fed NF, while NF fed *CG4607*KD flies

maintained their TAG stores. (G) HPD pre-fed flies had reduced TAG stores. (all experiments n=36 flies/genotype and

representative of two experiments) Significance between genotypes was calculated using a one-way ANOVA , ** p<0.01, ***

p<0.001 (Prism).

Figure 5: *CG4607* controls glucose utilisation and uptake. (A-C) All the following experiments were performed on

n= 40 control and *ubi*-GAL4>*CG4607* KK104152 males as previously described (Francis *et al.* 2019) .(A) Glucose oxidation is

reduced after 4 hours of administrating C14 labelled glucose. (B) C14 incorporation into TAGs is reduced in *CG4607*KD flies. (C)

The accumulation of blue dye, which indicates food intake, is not significantly different between control and *CG4607*KD flies

during the assay. The data is from two independent experiments. Significance between genotypes was calculated using a student's

t-test, ** p<0.01, **** p<0.0001 (Prism). (D,E) Micrographs of fixed and immuno-stained HeLa cells were transfected with

pCMV-CG4607-mRuby3 (red), immunostained with anti-LAMP1 antibody (green) and counterstained with Hoechst (blue). (D,

inset (E)) CG4607-mRuby3 colocalises with Lamp1 expressing vesicles.

REFERENCES
513

Bjedov I., J. M. Toivonen, F. Kerr, C. Slack, J. Jacobson, *et al.*, 2010 Mechanisms of life span extension by

rapamycin in the fruit fly Drosophila melanogaster. Cell Metab 11: 35.

https://doi.org/10.1016/j.cmet.2009.11.010

Brand A., and N. Perrimon, 1993 Targeted gene expression as a means of altering cell fates and generating

dominant phenotypes. Dev Camb Engl 118: 401–15.

Carayannopoulos M. O., M. Chi, Y. Cui, J. M. Pingsterhaus, R. A. McKnight, *et al.*, 2000 GLUT8 is a

glucose transporter responsible for insulin-stimulated glucose uptake in the blastocyst. Proc National Acad

Sci 97: 7313–7318. https://doi.org/10.1073/pnas.97.13.7313

Chandegra B., J. Tang, H. Chi, and N. Alic, 2017 Sexually dimorphic effects of dietary sugar on lifespan,

feeding and starvation resistance in Drosophila. Aging 9: 2521. https://doi.org/10.18632/aging.101335

Chng W., V. Hietakangas, and B. Lemaitre, 2017 Physiological Adaptations to Sugar Intake: New

Paradigms from Drosophila melanogaster. Trends Endocrinol Metabolism 28: 131–142.

https://doi.org/10.1016/j.tem.2016.11.003

Crosby M. A., S. L. Gramates, G. dos Santos, B. B. Matthews, S. E. Pierre, *et al.*, 2015 Gene Model

Annotations for Drosophila melanogaster: The Rule-Benders. G3 Genes Genomes Genetics 5: 1737–1749.

https://doi.org/10.1534/g3.115.018937

Diril M., S. Schmidt, M. Krauß, V. Gawlik, H. Joost, *et al.*, 2009 Lysosomal localization of GLUT8 in the

testis – the EXXXLL motif of GLUT8 is sufficient for its intracellular sorting via AP1- and AP2-mediated

interaction. Febs J 276: 3729–3743. https://doi.org/10.1111/j.1742-4658.2009.07089.x

Djawdan M., A. K. Chippindale, M. R. Rose, and T. J. Bradley, 1998 Metabolic Reserves and Evolved

Stress Resistance in Drosophila melanogaster. Physiol Zool 71: 584–594. https://doi.org/10.1086/515963

DOEGE H., A. BOCIANSKI, H.-G. JOOST, and A. SCHÜRMANN, 2000 Activity and genomic

organization of human glucose transporter 9 (GLUT9), a novel member of the family of sugar-transport

facilitators predominantly expressed in brain and leucocytes. Biochem J 350: 771–776.

https://doi.org/10.1042/bj3500771

Drysdale R. A., M. A. Crosby, and F. Consortium, 2005 FlyBase: genes and gene models. Nucleic Acids

Res 33: D390–D395. https://doi.org/10.1093/nar/gki046

Dubois L., K. Kyvik, M. Girard, F. Tatone-Tokuda, D. Perusse, *et al.*, 2012 Genetic and environmental

contributions to weight, height, and BMI from birth to 19 years of age: an international study of over

12,000 twin pairs. Plos One 7: e30153. https://doi.org/10.1371/journal.pone.0030153

Efeyan A., and D. M. Sabatini, 2013 Nutrients and growth factors in mTORC1 activation. Biochem Soc T

41: 902–905. https://doi.org/10.1042/bst20130063

Feng Y., S. M. Hartig, J. E. Bechill, E. G. Blanchard, E. Caudell, *et al.*, 2010 The Cdc42-interacting Protein-

4 (CIP4) Gene Knock-out Mouse Reveals Delayed and Decreased Endocytosis. J Biol Chem 285: 4348–

4354. https://doi.org/10.1074/jbc.m109.041038

Francis D., J. R. Krycer, G. J. Cooney, and D. E. James, 2019 A modified gas-trapping method for high-

throughput metabolic experiments in Drosophila melanogaster. Biotechniques. https://doi.org/10.2144/btn-

2019-0062

Gallagher M. D., and A. S. Chen-Plotkin, 2018 The Post-GWAS Era: From Association to Function. Am J

Hum Genetics 102: 717–730. https://doi.org/10.1016/j.ajhg.2018.04.002

Garlapow M. E., W. Huang, M. T. Yarboro, K. R. Peterson, and T. F. Mackay, 2015 Quantitative Genetics

of Food Intake in Drosophila melanogaster. Plos One 10: e0138129.

https://doi.org/10.1371/journal.pone.0138129

Geissmann Q., L. Rodriguez, E. J. Beckwith, and G. F. Gilestro, 2019 Rethomics: An R framework to

analyse high-throughput behavioural data. Plos One 14: e0209331.

https://doi.org/10.1371/journal.pone.0209331

Gibson D. G., L. Young, R.-Y. Chuang, C. J. Venter, C. A. III, *et al.*, 2009 Enzymatic assembly of DNA

molecules up to several hundred kilobases. Nat Methods 6: nmeth.1318. https://doi.org/10.1038/nmeth.1318

Harbison S. T., L. J. McCoy, and T. F. Mackay, 2013 Genome-wide association study of sleep in Drosophila

melanogaster. Bmc Genomics 14: 281. https://doi.org/10.1186/1471-2164-14-281

Harshman, Hoffmann, and Clark, 1999 Selection for starvation resistance in Drosophila melanogaster:

physiological correlates, enzyme activities and multiple stress responses. J Evolution Biol 12: 370–379.

https://doi.org/10.1046/j.1420-9101.1999.00024.x

Hoffmann A. A., and L. G. Harshman, 1999 Desiccation and starvation resistance in Drosophila: patterns of

variation at the species, population and intrapopulation levels. Heredity 83: 6886490.

https://doi.org/10.1046/j.1365-2540.1999.00649.x

Johnson E. C., N. Kazgan, C. A. Bretz, L. J. Forsberg, C. E. Hector, *et al.*, 2010 Altered Metabolism and

Persistent Starvation Behaviors Caused by Reduced AMPK Function in Drosophila. Plos One 5: e12799.

https://doi.org/10.1371/journal.pone.0012799

Keene A. C., E. R. Duboué, D. M. nald, M. Dus, G. Suh, *et al.*, 2010 Clock and cycle Limit Starvation-

Induced Sleep Loss in Drosophila. Curr Biol 20: 1209–1215. https://doi.org/10.1016/j.cub.2010.05.029

Krycer J. R., L.-E. Quek, D. Francis, D. J. Fazakerley, S. D. Elkington, *et al.*, 2019 Lactate production is a

prioritised feature of adipocyte metabolism. J Biol Chem jbc.RA119.011178.

https://doi.org/10.1074/jbc.ra119.011178

Lee K., S. J. Simpson, F. J. Clissold, R. Brooks, W. J. Ballard, *et al.*, 2008 Lifespan and reproduction in

Drosophila: New insights from nutritional geometry. Proc National Acad Sci 105: 2498–2503.

https://doi.org/10.1073/pnas.0710787105

Lee M., S. Ha, J. Kim, A. Koh, C. Lee, *et al.*, 2009 Glycolytic Flux Signals to mTOR through

Glyceraldehyde-3-Phosphate Dehydrogenase-Mediated Regulation of Rheb ▽. Mol Cell Biol 29: 3991–4001.

https://doi.org/10.1128/mcb.00165-09

Lee K., and T. Jang, 2014 Exploring the nutritional basis of starvation resistance in Drosophila

melanogaster. Funct Ecol 28: 1144. https://doi.org/10.1111/1365-2435.12247

Li C., Y. Zhang, X. Yun, Y. Wang, M. Sang, *et al.*, 2014 Methuselah-like genes affect development, stress

resistance, lifespan and reproduction in Tribolium castaneum. Insect Mol Biol 23: 587–597.

https://doi.org/10.1111/imb.12107

Lisinski I., A. Schurmann, H. Joost, S. Cushman, and H. Al-Hasani, Targeting of GLUT6 (formerly

GLUT9) and GLUT8 in rat adipose cells. The Biochemical journal 358: 517.

Mackay T. F., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles, *et al.*, 2012 The Drosophila

melanogaster Genetic Reference Panel. Nature 482: 173. https://doi.org/10.1038/nature10811

Maedera S., T. Mizuno, H. Ishiguro, T. Ito, T. Soga, *et al.*, 2019 GLUT6 is a lysosomal transporter that is

regulated by inflammatory stimuli and modulates glycolysis in macrophages. Febs Lett 0: 195–208.

https://doi.org/10.1002/1873-3468.13298

Markow T., 2015 The secret lives of Drosophila flies. Elife 4: e06793. https://doi.org/10.7554/elife.06793

Matsuda H., T. Yamada, M. Yoshida, and T. Nishimura, 2015 Flies without Trehalose. J Biol Chem 290:

1244–1255. https://doi.org/10.1074/jbc.m114.619411

Mattila J., A. Bremer, L. Ahonen, R. Kostiainen, and O. Puig, 2009 Drosophila FoxO Regulates Organism

Size and Stress Resistance through an Adenylate Cyclase ▽ †. Mol Cell Biol 29: 5357–5365.

https://doi.org/10.1128/mcb.00302-09

May C. E., A. Vaziri, Y. Lin, O. Grushko, M. Khabiri, *et al.*, 2019 High Dietary Sugar Reshapes Sweet

Taste to Promote Feeding Behavior in Drosophila melanogaster. Cell Reports 27: 1675-1685.e7.

https://doi.org/10.1016/j.celrep.2019.04.027

Mayer A. L., C. B. Higgins, M. R. Heitmeier, T. E. Kraft, X. Qian, *et al.*, 2016 SLC2A8 (GLUT8) is a

mammalian trehalose transporter required for trehalose-induced autophagy. Sci Rep-uk 6: 38586.

https://doi.org/10.1038/srep38586

Mony V. K., S. Benjamin, and E. J. O'Rourke, 2016 A lysosome-centered view of nutrient homeostasis.

Autophagy 12: 619–631. https://doi.org/10.1080/15548627.2016.1147671

Mueckler M., and B. Thorens, 2013 The SLC2 (GLUT) family of membrane transporters. Mol Aspects Med

34: 121–138. https://doi.org/10.1016/j.mam.2012.07.001

Narita H., K. Tanji, Y. Miki, F. Mori, and K. Wakabayashi, 2019 Trehalose intake and exercise upregulate a

glucose transporter, GLUT8, in the brain. Biochem Bioph Res Co 514: 672–677.

https://doi.org/10.1016/j.bbrc.2019.05.017

Nelson C. S., J. N. Beck, K. A. Wilson, E. R. Pilcher, P. Kapahi, *et al.*, 2016 Cross-phenotype association

tests uncover genes mediating nutrient response in Drosophila. Bmc Genomics 17: 867.

https://doi.org/10.1186/s12864-016-3137-9

Parra-Peralbo E., and J. Culi, 2011 Drosophila Lipophorin Receptors Mediate the Uptake of Neutral Lipids

in Oocytes and Imaginal Disc Cells by an Endocytosis-Independent Mechanism. Plos Genet 7: e1001297.

https://doi.org/10.1371/journal.pgen.1001297

Pfeiffenberger C., B. C. Lear, K. P. Keegan, and R. Allada, 2010 Locomotor activity level monitoring using

the Drosophila Activity Monitoring (DAM) System. Cold Spring Harb Protoc 2010: pdb.prot5518.

https://doi.org/10.1101/pdb.prot5518

Reiter L., L. Potocki, S. Chien, M. Gribskov, and E. Bier, 2001 A systematic analysis of human disease-

associated gene sequences in Drosophila melanogaster. Genome Res 11: 1114.

https://doi.org/10.1101/gr.169101

Robinson S. W., P. Herzyk, J. A. Dow, and D. P. Leader, 2013 FlyAtlas: database of gene expression in the

tissues of Drosophila melanogaster. Nucleic Acids Res 41: D744–D750. https://doi.org/10.1093/nar/gks1141

Rodríguez-Vázquez M., D. Vaquero, E. Parra-Peralbo, J. E. Mejía-Morales, and J. Culi, 2015 Drosophila

Lipophorin Receptors Recruit the Lipoprotein LTP to the Plasma Membrane to Mediate Lipid Uptake. Plos

Genet 11: e1005356. https://doi.org/10.1371/journal.pgen.1005356

Sales N., P. Pelegrini, and M. Goersch, 2014 Nutrigenomics: Definitions and Advances of This New

Science. J Nutrition Metabolism 2014: 202759. https://doi.org/10.1155/2014/202759

Sano H., A. Nakamura, M. J. Texada, J. W. Truman, H. Ishimoto, *et al.*, 2015 The Nutrient-Responsive

Hormone CCHamide-2 Controls Growth by Regulating Insulin-like Peptides in the Brain of Drosophila

melanogaster. Plos Genet 11: e1005209. https://doi.org/10.1371/journal.pgen.1005209

Schindelin J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, *et al.*, 2012 Fiji: an open-source

platform for biological-image analysis. Nat Methods 9: 676. https://doi.org/10.1038/nmeth.2019

Skorupa D. A., A. Dervisefendic, J. Zwiener, and S. D. Pletcher, 2008 Dietary composition specifies

consumption, obesity, and lifespan in Drosophila melanogaster. Aging Cell 7: 478–490.

https://doi.org/10.1111/j.1474-9726.2008.00400.x

Stunkard A., T. Foch, and Z. Hrubec, A twin study of human obesity. JAMA 256: 51.

Tang H., Y. Wang, S. Wang, M. Wu, S. Lin, *et al.*, 2011 Atg1-mediated myosin II activation regulates

autophagosome formation during starvation-induced autophagy. Embo J 30: 636–651.

https://doi.org/10.1038/emboj.2010.338

Unckless R. L., S. M. Rottschaefer, and B. P. Lazzaro, 2015 A genome-wide association study for

nutritional indices in Drosophila. G3 Genes Genomes Genetics 5: 417.

https://doi.org/10.1534/g3.114.016477

Yasugi T., T. Yamada, and T. Nishimura, 2017 Adaptation to dietary conditions by trehalose metabolism in

Drosophila. Sci Rep-uk 7: 1619. https://doi.org/10.1038/s41598-017-01754-9

Zhao H., M. Tang, M. Liu, and L. Chen, 2018 Glycophagy: An emerging target in pathology. Clin Chim

Acta 484: 298–303. https://doi.org/10.1016/j.cca.2018.06.014

Zobel T., K. Brinkmann, N. Koch, K. Schneider, E. Seemann, *et al.*, 2015 Cooperative functions of the two

F-BAR proteins Cip4 and Nostrin in the regulation of E-cadherin in epithelial morphogenesis. J Cell Sci

128: 499–515. https://doi.org/10.1242/jcs.155929

ubi>Control NF ■ ubi>CG4607 KD HCD
ubi>CG4607 KD NF ■ ubi>Control HCD

Table 1: Analysis of the mean starvation resistance data on all diets.

Diet	Mean	SEM	Variance (Genetic)	Broad Sense Heritability
NF	51.46	1.15	0.086	19%
HCD	68.88	1.64	0.364	50%
HFD	55.10	1.09	0.089	20%
HPD	36.77	0.97	0.671	65%

A summary table of the mean starvation data, SEM, genetic and environmental variance from the log transformed starvation data across all the DGRP lines on each diet.

Estimates of genetic variance and heritability are derived from linear mixed models of log survival (see supplementary table S2).

Table 2: GO term analysis of human orthologs of diet-responsive genes

HPD "Down"						
GO:ID	p.value	adj.p.value	Genes In	Genes Out	TERM	ONTOLOGY
GO:0004888	1.84E-08	1.07E-04	17	143	transmembrane signaling receptor activity	MF
GO:0038023	1.78E-07	3.44E-04	19	213	signaling receptor activity	MF
GO:0060089	1.57E-07	3.44E-04	20	234	molecular transducer activity	MF

GO ID	p.value	adj.p. value	Genes In	Genes Out	TERM	ONTOL OGY
GO:0005 272	9.34E-07	1.35E -03	6	12	sodium channel activity	MF
GO:0015 144	2.22E-06	2.57E -03	5	7	carbohydrate transmembrane transporter activity	MF
GO:0005 887	6.56E-06	6.33E -03	22	357	integral component of plasma membrane	CC

HFD "DOWN"						
GO ID	p.value	adj.p. value	Genes In	Genes Out	TERM	ONTOL OGY
GO:0015 294	7.51E-07	1.09E -03	9	30	solute:cation symporter activity	MF
GO:0038 023	3.10E-07	1.09E -03	22	210	signaling receptor activity	MF
GO:0043 410	7.25E-07	1.09E -03	19	168	positive regulation of MAPK cascade	BP
GO:0060 089	3.78E-07	1.09E -03	23	231	molecular transducer activity	MF
GO:0019 199	1.25E-06	1.44E -03	7	15	transmembrane receptor protein kinase activity	MF
GO:0015 370	1.64E-06	1.58E -03	8	24	solute:sodium symporter activity	MF
GO:0004 714	2.48E-06	2.05E -03	6	10	transmembrane receptor protein tyrosine kinase activity	MF

GO ID	p.value	adj.p. value	Genes In	Genes Out	TERM	ONTOLOGY
GO:0034 308	3.42E-06	2.47E-03	8	27	primary alcohol metabolic process	BP
GO:0004 888	6.95E-06	4.02E-03	16	144	transmembrane signaling receptor activity	MF
GO:0043 408	6.91E-06	4.02E-03	21	237	regulation of MAPK cascade	BP
HFD "UP"						
GO ID	**p.value**	**adj.p. value**	**Genes In**	**Genes Out**	**TERM**	**ONTOLOGY**
GO:0004 180	5.29E-09	3.06E-05	5	11	carboxypeptidase activity	MF
GO:0008 238	3.72E-06	1.08E-02	5	50	exopeptidase activity	MF
HCD "DOWN"						
GO ID	**p.value**	**adj.p. value**	**Genes In**	**Genes Out**	**TERM**	**ONTOLOGY**
GO:0034 035	2.43E-07	7.02E-04	5	9	purine ribonucleoside bisphosphate metabolic process	BP
GO:0050 427	2.43E-07	7.02E-04	5	9	3'-phosphoadenosine 5'-phosphosulfate metabolic process	BP
GO:0008 146	2.32E-06	4.29E-03	5	16	sulfotransferase activity	MF
GO:0030 902	2.97E-06	4.29E-03	8	76	hindbrain development	BP
GO:0051 923	5.89E-06	6.81E-03	4	8	sulfation	BP

			HCD "UP"			
GO ID	**p.value**	**adj.p. value**	**Genes In**	**Genes Out**	**TERM**	**ONTOL OGY**
GO:0005 201	1.81E-08	1.05E -04	5	38	extracellular matrix structural constituent	MF
GO:0005 581	5.98E-08	1.73E -04	4	16	collagen trimer	CC
GO:0005 604	7.12E-07	1.37E -03	4	32	basement membrane	CC
GO:0062 023	2.65E-06	3.83E -03	5	110	collagen-containing extracellular matrix	CC
GO:0030 198	5.63E-06	5.18E -03	5	129	extracellular matrix organization	BP
GO:0031 012	6.27E-06	5.18E -03	5	132	extracellular matrix	CC
GO:0044 420	5.95E-06	5.18E -03	3	15	extracellular matrix component	CC

Tables show the human orthologs of Drosophila genes associated with a predicted decrease (down) or increase (up) in starvation resistance upon HFD,HPD, or HCD exposure. GO ID with a p-value less than p<1E-5 were selected.

Table 3: Validated candidate genes.

Gene/ Ortholog[a]	SNP Class	Manova p-value[b]	Diet[c]	vHCD[d]	vHPD[e]	vHFD[f]	G X D pValue[g]	Function[h]
CG8311/ DOLK	NON_SYN ONYMOU S_CODIN G	3.70E-10	HCD	0.17	-0.67	-0.47	7.70E- 01	protein glycosylation
otp/OTP	NON_SYN ONYMOU S_CODIN G	2.40E-07	HFD	0.8	0.1	-0.07	6.60E- 01	DNA binding

LpR2/ VLDLR	INTRON	4.20E-06	HFD	0.13	0.31	-0.12	6.00E-01	lipid transport
sqa/ MYLK3	INTRON	8.20E-09	HFD	0.48	0.22	0.15	1.80E-01	starvation-induced autophagy
mthl3/ ADGRG7	DOWNSTREAM	6.90E-07	HFD	-0.06	N/A	-0.29	1.20E-01	response to starvation, determination of adult lifespan
hh/Shh	INTRON	1.80E-05	HFD	-0.04	N/A	-0.43	7.10E-02	segment polarity determination, stem cells maintenance and cell migration
hppy/ MAP4K3	INTRON	4.20E-06	HCD	-0.55	N/A	-0.3	2.30E-02	triglyceride homeostasis, regulation of TOR signaling, behavioral response to ethanol
Cip4/ TRIP10	INTRON	1.60E-06	HFD	-0.31	0.34	-0.94	1.30E-02	lipid binding
Gyc88E/ GUCY1B1	NON_SYNONYMOUS_CODING	1.10E-06	HPD	0.68	0.02	0.19	1.10E-02	neuronal oxygen detector, activated at reduced oxygen levels
CG3339/DNAH9	NON_SYNONYMOUS_CODING	3.70E-07	HFD	-0.11	-0.07	0.44	4.10E-03	dynein light chain binding
CG43345/ PLD3-4	NON_SYNONYMOUS_CODING	6.70E-08	HFD	0.42	-0.51	-0.67	1.10E-03	Phospholipase D/Transphosphatidylase;

CCHa2-R/BRS3	INTRON	1.10E-05	HFD	-0.01	0.15	-0.77	2.80E-04	neuropeptide signaling pathway
CG9674/ DPYD	UTR_5_PRIME	5.40E-07	HCD	-1.16	0.19	-0.23	1.50E-04	glutamate synthase (NADH) activity
T3dh/ ADHFE1	NON_SYNONYMOUS_CODING	2.60E-06	HCD	0.1	-0.24	-0.33	1.10E-05	alcohol dehydrogenase (NAD) activity
fd96Ca/ FOXB1	NON_SYNONYMOUS_CODING	9.20E-07	HFD	-0.4	-0.45	-0.99	1.00E-10	regulation of transcription
CG4607/ SLC2A6	UPSTREAM	2.20E-05	HPD	DOA	-1.29	-0.11	0.00E+00	glucose transmembrane transporter activity

a. Gene symbol/human ortholog

b. the significance from the multivariate anova of the diet responsive SNP

c. the diet upon which the gene is predicted to affect starvation resistance

d-f. the fold change in starvation resistance (SRcontrol $_{Diet/NF}$ / SRKD $_{Diet/NF}$)

g. The significance of the gene by diet interaction for each gene and it's fold change on the predicted diet (from d).

h. The functional annotation of the gene from www.flybase.org