Title: **Relationships between landscape structure and the prevalence of two tick-borne infectious**

agents, *Anaplasma phagocytophilum* and *Borrelia burgdorferi* sensu lato, in small mammal

communities

Running title:

Landscape structure and prevalence of two tick-borne infections in small mammals

Authors:

Grégoire Perez[1, 2, *]

Suzanne Bastian[2]

Amélie Chastagner[3, 4, §]

Albert Agoulon[2]

Yann Rantier[1]

Gwenaël Vourc'h[3]

Olivier Plantard[2]

Alain Butet[1]

[1] UMR CNRS 6553–Université de Rennes 1, ECOBIO, Campus de Beaulieu, 35042 Rennes, France

[2] BIOEPAR, INRA, Oniris, 44307, Nantes, France

[3] EPIA, INRA, VetAgro Sup, 63122, Saint-Genès-Champanelle, France

[4] Evolutionary Ecology Group, University of Antwerp, Campus Drie Eiken, Universiteitsplein 1, 2610,

Wilrijk, Belgium

* Corresponding author. E-mail: perez.gregoire[at]gmail.com

ORCID : 0000-0001-8861-4856

Current affiliation: UMR BIPAR, INRA-Anses-ENVA, 94701 Maisons-Alfort, France

[§] Current affiliation: ANSES, Laboratoire de Ploufragan-Plouzané-Niort, 22440 Ploufragan, France

Abstract

Context

By modifying ecosystems, land cover changes influence the emergence, the spread and the incidence of vector-borne diseases.

Objective

This study aimed at identifying associations between landscape structure and the prevalence of two tick-borne infectious agents, *Anaplasma phagocytophilum* and *Borrelia burgdorferi* s.l., in small mammal communities.

Methods

Small mammals were sampled in 24 sites along a gradient of woodland fragmentation and hedgerow network density, and screened for infectious agents with real-time PCR techniques. For each site, structural variables (composition and configuration) of the surrounding landscape at various scales (0-500 m) and variables of wooded habitats connectivity based on graph theory and least cost path distances for the two dominant species, bank voles (*Myodes glareolus*) and wood mice (*Apodemus sylvaticus*), were computed.

Results

The *A. phagocytophilum* prevalence increased with wooded habitats cover (0-500 m), likely through host population size, and increased slightly with bank vole abundance, which has a higher reservoir competence than wood mouse. The *B. burgdorferi* s.l. prevalence increased with wooded ecotones only at local scales (50-100 m). Wooded habitats connectivity measures did not improve models built with simple land cover variables. A more marked spatial pattern was observed for the prevalence of *A. phagocytophilum* than *B. burgdorferi* s.l..

Conclusions

This study highlights the interest of considering together the ecology of infectious agents (e.g. host specificity) and the host species community ecology to better understand the influence of the landscape structure on the spatial distribution of vector-borne infectious agents.

Keywords

Tick-borne diseases; Small mammal community; Landscape connectivity; Graph theory; Least cost paths

Introduction

Landscape changes are suspected to cause infectious disease emergence or reemergence worldwide (Jones et al. 2008). Several authors depicted the complexity of the interactions between the landscape, hosts, vectors and humans, which drive the emergence of these diseases (Lambin et al. 2010). The spread of infectious agents in ecosystems relies partly on their life history traits shaped by their evolutionary histories (e.g. transmission modalities). Because species within communities exhibit various levels of reservoir or vector competence, the transmission depends on hosts and vectors diversity and abundance (Keesing et al. 2009).

Landscape structure (composition and configuration), by filtering and regulating richness, abundance and dispersal of species in hosts and vectors communities, may influence the transmission rate and the diffusion of infectious agents, which ultimately determine their distribution and prevalence (Suzán et al. 2015). For instance, an increase in the proportion of non-reservoir hosts may induce an increased number of missed transmissions between reservoir hosts, or an increased number of wasted bites (failed transmission events) by vectors. This may reduce the number of transmission events and has a so called "dilution effect" on prevalence (Ostfeld and Keesing 2000; Clay et al. 2009). Conversely, an increase in the proportion of competent hosts may increase the number of transmission events or efficient bites by vectors (Roche et al. 2013). When less competent species are more prone to local extinction, this non-random species loss in host communities seems to be a driver of the increased direct and vector-borne infectious disease transmission (Ostfeld and LoGiudice 2003; Johnson et al. 2013; Roche et al. 2013).

Habitat connectivity partly shapes hosts community structure. Large-bodied species, with small clutch/litter size and long gestation/incubation time are supposed to be more sensitive to habitat fragmentation because they experience higher local extinction risk. Conversely, small-bodied species, with large clutch/litter size and short incubation/gestation time can persist in fragmented habitat landscapes. These species need less habitat surface and benefit from a competition and predation release in these landscapes that are less favorable to predators and specialist species (Nupp and Swihart 1998, 2000; Gehring and Swihart 2003; Ferrante et al. 2017). As it happens, the latter kind of species also seem to be better reservoirs for infectious agents (Gottdenker et al. 2012; Huang et al. 2013; Ostfeld et al. 2014). Thus, in fragmented habitat landscapes, one could expect an amplification of infection prevalence supported by a higher density of more competent hosts (Allan et al. 2003; Rubio et al. 2014). However, habitat connectivity can also indirectly influence the risk of local extinction of infectious agents by shaping hosts – and vectors – population size, as demonstrated on Hantavirus in bank voles (Guivier et al. 2011). Land use is highly changing in Europe (agricultural intensification in some areas and rewilding in others; Reidsma et al. 2006). Thus, more studies explicitly assessing how landscape composition, configuration, and the resulting functional

connectivity influence interactions and abundances of hosts, vectors and infectious agents to better

understand infectious diseases risk are needed.

Small mammals are reservoir hosts of many tick-borne infectious agent taxa: protozoan (e.g. *Babesia*

sp., Hersh et al. 2012), bacteria (e.g. *Borrelia* sp., Gern et al. 1998; *Anaplasma* sp., Stuen et al. 2013)

and viruses (e.g. tick-borne encephalitis virus, Mansfield et al. 2009). Small mammal communities

composition is known to vary according to landscape features like habitat patch size, shape and

isolation (Nupp and Swihart 2000; Michel et al. 2007) and to the constraints that the landscape matrix

imposes on the displacements of small mammals (Szacki et al. 1993).

Tick displacements *per se* are reduced and their dispersal occurs mostly while feeding on their hosts

(Araya-Anchetta et al. 2015). Hard ticks (Arthropoda, Arachnida, Ixodidae) need to have a first blood

meal on a vertebrate host to molt from larva to nymph, a second one to molt from nymph to adult then

adult females need a third one to produce eggs once fertilized by adult males. Tick-borne infectious

agents can be acquired by ticks or transmitted to hosts at each blood meal. They disperse either by the

dissemination of ticks getting infected while feeding on infected hosts, or by already infected ticks

while attached on hosts. The dispersal distance of infectious agents depends on home ranges and

competences of tick hosts (Kurtenbach et al. 2002b). Thus, landscape features affecting host

communities and/or their dispersal are of paramount importance in the distribution and the local

prevalence of tick-borne infections.

This study focused on *Anaplasma phagocytophilum* and *Borrelia burgdorferi* sensu lato (s.l.). The

A. phagocytophilum bacteria ecotype associated to small mammals is transmitted mostly by

I. trianguliceps and maybe other endophilic (burrow dwelling) tick species (Bown et al. 2009;

Blaňarová et al. 2014; Jahfari et al. 2014). Infections in small mammals last about a couple of months

(Bown et al. 2003). No transovarial transmission (from engorged females to their offspring) of these

bacteria is known in *Ixodes* ticks (Stuen et al. 2013).

In Europe, several genospecies of the *B. burgdorferi* s.l. complex can be hosted by small mammals:

B. afzelli, *B. bavariensis* (formerly *garinii* OspA serotype 4 strain), *B. bissetti*, *B. burgdorferi* sensu

stricto (s.s.), and *B. spielmani* (Kurtenbach et al. 2002a, 2006; Margos et al. 2009; Coipan and Sprong

2016). However, host ranges of some of these genospecies include also larger mammals like

hedgehogs (*Erinaceus europaeus*) and squirrels (*Sciurus vulgaris*) (Skuballa et al. 2012; Pisanu et al.

2014). The *B. burgdorferi* s.l. bacteria can be transmitted by several tick species, but the exophilic

(questing for host on the vegetation) tick species *I. ricinus* is assumed to be its main vector in Europe

(Rizzoli et al. 2011). Small mammals are major hosts for *I. ricinus* larvae and only occasional hosts for

nymphs, but they also host endophilic tick species like *I. trianguliceps* and *I. acuminatus* at all life

stages (Boyard et al. 2008; Bown et al. 2008; Hofmeester et al. 2016; Perez et al. 2017). Therefore, the

maintenance of the zoonotic cycle of these bacteria may partly rely on other tick species (Hubbard et

al. 1998; Heylen et al. 2013; Szekeres et al. 2015). Infections in small mammals are lifelong (Gern et

al. 1994; Humair et al. 1999). Transovarial transmission is assumed to be null or rare (Rollend et al.

2013).

Here, we investigated at a local scale how landscape composition, configuration, and resulting habitat

connectivity for small mammals, directly or *via* relationships with species abundances, can explain the

prevalence of *A. phagocytophilum* and *B. burgdorferi* s.l. in small mammals. We specifically tested

the three following hypothesis on the prevalence drivers:

(**H1**) Host population size: prevalence would be higher in sites supporting larger host

populations (i.e. in wooded habitat patches either of large area, with more surrounding wooded

habitats and/or more connected).

(**H2**) Overall host community competence: the landscape would indirectly influence the

prevalence by acting on the relative abundance of host species exhibiting different reservoir

competence, modifying the overall community competence.

(**H3**) Infectious agents' specificity: a stronger relationship between the prevalence and

landscape variables would be observed for the *A. phagocytophilum* ecotype specific to small

mammals, than for the *B. burgdorferi* s.l. group, which shows a wider range of host and of vector

species.

Materials and Methods

Study area

The study took place in the 'Zone Atelier Armorique' (Brittany, France), a 132 km² Long-Term

Ecological Research (LTER) area labeled by the CNRS (Centre National de la Recherche Scientifique)

and belonging to the International LTER network. This area includes different landscapes with various

land uses, from deciduous mixed forest, to livestock-crop mixed farming landscapes, to cereal-

oriented (mainly cereals and maize) open farming landscapes. Compared to mixed farming landscapes,

open farming landscapes have fewer (and smaller) woodlots (area: 10% vs 15%), fewer grasslands

(area: 20% vs. 40%), larger crops patches (mean area: 2.3 ha vs. 1.3 ha), and looser hedgerow

networks (50 vs. 115 m/ha) (Millán de la Peña et al. 2003; Al Hassan et al. 2012; Perez et al. 2016).

Sampling strategy

Wooded habitats support higher abundances of small mammals, better abiotic conditions for ticks

(moisture) and are as such supposed to be key habitats for small mammals-ticks interactions (Boyard

et al. 2008). For these reasons, 24 sampling sites were selected in wooded habitat patches in various

landscape contexts (Perez et al. 2016). The 24 sampling sites, at least 500 m apart from each other to

ensure their spatial independence, were placed as follows: six in the forest core; six in forest edge

landscapes; six in mixed farming landscapes; and six in open farming landscapes. In each of the two

agricultural landscape types, three sites were selected along hedgerows (six sites overall) and three

along woodlots edges (six sites overall). At forest edges and in agricultural landscapes, the sampling

sites (n = 18) were selected at the ecotones between wooded habitats and grasslands, as meadows are

less unfavorable to ticks than cultivated plots. Because small mammals start breeding in spring and

populations peak generally in autumn, we sampled in May-June and in October in 2012 and 2013.

These periods of the year also correspond to the main and the secondary activity peak of *I. ricinus*

nymphs, respectively, and conversely for *I. trianguliceps* adult females, while the nymphs of this latter

species are active from summer to mid-autumn (Randolph 1975a).

Small mammal sampling and ethic statement

Small mammals were trapped using French model (INRA) live traps with wooden dormitory boxes.

These traps can catch small mammals from 4 to 40 grams, including shrews and small rodents. Trap-

lines (100 m-long) were constituted of 34 traps (three meters apart) baited with a mix of commercial

seeds for pet rodents (sunflower, wheat, etc.), dry cat food, and a piece of apple for water supply.

Traps were checked in the morning after 24 and 48 hours. Animals were brought to the field lab to be

identified at the species level, euthanized by pentobarbital injection, weighted, sexed and dissected.

Traps were designed to limit as much as possible the stress or injury of the animals. The bait was

designed to obtain an optimal catch rate across small mammal species and a good survival of

individuals. Targeted animals were not protected species and thus no special authorization was needed,

according to the French law in force. All individuals were euthanized by authorized experimenters

according to current French law and to the European guidelines on the use of animals in science

(Close et al. 1997).

Molecular detection of infectious agents

DNA was extracted from small mammal spleens and ears with Macherey-Nagel NucleoSpin Tissue

kits (Chastagner et al. 2016). The screening for *A. phagocytophilum* was performed on DNA extracts

from small mammal spleens by real-time PCR targeting the *msp2* genes according to the protocol of

Courtney et al. (2004). The screening for *B. burgdorferi* s.l. was performed on DNA extracts from

small mammal ears by real-time PCR in SybrGreen according to the protocol of Halos et al. (Halos et

al. 2010). The *B. burgdorferi* s.l. prevalence was low and, for technical reasons, identifying them to

the geno-species level (genetically discriminated *Borrelia* species) was not possible (data not shown).

Thus, all the *B. burgdorferi* s.l. genospecies were combined hereafter. Although all geno-species

which can infect small mammals do not share the same host range, *B. afzelii* is known to be

predominant in the captured species (Kurtenbach et al. 2002a; Marsot et al. 2013). Therefore, this

simplification is assumed not to bias results. As temporal the variation of prevalence has been studied

elsewhere (Perez et al. 2017), the data from all the sampling sessions were grouped for each site to

study only the spatial variation hereafter.

Landscape variables

Landscape data

The used landscape data were: a shapefile of the land cover of 2012 kindly provided by the 'LETG-

COSTEL-Rennes' lab *via* the 'Zone Atelier Armorique'; a road network shapefile provided by the

'Institut National de l'Information Géographique et Forestière' (IGN); and a 5 m-resolution raster file

of the land cover of 2010 from Gil Tena et al. (2014) for woodland and hedgerows. The land covers

were characterized as "woodland", "hedgerows" (abandoned lands and hedgerows extracted from the

5 m-resolution raster file), "grassland", "crops", "roads" (ranked according to the IGN's 'road

importance index'), "buidings" and "water areas". All files were aggregated with ArcGIS 10.3.1 for

Desktop, Esri ®, into a single 5 m-resolution raster file that was used in subsequent analyses.

Landscape composition and configuration variables

The proportion of wooded habitats ('Wooded' hereafter), computed as the proportion of

"woodland"/"hedgerows" pixels, was chosen as a landscape composition variable representative of the

amount of favorable permanent habitats surrounding trap-lines. The proportion of wooded ecotones

surrounding trap-lines ('Ecotones' hereafter), computed as the proportion of 'wooded habitat' and

"grassland"/"crops"/"roads" adjacent pixels (pixel couples), was chosen as a landscape configuration

variable. Such ecotones are known to be favorable to generalist species and are places where species

from different habitat can co-occur (Meunier et al. 1999; Ouin et al. 2000; Boyard et al. 2008). This

configuration variable depends on landscape heterogeneity, and is maximal in landscapes with

moderate woodland fragmentation.

The relationships between landscape variables and small mammal abundances or infection prevalence

were evaluated at various spatial scales by computing these variables in circular zones of various radii

around trap-line centres with Chloe2012 software (Boussard and Baudry 2014). Because trap-lines

were 100 m-long, the first circular 50 m-radius zones were considered as informative on trapping

habitats only ('local'). Larger circular zones with an additional 50, 100, 200, and 500 m radius were

considered.

Landscape habitat connectivity metrics

The dispersal of the studied infectious agents is known to rely mostly on larval ticks, which acquire
them by feeding on an infected small mammal and falling off in the latter's home range. The dispersal
of infectious agents by young small mammals, which have not yet been highly exposed to ticks and
are thus yet unlikely infected, can be considered as negligible (Randolph 1975b; Sinski et al. 2006;
Kallio et al. 2014; Perez et al. 2017). Furthermore, despite *I. ricinus* nymphs can disperse bacteria on
longer distances attached on hosts with larger home ranges, small mammals are unlikely to be infected
by *I. ricinus* adult females because they rarely host them. For these reasons, we performed the
functional connectivity analysis at the scale of small mammals' home ranges.

Thus, we modelled the functional connectivity of the landscape for small mammals to evaluate its
relationship with their infection prevalence. We used the difference in Probability of Connectivity
metric ('dPC'), which is an integrative measure of reachable habitats based on a negative exponential
function (Saura and Pascual-Hortal 2007). The three dPC fractions were also used separately: the
intra-patch connectivity ('dPCintra'; no inter-patch displacement/dispersal), which is dependent of
patch area only, so this latter variable was used instead for simplification purpose ('Area'); the inter-
patch connectivity ('dPCflux') and the "stepping stone" connectivity ('dPCconnector'; connecting
other patches).

Land cover cost files were built only for the two dominant rodent species with contrasted habitat
preference: wood mice and bank voles (Boyard et al. 2008). Because these species move preferentially
along linear structures or across the shortest path between suitable habitat patches (Zhang and Usher
1991), the functional distance between two habitat patches was set using least cost path distances. For
"woodland", "grassland" and "crops", costs were assigned like the inverse of the relative species
abundance from the literature (Ouin et al. 2000; Tattersall et al. 2002; Boyard et al. 2008; Vourc'h et
al. 2008). The reference cost for "woodland" was one. For wood mice and bank voles, the chosen
costs were respectively four and fifteen for "grassland", and five and fifteen for "crops". A maximum
cost of 100 was assigned arbitrary to "water areas" and "buildings". Roads are repulsive habitats for
small mammals because they impede their displacements (Mader 1984; Rico et al. 2007). They also
cause a traffic-related mortality (Ruiz-Capillas et al. 2015). Eskilkdsen (2010) reported that finding
bank voles were 15 times less likely on the opposite side of a barrier (forest path or road) than on the
same side. Thus, a gradient of costs from 15 (countryside and forest tracks) to 75 (expressways) was
assigned to "roads" according to the 'road importance index'.

The median distance travelled by an individual was modelled using negative exponential functions
based on data from the same landscapes (Papillon et al. 2002), and from a forest landscape of the
Berkshire, United-Kingdom (Kikkawa 1964). In consistency with the lower mobility of bank voles
compared to wood mice, these distances were estimated at 25 and 100 m, respectively (Zhang and
Usher 1991). The nodes were defined as patches with an area of at least 0.025 ha, corresponding to the
minimum home range size of the studied species (Kikkawa 1964). Graphs were created based on least

cost path networks for each species. To compute connectivity measures with 'isolation by distance'

models, graphs based on Euclidian distance were also created for displacement/dispersal distances

corresponding to half the radius used to compute land cover variables: 25, 50, 100, and 250 m. Nodes

were adjacent woodland habitats pixels ("woodland" and "hedgerows").

Connectivity measures were computed at the patch level for each graph. Because habitat patch areas

and 'dPC' values were over-dispersed, those variables were log-transformed hereafter. Least cost

paths, graphs, and connectivity measures were computed using Graphab 1.2.3 (Foltête et al. 2012). An

example of resulting graphs is shown in **Appendix 1**. A summary of all the landscape variables is

shown in **Table 1**.

Data analyses

To determine whether prevalence in wood mice and bank voles should be modelled separately, we

first tested, for each infection, the correlation between prevalence in each species and the correlation

between numbers of infected individuals of each species. Because of the non-normal distribution of

prevalence data (Shapiro-Wilk test: $p < 0.05$), we used Spearman's correlation tests.

To assess indirect effects of landscape structure on prevalence *via* the abundances of small mammal

species, the relationship between these latter and landscape variables was tested. First, to determine

whether wood mouse and bank vole abundances could be modelled separately, their correlation was

tested through Pearson's correlation test. Then, to choose the most appropriate error distribution of

species abundances, Akaike Information Criterion corrected for finite sample size (AICc; Hurvich and

Tsai 1989) values of the null models fitted with a normal error distribution and with a Poisson error

distribution were compared. The normal error distribution had a better support (lower AICc-value) for

wood mouse abundance (AICc-normal = 180.2, AICc-Poisson = 232.0) and for bank vole abundance

(AICc-normal = 141.6, AICc-Poisson = 154.9). Species abundances were thus modelled using Linear

Models (LMs).

The variables were first selected in single explanatory variable models ($p < 0.1$). Then, multiple

explanatory variable LMs were built with the selected variables. To avoid collinearity problems,

models with correlated variables ($|r| > 0.5$) were excluded. The models were then ranked based on

AICc values, and the significance of the variable in the best models (ΔAICc < 2) was evaluated with

type II ANOVAs ($p < 0.05$).

To assess whether prevalence were related to landscapes variables or species abundances, prevalence

in small mammals per site was modelled using binomial Generalized Linear Models (GLMs) with the

same explanatory landscape variables as above, with addition of wood mouse abundance (N.As), bank

vole abundance (N.Mg), and the ratio between the bank vole abundance and the wood mouse

abundance (ratio.Mg:As). Abundances were expressed as the total number of captured individuals per

site. The variables were first selected in single explanatory variable models (p < 0.1). Another AICc-

based model ranking and variable significance evaluation was finally performed on multiple

explanatory variable GLMs, as described above.

All statistical analyses were performed using R software Version 1.2.5001 (R Development Core

Team 2019). The R-package 'car' was used for ANOVAs, 'MuMIn' for model selections, and 'psych'

for pairwise correlation tests between explanatory variables. The absence of autocorrelation in the

dependent variables was checked using the 'correlog' function of the 'ncf' R-package.

Results

A total of 612 small mammals belonging to five species were caught during the two sampling years

(see **Table 2** for detailed results; the whole data set is available in supplementary material). Wood

mice (*Apodemus sylvaticus*) and bank voles (*Myodes glareolus*) were largely dominant (74.2% and

24.3% of captured animals, respectively) and found in all the 24 sites. We captured nine individuals of

three other species on seven sites: four field voles (*Microtus agrestis*) on three sites, two common pine

voles (*Microtus subterraneus*) on two sites, and three crowned shrews (*Sorex coronatus*) on three

sites. A maximum of four species were present in a single forest core site with all species excepting

the common pine vole.

Wood mouse abundance and bank vole abundance were not correlated (rho = 0.229, p = 0.281) and

thus have been analysed separately hereafter. The best model for wood mouse abundance included

only the proportion of ecotones in 500 m-radius zones (R^2 = 0.290, p = 0.003) with a positive

relationship. Bank vole abundance was not significantly related to any landscape variable (p > 0.05).

Results for A. phagocytophilum

The analyses of *A. phagocytophilum* prevalence were based on the PCR results for 452 wood mice,

147 bank voles, four field voles, two common pine voles and three crowned shrews. Twenty wood

mice, twenty bank voles and two crowned shrews were positive (**Table 2**). The number of positive

wood mice and bank voles per site were significantly correlated (rho = 0.425, p = 0.038). However,

the prevalence were not (rho = 0.291, p = 0.177). The results of the selection procedure of the GLMs

of *A. phagocytophilum* prevalence as a function of landscape variables and species abundance

variables are detailed in **Table 3**.

The most supported GLM of *A. phagocytophilum* prevalence in all small mammals included bank vole

abundance, (**Figure 1a**), and the proportion of wooded habitats in 50 m-radius zones, both displaying

a significant positive relationship with the prevalence (**Figure 1a** and **1b**, respectively). The most

supported GLM of prevalence in wood mice included bank vole abundance and the proportion of

wooded habitats in the trapping habitats, both displaying a significant positive relationship (**Figure 1c**

and **1d**). The ratio between bank vole and wood mouse abundances was significantly positively related

to the prevalence in wood mice in the univariate model (not shown). The most supported GLM of

prevalence in bank voles included the bank vole abundance, the proportion of wooded habitats in

50 m-radius zones, and the proportion of ecotones in 200 m-radius zones, all displaying a significant

positive relationships with the prevalence (**Figure 1e**, **1f**, and **1g**, respectively).The relationship

between prevalence and ecotones was negative in single explanatory variable models considering all

sites, but positive when excluding forest sites (not shown). Thus, the apparent contradiction between

single and multiple explanatory variables models likely results from the reduction of ecotones length

in highly wooded landscapes (see the prevalence as a function of both landscape variables in

Figure 1h).

Results for B. burgdorferi *s.l.*

The analyses of *B. burgdorferi* s.l. prevalence were based on the PCR results for 450 wood mice, 147

bank voles, four field voles, two common pine voles and three crowned shrews. Fifteen wood mice,

ten bank voles and one common pine vole were positive (**Table 2**). The numbers of positive wood

mice and bank voles per site were not significantly correlated neither was the prevalence ($p > 0.05$).

The results of the selection procedure of the GLMs of *B. burgdorferi* s.l. prevalence as a function of

landscape variables and species abundance variables are detailed in **Table 4**.

For all small mammals, the only significant variable in GLMs of *B. burgdorferi* s.l. prevalence

included the proportion of ecotones in 50 m-radius zones, displaying a significant positive relationship

(**Figure 2a**). For prevalence in wood mice separately, no landscape variable displayed a significant

relationship. The most supported GLM of prevalence in bank voles included only the proportion of

ecotones in 100 m-radius zones, displaying a significant positive relationship (**Figure 2b**). This

variable computed in the trapping habitats was also significant. The proportion of wooded habitats in

the trapping habitats and in 50 m-radius zones displayed a significant negative relationship (not

shown). No species abundances variable was significantly related to any prevalence.

Discussion

In this study, we investigated the possible associations between landscape structure, abundances of

small mammal species and the prevalence of two tick-borne infectious agents in small mammals, to

better understand the spread of these infectious agents in the landscape. The small mammal

community at the woodland/hedgerows-grassland ecotones and in forest sites was dominated by the

wood mouse and the bank vole, two rodent species with different habitats use, and as such interesting

to compare at this landscape scale. However, we found only that the wood mouse abundance was

375 positively correlated to the proportion of ecotones in 500 m-radius zones. This finding is consistent

376 with the propensity of this generalist species to use various complementary habitats (Tew and

377 Macdonald 1994; Ouin et al. 2000). Despite bank vole abundance was not associated to any landscape

378 variable here, in a previous study considering seasonal variation, negative associations with two

379 landscape structural connectivity variables was observed in spring (Perez et al. 2016). Such

380 relationships, also reported in several other studies, are likely caused by a released predation pressure

381 in fragmented landscapes (Szacki 1987; van Apeldoorn et al. 1992; Paillat and Butet 1996; Michel et

382 al. 2007).

384 *Prevalence of* Anaplasma phagocytophilum

385 The numbers of positive individuals as well as the prevalence in wood mice and bank voles were

386 correlated, suggesting a common infection factor and/or an increased exposition of one species in the

387 presence of the other. We observed a higher prevalence in bank voles and a positive, despite weak,

388 relationship between bank vole abundance and the prevalence in all small mammals and in species

389 separately. Although not related to landscape variables, these results support an effect of relative host

390 species abundances on prevalence (**H2**): bank vole likely acts as an amplification host species for the

391 transmission of these bacteria, while wood mouse acts, comparatively, rather as a dilution host species

392 (Rosso et al. 2017; Perez et al. 2017).

393 Several studies demonstrated that the prevalence of directly transmitted infectious agents increased

394 with host population size, and decreased with population isolation as a consequence of a higher local

395 extinction risk (Begon et al. 2003; Guivier et al. 2011). Here, the proportion of wooded habitats in the

396 surrounding landscape was positively associated to *A. phagocytophilum* prevalence in all small

397 mammals and in species separately. More wooded landscapes probably support larger and more

398 connected populations of bank voles resulting in a lower local extinction probability in a given habitat

399 patch (Paillat and Butet 1996). Consequently, a lower local extinction probability of their parasites

400 (including small mammal specialist tick species like *I. trianguliceps* and *I. acuminatus*), and of the

401 infectious agents they could transmit is expected (**H1**).

403 *Prevalence of* Borrelia burgdorferi *sensu lato*

404 The *B. burgdorferi* s.l. prevalence in small mammals, particularly in bank voles, was positively related

405 to the proportion of ecotones within 100 m. These results suggest an enhanced transmission of these

406 bacteria at the interface between wooded and open (crops and grassland) habitats and therefore in

407 fragmented landscapes where such ecotones are frequent. A possible explanation of increased

408 *B. burgdorferi* s.l. prevalence in such landscapes is an increased density of small mammals and other

409 hosts, causing an increased density of *I. ricinus* nymphs, resulting in an amplification of the

transmission cycle (Agoulon et al. 2012; Li et al. 2012; Cayol et al. 2018). However, in previous

studies, no relationship was observed between the abundance of *I. ricinus* nymphs and

B. burgdorferi s.l. prevalence, and tick density was generally higher in woodlands (Boyard et al. 2008;

Hoch et al. 2010; Perez et al. 2017). An alternative explanation is an increased density of hosts

competent for the same geno-species than small mammals (Vourc'h et al. 2008).

The mobility of the wood mouse and its ability to use crops and grassland could explain the absence of

spatial pattern in the *B. burgdorferi* s.l. prevalence of this species. Several studies showed that wood

mice, more prone to host *Ixodes* sp. larvae than bank voles, may yield more nymphs infected by those

bacteria despite displaying lower bacterial loads (Humair et al. 1993, 1999). Thus, wood mice might

act as medium distance dispersers (at least up to 500 m) of infected *Ixodes* sp. larvae that become

infective nymphs, as previously suggested in other studies (Boyard et al. 2008; Gassner et al. 2008).

These results support the role of host specificity of infectious agents on their distribution patterns

(**H3**).

For both infections, prevalence models were not improved by using wooded habitats connectivity

measures compared to simpler landscape variables. In our study area, variations in wooded habitats

connectivity were maybe not contrasted enough to be captured (Michel et al. 2007). Alternatively,

within each considered land cover, the influence of habitat quality and heterogeneity on landscape

connectivity for small mammals might have been overlooked and should be accounted for in higher

resolution models and/or habitat quality based landscape connectivity models (Mortelliti et al. 2010).

It would also be interesting to account for other potential factors affecting small mammal communities

and their tick-borne infection prevalence, like predation pressure (Hofmeester et al. 2017).

Finally, species which were rare in the sampled habitats (crowned shrew, field vole and common pine

vole) can also host *A. phagocytophilum* and/or *B. burgdorferi* s.l. (Bown et al. 2008, 2009, 2011).

Further studies including habitats where these species can be more abundant (e.g. fallow lands, fields'

margins, clear cuts) and focusing on more hosts species, especially for *B. burgdorferi* s.l., would allow

a better overall view of the distribution and the spread of these infectious agents in the landscape.

Conclusion

The *A. phagocytophilum* prevalence could be substantially explained by a positive association with the

proportion of wooded habitats in the surrounding landscape at least up to 500 m, validating the host

population size hypothesis (**H1**). The *B. burgdorferi* s.l. prevalence could be explained only by the

presence of ecotones within 100 m. Despite a weak positive relationship between bank vole abundance

and *A. phagocytophilum* prevalence, no indirect effect of landscape variables on this prevalence has

been detected, questioning the overall host community competence hypothesis (**H2**). Indeed, only

wood mice abundance was significantly linked to our landscape variables, but at larger scales than the

relationships between landscape variables and the prevalence. The prevalence of the two infections

displayed contrasted spatial patterns, a difference which likely results from the wider range of hosts

and vectors of *B. burgdorferi* s.l., as predicted by the infectious agent specificity hypothesis (**H3**). As a

whole, our results demonstrate the interest of integrating complementary approaches, such as

landscape ecology and community ecology, to better understand the dynamics and the spatial

distribution of tick-borne infectious agents to prevent epidemiological risks at the scale of agricultural

landscapes.

Competing interests

The authors declare no competing interests

Authors' contributions

GP, SB, AA, GV, OP and AB designed the study. GP, SB, AA, YR, OP and AB participated to the

small mammal field sampling. AC performed most of the DNA extractions and the molecular

analyses. YR managed the GIS data. GP performed all data analyses and drafted the manuscript. All

authors read, commented and approved the manuscript.

Acknowledgement

We are very grateful to Agnès Bouju, Floriane Boullot, Axelle Durand, Mathieu Gonnet, Olivier

Jambon, Maggy Jouglin, Emmanuelle Moreau, Pranav Pandit, and Ionut Pavel who helped in

sampling and preparing small mammal tissues before molecular analyses; to Séverine Barry, Amélie

Cohadon, Angélique Pion, and Valérie Poux who helped in the lab for the molecular detection of

infectious agents; and to Nelly Dorr and Isabelle Lebert who managed the data base of the OSCAR

project (https://www6.inra.fr/oscar/). We thank the 'Zone Atelier Armorique' (https://osur.univ-

rennes1.fr/za-armorique/) for providing the GIS data and for the access to its field facilities. We thank

Henri Lemercier and Benoît Chevallier from the 'Office National des Forêts' for facilitating the access

to the Villecartier forest. The 'Tiques et Maladies à Tiques' team of the 'Réseau Ecologie des

Interactions Durables' group, supported by the INRA and the CNRS gave a rich thinking environment.

This work was funded by the French National Research Agency (ANR-11-Agro-001-04; call for

Proposal 'Agrobiosphere', OSCAR project). This work is part of the PhD of GP, which was supported

by a fellowship from the Brittany region, France. The funders had no role in study design, data

collection and interpretation, or the decision to submit the work for publication.

References

Agoulon A, Malandrin L, Lepigeon F, et al (2012) A vegetation index qualifying pasture edges is
related to *Ixodes ricinus* density and to *Babesia divergens* seroprevalence in dairy cattle herds.
Vet Parasitol 185:101–109.

Al Hassan D, Georgelin E, Delattre T, et al (2012) Does the presence of grassy strips and landscape
grain affect the spatial distribution of aphids and their carabid predators? Agric For Entomol
15:24–33.

Allan BF, Keesing F, Ostfeld RS (2003) Effect of forest fragmentation on Lyme disease risk. Conserv
Biol 17:267–272.

Araya-Anchetta A, Busch JD, Scoles GA, Wagner DM (2015) Thirty years of tick population
genetics: A comprehensive review. Infect Genet Evol 29:164–179.

Begon M, Hazel SM, Telfer S, et al (2003) Rodents, cowpox virus and islands: densities, numbers and
thresholds. J Anim Ecol 72:343–355.

Blaňarová L, Stanko M, Carpi G, et al (2014) Distinct *Anaplasma phagocytophilum* genotypes
associated with *Ixodes trianguliceps* ticks and rodents in Central Europe. Ticks Tick Borne Dis
5:928–938.

Boussard H, Baudry J (2014) Chloe2012: a software for landscape pattern analysis
(http://www.rennes.inra.fr/sad/Outils-Produits/Outils-informatiques/Chloe). 41 p.

Bown KJ, Begon M, Bennett M, et al (2003) Seasonal dynamics of *Anaplasma phagocytophila* in a
rodent-tick (*Ixodes trianguliceps*) system, United Kingdom. Emerg Infect Dis 9:63–70.

Bown KJ, Lambin X, Ogden NH, et al (2009) Delineating *Anaplasma phagocytophilum* ecotypes in
coexisting, discrete enzootic cycles. Emerg Infect Dis 15:1948–1954.

Bown KJ, Lambin X, Telford G, et al (2011) The common shrew (*Sorex araneus*): a neglected host of
tick-borne infections? Vector-borne Zoonotic Dis 11:947–953.

Bown KJ, Lambin X, Telford GR, et al (2008) Relative importance of *Ixodes ricinus* and *Ixodes
trianguliceps* as vectors for *Anaplasma phagocytophilum* and *Babesia microti* in field vole
(*Microtus agrestis*) populations. Appl Environ Microbiol 74:7118–7125.

Boyard C, Vourc'h G, Barnouin J (2008) The relationships between *Ixodes ricinus* and small mammal
species at the woodland-pasture interface. Exp Appl Acarol 44:61–76.

Cayol C, Jääskeläinen A, Koskela E, et al (2018) Sympatric *Ixodes*-tick species: pattern of distribution
and pathogen transmission within wild rodent populations. Sci Rep 8:16660.

Chastagner A, Moinet M, Perez G, et al (2016) Prevalence of *Anaplasma phagocytophilum* in small
rodents in France. Ticks Tick Borne Dis 7:988–991.

Clay CA, Lehmer EM, Jeor SS, Dearing MD (2009) Testing mechanisms of the dilution effect: Deer
      mice encounter rates, sin nombre virus prevalence and species diversity. Ecohealth 6:250–259.

Close B, Banister K, Baumans V, et al (1997) Recommendations for euthanasia of experimental
      animals: Part 2. Lab Anim 31:1–32.

Coipan EC, Sprong H (2016) Ecology of *Borrelia burgdorferi* sensu lato. In: Braks MAH, van Wieren
      SE, Takken W, Sprong H (eds) Ecology and prevention of Lyme borreliosis. Wageningen
      Academic Publishers, Wageningen, pp 41–61

Courtney JW, Kostelnik LM, Zeidner NS, Massung RF (2004) Multiplex real-time PCR for detection
      of *Anaplasma phagocytophilum* and *Borrelia burgdorferi*. J Clin Microbiol 42:3164–3168.

Eskildsen A (2010) Effects of resource abundance on habitat selection and spatial behavior of the bank
      vole (*Myodes glareolus*). Master's Thesis, University of Copenhagen

Ferrante M, González E, Lövei GL (2017) Predators do not spill over from forest fragments to maize
      fields in a landscape mosaic in central Argentina. Ecol Evol 7:1–9.

Foltête JC, Clauzel C, Vuidel G (2012) A software tool dedicated to the modelling of landscape
      networks. Environ Model Softw 38:316–327.

Gassner F, Verbaarschot P, Smallegange RC, et al (2008) Variations in *Ixodes ricinus* density and
      Borrelia infections associated with cattle introduced into a woodland in The Netherlands. Appl
      Environ Microbiol 74:7138–44.

Gehring TM, Swihart RK (2003) Body size, niche breadth, and ecologically scaled responses to
      habitat fragmentation: mammalian predators in an agricultural landscape. Biol Conserv 109:283–
      295.

Gern L, Estrada-Peña A, Frandsen F, et al (1998) European reservoir hosts of *Borrelia burgdorferi*
      sensu lato. Zentralbl Bakteriol 287:196–204.

Gern L, Siegenthaler M, Hu CM, et al (1994) *Borrelia burgdorferi* in rodents (*Apodemus flavicollis*
      and *A. sylvaticus*): Duration and enhancement of infectivity for *Ixodes ricinus* ticks. Eur J
      Epidemiol 10:75–80.

Gil-Tena A, Nabucet J, Mony C, et al (2014) Woodland bird response to landscape connectivity in an
      agriculture-dominated landscape: a functional community approach. Community Ecol 15:256–
      268.

Gottdenker NL, Chaves LF, Calzada JE, et al (2012) Host life history strategy, species diversity, and
      habitat influence *Trypanosoma cruzi* vector infection in changing landscapes. PLoS Negl Trop
      Dis 6:e1884, 11p.

Guivier E, Galan M, Chaval Y, et al (2011) Landscape genetics highlights the role of bank vole

metapopulation dynamics in the epidemiology of Puumala hantavirus. Mol Ecol 20:3569–3583.

Halos L, Bord S, Cotté V, et al (2010) Ecological factors characterizing the prevalence of bacterial
tick-borne pathogens in *Ixodes ricinus* ticks in pastures and woodlands. Appl Environ Microbiol
76:4413–4420.

Hersh MH, Tibbetts M, Strauss M, et al (2012) Reservoir competence of wildlife host species for
Babesia microti. Emerg Infect Dis 18:1951–1957.

Heylen D, Tijsse E, Fonville M, et al (2013) Transmission dynamics of *Borrelia burgdorferi* s.l. in a
bird tick community. Environ Microbiol 15:663–673.

Hoch T, Monnet Y, Agoulon A (2010) Influence of host migration between woodland and pasture on
the population dynamics of the tick *Ixodes ricinus*: a modelling approach. Ecol Modell
221:1798–1806.

Hofmeester TR, Coipan EC, Van Wieren SE, et al (2016) Few vertebrate species dominate the
Borrelia burgdorferi s.l. life cycle. Environ Res Lett 11:043001, 16p.

Hofmeester TR, Jansen PA, Wijnen HJ, et al (2017) Cascading effects of predator activity on tick-
borne disease risk. Proc R Soc B Biol Sci 284:20170453.

Huang ZYX, de Boer WF, van Langevelde F, et al (2013) Species' life-history traits explain
interspecific variation in reservoir competence: a possible mechanism underlying the dilution
effect. PLoS One 8:e54341, 6p.

Hubbard MJ, Baker AS, Cann KJ (1998) Distribution of *Borrelia burgdorferi* s.l. spirochaete DNA in
British ticks (Argasidae and Ixodidae) since the 19th century, assessed by PCR. Med Vet
Entomol 12:89–97.

Humair PF, Rais O, Gern L (1999) Transmission of *Borrelia afzelii* from *Apodemus* mice and
Clethrionomys voles to *Ixodes ricinus* ticks: differential transmission pattern and overwintering
maintenance. Parasitology 118:33–42.

Humair PF, Turrian N, Aeschlimann A, Gern L (1993) *Borrelia burgdorferi* in a focus of Lyme
borreliosis: epizootiologic contribution of small mammals. Folia Parasitol (Praha) 40:65–70.

Hurvich CM, Tsai C-L (1989) Regression and time series model selection in small samples.
Biometrika 76:297–307.

Jahfari S, Coipan EC, Fonville M, et al (2014) Circulation of four *Anaplasma phagocytophilum*
ecotypes in Europe. Parasites and Vectors 7:365.

Johnson PTJ, Preston DL, Hoverman JT, Richgels KLD (2013) Biodiversity decreases disease through
predictable changes in host community competence. Nature 494:230–233.

Jones KE, Patel NG, Levy MA, et al (2008) Global trends in emerging infectious diseases. Nature

451:990–993.

Kallio ER, Begon M, Birtles RJ, et al (2014) First Report of *Anaplasma phagocytophilum* and *Babesia*
microti in Rodents in Finland. Vector-borne Zoonotic Dis 14:389–393.

Keesing F, Brunner J, Duerr S, et al (2009) Hosts as ecological traps for the vector of Lyme disease.
Proc R Soc B Biol Sci 276:3911–3919.

Kikkawa T (1964) Movement, activity and distribution of the small rodents *Clethrionomys glareolus*
and *Apodemus sylvaticus* in Woodland. J Anim Ecol 33:259–299.

Kurtenbach K, De Michelis S, Etti S, et al (2002a) Host association of *Borrelia burgdorferi* sensu
lato–the key role of host complement. Trends Microbiol 10:74–79.

Kurtenbach K, De Michelis S, Sewell HS, et al (2002b) The key roles of selection and migration in the
ecology of Lyme borreliosis. Int J Med Microbiol 291:152–154.

Kurtenbach K, Hanincová K, Tsao JI, et al (2006) Fundamental processes in the evolutionary ecology
of Lyme borreliosis. Nat Rev Microbiol 4:660–669.

Lambin EF, Tran A, Vanwambeke SO, et al (2010) Pathogenic landscapes: interactions between land,
people, disease vectors, and their animal hosts. Int J Health Geogr 9:54, 13p.

Li S, Hartemink N, Speybroeck N, Vanwambeke SO (2012) Consequences of landscape fragmentation
on Lyme disease risk: a cellular automata approach. PLoS One 7:e39612.

Mader H (1984) Animal habitat isolation by roads and agricultural fields. Biol Conserv 29:81–96.

Mansfield KL, Johnson N, Phipps LP, et al (2009) Tick-borne Encephalitis Virus - A Review of an
Emerging Zoonosis. J Gen Virol 90:1781–94.

Margos G, Vollmer SA, Cornet M, et al (2009) A new *Borrelia* species defined by multilocus
sequence analysis of housekeeping genes. Appl Environ Microbiol 75:5410–5416.

Marsot M, Chapuis JL, Gasqui P, et al (2013) Introduced Siberian Chipmunks (*Tamias sibiricus*
barberi) Contribute More to Lyme Borreliosis Risk than Native Reservoir Rodents. PLoS One
8:1–8.

Meunier FD, Corbin J, Verheyden C, Jouventin P (1999) Effects of landscape type and extensive
management on use of motorway roadsides by small mammals. Can J Zool 77:108–117.

Michel N, Burel F, Legendre P, Butet A (2007) Role of habitat and landscape in structuring small
mammal assemblages in hedgerow networks of contrasted farming landscapes in Brittany,
France. Landsc Ecol 22:1241–1253.

Millán de la Peña N, Butet A, Delettre Y, et al (2003) Response of the small mammal community to
changes in western French agricultural landscapes. Landsc Ecol 18:265–278.

Mortelliti A, Amori G, Boitani L (2010) The role of habitat quality in fragmented landscapes: a
conceptual overview and prospectus for future research. Oecologia 163:535–547.

Nupp TE, Swihart RK (1998) Effects of forest fragmentation on populations attributes of white-footed
mice and eastern chipmunks. J Mammal 79:1234–1243.

Nupp TE, Swihart RK (2000) Landscape-level correlates of small-mammal assemblages in forest
fragments of farmland. J Mammal 81:512–526.

Ostfeld RS, Keesing F (2000) Biodiversity and disease risk: the case of Lyme disease. Conserv Biol
14:722–728.

Ostfeld RS, Levi T, Jolles AE, et al (2014) Life history and demographic drivers of reservoir
competence for three tick-borne zoonotic pathogens. PLoS One 9:e107387, 8p.

Ostfeld RS, LoGiudice K (2003) Community disassembly, biodiversity loss, and the erosion of an
ecosystem service. Ecology 84:1421–1427.

Ouin A, Paillat G, Butet A, Burel F (2000) Spatial dynamics of wood mouse (*Apodemus sylvaticus*) in
an agricultural landscape under intensive use in the Mont Saint Michel Bay (France). Agric
Ecosyst Environ 78:159–165.

Paillat G, Butet A (1996) Spatial dynamics of the bank vole (Clethrionomys glareolus) in a fragmented
landscape. Acta Œcologica 17:553–559.

Papillon Y, Buffiere L, Butet A (2002) Rhodamine B as collective marker for small mammals. Acta
Theriol (Warsz) 47:491–497.

Perez G, Bastian S, Agoulon A, et al (2016) Effect of landscape features on the relationship between
Ixodes ricinus ticks and their small mammal hosts. Parasit Vectors 9:20.

Perez G, Bastian S, Chastagner A, et al (2017) Ecological factors influencing small mammal infection
by *Anaplasma phagocytophilum* and *Borrelia burgdorferi* s.l. in agricultural and forest
landscapes. Environ Microbiol 19:4205–4219.

Pisanu B, Chapuis JL, Dozières A, et al (2014) High prevalence of *Borrelia burgdorferi* s.l. in the
European red squirrel *Sciurus vulgaris* in France. Ticks Tick Borne Dis 5:1–6.

R Development Core Team (2019) R v.3.6.1. R Foundation for Statistical Computing.

Randolph SE (1975a) Seasonal dynamics of a host-parasite system: *Ixodes trianguliceps* (Acarina:
Ixodidae) and its small mammal hosts. J Anim Ecol 44:425–449.

Randolph SE (1975b) Patterns of Distribution of the Tick *Ixodes trianguliceps* Birula on Its Hosts. J
Anim Ecol 44:451–474.

Reidsma P, Tekelenburg T, vand den Berg M, Alkemade R (2006) Impacts of land-use change on
biodiversity: An assessment of agricultural biodiversity in the European Union. Agric Ecosyst

Environ 114:86–102.

Rico A, Kindlmann P, Sedláček F (2007) Barrier effects of roads on movements of small mammals.
Folia Zool 56:1–12.

Rizzoli A, Hauffe HC, Carpi G, et al (2011) Lyme borreliosis in Europe. Eurosurveillance 16:1–8.

Roche B, Rohani P, Dobson AP, Guégan J-F (2013) The impact of community organization on vector-
borne pathogens. Am Nat 181:1–11.

Rollend L, Fish D, Childs JE (2013) Transovarial transmission of *Borrelia* spirochetes by *Ixodes*
scapularis: A summary of the literature and recent observations. Ticks Tick Borne Dis 4:46–51.

Rosso F, Tagliapietra V, Baráková I, et al (2017) Prevalence and genetic variability of *Anaplasma*
phagocytophilum in wild rodents from the Italian alps. Parasit Vectors 10:293.

Rubio A, Ávila-Flores R, Suzán G (2014) Responses of small mammals to habitat fragmentation:
epidemiological considerations for rodent-borne hantaviruses in the Americas. Ecohealth 526–
533.

Ruiz-Capillas P, Mata C, Malo JE (2015) How many rodents die on the road? Biological and
methodological implications from a small mammals' roadkill assessment on a Spanish
motorway. Ecol Res 30:417–427.

Saura S, Pascual-Hortal L (2007) A new habitat availability index to integrate connectivity in
landscape conservation planning: Comparison with existing indices and application to a case
study. Landsc Urban Plan 83:91–103.

Sinski E, Pawelczyk A, Bajer A, Behnke JM (2006) Abundance of wild rodents, ticks and
environmental risk of Lyme borreliosis: A longitudinal study in an area of Mazury Lakes district
of Poland. Ann Agric Environ Med 13:295–300.

Skuballa J, Petney T, Pfäffle M, et al (2012) Occurrence of different *Borrelia burgdorferi* sensu lato
genospecies including *B. afzelii*, *B. bavariensis*, and *B. spielmanii* in hedgehogs (*Erinaceus* spp.)
in Europe. Ticks Tick Borne Dis 3:8–13.

Stuen S, Granquist EG, Silaghi C (2013) *Anaplasma phagocytophilum*-a widespread multi-host
pathogen with highly adaptive strategies. Front. Cell. Infect. Microbiol. 4:1–33.

Suzán G, García-Peña GE, Castro-Arellano I, et al (2015) Metacommunity and phylogenetic structure
determine wildlife and zoonotic infectious disease patterns in time and space. Ecol Evol 5:865–
873.

Szacki J (1987) Ecological Corridor as a Factor Determining the Structure and Organization of a Bank
Vole Population. Acta Theriol (Warsz) 32:31–44.

Szacki J, Babińska-werka J, Liro A (1993) The influence of landscape spatial structure on small

mammal movements. Acta Theriol (Warsz) 38:113–123.

Szekeres S, Coipan EC, Rigó K, et al (2015) Eco-epidemiology of *Borrelia miyamotoi* and Lyme

borreliosis spirochetes in a popular hunting and recreational forest area in Hungary. Parasites and

Vectors. doi: 10.1186/s13071-015-0922-2

Tattersall FH, Macdonald DW, Hart BJ, et al (2002) Is habitat linearity important for small mammal

communities on farmland? J Appl Ecol 39:643–652.

Tew TE, Macdonald DW (1994) Dynamics of space use and male vigour amongst wood mice,

Apodemus sylvaticus, in the cereal ecosystem. Behav Ecol Sociobiol 34:337–345.

van Apeldoorn RC, Oostenbrink WT, van Winden A, van der Zee FF (1992) Effects of fragmentation

on the bank vole, *Clethrionomys glareolus*, in an agricultural landscape. Oikos 65:265–274.

Vourc'h G, Boyard C, Barnouin J (2008) Mammal and bird species distribution at the woodland-

pasture interface in relation to the circulation of ticks and pathogens. Ann N Y Acad Sci

1149:322–325.

Zhang Z, Usher MB (1991) Dispersal of wood mice and bank voles in an agricultural landscape. Acta

Theriol (Warsz) 36:239–245.

Tables and figure captions

Table 1: Landscape variables used in the study for the model selection procedures.

Variable	Scale (m)[a]	Unit	Mean	Median	Range (min; max)
Proportion of wooded habitats					
Wooded.local	0	Proportion of pixels	0.569	0.557	0.051;1.000
Wooded.50m	50	Proportion of pixels	0.511	0.495	0.053; 0.986
Wooded.100m	100	Proportion of pixels	0.487	0.441	0.039; 0.988
Wooded.200m	200	Proportion of pixels	0.453	0.352	0.042; 0.990
Wooded.500m	500	Proportion of pixels	0.420	0.3	0.034; 0.970
Proportion of ecotones					
Ecotones.local	0	Proportion of pixel couples	0.058	0.051	0.000; 0.132
Ecotones.50m	50	Proportion of pixel couples	0.048	0.042	0.021; 0.090
Ecotones.100m	100	Proportion of pixel couples	0.042	0.041	0.017; 0.075
Ecotones.200m	200	Proportion of pixel couples	0.034	0.030	0.014; 0.057
Ecotones.500m	500	Proportion of pixel couples	0.027	0.027	0.012; 0.047
Connectivity measures					
log(Area)	Patch	m²	11.15	11.14	6.48; 14.57
log(dPC.Euclidian.25m)	25	none	6.17	5.36	-2.00; 12.73
log(dPC.Euclidian.50m)	50	none	6.26	5.16	-2.08; 12.73
log(dPC.Euclidian.100m)	100	none	6.35	5.08	-2.12; 12.71
log(dPC.Euclidian.250m)	250	none	6.62	5.42	-1.82; 12.72
log(dPC.lcp.As.100m)[b]	100	none	6.52	5.61	-2.01; 12.79
log(dPC.lcp.Mg.25m)[c]	25	none	5.99	5.84	-3.20; 12.76
log(dPCflux.Euclidian.25m)	25	none	5.09	3.61	-3.09; 12.26
log(dPCflux.Euclidian.50m)	50	none	5.51	3.96	-2.63; 12.43
log(dPCflux.Euclidian.100m)	100	none	5.77	4.12	-2.57; 12.49
log(dPCflux.Euclidian.250m)	250	none	6.22	4.77	-2.05; 12.49
log(dPCflux.lcp.As.100m)[b]	100	none	5.63	4.17	-2.81; 12.29
log(dPCflux.lcp.Mg.25m)[c]	25	none	2.03	1.36	-10.78;10.21
log(dPCconnector.Euclidian.25m+1)	25	none	0.62	0.0000	0.000; 3.89
log(dPCconnector.Euclidian.50m+1)	50	none	1.56	0.0000	0.000; 7.17
log(dPCconnector.Euclidian.100m+1)	100	none	2.08	0.0121	0.000; 9.22
log(dPCconnector.Euclidian.250m+1)	250	none	2.78	0.2520	0.000; 10.35
log(dPCconnector.lcp.As.100m+1)[b]	100	none	1.04	0.0801	0.000; 7.98
log(dPCconnector.lcp.Mg.25m+1)[c]	25	none	5.79^{e-3}	0.0000	0.000; 0.109

[a] Radius of computing zones (proportions) or median movement distances (connectivity).

[b] As: with the friction coefficients set for *Apodemus sylvaticus*.

[c] Mg: with the friction coefficients set for *Myodes glareolus*.

Table 2: Results of *Anaplasma phagocytophilum* and *Borrelia burgdorferi* s.l. PCR detection per site over the two-year sampling (2012-2013).

Site	Landscape context[a]	Asyl Nbr[c]	Aph	Bb sl	Mgla Nbr[c]	Aph	Bb sl	Magr Nbr[c]	Aph	Bb sl	Msub Nbr[c]	Aph	Bb sl	Scor Nbr[c]	Aph	Bb sl	Total Nbr[c]	Aph	Bb sl
1	FC	11	1	0	2	0	0										13	1	0
2	FC	9	0	0	10	1	0										19	1	0
3	FC	9	0	0	2	1	0										11	1	0
4	FC	19	2	1	11	9	0	2	0	0				1	1	0	33	12	1
5	FC	19	2	1	10	2	1	1	0	0							30	4	2
6	FC	19[d]	1	1	1	0	0										20	1	1
7	FE	18	2	0	13	1	0	1	0	0							32	3	0
8	FE	32	2	2	3	1	0										35	3	2
9	FE	20	0	0	11	1	1										31	1	1
10	FE	18	1	1	1	0	0				1	0	0				20	1	1
11	FE	2	0	0	1[d]												3	0	0
12	FE	8[d]	0	0	2	0	0										10	0	0
13	ME	21	2	0	6[d]	0	0										27	2	0
14	ME	32	3	2	12	4	1										44	7	3
15	ME	24	0	0	7	0	1										31	0	1
16	MH	29	2	1	13	0	2										42	2	3
17	MH	21	0	0	6	0	1										27	0	1
18	MH	40[e]	0	1	4	0	0				1	0	1				45	0	2
19	OE	32	0	1	4	0	0										36	0	1
20	OE	9	0	0	5	0	0										14	0	0
21	OE	17	0	2	4	0	1										21	0	3
22	OH	16	0	1	6	0	0							1	0	0	23	0	1
23	OH	23	0	1	8	0	2										31	0	3
24	OH	6	2	0	7	0	0							1	1	0	14	3	0
Total	**24**	454	20	15	149	20	10	4	0	0	2	0	1	3	2	0	612	42	26
% of total		74.2	3.28	2.47	24.3	3.28	1.64	0.65	0	0	0.33	0	0.16	0.49	0.33	0	100	6.91	4.29
Prevalence (%)			4.42	3.33		13.6	6.80		0	0		0	50.0		66.7	0		6.91	4.29

[a] FC: forest core; FE: forest edge; ME: mixed farming at woodlot edge; MH: mixed farming along hedgerows; OE: open farming at woodlot edge; and OH: open farming along hedgerows.

[b] Asyl: *Apodemus sylvaticus*; Mgla: *Myodes glareolus*; Magr: *Microtus agrestis*; Msub: *M. subterraneus*; and Scor: *Sorex coronatus*.

[c] Total number of individuals over the two-year sampling.

[d] Including one individual not tested.

[e] Including two individuals not tested for *B. burgdorferi* s.l. only.

711
712

Table 3: Best binomial Generalized Linear Models of *Anaplasma phagocytophilum* prevalence as a function of landscape variables and species abundance variables according to the AICc-based selection procedure.

All small mammals					
Model set	Rank	Variables and their significativity	AICc	ΔAICc	R²
Null model			106.9		
Landscape	1	**Wooded.50m*****	86.5		0.332
Community	1	**N.Mg*****	96.3		0.189
	2	**N.Mg*** + Ratio.Mg:As	96.6	0.33	0.223
Community and landscape	1	**Wooded.50m*** + N.Mg*****	77.7		0.498
	2	**Wooded.local*** + N.Mg****	78.4	0.68	0.488

Wood mice (*Apodemus sylvaticus*)					
Model set	Rank	Variables and their significativity	AICc	ΔAICc	R²
Null model			61.5		
Landscape	1	**Wooded.50m***	59.0		0.145
	2	**Wooded.local***	59.1	0.05	0.143
	3	Wooded.100m°	60.4	1.32	0.105
	4	Ecotones.100m°	60.4	1.32	0.105
	5	log(Area)°	60.6	1.59	0.097
Community	1	**Ratio.Mg:As***	59.0		0.146
	2	N.Mg°	60.4	1.42	0.101
Community and landscape	1	**Wooded.local* + N.Mg***	57.8		0.261
	2	**Wooded.50m*** + N.Mg°	58.2	0.40	0.250
	3	**Wooded.local*** + Ratio.Mg:As°	58.3	0.41	0.248
	4	Wooded.50m° + Ratio.Mg:As°	58.5	0.06	0.243
	5	log(Area) + **Ratio.Mg:As***	59.5	1.07	0.225

Bank voles (*Myodes glareolus*)					
Model set	Rank	Variables and their significativity	AICc	ΔAICc	R²
Null model			73.0		
Landscape	1	**Ecotones.200m* + Wooded.50m*****	52.1		0.477
	2	Ecotones.200m° + **Wooded.100m*****	52.9	0.87	0.461
	3	**Ecotones.200m* + Wooded.200m*****	52.9	0.89	0.461
	4	**Ecotones.local*****	53.4	1.29	0.405
	5	**Wooded.100m*****	54.0	1.94	0.393
Community	1	**N.Mg***	70.2		0.096
Community and landscape	1	**Ecotones.200m** + Wooded.50m*** + N.Mg***	48.3		0.600

Only models with ΔAICc < 2 are shown. Models with connectivity variables that performed worse than models with the wooded patch area have been discarded. Significant codes are: "***" when $p \leq 0.001$; "**" when $0.001 < p \leq 0.01$; "*" when $0.01 < p \leq 0.05$; "°" when $0.05 < p \leq 0.1$; otherwise, $0.1 < p$. Significant variables at $p < 0.05$ are in bold.

716
717

Table 4: Best binomial Generalized Linear Models of *Borrelia burgdorferi* s.l. prevalence in small mammals as a function of landscape variables and species abundance variables according to the AICc-based selection procedure.

Model set	Rank	Variables and their significativity	AICc	ΔAICc	R²
		All small mammals			
Null model			56.7		
Landscape	1	**Ecotones.50m***	55.2	0.00	0.201
	2	Ecotones.100m°	55.4	0.25	0.188
Community	None				

Model set	Rank	Variables and their significativity	AICc	ΔAICc	R²
		Wood mice (*Apodemus sylvaticus*)			
Null model			43.0		
Landscape	None				
Community	None				

Model set	Rank	Variables and their significativity	AICc	ΔAICc	R²
		Bank voles (*Myodes glareolus*)			
Null model			35.2		
Landscape	1	**Ecotones.100m***	33.3	0.00	0.252
	2	**Ecotones.local***	33.6	0.30	0.235
	3	**Woodes.local***	33.6	0.31	0.234
	4	**Wooded.50m***	33.7	0.40	0.229
	5	dPC.250m°	34.0	0.65	0.214
	6	Ecotones.50m°	34.0	0.67	0.213
	7	dPC.25m°	34.0	0.68	0.212
	8	Wooded.100m°	34.1	0.78	0.206
	9	dPC.50m°	34.1	0.79	0.206
	10	dPC.100m°	34.1	0.81	0.205
	11	dPC.250m + Ecotones.100m	34.2	0.85	0.350
	12	dPCflux.250m°	34.2	0.86	0.202
	13	dPC.Mg°	34.3	0.94	0.197
	14	dPCconnector.250m°	34.3	0.97	0.196
	15	Ecotones.100m + Wooded.50m	34.3	1.01	0.347
	16	log(Area)°	34.3	1.02	0.193
Community	None				

Only models with ΔAICc < 2 are shown. Models with connectivity variables that performed worse than models with the wooded patch area have been discarded. Significant codes are: "***" when $p \leq 0.001$; "**" when $0.001 < p \leq 0.01$; "*" when $0.01 < p \leq 0.05$; "°" when $0.05 < p \leq$; otherwise, $0.1 < p$. Significant variables at $p < 0.05$ are in bold.

720
721

Figure 1: The *Anaplasma phagocytophilum* prevalence in small mammals as a function of landscape variables and species abundance variables.

The *A. phagocytophilum* prevalence, expressed as the frequency of infected individuals per site, in all small mammals (**a** and **b**), in wood mice (**c** and **d**), and in bank voles (**e, f, g** and **h**). Fitted binomial regression curves, according to the best multiple explanatory variables binomial Generalized Linear Models, are shown with 95% confidence intervals (light grey, except **h**). Prevalence is shown as a function of bank vole abundance (**a, c**, and **e**), as a function of the proportion of wooded habitats in 50 m-radius zones (**b, f** and **h**) or in the trapping habitats (**d**), and as a function of the proportion of ecotones in 200 m-radius zones (**g** and **h**). Disks areas are proportional to samples sizes (see Table 2).

Figure 2: The *Borrelia burgdorferi* sensu lato prevalence in small mammals as a function of landscape variables.

The *B. burgdorferi* s.l. prevalence, expressed as the frequency of infected individuals per site, in all small mammals (**a**), and in bank voles (**b**). Fitted binomial regression curves, according to the best single explanatory variable binomial Generalized Linear Models, are shown with 95% confidence intervals (light grey). Prevalence is shown as a function of the proportion of ecotones in 50 m-radius zones (**a**) and in 100 m-radius zones (**b**). Disks areas are proportional to samples sizes (see Table 2).

Appendix 1: Example of the habitats landscape connectivity analyses.

Final aggregated land cover 5 m-resolution raster file (**a**) and 'dPC' ('difference in Probability of Connectivity') metric computed on different graphs (**b, c**, and **d**). The sampling patch on the right (surrounded in blue) appears moderately connected with Euclidian distances (**b**), while it is weakly connected when weighted by least cost paths (**c** and **d**) because it is surrounded by a river at North and by roads on other sides (see **a**). The sampling patch on the left is moderately connected for the wood mouse (**c**) while it is weakly connected for the bank vole (**d**) because it is separated to other wooded habitat patches by large grassland or crops patches resulting in fewer connections for this latter species. See **Materials and Methods** for more details.

Figure 1

Figure 2

a: Aggregated 5 m-resolution land cover raster

0   0.25   0.5   1
Kilometres
N

b: Patches' connectivity (dPC) based on Euclidan distances (250 m)

c: Least cost paths and patches' connectivity (dPC) for the Wood mouse

d: Least cost paths and patches' connectivity (dPC) for the Bank vole

Legend

Land covers for a.
- Woodland and hedgerows
- Grassland
- Crops
- Roads
- Buildings and water areas

Sampling sites
- Trap lines (**a**)
- Sampling patches (**a**, **b**, **c**, and **d**)

Connectivity measures (dPC) for b, c and d.
- High connectivity
- Low connectivity
- Least cost paths

Appendix 1