Mutational analysis reveals how the essential interactions of a ClpP

proteolytic system regulate the apicoplast proteome in malaria

parasites

Florentin A.[1,2], Stephens D.R.[2,3], Brooks C.F.[1,2], Baptista R.P.[2,4], Muralidharan V[1,2,]

[#]

[1]Department of Cellular Biology
[2]Center for Tropical and Emerging Global Diseases
[3]Department of Pharmaceutical and Biomedical Sciences, College of Pharmacy
[4]Institute of Bioinformatics
University of Georgia, Athens, GA 30602, USA

[#]Address correspondence to: vasant@uga.edu

Abstract

The human malaria parasite, *Plasmodium falciparum*, contains a plastid called the apicoplast that functions to produce essential metabolites. Most of apicoplast proteins are encoded by the nuclear genome and it is unclear how the apicoplast controls its own proteome. Here, we investigated the apicoplast caseinolytic-protease (Clp) system and how it regulates organelle proteostasis. Our data show that interfering with expression, activity or interactions of the Clp protease (PfClpP), chaperone (PfClpC), non-catalytic subunit (PfClpR), or potential substrate adaptor molecule (PfClpS) results in parasite death because of apicoplast loss. Using conditional null-mutants, we demonstrated that PfClpP is essential due to its role in apicoplast biogenesis. Conditional knockdown mutants of PfClpP revealed the robustness of its proteolytic activity and its interaction with the chaperone PfClpC. A CRISPR/Cas9 based system was developed and used for the expression of a PfClpC variant with a mutation in the putative protease binding site, which led to the discovery of its essential interaction with PfClpP. A combination of several tagged Clp proteins was used for affinity purification and determined complex composition as well as identify putative substrates. Expression of a catalytically-dead variant of PfClpP in the background of PfClpP conditional mutants showed that PfClpP oligomerizes as a zymogen and that the pro-domain is removed via trans-autocatalysis. Conditional knockdown mutants of PfClpS demonstrated its essential function in plastid biogenesis and its interaction with PfClpR as well as putative substrates. This comprehensive study reveals the molecular interactions driving the function of the apicoplast PfClpP/R/C/S system and demonstrates its central role in the biogenesis of the plastid in malaria parasites.

Introduction

The deadly malaria parasite, *Plasmodium falciparum*, has developed resistance to all currently available therapies, making the discovery of new drug targets a top priority. Like most other apicomplexans, this eukaryotic parasite contains a plastid called the apicoplast[1]. This unique organelle evolved via a two-step endosymbiosis[2]; in the primary endosymbiotic event, a cyanobacterium was incorporated into a eukaryotic cell to form the modern chloroplast. During the secondary endosymbiotic event, a photosynthetic red alga was further taken up by a protist and led to the formation of a secondary plastid[3]. Although not photosynthetic, the apicoplast harbors essential prokaryotic metabolic pathways that are essential to the parasite throughout its complex life cycle[4]. In a sharp contrast with its human host, the *Plasmodium* apicoplast shares molecular features with prokaryotes, plants and parasites, and therefore has the potential to encompass multiple parasite-specific drug targets[5,6]. Indeed, drugs that target apicoplast biology are in clinical use[7,8]. Most of those drugs (e.g. Doxycycline, Clindamycin) target the prokaryotic protein synthesis machinery in the apicoplast[9,10]. However, less than 10% of the apicoplast proteome is encoded by its own genome. The vast majority of the hundreds of apicoplast proteins are encoded in the nuclear genome and are transported to the organelle via the secretory pathway[11-13]. Available data suggest that the apicoplast does not control the translation of these nuclear encoded proteins, and, in fact, parasites without an apicoplast continue to express and accumulate these proteins in vesicle-like structures in the cytoplasm[14,15].

Due to the inability of the apicoplast to control its own protein synthesis, it is likely that it maintains a stable proteome through protein degradation. This requires an organelle specific proteolytic machinery that has not yet been identified. We hypothesize that this function is executed by Clp (Caseinolytic protease) proteins. This family of proteins consists of ClpP proteases that form multi-subunit proteolytic complexes, though the complex composition varies widely between different species and organelles[16,17]. The ClpP proteases associate with Clp ATPase chaperones that unfold and feed substrates into the ClpP barrel-like cavity for degradation[18,19]. In bacteria, they play pivotal roles in cell division, transport, stress response and virulence[20]. In plants chloroplasts, Clp proteins regulate the levels and activities of

numerous metabolic enzymes and thus control chloroplast metabolism and differentiation[21].

Some of these metabolic pathways, such as isoprenoids biosynthesis, are conserved and

essential in the apicoplast.

Several putative Clp genes have been localized to the apicoplast of *P. falciparum*, but little is

known about their roles in apicoplast biology or their essentiality for parasite asexual life

cycle[22]. The putative *Plasmodium* Clp genes differ significantly from their bacterial orthologs

and it is unclear whether they interact or even form a complex. They also include a putative

noncatalytic subunit termed PfClpR that is absent in most bacteria[23]. We have previously

shown that the *Plasmodium* ClpC chaperone (PfClpC) is essential for parasite viability and

apicoplast biogenesis[15]. Here, we report the development and application of advanced

molecular genetic tools in a clinically important non-model organism to gain detailed

mechanistic insights into plastid Clp function and how this drives parasite biology. Collectively,

these data demonstrate that the Clp complex and interactions within this proteolytic system

acts as an essential nexus regulating apicoplast biogenesis.

Results and Discussion

PfClpP is essential for apicoplast biogenesis and survival of malaria parasites

In order to assess the biological role of the *Plasmodium* ClpP homolog, PfClpP, and test its potential as a drug target, we attempted to generate a null mutant, replacing the *pfclpp* gene with a drug marker using CRISPR/Cas9 gene editing (Figure 1A). Despite PCR evidence for successful integration into the *pfclpp* locus (Supplementary Figure 1), we repeatedly failed to retrieve live parasites following drug selection. This suggested that PfClpP is essential for parasite viability and we therefore decided to repeat the transfection in the presence of isopentenyl pyrophosphate (IPP). IPP is a small metabolite produced by the apicoplast which was shown to be the only essential function of the organelle during the asexual blood stages[14]. Transfection in the presence of IPP yielded live PfClpP knockout parasites (PfClpP^{KO}) on the first attempt, and integration was confirmed by PCR (Figure 1B). Using an anti-PfClpP antibody we verified that PfClpP^{KO} mutants do not express PfClpP (Figure 1C). Removal of IPP from the culturing media resulted in the rapid death of the PfClpP^{KO} parasites (Figure 1D). Immuno-fluorescence assays (IFA) revealed that apicoplast proteins such as Acyl Carrier Protein (ACP) and Chaperonin 60 (Cpn60) lost their typical apicoplast localization, and instead appear in vesicle-like structures (Figure 1E). Such structures are indicative of damage to organelle integrity. Indeed, quantitative Real Time PCR (qRT-PCR) showed that the entire apicoplast genome disappeared in PfClpP^{KO} mutants, while the mitochondria genome was unaffected (Figure 1F). Collectively, these data demonstrate that PfClpP is essential for parasite viability, and its function is required for the biogenesis of the apicoplast organelle.

PfClpP knockdown reveals robust enzymatic activity *in vivo*

Next, we wanted to investigate the molecular mechanisms of PfClpP function. To do that, we tagged the gene with a V5 tag and the *tetR*-aptamer conditional knockdown (KD) system, enabling translational repression of PfClpP expression when anhydrotetracycline (aTc) is removed from the culturing media[24]. Using CRISPR/Cas9, we incorporated the *tetR*-aptamer regulatory cassette at the 3'end of the *pfclpp* gene, creating the PfClpP^{apt} parasite line (Figure 2A). Using PCR analysis, we verified correct integration into the *pfclpp* locus (Figure 2B), and by

IFA we confirmed its apicoplast localization (Figure 2C). Western blot analysis revealed the
expected processing pattern of PfClpP; cytoplasmic fraction (I); apicoplast-localized zymogen
(II) and a mature protease after proteolytic removal of the pro-domain (III) (Figure 2D)[15,22].
Inducing knockdown by removing aTc resulted in a significant reduction (>95%) in PfClpP
protein levels, however this reduction had no effect on parasite growth (Figure 2D and E). This
indicated that residual PfClpP protein levels were sufficient to maintain biologically functional
enzymatic activity without any deleterious effects. The fact that PfClpP is essential for
apicoplast function, and its maturation is proteolytically controlled, raised the possibility that it
is post-translationally regulated.

The PfClpP and PfClpR bind each other and share common interacting partners
Another putative member of the plastid Clp complex and a potential regulator of PfClpP
activity is PfClpR, a potential non-catalytic subunit in the Clp complex[22,23]. We therefore
attempted to tag PfClpR with the V5 tag and the *tetR* aptamer knockdown system
(Supplementary Figure 2A). Tagging *pfclpr* locus has been challenging[15] and after multiple
attempts, we finally succeeded in tagging the locus with the *tetR* aptamer. However, PCR
analysis of the aptamer repeats region in PfClpR[apt] parasites, revealed that the number of
aptamer repeats decreased during drug selection from 10 to 7 (Supplementary Figure 2B). This
significantly compromised knockdown efficiency and prevented us from evaluating its effect
on parasite replication (Supplementary Figure 2C). We concluded that tagging this locus
confers a fitness cost and that, like PfClpP, PfClpR may be essential for plastid biogenesis.
Nevertheless, the PfClpR gene was correctly tagged with the V5 epitope and, like PfClpP, was
localized to the apicoplast (Supplementary Figure 2D). It is interesting to note that the non-
catalytic PfClpR comprises an inactive Clp protease domain, but does not have a pro-domain,
and migrates similarly on SDS-PAGE as the mature PfClpP (Supplementary Figure 2E).
In order to identify other potential Clp interactors and regulators, we performed
immunoprecipitation (IP) of PfClpP[apt] and PfClpR[apt] parasites using the V5 tag. For large scale
proteomic analyses, potential Clp interactors were isolated from non-synchronized PfClpP[apt]
and PfClpR[apt] parasites using anti-V5 antibody-conjugated beads. The parental line (3D7) was

used as a control, and all samples were analyzed by mass spectrometry. All detected proteins

were filtered for predicted apicoplast localization and the abundance of each apicoplast-

predicted protein was calculated and averaged between biological replicates. Using individual

peptide abundance values, we calculated a ratio between protein abundance in IP (PfClpP or

PfClpR) and the parental control and set a threshold of $\geq$5-fold enrichment (Figure 2F and G).

This resulted in a list of proteins that were either detected exclusively in the Clp IP or were at

least 5- fold or more abundant than in the control. PfClpP IP yielded 45 proteins that are

predicted to localize to the apicoplast, and 29 out of these passed the threshold and were

considered high-confidence PfClpP interactors (Figure 2F and supplementary table 1). For

PfClpR, out of 29 apicoplast-predicted proteins, only 8 passed the threshold and were

considered probable PfClpR interactors (Figure 2G and supplementary table 1). Importantly,

PfClpR appeared as a high-confidence interactor for PfClpP and vice versa, suggesting a

physical interaction between the two components of the Clp complex. Moreover, 7 out of the 8

PfClpR positive hits were also found in the PfClpP interactors list, further reinforcing them as

high-confidence interactors (Figure 2H). One of these seven proteins is chaperonin 60 (Cpn60),

which functions in folding of proteins transported to plastids[25]. The interaction of Cpn60 with

the Clp complex suggests that these proteins may function as a protein quality control

mechanism in the apicoplast to degrade damaged and misfolded proteins.

PfClpC interacts with mature PfClpP in the apicoplast

The IP approach relied primarily on strong protein-protein interactions and could not detect

weaker or transient interactions with potential biological significance. For example, the mass

spectrometry results did not identify any Clp chaperone that may interact with the complex.

While the *Plasmodium* genome does not encode typical bacterial chaperones such as ClpA or

ClpX, it does express an atypical AAA+ ATPase termed PfClpC with a potential tripeptide ClpP

binding sequence[15,22]. In order to study factors such as PfClpC, we wanted to express potential

PfClpP regulators *in vivo* in PfClpP[apt] parasites. For this purpose, we developed a novel,

modular tool for stable expression of tagged proteins in *P. falciparum*, that avoids the

inconsistent overexpression that occurs when using episomal plasmids. In this method, we use

CRISPR/Cas9 editing to insert the gene of interest into a specific genomic locus, where it is expressed under an endogenous promoter. We chose the *pfhsp110c* locus, that encodes an essential cytoplasmic protein that is consistently expressed throughout the parasite life cycle[26]. To test whether the Hsp110 system can drive expression of apicoplast localized proteins, we introduced a 2A skip peptide at the end of the gene, followed by a GFP reporter with an apicoplast transit peptide (tp) derived from acyl carrier protein (ACP)[11] (tp^{ACP}-GFP, Figure 3A). Transfection and integration into the into *the pfhsp110c* locus were efficient, and using IFA we confirmed robust GFP expression in the apicoplast of all clones isolated (Figure 3B). Western blot analysis revealed the typical double-band, consisting of a weak upper band of the cytoplasmic fraction and a more prominent lower band of the apicoplast localized GFP after removal of the transit peptide (Figure 3C). We concluded that this method works well for expression of apicoplast proteins, and can be used to study PfClpP-PfClpC interactions.

Using the Hsp110 expression system, we designed and expressed two isoforms of PfClpC in PfClpP^{apt} parasites (Figure 3A). The first was a copy of wild type PfClpC (PfClpC^{wt}) with a C-terminal 3xTy tag. We transfected PfClpC^{wt}, and confirmed its co-expression with PfClpP-V5 by western blot (Figure 3D, left lane) and their co-localization in the apicoplast by IFA (Figure 3F, upper panel). Importantly, PfClpP pulldown resulted in the co-immunoprecipitation of PfClpC^{wt}, indicating a physical interaction between the chaperone and the protease (Figure 3E, left). The reciprocal PfClpC pulldown revealed that the chaperone interacts with the mature protease (III) but not with the zymogen form (II) or the full length PfClpP with the transit peptide (I) (Figure 3E, right). This indicates that PfClpC and PfClpP interact only upon co-localization in the apicoplast and after the proteolytic maturation of the protease.

The Clp chaperone-protease interaction is essential for plastid biogenesis and parasite survival

The second PfClpC variant that we expressed via the Hsp110 system had a single point mutation in the 3rd amino acid of a Leu-Gly-Phe motif (aa 1101-1103) that may be required for binding PfClpP (F1103W, Figure 3A)[19]. Aside from that mutation, this construct, termed PfClpC^{F1103W}, was identical to the first (PfClpC^{wt}) and was similarly tagged with 3xTy and

expressed from the *pfhsp110c* locus in PfClpP^{apt} parasites, without interfering with the
endogenous *pfclpc* gene (Figure 3A). Unexpectedly, multiple attempts to express the
PfClpC^{F1103W} variant failed, suggesting that the mutated allele has a dominant negative effect
on parasite viability. To rule out a general cytotoxic effect of transfection or any putative lethal
consequences of interfering with PfHsp110c expression, we repeated transfections in the
presence of the essential apicoplast metabolite, IPP. Transfection of PfClpC^{F1103W} in the
presence of IPP has enabled us to retrieve live parasites. Western blot analysis revealed that in
these parasites, all processing of PfClpP was abolished and the only observed band was the
full-length cytoplasmic fraction (Figure 3D, right lane). Using microscopy, we observed that
both PfClpP^{apt} and PfClpC^{F1103W} did not appear in their typical apicoplast localization, and
instead form vesicle-like structures indicative of apicoplast damage (Figure 3F, lower panel).
Clones were isolated but their growth was entirely dependent on IPP, and removing it led to
parasites death (Figure 3G). Subsequently, quantitative real time PCR analysis revealed that
the apicoplast genome disappeared from PfClpC^{F1103W} parasites (Figure 3H). These results
indicate that interfering with the interaction between the PfClpC chaperone and the PfClpP
protease inhibits complex function. Furthermore, our data suggest that PfClpC dynamically
interacts with PfClpP and this essential interaction may be required for degradation of
substrates to ensure apicoplast biogenesis. To our knowledge, this is the first time that this
interaction has been shown to be required for any plastid biogenesis.

PfClpP oligomerizes as a zymogen and matures via trans-autoprocessing
The *Plasmodium* ClpP differs structurally from its bacterial orthologues primarily because it
contains a transit peptide and a pro-domain (Figure 2D), both of which are removed at
subsequent maturations steps. Like other proteins that are transported to the plastid, PfClpP
possesses a transit peptide that is cleaved by a putative stromal peptide peptidase in the
apicoplast[12]. A second cleavage event is required for the removal of the pro-domain through an
unknown mechanism. It is possible that another protease is responsible for PfClpP pro-domain
cleavage or that this is an autocatalytic event. It is unknown how these processing events
influence complex assembly and maturation of the protease. To address all of these questions

and to understand the mechanism of PfClpP oligomerization, we specifically interfered with

the proteolytic activity of PfClpP *in vivo*.

For this purpose, we designed a PfClpP dead-protease mutant (termed PfClpP^{DEAD}) with a

point mutation in the protease active site (Ser264Ala) that renders it catalytically inactive[27]

(Figure 4A). It also includes the swapping of two residues (Glu308Arg/ Arg285Glu) of an ion pair

at the interface of the ClpP subunits[28]. The purpose of this switch is to reduce the affinity

between PfClpP^{DEAD} and the endogenous PfClpP in order to minimize a potential dominant

negative effect which may prevent us from studying PfClpP maturation and oligomerization.

We used the Hsp110 expression system to express this Ty-tagged PfClpP^{DEAD} mutant in

PfClpP^{apt} parasites (Figure 4A). The PfClpP^{DEAD} mutant co-localized in the apicoplast with

PfClpP^{apt} (Figure 4B). Western blot analysis showed that the PfClpP^{DEAD} is processed similarly

to the wild type PfClpP^{apt} to produce the cytoplasmic fraction (I); apicoplast-localized zymogen

(II) and active protease without the pro-domain (III) (Figure 4C). However, a close examination

revealed that the proteolytic maturation of the endogenous PfClpP^{apt} was reduced in the

presence of the PfClpP^{DEAD} variant. In the parental PfClpP^{apt} parasite line, the majority of the

apicoplast localized PfClpP zymogen was cleaved (fraction III, see 'parental' in Figure 4C). In

contrast, the processing of the same endogenous V5 tagged PfClpP was inhibited in the

PfClpP^{DEAD} parasites (Figure 4C). Quantification of the ratio between the processed (III) and the

zymogen (II) fractions, revealed a decrease of about 70% in the proteolytic processing of

PfClpP^{apt} (Figure 4D).

The shift in the processing rate could be explained by competitive binding between the two

PfClpP alleles. To test this model, we co-immunoprecipitated the endogenous PfClpP^{apt} (V5

tag) and the PfClpP^{DEAD} (Ty tag). These experiments revealed that the two PfClpP alleles bind

each other, indicating that they hetero-oligomerize into a mixed complex (Figure 4E). This was

unexpected due to the swapping of the two residues (Glu308Arg/ Arg285Glu) of the ion pair at

the interface of the ClpP subunits. This cross-binding revealed several molecular features of

PfClpP activation; the ability of each protease variant to co-immunoprecipitate not only the

mature form (III) but also the zymogen (II) suggests that complex assembly precedes the final

processing step of removing the pro-domain (Figure 4E and 4H).

PfClpP protease activity is essential for complex function

The implication of the interaction between the wild-type and dead protease mutant became
apparent during PfClpP^{apt} knockdown; reducing the levels of the endogenous active protease
did not affect PfClpP^{DEAD} expression levels but it drastically inhibited its processing (Figure 4F).
The inability of the parasites to remove the pro-domain from the dead protease in the absence
of an active PfClpP, suggests that *in vivo* the PfClpP zymogen matures through trans-
autoprocessing (Figure 4H). These data further suggest that the PfClpP zymogen (II) is inactive
and this trans-autoprocessing is essential for PfClpP activation. These data support the model
that PfClpP zymogen forms oligomers, which may be required for trans-autoprocessing and
that the mechanism of maturation and oligomerization for plastid Clp proteases significantly
differs from their bacterial counterparts. Whether these are general features of ClpP zymogens
expressed in other plastids such as plant chloroplasts remains to be explored. Future structural
studies with the ClpP dead zymogens may reveal additional features such as whether the
zymogen is required to ensure proper trans-autocatalytic activity.

Importantly, the robust activity of PfClpP enabled the PfClpP^{DEAD} parasites to grow normally
despite the presence of the mixed complexes. However, the presence of the PfClpP^{DEAD}
mutant rendered them more sensitive to perturbations in PfClpP levels, and under these
settings, knockdown of endogenous PfClpP^{apt} via removal of aTc resulted in parasite death
(Figure 4G). Thus, the expression of PfClpP^{DEAD} under normal PfClpP levels revealed the
mechanisms of PfClpP oligomerization and maturation (Figure 4H). Upon PfClpP knockdown,
the lethal effects of the dead-protease proved that it is the capacity to degrade substrates that
is required for organelle biogenesis and parasite survival (Figure 4H).

The adaptor protein PfClpS is essential for plastid biogenesis and parasite survival

Our data show that the proteolytic activity of the apicoplast Clp complex is central to its
essential role in regulating plastid biogenesis. Therefore, it is essential to understand the
mechanisms by which the apicoplast ClpP/R/C complex recognizes its substrates.

Bioinformatic analysis have identified a putative substrate adaptor protein termed PfClpS, but

its localization and physiological functions have not yet been determined[29].

Therefore, we tagged PfClpS with a V5 tag and the *tetR*-aptamer knockdown system to

generate PfClpS[apt] parasites, and confirmed integration by PCR (Figure 5A). We observed co-

localization of PfClpS and the apicoplast marker ACP (Figure 5C). Removal of aTc from

PfClpS[apt] parasites led to a significant protein knockdown (Figure 5B). As a consequence, aTc

removal inhibited parasite growth, demonstrating that PfClpS is essential for parasite asexual

replication (Figure 5D). Importantly, this growth inhibition was completely rescued by addition

of IPP, linking the essential function of PfClpS to the apicoplast (Figure 5D). Microscopy

revealed that apicoplast proteins such as ACP accumulated in vesicle-like structures, and

quantitative Real Time PCR confirmed that the apicoplast genome disappears upon PfClpS

knockdown (Figure 5C and E). Collectively, these data show that PfClpS activity is essential for

parasite viability and, similar to other Clp proteins, it is required for apicoplast biogenesis.

PfClpS associates with the Clp complex and putative substrates

To examine the functional interaction between PfClpS and the Clp complex, we used the

PfClpS[apt] parasites to pull down potential interactors. The immunoprecipitation and proteomic

analyses were performed as described earlier for PfClpP and PfClpR. Out of 39 identified

potential PfClpS interactors, 20 hits passed the high-confidence threshold, including PfClpR

(Figure 5F and Supplementary Table 1). Due to the predicted function of PfClpS, this list may

be enriched with putative Clp substrates. The PfClpS phenotypic data suggest that the

substrates selected for degradation by PfClpS are essential for apicoplast biogenesis. Including

the 20 PfClpS binding proteins to the PfClpP and PfClpR interactors that were identified

earlier, resulted in a list of 35 potential Clp interactors (Supplementary table 1). Interestingly,

we found a high degree of overlap between hits, with roughly 50% of them shared between at

least 2 different IPs (PfClpP/R/S) (Figure 5G). Of a specific interest, was a small group of 5 high-

confidence hits that were common between all 3 IPs (Figure 5G, Supplementary table 1). These

predicted apicoplast proteins include a thioredoxin peroxidase, a putative RNA

methyltransferase and a putative acyl transferase (Figure 2H and Supplementary table 1).

Interestingly, these Clp interactors involved in very distinct metabolic pathways (protein
folding, RNA metabolism and lipid biosynthesis), thus emphasizing the role of the *Plasmodium*
Clp complex as a master regulator of apicoplast biology.
312
A model for proteostasis regulation by the apicoplast Clp complex
To conclude, we propose a model in which the apicoplast organelle maintains a stable
proteome via protein degradation. This central function is required, among other things, for
organelle biogenesis, and is executed and regulated by the proteolytic Clp complex (Figure
6A). As a consequence, interfering with the function of different subunits of the complex leads
to organelle loss and parasite death. In the core of the Clp complex is PfClpP, a conserved
potent serine protease (Figure 6B). PfClpP is transported to the apicoplast as a zymogen,
which hetero-oligomerizes into a complex, and then matures into the active protease through
trans-autocatalytic processing. PfClpR is a non-catalytic subunit that physically interacts with
other complex subunits and may play a regulatory role. PfClpC is the chaperone that unfolds
proteins and feed them into the proteolytic core. It interacts with the mature PfClpP but not
with the zymogen, suggesting that it binds at a later step during complex assembly.
Nevertheless, this interaction is crucial for complex activity and parasite survival. Finally, a
small adaptor molecule, PfClpS, binds specific substrates and delivers them to the complex.
Targeting protein synthesis in the apicoplast is clinically effective, as manifested by the clinical
use of antibiotics such as doxycycline as antimalarials. However, such drugs inhibit the
production of only a small fraction of apicoplast proteins, as the majority of them are
transcribed by the cell nucleus and translated by the eukaryotic ribosomes in the parasite
cytoplasm. Thus, a broader and potentially more efficient way to disrupt the apicoplast
proteome would be to target its prokaryotic Clp degradation system. Indeed, due to their
central function in prokaryotes, the bacterial ClpP homologs lie in the center of various drug
discovery programs that identified potent inhibitors[30-34]. Ongoing work testing the
effectiveness and specificity of these bacterial inhibitors against PfClpP may lead to the
identification of antimalarials[15]. Our data and model suggest that small molecules inhibiting
PfClpP activity will have a dual effect of reducing substrate cleavage as well as preventing the

further production of the mature and active PfClpP. Future work is required to identify such

potent inhibitors because our data show, similar to its bacterial orthologs, the *Plasmodium* Clp

complex is an excellent drug target.

Materials & Methods

Plasmids construction

Genomic DNA was isolated from *P. falciparum* using the QIAamp DNA blood kit (QIAGEN). All constructs utilized in this study were confirmed by sequencing. PCR products were inserted into the respective plasmids using Sequence and Ligation Independent Cloning (SLIC). Briefly, insert and cut vector were mixed with a T4 DNA polymerase and incubated for 2.5 minutes at room temperature, followed by 10 minutes incubation on ice and then transformed into bacteria. All restriction enzymes used for this study were purchased from New England Biolabs. All oligonucleotides used in this study are in Supplemental Table 2.

For the generation of PfClpP[KO] donor plasmid, 500 bp from the 5'UTR of *pfclpp* (PF3D7_0307400) were amplified using primers 1+2 and 500 bp from the 3'UTR of *pfclpp* were amplified using primers 3+4. The human dihydrofolate reductase (hDHFR) drug selection cassette was amplified from pL6 vector using primers 5+6. All three pcr products were inserted into a TOPO cloning vector (ThermoFisher). Primer 7 was used in combination with primer 6 or 4 to test for accurate integration in the parasite following transfections. To increase efficiency of CRISPR/Cas9 mediated-integration, two guide RNAs were used, targeting both ends of the PfClpP gene. For expression of a C-term PfClpP guide RNA, oligos 8+9 were inserted into pUF1-Cas9-guide, and for expression of the N-term PfClpP guide RNA, oligos 10+11 were independently inserted into pUF1-Cas9-guide as previously described[15,35].

For the generation of the PfClpP[apt] conditional mutant, 500 bps from the 3'UTR and from the C-term of *pfclpp* gene were amplified pClpP-glmS vector[15] using primers 18+19 and 20+21, respectively. The two products were conjugated together using PCR-sewing and inserted as one piece into pMG74-tetR-Dozi vector[24]. For expression of a C-term ClpP guide RNA, oligos 8+9 were inserted into pUF1-Cas9-guide.

For the generation of the PfClpR[apt] conditional mutant, 500 bps from the 3'UTR and from the C-term of *pfclpr* gene (PF3D7_1436800) were amplified from pClpR-glmS vector[15] using primers 33+34 and 35+21, respectively. The two products were conjugated together using PCR-sewing and inserted as one piece into pMG74-tetR-Dozi vector[24]. Primer 36 was used in combination with primer 34 or 38 to test for accurate integration in the parasite following

transfections. Evaluating aptamer repeats numbers was done using primers 37+38. For

expression of a C-term PfClpR guide RNA, oligos 39+40 were inserted into pUF1-Cas9-guide.

For the generation of the PfClpS^{apt} conditional mutant, 500 bps from the 3'UTR and from the

C-term of *pfclps* gene (PF3D7_1320100) were amplified from gDNA using primers 41+42 and

43+44, respectively. The two products were conjugated together using PCR-sewing and

inserted as one piece into pMG74-tetR-Dozi vector[24]. Primer 45 was used in combination with

primer 42 or 38 to test for accurate integration in the parasite following transfections. For

expression of a C-term PfClpS guide RNA, oligos 46+47 were inserted into pUF1-Cas9-guide.

Primers 24+25 and 25+26 were used to test for accurate integration in the parasite following

transfections.

For the generation of the Hsp110 expression system, we designed a repair plasmid that

contains homology sequences from the *pfhsp110c* gene (PF3D7_0708800). The repair

sequences include the last 429 bp from *pfhsp110c* gene, followed by a 2A skip peptide, a

modular tagging cassette, and the first 400 bps from *pfhsp110c* 3'UTR. The modular tagging

cassette comprised of a multiple cloning site with several optional tags including mCherry, V5,

HA and Ty tags, allowing different tagging strategies using the same vector. The entire

construct was created using gene synthesis (GeneScript) and was cloned into a puc57

backbone to create the puc57-Hsp110 vector. For expression of a PfHsp110c guide RNA, oligos

22+23 were inserted into pUF1-Cas9-guide.

For the generation of tp^{ACP}-GFP parasites, tp^{ACP}-GFP was amplified from pCEN-tp^{ACP}-GFP

using primers 27+28, and was inserted into puc57-Hsp110 vector (cut with MfeI and SpeI) using

SLIC. For the generation of PfClpC^{wt} parasites, PfClpC (PF3D7_1406600) was amplified from

gDNA using primers 29+32. To introduce the point mutation required for the generation of

PfClpC^{F1103W} parasites, PfClpC was amplified from gDNA in two pieces, using primers 29+30

and 31+32. The two pieces were PCR-sewed together and inserted into puc57-Hsp110 vector

(cut with MfeI and NheI) using SLIC.

For the generation of PfClpP^{DEAD} parasites, we introduced the following point mutations into

the Open Reading Frame of PfClpP: Ser264Ala, Glu308Arg and Arg285Glu. The gene was

synthesized (GeneScrpit) and was inserted into puc57-Hsp110 vector (cut with MfeI and NheI)

using SLIC.

Cell culture and transfections

Parasites were cultured in RPMI medium supplemented with Albumax I (Gibco) and

transfected as described earlier[15,36,37].

To generate PfClpP^{KO} parasites, a mix of three plasmids was transfected into 3D7 parasites in

the presence or absence of 200uM Isopentynyl Pyrophosphate (IPP, Isoprenoids LC). The

plasmid mix contained the PfClpP^{KO} donor plasmid and the two pUF1-Cas9-PfClpP-guide (N-

terminal and C-terminal guides). Drug pressure was applied 48 hours post transfection using

2.5 nM WR99210 (Sigma) to select for parasites expressing hDHFR.

To generate PfClpP^{apt}, PfClpR^{apt} and PfClpS^{apt} parasites, a mix of two plasmids was transfected

into 3D7 parasites in the presence of 0.5 μM anhydrotetracycline (aTc, Cayman Chemicals).

The plasmid mix contained 50 μg of pUF1-Cas9-guide (expressing the relevant guide:

PfClpP/R/S) and 50 μg of the pMG74 based donor plasmid (with the related homology regions).

Drug pressure was applied 48 hours post transfection using 2.5 μg/ml Blasticidin (BSD, Sigma).

For each transfection, at least 2 clones were isolated via limiting dilution and used in all

subsequent experiments.

To generate tp^{ACP}-GFP, PfClpP^{DEAD}, PfClpC^{wt} and PfClpC^{F1103W} parasites, a mix of two plasmids

was transfected into PfClpP^{apt} parasites. For each different transfection the primers mix

contained 2 plasmids; 50 μg of pUF1-Cas9-PfHsp110c-guide and 50 μg of the relevant marker-

less puc57-Hsp110 repair plasmid (puc57-Hsp110-tp^{ACP}-GFP/ puc57-Hsp110-PfClpP^{DEAD}/ puc57-

Hsp110-PfClpC^{wt}/ puc57-Hsp110-PfClpC^{F1103W}, respectively). For PfClpC^{F1103W}, transfections

were performed in the presence or absence of 200 μM IPP (Isoprenoids LC). Drug pressure was

applied 48 hours post transfection, using 1μM DSM1 (BEI Resources), selecting only for Cas9

expression. DSM1 was removed from the culturing media once parasites clones were isolated

using limiting dilution.

Growth assays

For PfClpP^{apt}, PfClpP^{DEAD}, PfClpR^{apt} and PfClpS^{apt}, asynchronous parasite cultures were washed

5 times and incubated without aTc. For PfClpP^{KO} and PfClpC^{F1103W}, parasite cultures were

washed once and incubated without IPP. Throughout the course of the experiment parasites

were sub-cultured to maintain the parasitemia between 1-5% and parasitemia was monitored

every 24 hours via flow cytometry. Relative parasitemia at each time point was back calculated

based on actual parasitemia multiplied by the relevant dilution factors. Parasitemia in the

presence of aTc or IPP at the end of each experiment was set as the highest relative

parasitemia and was used to normalize parasites growth. Data were analyzed using Prism

(GraphPad Software, Inc.)

For IPP rescue ($PfClpS^{apt}$, $PfClpP^{KO}$ and $PfClpC^{F1103W}$) media was supplemented with 200 µM of

IPP (Isoprenoids LC) in PBS.

Western blot

Western blots were performed as described previously[38]. Briefly, parasites were collected and

host red blood cells were permeabilized selectively by treatment with ice-cold 0.04% saponin

in PBS for 10 min, followed by a wash in ice-cold PBS. Cells were lysed using RIPA buffer,

sonicated, and cleared by centrifugation at 4°C. The antibodies used in this study were mouse

anti-GFP, JL8 (Roche, 1:3000), mouse anti-Ty SAB4800032 (Sigma, 1:1000), rabbit anti-V5,

D3H8Q (Cell Signaling, 1:1000), mouse anti-V5, TCM5 (eBioscience[TM], 1:1000), mouse

monoclonal anti-PMV (from D. Goldberg, 1:400), rabbit polyclonal anti-EF1α (from D.

Goldberg, 1:2000) and rabbit polyclonal anti-PfClpP (from W. Houry, 1:4000). The secondary

antibodies that were used are IRDye 680CW goat anti-rabbit IgG and IRDye 800CW goat anti-

mouse IgG (LICOR Biosciences, 1:20,000). The Western blot images and quantifications were

processed and analyzed using the Odyssey infrared imaging system software (LICOR

Biosciences).

Microscopy and image processing

For IFA cells were fixed using a mix of 4% Paraformaldehyde and 0.015% glutaraldehyde and

permeabilized using 0.1% Triton-X100. Primary antibodies used are mouse anti Ty1

SAB4800032 (Sigma, 1:500), rabbit anti-V5, D3H8Q (Cell Signaling, 1:100), mouse anti-V5,

TCM5 (eBioscience[TM], 1:100), rabbit anti-Cpn60 (from B. Striepen, 1:1,000) and rabbit anti-ACP

(from G. Mcfadden, 1:5,000). Secondary antibodies used are Alexa Fluor 488 and Alexa Fluor

546 (Life Technologies, 1:100). Cells were mounted on ProLong Diamond with DAPI

(Invitrogen). For live cell imaging of tp^{ACP}-GFP parasites were incubated with 8 µM Hoechst

(ThermoFisher Scientific) in PBS. The imaging was performed using DeltaVision II microscope

system with an Olympus IX-71 inverted microscope using a 100X objective. All images were

collected as Z-stack, were deconvolved using the DVII acquisition software SoftWorx and

displayed as maximum intensity projection. Image processing, analysis and display were

preformed using SoftWorx and Adobe Photoshop. Adjustments to brightness and contrast

were made for display purposes.

Flow cytometry

Aliquots of parasite cultures (5µl) were stained with 8 µM Hoechst (ThermoFisher Scientific) in

PBS. The fluorescence profiles of infected erythrocytes were measured by flow cytometry on a

CytoFlex S (Beckman Coulter, Hialeah, Florida) and analyzed by FlowJo software (Treestar,

Inc., Ashland, Oregon). The parasitemia data were fit to standard growth curve using Prism

(GraphPad Software, Inc.).

Quantitative Real Time PCR

Asynchronous parasites (PfClpP^{KO} and PfClpC^{F1103W}) or Synchronized ring stage parasites

(PfClpS^{apt}) samples were collected and genomic DNA was purified using QIAamp blood kits

(Qiagen). Primers that amplify segments from genes encoded by nuclear or organelles

genomes were designed using RealTime qPCR Assay Entry (IDT). The following primer

sequences were used: *cht1* (nuclear)- 12+13; *tufA* (apicoplast)- 14+15; *cytb3* (mitochondria)-

16+17. Reactions contained template DNA, 0.5 µM of gene specific primers, and IQ^{TM} SYBR

Green Supermix (BIORAD). Quantitative real-time PCR was carried out in triplicates and was

performed at a 2-step reaction with 95^{o}C denaturation and 56^{o}C annealing and extension for 35

cycles on a CFX96 Real-Time System (BIORAD). Relative quantification of target genes was

determined using Bio-Rad CFX manager 3.1 software. Standard curves for each primers set

were obtained by using different dilutions of control gDNA as template, and were used to

determine primers efficiency. The organelle: nuclear genome ratio of mutant parasites was

calculated relative to that of the control. Unpaired t-test (Benjamini, Krieger, Yekutieli) was

used to calculate significance between control and mutants.

Immunoprecipitation

For the pull-down proteomic analysis of PfClpP^{apt}, PfClpR^{apt} and PfClpS^{apt} parasites, the

parental line (wild type clone 3D7) was used a control and all pulldown experiments were done

in duplicates using two different clones. Immunoprecipitation protocols were performed using

anti-V5 antibody as previously described[39]. Briefly, pellets from 10^9 parasites were isolated

using cold saponin and were lysed and sonicated in Extraction Buffer (40 mM Tris HCL pH 7.6,

150 mM KCL, and 1 mM EDTA) supplemented with 0.5% NP-40 (VWR) and HALT protease

inhibitor (Thermo). 10% of the sample was kept for later analysis (input sample). Rabbit anti-V5

(Cell Signaling, D3H8Q) antibody was crosslinked to DynabeadsTM protein G (Invitrogen)

beads using 5 mM BS3 crosslinker (CovaChem) and then incubated with the supernatant at

4°C. Washes were performed using a magnetic rack (Life Technologies). Samples were run on

SDS-page and gel slices were sent to mass spectrometry analyses.

For the co-Immunoprecipitation of PfClpP^{DEAD} and PfClpP^{apt}, 2 samples of PfClpP^{DEAD} clone C9,

each containing 10^8 parasites were isolated as described above. Pellets were incubated with

either anti-Ty antibody (SAB4800032, Sigma) or with anti-V5 antibody (Cell Signaling,

D3H8Q). Input and IP samples were loaded on SDS-page and blotted with anti-Ty antibody

(SAB4800032, Sigma) or with anti-V5 (Cell Signaling, D3H8Q).

For the co-Immunoprecipitation of PfClpC^{wt} and PfClpP^{apt}, 2 samples of PfClpC^{wt}, each

containing 10^9 parasites were isolated as described above. Pellets were incubated with either

anti-Ty antibody (SAB4800032, Sigma) or with anti-V5 (Cell Signaling, D3H8Q) antibody. Input

and IP samples were loaded on SDS-page and blotted with Ty antibody (SAB4800032, Sigma)

or with anti-V5 (Cell Signaling, D3H8Q).

Mass spectrometry and data analysis

PfClpP^{apt}, PfClpR^{apt} and PfClpS^{apt} samples were sent to the proteomics shared resources in

Fred Hutchinson Cancer Research Center and were run on the OrbiDEAD Elite. The data were

searched using Proteome Discoverer 2.2 against the UP000001450 *Plasmodium falciparum*

(Uniprot Nov 2018) database including common contaminants using Sequest HT and

Percolator for validation. The search results were filtered for high confidence, with a strict 1%

false discovery rate.

All detected proteins were analyzed for predicted apicoplast localization using four different

bioinformatic algorithms: (i)PATS[40]; (ii)ApicoAP[41]; (iii)PlasmoAP[13]; and (iv) PlastNN[42]. We then

compared results from all four algorithms and used known apicoplast proteins (such as Clp

proteins) to estimate false-negative and false-positive rates for each algorithm. Based on these

data, we chose to filter for apicoplast prediction using PlasmoAP, since it had a lower false-

negative rate than PlastNN (higher sensitivity) and higher accuracy than the first two

algorithms. The abundance of each apicoplast-predicted protein was calculated by summing

the total intensities (MS1 values) of all matched peptides for each selected protein, and

normalizing by the total summed intensity of all matched peptides in the sample, as previously

described[42]. The resulted calculated value was then averaged between replicates. To filter for

high confidence hits, we calculated the abundance ratio between Clp IP (PfClpP/ PfClpR/

PfClpS) and the parental control, and set a threshold of $\geq$5 fold enrichment.

ACKNOWLEDGMENTS

We thank Geoffrey McFadden for anti-ACP antibody, Boris Striepen for anti-CPN60 antibody, Walid Houry for anti-PfClpP antibody, and Dan Goldberg for anti-PMV, and anti-EF1α antibodies; Julie Nelson at the CTEGD Cytometry Shared Resource Laboratory for help with flow cytometry and analysis; and Muthugapatti Kandasamy at the Biomedical Microscopy Core at the University of Georgia for help with microscopy. We acknowledge assistance of the Proteomics Resource at Fred Hutchinson Cancer Research Center for mass spectrometry. This work was supported the American Heart Association Postdoctoral Fellowship to A. F., and the U.S. National Institutes of Health (R21AI128195) to V. M.

Figure 1

A. Generation of PfClpP knockout (PfClpP^{KO}) parasites. CRISPR/Cas9 and two guide RNAs
(gRNA) targeting the N- and the C-termini of PfClpP were used to facilitate integration.
The repair plasmid contained 500bps homology regions to the 5'UTR and the 3'UTR of the
pfclpp locus flanking the human dihydrofolate reductase selection marker (hDHFR).
Through double crossover integration and drug selection, the gene was replaced with the
drug marker. This strategy failed to retrieve live parasites unless IPP was added during drug
selection.

B. PCR test confirming hDHFR integration at the *pfclpp* locus in the presence of IPP. Genomic
DNA was purified from transfected parasites. Primers 1+2 (see methods) were used to
amplify an integration specific product. Primers 1+3 were used to amplify the region
between the 5'UTR and 3'UTR of *pfclpp*, as illustrated in A. A shift of 1 KB corresponds to
the integration of the hDHFR cassette.

C. Western blot of parasite lysates from parental line and PfClpP^{KO} parasites probed with
antibodies against PfClpP and Plasmepsin V (PMV, loading control), confirming loss of
PfClpP expression in the null mutants. The protein marker sizes that co-migrated with the
probed protein are shown on the left.

D. IPP was removed from PfClpP^{KO} parasites and parasitemia was monitored every 24 hours
over 4 days via flow cytometry. Data are represented as mean ± S.E.M. (n=3 technical
replicates). One representative experiment out of three biological replicates is shown.

E. Immunofluorescence imaging of fixed PfClpP^{KO} parasites stained with antibodies against
ACP (Acyl Carrier Protein), and DAPI (upper panel) or Cpn60 (Chaperonin 60) and DAPI
(lower panel). Both ACP and Cpn60 lose their typical apicoplast localization in PfClpP^{KO} and
appear in vesicle-like structures. Z-stack images were deconvolved and projected as a
combined single image. Representative images from biological replicates are shown. Scale
bar, 5µm.

F. Genomic DNA samples were collected from PfClpP^{KO} and control parasites for quantitative
Real Time PCR analysis. Apicoplast: nuclear genome ratio was calculated for each sample.
Mitochondria: nuclear genome ratio served as a control. Genome ratios were normalized to

control parasites. Apicoplast genome is completely lost in PfClpP^{KO} mutants, while the

mitochondria genome was unaffected. One representative experiment (out of 3 biological

replicates) is shown. Data are represented as mean ± S.E.M (n=3 technical replicates,

unpaired t-test, P value =0.00013)

Figure 2

A. Diagram showing integration of the *tetR*-aptamer system into the *pfclpp* locus. The Cas9

nuclease together with a PfClpP-specific gRNA introduces a double-stranded break at the

C-terminus of the *pfclpp* gene. The repair plasmid provides the template for double-

crossover homologous recombination, introducing a 3xV5 tag and the *tetR-Dozi* cassette,

including 10 aptamer repeats.

B. PCR test confirming *tetR-Dozi* integration at the *pfclpp* locus. gDNA was purified from 2

independently selected mutant clones (1B7 and 2H1). Primers 1,2,3 (see methods) were

used to amplify the native locus (1+2) or an integration specific product (1+3), as illustrated

in the diagram in A.

C. Immunofluorescence imaging of fixed PfClpP^{apt} parasites clone 1B7 stained with antibodies

recognizing PfClpP (anti-V5, red), ACP (anti-Acyl Carrier Protein, green), and DAPI. Z-stack

images were deconvolved and projected as a combined single image. Representative

images from three biological replicates are shown. Scale bar, 5μm.

D. PfClpP^{apt} parasites were incubated without aTc for 2 days, and lysates were isolated every

24 hr. Western blot shows parasite lysates probed with antibodies recognizing PfClpP (anti-

V5) and Plasmepsin V (anti-PMV, loading control). The expected processing pattern of

PfClpP is depicted: cytoplasmic fraction (I); apicoplast-localized zymogen (II) and mature

protease after removal of the pro-domain (III). A significant reduction in PfClpP protein

levels in achieved after 24 hours. The protein marker sizes that co-migrated with the

probed protein are shown on the left.

E.   Two independent mutant clones (1B7 and 2H1) of PfClpP^{apt} parasites were grown with or

without 0.5 μM aTc, and parasitemia was monitored every 24 hr over 11 days via flow

cytometry. 100% of growth represents the highest value of calculated parasitemia (final

parasitemia in the presence of aTc). Data are fit to an exponential growth equation and are

represented as mean ± SEM (n= 3 technical replicates). One representative experiment out

of three biological replicates is shown.

F. Mass spectrometry analysis of PfClpP Immunoprecipitate, isolated from PfClpP^{apt} parasite

lysates using anti V5 antibody-conjugated beads. The parental line 3D7 was used a control

and pulldown experiments were done in duplicates using two independent PfClpP^{apt}

clones; 1B7 and 2H1. All proteins detected in both biological replicates were filtered for

predicted apicoplast localization. The abundance of each apicoplast-predicted protein was

calculated and averaged between replicates. To filter for high confidence hits, we set a

threshold of $\geq$5 fold enrichment in Clp IP vs. parental control. For PfClpP, 29 out of 45

proteins have passed the threshold and were defined as high-confidence hits, including

PfClpR.

G. PfClpR was immunoprecipitated from lysates of PfClpR^{apt} parasites using anti V5 antibody-

conjugated beads. Biological replicates were performed for both parental control and

PfClpR^{apt} parasites (Clones A10 and B5). The sample analysis and data were processed as in

F and 8 out of 29 hits passed the threshold, including PfClpP.

H. Venn diagram and a table showing the overlap between proteins identified in PfClpP and

PfClpR pulldowns. Out of the 8 PfClpR positive hits, 7 were also found in the PfClpP list,

outlining them as high confidence interactors.

Figure 3

A. Diagram showing integration and expression through the Hsp110 expression system. Cas9

nuclease together with a gRNA targeting the *pfhsp110c* locus, introduces a double-

stranded break at the c-terminus of the *pfHsp110c* gene. The repair plasmid provides two

homology regions for homologous recombination, flanking a 2A skip peptide and a tagged

gene of interest. Using this method, we introduced an apicoplast-localized GFP, a copy of

wild type PfClpC tagged with 3xTy (PfClpC^{wt}) and a PfClpC variant with a mutation in the

putative PfClpP binding site (PfClpC^{F1103W}).

B. Live fluorescence microscopy visualizing the apicoplast localized GFP (tp^{ACP}-GFP)

expressed through the *pfHsp110c* locus in PfClpP^{apt} parasites. Parasite DNA was stained

with Hoechst. Representative image from biological replicates is shown.

C. Western blot of parasite lysates from tp^{ACP}-GFP parasites probed with antibodies against

GFP and EF1α (loading control). The protein marker sizes that co-migrated with the probed

protein are shown on the left.

D. Western blot of parasite lysates from PfClpC^{wt} and PfClpC^{F1103W} parasites probed with

antibodies against V5 (PfClpP, red) and Ty(PfClpC, green). Loss of all processing of PfClpP

in PfClpC^{F1103W} parasites is due to apicoplast loss. The protein marker sizes that co-

migrated with the probed protein are shown on the left.

E. Co-Immunoprecipitation of PfClpP and PfClpC in PfClpP^{apt}; PfClpC^{wt} parasites. 10^9

parasites were isolated and pellets were incubated with either anti-V5 antibody-conjugated

beads (PfClpP pulldown, left) or anti-Ty antibody-conjugated beads (PfClpC pulldown,

rights). Input and IP samples were loaded on SDS-page and blotted with anti-Ty (PfClpC)

and anti-V5 (PfClpP) antibodies. An interaction between the protease and the chaperone is

observed. The residual heavy chain of the anti-Ty antibody that eluted with the IP sample

despite the cross linker is also denoted (*).

F. Immunofluorescence microscopy of fixed PfClpC^{wt} and PfClpC^{F1103W} parasites stained with

antibodies against V5 (PfClpP, red), Ty (PfClpC, green), and DAPI (DNA). The two proteins

co-localize in the apicoplast in PfClpC^{wt} parasites but appear in vesicle-like structures in

PfClpC^{F1103W} parasites. Z-stack images were deconvolved and projected as a combined

single image. Representative images from one of two biological replicates are shown.

Scale bar, 5μm.

G. IPP was removed from PfClpC^{F1103W} parasites and parasitemia was monitored every 24

hours over 3 days via flow cytometry. 100% of growth represents the highest value of

calculated parasitemia (final parasitemia in the presence of IPP). Data are represented as

mean ± S.E.M. (n=3). One representative experiment out of two biological replicates is

shown.

G. Genomic DNA samples were collected from PfClpC^{F1103W} and control parasites for

quantitative Real Time PCR analysis. Apicoplast: nuclear genome ratio was calculated for

each sample. Mitochondria: nuclear genome ratio served as a control. Genome ratios were

normalized to control parasites. Data are represented as mean ± S.E.M (n=3 technical

replicates, unpaired t-test, P value=0.00025). One representative experiment out of two

biological replicates are shown.

Figure 4

A. Diagram showing integration and expression of 3xTy-tagged PfClpP^{DEAD} variant into the

pfHsp110c locus. Cas9 nuclease in complex with a gRNA targeting this locus introduces a

double-stranded break at the C-terminus of the *pfHsp110c* gene. The repair plasmid

provides two homology regions for homologous recombination, flanking a 2A skip peptide

and the 3xTy-tagged PfClpP^{DEAD}. This catalytically-inactive PfClpP variant has a point

mutation in the protease active site (Ser264Ala).

B. Immunofluorescence microscopy of fixed PfClpP^{apt}; PfClpP^{DEAD} parasites stained with

antibodies against V5 (endogenous PfClpP^{apt}, red), Ty (PfClpP^{DEAD}, green) and DAPI (DNA).

The two proteins co-localize in the apicoplast. Z-stack images were deconvolved and

projected as a combined single image. Representative image from one of two biological

replicates is shown. Scale bar, 5μm.

C. Western blot of parental PfClpP^{apt} and PfClpP^{apt};PfClpP^{DEAD} parasite lysates probed with

antibodies against V5 (endogenous *wild type* PfClpP, red) and Ty (PfClpP^{DEAD}, green). The

identical migration pattern indicates that PfClpP^{apt} and PfClpP^{DEAD} are similarly processed.

The protein marker sizes that co-migrated with the probed protein are shown on the left.

D. The ratio between the mature and zymogen forms of endogenous PfClpP (V5-tagged) is

decreased upon PfClpP^{DEAD} expression, indicating that the dead variant interferes with the

processing of the endogenous protease. Quantification of the ratio of II and III fractions of

V5 tagged PfClpP from 3 biological replicates (p-value =0.0007, unpaired t-test).

E. Co-Immunoprecipitation of PfClpP^{apt} (V5-tagged) and PfClpP^{DEAD} (Ty-tagged). 10^8

parasites were isolated and pellets were incubated with either anti-V5 antibody-conjugated

beads (left) or anti-Ty antibody-conjugated beads (right). Input and IP samples were loaded

on SDS-page and blotted with anti-Ty and anti-V5 antibodies. The robust interaction

between the two Clp protease variants suggests that they hetero-oligomerize into a mixed

complex.

F. PfClpP^{apt};PfClpP^{DEAD} parasites were incubated without aTc for 24 hours, and lysates were

isolated. Western blot of these lysates was probed with antibodies against V5 (endogenous

PfClpP, red) and Ty (PfClpP^{DEAD}, green). PfClpP^{apt} knockdown results in the disappearance

of the processed form of PfClpP^{DEAD}, suggesting that the protease matures through

trans-autoprocessing.

G. PfClpP^{apt};PfClpP^{DEAD} parasite clones (C9 and D10) were grown with or without aTc, and

parasitemia was monitored every 24 hr over 11 days via flow cytometry. 100% of growth

represents the highest value of calculated parasitemia (final parasitemia in the presence of

aTc). Data are represented as mean ± SEM (n=3 technical replicates). One representative

experiment out of two biological replicates is shown.

H. A model illustrating the mechanisms of PfClpP oligomerization and maturation. PfClpP

oligomerizes as a zymogen and subsequently matures by removing its pro-domain

(denoted with *) via trans-autoprocessing. In the presence of the PfClpP^{DEAD} variant, a

mixed complex is assembled and due to the robustness of PfClpP proteolytic activity, there

is no effect on parasite growth. Upon aTc removal expression levels of endogenous PfClpP

are dramatically reduced, and predominantly PfClpP^{DEAD} zymogen complexes are formed.

Consequently, substrate degradation is inhibited and the parasites die.

Figure 5

A. Upper panel: Diagram showing integration of the *tetR*-aptamer cassette into the *pfclps* locus. Cas9 nuclease with the gRNA introduces a double-stranded break at the beginning of the C-terminus of the *pfclps* gene. The repair plasmid provides homology regions for double-crossover homologous recombination, introducing a 3xV5 tag and the *tetR*-aptamer cassette. Lower panel: PCR test confirming *tetR*-aptamer integration at the *pfclps* locus. gDNA was purified from transfected parasites. Primers 1+2 or 1+3 (see methods) were used to amplify the native locus or an integration specific band, as illustrated in the diagram.

B. PfClpS^{apt} parasites were incubated without aTc for 4 days, and lysates were isolated every 24 hr. Western blot of parasite lysates probed with antibodies against V5 (PfClpS, red) and Plasmepsin V (PMV, loading control, green). A significant reduction in PfClpS protein levels in apparent after 72 hours. The protein marker sizes that co-migrated with the probed protein are shown on the left.

C. Quantification of PfClpS knockdown (as shown in B, n= 3 biological replicates). Values were normalized to PfClpS levels in the presence of aTc.

D. Immunofluorescence microscopy of fixed PfClpS^{apt} parasites that were incubated for 8 days with aTc (upper panel) or without aTc and supplemented with IPP (lower panel). Parasites were stained with antibodies against V5 (PfClpS, red), ACP (Acyl Carrier Protein, green), and DAPI (DNA). PfClpS localizes to the apicoplast when aTc is present. Upon aTc removal, PfClpS disappears and ACP appears in vesicle-like structures. Z-stack images were deconvolved and projected as a combined single image. Representative images from one of two biological replicates are shown. Scale bar, 5μm.

E. PfClpS^{apt} parasites (Clone A2) were grown with or without aTc and IPP, and parasitemia was monitored every 24 hr over 7 days via flow cytometry. Parasites grown without aTc exhibit a growth defect by day 4, and they die in the following days. Addition of IPP rescues the growth defect, indicating an apicoplast damage. 100% of growth represents the highest value of calculated parasitemia (final parasitemia in the presence of aTc). Data are

represented as mean ± SEM (n= 3 technical replicates). One representative experiment out

of three biological replicates is shown.

F. Synchronized PfClpS^{apt} parasites were grown in the absence of aTc and presence of IPP.

Genomic DNA samples were taken at the beginning of each replication cycle for real-time

qPCR analysis. Apicoplast: nuclear genome ratio was calculated every 48h. Mitochondria:

nuclear genome ratio served as a control. Genome ratios were normalized to parasites

grown in the presence of aTc. Data are represented as mean ± SEM (n= 3 technical

replicates). One representative experiment out of three biological replicates is shown.

G. Mass spectrometry analysis of PfClpS immunoprecipitated from PfClpS^{apt} parasites using

anti V5 antibody-conjugated beads. The parental line 3D7 was used a control and pulldown

experiments were done in duplicates. 10^9 parasites pellets were isolated and incubated

with anti-V5 crosslinked-beads. Samples were analyzed by mass spectrometry. All

detected proteins detected in biological replicates were filtered for predicted apicoplast

localization. The abundance of each apicoplast-predicted protein was calculated and

averaged between replicates. To filter for high confidence hits, we set a threshold of ≥5

fold enrichment in PfClpS vs. parental control. PfClpS IP yielded 30 apicoplast-localized

proteins, out of which 20 have passed the threshold and were defined as high-confidence

hits, including PfClpR.

H. Venn diagram showing the overlap between PfClpS, PfClpP and PfClpR high-confidence

interactors. Out of the 35-total high-confidence hits, 17 are shared between more than one

Clp IP and 5 proteins were identified in all 3 proteomic experiments

Figure 6

A. The Clp complex is a master regulator of apicoplast biology. Most of apicoplast proteins are encoded by the cell nucleus, and are transported to the organelle via the secretory pathway. According to this model, the apicoplast does not control their synthesis, and therefore a stable proteome in the organelle is via protein degradation. This essential function is required, among other things, for organelle biogenesis, and is executed and regulated by the proteolytic Clp complex. As a consequence, interfering with the function of different subunits of the complex leads to organelle loss and parasite death.

B. Molecular function of the apicoplast Clp system. In the core of the Clp complex is PfClpP, a potent serine protease. PfClpP oligomerizes as a zymogen and then matures through trans-autoprocessing. PfClpR is a non-catalytic subunit that interacts with other Clp proteins and may have a regulatory role. PfClpC is the chaperone that unfolds proteins and feed them into the PfClpP proteolytic core. A transient interaction between PfClpC and the mature PfClpP is crucial for complex stability and function. A small adaptor molecule, PfClpS, binds and selects specific substrates for delivery to the complex.

771
772
Supplementary Figure 1

A diagram depicting the generation of PfClpP knockout (PfClpP[KO]) parasites. Cas9 nuclease

and two guide RNAs targeting the N- and the C-termini of PfClpP were used to facilitate

integration. The repair plasmid contained 500bps homology regions to the 5'UTR and the

3'UTR of the *pfclpp* locus flanking a drug marker (hDHFR, human Dihydrofolate Reductase).

Through double crossover integration and drug selection, the gene was replaced with the

drug marker. Despite evidence for integration into the genomic locus (bottom), this

strategy failed to retrieve live parasites in the absence of IPP, indicating the PfClpP is

essential to parasite viability due to its apicoplast function.

Supplementary Figure 2

A. Diagram showing integration of the *tetR*-aptamer system into the *pfclpr* locus. Cas9

nuclease with a specific guide RNA introduces a double-stranded break at the C-terminus

of the *pfclpr* gene. The repair plasmid provides homology regions for double-crossover

homologous recombination, introducing a 3xV5 tag and the *tetR*-aptamer cassette.

PCR test confirming *tetR-aptamer* integration at the *pfclpr* locus. Genomic DNA was

purified from 2 mutant clones (A10 and B5). Primers 1,2,3 (see methods) were used to

amplify the native locus (1+2) or an integration specific product (1+3), as illustrated in the

diagram.

B. PCR test to evaluate aptamer repeat numbers in *tetR*-aptamer cassette in genomic DNA

isolated from PfClpP^apt, PfClpR^apt, and PfClpS^apt clones. The original plasmid contained 10

aptamer repeats, each 100pb long. Reduction in repeats numbers implies decreased

binding by the *tet*-Repressor which compromises knockdown efficiency. While PfClpP^apt

and PfClpS^{apt} clones show the expected 1 kb PCR product which corresponds to 10 aptamer

repeats, all PfClpR^{apt} clones show a 0.7 kb product, indicating the loss of 3 aptamer repeats.

C. Two PfClpR^{apt} clones (A5 and B10) were grown with or without 0.5 uM aTc, and parasitemia

was monitored every 24 hr over 11 days via flow cytometry. 100% of growth represents the

highest value of calculated parasitemia (final parasitemia in the presence of aTc). Data are

represented as mean ± SEM (n= 3 technical replicates). One representative experiment out

of two biological replicates is shown.

D. Immunofluorescence microscopy of fixed PfClpR^{apt} parasites stained with antibodies

against V5 (PfClpR^{apt}, red), ACP (Acyl Carrier Protein, green), and DAPI (DAPI). Z-stack

images were deconvolved and projected as a combined single image. Representative

images from one of two biological replicates are shown. Scale bar, 5μm.

E. PfClpR^{apt} lysates were isolated and probed with antibodies against V5 (PfClpR^{apt}, red), and

PMV (loading control). The doublet observed for PfClpR is typical for apicoplast proteins

and include a cytoplasmic fraction (I) and an apicoplast-localized protein after the transit

peptide had been removed (II). The protein marker sizes that co-migrated with the probed

protein are shown on the left.

Supplementary Figure 3

Immunoprecipitation of PfClpP[apt], PfClpR[apt] and PfClpS[apt] for proteomic analysis. 10^9 parasites

were isolated and 10% of the volume was saved as an input sample. Immunoprecipitation of

the tagged proteins from respective parasite lysates were performed using anti-V5 antibody-

conjugated beads. Input, IP and unbound samples were separated on SDS-PAGE and blotted

with anti-V5 antibodies and anti-PMV (Plasmpesin V, loading control). The heavy chain of the

antibody that eluted with some of the IP samples, despite cross linking is shown (*).

References

References

1.   McFadden, G. I. Plastid in human parasites. *Nature* **381,** 482–482 (1996).

2.   Köhler, S., Delwiche, C. F., Denny, P. W., Tilney, L. G., Webster, P., Wilson, R. J., Palmer,
J. D. & Roos, D. S. A plastid of probable green algal origin in Apicomplexan parasites.
Science **275,** 1485–1489 (1997).

3.   van Dooren, G. G. & Striepen, B. The algal past and parasite present of the apicoplast.
Annu. Rev. Microbiol. **67,** 271–289 (2013).

4.   Ralph, S. A., Foth, B. J., Hall, N. & McFadden, G. I. Evolutionary Pressures on Apicoplast
Transit Peptides. *Mol Biol Evol* **21,** 2183–2194 (2004).

5.   Fichera, M. E. & Roos, D. S. A plastid organelle as a drug target in apicomplexan
parasites. *Nature* **390,** 407–409 (1997).

6.   Erica L Dahl, P. J. R. Multiple Antibiotics Exert Delayed Effects against the Plasmodium
falciparum Apicoplast. *Antimicrobial Agents and Chemotherapy* **51,** 3485–3490 (2007).

7.   Borrmann, S., Issifou, S., Esser, G., Adegnika, A. A., Ramharter, M., Matsiegui, P.-B.,
Oyakhirome, S., Mawili-Mboumba, D. P., Missinou, M. A., Kun, J. F. J., Jomaa, H. &
Kremsner, P. G. Fosmidomycin-clindamycin for the treatment of Plasmodium
falciparum malaria. *J Infect Dis.* **190,** 1534–1540 (2004).

8.   Sponer, U., Prajakwong, S., Wiedermann, G., Kollaritsch, H., Wernsdorfer, G. &
Wernsdorfer, W. H. Pharmacodynamic interaction of doxycycline and artemisinin in
Plasmodium falciparum. *Antimicrobial Agents and Chemotherapy* **46,** 262–264 (2002).

9.   Dahl, E. L., Shock, J. L., Shenai, B. R., Gut, J., DeRisi, J. L. & Rosenthal, P. J. Tetracyclines
specifically target the apicoplast of the malaria parasite Plasmodium falciparum.
Antimicrobial Agents and Chemotherapy **50,** 3124–3131 (2006).

10.   Goodman, C. D., Su, V. & McFadden, G. I. The effects of anti-bacterials on the malaria
parasite Plasmodium falciparum. *Molecular and Biochemical Parasitology* **152,** 181–191
(2007).

11.   Waller, R. F., Reed, M. B., Cowman, A. F. & McFadden, G. I. Protein trafficking to the
plastid of Plasmodium falciparum is via the secretory pathway. *EMBO J* **19,** 1794–1802
(2000).

12.   van Dooren, G. G., Su, V., D'Ombrain, M. C. & McFadden, G. I. Processing of an

Apicoplast Leader Sequence inPlasmodium falciparum and the Identification of a

Putative Leader Cleavage Enzyme. *Journal of Biological Chemistry* **277,** 23612–23619

(2002).

13.   Foth, B. J., Ralph, S. A., Tonkin, C. J., Struck, N. S., Fraunholz, M., Roos, D. S., Cowman,

A. F. & McFadden, G. I. Dissecting Apicoplast Targeting in the Malaria Parasite

Plasmodium falciparum. *Science* **299,** 705–708 (2003).

14.   Yeh, E. & DeRisi, J. L. Chemical rescue of malaria parasites lacking an apicoplast defines

organelle function in blood-stage Plasmodium falciparum. *PLoS Biol* **9,** e1001138 (2011).

15.   Florentin, A., Cobb, D. W., Fishburn, J. D., Cipriano, M. J., Kim, P. S., Fierro, M. A.,

Striepen, B. & Muralidharan, V. PfClpC Is an Essential Clp Chaperone Required for

Plastid Integrity and Clp Protease Stability in Plasmodium falciparum. *Cell Rep* **21,** 1746–

1756 (2017).

16.   Olivares, A. O., Baker, T. A. & Sauer, R. T. Mechanistic insights into bacterial AAA+

proteases and protein-remodelling machines. *Nat Rev Micro* **14,** 33–44 (2016).

17.   Nishimura, K. & van Wijk, K. J. Organization, function and substrates of the essential Clp

protease system in plastids. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1847,**

915–930 (2015).

18.   Weber-Ban, E. U., Reid, B. G., Miranker, A. D. & Horwich, A. L. Global unfolding of a

substrate protein by the Hsp100 chaperone ClpA. *Nature* **401,** 90–93 (1999).

19.   Kim, Y.-I., Levchenko, I., Fraczkowska, K., Woodruff, R. V., Sauer, R. T. & Baker, T. A.

Molecular determinants of complex formation between Clp/Hsp100 ATPases and the

ClpP peptidase. *Nature Structural & Molecular Biology* **8,** 230–233 (2001).

20.   Frees, D., Gerth, U. & Ingmer, H. Clp chaperones and proteases are central in stress

survival, virulence and antibiotic resistance of Staphylococcus aureus. *International*

Journal of Medical Microbiology **304,** 142–149 (2014).

21.   Rodríguez-Concepción, M., D'Andrea, L. & Pulido, P. Control of plastidial metabolism by

the Clp protease complex. *J. Exp. Bot.* **70,** 2049–2058 (2018).

22.   Bakkouri, El, M., Pow, A., Mulichak, A., Cheung, K. L. Y., Artz, J. D., Amani, M., Fell, S.,
880   de Koning-Ward, T. F., Goodman, C. D., McFadden, G. I., Ortega, J., Hui, R. & Houry, W.
881   A. The Clp chaperones and proteases of the human malaria parasite Plasmodium
882   falciparum. *Journal of Molecular Biology* **404,** 456–477 (2010).
883   23.   Bakkouri, El, M., Rathore, S., Calmettes, C., Wernimont, A. K., Liu, K., Sinha, D., Asad,
884   M., Jung, P., Hui, R., Mohmmed, A. & Houry, W. A. Structural insights into the inactive
885   subunit of the apicoplast-localized caseinolytic protease complex of Plasmodium
886   falciparum. *Journal of Biological Chemistry* **288,** 1022–1031 (2013).
887   24.   Ganesan, S. M., Falla, A., Goldfless, S. J., Nasamu, A. S. & Niles, J. C. Synthetic RNA-
888   protein modules integrated with native translation mechanisms to control gene
889   expression in malaria parasites. *Nature Communications* **7,** 10727 (2016).
890   25.   Wilson, R. H. & Hayer-Hartl, M. Complex Chaperone Dependence of Rubisco Biogenesis.
891   *Biochemistry* **57,** 3210–3216 (2018).
892   26.   Muralidharan, V., Oksman, A., Pal, P., Lindquist, S. & Goldberg, D. E. Plasmodium
893   falciparum heat shock protein 110 stabilizes the asparagine repeat-rich parasite
894   proteome during malarial fevers. *Nature Communications* **3,** 1310 (2012).
895   27.   Flynn, J. M., Neher, S. B., Kim, Y.-I., Sauer, R. T. & Baker, T. A. Proteomic discovery of
896   cellular substrates of the ClpXP protease reveals five classes of ClpX-recognition signals.
897   *Mol. Cell* **11,** 671–683 (2003).
898   28.   Trentini, D. B., Suskiewicz, M. J., Heuck, A., Kurzbauer, R., Deszcz, L., Mechtler, K. &
899   Clausen, T. Arginine phosphorylation marks proteins for degradation by a Clp protease.
900   *Nature* **539,** 48–53 (2016).
901   29.   AhYoung, A. P., Koehl, A., Vizcarra, C. L., Cascio, D. & Egea, P. F. Structure of a putative
902   ClpS N-end rule adaptor protein from the malaria pathogen Plasmodium falciparum.
903   *Protein Science* **25,** 689–701 (2016).
904   30.   Gersch, M., Famulla, K., Dahmen, M., Göbl, C., Malik, I., Richter, K., Korotkov, V. S.,
905   Sass, P., Rübsamen-Schaeff, H., Madl, T., Brötz-Oesterhelt, H. & Sieber, S. A. AAA+
906   chaperones and acyldepsipeptides activate the ClpP protease via conformational
907   control. *Nature Communications* **6,** 6320 (2015).

31. Hackl, M. W., Lakemeyer, M., Dahmen, M., Glaser, M., Pahl, A., Lorenz-Baath, K.,
Menzel, T., Sievers, S., Böttcher, T., Antes, I., Waldmann, H. & Sieber, S. A. Phenyl
Esters Are Potent Inhibitors of Caseinolytic Protease P and Reveal a Stereogenic Switch
for Deoligomerization. *J. Am. Chem. Soc.* **137,** 8475–8483 (2015).

32. Brötz-Oesterhelt, H. & Sass, P. Bacterial caseinolytic proteases as novel targets for
antibacterial treatment. *International Journal of Medical Microbiology* **304,** 23–30 (2014).

33. Böttcher, T. & Sieber, S. A. β-Lactones as Specific Inhibitors of ClpP Attenuate the
Production of Extracellular Virulence Factors of Staphylococcus aureus. *J. Am. Chem.*
Soc. **130,** 14400–14401 (2008).

34. Conlon, B. P., Nakayasu, E. S., Fleck, L. E., LaFleur, M. D., Isabella, V. M., Coleman, K.,
Leonard, S. N., Smith, R. D., Adkins, J. N. & Lewis, K. Activated ClpP kills persisters and
eradicates a chronic biofilm infection. *Nature* **503,** 365–370 (2013).

35. Ghorbal, M., Gorman, M., Macpherson, C. R., Martins, R. M., Scherf, A. & Lopez-Rubio,
J.-J. Genome editing in the human malaria parasite Plasmodium falciparum using the
CRISPR-Cas9 system. *Nature Biotechnology* **32,** 819–821 (2014).

36. Cobb, D. W., Florentin, A., Fierro, M. A., Krakowiak, M., Moore, J. M. & Muralidharan, V.
The Exported Chaperone PfHsp70x Is Dispensable for the Plasmodium falciparum
Intraerythrocytic Life Cycle. *mSphere* **2,** e00363–17 (2017).

37. Kudyba, H. M., Cobb, D. W., Florentin, A., Krakowiak, M. & Muralidharan, V.
CRISPR/Cas9 Gene Editing to Make Conditional Mutants of Human Malaria Parasite P.
falciparum. *J Vis Exp* (2018). doi:10.3791/57747

38. Muralidharan, V., Oksman, A., Iwamoto, M., Wandless, T. J. & Goldberg, D. E.
Asparagine repeat function in a Plasmodium falciparum protein assessed via a
regulatable fluorescent affinity tag. *Proceedings of the National Academy of Sciences*
108, 4411–4416 (2011).

39. Kudyba, H. M., Cobb, D. W., Fierro, M. A., Florentin, A., Ljolje, D., Singh, B., Lucchi, N.
W. & Muralidharan, V. The endoplasmic reticulum chaperone PfGRP170 is essential for
asexual development and is linked to stress response in malaria parasites. *Cellular*
Microbiology **23,** e13042 (2019).

40.   Zuegge, J., Ralph, S., Schmuker, M., McFadden, G. I. & Schneider, G. Deciphering

938   apicoplast targeting signals--feature extraction from nuclear-encoded precursors of

939   Plasmodium falciparum apicoplast proteins. *Gene* **280,** 19–26 (2001).

940   41.   Cilingir, G., Broschat, S. L. & Lau, A. O. T. ApicoAP: the first computational model for

941   identifying apicoplast-targeted proteins in multiple species of Apicomplexa. *PLoS ONE*

942   **7,** e36598 (2012).

943   42.   Boucher, M. J., Ghosh, S., Zhang, L., Lal, A., Jang, S. W., Ju, A., Zhang, S., Wang, X.,

944   Ralph, S. A., Zou, J., Elias, J. E. & Yeh, E. Integrative proteomics and bioinformatic

945   prediction enable a high-confidence apicoplast proteome in malaria parasites. *PLoS Biol*

946   **16,** e2005895 (2018).

947