17a-Estradiol limits the impact of ibuprofen upon community

respiration by streambed biofilms in a sub-urban stream.

Peter McClean[1,2], William Ross Hunter[1,2]*

[1]University of Ulster, School of Geography and Environmental Science, Coleraine,

BT52 1SA, United Kingdom.

*Corresponding author. Email: w.hunter@ulster.ac.uk

Co-first author note: [1,2]These authors contributed equally to this work.

Acknowledgment

This work was completed by PMcC during his final year undergraduate research

project, supervised by WRH. It was funded through start-up funds provided to WRH

by the University of Ulster's School of Geography and Environmental Science. We

acknowledge fieldwork assistance by Ashley Williamson, and technical support in the

lab from Peter Devlin and Hugo McGrogan.

Disclaimer: We declare no conflicts of interest relating to this study.

Data Accessibility statement

Title: **17a-Estradiol limits the impact of ibuprofen upon community**

respiration by streambed biofilms in a sub-urban stream.

Abstract

Pharmaceuticals compounds such as the non-steroidal anti-inflammatory drug

ibuprofen and the artificial estrogen 17a-estradiol are contaminants of emerging

concern in freshwater systems. Globally, the use of these compounds is growing by

around ~3 % per year, yet we know little about how interactions between different

pharmaceuticals may affect aquatic ecosystems. Here we test how interactions

between ibuprofen and 17a-estradiol affect the growth and community metabolism of

streambed biofilms. We used contaminant exposure experiments to quantify how

these compounds affected biofilm growth (biomass), respiration and gross primary

production, both individually and in combination. Within our study, we found no

effects of either ibuprofen or 17a-estradiol on biofilm biomass (using ash free dry

mass as a proxy) or gross primary production. Ibuprofen significantly reduced biofilm

respiration. However, concomitant exposure to 17a-estradiol counteracted the

depressive effects ibuprofen upon biofilm metabolism. Our study, thus, demonstrates

that interactions between pharmaceuticals in the environment may have complex

effects upon microbial contributions to aquatic ecosystem functioning.

Key Words

17a-Estradiol; Biofilm; Contaminants of Emerging Concern; Ibuprofen; Microbial

Metabolism; Pharmaceuticals and Personal Care products

1. Introduction

Human pharmaceuticals and personal care products (PPCPS) are contaminants of emerging concern within the environment [1, 2]. Since the year 2000, pharmaceutical use has grown by approximately 3% per year globally and this predicted to increase further as human populations grow [3]. Removal of Pharmaceuticals and personal care products (PPCPs) via waste-water treatment plants (WWTPs) is inefficient leading to constant release of compounds which are often specifically designed specifically to produce physiological effects within an organism, at ultra-low (nano-molar) concentrations [2, 4]; compounds such as non-steroidal anti-inflammatory drugs (NSAIDs) (e.g. ibuprofen), antimicrobial compounds (e.g. triclosan, and trimethoprim) and endocrine inhibitors (e.g. Estradiol) into the aquatic environment [4-7]. Eco-toxicological studies reveal that PPCPs at environmental concentrations can have significant physiological effects on both aquatic fauna and microorganisms, with the potential to disrupt the functioning of aquatic ecosystems, alter carbon and nutrient cycling, and negatively affect water quality [8-12].

Headwater and lower order streams are the smallest tributaries of a river system, which are typically closest to the rivers' sources. In these streams aquatic biofilms attached to the streambed represent the dominant mode of microbial life [13, 14]. Biofilms are composed of consortia of bacteria and unicellular eukaryotic algae bound within a complex matrix of extracellular polymeric substances (EPS) and play a key role in the functioning of fluvial ecosystems, controlling both the transport and degradation of organic matter within a stream [14]. Rosi-Marshall et al. [10] revealed that aquatic PPCPs such as caffeine, cimetidine, ciprofloxacin, diphenhydramine, metformin and ranitidine and negative effects upon biofilm growth, respiration, and community composition. PPCPs, however, are diverse group of chemicals, which may interact with each other in a multitude of different, and often-unexpected ways [1, 10, 15]. Consequently, a mechanistic understanding of the interactions between different PPCPs is needed if we are to fully understand their environmental impacts.

Within the broad spectrum of PPCPs the non-steroidal anti-inflammatories (NSAIDs), such as ibuprofen, and endocrine disruptors, such as 17a-estradiol, represent some of the most commonly detected compounds in aquatic systems [1, 4, 9]. NSAIDs are known to have antimicrobial properties, with ibuprofen exhibiting potential as a biofilm control agent [12, 16-18]. Conversely, oestrogens and other endocrine

disruptors may adsorb onto microbial biofilms facilitating their biological degradation

[19-21]. As such, there is potential for antagonistic interactions between NSAIDs and

Endocrine Disruptors to affect the growth and metabolism of streambed biofilms.

Here we present the first data on how interactions between ibuprofen and 17a-

estradiol affect the growth and community metabolism of streambed biofilms. We

conducted in situ contaminant exposure experiments, following Costello et al. [22], to

test how chronic exposure to ibuprofen, and 17a-estradiol, both individually and in

combination, affected streambed biofilm growth, primary production and respiration.

2. Materials and Methods

All experiments were carried out between the 30[th] November 2018 and the 22[nd]

January 2019 in the Ballysally Blagh (Latitude: 55°08'45.1"N Longitude:

6°40'18.0"W), a ground-water fed second-order stream. The Ballysally Blagh is a

tributary of the lower River Bann (Northern Ireland), draining a mixed agricultural

(consisting of 21.9 % arable; 55.9 % grassland; 13.7 % heathland; 1.9 % woodland)

and urban (7.3 %) catchment of 14.2 km^2. The mean volumetric rate for water flow in

the Ballysally Blagh is 0.21 (± 0.27) m^3 s^{-1}, measured at a V-shaped weir [23] and

the stream is defined as eutrophic, with dissolved nitrate concentrations ranging

between 1.37 and 14.15 ml.l^{-1} and soluble reactive phosphorus concentrations

between 0.033 and 0.4 mg.l^{-1}. Water temperature at the study site was recorded at

1-hour intervals throughout the experiment using a HOBO MX2204 Bluetooth

temperature logger (Fig 1). Temperatures ranged between 9.35 ^{o}C and 5.16 ^{o}C, with

a mean temperature of 7.72 (± 0.85) ^{o}C recorded over the study period.

Contaminant exposure experiments were conducted following Costello et al. [22].

Briefly, forty 120 ml screw cap sample pots where filled with 2 % agar impregnated,

of which 10 were impregnated a 0.5 mmol.l^{-1} dose of ibuprofen, 10 with a 0.5 mmol.l$^-$

^{1} dose of 17a-estradiol, 10 with a 0.5 mmol.l^{-1} dose of both ibuprofen and 17a-

estradiol, and 10 received no pharmaceutical treatment (control). Both ibuprofen and

estradiol have relatively low solubility in water (21 mg.l^{-1} and 3.6 mg.l^{-1} respectively).

As such, stock solutions for each pharmaceutical treatment were made up in 70 %

ethanol, with 1 ml aliquots used to dose each contaminant exposure experiment and

the control treatments receiving a 1 ml aliquot of 70 % ethanol. Pre-combusted

Whatman® 45 mm GF/F filters were placed onto of the solid agar and secured using

the screw cap, to provide a substratum for streambed biofilm colonization.

Contaminant exposure experiments were then secured to four L-shaped metal bars

(dimensions) and deployed at 10 cm depth, in an area of turbulent flow (riffle) within

the stream.

Environmental chambers were assembled from two Curry's Essentials® C61CF13

chest freezers, with the power source re-routed through Inkbird ITC-308 Digital

Temperature Controller used to override the freezers internal thermostat. A single

Tetra HT50 (50 Watt) aquarium heater was also attached to the Inkbird temperature

controller of each unit to help stablise the internal temperature. Two NICREW

planted aquarium LED strip lights were attached to the lid, providing a source of

photosynthetically active radiation (– 106.0 µmol m^{-2} s^{-1}, measured using an Apogee

Instruments Photosynthetically Active Radiation Meter). Environmental chambers

were filled with 20 l of streamwater and the internal temperatures set to 7.7 ^{o}C. The

contaminant exposure experiments were left *in situ* for 54 days, after which they

were recovered from the stream and placed into one of the environmental chambers

and allowed to acclimate over 24 hours. During the acclimation period each

mesocosm was aerated using a Aquarline Hailea Aco-9630.

After the acclimation period, biofilm respiration and gross primary production were

determined by changes in oxygen consumption by enclosing each contaminant

exposure experiment into a sealed Perspex push core (height = 30 cm, internal

diameter = 7 cm) chambers containing xx l of streamwater and held at 7.7 ^{o}C in one

of the environmental chambers ([24, 25]. Biofilm respiration (R) were quantified by

measuring the change in oxygen concentrations over a one-hour period (oxygen

consumption in darkness (PAR ~ 0.0 µmol m^{-2} s^{-1}) using a Hach Sension 6 dissolved

oxygen meter. Net primary production (NPP) was then quantified by measuring the

change in oxygen concentration over a one 1-hour period, under artificial illumination

(PAR ~ 106.0 µmol m^{-2} s^{-1}). Biofilm Gross Primary Production (GPP) was calculated

from NPP and R as:

[1]   GPP = NPP – R

Microbial biomass within each Contaminant Exposure Experiment was quantified as

Ash Free Dry Mass of the GF/F filters. These were dried four 48 hours at 65 ^{o}C and

then subsequently combusted at 550 ^{o}C for 2 hours.

All data analysis was conducted in the R statistical computing environment using the

base and *ggplot2* packages [26, 27]. We tested for independent and combined

effects of ibuprofen and estradiol upon in microbial biomass (Ash Free Dry Weight),

Respiration and Net Ecosystem Production using two-way analysis of variance

(ANOVA). Post-hoc testing of significant interactions was conducted using Tukey's

test for Honest Significant Difference. All data were visually explored, to ensure they

conformed to the assumptions of normality and homoscedacity, following Zuur et al.

[28]. Microbial biomass data were $\log_{10}$ transformed to ensure the residuals of the

ANOVA model conformed to a normal distribution.

3. Results

Using ash free dry mass as a proxy for microbial biomass we detected no significant

effects of pharmaceutical exposure upon microbial biofilm growth (Fig 2 A;

Table 1 a). We detected a significant interaction between ibuprofen and estradiol

affecting microbial respiration (Fig 2 B; Table 1 b). Exposure to ibuprofen alone

depressed microbial oxygen consumption by ~ 38 %, whilst exposure to estradiol

alone resulted in a slight (non-significant) increase in oxygen consumption. In

combination, estradiol counteracted the depressive effect of ibuprofen upon of

microbial respiration. Gross Primary Production was negative in all treatments, with

no significant effects of the pharmaceutical treatments detected (Fig 2 C; Table 1 c).

4. Discussion

Our study demonstrates that interactions between an NSAID (ibuprofen) and an

endocrine disruptor (17a-estradiol) have a significant effect upon the metabolism of

streambed biofilms. Specifically, concomitant exposure to both 17a-estradiol and

estradiol reduces the depressive effect of ibuprofen upon biofilm respiration.

Ibuprofen is known to have antimicrobial properties and has been reported to inhibit

biofilm formation by both *Staphylococcus aureus* and *Escherichia coli* [16-18]. It is,

therefore, unsurprising that ibuprofen depressed microbial respiration within the

streambed biofilms. Estradiol has been observed to adsorb to microbial biofilms [19]

where it can then be used by the resident microorganisms as an organic matter

source [29, 30]. Consequently, biofilms have been proposed as a tool for the

removal of estradiol and other endocrine disruptors within wastewater treatment

facilities [31]. Furthermore, we propose that the sorption of estradiol to the biofilm

may protect the microbial cells, by reducing the space available within the EPS
matrix onto which ibuprofen molecules may bind. This mechanism, however,
remains speculative and would require investigation within controlled laboratory
experiments.

Given ibuprofen's potential as a biofilm control agent [12, 16-18], we were surprised
to observe that it had no effect upon biofilm biomass within our experiments. Ash
free dry mass is, however, a coarse method of estimating microbial biomass and so
may not be able to detect small changes in the biofilm. This is likely to be of
particular concern in urban and agricultural streams, where siltation may introduce a
significant bias into weight-based estimates of biomass. Visual methods, such as
microscopic cell counts [32], quantification of EPS polysaccharides [32, 33] or other
biomarkers, such as polar lipid fatty acids [34-37] may have provided a more
accurate proxy for biomass. Thus, we cannot reliably infer whether interactions
between ibuprofen and estradiol may have altered biofilm biomass within this study.

The negative values for GPP within the present study suggest that the biofilms were
net heterotrophic, relying on the supply of organic matter from the surrounding
environment to provide energy and nutrients for biofilm growth. This may reflect the
choice of agar as the carrier medium for the pharmaceuticals within the contaminant
exposure experiments. The agar releases a constant supply of dissolved organic
matter through the glass fibre filters [10, 22], which may generate favorable
microhabitat heterotrophic microorganisms. As such we were unable to determine
whether chronic pharmaceutical exposure had any effects upon photosynthetic
pathways within our biofilms.

Within this short paper we present preliminary results which demonstrate that
interactions between NSAIDs and endocrine disruptors have important implications
for aquatic ecosystem functioning during the winter period, when lower water
temperatures limit microbial activity within streambed biofilms [38]. Overall, this
suggests that PPCPs represent a major threat to ecosystem functioning in many
streams and rivers. The interactions between different PPCPs, however, are
potentially complex. Whilst identification of the underlying biochemical mechanisms
is beyond the scope of this study, our results highlight the need for detailed
ecotoxicological analysis of the multiple interactions between PPCPs and how this
effects microbial and faunal activity, in the environment.

5. References

1.   Gaston L, Lapworth DJ, Stuart M, Arnscheidt J. 2019. Prioritization Approaches

for Substances of Emerging Concern in Groundwater: A Critical Review.

Environmental Science & Technology 53:6107-6122.

2.   Rosi-Marshall EJ, Royer TV. 2012. Pharmaceutical compounds and ecosystem

function: an emerging research challenge for aquatic ecologists. *Ecosystems.*

15:867-880.

3.   Van Boeckel TP, Gandra S, Ashok A, Caudron Q, Grenfell BT, Levin SA,

Laxminarayan R. 2014. Global antibiotic consumption 2000 to 2010: an analysis of

national pharmaceutical sales data. *Lancet Infectious Diseases* 14:742-750.

4.   Álvarez-Muñoz D, Rodríguez-Mozaz S, Maulvault AL, Tediosi A, Fernández-

Tejedor M, Van den Heuvel F, Kotterman M, Marques A, Barceló D. 2015.

Occurrence of pharmaceuticals and endocrine disrupting compounds in

macroalgaes, bivalves, and fish from coastal areas in Europe. *Environmental*

Research 143:56-64.

5.   Gros M, Petrović M, Barcelo D. 2007. Wastewater treatment plants as a

pathway for aquatic contamination by pharmaceuticals in the Ebro river basin

(northeast Spain). *Environmental Toxicology and Chemistry* 26:1553-1562.

6.   Ruhí A, Acuña A, Barceló D, Huerta B, Mor J-, Rodríguez-Mozaz S, Sabater S.

2016. Bioaccumulation and trophic magnification of pharmaceuticals and endocrine

disruptors in a Mediterranean river food web. *Science of the Total Environment*

540:250-259.

7. Archer E, Petrie B, Kasprzyk-Hordern B, Wolfaardt GM. 2017. The fate of

pharmaceuticals and personal care products (PPCPs), endocrine disrupting

contaminants (EDCs), metabolites and illicit drugs in a WWTW and environmental

waters. *Chemosphere* 174:437-446.

8. Jobling S, Casey D, Rodgers-Gray T, Oehlmann J, Schulte-Oehlmann U,

Pawlowski S, Baunbeck T, Turner AP, Tyler CR. 2003. Comparative responses of

molluscs and fish to environmental estrogens and an estrogenic effluent. *Aquatic*

Toxicology 65:205-220.

9. Hernando MD, Mezcua M, Fernández-Alba AR, Barceló D. 2006.

Environmental risk assessment of pharmaceutical residues in wastewater effluents,

surface waters and sediments. *Talanta* 69:334-342.

10. Rosi-Marshall EJ, Kincaid DWL, Bechtold H, Royer TV, Rojas M, Kelly JJ.

2013. Pharmaceuticals suppress algal growth and microbial respiration and alter

bacterial communities in stream biofilms. *Ecological Applications* 23:583-593.

11. Drury B, Scott J, Rosi-Marshall EJK, J.J. 2013. Triclosan Exposure Increases

Triclosan Resistance and Influences Taxonomic Composition of Benthic Bacterial

Communities. *Environmental Science and Technology* 47:8923-8930.

12. Żur J, Piński A, Marchlewicz A, Hupert-Kocurek K, Wojcieszyńska D, Guzik U.

2018. Organic micropollutants paracetamol and ibuprofen: toxicity, biodegradation,

and genetic background of their utilization by bacteria. *Environmental Science and*

Pollution Research 25:21498-21524.

13.   Besemer K, Peter H, Logue JB, Langenheder S, Lindstrom ES, Tranvik LJ,

Battin TJ. 2012. Unraveling assembly of stream biofilm communities. *ISME Journal*

6:1459-1468.

14.   Battin TJ, Besemer K, Bengtsson MM, Romani AM, Packmann AI. 2016. The

ecology and biogeochemistry of stream biofilms. *Nature Reviews Microbiology*

14:251-263.

15.   Gerbersdorf SU, Cimatoribus C, Class H, Engesser K-, Helbich S, Hollert H,

Lange C, Kranert M, Metzger J, Nowak W, Seiler T-, Steger K, Steinmetz H,

Wieprecht S. 2015. Anthropogenic Trace Compounds (ATCs) in aquatic

habitats: Research needs on sources, fate, detection and toxicity to ensure timely

elimination strategies and risk management. *Environment International 79* 79:85-

105.

16.   Reśliński A, Dábrowiecki S, Głowacka K. 2015. The impact of diclofenac and

ibuprofen on biofilm formation on the surface of polypropylene mesh. *Hernia* 19:179-

185.

17.   Shah PAL, Marshall-Batty KR, Smolen JA, Tagaev JA, Chen Q, Rodesney

CA, Le HH, Gordon VD, Greenberg DE, Cannon CL. 2018. Antimicrobial Activity of

Ibuprofen against Cystic Fibrosis-Associated Gram-Negative Pathogens.

Antimicrobial Agents and Chemotherapy 62:e01574-17.

18.   Oliveira IM, Borges A, Borges F, Simões M. 2019. Repurposing ibuprofen to

control Staphylococcus aureus biofilms. *European Journal of Medicinal Chemistry*

166:197-205.

19.   Writer JH, Ryan JN, Keefe SH, Barb LB. 2012. Fate of 4-Nonylphenol and

17β-Estradiol in the Redwood River of Minnesota. *Environmental Science &*

Technology 46:860-868.

20.   Zhang X, Li Y, Liu B, Wang J, Feng C. 2014. The effects of estrone and 17β-

estradiol on microbial activity and bacterial diversity in an agricultural soil:

sulfamethoxazole as a co-pollutant. *Ecotoxicology and Environmental Safety*

107:313-320.

21.   Adeel M, Song S, Wang Y, Francis D, Yang Y. 2017. Environmental impact of

estrogens on human, animal and plant life: A critical review. *Environment*

International 99:107-119.

22.   Costello DM, Rosi-Marshall EJ, Shaw LE, Grace M, Kelly JJ. 2015. A novel

method to assess effects of chemical stressors on natural biofilm structure and

function. *Freshwater Biology* 61:2129-2140.

23.   National River Flow Archive. 2019. Station 203050: Ballysally Blagh at

University of Ulster. https://nrfa.ceh.ac.uk/data/station/spatial/203050.

24.   Bott TL, Brock JT, Cushing CE, S. V. Gregory SV, King D, Petersen RC. 1978.

A comparison of methods for measuring primary productivity and community

respiration in streams. *Hydrobiologia* 60:3-12.

25.   Fellows CS, Clapcott JE, Udy JW, Bunn SE, Harch BD, Smith MJ, Davies PM.

2006. Benthic Metabolism as an Indicator of Stream Ecosystem Health.

Hydrobiologia 572:71-87.

26.   R Development Core Team. 2009. R: A language and environment for

statistical computing.

27.   Wickham H. 2016. *ggplot2: Elegant Graphics for Data Analysis.* Springer-

Verlag New York.

28.   Zuur AF, Ieno EN, Elphick CS. 2010. A protocol for data exploration to avoid

common statistical problems. *Methods Ecology and Evolution.* 1:3-14.

29.   Stumpe B, Marschner B. 2009. Factors controlling the biodegradation of 17β-

estradiol, estrone and 17α-ethinylestradiol in different natural soils. *Chemosphere*

74:556-562.

30.   Ribeiro AR, Carvalho MF, Afonso CMM, Tiritan ME, Castro PML. 2010.

Microbial degradation of 17β -estradiol and 17α -ethinylestradiol followed by a

validated HPLC-DAD method. *Journal of Environmental Science and Health, Part B*

45:265-273.

31.   Pieper C, Rotard W. 2011. Investigation on the removal of natural and

synthetic estrogens using biofilms in continuous flow biofilm reactors and batch

experiments analysed by gas chromatography/mass spectrometry. *Water Research*

45:1105-1114.

32.   Grzegorczyk M, Pogorzelski SJ, Pospiech A, Boniewicz-Szmyt K. 2018.

Monitoring of Marine Biofilm Formation Dynamics at Submerged Solid Surfaces With

Multitechnique Sensors. *Frontiers in Marine Science* 5:363.

33.   Fish K, Osborn AM, Boxall JB. 2017. Biofilm structures (EPS and bacterial

communities) in drinking water distribution systems are conditioned by hydraulics

and influence discolouration. *Science of The Total Environment* 593-594:571-580.

34.   Middelburg JJ, Barranguet C, Boschker HTS, Herman PMJ, Moens T, Heip

CHR. 2000. The fate of intertidal microphytobenthos carbon: An in situ 13C-labeling

study. *Limnology and Oceanography.* 45:1224-1234.

35.   Frostegård A, Tunlid A, Bååth E. 2011. Use and misuse of PLFA

measurements in soils. *Soil Biology and Biochemistry.* 43:1621-1625.

36.   Hunter WR, Veuger B, Witte U. 2012. Macrofauna regulate heterotrophic

bacterial carbon and nitrogen incorporation in low-oxygen sediments. *ISME Journal*

6:2140-2151.

37.   Hunter WR, Jamieson A, Huvenne VAI, Witte U. 2013. Sediment community

responses to marine vs. terrigenous organic matter in a submarine canyon.

Biogeosciences. 10:67-80.

38.   Ylla I, Romaní AM, Sebater S. 2012. Labile and Recalcitrant Organic Matter

Utilization by River Biofilm Under Increasing Water Temperature. *Microbial Ecology*

64:593-604.

342 **Table**

343 Table 1. ANOVA summary tables of the effects of Ibuprofen and 17a-Estradiol upon
344 a) biomass (ash free dry weight), b) respiration and c) Gross Primary Production of
345 cultured streambed biofilms.
346

a) Biomass (Ash Free Dry Weight)					
	Df	SS	MS	F	p
Ibuprofen	1	0.001	0.00086	0.008	0.931
Estradiol	1	0.006	0.00586	0.051	0.822
Ibuprofen : Estradiol	1	0.151	0.15142	1.331	0.256
Residuals	36	4.097	0.11379		
b) Respiration					
	Df	SS	MS	F	p
Ibuprofen	1	6482	6482	41.13	**<0.001**
Estradiol	1	5085	5085	32.26	**<0.001**
Ibuprofen : Estradiol	1	2952	2952	18.73	**<0.001**
Residuals	36	5674	158		
c) Gross Primary Production					
	Df	SS	MS	F	p
Ibuprofen	1	40	40.4	0.037	0.848
Estradiol	1	2612	2611.7	2.403	0.130
Ibuprofen : Estradiol	1	2318	2318.1	2.133	0.153
Residuals	36	39121	1086.7		

 Figures

Figure 1. Change in water temperature over the biofilm growth period, measured at 1
hour intervals using a HOBO MX2204 Bluetooth temperature logger

Figure 2. Effects of Ibuprofen (⬥) and Estradiol (♀) upon the (A) biomass (ash free
dry weight), (B) respiration and (C) Gross Primary Production of cultured streambed
biofilms. Significance levels: ***$p < 0.001$; **$p < 0.01$; *$p < 0.05$; N/S $p > 0.05$.Where
significant interactions were identified, groups labelled with the same lowercase
letter are not significantly different ($p > 0.05$; Tukey's tests).