Perampanel rescues cortical gamma dysregulation associated with parvalbumin interneuron GluA2 upregulation in epileptic *Syngap1*[+/-] mice

Brennan J. Sullivan[1], Simon Ammanuel[2], Pavel A. Kipnis[1], Yoichi Araki[3], Richard L. Huganir[3], Shilpa D. Kadam[1,4*]

[1] Neuroscience Laboratory, Hugo Moser Research Institute at Kennedy Krieger, Baltimore, MD 21205, USA

[2] School of Medicine, University of California, San Francisco, 505 Parnassus Avenue, San Francisco, CA, 94143, USA.

[3] Department of Neuroscience, Kavli Neuroscience Discovery Institute, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

[4] Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA;

*Correspondence: kadam@kennedykrieger.org

Number of Pages: 38
Number of Figures: 10
Number of Supplemental Figures: 9
Number of Supplemental Video: 1
Number of Words in Main Text: 5969
Number of Words in Methods: 2304
Number of References: 68

Key words: Myoclonic seizures, sleep cycles, NREM, REM, interictal spikes, gamma oscillations, parvalbumin interneurons, AMPA receptors, Perampanel.

Abbreviations list: AMPAR (AMPA receptor), PV+IN (parvalbumin interneuron), IIS (interictal spike), BCX (barrel cortex), MCX (motor cortex), PMP (Perampanel), CA1 (cornu Ammonis 1), CA3 (cornu Ammonis 3), SR (stratum radiatum), SO (stratum oriens), SP (stratum pyramidale), MRD5 (mental retardation type 5), PSD (post-synaptic density), NREM (non-rapid eye movement), REM (rapid eye movement), EEG (electroencephalogram), EMG (electromyogram), NDD (neurodevelopmental disorder), TGR (theta-gamma ratio), IHC (immunohistochemical), FFT (fast Fourier transform).

Corresponding Author:
Shilpa D. Kadam, PhD
Neuroscience Laboratory, Hugo Moser Research Institute at Kennedy Krieger;
Department of Neurology, Johns Hopkins University School of Medicine,
707 North Broadway, 400H;
Baltimore, MD 21205
Phone: 443-923-2688, Fax: 443-923-2695,
E-mail: kadam@kennedykrieger.org

Abstract

Loss-of-function *SYNGAP1* mutations cause a neurodevelopmental disorder characterized by intellectual disability and epilepsy. SYNGAP1 is a Ras-GTPase-activating protein that underlies the formation and experience-dependent regulation of postsynaptic densities. The mechanisms that contribute to this proposed monogenic cause of intellectual disability and epilepsy remain unresolved. Here, for the first time we establish the phenotype of the epileptogenesis in a *Syngap1$^{+/-}$* mouse model using 24h video electroencephalogram/electromyogram (vEEG/EMG) recordings at advancing ages. A progressive worsening of seizures, interictal spike frequency, sleep dysfunction, and hyperactivity was identified in *Syngap1$^{+/-}$* mice. Interictal spike and seizures emerged predominantly during NREM in 24h vEEG of *Syngap1$^{+/-}$* mice and patients with *SYNGAP1* haploinsufficiency. In *Syngap1$^{+/-}$* mice, EEG spectral power analyses identified a significant loss of cortical gamma homeostasis during behavioral-state transitions from NREM to Wake and NREM to REM. The loss of gamma homeostasis was associated with a region and location specific significant increase in the somatic GluA2 AMPA receptor subunit expression in parvalbumin-positive (PV+) interneurons. Acute dosing with Perampanel, an FDA approved AMPA antagonist significantly rescued cortical gamma homeostasis, identifying a novel mechanism implicating Ca^{2+} impermeable AMPARs on PV+ interneurons underlying circuit dysfunction in *SYNGAP1* haploinsufficiency.

Introduction

SYNGAP1 codes for SYNGAP1, a Ras-GTPase-activating protein critical for the formation of postsynaptic densities (PSDs) and the experience-dependent AMPA receptor (AMPAR) insertion that underlies synaptic plasticity[1–5]. Mutations in *SYNGAP1* are prevalent in patients with schizophrenia, intellectual disability (ID), and autism spectrum disorder[6–8]. Severe loss-of-function *SYNGAP1* mutations result in haploinsufficiency and cause mental retardation type 5 (MRD5, OMIM#612621), a distinct generalized developmental and epileptic encephalopathy with ID, ataxia, severe behavioral problems, and a risk for autism [6–12].

The majority of patients with MRD5 have refractory epilepsy[12]. Poor quality sleep is highly prevalent in patients with neurodevelopmental disorders (NDDs) and epilepsy[13,14]. The relationship between epilepsy and dysfunctional sleep is a major focus of ongoing clinical and pre-clinical research[13–15]. For example, Rett syndrome (RTT) is an NDD with comorbidity of epilepsy and sleep dysfunction[16,17]. Research in pre-clinical models of RTT, as well as patients with RTT has identified translatable quantitative EEG (qEEG) biomarkers[17,18]. There is an urgent need to develop robust biomarkers for NDDs, as bench-to-bedside therapeutic strategies are severely hindered without validated quantitative outcome measures.

SYNGAP1 patients predominantly present with myoclonic, absence, or tonic-clonic seizures[8,12]. Clinical cohort studies reveal progressive epilepsy followed by spontaneous remission of epilepsy in a subset of SYNGAP1 patients in their late teens[12]. Children with epileptic encephalopathies have a slow regression of development that is primarily due to seizures, interictal spikes (IISs), or cortical dysrhythmia identified on EEG[19,20]. Case reports indicate that adult patients demonstrate gradual decline in cognitive abilities[21].

Mice with SYNGAP1 haploinsufficiency (*Syngap1+/-*) are relevant translational models

presenting with learning and memory deficits, abnormal dendritic spine dynamics, cortical

hyperexcitability, and precocious unsilencing of thalamocortical synapses during development

[22–24]. Currently, no studies have investigated the epileptogenesis, sleep, and underlying

mechanisms associated with MRD5. An understanding of the mechanisms that promote

spontaneous seizures and network dysfunction can help accelerate the discovery of novel

therapeutic strategies. For this purpose, a *Syngap1+/-* loss-of-function mouse model (exon 7

and 8 deletions) underwent 24h video vEEG/EMG recordings at advancing ages (P60-P120).

Results

***Syngap1+/-* mice have recurrent spontaneous seizures**

Patients with pathogenic *SYNGAP1* mutations have epilepsy[12]. The predominant

seizure phenotypes were absence, eyelid myoclonia with absences, and myoclonic seizures.

The myoclonic seizures were distinguished from atypical absences by pronounced myoclonic

jerks and a global spike wave discharge ~3Hz on EEG[12]. Continuous 24h vEEG/EMG

recordings at temporally advancing ages were utilized to investigate and identify spontaneous

seizures in 50% (n/n; 4/8) of *Syngap1+/-* mice (Fig. 1). At P60 all seizures were myoclonic (5/5);

80% (4/5) of these seizures started in non-rapid eye movement (NREM) at NREM-Wake

transitions (Fig. 1A). Myoclonic seizures were 34.4±5.9s in duration, with rhythmic spike-wave

discharges occurring ~3-4Hz frequency (Fig. 1A). The myoclonic seizures were distinguished

by their time-locked myoclonic jerks recorded on EMG with video confirmation of head, neck,

and upper shoulder myoclonic jerks (see Supplemental Video 1).

Multiple seizure phenotypes were identified in P120$^{+/-}$ mice. Myoclonic, generalized

tonic-clonic, and electrographic seizures were observed (Fig. 1B, 1C1, and Supplemental Fig.

2). The majority of seizures were electrographic seizures emerging during wake (n/n; 7/11),

while all myoclonic seizures occurred during NREM (3/11), and a single tonic-clonic seizure

occurred during NREM (1/11). Moreover, all myoclonic and tonic-clonic seizures occurred

during NREM in P120$^{+/-}$ mice, whereas all electrographic seizures occurred during wake (Fig

1C3). In P60$^{+/-}$ mice, all seizures were myoclonic and 4/5 occurred during NREM (Fig. 1C2).

From P60 to P120, the average seizure duration significantly increased from 27±0.63 to

106±22.7s respectively (Fig. 1C1). Together, these data suggest that myoclonic seizures had a

propensity to emerge during NREM and were of significantly shorter duration than all other

seizure phenotypes.

Recurrent generalized seizures in human *SYNGAP1* haploinsufficiency show affinity for

transition-states and clustering in NREM.

To evaluate the translational value of qEEG analysis of 24h EEGs in *Syngap1*$^{+/-}$ mice,

an IRB-approved study to acquire copies of 24h EEG recordings from children diagnosed with

pathogenic *SYNGAP1* mutation related ID has been recently initiated[25]. A 20h continuous EEG

which captured an entire cycle of overnight sleep in a 3 year old boy with a *de novo* intronic

variant in *SYNGAP1* associated with myoclonic and atonic absence seizures (Fig. 2) was

analyzed. The global seizures were bursts of 3-4Hz spike and slow waves that lasted ~0.5-3s

(Fig. 2A). The EEG was sleep-scored for wake, NREM, and REM using video and qEEG

spectral power analysis similar to our previously published protocols[17]. Ictal events occurred

both in wake and sleep. All ictal events observed during sleep occurred in NREM and

specifically at Wake-NREM and NREM-REM transitions (Fig. 2B). We identified a proclivity for

clustering of ictal events (i.e. inter-ictal durations <5min) that was predominant during NREM

(Fig. 2C & D). These analyses for the first time identify clustering of ictal-events during NREM

in a child with *SYNGAP1* related ID. The propensity of ictal events to occur at transition-states

is a novel finding and may be a biomarker for underlying mechanisms.

144   Progressive alterations in macro-sleep architecture

Sleep dysfunction is a common feature of NDDs and is also reported in patients with

SYNGAP1 mutations[12,21]. Quantitative assessments of the sleep problems in patients with

SYNGAP1 mutations are lacking; therefore, a robust assessment is warranted[25]. In this study,

the WT[+/+] sleep architecture was ultradian with 15.38±2.187 sleep cycles occurring over 24h

(Fig. 3A). WT[+/+] mice spent ~60% of the 24h period awake. During light vs. dark phase, WT[+/+]

mice spent ~50% and ~80% of the time awake (Fig. 3A - B2), respectively, demonstrating the

expected nocturnal predominance of activity in WT[+/+] mice.

Similar to WT[+/+] mice, P60[+/-] mice also spent significantly more time awake than asleep.

In contrast, the P120[+/-] mice lacked any significant differences between time spent awake and

asleep (Fig. 3B). Between groups, P60[+/-] and P120[+/-] mice spent less time awake than WT[+/+]

mice over the 24h recording. This decrease in wake duration (and subsequent increase in

sleep duration), was apparent during the light phase (Fig. 3B1). WT[+/+] mice did not

demonstrate any differences between time spent awake and asleep during the light phase. In

contrast, P60[+/-] and P120[+/-] mice spent significantly less time awake than asleep (Fig. 3B1).

Compared to WT[+/+] mice, P120[+/-] mice spent significantly less time awake.

During the dark phase, WT$^{+/+}$ mice spent significantly more time awake than asleep.

Nocturnal activity in WT$^{+/+}$ mice included nest building, exploration, and grooming behaviors.

All mice regardless of genotype or age spent a significantly greater amount of time awake than

asleep during the dark phase (Fig. 3B2). However, compared to WT$^{+/+}$ mice, P120$^{+/-}$ mice

spent significantly more time asleep in the dark phase (Fig. 3B2). These data suggest that

P120$^{+/-}$ mice spent significantly more time asleep than WT$^{+/+}$ mice.

In WT$^{+/+}$ mice, there were significantly fewer sleep/wake cycles (transitions from wake to

sleep) during the dark phase (9.56±1.98 cycles) compared to the light phase (22.67±9.56

cycles, Fig. 3C). During the dark phase, both P60$^{+/-}$ and P120$^{+/-}$ mice had more sleep/wake

cycles (16.33±3.06 and 16.22±3.08 cycles, respectively) than WT$^{+/+}$ mice (9.56±1.98 cycles),

however this increase was not significant. Sleep/wake cycles during the light phase were not

significantly different (Fig. 3C). In P60$^{+/-}$ and P120$^{+/-}$ mice, the difference between the numbers

of sleep/wake cycles during light vs. dark phase was lost due to these alterations (Fig. 3C).

WT$^{+/+}$ mice spent 6.99±0.3h in NREM and 1.76±0.2h in REM during the 24h recording.

Progressive alterations to macro-sleep architecture in P60$^{+/-}$ and P120$^{+/-}$ mice led to an

increase in the total duration of NREM (8.56±0.3 and 9.28±0.2h, respectively) and REM

(2.39±0.3 and 3.00±0.3h, respectively) when compared to WT$^{+/+}$ mice. The microarchitecture

of sleep (the percentages of NREM and REM that constitute sleep) in WT$^{+/+}$ mice was

80.27±1.7% NREM and 19.72±1.86% REM during the 24h recording (Fig. 3C1 & C2). P60$^{+/-}$

and P120$^{+/-}$ slept more than WT$^{+/+}$ mice, yet their microarchitecture of sleep was not

significantly different compared to WT$^{+/+}$ mice (Fig. 3C1 & C2). Therefore, all groups

maintained a sleep microarchitecture of ~80% NREM and ~ 20% REM.

IISs predominantly occurred in NREM sleep

Uncontrolled epilepsy is at least partially responsible for cognitive regression in children with epileptic encephalopathies[19,20]. IISs are reliable biomarkers of epileptogenesis that frequently occur on the EEGs of patients with epilepsy[26]. *Syngap1*[+/-] mice (n/n=8/8) presented IISs during both sleep when delta power and synchrony are high, and during wake when delta power and synchrony are relatively low (Fig. 4A).IISs significantly increased in frequency at P120[+/-] when compared to WT[+/+] mice (1-way ANOVA, $F_{2,19}=5.27$, P=0.015, p=0.013). This significant increase was driven by IISs that occurred during sleep (Fig. 4B) and was driven by light phase sleep (Fig. 4C). However, IISs during both light and dark phase sleep progressively increased in P120[+/-] mice (Fig. 4C). The frequency of IISs per hour of NREM and REM sleep was significantly greater in P120[+/-] mice (Fig. 4D). Therefore, taken together these data demonstrate that the overall number of IISs during the 24h recording period increased in P120[+/-] mice and these IISs mostly occurred during sleep in the light phase. Furthermore, even though the total duration of sleep increased significantly in P120[+/-] mice, (Fig. 3B) the rate of IIS occurrence per hour of NREM and REM sleep was significantly greater in P120[+/-] mice compared to WT[+/+] (Fig. 4D).

Emerging research utilizing intracranial recordings from epilepsy patients has demonstrated that regulation of brain activity occurs on long timescales[27]. Daily patterns of seizure occurrence demonstrate circadian rhythmicity and clustering organization[27]. In this study, we report the relationship between seizure occurrence and frequency distribution of IISs over 24h for all mice (Fig. 4E & 4F). Grouped data of IIS frequency averaged over 24h showed an escalation in IIS frequency at the transition between the dark and light phases associated with increased incidence of seizures in P120[+/-] mice which was not apparent in P60[+/-] mice.

Antagonizing AMPARs acutely with Perampanel (PMP) prevented seizures and resulted in a

non-significant decrease in IIS frequency (Supplementary Fig. 4). In summary, the data

suggest that IIS and seizure occurrence was greater during sleep in *Syngap1$^{+/-}$* mice.

Hyperactivity worsened with age

Video tracking software allows for high specificity and sensitivity when tracking mouse

motor activity over the 24h recordings. Total distance traveled by each mouse over the 24h

period (Fig. 5A1 - A3) was used to identify hyperactivity in *Syngap1$^{+/-}$* mice. Compared to

WT$^{+/+}$ mice (267.92±46.8m over 24h), P120$^{+/-}$ mice traveled greater distances (479.75±46.1m

over 24h) that was not detected in the P60$^{+/-}$ mice (288.22±32m over 24h; Fig. 5A2 & A3).

Since P120$^{+/-}$ mice spent more time asleep than WT$^{+/+}$ mice (Fig. 3), the distance traveled by

each mouse was normalized to the duration of time spent asleep. The running average of all

mice in each group highlighted increased motor activity of P120$^{+/-}$ mice at the end of the dark

phase when the majority of nesting behavior occurs[28] (Fig. 5B). The normalized distance

traveled by P120$^{+/-}$ mice was significantly greater than WT$^{+/+}$ and P60$^{+/-}$ mice (Fig. 5C). Taken

together these data suggest that P120$^{+/-}$ mice spent more time asleep but were hyperactive

when awake, as they travelled twice the distance compared to WT$^{+/+}$ mice over 24h. These

results are consistent with recent reports of home cage hyperactivity in male *Syngap1$^{+/-}$*

mice[29,30].

Activity-dependent theta modulation is impaired during wake in P120$^{+/-}$

Field potential recordings have observed cross–frequency coupling between theta and

gamma oscillations [31,32]. Specifically, gamma oscillations are nested within theta oscillations

[33,34]. In humans, theta-gamma coupling has shown high predictability for correct memory

retrieval [35–37]. Mouse studies utilizing optogenetics have demonstrated causal evidence for the

importance of theta-gamma coupling in cognitive function [38]. In WT$^{+/+}$ mice, theta and gamma

power demonstrated an inverse relationship during wake and sleep states (Supplementary Fig.

5). During wake, the theta-gamma ratio (TGR) was low in WT$^{+/+}$ mice, while during sleep the

TGR was high (Supplementary Fig. 5) [39,40]. This relationship between theta and gamma power

was lost in P120$^{+/-}$ as the TGR did not change according to wake or sleep state

(Supplementary Fig. 5C - D2). PMP rescued *Syngap1$^{+/-}$* TGR modulation to WT$^{+/+}$ levels.

To determine differences in theta and gamma power during transitions from stationary-

wake to active-wake, ten-minute epochs of transitions from stationary- to active-wake (five

minutes each) were analyzed (Fig. 6A). During stationary-wake, high theta and low gamma

power were apparent in WT$^{+/+}$ mice (Fig. 6A & B). However, during active-wake, WT$^{+/+}$ mice

transitioned to high gamma power and low theta power. This state-dependent (stationary- vs.

active-wake) transition was not apparent in P60$^{+/-}$ or P120$^{+/-}$ mice (Fig. 6B & C). Specifically,

theta power remained consistently high through transitions from stationary- and active-wake in

P120$^{+/-}$ mice. Gamma power was not significantly affected during these specific wake-state

transitions in P60$^{+/-}$ or P120$^{+/-}$ mice. PMP did not significantly modulate theta during transitions

from stationary-wake to active-wake (Fig. 6C).

Loss of behavioral-state homeostasis of cortical gamma

Gamma (35-50Hz) is low during sleep and high during wake[41–43]. High cognitive load

during wakefulness [41–43] is associated with increased gamma power. Accordingly, in WT$^{+/+}$

mice gamma power was high during wake and low during NREM sleep (Fig. 7A). P120$^{+/-}$ mice

did not demonstrate behavioral-state dependent gamma power modulation during Wake-

NREM transitions, as gamma power remained high during wake and NREM (Fig. 7B). PMP

rescued behavioral-state dependent gamma power modulation. PMP-treated *Syngap1$^{+/-}$* mice

had high gamma during wake and low gamma during NREM similar to WT$^{+/+}$ mice (Fig. 7C).

To analyze differences in Wake-NREM transitions over 24h, slopes for gamma power were

calculated for every Wake-NREM and NREM-Wake transition (see Methods). In WT$^{+/+}$ mice,

the slope of gamma power during Wake-NREM transitions was negative as gamma power

decreased during sleep compared to the previous wake. During NREM-Wake transitions, the

slope of gamma power was positive in WT$^{+/+}$ mice as gamma power increased from NREM to

wake (Fig. 7D1). The negative slopes of Wake-NREM transitions and the positive slopes of

NREM- Wake transitions were significantly different in WT$^{+/+}$ mice (Fig. 7D1). P120$^{+/-}$ mice

failed to show significant differences between wake and sleep transitions as gamma power

remained high during NREM (Fig. 7D2). PMP rescued the lack of significant differences

between the slopes of gamma power during Wake-NREM and NREM-Wake transitions in

Syngap1$^{+/-}$ mice (Fig. 7D3). The average gamma slopes for Wake-NREM and NREM-Wake

transitions were calculated to compare groups (Supplemental Fig. 6). Mean gamma slopes for

NREM-Wake and Wake-NREM in WT$^{+/+}$ mice were significantly different. In P120$^{+/-}$ mice

gamma slopes between NREM-Wake and Wake-NREM were not significantly different. PMP

rescued the lack of significant differences between NREM-Wake and Wake-NREM in P120$^{+/-}$

mice (Supplemental Fig. 6).

Frequency analysis of gamma power (35-50Hz) identified differences between NREM

and wake in WT$^{+/+}$ mice (Supplemental Fig. 8). Gamma power from 35-45Hz was significantly

higher during wake compared to NREM in WT$^{+/+}$ mice. In contrast, in P120$^{+/-}$ mice, wake and

NREM gamma power from 35-45Hz was not significantly different (Supplemental Fig. 8).

Therefore, *Syngap1*[+/-] mice demonstrated high gamma power during sleep and a loss of state-

dependent gamma modulation. Antagonizing AMPARs with PMP rescued state-dependent

gamma modulation.

During NREM the majority of brain activity is synchronous slow wave activity with low

gamma power [41–43]. In contrast, gamma power increases during the asynchronous activity of

REM[41–43]. In WT[+/+] mice, NREM-REM transitions underwent an increase in gamma, decrease

in theta, and decrease in delta power (Fig. 8A & B). P60[+/-] mice had a significantly greater

decrease in delta from NREM to REM compared to WT[+/+] mice. However, the significant

decrease in delta was not apparent in P120[+/-] mice. In contrast, a significantly lower rate of

percent change in gamma power during NREM-REM transitions was apparent in P120[+/-] mice.

Gamma power remained high during both NREM and REM (Fig. 8A & B) in P120[+/-] mice. The

impairment of gamma power modulation from NREM to REM worsened with age, and PMP

rescued the modulation of gamma power (Fig. 8C).

290 Deficits in PV+IN innervation in prefrontal cortex

Majority of perisomatic inhibition is mediated by parvalbumin interneurons (PV+INs)[41],

which plays a significant role in cortical circuits and in the modulation of cortical gamma

oscillations[44]. The novel finding of consistently high gamma power in *Syngap1*[+/-] mice directed

the investigation of their density distribution in the cortex, hippocampus, and prefrontal

cortices. Immunohistochemical (IHC) probes quantifying the PV+IN distributions did not show

significant differences in overall densities in the cortex and hippocampus. Layer-specific

analyses detected significantly higher numbers of PV+INs in the SR and SO regions of the

CA3 and CA1 regions respectively (*t-test*, t_{14}=-2.479, p=0.027, and t_{14}=-2.751, p=0.016

respectively). Quantification of overall counts of PV+ puncta identified significant deficits in the

DP cortex in *Syngap1*[+/-] mice (Fig. 9A & B) but not in the PrL cortex (data not shown).

Additionally, a significant decrease in perisomatic PV+ puncta was identified in the DP cortex

of *Syngap1*[+/-] mice (Fig. 9B). No significant differences in mean GluA2 fluorescence intensity

on PV+INs or non-PV+ neurons were detected in the DP cortex (Fig. 9C1 & 9C2). These

findings indicate that deficits in PV+ innervation were region-specific in *Syngap1*[+/-] mice even

within the prefrontal cortex.

The significant difference between counts of PV+ puncta in the DP cortex (Fig. 9)

warranted the analysis of PV+ puncta counts in other regions of interest (ROIs). Unlike the

prefrontal cortex, no significant differences in PV+ puncta count were noted in either the

sensory or motor cortices (Fig. 10A2 & B2). A layer-specific investigation of the barrel cortex

(BCX) and motor cortex (MCX) demonstrated that PV+ puncta counts were not significantly

different between layers 2-3 and 5-6 in the BCX and MCX in both the total counts within the

whole ROIs and the somas of non-PV+ cells (Fig. 10A1, A2 & Fig. 10B1, B2). In the

hippocampus, PV+ puncta counts were not significantly different in CA1 or CA3 (Fig. 10C1).

Normalized PV+ puncta counts were not significantly different between the SP of WT[+/+] and

P120[+/-] mice in CA1 and CA3 (data not shown, *t-test*, t_{22}=0.2861, p=0.7775, and t_{31}=0.8392,

p=0.4078 respectively). SO and SR were also examined, and PV+ puncta counts were not

significantly different in CA1 and CA3 in the SO (data not shown, *t-test*, t_{22}=1.125, p=0.2725,

and t_{31}=0.5407, p=0.5926 respectively) and the SR (data not shown, *t-test*, t_{22}=0.8627,

p=0.3976, and t_{30}=0.3402, p=0.7361 respectively) between WT[+/+] and P120[+/-] mice. Since all

Syngap1[+/-] mice showed epileptiform activity, the hippocampi were investigated for mossy fiber

sprouting. No evidence for mossy fiber sprouting was detected in this model (see

Supplemental Fig. 9). In summary, PV+IN deficits in innervation were only observed in the DP

cortex.

Increased GluA2 expression on PV+ soma in somatosensory but not motor cortex

A rich diversity of GABAergic neurons shape the spatiotemporal dynamics of cortical

circuit outputs by elegant inhibitory control mechanisms [45,46]. Identification of the dysregulated

cortical gamma and its acute rescue by PMP directed the quantification of AMPA-GluA2 on

PV+INs and non-PV+ neurons in these regions. Compared to neurons, PV+INs are known to

have a lower expression profile for the GluA2 subunit[47]. The findings (Fig. 10) highlighted a

region-specific alteration for the significant upregulation of GluA2 in both PV+INs and non-PV+

neurons (for GluA2 antibody verification see Supplemental Fig. 9). The sensory and motor

cortices in the *Syngap1*[+/-] mice were distinctly different. PV+INs in the both 2-3 and 5-6 cortical

layers of the BCX showed significant upregulation, (Fig. 10A2) which was absent in the MCX

(Fig. 10B2). Significant GluA2 upregulation in non-PV+ sampled cells in both ROIs was evident

in layer 2-3 of the BCX and layer 5-6 of the MCX (Fig. 10B2). The mean GluA2 fluorescence

intensity on PV+INs was also significantly higher in the CA3 but not in the CA1 (Fig. 10C2). For

non-PV+ somas, the mean GluA2 fluorescence intensity was significantly lower in the CA1 but

significantly higher in the CA3 neurons (Fig. 10C2). Therefore, the results identified a

significant dysregulation of GluA2 expression in PV+INs, a population of INs known to

contribute significantly in cortical gamma oscillations [44]. The regional differences highlight the

complexity of the outcomes *Syngap1*[+/-] haploinsufficiency can exert in different circuits.

Additionally, a higher mean GluA2 fluorescence intensity was observed in the MCX

compared to the BCX. In WT$^{+/+}$ mice, the mean GluA2 fluorescence intensity in MCX neurons

was significantly higher than BCX neurons both for layers 2/3 and 5/6 (Fig. 10A2 vs. 10B2, *t-*

test, t_{117}=10.6, p<0.0001, and t_{118}=12.94, p<0.0001 respectively). This relationship was

preserved in P120$^{+/-}$ mice, as the mean GluA2 fluorescence intensity in MCX neurons was also

significantly higher than in BCX neurons in both layers 2/3 and 5/6 (Fig 10A2 vs. 10B2, *t-test*,

t_{119}=11.31, p<0.0001, and t_{126}=14.18, p<0.0001 respectively).

Discussion

In this study, we used long-term EEG monitoring and qEEG protocols developed

specifically to identify biomarkers underlying epileptogenesis in mouse models of acquired and

genetic NDDs[17,18,48]. The results showed a progressive worsening in epileptogenesis

associated with significant alterations in macro- and micro-sleep architecture, IIS frequency,

and impaired behavioral-state dependent cortical activity during wake- and sleep-state

transitions. The IISs and seizures in the *Syngap1*$^{+/-}$ mice predominantly arise during NREM

sleep. In the human overnight EEG analysis, identification of ictal clusters during NREM were

a novel finding that support the translational validity of this *Syngap1*$^{+/-}$ mouse model. These

results highlight the similarities in seizure phenotypes in this mouse model to phenotypes

described in the clinic. Seizures are highly prevalent in patients with MRD5, therefore these

findings provide a valuable contribution to help further understand the significance of *Syngap1*

in epileptogenesis. While multiple groups have investigated the intellectual disability in

Syngap1 heterozygous mice, this study is the first thorough analysis of the epilepsy in a model

of *Syngap1* haploinsufficiency. Currently, it is unknown if other models of *Syngap1*

haploinsufficiency that differ by mutation site or mutation strategy (i.e. germ-line vs. Cre-

recombinase) have recurrent spontaneous seizures.

Spectral power analysis of the mouse EEGs quantified the cortical gamma homeostasis

during behavioral-state transitions and identified a significant disruption in *Syngap1*[+/-] mice.

The rescue of this biomarker with acute dosing by PMP identified a critical role of AMPARs in

gamma homeostasis during such transitions. Given the novel finding of acute rescue of

gamma homeostasis by an AMPAR antagonist PMP and previous work implicating severe

disruption of gamma oscillations associated with induced increase of GluA2 expression in

INs[49]; the Ca^{2+} impermeable AMPA subunit became a primary focus. This approach identified

excessive GluA2-AMPAR expression on the somas of PV+INs that was location- and circuit-

specific indicating a significant role of PV+INs in MRD5 pathology. This study also found

abnormal cortical activity in theta frequencies during transitions from stationary-wake to active-

wake. As theta-gamma coupling is essential for memory retrieval, abnormal theta and gamma

homeostasis pose a potential relevant biomarker for memory impairment that has been

previously verified in human subjects and optogenetic rodent studies [35–38].

To date, there is no clear correlation between *SYNGAP1* mutation location and

phenotypic severity. However, mutations in exons 8-15 seem to be associated with seizures

more pharmacoresistant than with mutations in exons 4-5 [11]. In this preclinical model, deletions

in *Syngap1* exons 8 and 9 resulted in half of the *Syngap1*[+/-] mice presenting with a seizure

during the 24 recording period indicating a high seizure frequency. However, 100% of the

Syngap1[+/-] mice presented progressively increasing IIS frequency and sleep dysfunction.

These findings provide a potential mechanism for the reported decline of cognitive ability

during the developmental trajectory of MRD5 into adulthood [21].

Wakefulness is associated with net-synaptic potentiation, whereas sleep favors global

synaptic depression thereby preserving an overall balance of synaptic strength [50]. AMPAR

levels are high during wakefulness and low during sleep. Therefore, consistently high AMPAR

levels in PV+INs would perturb this balance toward impaired signal processing and learning

disability. qEEG analyses revealed significant loss in cortical gamma homeostasis associated

with behavioral-state transitions both in wake and sleep that could impede cortical information

processing. Gamma oscillations [51,52] are known to heavily depend upon AMPAR kinetics in

interneurons, especially for long-range synchrony. We showed that acute treatment with PMP,

an AMPAR antagonist, significantly rescued many of the identified EEG biomarkers along with

the disrupted cortical gamma oscillations. IHC analyses observed and quantified a significant

increase in GluA2-AMPAR expression onto the somas of cortical and hippocampal PV+INs in

a region-specific manner. The significant rescue of the cortical gamma homeostasis by acute

dosing of PMP when compared to the weaker effect on IIS suppression may indicate a novel

role for PMP unrelated to its FDA approved anti-seizure role. Additionally, the findings indicate

the necessity to develop GluA2-AMPAR selective antagonists, which currently do not exist.

These analyses also identified poor PV+ interneuron innervation in the prefrontal cortex that

was not detected in the other ROIs evaluated both in the cortex and hippocampus. This finding

in the prefrontal cortex is supported by previously published reports [53].

Increasingly, brain oscillations are being used to understand complex neuropsychiatric

disorders. Gamma (35–50Hz) oscillations have warranted special attention due to their

association with higher order cognitive processes including sensory processing, attention,

working memory, and executive functioning. Activation of GABAergic interneuron networks

has been shown to produce gamma oscillations (~40Hz) in both the hippocampal and

neocortical networks. Given the critical role of PV+INs in cortical gamma oscillations [44,54] and

the role of AMPARs in homeostatic synaptic plasticity [55], the loss of gamma homeostasis

identified for *Syngap1* haploinsufficiency during periods commonly associated with intense

synaptic plasticity [56] provide novel insights into the associated intellectual disabilities.

GluA2 subunits confer Ca^{2+} impermeability to the AMPAR, a slow EPSP in excitatory

neurons, and are expressed at low levels in GABAergic interneurons [51]. AMPA-mediated

currents rise and decay faster in interneurons partly because of these differences in subunit

profiles [57,58]. A genetically altered GAD-GluR-B (GluA2) mouse expressing high levels of the

GluA2 subunit in GAD+ cells exhibited a severe disruption in gamma oscillations between

spatially separated sites and significant changes in interneurons firing patterns[51]. PV+INs

receive convergent excitatory input from principal neurons, and inhibitory input primarily from

other PV+INs. Experiments that have causally tested stimulation of PV+ cells *in vivo* in the

BCX by ontogenetic manipulation selectively amplified gamma oscillations[59]. Importantly, this

activation suppressed the sensory processing in nearby excitatory neurons within the BCX.

Findings of increased GluA2 AMPARs on PV+ soma both in layer 2/3 and 5/6 of the BCX

would predict similar suppression of sensory processing in the BCX. Findings supporting this

hypothesis have recently been documented in the BCX of a mouse model of *Syngap1*

haploinsufficiency[60]. GluA2 data for PV+INs in the BCX but not in the MCX, highlight the

divergent circuit- and location-specific effects of *Syngap1* haploinsufficiency. Similarly

significantly higher mean GluA2 fluorescence intensity in layer 5-6 motor output neurons in the

MCX highlight location-specific effects for non-PV+ neurons. These data also indicate that

PV+IN hyperexcitability or lack thereof may be one of the underlying factors that determine

circuit outcomes to motor vs. sensory inputs. An increase of GluA2 expression in PV+INs in

the BCX could implicate recruitment of excessive PV+IN mediated inhibition resulting in

cortical hyperexcitability via rebound excitation. This hypothesis may underlie the clinical

reports of reflex seizures in MRD5 emerging during activities associated with increased cortical

gamma synchronicity [12,61].

Morphology of PV+INs has been identified as unipolar vs. multipolar [62] and little is

known about the differences in AMPA-mediated currents in these subtypes. GluA2-lacking

AMPARs promote anti-Hebbian long-term plasticity [63] which is critical for projection

interneurons that functionally connect spatially distant circuits during development. Increase of

GluA2-AMPARs in PV+IN soma could be one novel mechanism underlying autistic-like

behaviors, ID, and seizures in *SYNGAP1* haploinsufficiency. Little is known of the

developmental and regional expression profiles of SYNGAP1 isoforms, however it is known is

that they can exert opposing effects on synaptic strength[64]. Therefore, region-specific and

isoform-specific haploinsufficiency could confer vastly disparate and sometimes opposing

circuit-specific outcomes within the same brain. The findings of this study and previous reports

[65] support this conclusion.

Synaptic strength is an important factor in learning and memory that is regulated by

complex biochemical machinery at the PSD. There are currently no evidence-based clinical

interventions that can directly modulate SYNGAP1 function. One of the prominent roles of

SYNGAP1 is to regulate synaptic plasticity, and this process is heavily involved in both

epileptogenesis and sleep homeostasis. Therefore, using validated qEEG biomarkers for both

epileptogenesis and circuit dysfunction in preclinical models could be a sound translational

approach. Investigation of circuit function in intact brains as they transition from active

exploratory, inactive-wake, and REM/NREM sleep states is an exciting frontier. These are

behavioral-state transitions that are most reliant upon synaptic plasticity.

Methods

All animal care and procedures were carried out in accordance with the

recommendations in the Guide for Care and Use of Laboratory Animals of the National

Institutes of Health. All protocols used in this study were approved by the Committee on the

Ethics of Animals Experiments of the Johns Hopkins University. In *Syngap1*[+/-] mice, exons 7

and 8 (C2 domain) were replaced with a neomycin resistance gene in the opposite direction to

eliminate all possible slice variants[24]. All mice were single-housed at P46 (1 week before

electrode implantation surgeries) in polycarbonate cages with food and water provided *ad*

libitum, on a 12h light-dark cycle. All efforts were made to minimize animal suffering and the

total number of animals used.

EEG surgery and 24h video-EEG/EMG recording

Surgical procedures implemented in this study are previously described [66]. All surgical

procedures were performed under isoflurane anesthesia (4%-2.5%). 7 days (±1 day) before

recording, mice were surgically implanted with subdural EEG electrodes and suprascapular

EMG electrodes (Fig. 1A & C). EEG electrodes utilized coordinates from bregma for consistent

placement [18].

After recovering from electrode implantation surgery (7±1 day) mice were placed in a

recording chamber with food and water provided *ad libitum*. WT[+/+] and *Syngap1*[+/-] mice

underwent 24h vEEG/EMG at younger (P60) and older (P90-120) ages (see Supplementary

Fig. 1 for sample size). All 24h vEEG/EMG were recorded using Sirenia Acquisition software

(Pinnacle Technology Inc., Lawrence, KS, USA). Each chamber had a top-view infrared

camera and a tethered pre-amplifier connected to a commutator. Connecting the commutator

to the implanted head cap allowed for unrestricted movement within the recording chamber

during recording. EEG recordings were acquired at 400Hz with a 100 gain, band passed from

0.5 to 50Hz [67]. Before each recording, all mice received a new piece of nesting material and a

small piece of nesting from their home-cage. Data from all wild type recordings at P60, 90 and

120 were pooled together (see Supplementary Fig. 4) for analysis, P90 *Syngap1*$^{+/-}$ (small

sample size; n=2) recordings were pooled with P120$^{+/-}$ mouse recordings. At P121, half of the

mice in the study underwent another 24h recording after being administered PMP (2mg/kg;

intraperitoneal) at 10am and again at 6pm in the 24h recording period.

Hypnogram and EEG power analysis

EEG data were manually scored for wake, NREM, and REM EEG states in 10s epochs by a

scorer blinded to genotype and sex. All EEG artifacts were identified after EEG and video

confirmation. Spectral power data for these identified epochs was excluded from further

analyses to avoid contamination of data from background EEG. qEEG spectral analysis

utilized delta (0.5-4.0Hz), theta (5.5-8.0Hz), alpha (8.0-13.0Hz), beta (13.0-30Hz), and gamma

(35-50Hz) frequencies. The data underwent fast Fourier transform (FFT) and were exported for

all respective frequencies from an automated analysis module within Sirenia Sleep Pro

(Pinnacle Technology Inc., Lawrence, KS, USA).

Seizure and IIS scoring

Spontaneous seizures were identified by manual review of all 24h EEGs. Identified EEG

seizure activity was then further classified by phenotype (review of synchronous video and

EMG) and duration. EEG epochs containing seizure activity were also excluded from spectral

power analyses of background EEG and spike frequency analysis. IISs were visually analyzed

on raw EEG using conservative parameters obtained from obvious pre- and post-ictal spikes

within the same dataset. Each subject's mean EEG amplitude during wake was used to

determine the minimum absolute value for IIS identification. 1X of the absolute value was the

criterion for the positive inflection, and 2X of the absolute value was the criterion for the

negative inflection. If the positive and negative inflection criteria were met, the synchronous

video was analyzed to eliminate artifact contamination (e.g. scratching, grooming or drinking

water). IISs were manually scored for every 5s EEG epoch to enhance the resolution of IIS

frequency detection.

516   Gamma AUC and slope

Spectral power data for each mouse was exported for area under the curve (AUC) analysis

using trapezoidal summations[48,66]. FFT generated spectral power data for each mouse

underwent AUC analysis in R statistical software with automated code.

To analyze differences in gamma power transitions during changes in behavioral states

(i.e.; from sleep to wake and wake to sleep), only behavioral states that lasted ≥6min were

used. Two one-minute averages of gamma power AUC were calculated, the first occurring

right before the behavioral-state transition and the second occurring 5min after the behavioral-

state transition. The slope between these two gamma power averages was then calculated by

automated code for each behavioral-state transition in R statistical software[66].

$$slope = \frac{y_2 - y_1}{x_2 - x_1}$$

527 Where y_2 = gamma power average after transition, y_1 = gamma power average before

528 transition, x_2 = 6min, and x_1 = 1min.

530 Theta-gamma ratio and activity dependent theta-gamma modulation

Theta-Gamma Ratio (TGR) was calculated for every 10s epoch for the 24h recording

period. To analyze differences between wake and sleep TGR changes between WT[+/+] and

Syngap1[+/-] mice, the percent change in TGR from wake to sleep was calculated. Wake activity

dependent theta-gamma modulation was analyzed by pulling transitions of equal duration

(10min total) from stationary-wake to active-wake (5 min each; verified by EMG and video).

Respective powers were summed according to their activity state, and then computed for

percent change. NREM-REM dependent theta-gamma modulation was analyzed by examining

NREM-REM transitions of equal duration (100s total). Respective powers were summed

according to their sleep state and percent change was calculated.

541 Analysis of human EEG recording

Continuous overnight EEGs are being acquired with parent consent under approved

IRB protocol [25].   The raw EEG recording analyzed here was imported into Brainstorm

(Brainstorm, CA, USA). The EEG channels F4, C4 and O2 were re-referenced to A1 or A2

and exported in the European Data File (EDF) format, de-identified and assigned an ID

number. A bandpass filter of 0-80Hz was used. Automated spectral analysis was conducted

(delta 0.5–4Hz, theta 5.5–8.5Hz, alpha 8–13Hz, beta 13–30Hz and gamma 35–80Hz) using

the Sirenia Sleep score module and spectral powers were calculated for every 10s epoch of

the EEG. Each epoch was manually scored as Wake or Sleep (NREM or REM) using video

and delta power during slow-wave sleep in NREM using a previously published protocol[17].

These analyses were performed blind to the time points of the ictal events on the raw EEG.

Ictal events were identified on original raw EEG recorded as a 24-channel digital video EEG

using an international 10-20 system of electrode placement. Eye monitors and one channel

EKG recordings were also available for review. The raster plot for identified ictal events on this

20h long recording was then superimposed over the hypnogram generated for the same EEG

recording.

PV and GluA2 immunofluorescence acquisition and quantification

At P127 (P120+ 7 days) mice were anesthetized with chloral hydrate (300mg/ml; IP)

before transcardiac perfusion with ice-cold PBS followed by formalin (10%). Brains were then

cryoprotected and stored at -80°C before cryosectioning. All coronal sections were cut at 14um

and mounted onto Permafrost™ microscope slides. The primary antibodies used were anti-

GluA2 (ThermoFisher Scientific; RRID: AB_2533058; diluted 1:100), anti-parvalbumin (Swant;

RRID: AB_2313848; diluted 1:5000), and anti-ZnT3 (Synaptic Systems; RRID: AB_2189664;

diluted 1:100). Hoechst 33342 fluorescent dye (ThermoFisher Scientific; Cat. H3570) was used

to label cell nuclei. The secondary antibodies used were: donkey anti-rabbit Alexa 488

(ThermoFisher Scientific; RRID: AB_2535792; diluted 1:500), and goat anti-mouse Alexa 594

(ThermoFisher Scientific; RRID: AB_2556549; diluted 1:500). For GluA2 antibody verification,

P60 GluA2 KO mice (Richard Huganir, Johns Hopkins University) and age matched littermates

were perfused with PBS and 4% PFA then sectioned at 50μm (Supplemental Fig. 9).

Fluorescent images were acquired with Axiovision Software 4.6 and Apotome® (Carl

Zeiss, Jena, Germany). Images from all regions of interest (ROIs): hippocampal regions CA1

and CA3, MCX, BCX, and prefrontal cortex regions DP and PrL, were acquired at both 10X

and 40X magnifications for all ROIs. The 10X images were acquired as Z-stack mosaics of

four 3μm steps, while 40X images were acquired as Z-stack mosaics of four 1μm steps. For

10X and 40X images, two images per animal were analyzed from each ROI. All ZnT3 and

GluA2 KO stained hippocampi were imaged as 10X mosaics same as above. All data analyses

were performed using the ImageJ software[68]. All data were acquired and analyzed by a scorer

blinded to both genotype and sex.

All Z-stack images were fused with equal number of step sizes (four 3μm steps for 10X

images and four 1μm steps for 40X images) and split by RGB channel in ImageJ. PV+ Alexa

488 immunofluorescence identified both PV+IN cell bodies and PV+ puncta representing their

presynaptic terminals innervating non-PV cells in the same ROIs. PV+ puncta around two

randomly selected non-PV+ cells per ROI were analyzed. In this study, non-PV+ cells

represented neurons that were selected by the large size of their nuclei with multiple PV+

puncta representing PV innervation. A low-pass filter was used to exclude non-specific labeling

of structures larger than 10 pixels (such as blood vessels and RBC), and a high-pass filter was

used to exclude structures smaller than 2 pixels (such as nonspecific fluoropore staining on the

surface of the brain section). The filtered PV immunofluorescence was then converted to a

binary image and PV+ puncta counts were quantified using ImageJ. Total PV+ puncta counts

were quantified using ImageJ. For the stratum oriens (SO), stratum pyramidale (SP), and

stratum radiatum (SR) layers were delineated and quantified individually for hippocampal

images. PV+ puncta were expected to be higher in regions of greater neuron density, such as

the SP layer in the CA1 and CA3. Therefore, the PV+ puncta counts were additionally

normalized by each sub-ROI area of SP, SO, and SR for every hippocampal image. The

somas of non-PV+ cells were circumscribed using an elliptical selection tool to demarcate the

neuronal cell body, and puncta immediately around the soma were manually counted. Counts

of PV+ puncta on non-PV+ neurons were normalized to their respective cell areas.

To quantify GluA2 expression on PV+INs, mean somatic GluA2 fluorescence (Alexa

594) intensity was quantified for all PV+INs within each ROI. PV+INs were sub-classified by

their location in layers 2/3 vs. 5/6 in MCX and BCX images. In hippocampal images, PV+INs

were classified by their location in CA1 vs. CA3 and by layer specificity in the SO, SP, or SR.

Two randomly selected non-PV+ cells were quantified per ROI in the MCX and BCX. In

hippocampal images, two non-PV+ cells were quantified only for the SP. Mean somatic GluA2

fluorescence intensity was quantified as the average pixel intensity (0-255) around each cell

body.

Statistical analyses

Statistical tests were performed using SPSS24 (IBM, Armonk, NY, USA) and Prism 7.0

(GraphPad Software, La Jolla, CA, USA). All 1-way and 2-way ANOVAs were performed with

Bonferroni's post-hoc corrections. Independent *t*-tests were two-way, and all data reported as

means were ± 1 S.E.M. Analysis of seizure durations between P60$^{+/-}$ and P120$^{+/-}$ was an

independent two-way *t*-test in Prism 7.0. Analyses of macro and micro-sleep architecture were

2-way ANOVAs with Bonferroni post-hoc corrections in Prism 7.0. IIS frequency and motor

hyperactivity analyses were 1-way ANOVAs with Bonferroni post-hoc corrections. The running

weighted averages of IIS frequency and hyperactivity were plotted using the Lowess method in

Prism 7.0.

Theta power during transitions from inactive to active wake were analyzed as 1-way

ANOVAs and representative traces of theta-gamma modulation were plotted using the Lowess

method in Prism 7.0. Representative TGRs underwent a min-max normalization and were

plotted in Prism 7.0. Average % increase in TGR from wake to sleep were analyzed as a 1-

way ANOVA with Bonferroni post-hoc correction in Prism 7.0.

Differences within NREM-wake and wake-NREM gamma slopes for each group were

analyzed as independent two-way *t*-test in Prism 7.0. To identify differences between NREM-

wake and wake-NREM between groups, the average gamma slopes for each transition state

were analyzed as a 2-way ANOVA with Bonferroni post-hoc correction in Prism 7.0. Gamma

frequency plots of Wake, REM, and NREM were plotted using the Lowess method in Prism

7.0. Wake and NREM spectral frequency were analyzed as two-way independent *t*-tests in

Prism 7.0.

Mean PV+IN puncta counts and mean GluA2 florescence intensity from 40X Z-stack

images were analyzed in Prism 7.0 as two-way independent *t*-tests.

Acknowledgements:

We thank the families who have consented to participate in the post-hoc analyses of overnight EEGs recorded in their children with identified pathogenic *SYNGAP1* mutations. Research reported in this publication was supported by the Eunice Kennedy Shriver National Institute of Child Health and Human Development of the National Institutes of Health under Award Number R01HD090884 (SDK). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Author Contributions

S.D.K. designed the experiments. S.D.K and R.L.H conceptualized the project. S.D.K. and B.J.S performed experiments. Y.A. managed mouse colonies. B.J.S., S.A., and P.A.K. analyzed data. S.D.K. and B.J.S interpreted the data. S.D.K and B.J.S wrote the manuscript. S.D.K., B.J.S., and P.A.K. edited and reviewed the manuscript.

Competing Interests

The authors declare no competing interests.

Corresponding author

Correspondence to S. D. Kadam.

Fig. 1. *Syngap1$^{+/-}$* **mice are epileptic with multiple seizure phenotypes (A)** Representative EEG trace (0.5-50Hz) of a spontaneous myoclonic seizure at P60 during NREM sleep. Expanded timescale of recording showed the prototypical time-locked myoclonic jerks that were recorded on EMG (see Supplementary Video 1). **(B)** Representative spontaneous tonic-clonic seizure at P120 during NREM sleep. Expanded timescale of recording showed tonic-clonic discharges on EEG and EMG at beginning of the seizure which progressed to burst discharges on EEG associated with tonic spasm on EMG. **(C1)** The average duration of seizures significantly increased from P60 to P120 (*t-test*, t_{14}=2.307, p=0.037). **(C2)** At P60 all seizures were myoclonic and the majority occurred during NREM (4/5). **(C3)** At P120, all myoclonic (3/11) and tonic-clonic (1/11) seizures occurred during NREM, whereas all electrographic seizures (see Supplementary Figure 2) occurred during wake (7/11).

Fig. 2. Patient with pathogenic *SYNGAP1* mutation exhibiting multiple ictal events on EEG over a 20h period (A) Representative ~3Hz short duration epileptiform events during wake and NREM sleep were identical. **(B)** Hypnogram for the 20h recording. All ictal events were identified and overlaid on the hypnogram. Obvious clustering of events was noted during NREM sleep. **(C)** 5min EEG trace showed four clustered ictal events (red arrowheads) occurring within a 5min period during the last NREM cycle before wake. Ictal events occurring at transitions between wake-sleep and REM-NREM were also noted. **(D)** Bar graphs grouping clustered vs. non-clustered ictal events for wake and NREM showed proclivity for clustering during NREM. Seizure rates were normalized per hour of EEG recording for the raster were: wake=1.27/h; NREM=2.68/h and REM=0/h for this recording.

Fig. 3. Progressive alterations in macro-sleep architecture (A) 24h hypnograms of ultradian rhythms over the diurnal light/dark phase in WT$^{+/+}$, P60$^{+/-}$, and P120$^{+/-}$ mice. Regardless of age, all WT$^{+/+}$ mice had consistent macro- and micro-sleep architectures (Supplemental Fig. 3). **(B)** All WT$^{+/+}$ mice spent significantly more time awake than asleep over the 24h period (WT$^{+/+}$ Wake vs. WT$^{+/+}$ Sleep: 2-way ANOVA, $F_{2,42}$=42.03, P<0.0001, p<0.0001). P60$^{+/-}$ mice also spent significantly more time awake than asleep (P60$^{+/-}$ Wake vs. P60$^{+/-}$ Sleep: 2-way ANOVA, $F_{2,42}$=42.03, P<0.0001, p=0.002), however P120$^{+/-}$ mice lacked any significant differences between wake and sleep durations over the 24h period. Between groups, time spent awake significantly decreased in P60$^{+/-}$ (WT$^{+/+}$ Wake vs. P60$^{+/-}$ Wake: 2-way ANOVA, $F_{2,42}$=42.03, P <0.0001, p=0.003) and P120$^{+/-}$ mice (WT$^{+/+}$ Wake vs. P120$^{+/-}$ Wake: 2-way ANOVA, $F_{2,42}$=42.03, P <0.0001, p<0.0001). **(B1)** During the 12h light phase both P60$^{+/-}$ and P120$^{+/-}$ mice spent more time asleep than awake (P60$^{+/-}$ Light Wake vs. P60$^{+/-}$ Light Sleep: 2-way ANOVA, $F_{2,42}$=29.02, P<0.0001, p<0.0001; P120$^{+/-}$ Light Wake vs. P120$^{+/-}$ Light Sleep, p<0.0001). During the light phase, the amount of time P120$^{+/-}$ mice spent awake was significantly less than WT$^{+/+}$ (WT$^{+/+}$ Light Wake vs. P120$^{+/-}$ Light Wake: $F_{2,42}$=29.02, P <0.0001, p<0.0001). **(B2)** WT$^{+/+}$, P60$^{+/-}$, and P120$^{+/-}$ mice all spent significantly more time awake than asleep during the dark phase (WT$^{+/+}$ Dark Wake vs. WT$^{+/+}$ Dark Sleep: 2-way ANOVA, $F_{2,42}$=10.95, P=0.0001, p<0.0001; P60$^{+/-}$ Dark Wake vs. P60$^{+/-}$ Dark Sleep: p<0.0001, P120$^{+/-}$ Dark Wake vs. P120$^{+/-}$ Dark Sleep: p<0.0001). Between groups, P120$^{+/-}$

mice spent significantly less time awake during the dark phase than WT$^{+/+}$ (WT$^{+/+}$ Dark Wake
vs. P120$^{+/-}$ Dark Wake: $p<0.0001$, P120$^{+/-}$ Dark Wake vs. P120$^{+/-}$ Dark Sleep: $p=0.0256$). **(C)**
WT$^{+/+}$ mice had significantly fewer cycles (transitions between sleep and wake) during the dark
phase than the light phase (WT$^{+/+}$ Light cycles vs. WT$^{+/+}$ dark cycles: 2-way ANOVA,
$F_{2,42}=1.547$, $P=0.0001$, $p=0.0066$. **(C1 & C2)** This significant difference in the number of
sleep/wake cycles detected in WT$^{+/+}$ was lost in P60$^{+/-}$ and P120$^{+/-}$ mice. The micro-
architecture of both NREM and REM cycles during sleep was not significantly different
between groups. ($p<0.05$ *, $p<0.01$ **; P120 $p<0.001$ ***; $p<0.05$ #, $p<0.01$ ##; $p<0.001$ ###,
post-hoc Bonferroni).
715

Fig. 4. Interictal spiking during NREM (A) Representative 30s EEG trace (0.5-50Hz) during
wake showing IISs (n=2) associated with low amplitude background delta power (0.5-4Hz).
During NREM sleep, a representative 30s EEG trace showing IISs (n=5) with high amplitude
background delta power (0.5-4Hz) associated with slow-wave-sleep. **(B)** Number of spikes
during sleep significantly increased in P120$^{+/-}$ compared to WT$^{+/+}$ mice (WT$^{+/+}$ Sleep Spikes vs.
P120$^{+/-}$ Sleep Spikes: 1-way ANOVA, $F_{2,\ 19}=5.36$, $P=0.014$, $p=0.012$) **(C)** Number of spikes
significantly increased during both light and dark sleep in P120$^{+/-}$ mice (WT$^{+/+}$ vs. P120$^{+/}$: 1-way
ANOVA, Light Sleep Spikes: $F_{2,19}=5.386$, $P=0.014$, $p=0.012$, Dark Sleep Spikes:$F_{2,19}=4.838$,
$P=0.02$, $p=0.018$). **(D)** IISs significantly increased with age during NREM (WT$^{+/+}$ NREM spikes
vs. P120$^{+/-}$ NREM spikes: 1-way ANOVA, $F_{2,\ 19}=5.26$, $P=0.015$, $p=0.013$) and REM (WT$^{+/+}$ vs.
P120$^{+/}$:1-way ANOVA, $F_{2,\ 19}=3.458$, $P=0.052$, $p=0.050$). **(E & F)** Running averages of IISs
(black traces) and seizure frequency (red lines) over 24h. ($p<0.05$ *, $p<0.01$ **; $p<0.001$ ***;
post-hoc Bonferroni).
729

Fig. 5. Progressive increase of motor activity in _Syngap1$^{+/-}$_ mice (A1) Image from the
infrared top-view camera used for motion tracking. **(A2 & A3)** Representative traces of WT$^{+/+}$
and P120$^{+/-}$ activity over 24h within individual recording chambers. **(B)** Running averages of
distance traveled over 24h by all WT$^{+/+}$, P60$^{+/-}$, and P120$^{+/-}$ mice. **(C)** P120$^{+/-}$ showed
significantly higher motor activity than WT$^{+/+}$ and P60$^{+/-}$ during the exploration and nesting
phases (WT$^{+/+}$ vs. P120$^{+/-}$: 1-Way ANOVA; $F_{2,20}=20.18$ $P<0.001$ $p<0.001$, P120$^{+/-}$ vs. P60$^{+/-}$:
$p<0.001$). ($p<0.05$ *, $p<0.01$ **; $p<0.001$ ***; post-hoc Bonferroni).
737

Fig. 6. Impaired theta power during transitions from stationary-wake to active-wake (A)
EEG and EMG spectral power heat maps of activity-dependent transitions from stationary-
wake to active-wake in WT$^{+/+}$ mice **(B)** Representative traces of theta-gamma modulation in
WT$^{+/+}$ and P120$^{+/-}$ mice during transitions from stationary-wake to active-wake (for further
analyses of theta-gamma interactions see supplemental Fig. 5). **(C)** The lack of theta
modulation during transitions progressively worsened in P120$^{+/-}$ (WT$^{+/+}$ theta vs. P120$^{+/-}$ theta
% change: 1-way ANOVA, $F_{2,19}=12.123$, $P<0.0001$, $p<0.0001$). PMP administration did not
significantly modulate theta during transitions from stationary-wake to active-wake (WT$^{+/+}$ theta
vs. PMP$^{+/-}$ theta % change: 1-way ANOVA, $F_{2,19}=12.123$, $P<0.0001$, $p=0.017$). ($p<0.05$ *,
$p<0.01$ **; $p<0.001$ ***; post-hoc Bonferroni).
748

Fig. 7. Gamma power modulation during transitions between Wake-NREM and NREM-
Wake (A) Gamma power modulation over 24h period showed high gamma power during wake
and low gamma power during NREM. **(A1 - C1)** Show expanded time scales of data in A-C
(indicated by red dotted boxes) show **(A1)** a gradual fall of gamma power during transition
Wake-NREM and gradual rise of gamma power during transition NREM-Wake. **(B)** Gamma
power in the P120$^{+/-}$ mice failed to transition during Wake-NREM and NREM-Wake, expanded
time scale **(B1)** demonstrates the failure of gamma power attenuation during the NREM cycle.
(C) In PMP-administered P120$^{+/-}$ mice these behavioral-state transitions in gamma power were
significantly rescued **(C1)** expanded time scale. **(D)** Cumulative frequency graphs of positive
NREM-Wake gamma slopes and negative Wake-NREM slopes in WT$^{+/+}$mice **(D1)** Gamma
slopes were significantly different (*t-test*, t_{275}=10.59, p<0.001) between the two transition
states. **(D2)** No behavioral-state dependent modulation of gamma power occurred in P120$^{+/-}$
mice (*t-test*, t_{327}=0.3488, p=0.7275). **(D3)** PMP administration rescued gamma power
behavioral-state dependent modulation (*t-test*, t_{156}= 9.854, p<0.0001). (p<0.05 *, p<0.01 **;
p<0.001 ***). (see supplemental Fig. 6 for frequency distribution of gamma slopes for groups
and 2-way ANOVA for means of gamma slopes; see supplemental Fig. 7 for no significant
effect of PMP on WT$^{+/+}$ gamma modulation at P121).
768

Fig. 8. PMP rescued gamma modulation during NREM-REM transitions (A)
Representative heat maps (100s each) demonstrated an increase in gamma and decrease in
theta and delta power during NREM-REM transitions in WT$^{+/+}$ mice. In P120$^{+/-}$ mice both delta
and theta power decreased during NREM-REM transitions similar WT$^{+/+}$ mice. In contrast to
WT$^{+/+}$ mice, gamma power decreased during NREM-REM transitions. **(B)** Delta power during
NREM-REM transitions was significantly attenuated in P60$^{+/-}$ mice (WT$^{+/+}$ Delta vs. P60$^{+/-}$ Delta
% change: 1-way ANOVA, $F_{2,97}$=2.066, P=0.005, p=0.018) but was not attenuated in P120$^{+/-}$
mice (WT$^{+/+}$ Delta vs. P120$^{+/-}$ Delta % change 1-way ANOVA, $F_{2,97}$=2.066, P=0.005, p=1.00).
NREM-REM gamma modulation was significantly attenuated at P120$^{+/-}$ (WT$^{+/+}$ Gamma vs.
P120$^{+/-}$ Gamma % change 1-way ANOVA, $F_{2,97}$=6.821, P=0.002, p=0.001). **(C)** After PMP
administration, gamma modulation was not significantly different from WT$^{+/+}$ mice (p<0.05 *,
p<0.01 **; p<0.001 ***; post-hoc Bonferroni).
781

Fig. 9. PV+IN deficits in pre-frontal cortex (A) Representative 40X Z-stack fused image from
the DP cortex in WT$^{+/+}$ and *Syngap1$^{+/-}$* mice. **(B)** PV immunofluorescence represented by PV+
puncta showed a significant reduction of PV innervation in the DP cortex. In *Syngap1$^{+/-}$* mice,
total PV+ puncta counts were significantly lower (*t-test,* t_{33}=2.99, p=0.0051). **(C1 & C2)** PV+
puncta counts onto non-PV+ somas were also significantly lower in the DP cortex (*t-test*,
t_{142}=3.25, p=0.0014). There were no significant differences in mean GluA2 fluorescence
intensity on PV+IN and non-PV+ neurons in the DP cortex. Similar analyses done in the PrL
found no significant differences between genotypes. (p<0.05 *, p<0.01 **; p<0.001 ***).
790

Fig. 10. Increased mean GluA2 fluorescence intensity on PV+IN soma are region-specific (A1) Representative 40X Z-stack images of WT$^{+/+}$ and P120$^{+/-}$ mice from BCX, layers 2-3 and 5-6. White arrow heads in A1 indicate PV+IN shown in A2 and A3 for BCX layers 2-3 and 5-6, respectively. Total PV+ puncta counts and somatic PV+ puncta counts were not significantly different in the BCX. Representative 40X Z-stack images of mean GluA2 fluorescence intensity on PV+IN and non-PV+ neurons for BCX layers 2-3 and 5-6 **(A2 & A3)**. P120$^{+/-}$ mean GluA2 fluorescence intensity was significantly higher on PV+IN in both BCX layers 2-3 (*t-test*, $t_{63}=2.437$ p=0.0176) and 5-6 (*t-test*, $t_{70}=2.219$, p=0.0297). P120$^{+/-}$ non-PV+ neurons in BCX layer 2-3 also had a significant increase in mean GluA2 fluorescence intensity (*t-test*, $t_{110}=2.405$, p=0.0178). **(B1)** Representative 40X Z-stack images from MCX, layers 2-3 and 5-6, total PV+ puncta counts and somatic PV+ puncta counts were not significantly different in the MCX. White arrow heads in B1 indicate PV+IN shown in B2 and B3 for MCX layers 2-3 and 5-6, respectively. **(B2 & B3)** Mean GluA2 fluorescence intensity was not significantly different for PV+IN in the MCX layers 2/3 and 5/6. P120$^{+/-}$ mean GluA2 fluorescence intensity was significantly increased in PV+IN of MCX layers 5/6 (*t-test*, $t_{126}=2.707$, p=0.0077). **(C1)** Representative 40X Z-stack images from hippocampal CA1 and CA3. White arrow heads in C1 indicate PV+IN shown in C2 and C3 for CA1 and CA3, respectively. Total PV+ puncta counts and somatic PV+ puncta counts were not significantly different in CA1 or CA3. Representative 40X Z-stack images of mean GluA2 fluorescence intensity on PV+INs **(C2)** and non-PV+ neurons **(C3)**. CA3 P120$^{+/-}$ PV+INs showed an increase in mean GluA2 fluorescence intensity (*t-test*, $t_{52}=2.622$, p=0.0114; **(C2).** In non-PV+ neurons, CA3 mean GluA2 fluorescence intensity was also significantly increased in P120$^{+/-}$ mice (*t-test*, $t_{64}=2.428$, p=0.018; C3). In contrast, mean GluA2 fluorescence intensity in CA1 P120$^{+/-}$ non-PV+ neurons was significantly lower than in WT$^{+/+}$ mice (*t-test*, $t_{48}=2.582$, p=0.0129; C3). (p<0.05 *, p<0.01 **; p<0.001 ***).

Supplemental Fig. 1 Experimental paradigm to evaluate epilepsy in *Syngap*1$^{+/-}$ mice (A) *In vivo* EEG recordings of mice at P60, P90, P120, and P121. **(B)** At P121 7 out of 16 mice received Perampanel (2mg/kg I.P.) at 10am and 6pm during the additional 24h recording period. (Schematic generated with help of Biorender) **(C)** Location of electrode placement. **(D)** Table for sample sizes of mice in the study and samples sizes of all the 24h recordings analyzed in this study.

Supplemental Fig. 2 Spontaneous electrographic seizures during wake in *Syngap1*$^{+/-}$ mice (A) A 10min representative EEG trace (0.5-50Hz) of a spontaneous electrographic seizure during wake in a P120$^{+/-}$ mouse (red arrows denote the start and end of the electrographic seizure). Synchronous EMG recording demonstrated attenuation of EMG activity associated with freezing behavior in the mouse noted on video. This phenotype was in contrast to short duration myoclonic seizures (Fig. 1) with ~3Hz spike-wave discharges on EEG that showed time-locked myoclonic jerks. All electrographic seizures emerged during wake, whereas myoclonic seizures predominantly emerged during NREM. **(B)** Expanded time scale showed ictal activity on EEG recordings with concurrent silent EMG during the freezing behavior.

Supplemental Fig. 3 No differences in macro- or micro-sleep architecture of WT$^{+/+}$ mice between the ages of P60 and P120 (A) 24h hypnograms of ultradian rhythms over the diurnal light/dark phase in WT$^{+/+}$ mice at P60 (P60$^{+/+}$) and P120 (P120$^{+/+}$). **(B)** P60$^{+/+}$ and P120$^{+/+}$ mice spent significantly more time awake than asleep over the 24h period (P60$^{+/+}$ Wake vs. P60$^{+/+}$ Sleep: 2-way ANOVA, $F_{1,12}=162.1$, $P<0.0001$, $p<0.0001$; P120$^{+/+}$ Wake vs. P120$^{+/+}$ Sleep: 2-way ANOVA, $F_{1,12}=162.1$, $P<0.0001$, $p<0.0001$). **(B1)** P60$^{+/+}$ and P120$^{+/+}$ mice both did not have any significant differences between wake and sleep durations during the light phase. **(B2)** P60$^{+/+}$ and P120$^{+/+}$ mice spent significantly more time awake than asleep during the dark phase (P60$^{+/+}$ Wake vs. P60$^{+/+}$ Sleep: 2-way ANOVA, $F_{1,12}=216.9$, $P<0.0001$, $p<0.0001$; P120$^{+/+}$ Wake vs. P120$^{+/+}$ Sleep: 2-way ANOVA, $F_{1,12}=216.9$, $P<0.0001$, $p<0.0001$). **(C)** P60$^{+/+}$ and P120$^{+/+}$ mice did not have significant differences in the number of sleep/wake cycles during light vs. dark phase. **(C1 & C2)** The micro-architecture for the percent of both NREM and REM cycles during sleep showed an 80-20% split respectively, similar to WT$^{+/+}$ mice, was and not significantly different betweenP60$^{+/+}$ and P120$^{+/+}$ mice. ($p<0.05$ *, $p<0.01$ **; P120 $p<0.001$ ***, post-hoc Bonferroni).

Supplemental Fig. 4 IISs frequency distributions before and after PMP administration (A1 & B1) Representative frequency distributions of IIS (grey lines) and seizures (red lines) over 24h of EEG recording in two P120$^{+/-}$ mice. **(A2 & B2)** Two doses of PMP (2mg/kg I.P.) administered during the EEG recording on P121 prevented seizure occurrence in *Syngap$^{+/-}$* mice. PMP also reduced IIS rates (counts stated in each panel) which did not reach significance with the acute dosing protocol of PMP used in this study **(A2 & B2)**.

Supplemental Fig. 5 Dissociation of theta-gamma interaction during wake vs. sleep states (A) 24h spectral power analyses of theta and gamma power during wake and sleep. Theta and gamma power during wake and sleep were normalized to their respective durations for each mouse. Theta power increased significantly during wake in P120$^{+/-}$ mice (WT$^{+/+}$ Wake Theta vs. P120$^{+/-}$ Wake Theta: 1-way ANOVA, $F_{2,38}=11.64$, $P=0.0001$, $p=0.0028$). Gamma power during wake was significantly increased in P60$^{+/-}$ mice (WT$^{+/+}$ Wake Gamma vs. P60$^{+/-}$ Wake Gamma: 1-way ANOVA, $F_{2,38}=11.64$, $P=0.0001$, $p=0.021$); this significance was lost in P120$^{+/-}$ mice when compared to WT$^{+/+}$ mice. **(B)** During sleep, there were no significant differences in either theta or gamma power compared to WT$^{+/+}$ mice. However, during sleep a general increase in gamma power was apparent in P60$^{+/-}$ and P120$^{+/-}$ mice. **(C)** Representative 24h theta-gamma ratio plots during wake and sleep in a WT$^{+/+}$, P120$^{+/-}$, and PMP-administered (2mg/kg I.P. inj; B.D.) mouse. WT$^{+/+}$ mice had low theta power with high gamma power during wake states, resulting in a low theta-gamma ratio (TGR= 1.4). During sleep, WT$^{+/+}$ mice had high theta power with low gamma power, resulting in a high TGR during sleep (TGR=4.4). P120$^{+/-}$ mice demonstrated high gamma during both wake and sleep, resulting in similar TGRs during both wake and sleep states. **(D1)** The change in TGR from wake to sleep states was significantly reduced in P120$^{+/-}$ mice (WT$^{+/+}$ TGR % vs. P120$^{+/-}$ TGR%:1-way ANOVA, $F_{2,16}=6.456$, $P=0.0088$, $p=0.0076$). **(D2)** *Syngap1$^{+/-}$* mice treated with PMP were not significantly different than WT$^{+/+}$ mice. ($p<0.05$ *, $p<0.01$ **; P120 $p<0.001$ ***, post-hoc Bonferroni).

Supplemental Fig. 6 Frequency distribution of gamma slopes during Wake-NREM and
NREM-Wake transitions (A) The majority of gamma slopes during NREM-Wake transitions
(red lines) were positive in WT$^{+/+}$ mice, as gamma increased from NREM to wake. In contrast,
the majority of gamma slopes during Wake-NREM transitions (blue lines) were negative, as
gamma decreased from wake to NREM. In WT$^{+/+}$, the slope of gamma during Wake-NREM
and NREM-Wake transitions were state-dependent. **(B)** In P60$^{+/-}$ mice, the differences
between state-dependent gamma slopes were state-dependent. **(C)** In contrast, gamma slopes
were not state-dependent in P120$^{+/-}$ mice. **(D)** Administration of PMP restored state-dependent
gamma slopes as the gamma slopes were positive during NREM-Wake transitions and
negative during Wake-NREM transitions. **(E)** In WT$^{+/+}$ mice mean gamma slopes for NREM-
Wake and Wake-NREM transitions were significantly different (WT$^{+/+}$ NREM to Wake vs. WT$^{+/+}$
Wake to NREM: 2-way ANOVA, $F_{3,963}=28.24$, $P<0.0001$, $p<0.0001$). Further, both NREM-
Wake and Wake-NREM gamma slopes in WT$^{+/+}$ mice were significantly different than those in
P120$^{+/-}$ mice (WT$^{+/+}$ Wake vs. P120$^{+/-}$ Wake: 2-way ANOVA, $F_{3,963}=28.24$, $P<0.0001$,
$p<0.0001$; WT$^{+/+}$ NREM vs. P120$^{+/-}$ NREM, $p<0.0001$). Gamma slopes of NREM-Wake and
Wake-NREM transitions in P60$^{+/-}$ mice remained significantly different (P60$^{+/-}$ Wake vs. P60$^{+/-}$
NREM: 2-way ANOVA, $F_{3,963}=28.24$, $P<0.0001$, $p=0.0313$). In P120$^{+/-}$ mice the gamma slope
differences between NREM-Wake and Wake-NREM transitions were completely lost.
Administration of PMP restored the significant differences between Wake-NREM and NREM-
Wake transitions to WT$^{+/+}$ levels (PMP$^{+/-}$ Wake vs. PMP$^{+/-}$ NREM: 2-way ANOVA, $F_{3,963}=28.24$,
$P<0.0001$, $p<0.0001$, post-hoc Bonferroni).
901
Supplemental Fig. 7 PMP treatment did not significantly alter cortical gamma spectral
power transitions between wake and sleep at P120 (A) 24h WT$^{+/+}$ mice treated with PMP
have high gamma during wake and low gamma during NREM, similar to untreated WT$^{+/+}$ mice
(Fig. 7). **(A1)** Expanded time scales shows gradual fall of gamma during Wake-NREM
transitions and gradual rise of gamma during NREM-Wake transitions**. (B)** Cumulative
frequency plots of positive NREM to Wake slope (solid blue line) and negative Wake to NREM
slope (dashed line), these slopes were significantly different (*t-test*, $t_{59}=10.5$, $p=0.0001$).
($p<0.05$ *, $p<0.01$ **; P120 $p<0.001$ ***).
910
Supplemental Fig. 8 Gamma frequency analysis during wake and NREM (A1) Respective
frequency analysis (0.5-50Hz) of a WT$^{+/+}$ mouse during wake, NREM, and REM. **(A2)** The
gamma power (35-50Hz) of the respective WT$^{+/+}$ mouse was significantly higher during wake
than NREM (Wake vs. NREM: *t-test*, $t_{14.02}=300$, $P<0.0001$). **(B1)** WT$^{+/+}$ mice had significantly
higher gamma power (35-50Hz) during wake than NREM (WT$^{+/+}$ Wake vs. WT$^{+/+}$ NREM: *t-test*,
$t_{19.65}=300$, $P<0.0001$). **(B2)** P120$^{+/-}$ mice had significantly different gamma power from 35-50Hz
during NREM than wake (P120$^{+/-}$ Wake vs. P120$^{+/-}$ NREM: *t-test*, $t_{4.25}=300$, $P<0.0001$). **(C1)**
Lower frequency gamma from 35-45Hz in wake and NREM was significantly different in WT$^{+/+}$
mice (WT$^{+/+}$ 35-50Hz Wake vs. WT$^{+/+}$ 35-50Hz NREM: *t-test*, $t_{24.84}=200$, $P<0.0001$). **(C2)** In
P120$^{+/-}$ mice, lower gamma frequency (35-45Hz) remained high during NREM. P120$^{+/-}$ mice
did not have any significant differences in gamma power (35-45Hz) between wake and NREM
(P120$^{+/-}$ 35-50Hz Wake vs. P120$^{+/-}$ 35-50Hz NREM: *t-test*, $t_{15.31}=198$, $P<0.0001$). ($p<0.05$ *,
$p<0.01$ **; P120 $p<0.001$ ***).
924

Supplemental Fig. 9 *Syngap1*^{+/-} mice did not show evidence for mossy fiber sprouting
and GluA2 antibody validation for IHC. (A) Representative 10X hippocampal section of a
WT$^{+/+}$ mouse with ZnT3 staining (green) to label mossy fibers, an established marker of
hippocampal epileptogenesis, with representative expansions of CA3 (left) and dentate gyrus
(DG; right). **(B)** Representative 10X hippocampal section of a P120$^{+/-}$ mouse demonstrated the
absence of mossy fiber sprouting in both the DG and CA3. (C) Representative 10X
hippocampal section of a WT$^{+/+}$ mouse with GluA2 positive fluorescence and a GluA2$^{-/-}$ mouse
(Richard Huganir, Johns Hopkins University) lacking GluA2 fluorescence.
933
934

References

1.   Chen, H.-J., Rojas-Soto, M., Oguni, A. & Kennedy, M. B. A Synaptic Ras-GTPase Activating
Protein (p135 SynGAP) Inhibited by CaM Kinase II. *Neuron* **20**, 895–904 (1998).
2.   Kim, J. H., Liao, D., Lau, L. F. & Huganir, R. L. SynGAP: a synaptic RasGAP that associates
with the PSD-95/SAP90 protein family. *Neuron* **20**, 683–691 (1998).
3.   Rumbaugh, G., Adams, J. P., Kim, J. H. & Huganir, R. L. SynGAP regulates synaptic strength
and mitogen-activated protein kinases in cultured neurons. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 4344–
4351 (2006).
4.   Araki, Y., Zeng, M., Zhang, M. & Huganir, R. L. Rapid dispersion of SynGAP from synaptic
spines triggers AMPA receptor insertion and spine enlargement during LTP., Rapid Dispersion of
SynGAP from Synaptic Spines Triggers AMPA Receptor Insertion and Spine Enlargement during LTP.
Neuron Neuron **85**, **85**, 173, 173–189 (2015).
5.   Zeng, M., Bai, G. & Zhang, M. Anchoring high concentrations of SynGAP at postsynaptic
densities via liquid-liquid phase separation. *Small GTPases* 1–9 (2017).
doi:10.1080/21541248.2017.1320350
6.   Purcell, S. M. *et al.* A polygenic burden of rare disruptive mutations in schizophrenia. *Nature*
506, 185–190 (2014).
7.   Hamdan, F. F. *et al.* Mutations in SYNGAP1 in autosomal nonsyndromic mental retardation. *N.*
Engl. J. Med. **360**, 599–605 (2009).
8.   Hamdan, F. F. *et al.* De novo SYNGAP1 mutations in nonsyndromic intellectual disability and
autism. *Biol. Psychiatry* **69**, 898–901 (2011).
9.   Berryer Martin H. *et al.* Mutations in SYNGAP1 Cause Intellectual Disability, Autism, and a
Specific Form of Epilepsy by Inducing Haploinsufficiency. *Hum. Mutat.* **34**, 385–394 (2012).
10.   Carvill, G. L. *et al.* Targeted resequencing in epileptic encephalopathies identifies *de novo*
mutations in *CHD2* and *SYNGAP1*. *Nat. Genet.* **45**, 825–830 (2013).
11.   Mignot, C. *et al.* Genetic and neurodevelopmental spectrum of SYNGAP1-associated intellectual
disability and epilepsy. *J. Med. Genet.* **53**, 511–522 (2016).
12.   Vlaskamp, D. R. M. *et al.* SYNGAP1 encephalopathy: A distinctive generalized developmental
and epileptic encephalopathy. *Neurology* **92**, e96–e107 (2019).
13.   Jain, S. V. & Kothare, S. V. Sleep and Epilepsy. *Semin. Pediatr. Neurol.* **22**, 86–92 (2015).
14.   Robinson-Shelton, A. & Malow, B. A. Sleep Disturbances in Neurodevelopmental Disorders.
Curr. Psychiatry Rep. **18**, 6 (2015).
15.   Malow, B. A. Sleep disorders, epilepsy, and autism. *Ment. Retard. Dev. Disabil. Res. Rev.* **10**,
122–125 (2004).
16.   Glaze, D. G. Neurophysiology of Rett syndrome. *J Child Neurol* **20**, 740–746 (2005).
17.   Ammanuel, S. *et al.* Heightened Delta Power during Slow-Wave-Sleep in Patients with Rett
Syndrome Associated with Poor Sleep Efficiency. *PLoS One* **10**, e0138113- (2015).
18.   Johnston, M. V. *et al.* Twenty-four hour quantitative-EEG and in-vivo glutamate biosensor
detects activity and circadian rhythm dependent biomarkers of pathogenesis in Mecp2 null mice. *Front*
SystNeurosci **8**, 118- (2014).
19.   Holmes, G. L. Effects of seizures on brain development: lessons from the laboratory. *Pediatr.*
Neurol. **33**, 1–11 (2005).

20.   Nabbout, R. & Dulac, O. Epileptic encephalopathies: a brief overview. *J. Clin. Neurophysiol.*
Off. Publ. Am. Electroencephalogr. Soc. **20**, 393–397 (2003).
21.   Prchalova, D. *et al.* Analysis of 31-year-old patient with SYNGAP1 gene defect points to
importance of variants in broader splice regions and reveals developmental trajectory of SYNGAP1-
associated phenotype: case report. *BMC Med. Genet.* **18**, (2017).
22.   Clement, J. P., Ozkan, E. D., Aceti, M., Miller, C. A. & Rumbaugh, G. SYNGAP1 Links the
Maturation Rate of Excitatory Synapses to the Duration of Critical-Period Synaptic Plasticity. *J.*
Neurosci. **33**, 10447–10452 (2013).
23.   Clement, J. P. *et al.* Pathogenic SYNGAP1 mutations impair cognitive development by
disrupting maturation of dendritic spine synapses. *Cell* **151**, 709–723 (2012).
24.   Kim, J. H., Lee, H.-K., Takamiya, K. & Huganir, R. L. The role of synaptic GTPase-activating
protein in neuronal development and synaptic plasticity. *J. Neurosci. Off. J. Soc. Neurosci.* **23**, 1119–
1124 (2003).
25.   SYNGAP1 Research. Available at: https://www.kennedykrieger.org/research/participate-in-
research. (Accessed: 1st July 2019)
26.   Staley, K. J., White, A. & Dudek, F. E. Interictal spikes: harbingers or causes of epilepsy?
Neurosci. Lett. **497**, 247–250 (2011).
27.   Baud, M. O. *et al.* Multi-day rhythms modulate seizure risk in epilepsy. *Nat. Commun.* **9**, 88
(2018).
28.   Bains, R. S. *et al.* Assessing mouse behaviour throughout the light/dark cycle using automated
in-cage analysis tools. *J. Neurosci. Methods* **300**, 37–47 (2018).
29.   Nakajima, R. *et al.* Comprehensive behavioral analysis of heterozygous Syngap1 knockout mice.
Neuropsychopharmacol. Rep. (2019). doi:10.1002/npr2.12073
30.   Komiyama, N. H. *et al.* SynGAP Regulates ERK/MAPK Signaling, Synaptic Plasticity, and
Learning in the Complex with Postsynaptic Density 95 and NMDA Receptor. *J. Neurosci.* **22**, 9721–
9732 (2002).
31.   Belluscio, M. A., Mizuseki, K., Schmidt, R., Kempter, R. & Buzsáki, G. Cross-frequency phase-
phase coupling between θ and γ oscillations in the hippocampus. *J. Neurosci. Off. J. Soc. Neurosci.* **32**,
423–435 (2012).
32.   Buzsáki, G., Anastassiou, C. A. & Koch, C. The origin of extracellular fields and currents —
EEG, ECoG, LFP and spikes. *Nat. Rev. Neurosci.* **13**, 407–420 (2012).
33.   Bragin, A. *et al.* Gamma (40-100 Hz) oscillation in the hippocampus of the behaving rat. *J.*
Neurosci. Off. J. Soc. Neurosci. **15**, 47–60 (1995).
34.   Soltesz, I. & Deschênes, M. Low- and high-frequency membrane potential oscillations during
theta activity in CA1 and CA3 pyramidal neurons of the rat hippocampus under ketamine-xylazine
anesthesia. *J. Neurophysiol.* **70**, 97–116 (1993).
35.   Axmacher, N. *et al.* Cross-frequency coupling supports multi-item working memory in the
human hippocampus. *Proc. Natl. Acad. Sci.* **107**, 3228–3233 (2010).
36.   Demiralp, T. *et al.* Gamma amplitudes are coupled to theta phase in human EEG during visual
perception. *Int. J. Psychophysiol. Off. J. Int. Organ. Psychophysiol.* **64**, 24–30 (2007).
37.   Köster, M., Friese, U., Schöne, B., Trujillo-Barreto, N. & Gruber, T. Theta-gamma coupling
during episodic retrieval in the human EEG. *Brain Res.* **1577**, 57–68 (2014).
38.   Kim, H., Ährlund-Richter, S., Wang, X., Deisseroth, K. & Carlén, M. Prefrontal Parvalbumin
Neurons in Control of Attention. *Cell* **164**, 208–218 (2016).
39.   Lisman, J. E. & Jensen, O. The Theta-Gamma Neural Code. *Neuron* **77**, 1002–1016 (2013).

40. Moretti, D. V. *et al.* Increase of theta/gamma ratio is associated with memory impairment. *Clin.*
Neurophysiol. **120**, 295–303 (2009).
41. Buzsaki, G. & Wang, X. J. Mechanisms of gamma oscillations. *Annu. Neurosci* **35**, 203–225
(2012).
42. Valderrama, M. *et al.* Human Gamma Oscillations during Slow Wave Sleep. *PLoS ONE* **7**,
(2012).
43. Scheffzük, C. *et al.* Selective coupling between theta phase and neocortical fast gamma
oscillations during REM-sleep in mice. *PloS One* **6**, e28489 (2011).
44. Sohal, V. S., Zhang, F., Yizhar, O. & Deisseroth, K. Parvalbumin neurons and gamma rhythms
enhance cortical circuit performance. *Nature* **459**, 698–702 (2009).
45. Tremblay, R., Lee, S. & Rudy, B. GABAergic Interneurons in the Neocortex: From Cellular
Properties to Circuits. *Neuron* **91**, 260–292 (2016).
46. Huang, Z. J. & Paul, A. The diversity of GABAergic neurons and neural communication
elements. *Nat. Rev. Neurosci.* 1 (2019). doi:10.1038/s41583-019-0195-4
47. Akgül, G. & McBain, C. J. Diverse roles for ionotropic glutamate receptors on inhibitory
interneurons in developing and adult brain. *J. Physiol.* **594**, 5471–5490 (2016).
48. Adler, D. A. *et al.* Circadian cycle-dependent EEG biomarkers of pathogenicity in adult mice
following prenatal exposure to in utero inflammation. *Neuroscience* (2014).
49. Fuchs, E. C. *et al.* Genetically altered AMPA-type glutamate receptor kinetics in interneurons
disrupt long-range synchrony of gamma oscillation. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 3571–3576
(2001).
50. Vyazovskiy, V. V., Cirelli, C., Pfister-Genskow, M., Faraguna, U. & Tononi, G. Molecular and
electrophysiological evidence for net synaptic potentiation in wake and depression in sleep. *Nat.*
Neurosci. **11**, 200–208 (2008).
51. Fuchs, E. C. *et al.* Genetically altered AMPA-type glutamate receptor kinetics in interneurons
disrupt long-range synchrony of gamma oscillation. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 3571–3576
(2001).
52. Fuchs, E. C. *et al.* Recruitment of Parvalbumin-Positive Interneurons Determines Hippocampal
Function and Associated Behavior. *Neuron* **53**, 591–604 (2007).
53. Berryer, M. H. *et al.* Decrease of SYNGAP1 in GABAergic cells impairs inhibitory synapse
connectivity, synaptic inhibition and cognitive function. *Nat. Commun.* **7**, 13340 (2016).
54. Le Magueresse, C. & Monyer, H. GABAergic Interneurons Shape the Functional Maturation of
the Cortex. *Neuron* **77**, 388–405 (2013).
55. Altimimi, H. F. & Stellwagen, D. Persistent Synaptic Scaling Independent of AMPA Receptor
Subunit Composition. *J. Neurosci.* **33**, 11763–11767 (2013).
56. Turrigiano, G. Too Many Cooks? Intrinsic and Synaptic Homeostatic Mechanisms in Cortical
Circuit Refinement. *Annu. Rev. Neurosci.* **34**, 89–103 (2011).
57. Jonas, P. Differences in Ca2+ permeability of AMPA-type glutamate receptor channels in
neocortical neurons caused by differential GluR-B subunit expression. *Neuron* **12**, 1281–1289 (1994).
58. Hu, H., Gan, J. & Jonas, P. Fast-spiking, parvalbumin+ GABAergic interneurons: From cellular
design to microcircuit function. *Science* **345**, 1255263–1255263 (2014).
59. Cardin, J. A. *et al.* Driving fast-spiking cells induces gamma rhythm and controls sensory
responses. *Nature* **459**, 663–667 (2009).
60. Michaelson, S. D. *et al.* SYNGAP1 heterozygosity disrupts sensory processing by reducing
touch-related activity within somatosensory cortex circuits. *Nat. Neurosci.* **21**, 1 (2018).

61.   Striano, P. & Huppke, P. A synaptic protein defect associated with reflex seizure disorder.
Neurology 10.1212/WNL.000000000006720 (2018). doi:10.1212/WNL.000000000006720
62.   Blatow, M. *et al.* A Novel Network of Multipolar Bursting Interneurons Generates Theta
Frequency Oscillations in Neocortex. *Neuron* **38**, 805–817 (2003).
63.   Kullmann, D. M. & Lamsa, K. Roles of distinct glutamate receptors in induction of anti-Hebbian
long-term potentiation. *J. Physiol.* **586**, 1481–1486 (2008).
64.   McMahon, A. C. *et al.* SynGAP isoforms exert opposing effects on synaptic strength. *Nat.*
Commun. **3**, 900 (2012).
65.   Creson, T. K. *et al.* Re-expression of SynGAP protein in adulthood improves translatable
measures of brain function and behavior. *eLife* **8**, (2019).
66.   Kang, S. K. *et al.* Sleep dysfunction following neonatal ischemic seizures are differential by
neonatal age of insult as determined by qEEG in a mouse model. *Neurobiol. Dis.* **116**, 1–12 (2018).
67.   Kadam, S. D. *et al.* Methodological standards and interpretation of video-
electroencephalography in adult control rodents. A TASK1-WG1 report of the AES/ILAE Translational
Task Force of the ILAE. *Epilepsia* **58 Suppl 4**, 10–27 (2017).
68.   ImageJ Plugins. *ImageJ* Available at: https://imagej.nih.gov/ij/plugins/index.html. (Accessed: 1st
July 2019)

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

Figure 1

1120

Figure 2

Figure 3

1130

1131

1132

1133

1134

1135

1136

Figure 4

Figure 5

Figure 6

1150

1151

1152

Figure 7

1159

Figure 8

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

Figure 9

Figure 10

Supplemental Figure 1

A. Surgery   P60   P90   P120   P121

B. Electrode Location

1 Anchor   3 EEG1

2 Ground/Reference   4 EEG2

C. PMP I.P. 2mg/kg   10am   6pm

D.

	Control	SynGAP
Male	4	4
Female	4	4
Total	8	8
P60 24h EEG	4	6
P90-120 24h EEG	5	9
Total	9	15
P120 PMP 24h EEG	3	4

Supplemental Figure 2

Supplemental Figure 3

Supplemental Figure 4

Supplemental Figure 5

Supplemental Figure 6

Supplemental Figure 7

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

Supplemental Figure 8