Pretreatment with yeast derived complex dietary-polysaccharide leads to suppressed gut inflammation, altered microbiota composition and increased immune regulatory short-chain fatty acid production in C57BL/6 mice

Radhika Gudi[1*], Jada Suber[1*], Robert Brown[1*], Benjamin M. Johnson[1*], and Chenthamarakshan Vasu[1,5]

[1]Department of Microbiology and Immunology, College of Medicine, Medical University of South Carolina, Charleston, SC-29425

Running title: Dietary polysaccharide on gut inflammation

*contributed equally

[5]Address Correspondence: Chenthamarakshan Vasu
Microbiology and Immunology
Medical University of South Carolina
173 Ashley Avenue, BSB208E
Charleston, SC-29425
Phone: 843-792-1032, Fax: 843-792-9588
Email: vasu@musc.edu

Conflict of Interest: Authors do not have any conflict(s) of interest to disclose.

ABSTRACT

Background: β-Glucans (BGs), a group of complex non-digestible polysaccharides, are

considered to have beneficial health effects due to their immune modulatory properties and are

considered as dietary supplements. However, the impact of oral administration of high-pure, well-

defined BGs on gut inflammation, and the influence of intestinal microbiota and short-chain fatty

acid (SCFA) on the therapeutic effect are largely unknown.

Objectives: The aim of this study, using a mouse model of chemical induced colitis, was to

investigate the impact of oral administration of high-pure yeast BG (YBG) on the susceptibility to

colitis, gut immune function, and structure and function of microbiota.

Methods: To determine the impact of oral administration of YBG on colitis susceptibility, eight

week old C57BL/6 (B6) mice were pre-treated with YBG (250 µg/mouse/day) and given dextran

sulfate sodium (DSS) in drinking water (2.5% w/v) and examined for the symptoms and features

of colitis. To assess the effect of oral administration of YBG on gut mucosa and microbiota

structure and function, and gut immune regulation, we determined the microbiota composition,

fecal SCFA levels, and intestinal T cell phenotype and cytokine secretion. The role of gut

microbiota in YBG treatment induced modulation of gut inflammation and immune function were

determined in B6 mice treated with broad-spectrum antibiotic cocktail (1 g/L ampicillin, 0.5 g/L

vancomycin, 1 g/L neomycin, and 1 g/L metronidazole) in drinking water.

Results Compared to untreated mice, B6 mice that received prolonged pre-treatment with YBG

showed diminished severity of different features of DSS-induced colitis including overall loss of

body weight ($P<0.001$), shortening of colon ($P=0.016$) and histopathology ($P=0.01$). However,

high-pure YBG has no beneficial effect in terms of suppressing colitis severity when consumed

only during the disease stage. Compared to untreated controls, YBG pre-treated mice showed

higher regulatory T cell (Treg) frequencies ($P=0.043$) in the gut mucosa, a shift in the abundance

of gut microbiota towards polysaccharide-fermenting bacterial phyla Bacteroides ($P=0.049$) and

Verrucomicrobia (Mean±SD: control=13.0±0.33 vs YBG=10.9.7±0.69) and diminished

Firmicutes ($P<0.001$) and Proteobacteria ($P<0.001$), and significantly higher production of SCFA

such as acetic acid ($P=0.016$), propionic acid ($P=0.026$) and butyric acid ($P=0.013$). Depletion of

gut microbiota in YBG-fed B6 mice using broad spectrum antibiotics caused not only elimination

of YBG treatment associated SCFA production and Treg increase, but also profound aggravation

of the pathological features of colitis such as loss of body weight ($P<0.01$) and colonic

inflammation ($P=0.04$) compared to that of YBG treated control mice.

Conclusions: Oral consumption of high-pure BG promotes a healthy gut homeostasis and immune

regulation, and minimizes susceptibility to DSS induced colitis in B6 mice in a microbiota (and

microbial SCFA) - dependent manner. On the contrary, YBG consumption when gut mucosa and

microbiota are compromised not only reverses this protection but also increases the susceptibility

to gut inflammation and disease severity, perhaps through its direct interaction with gut immune

cells. In conclusion, while YBG consumption may be beneficial for gut health and to prevent gut

inflammation in healthy individuals and under intact microbiota, this immune stimulatory dietary

supplement may not have any health benefits in individuals with active gut inflammation and could

cause adverse effect in those who are on oral antibiotics.

Key words: Yeast β-glucan; gut mucosa, Gut microbiota; short-chain fatty acid, colitis, gut

inflammation, immune modulation; immune regulation.

INTRODUCTION

Dietary approaches have increasingly been considered for modulating immune function, and preventing autoimmunity and other clinical conditions. β-glucans (BGs) are non-digestible complex dietary polysaccharides (CDPs) of varying structures (β-1,3-D-glucan, β-1,3/1,6-D-glucan, β-1,3/1,4-D-glucan and β-1,4-D-glucan) commonly found in plants, cereals, bacteria, yeast, and fungi (1). While BGs can potentiate the function of immune system directly (2, 3), BG-containing preparations have been promoted also as daily dietary supplements to enhance metabolic functions, and as prebiotics (4). On the other hand, clinical studies using YBG preparations reported conflicting outcomes in terms of their potential benefits as dietary supplements (5-9). Studies using animal models to determine the potential of BG preparations and BG containing diets to modulate gut inflammation have also produced conflicting outcomes (10-15). Furthermore, inconsistent results on colitis susceptibility of Dectin-1 (BG interacting receptor) knockout mice have been reported (16, 17).

Differences observed in the impact of BG dietary supplements on host health could be due multiple factors including 1) the purity and/or the structure of the BG employed, 2) degree of direct interactions of those BG preparations with host receptors, 3) possible association between gut microbes and BG induced effects in the host, and differences in the structure and function of basal gut microbiota. Determination of the true effect of oral consumption of BGs on the host, thus, requires systematic pre-clinical studies using well-defined high-pure BGs. Further, studying the direct immune modulatory effect of BGs on gut mucosa and the indirect, microbiota-dependent, effect is necessary to better understand mechanisms of immune modulation induced these dietary complex polysaccharides. Such studies will also help explain the contradictory effects of these agents on the host. For this study, we hypothesized that oral administration of a highly purified BG [β-1,3-linked D-glucose molecules (β-1,3-D-glucan) with β-1,6-linked side chains] from

Baker's yeast (*Saccharomyces cerevisiae*) (YBG) prevents autoimmunity (18, 19) can enhance

immune modulation by directly interacting with the gut mucosa and by serving as a fermentation

substrate for gut commensals that produce immune regulatory metabolites. Therefore, we

examined the impact of oral administration of high pure YBG on gut inflammation employing

DSS induced colitis in C57BL/6 (B6) mice as the model, and assessed the contribution of gut

microbiota to YBG consumption associated enhanced gut immune regulation and suppression of

colitis susceptibility.

Materials and Methods

Mice

C57BL/6 (B6) mice were originally purchased from the Jackson laboratory. Foxp3-GFP-knockin

(Foxp3-GFP) mice in B6 background were provided by Dr. Vijay Kuchroo (Harvard Medical

School, MA). Breeding colonies of these strains were established and maintained in the pathogen-

free facility of Medical University of South Carolina (MUSC). Mice were given autoclaved rodent

diet (20), which contains 18.6% (wt:wt) protein, 6.2% (wt:wt) fat, and 44.2% (wt:wt)

carbohydrates (Harlan Laboratories, Teklad Global 18% Protein Rodent diet no. 2018), and

autoclaved water ad libitum. All animal studies were approved by the animal care and use

committee at MUSC. Since B6 mice are highly susceptible to DSS induced distal gut inflammation

and has been widely used as a model of ulcerative colitis (UC) (21-24), we employed this strain

for determining the impact of YBG on DSS induced colitis. While most experiments described in

this study used age matched females for control and test groups, our initial experiments showed

YBG treatment induced similar protection of male B6 mice from colitis. However, since mortality

was as high as 50% by day 10 with control males with DSS colitis, to be able to compare similar

number of controls and YBG treated mice side-by-side, we preferred females for this study. B6-

Foxp3-GFP report mice that express GFP under Foxp3 promoter and only in regulatory T cell were

used in some experiments to visualize and enumerate Tregs by FACS unambiguously.

YBG and other reagents

Purified YBG (glucan from Baker's yeast, *S. cerevisiae*), ≥98% pure, was purchased from Sigma-

Aldrich and the purification method has been described before (18). This agent was tested for

specific activity using thioglycolate-activated macrophages as described before by us and others

(25-27). PMA, ionomycin, Brefeldin A, and purified and conjugated antibodies were purchased

from Sigma-Aldrich, BD Biosciences, eBioscience, Invitrogen, R&D Systems, and Biolegend

Laboratories. In some experiments, intestinal epithelial cells were depleted by Percoll gradient

centrifugation (28) or using A33 antibody (Santa Cruz Biotechnology) and anti-rabbit IgG-

magnetic beads (29, 30). Oligonucleotide probes for qPCR were custom synthesized by Thermo

Fisher. Magnetic bead-based multiplex assay kits were purchased from Invitrogen. Additional

details of commercial products used in this study are listed in supplemental table 1.

YBG treatment

Eight week old B6 mice or B6-Foxp3-GFP mice were given YBG suspension in saline (250

μg/mouse/day) for up to 30 days by oral gavage. Mice that were subjected to colitis induction using

DSS, as described below, continued to receive YBG or saline until the mice were euthanized on

day 40. In some experiments, YBG and saline treatments were initiated only 1 day before initiating

the DSS treatment to determine the effect of treatment initiation at disease stage. This dose of YBG

was considered because it is a human relevant dose of BG dietary supplements (500 to 1000

mg/day) equaling approximately 10 mg/kg body weight. Further, as mentioned in our previous

report (29), lower doses or short-term treatments do not produce observable immune modulation

in the gut mucosa. We also found that even a very short-term (3 day) treatment with higher doses

such as 2000 μg/mouse/day produce pro-inflammatory response in the gut mucosa in Dectin-1

dependent manner (29); hence a relatively lower daily dose was considered for this study.

Induction of DSS colitis and disease evaluation

To induce colitis, mice were maintained on drinking water containing 2.5% (w/v) DSS salt

(36,000-50,000 MW) for 5 consecutive days. After 5 days, mice were switched to regular water

until euthanasia on day 10, post-DSS treatment initiation. Mice were weighed, and examined for

the stool consistency and presence of occult blood every day, and the severities were scored as

described before (31). In some experiments, mice were given a cocktail of antibiotics starting 5

days before DSS treatment. Mice were euthanized 10 days post-DSS treatment initiation, intestinal

tissues were collected, the colon lengths were measured, and distal colon tissues were subjected to

histochemical staining to determine the severity of inflammation. Distal colon tissues were fixed

in 10% formaldehyde, 5-μm paraffin sections were cut, and stained with hematoxylin and eosin.

Stained sections were analyzed using the 0-6 grading criteria described before (32) for

inflammatory cell infiltrate, epithelial changes and mucosal architecture, and the values were

averaged to calculate the overall inflammation score of individual mice.

Immunological assays and qPCR

Cells from the small intestinal lamina propria (SiLP), large intestinal lamina propria (LiLP) and

mesenteric lymph node (MLN) were examined for immune cell phenotype and/or cytokine profiles

by FACS and multiplex assay respectively as described in our recent reports (29, 30, 33). Briefly:

for determining Treg frequencies, freshly isolated cells were stained for CD4 and Foxp3. For

determining cytokine positive T cells frequencies, cells were cultured in the presence or PMA,

Ionomycin and Brefeldin A for 4 h, before staying for CD4 and intracellular cytokines. For

determining secreted cytokine levels, cells were cultured in the presence of anti-CD3 antibody for

24 h and the spent media were subjected to multiplex assay and read using FlexMap 3D instrument

(Luminex). SYBR green based qPCR assay was run in a CFX96 Touch real time PCR machine

(BioRad) for determining cytokine gene expression levels in the RNA prepared from

cryopreserved intestinal tissues.

16S rRNA gene targeted sequencing and bacterial community profiling

Total DNA was prepared from the fecal pellets for 16S rRNA gene targeted sequencing and

bacterial community profiling as detailed in our previous reports (19, 34, 35) with minor

modifications. Briefly, the sequencing reads were fed into the Metagenomics application of

BaseSpace (Illumina) for performing taxonomic classification of 16S rRNA targeted amplicon

reads using an Illumina-curated version of the GreenGenes taxonomic database which provided

raw classification output at multiple taxonomic levels. The sequences were also fed into QIIME

open reference operational taxonomic units (OTU) picking pipeline (36) using pre-processing

application of BaseSpace. The OTUs were compiled to different taxonomical levels based upon

the percentage identity to GreenGenes reference sequences (i.e. >97% identity) and the percentage

values of sequences within each sample that map to respective taxonomical levels were calculated.

The OTUs were also normalized and used for metagenomes prediction of Kyoto Encyclopedia of

Genes and Genomes (KEGG) orthologs employing PICRUSt as described before (37-40). The

predictions were summarized to multiple levels and functional categories were compared among

control and YBG fed groups using the statistical Analysis of Metagenomic Profile Package

(STAMP) as described before (38, 41). STAMP and web-based MicrobiomeAnalyst (42)

applications were also employed for visualization and/or statistical analysis of microbiome data.

Depletion of gut microbiota

Mice were given a broad-spectrum antibiotic cocktail (ampicillin (1 g/l), vancomycin (0.5 g/l),

neomycin (1 g/l), and metronidazole (1 g/l) -containing drinking water during the last 5 days of

YBG treatment to deplete gut microbiota as described in our recent report (29). Fecal pellet

suspensions were cultured under aerobic and anaerobic conditions to assess microbiota depletion

(29). In some experiments, antibiotic treatment was continued during colitis induction using DSS.

Determination of fecal SCFA levels

Fresh fecal samples were collected from individual mice, stored at -80°C, and shipped on dry ice

to Microbiome Insights (Vancouver, Canada) for SCFA analysis. SCFA were detected by this

service provider using gas chromatography (Thermo Trace 1310), coupled to a flame ionization

detector, using Thermo TG-WAXMS A GC Column by following a previously described method

(43).

Statistical analysis

Mean, SD, and statistical significance (*P-value*) were calculated, graphic visualizations were made

using Excel (Microsoft), Prism (GraphPad), Morpheus (Broad Institute), STAMP (38, 41) and/or

MicrobiomeAnalyst (42) applications. Briefly, unpaired, two-tailed *t*-test or two-tailed non-

parametric Mann-Whitney test was employed for most assays where test (YBG fed or microbiota

depleted) group samples were compared to that of respective control (saline fed or those with intact

microbiota) group, unless specified here. For DSS colitis associated weight loss, stool consistency

and stool levels of blood, test group mean of each time point was compared to the respective time-

point control group mean by unpaired, two-tailed *t*-test. To compare the number of mice with grade

≤3 and ≥4 colon histopathology scores in test and control groups, two-tailed Fishers' exact

contingency test was employed. Log-rank test was employed for comparing the survival rates of

YBG treated mice with that of controls. Statistical analysis of microbiota at genus level as well as

the number of sequences representing (predicted) specific metabolic pathways were done in

STAMP employing two-sided Welch's *t*-test and FDR corrected using Benjamini and Hochberg

approach, and the heatmap visualization was done in Morpheus. MicrobiomeAnalyst was

employed for generating principal component (PC) analysis representing β-diversity (Bray Curtis

distance) and α-diversity / species richness / Chao1 estimator plots, and for statistical significances

by permutational multivariate analysis of variance and Mann-Whitney approaches respectively. A

p-value of ≤0.05 was considered significant.

Results

Prolonged pretreatment with YBG diminishes susceptibility to DSS induced colitis in B6

mice.

Eight week old female B6 mice were orally administered with YBG and subjected to colitis

induction using DSS as shown in **Fig. 1A**, and examined for gut inflammation associated features

to determine the impact of pre-treatment with YBG on colitis susceptibility. As observed in **Fig.**

1B, YBG treated mice were relatively less susceptible to DSS colitis associated weight loss

compared to control mice. Significant difference in the body weight between control and YBG

treated groups was observed starting day 6 (P=0.027) and peaked at day 10 (P<0.001). As shown

in **Fig. 1C**, YBG treated mice also showed significant resistance to colitis associated shortening

of the colon compared to that of control mice (P=0.016). Overall DSS induced histological injury

scores of YBG treated mice was significantly lower (P=0.016) as compared to that of control group

(Fig. 1D). The colon of DSS-treated control mice exhibited more severe inflammatory infiltrates

and ulceration compared to YBG treated mice. Further, compared to DSS treated control mice,

better stool consistency **(supplemental Fig. 1A)** and lesser fecal blood **(supplemental Fig. 1B)**

were detected in YBG-fed mice that were treated with DSS at the time-point in which these

features were most severe. Of note, prolonged pre-treatment of male B6 mice with YBG produced

similar protection to that of B6 female counterparts, but control group showed 50% mortality prior

to termination of the experiment (not shown). Further, we also observed that mice that were given

YBG only during colitis induction did not show any difference in the severity of colitis associated

parameters compared to control mice **(Supplemental Fig. 2)**.

YBG treated and control B6 mice with colitis showed distinct immune characteristics

Since B6 mice that were pretreated with YBG showed less severe colon inflammation, T cells

from their gut associated mesenteric lymph nodes (MLNs) were examined for phenotypic

properties. As shown in **Fig. 2A** and **supplemental Fig. 3**, Foxp3+ T regulatory cell (Treg)

frequencies were significantly higher in the MLN of YBG-fed mice (P=0.0043) compared to

control mice. FACS analyses revealed significantly higher frequencies of IL10+ (P=0.0043), IL4+

(P=0.041) and IL9+ (P=0.015) and lower IFNγ+ (P=0.0087), IL17+ (P=0.0022) and IL22+

(P=0.0152) T cell frequencies in YBG-fed mice compared to controls. To validate these

observations, the levels of cytokines secreted by MLN cell were determined. Multiplex cytokine

assay revealed significantly higher secretion of IL10+ (P=0.0087), IL4+ (P=0.0152) and IL9+

(P=0.041) and lower IFNγ+ (P=0.0087), IL17+ (P=0.041) and IL22+ (P=0.026) by MLN cells

from YBG fed mice, compared to that of control mice, upon *ex vivo* activation using anti-CD3

antibody **(Fig. 2B)**.

Prolonged YBG treatment results in changes in the fecal microbiota composition in B6 mice

Since only prolonged with YBG prior to DSS treatment reduced the susceptibility to colitis in B6

mice, we determined if YBG treatment alters gut microbiota structure and function in them.

Compilation of OTU data generated from 16S rRNA sequences of fecal samples to different

taxonomical levels showed that, compared to the pre-treatment time-point, abundances of major

phyla changed significantly in YBG-fed mice, but not in control mice (supplemental Fig. 4A).

YBG treatment resulted in increase in the abundances of Bacteroidetes (P=0.049), and

Verrucomicrobia (Mean±SD: control=13.0±0.33 vs YBG=10.9.7±0.69) phyla members and

decrease in Firmicutes ($P<0.001$) and Proteobacteria ($P<0.001$) in their fecal samples.

Interestingly, at genus level, abundances of many major microbial communities are altered

significantly after YBG treatment (**Fig. 3A**). Compared to pre-treatment time-point, significant

reductions in the abundance of microbial communities belonging to multiple genera including

Ruminococcus (phylum: Firmicutes; P=0.003), *Lactobacillus* (phylum: Firmicutes; P=0.02),

Oscillospira (phylum: Firmicutes; $P<0.001$), *Dehalobacterium* (phylum: Firmicutes; $P<0.006$),

and *Odoribacter* (phylum: Firmicutes; $P<0.001$) were observed after YBG treatment. On the other

hand, abundance of microbial communities belonging to many genera including

Parabacteroidetes (phylum: Bacteroidetes; $P<0.005$), *Prevotella* (phylum: Bacteroidetes;

$P<0.002$), *Sutterella* (phylum: Proteobacteria; $P<0.007$), and *Bifidobacterium* (phylum:

Actinobacteria; P=0.048) were increased after YBG treatment. Interestingly fecal samples

collected after YBG treatment showed lower gut microbial α-diversity / species richness

($P<0.001$) compared to pretreatment time-point (Supplemental Fig. 4B). Further, β-diversity (Bray

Curtis distance) analysis of fecal microbiota showed significant distance (P=0.007) in the

clustering / separation of samples collected before and after YBG treatment (Supplemental Fig.

4C). As shown in **Fig. 3C**, OTU-based prediction of metabolic functions of gut microbes revealed

overrepresentation of many pathways including that of carbohydrate metabolism (P=0.031),

glycan biosynthesis and metabolism (P=0.027) and biosynthesis of secondary metabolites

(P=0.01) in B6 mouse gut microbiota after YBG treatment compared to pre-treatment time-point.

YBG treatment results in modulation of small and large intestinal immune phenotype in B6 mice

Since YBG treatment altered gut microbiota and suppressed colitis susceptibility in B6 mice, we determined if YBG treatment alone has an impact on gut immune phenotype. Supplemental fig. 5 shows that the primarily small intestine (ileum), but not large intestine (colon), of mice that received YBG for 30 consecutive days expressed significantly higher levels of pro-inflammatory cytokine TNFα ($P=0.0039$) and immune regulatory enzyme Raldh1A2 ($P=0.001$) compared to that of untreated counterparts. On the other hand, significantly higher levels of IL10 expression was detected in both ileum ($P=0.0017$) and colon ($P=0.0042$) of YBG treated mice compared to controls. We then determined the T phenotype of gut mucosa in mice that received YBG for 30 consecutive days by assessing the Foxp3+ T cell frequencies and cytokine production. As shown in supplemental Fig. 6A and **Fig. 4A**, significantly higher frequencies of Foxp3+ T cells were detected in mesenteric LNs (MLNs) ($P=0.016$), small intestine lamina propria (SiLP) ($P=0.0079$) and large intestinal lamina propria (LiLP) ($P=0.0159$) of YBG treated mice compared to controls. Further, compared to that of controls, small intestinal cells from YBG treated mice produced significantly higher amounts of IL10 ($P=0.016$) and IL17 ($P=0.0044$). On the other hand, colonic immune cells of YBG treated mice produced higher amounts of IL10 ($P=0.0079$), but not IL17 or IFNγ.

We then determined the influence of microbiota on YBG consumption-associated immune modulation in B6 mice after depleting gut bacteria using broad-spectrum antibiotics (Supplemental fig. 7). Depletion of gut bacteria alone caused reduction in Foxp3+ T cell frequencies in both control and YBG recipient mice (supplemental Fig. 6B and **Fig. 4B**), relative to their counterparts with intact microbiota (shown in Fig. 4A). Further, SiLP cells, but not LiLP cells, from microbiota-

depleted YBG recipient mice showed relatively higher frequencies of Foxp3+ T cells ($P=0.0041$)

compared to their control counterparts. On the other hand, small intestine of antibiotic-treated

YBG-recipients showed higher IL10 ($P=0.0087$), IL17 ($P=0.0159$) and IFNγ ($P=0.041$)

production, compared to microbiota-depleted controls. Immune cells from large intestine of YBG-

treated microbiota-depleted mice produced significantly higher amounts of IL17 ($P=0.0034$) and

IFNγ ($P=0.0039$), but not IL10, compared to control counterparts.

YBG treatment results in increased immune regulatory SCFA production in B6 mice

Since fermentation of non-digestible sugars by colonic bacteria can result in the generation of

metabolites including SCFA, fecal SCFA levels in control and YBG treated B6 mice with intact

and depleted gut microbiota were determined. As observed in Fig. 5A, significantly higher fecal

acetic acid ($P=0.016$), propionic acid ($P=0.026$) and butyric acid ($P=0.013$), and lower valeric

acid ($P=0.036$) levels were detected in YBG-treated mice compared to their control counterparts.

On the other hand, depletion of gut microbiota not only resulted in the elimination of YBG-

treatment associated effect on SCFA production, but caused profound suppression of overall SCFA

levels in both control and YBG treated mice (Fig. 5B) compared to that of mice with intact

microbiota (Fig. 5A).

YBG treatment associated protection of B6 mice from colitis is gut microbe-dependent

To determine if the YBG treatment associated protection of B6 mice from colitis is microbiota

dependent, YBG treated and control B6 mice were given broad spectrum antibiotics to deplete gut

microbiota and given DSS to induce colitis as depicted in **Fig. 6A**. As observed in **Fig. 6B and C**,

control and YBG treated mice with intact gut microbiota showed weight loss and disease severity

comparable to that of Fig. 1. Importantly, control mice with microbiota depletion showed relatively

lesser weight loss and disease severity compared to their control counterparts, and the difference

was statistically significant (P=0.012) only at 10 week time point (**Fig. 6B; left panel**). However,

colon inflammation scores were not statistically significant in these mice (**Fig. 6C; left panel**). On

the other hand, YBG-treated, microbiota-depleted mice showed significantly higher weight loss

and disease severity starting at day 6 time-point (P=0.016) and more severe thereafter (P<0.01)

compared to untreated mice with intact microbiota. Further, YBG treated, microbiota depleted

mice showed higher colon inflammation (P=0.047) compared to control mice **(Fig. 6C; right**

panel). Examination of cytokine profiles of immune cells from the colons revealed significantly

higher production of pro-inflammatory IFNγ (P<0.001) and IL17 (P<0.001), and comparable anti-

inflammatory IL10, production in YBG-treated, microbiota depleted mice compared to YBG

treated mice with intact gut microbiota **(Fig. 6D)**. However, cells from microbiota depleted control

mice showed diminished IFNγ production compared to cells from controls with intact microbiota

(P<0.001).

Discussion

Literature suggests that certain complex dietary polysaccharides such as BGs of different microbial

and plant origin affect the host immune function (44-48). While previous studies using various BG

preparations and BG rich diets have shown that they have the ability to suppress gut inflammation

in colitis models (10-14), others have shown that BGs aggravate the colitis severity (15). Further,

while it has been shown that genetic deletion of a BG interacting receptor, Dectin-1, does not affect

the course of murine experimental colitis, another group found Dectin-1 deficiency increased

susceptibility to colitis (16, 17). Hence, although BGs from different sources can vary structurally

and functionally (1), the true impact of orally administering BGs on gut inflammation and the

associated mechanisms is unknown, and the previously reported outcomes are non-conclusive

without the use of well-defined, high pure, BGs such as the YBG used in this study. Here, we show

that prolonged oral administration of high pure YBG prior to DSS treatment to induce colitis

results in significantly lowered severity of gut inflammation in B6 mice. This diminished

susceptibility to DSS colitis is associated altered gut microbiota composition and function favoring

polysaccharide metabolism, higher immune regulatory SCFA production, and enhanced gut

immune regulation.

Intriguingly, we found that treatment with high pure YBG at disease induction stage alone has no

beneficial effects, but only prolonged oral pretreatment with this agent suppresses colitis

susceptibility. This shows that well defined BGs can promote protection from colitis, but it

depends on gradually acquired enhanced immune regulation, perhaps mediated by altered / YBG

shaped gut microbiota. In fact, although the role of immune regulation promoted directly by YBG

in the gut mucosa cannot be ruled out, we found that prolonged oral treatment using YBG causes

changes in the gut microbial composition. YBG consumption resulted in reduced abundance of

Firmicutes and increased abundance of Bacteroidetes and Verrucomicrobia phyla, which include

many polysaccharide fermenting bacteria (49, 50). However, YBG treatment also diminished gut

microbial diversity / species richness, suggesting that YBG treatment induced effect is due to

selective enrichment of microbial communities that promote immune regulation, but not by

increasing the diversity of gut microbiota. Further, predictive functional profiling of fecal

microbiota showed overrepresentation of metabolic pathways linked to carbohydrate metabolism,

glycan biosynthesis and metabolism, and biosynthesis of secondary metabolites after YBG

treatment compared to pre-treatment time-point. More importantly, our novel observations that

fecal levels of immune regulatory microbial SCFA (51, 52) such as acetic acid, propionic acid

and butyric acid produced by gut microbiota are higher in YBG treated mice suggest that altered

microbiota is functionally superior in terms of their ability to promote immune regulation. This

notion has been validated by diminishes levels of these SCFAs, Treg frequencies in the gut

mucosa, and elimination of YBG treatment induced effect upon depletion of gut microbiota.

Further, the need for prolonged pre-treatment, which could potentially help gradual shaping of gut

microbiota structure and function, for protection from colitis is another indication that altered gut

microbiota is the key contributor of protection from colitis in YBG fed mice.

Colonic microbial communities can ferment non-digestible dietary polysaccharides to use them as

energy sources for their growth, and also produce small metabolites such as SCFA that enhance

gut immune regulation (16, 49, 51, 53-58). Previous studies have shown that gut colonization by

specific commensal microbial communities causes induction and expansion of Foxp3+ Tregs in

the intestinal and systemic compartments (59-61). Further, gut bacteria play a critical role in

maintaining peripheral tolerance and gut immune homeostasis (62, 63). However, it is not known

if BG shaped gut microbes promote Treg induction and/or expansion. Our observations that YBG

treatment increases the production of microbial SCFA, which are known to enhance gut integrity

and immune regulation (51, 55), suggest that microbiota does contribute to YBG treatment

associated increase in overall immune regulation. Importantly, abundances of microbes belonging

to polysaccharide degrading and well as host beneficial communities such as *Parabacteroides,*

Sutterella, Prevotella, Bifidobacterium and Akkermansia are increased upon prolonged YBG treatment.

Moreover, microbial depletion and reduced SCFA production resulted in increased production of

pro-inflammatory cytokines IFNγ and IL17 in the distal gut of YBG treated mice substantiating

the notion that microbes contributes to enhance immune regulation and diminished susceptibility

to colitis when pre-treated with this CDP.

Although this study is focused on the impact of YBG treatment on colitis susceptibility and distal

gut, we found that, unlike the large intestine, small intestine of YBG treated mice, with intact and

depleted microbiota, have relatively higher frequencies of Tregs compared to untreated controls.

This suggests that YBG-treatment associated increase in Treg abundance in small and large

intestinal compartments may involve different mechanisms. It is possible that while prolonged

treatment with YBG, at the employed human relevant dose, interacts directly with small intestinal

mucosa to enhance immune regulation, YBG treatment associated enhanced immune regulation in

the large intestine may be primarily microbiota (and SCFA) dependent. Of note, our recent report

showed that treatment of mice with a higher dose of this YBG caused strong pro-inflammatory

immune response in both small and large intestine (29), suggesting that use of higher doses of this

CDP is not advisable to ameliorate susceptibility to gut inflammation.

Although we previously reported that Dectin-1 dependent interaction of high pure YBG with

immune cells could promote a combination of both pro-inflammatory and immune regulatory

immune responses (19, 27) under normal circumstances, our current study suggest that such direct

interaction can produce primarily pathogenic response under depleted microbiota and pro-

inflammatory conditions such as a chemical insult using DSS. In fact it has been shown that the

degree and type of immune cell response to BGs depends on the cytokine milieu and

microenvironment (64). It is possible that YBG is degraded by colonic bacteria and produce

immune regulatory SCFA along with minimizing its direct interaction with large intestinal mucosa.

On the other hand, microbial depletion could diminish YBG degradation and SCFA production.

Further, DSS which causes epithelial damage can expose immune cells for gut mucosa for direct

interaction with YBG resulting in dominant pro-inflammation immune response. This is evident

from our observation that YBG treated, microbiota depleted mice presents more severe colitis

compared to YBG treated mice with intact microbiota, and control groups of mice with intact and

depleted microbiota.

Overall, our studies using YBG found that this CDP can promote gut immune regulation by

altering gut microbiota, increasing SCFA production, enhancing the overall immune regulation

and gut health under normal circumstances suggesting the prebiotic value of BGs. Importantly,

only pre-conditioning of the intestinal microenvironment in terms of altering gut microbiota

composition and function, which enhances SCFA production and immune regulation, by treating

with YBG diminishes susceptibility to chemical induced colitis. We show that YBG treatment has

no benefits in ameliorating the severity of ongoing gut inflammation, as in the case of UC. Hence,

with respect to the colitis, health benefits of YBG could be limited to those who are susceptible to,

but not yet developed, the disease rather than those who have established disease. We employed

microbiota depletion approach in our study primarily to understand the mechanism by which YBG

treatment enhances gut immune regulation and diminish colitis susceptibility. Nevertheless, our

observations suggest that consumption of YBG, as a dietary supplement, by those who are under

antibiotic therapy could produce adverse effects such as gut inflammation. Of note, utility of

antibiotic treatments have been considered for ameliorating the severity of gut inflammation and

for managing UC symptoms (65, 66). Our observations suggest the possibility that consumption

of YBG as a dietary supplement by UC patients who are under such therapy will result in

aggravated, rather than ameliorated, disease severity. In conclusion, this study, which used high

pure YBG, not only demonstrates the dietary supplement value of BGs in promoting gut health

and the associated mechanisms, but also show that these CDPs may not be beneficial in

suppressing ongoing gut inflammation and they produce adverse effects when used in combination

with antibiotic therapy.

Conflict of Interest statement: Authors do not have any conflict(s) of interest to disclose.

Author contribution: R.B., B.J, and J.S. researched and analyzed data, and R.G. researched and

analyzed the data and edited the manuscript; C.V. designed experiments, researched and analyzed

data, and wrote/reviewed/edited manuscript.

Acknowledgments: This work was supported by internal funds from MUSC, National Institutes

of Health (NIH) grants: R21AI133798 (NIAID), and R21AI133798 and R21AI136339-02S1

Administrate supplements (Office of Dietary Supplements) to CV. The authors are thankful to the

Histology core of Pathology department of MUSC for the histology service, and Genomic center

of MUSC for 16S rRNA gene sequencing service. Dr. Vasu is the guarantor of this work and as

such, has full access to all the data in the study and takes responsibility for integrity of the data

and accuracy of data analysis.

 Legends for Figures

FIGURE 1: Effect of pre-treatment with YBG (A) on induced colitis associated weight loss (B),

colon shortening (C) and colon inflammation (D) in B6 mice. (A) Cartoon depicting experimental

design is shown. (B) Body weights of individual animals were measured every day starting at day

30 and changes in percentage of body weights, relative to initial body weight, are shown. (C)

Images of representative colons (left panel) and colon lengths (right panel) of euthanized mice are

shown. (D) H&E stained distal colon sections were evaluated for inflammation and images of

representative sections (left panel) and inflammation severity scores (right panel) are shown. (B,

right panels of C, D) Mean±SDs (n=8). *, **,*** Different from control, P<0.05, P<0.01, P<0.001

respectively.

FIGURE 2: Effect of YBG pre-treatment on Treg (A) and cytokine expressing (B) CD4+ T

frequencies in, and cytokine secretion (C) by, MLN cells of B6 mice with DSS induced colitis.

(A,B) Frequencies of CD4+ cells in the MLN single cell suspension that express indicated factor

determined by FACS are shown. C) Concentrations of indicated cytokines secreted by MLN cells

upon *ex vivo* stimulation using anti-CD3 antibody are shown. Mean±SDs (n=5) are shown for all

panels. *,** Different from control, P<0.05, P<0.01 respectively.

FIGURE 3: Impact of oral administration of YBG on the composition of gut microbiota at genus

level (A) and predictive functions (B) in B6 mice. (A) OTU data of 16S rRNA gene sequences of

fecal pellet collected before (day 0) and 30 days after YBG treatment (day 30) were compiled to

genus level based upon the percentage identity to reference sequences (i.e. >97% identity) and

heatmap was generated using sequence counts of major genera (those with >1 % of total

sequences) that showed statistically significant differences. (B) OUT-biom data was used for

predicting gene functional categories using PICRUSt application and the level 2 categories of

KEGG pathway that showed statistically significant differences are shown. n=5 for both panels.

*, **,*** Different from control, $P<0.05$, $P<0.01$, $P<0.001$ respectively.

FIGURE 4: Impact of oral administration of YBG on the Treg frequency and cytokine secretion

profile of immune cells in B6 mice with intact (A) and depleted (B) gut microbiota. (A, B) Foxp3+

CD4 cell frequencies in gut associated lymphoid tissues (MLN, SiLP and LiLP) determined by

FACS (left panels) and the levels of indicated cytokines secreted by MLN upon stimulation using

anti-CD3 antibody determined by multiplex assay (right panels) are shown. Mean±SDs (n=5) are

shown for all panels. *,** Different from control, $P<0.05$, $P<0.01$ respectively.

FIGURE 5: Impact of oral administration of YBG on the fecal SCFA levels in B6 mice with intact

(A) and depleted (B) gut microbiota. (A, B) SCFA levels in fecal pellets collected from control

and YBG treated mice that were left alone or given broad spectrum antibiotics are shown.

Mean±SDs (n=5) are shown for all panels. *Different from control, $P<0.05$.

FIGURE 6: Effect of microbiota depletion (A) in DSS colitis induced weight loss (B), colon

inflammation (C) and cytokine secretion by intestinal immune cells (D) in control and YBG treated

B6 mice. (A) Cartoon depicting experimental design is shown. (B) Body weights of individual

animals were measured every day starting at day 30 and changes in percentage of body weights,

relative to initial body weight, are shown. (C) H&E stained distal colon sections were evaluated

for inflammation and the severity scores are shown. D) Concentrations of indicated cytokines

secreted by colonic immune cells upon *ex vivo* stimulation using anti-CD3 antibody are shown.

(B,C, D) Mean±SDs (n=5) are shown. *, **,*** Different from control, $P<0.05$, $P<0.01$, $P<0.001$

respectively.

References:

1.   Synytsya A, Novak M. Structural analysis of glucans. Ann Transl Med. 2014 Feb;2:17.
2.   Chen J, Seviour R. Medicinal importance of fungal beta-(1-->3), (1-->6)-glucans.
Mycological research. 2007 Jun;111:635-52.
3.   Vannucci L, Krizan J, Sima P, Stakheev D, Caja F, Rajsiglova L, Horak V, Saieh M.
Immunostimulatory properties and antitumor activities of glucans (Review). International journal
of oncology. 2013 Aug;43:357-64.
4.   El Khoury D, Cuda C, Luhovyy BL, Anderson GH. Beta glucan: health benefits in obesity
and metabolic syndrome. J Nutr Metab. 2012;2012:851362.
5.   Mosikanon K, Arthan D, Kettawan A, Tungtrongchitr R, Prangthip P. Yeast beta-Glucan
Modulates Inflammation and Waist Circumference in Overweight and Obese Subjects. J Diet
Suppl. 2017 Mar 4;14:173-85.
6.   Zou Y, Liao D, Huang H, Li T, Chi H. A systematic review and meta-analysis of beta-
glucan consumption on glycemic control in hypercholesterolemic individuals. Int J Food Sci Nutr.
2015;66:355-62.
7.   Fuller R, Moore MV, Lewith G, Stuart BL, Ormiston RV, Fisk HL, Noakes PS, Calder PC.
Yeast-derived beta-1,3/1,6 glucan, upper respiratory tract infection and innate immunity in older
adults. Nutrition. 2017 Jul - Aug;39-40:30-5.
8.   McFarlin BK, Carpenter KC, Davidson T, McFarlin MA. Baker's yeast beta glucan
supplementation increases salivary IgA and decreases cold/flu symptomatic days after intense
exercise. J Diet Suppl. 2013 Sep;10:171-83.
9.   Leentjens J, Quintin J, Gerretsen J, Kox M, Pickkers P, Netea MG. The effects of orally
administered Beta-glucan on innate immune responses in humans, a randomized open-label
intervention pilot-study. PloS one. 2014;9:e108794.
10.   Kamiya T, Tang C, Kadoki M, Oshima K, Hattori M, Saijo S, Adachi Y, Ohno N, Iwakura
Y. beta-Glucans in food modify colonic microflora by inducing antimicrobial protein, calprotectin,
in a Dectin-1-induced-IL-17F-dependent manner. Mucosal Immunol. 2018 May;11:763-73.
11.   Liu B, Lin Q, Yang T, Zeng L, Shi L, Chen Y, Luo F. Oat beta-glucan ameliorates dextran
sulfate sodium (DSS)-induced ulcerative colitis in mice. Food Funct. 2015 Nov;6:3454-63.
12.   Chung MY, Hwang JT, Kim JH, Shon DH, Kim HK. Sarcodon aspratus Extract
Ameliorates Dextran Sulfate Sodium-Induced Colitis in Mouse Colon and Mesenteric Lymph
Nodes. J Food Sci. 2016 May;81:H1301-8.
13.   Sun Y, Shi X, Zheng X, Nie S, Xu X. Inhibition of dextran sodium sulfate-induced colitis
in mice by baker's yeast polysaccharides. Carbohydr Polym. 2019 Mar 1;207:371-81.
14.   Zhou M, Wang Z, Chen J, Zhan Y, Wang T, Xia L, Wang S, Hua Z, Zhang J.
Supplementation of the diet with Salecan attenuates the symptoms of colitis induced by dextran
sulphate sodium in mice. Br J Nutr. 2014 May 28;111:1822-9.
15.   Heinsbroek SE, Williams DL, Welting O, Meijer SL, Gordon S, de Jonge WJ. Orally
delivered beta-glucans aggravate dextran sulfate sodium (DSS)-induced intestinal inflammation.
Nutr Res. 2015 Dec;35:1106-12.
16.   Heinsbroek SE, Oei A, Roelofs JJ, Dhawan S, te Velde A, Gordon S, de Jonge WJ. Genetic
deletion of dectin-1 does not affect the course of murine experimental colitis. BMC Gastroenterol.
2012 Apr 16;12:33.
17.   Iliev ID, Funari VA, Taylor KD, Nguyen Q, Reyes CN, Strom SP, Brown J, Becker CA,
Fleshner PR, et al. Interactions between commensal fungi and the C-type lectin receptor Dectin-1
influence colitis. Science. 2012 Jun 8;336:1314-7.

18. Bacon JS, Farmer VC, Jones D, Taylor IF. The glucan components of the cell wall of
baker's yeast (Saccharomyces cerevisiae) considered in relation to its ultrastructure. The
Biochemical journal. 1969 Sep;114:557-67.
19. Gudi R, Perez N, Johnson BM, Sofi MH, Brown R, Quan S, Karumuthil-Melethil S, Vasu
C. Complex dietary polysaccharide modulates gut immune function and microbiota, and promotes
protection from autoimmune diabetes. Immunology. 2019 Feb 3.
20. Chevrier G, Mitchell PL, Rioux LE, Hasan F, Jin T, Roblet CR, Doyen A, Pilon G, St-
Pierre P, et al. Low-Molecular-Weight Peptides from Salmon Protein Prevent Obesity-Linked
Glucose Intolerance, Inflammation, and Dyslipidemia in LDLR-/-/ApoB100/100 Mice. J Nutr.
2015 Jul;145:1415-22.
21. Mahler M, Bristol IJ, Leiter EH, Workman AE, Birkenmeier EH, Elson CO, Sundberg JP.
Differential susceptibility of inbred mouse strains to dextran sulfate sodium-induced colitis. Am J
Physiol. 1998 Mar;274:G544-51.
22. Taghipour N, Molaei M, Mosaffa N, Rostami-Nejad M, Asadzadeh Aghdaei H, Anissian
A, Azimzadeh P, Zali MR. An experimental model of colitis induced by dextran sulfate sodium
from acute progresses to chronicity in C57BL/6: correlation between conditions of mice and the
environment. Gastroenterol Hepatol Bed Bench. 2016 Winter;9:45-52.
23. Eichele DD, Kharbanda KK. Dextran sodium sulfate colitis murine model: An
indispensable tool for advancing our understanding of inflammatory bowel diseases pathogenesis.
World J Gastroenterol. 2017 Sep 7;23:6016-29.
24. DeVoss J, Diehl L. Murine models of inflammatory bowel disease (IBD): challenges of
modeling human disease. Toxicol Pathol. 2014 Jan;42:99-110.
25. Dennehy KM, Ferwerda G, Faro-Trindade I, Pyz E, Willment JA, Taylor PR, Kerrigan A,
Tsoni SV, Gordon S, et al. Syk kinase is required for collaborative cytokine production induced
through Dectin-1 and Toll-like receptors. European journal of immunology. 2008 Feb;38:500-6.
26. Dennehy KM, Willment JA, Williams DL, Brown GD. Reciprocal regulation of IL-23 and
IL-12 following co-activation of Dectin-1 and TLR signaling pathways. European journal of
immunology. 2009 May;39:1379-86.
27. Karumuthil-Melethil S, Gudi R, Johnson BM, Perez N, Vasu C. Fungal beta-glucan, a
Dectin-1 ligand, promotes protection from type 1 diabetes by inducing regulatory innate immune
response. J Immunol. 2014 Oct 1;193:3308-21.
28. Montufar-Solis D, Klein JR. An improved method for isolating intraepithelial lymphocytes
(IELs) from the murine small intestine with consistently high purity. J Immunol Methods. 2006
Jan 20;308:251-4.
29. Gudi R, Perez N, Johnson BM, Sofi MH, Brown R, Quan S, Karumuthil-Melethil S, Vasu
C. Complex dietary polysaccharide modulates gut immune function and microbiota, and promotes
protection from autoimmune diabetes. Immunology. 2019 May;157:70-85.
30. Sofi MH, Johnson BM, Gudi RR, Jolly A, Gaudreau MC, Vasu C. Polysaccharide A-
Dependent Opposing Effects of Mucosal and Systemic Exposures to Human Gut Commensal
Bacteroides fragilis in Type 1 Diabetes. Diabetes. 2019 Oct;68:1975-89.
31. Chassaing B, Aitken JD, Malleshappa M, Vijay-Kumar M. Dextran sulfate sodium (DSS)-
induced colitis in mice. Curr Protoc Immunol. 2014 Feb 4;104:Unit 15 25.
32. Erben U, Loddenkemper C, Doerfel K, Spieckermann S, Haller D, Heimesaat MM, Zeitz
M, Siegmund B, Kuhl AA. A guide to histomorphological evaluation of intestinal inflammation
in mouse models. Int J Clin Exp Pathol. 2014;7:4557-76.
33. Gudi RR, Karumuthil-Melethil S, Perez N, Li G, Vasu C. Engineered Dendritic Cell-
Directed Concurrent Activation of Multiple T cell Inhibitory Pathways Induces Robust Immune
Tolerance. Sci Rep. 2019 Aug 19;9:12065.

34.   Johnson BM, Gaudreau MC, Al-Gadban MM, Gudi R, Vasu C. Impact of dietary deviation
on disease progression and gut microbiome composition in lupus-prone SNF1 mice. Clin Exp
Immunol. 2015 Aug;181:323-37.
35.   Sofi MH, Gudi R, Karumuthil-Melethil S, Perez N, Johnson BM, Vasu C. pH of drinking
water influences the composition of gut microbiome and type 1 diabetes incidence. Diabetes. 2014
Feb;63:632-44.
36.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer
N, Pena AG, Goodrich JK, et al. QIIME allows analysis of high-throughput community sequencing
data. Nature methods. 2010 May;7:335-6.
37.   Wixon J, Kell D. The Kyoto encyclopedia of genes and genomes--KEGG. Yeast. 2000
Apr;17:48-55.
38.   Parks DH, Beiko RG. Identifying biologically relevant differences between metagenomic
communities. Bioinformatics. 2010 Mar 15;26:715-21.
39.   Hevia A, Milani C, Lopez P, Cuervo A, Arboleya S, Duranti S, Turroni F, Gonzalez S,
Suarez A, et al. Intestinal dysbiosis associated with systemic lupus erythematosus. mBio.
2014;5:e01548-14.
40.   Zhang H, Liao X, Sparks JB, Luo XM. Dynamics of gut microbiota in autoimmune lupus.
Applied and environmental microbiology. 2014 Dec 15;80:7551-60.
41.   Parks DH, Tyson GW, Hugenholtz P, Beiko RG. STAMP: statistical analysis of taxonomic
and functional profiles. Bioinformatics. 2014 Nov 1;30:3123-4.
42.   Dhariwal A, Chong J, Habib S, King IL, Agellon LB, Xia J. MicrobiomeAnalyst: a web-
based tool for comprehensive statistical, visual and meta-analysis of microbiome data. Nucleic
Acids Res. 2017 Jul 3;45:W180-W8.
43.   Zhao G, Nyman M, Jonsson JA. Rapid determination of short-chain fatty acids in colonic
contents and faeces of humans and rats by acidified water-extraction and direct-injection gas
chromatography. Biomed Chromatogr. 2006 Aug;20:674-82.
44.   Ramberg JE, Nelson ED, Sinnott RA. Immunomodulatory dietary polysaccharides: a
systematic review of the literature. Nutr J. 2010 Nov 18;9:54.
45.   Hong F, Yan J, Baran JT, Allendorf DJ, Hansen RD, Ostroff GR, Xing PX, Cheung NK,
Ross GD. Mechanism by which orally administered beta-1,3-glucans enhance the tumoricidal
activity of antitumor monoclonal antibodies in murine tumor models. Journal of immunology.
2004 Jul 15;173:797-806.
46.   Suzuki I, Tanaka H, Kinoshita A, Oikawa S, Osawa M, Yadomae T. Effect of orally
administered beta-glucan on macrophage function in mice. Int J Immunopharmacol. 1990;12:675-
84.
47.   Suzuki I, Hashimoto K, Ohno N, Tanaka H, Yadomae T. Immunomodulation by orally
administered beta-glucan in mice. Int J Immunopharmacol. 1989;11:761-9.
48.   Volman JJ, Ramakers JD, Plat J. Dietary modulation of immune function by beta-glucans.
Physiol Behav. 2008 May 23;94:276-84.
49.   Flint HJ, Scott KP, Duncan SH, Louis P, Forano E. Microbial degradation of complex
carbohydrates in the gut. Gut Microbes. 2012 Jul-Aug;3:289-306.
50.   Martinez-Garcia M, Brazel DM, Swan BK, Arnosti C, Chain PS, Reitenga KG, Xie G,
Poulton NJ, Lluesma Gomez M, et al. Capturing single cell genomes of active polysaccharide
degraders: an unexpected contribution of Verrucomicrobia. PloS one. 2012;7:e35314.
51.   Marino E, Richards JL, McLeod KH, Stanley D, Yap YA, Knight J, McKenzie C, Kranich
J, Oliveira AC, et al. Gut microbial metabolites limit the frequency of autoimmune T cells and
protect against type 1 diabetes. Nature immunology. 2017 May;18:552-62.

52.   D'Souza WN, Douangpanya J, Mu S, Jaeckel P, Zhang M, Maxwell JR, Rottman JB,
Labitzke K, Willee A, et al. Differing roles for short chain fatty acids and GPR43 agonism in the
regulation of intestinal barrier function and immune responses. PloS one. 2017;12:e0180190.
53.   Porter NT, Martens EC. The Critical Roles of Polysaccharides in Gut Microbial Ecology
and Physiology. Annu Rev Microbiol. 2017 Sep 8;71:349-69.
54.   Temple MJ, Cuskin F, Basle A, Hickey N, Speciale G, Williams SJ, Gilbert HJ, Lowe EC.
A Bacteroidetes locus dedicated to fungal 1,6-beta-glucan degradation: Unique substrate
conformation drives specificity of the key endo-1,6-beta-glucanase. The Journal of biological
chemistry. 2017 Jun 23;292:10639-50.
55.   Geirnaert A, Calatayud M, Grootaert C, Laukens D, Devriese S, Smagghe G, De Vos M,
Boon N, Van de Wiele T. Butyrate-producing bacteria supplemented in vitro to Crohn's disease
patient microbiota increased butyrate production and enhanced intestinal epithelial barrier
integrity. Sci Rep. 2017 Sep 13;7:11450.
56.   Turunen K, Tsouvelakidou E, Nomikos T, Mountzouris KC, Karamanolis D, Triantafillidis
J, Kyriacou A. Impact of beta-glucan on the faecal microbiota of polypectomized patients: a pilot
study. Anaerobe. 2011 Dec;17:403-6.
57.   Carlson JL, Erickson JM, Hess JM, Gould TJ, Slavin JL. Prebiotic Dietary Fiber and Gut
Health: Comparing the in Vitro Fermentations of Beta-Glucan, Inulin and Xylooligosaccharide.
Nutrients. 2017 Dec 15;9.
58.   Holscher HD. Dietary fiber and prebiotics and the gastrointestinal microbiota. Gut
Microbes. 2017 Mar 4;8:172-84.
59.   Furusawa Y, Obata Y, Fukuda S, Endo TA, Nakato G, Takahashi D, Nakanishi Y, Uetake
C, Kato K, et al. Commensal microbe-derived butyrate induces the differentiation of colonic
regulatory T cells. Nature. 2013 Dec 19;504:446-50.
60.   Ye J, Qiu J, Bostick JW, Ueda A, Schjerven H, Li S, Jobin C, Chen ZE, Zhou L. The Aryl
Hydrocarbon Receptor Preferentially Marks and Promotes Gut Regulatory T Cells. Cell Rep. 2017
Nov 21;21:2277-90.
61.   Chang PV, Hao L, Offermanns S, Medzhitov R. The microbial metabolite butyrate
regulates intestinal macrophage function via histone deacetylase inhibition. Proceedings of the
National Academy of Sciences of the United States of America. 2014 Feb 11;111:2247-52.
62.   Belkaid Y, Harrison OJ. Homeostatic Immunity and the Microbiota. Immunity. 2017 Apr
18;46:562-76.
63.   Steinmeyer S, Lee K, Jayaraman A, Alaniz RC. Microbiota metabolite regulation of host
immune homeostasis: a mechanistic missing link. Curr Allergy Asthma Rep. 2015 May;15:24.
64.   Municio C, Alvarez Y, Montero O, Hugo E, Rodriguez M, Domingo E, Alonso S,
Fernandez N, Crespo MS. The response of human macrophages to beta-glucans depends on the
inflammatory milieu. PloS one. 2013;8:e62016.
65.   Nitzan O, Elias M, Peretz A, Saliba W. Role of antibiotics for treatment of inflammatory
bowel disease. World J Gastroenterol. 2016 Jan 21;22:1078-87.
66.   Turner D, Levine A, Kolho KL, Shaoul R, Ledder O. Combination of oral antibiotics may
be effective in severe pediatric ulcerative colitis: a preliminary report. J Crohns Colitis. 2014
Nov;8:1464-70.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

A

B

Fig. 5

Fig. 6