1 **Systematic revision and redefinition of the genus *Scirrotherium* Edmund**

2 **and Theodor, 1997 (Cingulata, Pampatheriidae): Implications for the**

3 **origin of pampatheriids and the evolution of the South American lineage**

4 **including *Holmesina***

5

6 Kevin Jiménez-Lara [a, b]

7 [a] División de Paleontología de Vertebrados, Museo de La Plata, Facultad de Ciencias

8 Naturales y Museo, Universidad Nacional de La Plata, Paseo del Bosque s/n, B1900FWA

9 La Plata, Argentina.

10 [b] CONICET, Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina.

11 e-mail: kjimenezlara@fcnym.unlp.edu.ar

12

13 **Abstract**

14 The intrageneric relationships of the pampatheriid genus *Scirrotherium* and its affinities

15 with supposedly related genera, i.e., *Kraglievichia* and *Holmesina*, are revised through

16 parsimony phylogenetic analyses and new comparative morphological descriptions. For this

17 work, unpublished material of pampatheriids (numerous osteoderms, one partial skull and a

18 few postcranial bones) from Neogene formations of Colombia was analyzed. The results

19 show that *Scirrotherium* is paraphyletic if we include all its referred species, i.e.,

20 *Scirrotherium hondaensis*, *S. carinatum* and *S. antelucanus*. The species *S. carinatum* is

21 closer to *Kraglievichia paranensis* than to *S. hondaensis* or *S. antelucanus*, therefore the

new name *K. carinatum* comb. nov. is proposed. The relationship among *S. hondaensis* and

S. antelucanus could not be resolved, so these species should be designated in aphyly. In

spite of failing to recover *S. hondaensis* and *S. antelucanus* as one single clade, here is

preferred to maintain the generic name *Scirrotherium* in both species based on diagnostic

evidence. New emended diagnoses for *Scirrotherium*, *S. hondaensis* and *Kraglievichia* are

provided. The genus *Holmesina* was found to be monophyletic and positioned as the sister

clade of *Scirrotherium* + *Kraglievichia*. The evolutionary and biogeographic implications

of the new phylogeny and taxonomic re-arrangements are discussed. A possible geographic

origin of the family Pampatheriidae and *Scirrotherium* in low latitudes of South America as

early as Early Miocene times is claimed. The South American ancestor or sister taxon of

Holmesina is predicted to be morphologically more similar to *Scirrotherium* than to

Kraglievichia.

Keywords: Pampatheriidae

Scirrotherium

Kraglievichia

Holmesina

Great American Biotic Interchange

Neogene

1. Introduction

The pampatheriids (Pampatheriidae) are a morphologically conservative extinct clade of glyptodontoid cingulates (Xenarthra: Glyptodontoidea *sensu* McKenna and Bell 1997) with medium-to-large body sizes (Edmund 1985; Góis et al. 2013). They were distributed from the Neogene to the Early Holocene in numerous localities in South America (their native range), Central America, Mexico and the United States (Edmund 1985; Vizcaíno et al. 1998; Rincón et al. 2014; Góis et al. 2015 and references therein). As the modern armadillos (Dasypodidae), pampatheriids have a flexible carapace characterized by the presence of three transverse bands of imbricated osteoderms, which form a kind of "articulation" between the scapular and pelvic shields (Edmund 1985). The pampatheriids also have multiple features, especially in their skull and mandible, which, collectively, define them as the sister group of glyptodontids –Glyptodontidae (Gaudin 2004; Gaudin and Wible 2006; Billet et al. 2011; Delsuc et al. 2012), namely, a deep horizontal mandibular ramus, a laterally-directed zygomatic root, a transversely wide glenoid fossa, rugose pterygoids, among others (Gaudin and Wible, 2006).

The fossil record of Pampatheriidae is mainly represented by isolated specimens, most of which are osteoderms and, to a lesser extent, skulls, mandibles and postcranial bones. Relatively complete and articulated skeletons are uncommon (Edmund 1985; Góis 2013). Due to this fact, the systematics of this group has historically been based on osteoderm characters (Edmund 1985, 1987; Góis et al. 2013), as has often been the case with other cingulate clades. Overall, nearly 20 pampatheriid species and seven genera are known (Góis 2013). The latter conform two possible subfamilial lineages: (1) that including to the genera *Plaina* and *Pampatherium*; and (2) that comprising the genera *Scirrotherium*, *Kraglievichia* and *Holmesina* (Edmund 1985). However, there is no any published

phylogenetic analysis on the relationships among the different pampatheriid genera in the

scientific literature. Only Góis (2013) performed a phylogenetic analysis of these taxa, but

his results have not been published so far. In Góis's consensus tree, it was corroborated the

hypothesis on the two subfamilial lineages as suggested by Edmund (1985).

The genus *Scirrotherium* is the oldest undoubted pampatheriid (Góis et al. 2013; Rincón et

al. 2014) and one of the four Miocene genera (the others are *Kraglievichia* Castellanos

1927; *Vassallia* Castellanos, 1927; and *Plaina* Castellanos, 1937). This taxon was

originally described by Edmund and Theodor (1997) based on craniomandibular,

postcranial and osteoderm specimens collected in the Middle Miocene (Serravalian)

sedimentary sequence of the La Venta area, in southwestern Colombia. These authors

suggested that the type and only known species in that time, *Scirrotherium hondaensis*, has

plesiomorphic traits in its osteological morphology which are expected for its antiquity.

Additionally, they highlighted the morphological similarity of *S. hondaensis* with the

species *Vassallia minuta* (Late Miocene of southern and central South America; De Iullis

and Edmund 2002), more than with any other pampatheriid.

Later, Góis et al. (2013) described a second species for *Scirrotherium*, *S. carinatum*, from

the Late Miocene (Tortonian) of northeastern and southern Argentina and northwestern

Brazil. In northeastern Argentina (Province of Entre Ríos), *S. carinatum* is found in the

same basal stratigraphic levels ("Conglomerado Osífero", literally meaning 'bone-bearing

conglomerate') of the Ituzaingó Formation as the middle-sized pampatheriid *Kraglievichia*

paranensis (Góis et al. 2013; Scillato-Yané et al. 2013), a taxon clearly distinct but not

distantly related to *Scirrotherium*, as previously indicated. *Scirrotherium carinatum*, based

exclusively on osteoderms from different regions of the armored carapace, has an estimated

body size comparable or slightly smaller than that of *S. hondaensis* (Góis et al. 2013).

The phylogenetic analysis conducted by Góis (2013) recovered a polytomy involving *S.*

carinatum and *S. hondaensis*, one of these species or both being the sister taxon/taxa of the

clade *Kraglievichia + Holmesina* (except *H. floridanus*). Considering that *Scirrotherium* is

the oldest known pampatheriid, it is notorious the non-basal position of the *Scirrotherium*

species in the cladogram of Góis (2013). Instead, these species are closely allied with

terminal taxa, i.e., *Holmesina* spp. (Edmund 1985, 1987; Gaudin and Lyon 2017). If this

result is correct, it would indicate a significant ghost lineage at the base of Pampatheriidae.

Góis (2013) found phylogenetic support for *Scirrotherium* through one single

synapomorphy, i.e., the presence of deep longitudinal depressions in the osteoderms. This

is a distinctive feature of *S. carinatum* but the longitudinal depressions in *S. hondaensis* are

relatively shallow. Interestingly, this putative synapomorphy is actually shared by *K.*

paranensis. Góis (2013) explained the lack of phylogenetic resolution for *Scirrotherium* by

noting the fragmentary character of the available fossil specimens of *S. hondaensis*.

However, unlike the latter species, the skull, mandible and any postcranial bone of *S.*

carinatum are unknown (Góis et al. 2013).

Laurito and Valerio (2013) reported new pampatheriid material from the Late Miocene

(Tortonian to Messinian) of Costa Rica, which they assigned to a new species, *S.*

antelucanus. This species, the largest referred to as *Scirrotherium* so far (body size

comparable or slightly smaller than that of *K. paranensis*; Laurito and Valerio 2013), is

based on osteoderms and some postcranial bones (femoral fragments and metatarsals). The

occurrence of *S. antelucanus* in the Late Miocene of southern Central America suggests

that *Scirrotherium* took part in the late Cenozoic biotic interchanges of the Americas earlier

than any other pampatheriid (i.e., *Plaina*, *Pampatherium*, *Holmesina*; Woodburne 2010),

invading tropical North America ("North America" is defined here as all the continental

territories north of the ancient location of the main geographic barrier between the

Americas during the early Neogene, i.e., the Central American Seaway in northwestern

Colombia) before the definitive closing of the Panamanian Land Bridge (PLB) ca. 3 mya

(Schmidt 2007; Coates and Stallard 2013; O'dea et al. 2016; Jaramillo 2018).

Recently, the occurrence of isolated osteoderms designated as *Scirrotherium* sp. or cf.

Scirrotherium has been reported in several contributions on fossil vertebrate assemblages

from the Neogene of Venezuela and Peru. On the basis of these discoveries, the geographic

and chronological distribution of the genus has been expanded in such a way that this taxon

is now known for the Early and Late Miocene (Burdigalian and Tortonian) of northwestern

Venezuela (Rincón et al. 2014; Carrillo-Briceño et al. 2018) and Late Miocene (Tortonian)

of eastern Peru (Antoine et al. 2016).

Assuming all the previous taxonomic assignments are correct, the latitudinal range of

Scirrotherium, from southern Central America to Patagonia (southern Argentina), is the

widest latitudinal range of a Miocene pampatheriid, comparable only with those of the Plio-

Pleistocene forms *Pampatherium* and *Holmesina* (Scillato-Yané et al. 2005). This

biogeographic evidence provides support for the hypothesis that *Scirrotherium* inhabited

varied environments within its latitudinal range, and, consequently, that it probably had a

relatively high ecological flexibility (Góis et al. 2013).

Despite the progress in the systematic and biogeographic research on *Scirrotherium*, a new

reevaluation of several fundamental hypotheses about this taxon is needed, including its

taxonomic definition and evolutionary relationships with other pampatheriid genera. Using

parsimony phylogenetic analyses and comparative morphological descriptions of new

pampatheriid remains from the Neogene of Colombia, this contribution reevaluates the

taxonomic status of *Scirrotherium* and its relationships with supposedly allied genera, i.e.,

Kraglievichia and *Holmesina*. Accordingly, I suggest a new taxonomic and nomenclatural

reorganization, with emended diagnoses for *Scirrotherium* and *Kraglievichia*.

Finally, considering the systematic reanalysis, I develop a model of biogeographic

evolution for the lineage *Scirrotherium-Kraglievichia-Holmesina*. From this model, I draw

out new hypotheses on the geographic origin of Pampatheriidae and the late Cenozoic

dispersal events of pampatheriids to/from North America, including a possible re-entry

event to South America for the species *S. antelucanus*.

2. Material and methods

2.1. Taxonomic sampling

I studied 12 species of pampatheriids attributed to six different genera. These species, in

alphabetic order, are: *Holmesina floridanus* Robertson, 1976; *H. major* Lund, 1842; *H.

occidentalis* Hoffstetter, 1952; *H. paulacoutoi* Cartelle and Bohórquez, 1985; *H.

septentrionalis* Leidy, 1889; *Kraglievichia paranensis* Ameghino, 1888; *Pampatherium

humboldtii* Lund, 1839; *Plaina intermedia* Ameghino, 1888; *Scirrotherium antelucanus*

Laurito and Valerio, 2013; *S. carinatum* Góis, Scillato-Yané, Carlini and Guilherme, 2013;

S. hondaensis Edmund and Theodor, 1997; and *Vassallia minuta* Moreno and Mercerat,

1891. Unidentified pampatheriid material (MUN STRI 16718 and 38064; see the section

Institutional abbreviations) from the Castilletes Formation in Colombia (see below), which

is referred to as "Castilletes specimens", was also included in this selection.

Among the former nominal species, I follow Góis (2013) in considering *Vassallia maxima*

as a junior synonym of *Pl. intermedia*. The only species of *Holmesina* not included in this

study were *H. rondoniensis* Góis, Scillato-Yané, Carlini and Ubilla, 2012 and *H. cryptae*

Moura, Góis, Galliari and Fernandes 2019. In the case of *H. rondoniensis*, its exclusion is

based on a preliminary phylogenetic analysis in which it was identified as a "wildcard"

taxon obscuring phylogenetic resolution as a result of lack of information on the osteoderm

features of this species. On the other hand, the scientific article in which *H. cryptae* was

described has been very recently published (Moura et al. 2019), after the completion of this

work. Consequently, it was preferred to omit this species here. *Tonnicinctus mirus* Góis,

González Ruiz, Scillato-Yané and Soibelzon, 2015 was also not included in this analysis

given that this species is considered a highly divergent taxon without any apparent

substantial interest with respect to the systematic issues here addressed.

2.2. Morphological description of the specimens

The osteological morphology of the selected species was re-examined from direct

observations of specimens and published/unpublished descriptions (Simpson 1930,

Castellanos 1937; Edmund 1985, 1987; Edmund and Theodor 1997; Góis 2013; Góis et al.

2013; Laurito and Valerio 2013; Scillato-Yané et al. 2013; Góis et al. 2015; Gaudin andg

Lyon 2017). Naturally, according to the objectives of this research, during the revision of

material I focused on the species *S. antelucanus*, *S. carinatum* and *S. hondaensis*, and,

additionally, species of genera considered closely allied to *Scirrotherium*, i.e., *K.*

paranensis and *Holmesina* spp. (particularly *H. floridanus*; Appendix S1 of the

Supplementary Material).

On the other hand, new undescribed cranial, postcranial and osteoderm specimens were

also used to reevaluate the morphological variability of *Scirrotherium.* This material comes

from five Neogene geological units of Colombia (Fig. 1): (1) Castilletes Formation (Early

to Middle Miocene, late Burdigalian-Langhian), Municipality of Uribia, Department of La

Guajira; (2) La Victoria Formation (late Middle Miocene, Serravalian), Municipality of

Villavieja, Department of Huila; (3) Villavieja Formation (late Middle Miocene,

Serravalian), Municipality of Villavieja, Department of Huila; (4) Sincelejo Formation

(Late Miocene-Early Pliocene, Messinian-Zanclean), Municipality of Los Palmitos,

Department of Sucre; (5) Ware Formation (Late Pliocene, Piacenzian), Municipality of

Uribia, Department of La Guajira. For detailed lithological descriptions and

chronostratigraphic data on these formations, the reader is referred to the following

references: Moreno et al. (2015) for the Castilletes and Ware Formations; Guerrero (1997),

Flynn et al. (1997) and Anderson et al. (2016) for the La Victoria and Villavieja

formations; and Flinch (2003), Villarroel and Clavijo (2005), Bermúdez et al. (2009) and

Alfaro and Holz (2014) for the Sincelejo Formation. The new fossils are deposited in the

Paleontological Collection of the Museo Mapuka de la Universidad del Norte, Barranquilla,

Colombia, except those collected in the La Victoria and Villavieja Formations. The latter

are housed at the Museo de Historia Natural La Tatacoa, La Victoria Town, Municipality of

Villavieja, Department of Huila, Colombia.

Cranial measurements, all taken on the midline of the skull (dorsally or ventrally), follow

Góis (2013). The anatomical terminology for osteoderms is based on the proposals by Góis

et al. (2013). All the measurements were taken with a digital caliper.

2.3. Character matrix

I used exclusively cranial, dental and osteoderm characters, given that postcranial bones of

most species of Pampatheriidae are poorly known (Góis 2013). The character construction

was based on personal observations and previous quantitative and qualitative analyses of

the interspecific, intergeneric and familial morphological variability of pampatheriids (e.g.,

Edmund 1985, 1987; Góis 2013; Góis et al. 2013; Laurito and Valerio 2013; Gaudin and

Lyon, 2017). Overall, a matrix of 27 characters (Appendix S2 of the Supplementary

Material) was built and managed on Mesquite version 2.75 (Maddison and Maddison

2010). If present, parsimony-uninformative characters were used to define potential

autapomorphies of the taxa under study.

2.4. Cladistic analyses

Parsimony analyses under schemes of equal weights and implied weights (characters

reweighted *a posteriori*; see below) were performed in PAUP* version 4.0a142 (Swofford

2015). In both weighting schemes, the species *P. humboldtii*, *Pl. intermedia* and *V. minuta*

were selected as outgroup (sister group). This selection is based on the hypothesis about

subfamilial relationships of Pampatheriidae by Edmund (1985) and the phylogeny of Góis

(2013). All the characters were treated as unordered. The analyses were run with the branch

and bound search to estimate maximum parsimony trees. The criterion for character

optimization was DELTRAN (see Gaudin [2004] for justification of this configuration).

For reordering of branches, the algorithm of tree-bisection-reconnection (TBR) branch

swapping was used. The topological results of most parsimonious trees were summarized

through strict consensus trees.

In this work, the methodology of implied weights was intended to mitigate potential biases

arising from limited number of characters (especially osteoderm characters, as a

consequence of the evolutionary trend in Pampatheriidae towards a simplification of the

ornamentation in comparison with that in other cingulate clades, e.g., Glyptodontidae) and

the effect of homoplastic characters (Goloboff et al. 2008; Goloboff 2014). Characters were

reweighted using the rescaled consistency index (mean value) of the equally-weighted

parsimony analysis (see Ausich et al. 2015 and references therein for justification of the use

of rescaled consistency index for implied-weights parsimony analyses). A default concavity

value ($k = 3$) was selected (Goloboff et al. 2018). Three successive rounds of character

reweighting were needed until identical set of strict consensus trees were found in two

consecutive searches (Swofford and Bell 2017).

Node stability for the strict consensus trees was evaluated using bootstrap resampling

procedures (branch and bound search with 100 replicates). The software FigTree v1.4.3

(http://tree.bio.ed.ac.uk/software/figtree/) was used as a graphical viewer and editor for the

cladograms.

2.5. Taxonomic and nomenclatural criteria

I applied a taxonomic and nomenclatural criterion reasonably, but not strictly, constrained

by the phylogeny. This implies looking for a natural classification (i.e., based on

monophyletic groups) without ignoring possible limitations of the phylogenetic inference

related to the available information in the fossil record and major morphological gaps.

Additionally, open nomenclature was used to indicate taxonomic uncertainty when

necessary, following the general recommendations of Bengston (1988) and updated

definitions by Sigovini et al. (2016) for the qualifiers of this semantic tool of taxonomy.

2.6. Institutional abbreviations

CFM, Museo Nacional de Costa Rica, Colección de fósiles de la sección de Geología, San

José, Costa Rica; FMNH, Field Museum Natural History, Chicago, Illinois, USA; MACN,

Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Colección de

Paleovertebrados, Ciudad Autónoma de Buenos Aires, Argentina; MCL, Museu de

Ciências Naturais da Pontifícia Universidade Católica de Minas Gerais, Belo Horizonte,

Brazil; MG-PV, Museo Provincial de Ciencias Naturales Dr. Ángel Gallardo, Rosario,

Argentina; MHD-P, Museo Histórico Departamental de Artigas, Artigas, Uruguay; MLP,

Museo de La Plata, La Plata, Argentina; MUN STRI, Museo Mapuka de la Universidad del

Norte, Colección de paleontología, Barranquilla, Colombia; ROM, Royal Ontario Museum,

Toronto, Canadá; UCMP, University of California Museum of Paleontology, Berkeley,

California, USA; UF, Florida Museum of Natural History, Gainesville, Florida, USA;

UZM, Universitets Zoologisk Museum, Copenhagen, Denmark; VPPLT, Museo de Historia

Natural La Tatacoa, Colección de paleontología, La Victoria Town, Huila, Colombia.

2.7. Anatomical abbreviations

FL, frontal bone length; GFL, greatest femoral length; GSL, greatest skull length; LUR,

length of the upper teeth row; Mf, upper molariform; mf, lower molariform; NL, nasal bone

length; PAL, parietal bone length; PL, hard palate length; TTW, maximum width at the

third trochanter of the femur; DW, maximum width of the femoral distal epiphysis.

3. Results

The parsimony analysis with equal weights obtained 107 most parsimonious trees (MPTs),

each one of these with a tree length of 44 steps (consistency index = 0.909; retention index

= 0.907; rescaled consistency index = 0.825). The strict consensus tree from these MPTs

(Fig. 2 (A); tree length = 52; consistency index = 0.769; retention index = 0.721; rescaled

consistency index = 0.555) is not fully resolved because it has two polytomies. One of these

polytomies involves the species *S. hondaensis*, *S. antelucanus* and *H. floridanus*, whereas

the other is formed by *H. septentrionalis*, *H. major*, *H. paulacoutoi* and *H. occidentalis*.

Three clades were recovered (excluding that of the entire ingroup): (1) All the ingroup taxa

except "Castilletes specimens"; (2) *S. carinatum* + *K. paranensis*; and (3) *Holmesina* spp.

except *H. floridanus*. On the other hand, the parsimony analysis with implied weights

yielded 30 most parsimonious trees (MPTs) with a tree length of 109 weighted steps

(consistency index = 0.982; retention index = 0.982; rescaled consistency index = 0.964).

The strict consensus tree from the MPTs (Fig. 2(B); tree length = 91; consistency index =

0.978; retention index = 0.980; rescaled consistency index = 0.959), like that produced by

the equally weighted approach, is not fully resolved. Again, two polytomies resulted, but in

this case the polytomy including *S. hondaensis*, *S. antelucanus* and *H. floridanus* was

altered. The latter taxon was placed as the basal-most *Holmesina* species. The polytomy

formed by *H. septentrionalis*, *H. major*, *H. paulacoutoi* and *H. occidentalis* was

unmodified. As a consequence of the relocation of *H. floridanus* within the topology, four

clades were recovered: (1) All the ingroup taxa except "Castilletes specimens"; (2) *S.*

carinatum + *K. paranensis*; (3) *Holmesina* spp.; and (4) *H. septentrionalis*, *H. major* + *H.*

paulacoutoi + *H. occidentalis*. According to the two schemes of weighting for the

parsimony analyses, *Scirrotherium* is paraphyletic if it is comprised of *S. antelucanus*, *S.*

hondaensis and *S. carinatum*. *Scirrotherium carinatum* is closer to *K. paranensis* than to *S.*

hondaensis or *S. antelucanus*. The relationship among *S. hondaensis* and *S. antelucanus* is

not resolved in either of the two strict consensus trees.

Under the equal weights analysis, all the nodes that include taxa of interest show (nearly)

significant stability (resampling frequencies ca. 70 or greater). As expected, a similar result

was obtained in the bootstrap resampling under implied weights, but with improved

frequencies for nearly all the branches (except the basal-most branch, whose frequency

decreased slightly from that of the former bootstrapping [85 to 83]).

4. Systematic paleontology

Xenarthra Cope, 1889

Cingulata Illiger, 1811

Glyptodontoidea Gray, 1869

Pampatheriidae Paula Couto, 1954

Scirrotherium Edmund and Theodor, 1997

LSID. urn:lsid:zoobank.org:act:313358B5-3B1F-4902-8C2E-BB07CFCBEE18

Type species: *Scirrotherium hondaensis* Edmund and Theodor, 1997 by original

designation.

Emended diagnosis: A pampatheriid of small-to-middle body size that can be

distinguished from other pampatheriids by the following combination of features: thin non-

marginal fixed osteoderms (ca. 3.5–7 mm in thickness); slightly to moderately rough

external surface of osteoderms; external surface of osteoderms with a sharp and uniformly

narrow longitudinal central elevation; longitudinal central elevation from superficial to

well-elevated; (very) shallow longitudinal depressions with gentle slope towards the

marginal elevations; usually one single, transversely elongated row of large foramina in the

anterior margin of fixed osteoderms; maximum number of foramina per row between 6 and

11.

Remarks: The taxonomic status of *Scirrotherium* is saved from invalidity by paraphyly by

exclusion of the species '*S.*' *carinatum* from the genus (see below). However, according to

the preferred phylogenetic hypothesis presented here, i.e., the strict consensus tree from the

parsimony analysis under implied weights (Fig. 2(B)), the other two referred species of

Scirrotherium, *S. antelucanus* and *S. hondaensis*, should be designated in aphyly because

they do not have resolved relationship between them (see Ebach and Williams 2010 for

details about the phylogenetic concept of aphyly). Until new evidence becomes available,

maintenance of the taxonomic validity of *Scirrotherium*, as defined here, is based on the

emended diagnosis of this taxon, which is partially built from ambiguous synapomorphies,

as well as from qualitative and quantitative morphological differences with other genera.

Scirrotherium hondaensis Edmund and Theodor, 1997

LSID. urn:lsid:zoobank.org:act:E3B83181-91D6-44C8-90C0-BBAACEC2CDEE

Holotype: UCMP 40201, incomplete skull and left hemimandible (Edmund and Theodor,

1997).

Type locality and horizon: Municipality of Villavieja, Department of Huila, Colombia. La

Victoria Formation, upper Middle Miocene, Serravalian (Edmund and Theodor, 1997).

Emended differential diagnosis. Pampatheriid of small body size that differs from other

pampatheriids based on this unique combination of characters: external surface of

osteoderms with ornamentation (especially the longitudinal central elevation and marginal

elevations), in general terms, more protuberant than in *S. antelucanus*, but less than in

Kraglievichia; size range of fixed osteoderms smaller than in *S. antelucanus* and similar to

that in *Kraglievichia carinatum* comb. nov. (= ‘*S.*’ *carinatum*; Góis et al. 2013; see below);

fixed osteoderms generally thicker than in *K. carinatum* comb. nov. but less than in *K.*

paranensis, similar to *S. antelucanus*; anterior foramina smaller than in *S. antelucanus*;

anterior foramina in fixed osteoderms usually aligned in one individual row, although

infrequently these osteoderms show an extra, short or reduced row of anterior foramina;

two rows of anterior foramina in mobile osteoderms, similar to *Vassallia* (Góis 2013); mf9

incipiently bilobed; frontals prominently convex in lateral view, with this convexity

positioned posterior to the insertion of the anterior root of the zygomatic arch; anterior root

of the zygomatic arch posterolaterally projected with respect to the main body of maxilla.

Referred material: VPPLT 004, several fixed osteoderms; VPPLT 264, several fixed

osteoderms and one semi-mobile osteoderm; VPPLT 348, tens of fixed and (semi) mobile

osteoderms; VPPLT 701, several fixed osteoderms; VPPLT 706, one anterior skull, one

femoral diaphysis, one ulna without distal epiphysis, several vertebrae and numerous fixed

and (semi) mobile osteoderms; VPPLT 1683 - MT 18, several fixed and (semi) mobile

osteoderms; UCMP 39846, one proximal femoral epiphysis, one left calcaneum and one

left astragalus. All the osteoderms referred to as *S. hondaensis* are illustrated in Fig. 3.

Other important specimens are illustrated in Figs. 4–7 and 9(C).

Occurrence: VPPLT 004, 264, 701, 706 and (partially) 1683 - MT 18 were collected in the

La Victoria Formation, upper Middle Miocene (Serravalian; see Figs. 3–6 for more details

on the stratigraphic provenance of individual specimens), while the UCMP 39846 and part

of VPPLT 1683 - MT18 comes from the Villavieja Formation, upper Middle Miocene

(Serravalian).

Description: For the original and detailed description of this species, including its

osteoderms, see Edmund and Theodor (1997). See the Tables 1 and 2 for an updated

compilation of osteoderm measurements of the *Scirrotherium* species and comparisons

with those of related taxa. Below there are descriptions of osteological structures and traits

incompletely known or unknown for *S. hondaensis* so far.

Skull: The holotype of *S. hondaensis* UCMP 40201 includes a very fragmentary

skull. This specimen does not preserve the anterior end of the rostrum, much of the orbit

(both dorsally and ventrally), part of the upper dental series, ear region, braincase and

occipital region. In comparison, the skull VPPLT 706 (Fig. 4) described here, is more

complete, despite the fact that it is also missing some structures. This new, small skull (see

Table 3 for morphometric comparisons) is relatively well preserved from the orbit to the

anterior end of rostrum, except for the anterior zygomatic arch and nasal bones. It also has a

less deformed rostrum than the holotypic skull. The general aspect of the new skull is

similar to those of other pampatheriids. In lateral view, it is markedly depressed towards its

anterior end. In dorsal view, it is also tapered towards its anterior tip, where it ends abruptly

(Castellanos, 1937). Proportionally, the rostrum is shorter than that of *K. paranensis* and

even more than that of *H. floridanus*. In lateral view, the facial process of premaxilla is less

well defined than that of *H. floridanus*, although the premaxilla-maxilla suture has a convex

form like the latter species (Gaudin and Lyon, 2017). The antorbital fossa is arranged more

vertically than those of *K. paranensis* and *H. floridanus*. The lacrimal is, proportionally, the

largest among pampatheriids. This bone precludes the frontomaxillary contact (restricted

contact in the skull of *K. paranensis*). The dorsal contribution of the lacrimal to the orbit is,

proportionally, greater than in *H. floridanus* but similar to that in *K. paranensis*. The

anterior root of zygomatic arch is projected posterolaterally, unlike other pampatheriids

whose skull is known, where the anterior root projects laterally. The frontals show a

conspicuous convexity in a position posterior to the insertion of the anterior root of

zygomatic arch. In dorsal view, the frontals are more anteroposteriorly elongated and more

laterally expanded than in *K. paranensis*, but are similar to those of *Holmesina* spp. In

ventral view, the hard palate has a wide aspect, since the rostrum is shortened in

comparison with other pampatheriids. Only two anterior molariforms are preserved (the left

Mf1 and the right Mf2), so inferences about upper dentition are made from the alveoli. The

upper dental series, as in all the members of the family, is comprised of by nine

molariforms. Of these, the last five (Mf5-Mf9) are bilobed. The anterior molariforms (Mf1-

Mf4) converge anteriorly, but do not imbricate. The latter teeth are rounded to elliptical,

similarly to the condition observed in *H. floridanus*. They also are less mesiodistally

elongated than in *K. paranensis*. The molariforms with greatest occlusal area are the Mf5

and Mf6. The occlusal area of the upper molariforms decrease distally from the fifth and

sixth molariforms to the ninth, as in all the pampatheriids. The Mf9 is the smallest of lobed

molariforms and has the least degree of lobulation (elliptical shape for this tooth in the type

material of *S. hondaensis*, according to Edmund and Theodor 1997). In ventral view,

VPPLT 706 is characterized by a gradual transverse widening of the palatal portion of the

maxilla from the level of the anterior border of the Mf5. The anterior portion of the

palatines is preserved up to a level slightly posterior to the Mf9. The maxilla-palatine suture

is not recognizable.

Femur: This bone in *S. hondaensis* was largely unknown so far, except for a pair of

epiphyses (proximal and distal) from a left femur in the UCMP collections (UCMP 39846).

VPPLT 706 preserves a left femur (Fig. 5(A–D)) without epiphyses (apparently it is not the

same bone from which the previously noted epiphyses came). Thus, the description of all

these anatomical elements allows for a reconstruction of most aspects of femoral anatomy.

The estimated proximo-distal length of this bone is ca. 162 mm, and its transverse width at

the third trochanter is 27.6 mm. These morphometric values are the smallest known for

femora of Pampatheriidae (Table 4). They are comparable only to those of MLP 69-IX-8-

13A which was referred to as *K*. cf. *paranensis* (Góis 2013; Scillato-Yané et al. 2013). The

femoral head is hemispheric and the greater trochanter is less high than that of *K*. cf.

paranensis, but similar to the condition observed in *H. floridanus*. However, the greater

trochanter has a more tapered proximal end than in the latter species. In *S. hondaensis*, the

lesser trochanter is less mediolaterally expanded than in *K.* cf. *paranensis*. The femoral

diaphysis is less curved mediolaterally than in *K.* cf. *paranensis*, similar to that of *H.*

floridanus. The laterodistal border of the femur is more curved than in *H. floridanus*,

similar to that of *K.* cf. *paranensis*. The third trochanter is, proportionally, larger than that

of *K.* cf. *paranensis*, but is smaller than that of *H. floridanus*. The patellar facets are less

defined or delimited than those of *K.* cf. *paranensis*. In *S. hondaensis* these facets are

oriented toward the center of the anterior surface of the distal epiphysis, rather than

laterally as in *K.* cf. *paranensis* and *H. floridanus*.

Ulna: This bone is described here for the first time in *S. hondaensis*. A right ulna

(Fig. 5(E–H)) missing part of the diaphysis and the distal epyphysis is preserved in VPPLT

706. The estimated proximo-distal length is 89.5 mm. The olecranon is elongated and

protuberant. In medial view, it is less proximally tapered than that of *H. floridanus*. The

lateral entrance to the trochlear notch is very restricted, similarly to *Holmesina*. Likewise, it

is less proximo-distally elongated than that of *H. floridanus*. Proximally, at the level of the

trochlear notch, the posterior border is uniformly convex, not slightly concave as it is in *H.*

floridanus. The depression for the insertion of the anconeus muscle is deep and proximally

located, like that of *Holmesina*.

Vertebrae: Several vertebrae are also preserved in VPPLT 706 (Fig. 6). One of

these is a thoracic vertebra and five are caudal vertebrae, of which four are articulated in

two pairs and one is an isolated distal caudal vertebra. The body of the thoracic vertebra is

eroded anteriorly, as are the anterior zygapophyses. Posteriorly, the vertebral body has an

outline similar to that of other pampatheriids. Notably, two ventrolateral apophyses are

projected from the vertebral body. Although fragmented, the neural spine of the same

vertebra appears to be proportionally shorter than in *H. floridanus*. The anterior caudal

vertebrae have a posteriorly oriented and tall neural spine. The transverse processes are

relatively short.

Astragalus: Edmund and Theodor (1997) mentioned the existence of numerous

undetermined postcranial elements whose description was to be postponed to a subsequent

publication. However, that description was never published. This postcranial material from

the UCMP collections includes a left astragalus (UCMP 39846; Fig. 7(A–D)). In dorsal

view, the lateral trochlea is considerably larger than the medial trochlea. The astragalar

head is bulging, spherical, and almost uniformly convex. There is a shallow concavity in

the dorsal margin of the astragalar head whose function has been not determined, but it

could be for tendinous insertion. The astragalar neck is well-differentiated, similar to that in

Holmesina. In ventral view, the facets of articulation with the calcaneum, i.e., ectal and

sustentacular, are widely separated, as one would expect by observing their counterparts on

the calcaneum (Edmund 1987). The ectal facet is noticeably larger than the sustentacular

facet, in contrast to the condition in *H. floridanus*. The ectal facet is kidney-shaped and the

sustentacular facet has a sub-oval shape. Both of them are concave, especially the ectal

facet, which is very deep. The sustentacular facet is located in a central position within the

astragalar neck.

Calcaneum: This bone is other postcranial element not described by Edmund and

Theodor (1997) for *S. hondaensis*. UCMP 39846 is a well-preserved left calcaneum (Fig.

7(E–F)). It has proximo-distal length of 54.12 mm and a width at the level of facets (ectal

and sustentacular) of ca. 10.2 mm. These values are the smallest known for calcanei

referred to as Pampatheriidae. The only species whose calcaneum is comparable in size to

that of *S. hondaensis* is *H. floridanus*. The calcaneum of the latter species is slightly more

proximo-distally elongated than in *S. hondaensis*, but it is roughly two times wider at the

level of the ectal and sustentacular facets. This means that the calcaneum of *H. floridanus* is

more robust than that of *S. hondaensis*. The calcaneal head is anteroposteriorly elongated,

like in *H. floridanus* and unlike the proportionally short calcaneal head of *H.*

septentrionalis. The anterior end of the calcaneal head is less shortened than that of

Holmesina. The calcaneum of *S. hondaensis* shows no contact between the ectal and

sustentacular facets, similar to the condition in species of *Holmesina* other than *H.*

floridanus (Góis 2013). These facets are slightly convex and they are separated by a

moderately deep and very wide groove, i.e., the sulcus tali (see below). Like *H. floridanus*,

the facets are highly asymmetrical, but the condition in *S. hondaensis* is even more

exaggerated, as the ectal facet is much larger than the sustentacular facet. As in the

astragalus described above, the ectal facet is kidney-shaped and the sustentacular facet is

sub-oval. The ectal facet is located at an oblique angle with respect to the long axis of the

tuber calcanei, unlike that of *H. floridanus*. Like other pampatheriids, the calcaneal

sustentacular facet of *S. hondaensis* is located anterior to the anterior border of the ectal

facet. However, this facet is even more anteriorly placed in *S. hondaensis* than in other

pampatheriid species as consequence of its particularly wide sulcus tali. Posteriorly, the

calcaneal tuber is not massive in comparison with Pleistocene species of *Holmesina* (e.g.,

H. septentrionalis), but rather mediolaterally compressed, particularly towards its dorsal

side.

Scirrotherium antelucanus Laurito and Valerio, 2013

LSID. urn:lsid:zoobank.org:act:225CD304-3B63-4B55-B8B8-33B46C90A194

Holotype. CFM-2867, mobile osteoderm (Laurito and Valerio, 2013).

Type locality and horizon. San Gerardo de Limoncito, County of Coto Brus, Province of Puntarenas, Costa Rica. Upper Curré Formation, Upper Miocene (Laurito and Valerio). For further information about the stratigraphic position of the Curré Formation, see these references: Lowery 1982; Yuan 1984; Rivier 1985; Kolarsky et al. 1995; Alvarado et al. 2009; Aguilar et al. 2010; Obando 2011. There are no published absolute ages for this geological unit.

Diagnosis. Unmodified (see Laurito and Valerio, 2013; p. 47).

Referred material. MUN STRI 36880, an isolated fixed osteoderm (Fig. 8).

Occurrence. Upper Sincelejo Formation, Upper Miocene to Pliocene (Messinian to Zanclean). El Coley Town, Municipality of Los Palmitos, Department of Sucre, Colombia. For further information about the stratigraphic position of the Sincelejo Formation, see these references: Flinch 2003; Villarroel and Clavijo 2005; Bermúdez et al. 2009; and Alfaro and Holz 2014. There are no published absolute ages for this geological unit.

Remarks. The fixed osteoderm MUN STRI 36880, possibly from the of the pelvic shield, is assigned to the species *S. antelucanus* on the basis of the following observations: (I) the area and thickness of this osteoderm (linear measurements in millimetres: anteroposterior length = 34.91; transverse width = 24; thickness = 4.45; approximate area = 837.8 mm^2) are

within the range of variability for comparable osteoderms of *S. antelucanus* and exceed the

known values of area for most of the same kind of osteoderms for *S. hondaensis*; (II) the

external surface is relatively smooth; (III) the anterior margin is wide; (IV) the anterior

foramina are larger (2–3 millimetres of diameter) than in *S. hondaensis*, like *S. antelucanus*

from Costa Rica; (V) the number of anterior foramina (9) is within the range of variability

for *S. antelucanus* (7–10 for quadrangular osteoderms, like the specimen here described),

but greater than the range for *S. hondaensis*; (VI) poorly elevated longitudinal central

elevation, like in some osteoderms of *S. antelucanus* (the longitudinal central elevation is

generally more elevated in *S. hondaensis*; see Laurito and Valerio 2013).

aff. *Scirrotherium*

Referred material: MUN STRI 16718 (Fig. 9(A)), fixed osteoderm of the scapular shield;

MUN STRI 38064 (Fig. 9(E)), undetermined fixed osteoderm; MUN STRI 16719 (Fig.

9(G)), mobile osteoderm fragmented in its anterior margin.

Occurrence: Castilletes Formation, upper Lower Miocene to lower Middle Miocene, upper

Burdigalian to Langhian). Localities of Makaraipao, Kaitamana and Patajau Valley

(localities with numbers 390093, 430202 and 390094 in Moreno et al. 2015, respectively),

Municipality of Uribia, Department of La Guajira, Colombia.

Description: The fixed osteoderm of the scapular shield MUN STRI 16718 (Fig. 9(A)) is

relatively large and has a pentagonal outline. Its linear measurements in millimetres are:

anteroposterior length = 45.02; transverse width = 33.41; thickness = 6.66. These values

imply that this osteoderm has greater area than any other known area size for osteoderms

referred to as *Scirrotherium* (Table 1), including those of the osteoderms of the larger

Scirrotherium species, i.e., *S. antelucanus* (see Appendix 1 in Laurito and Valerio 2013).

Rather, this osteoderm is similar in size to those reported for *H. floridanus*. The external

surface of the osteoderm MUN STRI 16718 is punctuated by numerous diminutive pits,

like *S. hondaensis* and *S. antelucanus*. The surface does not have a recognizable

longitudinal central elevation nor longitudinal depressions, so that the osteoderm has a

flattened appearance, similar to that of several osteoderms of *S. antelucanus* (Laurito and

Valerio 2013). In contrast, the marginal elevations are easily identifiable. These ridges are

relatively low and narrow. There are foramina with a nearly homogeneous large size in the

anterior margin. They are aligned in two well defined rows. The most anterior row has five

foramina and the posterior one has six. Collectively, the two rows of foramina rows are

equivalent to ca. 25% of the anteroposterior length of the osteoderm. In *S. hondaensis*, the

rows of foramina in fixed osteoderm, when present, this percentage comprises less than

20%.

The osteoderm MUN STRI 38064 (Fig. 9(E)) does not appears to be a non-marginal

osteoderm. It has a trapezoidal outline and the following measurements in millimetres:

anteroposterior length = 39.08; transverse width = 39.55; thickness: 5.98. These values are

within the range of variability of *S. antelucanus*. This osteoderm has two long rows of

anterior foramina in which the posterior row seems to extend partially over the anterior

lateral margins, unlike the anterior foramina row(s) in *S. hondaensis* and *S. antelucanus*.

The most anterior row of foramina is formed by eight foramina and the posterior row has

11 foramina. In both of these rows, the foramina are of similar size, although a few are

smaller. The rows of foramina diverge on the left lateral margin and within the resultant

space between these rows is located a large and isolated foramen. This osteoderm does not

have a recognizable longitudinal central elevation nor longitudinal depressions, i.e., it is

flattened. Its marginal elevations are narrow and poorly elevated. The foramina of the

lateral margins are smaller than most of anterior foramina. As a consequence of

preservation factors, the pits on the external surface are not present.

The osteoderm MUN STRI 16720 is a partial mobile osteoderm (Fig. 9(G)) with an

elongated rectangular shape. Its linear measurements in millimetres are: anteroposterior

length (incomplete by fragmentation) = 45.68; transverse width = 30.69; thickness = 6.96.

The external surface is convex and without a longitudinal central elevation or longitudinal

depressions. The anterior margin shows a set of foramina not clearly aligned in rows.

Remarks: With current evidence, the osteoderms MUN STRI 16718 and MUN STRI

38064 could not be confidently assigned to *Scirrotherium*. This taxonomic decision is

supported by several arguments. First, morphologically, the osteoderms referred here to aff.

Scirrotherium are more similar to those of *S. hondaensis* and *S. antelucanus* than to any

other osteoderms of known pampatheriids. The osteoderms of aff. *Scirrotherium* differ with

respect to the osteoderms of *S. hondaensis* and *S. antelucanus* in three characteristics: (I) a

greater number of anterior foramina and/or greater development of two rows from these

foramina; (II) longitudinal central elevation possibly absent, i.e., the presence of a flattened

external surface; (III) larger maximum osteoderm area. Of these features, the third one (III)

is the least ambiguous, i.e., the maximum area of fixed osteoderms exceeds those of the

osteoderms of *S. hondaensis* and *S. antelucanus*. Comparatively, the first and second (I and

II) characteristics are more ambiguous considering that similar conditions were also

observed in *S. hondaensis* and *S. antelucanus*. These conditions are described as follows.

Some infrequent osteoderms of *S. hondaensis* have two anterior rows of foramina, of which

the anterior row is comparatively less developed (i.e., with smaller and fewer foramina)

than in aff. *Scirrotherium*. Additionally, in *S. hondaensis* and, particularly in *S.*

antelucanus, some osteoderms have a flattened or even a missing longitudinal central

elevation. These observations imply limitations on the taxonomic resolution, especially

considering that the material on which aff. *Scirrotherium* is based is scarce and does not

allow comparisons of a representative series of osteoderms encapsulating the

morphological variability within the carapace of this animal.

Kraglievichia Castellanos, 1927

LSDI. urn:lsid:zoobank.org:act:92C8B169-4F79-467E-B951-EF1DE6E327B1

Type species: *Kraglievichia paranensis* Ameghino, 1883

Other referred species: *Kraglievichia carinatum* comb. nov. (= *Scirrotherium carinatum*

Góis, Scillato-Yané, Carlini and Guilherme, 2013; see below)

Emended differential diagnosis: Small-to-middle sized pampatheriid characterized by

fixed osteoderms with ornamentation (particularly the longitudinal central elevation) more

conspicuous than in any other pampatheriid; anteriorly wide and posteriorly tapered

longitudinal central elevation; very deep longitudinal depressions; highly elevated and

frequently blunt marginal elevations, even flattened towards their top; external surface of

osteoderms generally rougher than in *Scirrotherium* but less than in *Holmesina*.

Kraglievichia carinatum comb. nov.

2013. *Scirrotherium carinatum* – Góis, Scillato-Yané, Carlini and Guilherme, Fig. 4.

Holotype: MLP 69-IX-8-13-AB, a mobile osteoderm (Góis et al. 2013).

Type locality and horizon: Paraná River cliffs, Province of Entre Ríos, Argentina.

Ituzaingó Formation, Upper Miocene, Tortonian (Góis et al. 2013).

Differential diagnosis: Unmodified (see Góis et al. 2013, p. 182).

Referred material: The holotype, paratypes and part of the hypodigm of this species (see

Fig. 10 and Appendix S1 of the Supplementary Material).

Discussion. In their descriptive work on *K. carinatum* comb. nov., Góis et al. (2013) did

not explicitly justify the inclusion of this species within *Scirrotherium*. Interestingly, part of

the material assigned to the taxon they create, coming from northwestern Brazil (Solimões

Formation), was previously referred to as *Kraglievichia* sp. by several researchers,

including Góis himself (Góis et al. 2004; Góis 2005; Cozzuol 2006; Latrubesse et al. 2010;

Góis et al. 2013). However, Góis et al. (2013) refuted the original taxonomic assignment,

arguing that it was erroneous, although they did not offer any concrete support for their

decision. In the absence of a phylogenetic analysis in Góis et al. (2013), we could assume

by default that these authors included *K. carinatum* comb. nov. within *Scirrotherium*

because they considered osteoderm features of this species at least compatible with the

generic diagnosis proposed by Edmund and Theodor (1997). Based on morphological

similarity, Góis and colleagues hypothesized closer affinities between *K. carinatum* comb.

nov. and *S. hondaensis* than those between *K. carinatum* comb. nov. and *K. paranensis*.

First, it is necessary to analyse in detail each of the osteoderm features of *K. carinatum*

comb. nov. in relation to the original diagnosis of *Scirrotherium*. According to Edmund and

Theodor (1997), the fixed osteoderms of *Scirrotherium* have a small (not specified) number

of large piliferous foramina on the anterior margin. These foramina are well spaced but

interconnected by a distinct channel. This is observed both in *K. carinatum* comb. nov. and

S. hondaensis. Likewise, the presence of continuous marginal elevations, posteriorly

confluent with the longitudinal central elevation, is a trait also shared by the two compared

species. Finally, the relative osteoderm size of *K. carinatum* comb. nov. is small relative to

other pampatheriids, which is in line with the original diagnosis for *Scirrotherium*.

Therefore, the osteoderm features of *K. carinatum* comb. nov. are compatible with those

mentioned in the diagnosis for *Scirrotherium* by Edmund and Theodor (1997). However,

this does not necessarily imply that the taxonomic allocation of *K. carinatum* comb. nov. to

the genus *Scirrotherium* is correct. In fact, there are several reasons to consider this

assignment is unreliable. Initially, the diagnosis of Edmund and Theodor (1997) contained

only three allegedly diagnostic osteoderm features, including the relative osteoderm size.

Furthermore, and more importantly, these "diagnostic features" do not allow definitive

discrimination between *Scirrotherium* and any other genus of pampatheriids. Indeed, in

their analysis, Góis et al. (2013) accept that, for instance, *Vassallia minuta* also shares the

presence of fixed osteoderms with a small number of large anterior foramina, which are

well spaced and connected by a canal. Independently, Laurito and Valerio (2013) also

highlighted the non-diagnostic nature for *Scirrotherium* of the latter osteoderm trait. The

other osteoderm features under consideration, i.e., the posterior confluence of the marginal

elevations with the longitudinal central elevation and the small osteoderm size, are also of

ambiguous diagnostic value. For instance, the confluence of marginal elevations and

longitudinal central elevation is also found in *K. paranensis*, a pampatheriid clearly

different from *Scirrotherium*. And although apparently informative on body-size trends of

some individual pampatheriid lineages (e.g., *Holmesina* spp.) and useful as discriminant

factor between species (Góis et al. 2013; Laurito and Valerio 2013), the relative osteoderm

size is not necessarily sufficient to make taxonomic assignments to the genus level among

pampatheriids as a whole (see below). In this sense, it is worth noting the potentially

conflicting taxonomic conclusions that could be reached using osteoderm-inferred relative

body size versus those based on non-osteoderm evidence. For example, MLP 69-IX-8-13A,

a femur belonging to an adult individual from the Ituzaingó Formation, was assigned to *K.*

cf. *paranensis* by Scillato-Yané et al. (2013), is comparable in size to that of the small

pampatheriid *S. hondaensis*. Therefore, it is probable that the referred femur does not

belong to the medium-to-large sized *K. paranensis*, although it is reasonable to include it in

the genus *Kraglievichia* (as the authors decided). However, Scillato-Yané et al. (2013) did

not discuss the possibility that the material assigned to *K.* cf. *paranensis*, particularly MLP

69-IX-8-13A, might be related to the (partially) co-occurrent species of *K. paranensis*, i.e.,

K. carinatum comb. nov., a pampatheriid whose small body size is fully compatible with

the small size of that femur. In other words, like Góis et al. (2013), they did not seriously

consider the hypothesis of *K. carinatum* comb. nov. as a small species of *Kraglievichia*,

rather than a species belonging to *Scirrotherium*.

Again analysing the original diagnosis of *Scirrotherium* by Edmund and Theodor (1997), it

should be regarded as ambiguous and hardly useful to differentiate this genus from other

genera in Pampatheriidae, at least with respect to osteoderm traits. It is likely that these

supposedly diagnostic features are actually symplesiomorphies for the entire family or, at

most, a hypothetical subfamilial lineage. This means that Góis et al. (2013) did not have a

sufficiently robust diagnosis of *Scirrotherium* to confidently assign *K. carinatum* comb.

nov. to this genus. Alternatively, they may have noted morphological similarity between

osteoderms of *K. carinatum* comb. nov. and *S. hondaensis* from features not included in the

diagnosis by Edmund and Theodor (1997). However, Góis et al. (2013) only listed

morphological differences between these species and virtually did not mention any

similarity for them, except for potentially equivocal resemblance as that indicated by

relative osteoderm size (i.e., small osteoderm sizes in comparison with those of *K.*

paranensis and *Plaina*). The lack of usefulness of the relative osteoderm size for generic

assignation is further supported by the osteoderm morphometric analysis in Góis et al.

(2013, p. 185), whose resulting PCA and CCA plots show that, despite the similarity in

relative osteoderm size, *K. carinatum* comb. nov. is located far from *S. hondaensis* (which

is closer to *V. minuta*, a taxon apparently related to other main lineage within

Pampatheriidae, i.e., *Plaina-Pampatherium*) and *K. paranensis* in morphospace.

Summarizing, there is little justification by Góis et al. (2013) on their taxonomic decision

to including *K. carinatum* comb. nov. within *Scirrotherium*. Considering the morphological

conservatism of Pampatheriidae, the observation of a common general morphological

pattern between *K. carinatum* comb. nov. and *S. hondaensis* does not necessarily imply the

grouping of these species under the same generic taxon, least of all by omitting the

taxonomic significance of striking similarities in osteoderm ornamentation between *K.*

carinatum comb. nov. and the better known taxon *K. paranensis*. Furthermore, we should

note that Góis and colleagues, in their work on *K. carinatum* comb. nov., did not make

morphological comparisons including to *S. antelucanus*, a species more similar in

osteoderm features to *S. hondaensis* (i.e., the type species of *Scirrotherium*). The species *S.*

antelucanus was described on a scientific article (Laurito and Valerio 2013) published

nearly simultaneously, but later, to that of *K. carinatum* comb. nov. Therefore, Góis et al.

(2013) did not know about the existence of *S. antelucanus* when they performed their

systematic analysis ("Until the present study, *S. hondaensis* was the only known species of

this genus"; Góis et al. 2013, p. 177), so that their taxonomic assignment of *K. carinatum*

comb. nov. to *Scirrotherium* was likely the result of limited notions on the morphological

variability and diversity of *Scirrotherium* in northern South America and southern Central

America.

In this work I decide to assign *K. carinatum* comb. nov. to the genus *Kraglievichia* based

on the results of phylogenetic analyses that I designed considering the hypothesis of

Edmund (1985) on the probable subfamilial relationships within Pampatheriidae, which

implicitly sustains that the creation of supraspecific taxa from osteoderm evidence should

be determined –with the prerequisite of morphological similarity- by the degree of

development of the ornamentation. Understanding that *K. carinatum* comb. nov. has

morphologically similar osteoderms to those of *K. paranensis* (apart from relative

osteoderm size) and has one of the more conspicuous, protuberant osteoderm

ornamentations among pampatheriids, along with *K. paranensis*, as acknowledged by Góis

et al. (2013) themselves, this means that *K. carinatum* comb. should be considered closely

related to *K. paranensis* and therefore they both should also be included in the same genus,

i.e., *Kraglievichia*.

5. Discussion

5.1. Systematic implications

This systematic analysis is the first attempt to test the intergeneric relationships and internal

structure of the genus *Scirrotherium* with its three previously referred species, i.e., *S.*

hondaensis (type species), '*S.*' *carinatum* (= *K. carinatum* comb. nov.) and *S. antelucanus*.

The two strict consensus trees from the distinct character weighting schemes (equal and

implied weights) show very similar results. However, the preferred phylogenetic hypothesis

is that supported by the implied weights analysis. According to Goloboff et al. (2018), the

parsimony analysis under implied weights outperforms equal weighting and the model-

based methods. Beyond this preference for a particular hypothesis (further supported

below), both resultant trees agree that the species referred to as *Scirrotherium* are not

monophyletic and, consistently, from a diagnostic point of view only *S. hondaensis* and *S.*

antelucanus appears as those actually referable to *Scirrotherium*. The relationship between

S. hondaensis and *S. antelucanus* could not be confidently resolved, despite the inclusion of

new osteoderm characters (the only ones comparable between these species so far) in these

parsimony analyses. Consequently, a paraphyletic relationship among *S. hondaensis* and *S.*

antelucanus should not be rule out. In conjunction, these results suggest the need of

information on craniomandibular or dental characters for *S. antelucanus* to further test the

affinities of this species with respect to *S. hondaensis*. Until new phylogenetic evidence is

available, *Scirrotherium* is maintained as taxonomically valid using a new, emended

diagnosis which focus (in addition to specific osteoderm similarities between *S. hondaensis*

and *S. antelucanus*) on its lesser degree of development of the osteoderm ornamentation in

comparison with those in *Holmesina* and, particularly, *Kraglievichia*, but greater degree of

development than in the *Plaina-Pampatherium* lineage. This new diagnosis replaces the

original and now inadequate diagnosis by Edmund and Theodor (1997).

Unlike the unpublished phylogeny of Góis (2013), the phylogenetic position of '*S.*'

carinatum was resolved here, i.e., this species is the sister taxon of *K. paranensis*.

Therefore, it is proposed the new name *K. carinatum* comb. nov. Despite Góis (2013) did

not recover as a clade to *S. hondaensis* and *K. carinatum* comb. nov., as expected if both

these species were assigned to *Scirrotherium*, he presented one supposed synapomorphy

that join them, i.e., very deep longitudinal depressions, "in particular in *S. carinatum*" (Góis

2013, p. 215). This feature is not a confident support for grouping *S. hondaensis* and *K.*

carinatum comb. nov. because the deepest longitudinal depressions in this pampatheriid

lineage are found in *K. carinatum* comb. nov. and *K. paranensis*, not in *S. hondaensis*. The

new emended differential diagnosis for *Kraglievichia* acknowledges the highly sculpted

external osteoderm surface documented on this taxon, which is more protuberant than in the

Plio-Pleistocene genus *Holmesina*. This diagnosis provides an updated and concise

description of useful osteoderm features to distinguish *Kraglievichia* from other genera

within Pampatheriidae. It is important to note that the species *K. paranensis* has several

autapomorphies (see Appendix S4 of the Supplementary Material) which, in addition to the

relative osteoderm size, need to be compared in the future with homologous, unknown

endoskeletal traits in *K. carinatum* comb. nov. in order to test the phylogenetic affinity of

these two species as inferred from osteoderm traits. Provisionally, the difference in relative

osteoderm size (consequently also in relative body size) and some morphological

differences between *K. carinatum* comb. nov. and *K. paranensis*, as noted by Góis et al.

(2013), may be linked to distinct ontogenetic growth trajectories in these species, with *K.*

carinatum comb. nov. representing the plesiomorphic condition (i.e., small body size; see

Sánchez-Villagra 2012 for a discussion on the implications for taxonomy of the ontogenetic

growth in extinct species).

Other important difference here with respect to the phylogeny of Góis (2013) is that *K.*

paranensis was not recovered as the one single sister taxon of *Holmesina* spp. (except *H.*

floridanus). Instead, it is part of a group additionally formed by *S. hondaensis*, *S.*

antelucanus and *K. carinatum* comb. nov. Together, these taxa are the sister clade of

Holmesina. This means that we do not have evidence of direct ancestral forms to

Holmesina yet. However, despite some striking differences, it is remarkable the greater

morphological similarity of the osteoderm ornamentation and several cranial features

between *Holmesina* (especially *H. floridanus*) and *Scirrotherium*, rather than with those of

Kraglievichia. The recognizable similarities between *H. floridanus* and *S. hondaensis*, and

at the same time differences with *K. paranensis*, include less protuberant osteoderm

ornamentation; occurrence of uniformly narrow longitudinal central elevation in fixed

osteoderms; more robust skull; more expanded frontals; the plane of the frontals forming a

reflex angle with that of the parietals; less anteroposteriorly elongated upper teeth; and the

first two upper molariforms less obliquely oriented with respect to the midline of the hard

palate.

Góis (2013) found that *Holmesina* is non-monophyletic due to the phylogenetic position of

H. floridanus with respect to the other *Holmesina* species. This result coincides with that

recovered here from the parsimony analysis with equal weights. *Holmesina floridanus* has

several plesiomorphic features in comparison with the remaining *Holmesina* species, e.g.,

less protuberant ornamentation, less rough external surface of the osteoderms and less

dorsally situated basicranium with respect the palatal plane. However, the confluent

arrangement of the calcaneal facets of the astragalus in *H. floridanus* suggests that this

species might not be directly related to any known South American pampatheriid (Edmund

1987). Likewise, Gaudin and Lyon (2017) have recently found potential support for the

monophyly of *Holmesina* from craniomandibular specimens. Therefore, the position of *H.*

floridanus in a polytomy with *S. hondaensis* and *S. antelucanus* in the analysis with equal

weights is explained from the lack of resolution (relatively small character sampling by a

fragmentary fossil record and homoplastic noise), not as product of a "real" non-

monophyletic relationship with the remaining *Holmesina* species. Conversely, the

Holmesina is recovered as monophyletic by the implied weights analysis. As expected, in

this topology *H. floridanus* is the most basal species of *Holmesina*. In both strict consensus

trees, *H. septentrionalis*, the other North American species, is grouped together with all the

South American *Holmesina* species (except *H. rondoniensis* and *H. cryptae*, both excluded

in this study).

I abstained from revising the diagnosis of *Holmesina* because this is considered beyond the

intended objectives of this work. However, six putative synapomorphies (unambiguous and

ambiguous) are proposed (or further supported) for this genus: (1) Anterior and lateral

margins with elongated, strong bone projections as radii directed from the external border

of the central figure towards the osteoderm borders in non-marginal fixed osteoderms; (2)

anteriorly convergent, nearly in contact medial processes of premaxillae; (3) length of

nasals greater than 30% of the maximum anteroposterior length of the skull; (4)

conspicuous and anteroposteriorly elongated maxillary ridge (Gaudin and Lyon 2017); (5)

relatively small lacrimal; and (6) bilobed posterior upper molariforms (Mf5-Mf9) with

incipient trilobulation.

5.2. Evolutionary and biogeographic implications

Scirrotherium is a pampatheriid genus from the Early Miocene-Late Pliocene of northern

South America and southern Central America. This taxon, along with *Kraglievichia*, forms

the sister group of *Holmesina* (Fig. 11), a pampatheriid probably originated in tropical

southern North America (Mexico? see Woodburne 2010). However, based on the

osteological comparisons presented above, which are expanded with respect to those of

Edmund (1987), the hypothetical South American ancestor or sister taxon of *Holmesina*

probably was morphologically generalized, more similar to *Scirrotherium* or aff.

Scirrotherium than to *Kraglievichia*. This interpretation is in line with that of Edmund

(1987), according to which the calcaneo-astragalar articulation of *S. hondaensis* challenges

the hypothesis that this pampatheriid is ancestral to *H. floridanus*, but the ornamentation

pattern of the osteoderms suggests "at least some degree of relationship" with the latter

species. Anatomically, *Kraglievichia* should be considered a highly divergent taxon,

especially to taking into account its Miocene age. According to Edmund (1987, p. 16), "the

osteoderms [of *H. floridanus*] are quite dissimilar to those of *Kraglievichia*". This

interpretation is shared in this work and contrasts with that of Scillato-Yané et al. (2005),

which suggested that *Holmesina* originated from a hypothetical South American basal form

of *Holmesina* or *Kraglievichia*. It also is opposed to Simpson (1930) and to the phylogeny

of Góis (2013) in which *Kraglievichia* is the only sister taxon of *Holmesina*.

The earliest record of *Scirrotherium*, here treated as tentative by scarce and poorly

preserved material, comes from the Early Miocene (late Burdigalian) of northwestern

Venezuela (Rincón et al. 2014; see below). Independently from the validity of occurrence

of *Scirrotherium* in an Early Miocene locality of northern South America, this record

represents the oldest pampatheriid reported in the scientific literature so far. This

improvement makes the fossil record of Pampatheriidae more congruent with the expected

time of origination of this family, i.e., Late Oligocene-Early Miocene, according to the very

few available time-calibrated phylogenies including representatives of Pampatheriidae and

its sister group, Glyptodontidae (e.g., Fernícola 2008; Billet et al. 2011). Apparently, the

Early Miocene Venezuelan pampatheriid indicates an origin of these xenartrans in low

latitudes in South America. However, this hypothesis could be defied by a possible Late

Eocene pampatheriid of Argentina, which has been not formally described and published

yet (Góis 2013).

Beyond the geographic origin of Pampatheriidae, northern South America seems to have

been a critical area for the early diversification of, at least, the lineage including

Scirrotherium. This genus probably differentiated at least as early as the late Early

Miocene-early Middle Miocene (late Burdigalian-Langhian) in northernmost South

America. The former evolutionary inference is consistent with the late Early Miocene

record referred to as *Scirrotherium* from Venezuela (Rincón et al. 2014).

Collectively, *Scirrotherium* and *Kraglievichia* occupied a large area in South America

during the Neogene (Fig. 12). The geographic range of *Scirrotherium* was more restricted

than that of *Kraglievichia*, comprising only tropical low latitudes, instead of a wide

latitudinal range, as suggested by Góis et al. (2013). The revaluated distributional pattern of

Scirrotherium is comparable to that of the glyptodontid *Boreostemma*, which is recorded

from the Middle Miocene to the Late Pliocene of Colombia and Venezuela (Carlini et al.

2008; Zurita et al. 2016). In contrast, the distributional range of *Kraglievichia* is similar to

that of other Miocene xenarthran taxa at the generic and specific level, which occurred in

southern South America and northwestern Brazil, but not in the northern or northwestern

end of South America (see Ribeiro et al. 2014).

Overall, this evidence indicating biogeographic divergence of northwesternmost South

America as an independent faunal province from the late Early Miocene to Middle

Miocene, and possibly into the Late Miocene or even the Early Pliocene, is consistent with

the results of multiple analyses of the South American terrestrial mammal fossil record for

the Neogene (Patterson and Pascual 1968; Cozzuol 2006; Ortiz-Jaureguizar and Cladera

2006; Croft 2007; Carrillo et al. 2015; Rincón et al. 2016; Kerber et al. 2017; Brandoni et

al. 2019). Apparently, the existence of one or several strong geographic and/or ecoclimatic

barriers (e.g., the Pebas Mega-Wetland System, whose expansion climax coincides with the

Middle Miocene) in northern South America would explain that regional endemism pattern

(MacFadden 2006; Croft 2007; Salas-Gismondi et al. 2015; Jaramillo et al. 2017). At the

same time, the development of a late Early-to-Middle Miocene biogeographic divergence

between northwesternmost South America and the rest of this continent may account for

the evolutionary divergence of *Scirrotherium* and *Kraglievichia*.

In the Late Miocene, without a completely formed Panamanian Land Bridge (O'dea et al.

2016), *Scirrotherium* expanded its geographic range to southern Central America (Fig. 12),

suggesting a possible ephemeral land connection or, more likely, overwater dispersal

between South America and Central America (maybe via rafting mechanism; efficient

active swimming of pampatheriids in a marine channel seems highly improbable). This is

the earliest dispersal event of a pampatheriid to North America (see below). The Central

American species of *Scirrotherium*, *S. antelucanus*, is larger than *S. hondaensis*, but

comparable or even smaller than aff. *Scirrotherium*. From available evidence, it is not

possible to determinate the most probable area of evolutionary differentiation of *S.*

antelucanus, but now there is support for occurrence of this species in the late Neogene of

northwestern South America, specifically in the Department of Sucre, Colombia.

The South American record of *S. antelucanus* is probably several million years younger (3–

5 my) than the Central American record. However, given the lack of absolute dating for the

fossil-bearing stratigraphic levels and the occurrence of Late Miocene strata in the same

geological unit (Sincelejo Formation) where comes the material here assigned to *S.*

antelucanus in Colombia, it should be recognized a significant age uncertainty for the new

South American record of this species. In any case, this age is considered may be Early

Pliocene or, alternatively, Latest Miocene from the stratigraphic position of the fossil-

bearing horizons (Villarroel and Clavijo, 2005; Bermúdez et al. 2009; Alfaro and Holz

2014; Bernal-Olaya et al. 2015; Córtes et al. 2018), as well as from associated

palynomorphs (Silva et al. 2012; B. Fernandes and C. Jaramillo, pers. comm. 2014).

The biogeographic correlation across the Isthmus of Panama using *S. antelucanus* has

insightful implications for the understanding of the late Cenozoic intercontinental migratory

dynamics in the Americas, including the Great American Biotic Interchange (GABI) (Webb

2006; Woodburne et al. 2006; Woodburne 2010; Cione et al. 2015; Bloch et al. 2016).

Noteworthy, this is the first transisthmian biogeographic correlation for a Neogene

terrestrial mammal at the level of species; furthermore, it is the first short-distance

intercontinental correlation (i.e., adjacent to the Central American Seaway) with high

taxonomic resolution for Neogene land mammals of the Americas; and, finally, it

constitutes the first evidence of a distributional pattern congruent with a re-entry event to

South America by a pre-Pleistocene xenarthran.

We know a few biogeographic correlations across the Isthmus of Panama which are based

on records at generic level of Neogene and Pleistocene land mammals, as well as a very

few records at species level of the latter epoch. The Neogene biogeographic correlations

include the pampatheriid genera *Plaina*, in Mexico and central-southwestern South

America; *Pampatherium*, in Mexico and southeastern South America; and *Holmesina*, in

the United States, Mexico and El Salvador, as well as in northwestern and southeastern

South America (Woodburne 2010). At the level of species, for instance, the Pleistocene

megatheriine *Eremotherium laurillardi* has occurrence in both sides of the Isthmus of

Panama in North- and South America (Cartelle and De Iuliis 1995, 2006; Tito 2008;

McDonald and Lundelius, E. L. Jr. 2009; Martinelli et al. 2012; Cartelle et al. 2015).

The record in South America of *S. antelucanus* increase the taxonomic resolution of

transisthmian biogeographic correlations of Neogene land mammals, opening the

possibility of new correlations of this kind and their biostratigraphic application in circum-

Caribbean basins, in a similar way as envisioned by the renowned American

palaeontologist Ruben A. Stirton from his revision of the fossil mammal remains of "La

Peñata fauna" (Stirton 1953), the vertebrate fossil association where comes the new record

of *S. antelucanus*. This translates into direct correlation of Land Mammal Ages (in this

case, SALMA and NALMA) from migrant mammals which are shared at species level by

both South and North America. Using to *S. antelucanus*, this would mean exists a support

for faunal, not necessarily chronological, correlation of the early Hemphillian and

Montehermosan mammal (xenarthran) assemblages in North America and South America,

respectively (see Laurito and Valerio, 2013). Naturally, any compelling intercontinental

faunal correlation requires more than one taxonomic element for support. The direct

intercontinental faunal correlations from Cenozoic land mammals between South and North

America are still underdeveloped in comparison with those between other continents (e.g.,

North America and Europe or North America and Asia; Woodburne and Swisher, C. C. III

1995; Beard and Dawson 1999; Bowen et al. 2002)

Additionally, the transisthmian correlation of *S. antelucanus* allows to increase the

geographic resolution in the detection of intercontinental migrations of late Cenozoic land

mammals, which are restricted mainly to large and middle distance correlations for the

Neogene record (e.g., Mexico-southern South America; Woodburne 2010). This pattern has

prevented the exploration of possible early or intermediate phases of

anagenetic/cladogenetic events in late Cenozoic Interamerican migrant taxa, which in turn

it is reflected in the fact that we are detecting "suddenly" well-differentiated terminal taxa

(e.g., *Holmesina*) in marginal, distant areas with respect to the Central American Seaway

and adjacent terrains (Cione et al. 2015 and references therein).

On the other hand, the new transisthmian correlation here presented suggests a possible

Neogene re-entry event by a xenartran to South America after its evolutionary

differentiation in North America (Fig. 12). The confirmation of this depends on a confident

determination of the differentiation area for *S. antelucanus*, i.e., if this species originated in

South America, the new record is explained more parsimoniously by population

maintenance in the ancestral area. Conversely, if this species originated in Central America

from a South American species of *Scirrotherium* as *S. hondaensis*, then we are considering

a re-entry event to South America. However, as mentioned above, it is not possible to

constrain much more than that at this moment. In any case, the possibility of a Neogene re-

entry event to South America by a xenarthran is compatible with the fact that we know

several of these events during the Pleistocene. Among these Pleistocene events, there are

several involved xenarthrans, including the pampatheriids *Holmesina* and *Pampatherium*,

the glyptodontid *Glyptotherium*, the pachyarmatheriid *Pachyarmatherium*, the dasypodid

Dasypus and the megatheriine *Eremotherium* (Woodburne et al. 2006; Woodburne 2010

and references therein).

On another note, the results of this work have evolutionary implications for the genus

Holmesina and the multiple Interamerican dispersal events of pampatheriids, including that

of *Scirrotherium* (discounting the non-confirmed re-entry event to South America). The

genus *Holmesina* has its oldest record (*Holmesina* sp.) in sedimentary rocks deposited

around the Pliocene-Pleistocene boundary (ca. 2.4 mya) in Florida, United States (Edmund

1987; Woodburne 2010 and references therein; Gaudin and Lyon 2017). This northward

dispersal event is part of the earliest phase of the GABI (GABI 1), in which additionally

participated other xenartrans as *Dasypus*, *Pachyarmatherium* and *Eremotherium*

(Woodburne 2010). Typically, *H. floridanus* has been considered the most basal among the

Holmesina species (Edmund 1987), as it is supported here. The hypothetical time-calibrated

phylogeny introduced in this work for Pampatheriidae (Fig. 11) suggests that exist a long

ghost lineage leading to *Holmesina*, from the Early Miocene (Burdigalian) until the Late

Pliocene. The improvement of the fossil record in northern South America, Central

America and Mexico will allow to advance in the recognition of probable direct ancestral

forms for *Holmesina*.

From the above analysis, a probable model of biogeographic evolution of *Holmesina* is as

follows (Fig. 12). A hypothetical pampatheriid close to *Holmesina* or even a hypothetical

Holmesina species basal with respect to *H. floridanus* dispersed to Central America,

Mexico and United States during the Pliocene (Early Pliocene according the time-calibrated

phylogeny). Once it was established the genus *Holmesina* in North America with *H.*

floridanus, the larger species *H. septentrionalis* diverged and differentiated in the Early

Pleistocene of southern United States. Later, *H. septentrionalis* expanded southward to

Mexico and Central America during the Early-Middle Pleistocene (Aguilar and Laurito

2009). In the Middle or early Late Pleistocene, possibly *H. septentrionalis* colonized South

America, where took place an important diversification, which was likely influenced by the

Late Pleistocene climatic changes (Scillato-Yané et al. 2005). This diversification gave

origin to the species *H. occidentalis*, *H. rondoniensis*, *H. cryptae*, *H. major* and the most

robust pampatheriid, *H. paulacoutoi* (Scillato-Yané et al. 2005; Moura et al. 2019).

As inferred from the phylogeny and derived interpretations here presented, the dispersal

events of *S. antelucanus* and *H. floridanus* to North America are independent of each other.

This means that the number of northward intercontinental dispersal events of pampatheriids

during the late Cenozoic actually is at least three, which in chronological order are: (1)

genus *Scirrotherium* (Late Miocene); (2) lineage *Plaina-Pampatherium* (Early Pliocene);

(3) genus *Holmesina* (undetermined Pliocene). From these events, only the latter, based on

the fossil record of *H. floridanus*, is included in the GABI. The remaining two events are

classified as part of the macroevolutionary invasion "wastebasket" called "Pre-GABI"

(literally, 'before the GABI'; Woodburne et al. 2006; Woodburne 2010; also named by

Cione et al. 2015 as "ProtoGABI"). In the lineage *Plaina-Pampatherium*, it was

differentiated one genus, *Pampatherium*, and at least three species (*P. mexicanum*, *P. typum*

and *P. humboldtii*, being the two latter recorded in South America). Meanwhile, the

northward dispersal event of *Scirrotherium* seems to give no origin to any other species

different to *S. antelucanus*. Only a confirmed southward intercontinental dispersal event of

the *Scirrotherium-Kraglievichia-Holmesina* clade has been well-established, i.e., that of

Holmesina to South America in the Middle or early Late Pleistocene (Aguilar and Laurito

2009). This event probably is not part of any of the GABI phases of Woodburne (2010) but

it appears to be chronologically located between the GABI 2 and 3.

As it has been shown, the study of more abundant and complete pampatheriid material

preserved in Neogene geological units of northern South America, in particular, and the

current Intertropical region of the Americas, in general, has the potential of provide us more

complex and interesting scenarios on the evolution of this glyptodontoid family and,

specifically, the genera *Scirrotherium* and *Holmesina*.

1001

6. Conclusion

The monophyly of *Scirrotherium* has been tested through parsimony phylogenetic analyses.

This taxon is recovered as paraphyletic. '*Scirrotherium*' *carinatum* forms a clade with

Kraglievichia paranensis and, therefore, here it is proposed the new name *K. carinatum*

comb. nov. The remaining referred species to *Scirrotherium*, *S. hondaensis* and *S.*

antelucanus, are designed in aphyly. The taxonomic validity of *Scirrotherium*, as defined

here, is maintained from diagnostic evidence. *Scirrotherium* is probably the sister taxon of

Kraglievichia, and these two genera form the sister clade of *Holmesina*. *Scirrotherium* has

occurrence from the late Early Miocene to Late Pliocene of northwestern South America

(Colombia and Venezuela) and the Late Miocene of southern Central America (Costa

Rica). A geographic origin of Pampatheriidae in northernmost South America is suggested

from the fossil record of *Scirrotherium* and a new time-calibrated phylogeny. *Scirrotherium*

also represents the earliest member of Pampatheriidae which participated in a dispersal

event to North America, specifically to the ancient Central American peninsula. This

dispersal event happened when the Panamanian Land Bridge was not fully formed yet. The

species *S. antelucanus* lived in Central America and northern Colombia during the late

Neogene. This is the first Interamerican biogeographic correlation of a Neogene land

mammal with high taxonomic resolution, i.e., at the species level. The record of *S.*

antelucanus in both sides of the ancient Central American Seaway is compatible with a

possible re-entry event of this pampatheriid to South America. In addition, *Scirrotherium* is

not probably the South American ancestor of the originally-endemic North American genus

Holmesina. In contrast with a previous hypothesis which argues that *Holmesina* may have

evolved from *Kraglievichia*, here it is suggested that there is no evidence of direct ancestral

forms of *Holmesina*, although the unknown South American ancestor of *Holmesina* may be

morphologically more similar to *Scirrotherium*.

1027

Acknowledgements

I am greatly indebted to all curators and managers of the collections visited for revision of

specimens during this work, especially to: Andrés Vanegas (Museo de Historia Natural La

Tatacoa, Colombia), Aldo Rincón (Museo Mapuka, Universidad del Norte, Colombia),

Jorge Moreno-Bernal (Museo Mapuka, Universidad del Norte), Carlos De Gracia (Museo

Mapuka, Universidad del Norte), Richard C. Hulbert Jr. (Florida Museum of Natural

History, USA), Kenneth Angielczyk (Field Museum, USA), Bill Simpson (Field Museum

of Natural History), Patricia Holroyd (University of California Museum of Paleontology,

USA) and Marcelo Reguero (Museo de La Plata, Argentina). I am also grateful to Carlos

Jaramillo (Smithsonian Tropical Research Institute, Panama) and his field work team from

allowing access to fossils they collected in the Department of La Guajira (Colombia).

Likewise, special thanks to Antonio Tovar (University of Sucre, Colombia) by inspiration

to explore the Sincelejo Formation; to my field assistants during this exploration in the

Department of Sucre (Colombia), Oscar Melendrez, Angel Cruz and Juan Pacheco. I would

like to thank César Laurito (Instituto Nacional de Aprendizaje de Costa Rica) and Edwin

Cadena (Universidad del Rosario, Colombia) for providing photos of osteoderms of the

species *S. antelucanus* and one photo of outcrops of the Castilletes Formation, respectively.

This work would not have been possible without logistic support and encourage by Alba

Lara, Carolay Jiménez and Mónica Montalvo. An anonymous reviewer provided comments

that made it possible to improve the manuscript. Financial assistance for this project was

provided by several institutions/agencies: CONICET (Internal Doctoral Fellowship); Field

Museum of Natural History (Science Visiting Scholarship); Florida Museum of Natural

History (International Travel Grant); University of California Museum of Paleontology

(Welles Fund); and The Paleontological Society (PalSIRP-Sepkoski Grant).

1052

Data archiving statement

Data for this study are available in the Dryad Digital Repository: [Intentionally blank]

The nomenclatural acts contained in this work are registered in ZooBank:

LSID. urn:lsid:zoobank.org:act:313358B5-3B1F-4902-8C2E-BB07CFCBEE18

LSID. urn:lsid:zoobank.org:act:E3B83181-91D6-44C8-90C0-BBAACEC2CDEE

LSID. urn:lsid:zoobank.org:act:225CD304-3B63-4B55-B8B8-33B46C90A194

LSID. urn:lsid:zoobank.org:act:92C8B169-4F79-467E-B951-EF1DE6E327B1

1060

References

Aguilar, D.H. Laurito, C.A., 2009. El armadillo gigante (Mammalia, Xenarthra,

Pampatheriidae) del río Tomayate, Blancano tardío-Irvingtoniano temprano, El

Salvador, América Central. Revista Geológica de América Central 41, 25–36.

doi:10.15517/rgac.v0i41.4176

Aguilar, T., Acevedo, B., Ulloa, A., 2010. Paleontología de una sección del río Corredores,

Formación Curré, Mioceno, Costa Rica. Revista Geológica de América Central 42,

43–75. doi:10.15517/rgac.v0i42.4170

Alfaro, E., Holz, M., 2014. Review of the chronostratigraphic charts in the Sinú-San Jacinto

Basin based on new seismic stratigraphic interpretations. Journal of South American

Earth Sciences 56, 139–169. doi:10.1016/j.jsames.2014.09.004

Alvarado, G.E., Barquero, R., Taylor, W., López, A., Cerdas, A., Murillo, J., 2009.

1073   Geología de la hoja general, Costa Rica. Revista Geológica de América Central 40,

1074   97–107. doi:0.15517/rgac.v0i40.4189

1075   Ameghino, F., 1888. Lista de las especies de mamíferos fósiles del Mioceno superior de

1076   Monte Hermoso hasta ahora conocidas. Obras Completas y Correspondencia

1077   Científica 5, 481–496.

1078   Anderson, V.J., Horton, B.K., Saylor, J.E., Mora, A., Tesón, E., Breecker, D.O., Ketcham,

1079   R.A., 2016. Andean topographic growth and basement uplift in southern Colombia:

1080   Implications for the evolution of the Magdalena, Orinoco, and Amazon river systems.

1081   Geosphere 12, 1235–1256. doi:10.1130/GES01294.1

1082   Antoine, P.O., Abello, M.A., Adnet, S., Sierra, A.J., Baby, P., Billet, G., Boivin, M.,

1083   Calderon, Y., Candela, M.A., Chabain, J., Corfu, F., Croft, D.D., Ganerød, M.,

1084   Jaramillo, C., Klaus, S., Marivaux, L., Navarrete, R.E., Orliac, M.J., Parra, F., 2016.

1085   A 60-million-year Cenozoic history of western Amazonian ecosystems in

1086   Contamana, eastern Peru. Gondwana Research 31, 30–59.

1087   doi:10.1016/j.gr.2015.11.001

1088   Ausich, W.I., Kammer, T.W., Rhenberg, E.C.,Wright, D.F., 2015. Early phylogeny of

1089   crinoids within the pelmatozoan clade. Palaeontology 58, 937–952.

1090   doi:10.1111/pala.12204

1091   Beard, K.C., Dawson, M.R., 1999. Intercontinental dispersal of Holarctic land mammals

1092   near the Paleocene/Eocene boundary: Paleogeographic, paleoclimatic and

biostratigraphic implications. Bulletin de la Société géologique de France 170, 697–

706.

Bengtson, P., 1988. Open nomenclature. Palaeontology 31, 223–227.

Bermúdez, H.D., Alvarán, M., Grajales, J.A., Restrepo, L.C., Rosero, J.S., Guzmán, C.,

Ruíz, E.C., Navarrete, R.E., Jaramillo, C., Osorno, J.F., 2009. Estratigrafía y

evolución geológica de la secuencia sedimentaria del Cinturón Plegado de San

Jacinto. Memorias XII Congreso Colombiano de Geología, 1–27.

doi:10.13140/2.1.5072.7367

Bernal-Olaya, R., Mann, P., Vargas, C.A., 2015. Earthquake, tomographic, seismic

reflection, and gravity evidence for a shallowly dipping subduction zone beneath the

Caribbean Margin of Northwestern Colombia, in: Bartolini, C., Mann, P. (Eds.),

Petroleum Geology and Potential of the Colombian Caribbean Margin. AAPG

Memoir 118, pp. 247–270. doi:10.1306/13531939M1083642

Billet, G., Hautier, L., De Muizon, C. and Valentin, X., 2011. Oldest cingulate skulls

provide congruence between morphological and molecular scenarios of armadillo

evolution. Proceedings of the Royal Society of London B, Biological Sciences 278,

2791–2797. doi:0.1098/rspb.2010.2443

Bloch, J.I., Woodruff, E.D., Wood, A.R., Rincon, A.F., Harrington, A.R., Morgan, G.S.,

Foster, D.A., Montes, C., Jaramillo, C.A., Jud, N.A., Jones, D.S., MacFadden, B.J.,

2016. First North American fossil monkey and early Miocene tropical biotic

interchange. Nature 533, 243–258. doi:10.1038/nature17415

Bowen, G.J., Clyde, W.C., Koch, P.L., Ting, S., Alroy, J., Tsubamoto, T., Wang, Y., Wang,

Y., 2002. Mammalian dispersal at the Paleocene/Eocene boundary. Science 295,

2062–2065. doi:10.1126/science.1068700

Brandoni, D., Ruiz, L.G.,Bucher, J., 2019. Evolutive Implications of *Megathericulus*

patagonicus (Xenarthra, Megatheriinae) from the Miocene of Patagonia Argentina.

Journal of Mammalian Evolution, 1–16. doi:10.1007/s10914-019-09469-6

Carlini, A.A., Zurita, A.E., Scillato-Yané, G.J., Sánchez, R., Aguilera, O.A., 2008. New

Glyptodont from the Codore Formation (Pliocene), Falcón State, Venezuela, its

relationship with the *Asterostemma* problem, and the paleobiogeography of the

Glyptodontinae. Paläontologische Zeitschrift 82, 139–152. doi:10.1007/BF02988405

Carrillo, J.D., Forasiepi, A., Jaramillo, C., Sánchez-Villagra, M.R., 2015. Neotropical

mammal diversity and the Great American Biotic Interchange: Spatial and temporal

variation in South America's fossil record. Frontiers in Genetics 5, 451, 1–11.

doi:10.3389/fgene.2014.00451

Carrillo-Briceño, J.D., Reyes-Cespedes, A.E., Salas-Gismondi, R., Sánchez, R., 2018. A

new vertebrate continental assemblage from the Tortonian of Venezuela. Swiss

Journal of Palaeontology, 1–12. doi:10.1007/s13358-018-0180-y

Cartelle, C., Bohórquez, G.A., 1985. *Pampatherium paulacoutoi*, uma nova

espécie de tatu gigante da Bahia, Brasil (Edentata, Dasypodidae). Revista Brasileira de

Zoologia 2, 229–254. doi:10.1590/S0101-81751983000400003

Cartelle, C., De Iuliis, G., 1995. *Eremotherium laurillardi*: The Panamerican late

Pleistocene megatheriid sloth. Journal of Vertebrate Paleontology 15, 830–841.

doi:10.1080/02724634.1995.10011265

Cartelle, C., De Iuliis, G., 2006. *Eremotherium laurillardi* (Lund) (Xenarthra,

Megatheriidae), the Panamerican giant ground sloth: Taxonomic aspects of the

ontogeny of skull and dentition. Journal of Systematic Palaeontology 4, 199–209.

doi:10.1017/S1477201905001781

Cartelle, C., De Iuliis, G., Pujos, F., 2015. *Eremotherium laurillardi* (Lund, 1842)

(Xenarthra, Megatheriinae) is the only valid megatheriine sloth species in the

Pleistocene of intertropical Brazil: A response to Faure et al., 2014. Comptes Rendus

Palevol 14, 15–23. doi:10.1016/j.crpv.2014.09.002

Castellanos, A., 1937. Anotaciones sobre la línea filogenética de clamiterios. Serie

Técnico-Científica de la Facultad de Ciencias Matemáticas, Físico-Químicas y

Naturales 8, 1–35. Rosario, Argentina.

Cione, A., Gasparini, G., Soibelzon, E., Soibelzon, L., Tonni, E., 2015. The Great

American Biotic Interchange. A South American perspective. Springer Briefs in

Earth System Sciences. South America and the Southern Hemisphere, Amsterdam,

Netherlands. doi:10.1007/978-94-017-9792-4

Coates, A.G., Stallard, R.F., 2013. How old is the Isthmus of Panama? Bulletin of Marine

Science 89, 801–813. doi:10.5343/bms.2012.1076

Cortes, J.E., Aguilera, R., Wilches, O., Osorno, J.F., Cortes, S.I., 2018. Organic

geochemical insights from oil seeps, tars, rocks, and mud volcanoes on the petroleum

1156  systems of the Sinú-San Jacinto basin, Northwestern Colombia. Journal of South

1157  American Earth Sciences 86, 318–341. doi: 10.1016/j.jsames.2018.06.007

1158 Cozzuol, M.A., 2006. The Acre vertebrate fauna: age, diversity, and geography. Journal of

1159  South American Earth Sciences 21, 185–203. doi:10.1016/j.jsames.2006.03.005

1160 Croft, D.A. 2007., The Middle Miocene (Laventan) Quebrada Honda fauna, southern

1161  Bolivia and a description of its notoungulates. Palaeontology 50, 277–303.

1162  doi:10.1111/j.1475-4983.2006.00610.x

1163 De Iuliis, G., Edmund, A.G., 2002. *Vassallia maxima* Castellanos, 1946 (Mammalia:

1164  Xenarthra: Pampatheriidae), from Puerta del Corral Quemado (late Miocene to early

1165  Pliocene), Catamarca Province, Argentina. Smithsonian Contributions to

1166  Paleobiology 93, 49–64.

1167 Delsuc, F., Superina, M., Tilak, M.K., Douzery, E.J., Hassanin, A., 2012. Molecular

1168  phylogenetics unveils the ancient evolutionary origins of the enigmatic fairy

1169  armadillos. Molecular Phylogenetics and Evolution 62, 673–680.

1170  doi:10.1016/j.ympev.2011.11.008

1171 Ebach, M.C., Williams, D.M., 2010. Aphyly: a systematic designation for a taxonomic

1172  problem. Evolutionary biology, 37 (2-3), 123–127. doi:10.1007/s11692-010-9084-5

1173 Edmund, A.G., 1985. The armor of fossil giant Armadillos (Pampatheriidae, Xenarthra,

1174  Mammalia). Texas Memorial Museum, Pearce-Sellards-Series 40, 1–20.

1175 Edmund, A.G., 1987. Evolution of the Genus *Holmesina* (Pampatheriidae, Mammalia) in

1176  Florida, with Remarks on Taxonomy and Distribution. Texas Memorial Museum,

1177  Pearce-Sellards-Series 45, 1–20.

Edmund, A.G., Theodor, J., 1997. A new giant Armadillo, in: Kay, R.F. Cifelli, R.L., Flynn

J.J., Madden, R. (Eds.), Vertebrate Paleontology of the Miocene Fauna of La Venta,

Colombia. Smithsonian Institution Press, Washington, pp. 227–232.

Fernícola, J.C., 2008. Nuevos aportes para la sistemática de los Glyptodontia Ameghino

1889 (Mammalia, Xenarthra, Cingulata). Ameghiniana, 45, 553–574.

Flinch, J.F., 2003. Structural evolution of the Sinú-Lower Magdalena area (northern

Colombia), in: Bartolini, C., Buffler, R.T., Blickwede, J. (Eds.), The Circum-Gulf of

Mexico and the Caribbean: Hydrocarbon Habitats, Basin Formation, and Plate

Tectonics, AAPG Memoir 79, pp. 776–796.

Flynn J.J., Guerrero, J., Swisher, C.C., III. 1997. Geochronology of the Honda Group, in:

Kay, R.F. Cifelli, R.L., Flynn J.J., Madden, R. (Eds.), Vertebrate Paleontology of the

Miocene Fauna of La Venta, Colombia. Smithsonian Institution Press, Washington,

pp. 44–59.

Gaudin, T.J., 2004. Phylogenetic relationships among sloths (Mammalia, Xenarthra,

Tardigrada): The craniodental evidence. Zoological Journal of the Linnean Society

140, 255–305. doi:10.1111/j.1096-3642.2003.00100.x

Gaudin, T.J., Wible, J.R., 2006. The phylogeny of living and extinct armadillos

(Mammalia, Xenarthra, Cingulata): A craniodental analysis, in: Carrano, M.T.,

Gaudin, T. J., Blob, R.W., Wible, J.R. (Eds.), Amniote Paleobiology: Perspectives

on the Evolution of Mammals, Birds and Reptiles. University of Chicago Press,

Chicago, pp. 153–198.

Gaudin, T.J., Lyon, L.M., 2017. Cranial osteology of the pampathere *Holmesina floridanus*

(Xenarthra: Cingulata; Blancan NALMA), including a description of an isolated

petrosal bone. PeerJ 5, e4022. doi:10.7717/peerj.4022

Góis, F. 2005. Estudo descritivo e geométrico dos Cingulata (Mammalia, Xenarthra) do

Neógeno e Quaternário da Amazônia Sul-Ocidental. Bachelor thesis, Universidade

Federal de Rondônia (unpubl.).

Góis, F. 2013. Análisis morfológico y afinidades de los Pampatheriidae (Mammalia,

Xenarthra). PhD thesis, Universidad Nacional de La Plata (unpubl.).

Góis, F., Scillato-Yané, G.J., Carlini, A.A., Ubilla, M., 2012. Una nueva especie de

Holmesina Simpson (Xenarthra, Cingulata, Pampatheriidae) del Pleistoceno de

Rondônia, sudoeste de la Amazonia, Brasil. Revista Brasileira de Paleontologia 15,

211–227.doi: 10.4072/rbp.2012.2.09

Góis, F., Scillato-Yané, G.J., Carlini, A.A., Guilherme, E., 2013. A new species of

Scirrotherium Edmund and Theodor, 1997 (Xenarthra, Cingulata, Pampatheriidae)

from the late Miocene of South America. Alcheringa: An Australasian Journal of

Palaeontology 37, 177–188. doi: 10.1080/03115518.2013.733510

Góis, F., Ruiz, L.R., Scillato-Yané, G.J., Soibelzon, E., 2015. A peculiar new

Pampatheriidae (Mammalia: Xenarthra: Cingulata) from the Pleistocene of Argentina

and comments on Pampatheriidae diversity. PloS One 10, e0128296.

doi:10.1371/journal.pone.0128296

Góis, F., Nascimento, E.R., Porto, A.S., Holanda, E.C., Cozzuol, M.A., 2004. Ocorrências

de Cingulata dos gêneros *Kraglievichia* e *Holmesina* do Terciário e Quaternário da

Amazônia Sul-Ocidental. Ameghiniana 49, 41.

Guerrero, J. 1997. Stratigraphy and sedimentary environments of the Honda Group in the

La Venta area. Miocene uplift of the Colombian Andes, in Kay, R.F., Cifelli, R.L.,

Flynn, J.J., Madden, R. (Eds.), Vertebrate Paleontology of the Miocene Fauna of La

Venta, Colombia. Smithsonian Institution Press, Washington, pp. 15–43.

Goloboff, P.A., 2014. Extended implied weighting. Cladistics 30, 260–272.

doi:10.1111/cla.12047

Goloboff, P.A., Torres, A., Arias, J.S., 2018. Weighted parsimony outperforms other

methods of phylogenetic inference under models appropriate for morphology.

Cladistics 34, 407–437. doi:10.1111/cla.12205

Goloboff, P.A., Carpenter, J.M., Salvador Arias, J., Miranda Esquivel, D.R., 2008.

Weighting against homoplasy improves phylogenetic analysis of morphological data

sets. Cladistics 24, 758–773. doi:10.1111/j.1096-0031.2008.00209.x

Hoffstetter, R., 1952. Les mammifères pleistocenes de la République de l'Équateur.

Mémoires Societé Géologique de France 66, 1–391.

Jaramillo, C., 2018. Evolution of the Isthmus of Panama: Biological, paleoceanographic,

and paleoclimatological implications, in: Hoorn, C., Antonelli, A. (Eds.), Mountains,

climate and biodiversity. John Wiley and Sons, Oxford, pp. 323–338.

Jaramillo, C., Romero, I., D'Apolito, C., Bayona, G., Duarte, E., Louwye, S., Escobar, J.,

Luque, J., Carrillo-Briceño, J.D., Zapata, V., Mora, A., Schouten, S., Zavada, M.,

Harrington, G., Wesseling, F.P., 2017. Miocene flooding events of western

Amazonia. Science advances 3, e1601693. doi:10.1126/sciadv.1601693

Kerber, L., Negri, F.R., Ribeiro, A.M., Nasif, N., Souza-Filho, J.P., Ferigolo, J., 2017.

Tropical fossil caviomorph rodents from the southwestern Brazilian Amazonia in the

context of the South American faunas: Systematics, biochronology, and

paleobiogeography. Journal of Mammalian Evolution 24, 57–70.

doi:10.1007/s10914-016-9340-2

Kolarsky, R.A., Mann, P., Montero, W., 1995. Island arc response to shallow subduction of

the Cocos Ridge, Costa Rica. Special papers of the Geological Society of America

295, 235–262. doi:10.1130/SPE295-p235

Latrubesse, E.M., Cozzuol, M.A., Silva-Caminha, S.A., Rigsby, C.A., Absy, M.L.,

Jaramillo, C., 2010. The late Miocene paleogeography of the Amazon Basin and the

evolution of the Amazon River system. Earth-Science Reviews 99, 99–124.

doi:10.1016/j.earscirev.2010.02.005

Laurito, C.A., Valerio, A.L., 2013. *Scirrotherium antelucanus*, una nueva especie de

Pampatheriidae (Mammalia, Xenarthra, Cingulata) del Mioceno Superior de Costa

Rica, América Central. Revista Geológica de América Central 49, 45–62.

doi:10.15517/rgac.v0i49.13101

Leidy, J., 1889. Fossil Vertebrates from Florida. Proceedings of the Academy of Natural

Sciences of Philadelphia 41, 96-97.

Lowery, B.J., 1982. Sedimentology and tectonic implications of the Middle to Upper

Miocene Curré Formation. M.S. thesis, Louisiana State University (unpub.).

Lund, P., 1839. Blik paa Braziliens Dyreverden for Sidste Jordomvaeltning. Anden

Afhandling: Pattedyrene. Det Kongelige Danske Videnskabernes

SelskbasNaturvidenskabelige og Mathematiske Afhandlinger 8, 61–44.

Lund, P., 1842. Blik paa Braziliens Dyreverden for Sidste Jordomvaeltning.

TredieAfhandling: Forsaettelse af Pattedyrene. Det Kongelige Danske

Videnskabernes SelskbasNaturvidenskabelige og Mathematiske Afhandlinger 8, 217–

272.

MacFadden, B.J., 2006. Extinct mammalian biodiversity of the ancient New World tropics.

Trends in Ecology and Evolution 21, 157–165. doi:10.1016/j.tree.2005.12.003

Maddison, W.P., Maddison, D.R., 2010. Mesquite: a modular system for evolutionary

analysis. Version 2.73. [http://mesquiteproject.org]

Martinelli, A.G., Ferraz, P.F., Cunha, G.C., Cunha, I.C., De Souza Carvalho, I., Ribeiro, L.

C.B., Macedo Neto, F., Lourencini Cavellani, C., Antunes Teixeira, V.P., Da Fonseca

Ferraz, M.L., 2012. First record of *Eremotherium laurillardi* (Lund, 1842)

(Mammalia, Xenarthra, Megatheriidae) in the Quaternary of Uberaba, Triângulo

Mineiro (Minas Gerais State), Brazil. Journal of South American Earth Sciences, 37,

202–207. doi: 10.1016/j.jsames.2012.03.006

McDonald, H.G., Lundelius, E.L., Jr., 2009. The giant ground sloth *Eremotherium*

laurillardi (Xenarthra, Megatheriidae), in: Albright, L.B. III (Ed.), Papers on

geology, vertebrate paleontology, and biostratigraphy in honor of Michael O.

Woodburne, Bulletin 65, Texas, pp. 407–421.

McKenna, M.C., Bell, S.K., 1997. Classification of Mammals Above the Species Level.

Columbia University Press, New York.

Moreno, E. P., Mercerat, A., 1891. Exploración arqueológica de la provincia de Catamarca:

Paleontología. Revista del Museo de La Plata, 1: 222–236.

Moreno, F., Hendy, A. J.W., Quiroz, L., Hoyos, N., Jones, D.S., Zapata, V., Zapata, S.,

Ballen, G.A., Cadena, E., Cárdenas, A.L., Carrillo-Briceño, J.D., Carrillo, J.D.,

Delgado-Sierra, D., Escobar, J., Martínez, J.I., Martínez, C., Montes, C., Moreno, J.,

Pérez, N., Sánchez, R., Suárez, C., Vallejo-Pareja, M.C., Jaramillo, C., 2015. Revised

stratigraphy of Neogene strata in the Cocinetas basin, La Guajira, Colombia. Swiss

Journal of Palaeontology 134, 5–43. doi: 10.1007/s13358-015-0071-4

Obando, L.G., 2011. Stratigraphy and tectonic of northeast part of Dota quadrangle (1:

50,000), Costa Rica. Revista Geológica de América Central 44, 71–82.

O'dea, A., Lessios, H.A., Coates, A.G., Eytan, R.I., Restrepo-Moreno, S.A., Cione, A.L.,

Collins, L.S., De Queiroz, A., Farris, D.W., Norris, R.D., Stallard, R.F., Woodburne,

M.O., Aguilera, O., Aubry, M-P., Berggren, W.A., Budd, A.F., Cozzuol, M.A.,

Coppard, S.E., Duque-Caro, H., Finnegan, S., Gasparini, G.M., Grossman, E.L.,

Jhonson K.G., Lloyd, D.K, Knowlton N., Leigh E.G., Leonard-Pingel J.S., Marko,

P.B., Pyenson, N.D., Rachello-Dolmen, P.G., Soibelzon, E., Soibelzon, L., Todd,

J.A., Vermeij, G.J., Jackson, J.B.C., 2016. Formation of the Isthmus of Panama.

Science advances 2, e1600883. doi: 10.1126/sciadv.1600883

Ortiz-Jaureguizar, E., Cladera, G.A., 2006. Paleoenvironmental evolution of southern South

America during the Cenozoic. Journal of Arid Environments 66, 498–532. doi:

10.1016/j.jaridenv.2006.01.007

Patterson, B., Pascual, R., 1968. The fossil mammal fauna of South America. The Quarterly

Review of Biology 43, 409–451.

Ribeiro, A.M., Madden, R.H., Negri, F.R., Kerber, L., Hsiou, A.S., Rodrigues, K.A., 2013.

Mamíferos fósiles y biocronología en el suroeste de la Amazonia, Brasil, in:

Brandoni, D., Noriega, J.I. (Eds.), El Neógeno de la Mesopotamia argentina.

Asociación Paleontológica Argentina, Publicación Especial 14, Buenos Aires, pp.

207–221.

Rincón, A.D., Solórzano, A., Benammi, M., Vignaud, P., McDonald, H.G., 2014.

Chronology and geology of an Early Miocene mammalian assemblage in North of

South America, from Cerro La Cruz (Castillo Formation), Lara State, Venezuela:

Implications in the changing course of Orinoco River hypothesis. Andean geology

41, 507–528. doi: 10.5027/andgeoV41n3-a02

Rincón, A.D., Solórzano, A., Macsotay, O., McDonald, H.G., Núñez-Flores, M., 2016. A

new Miocene vertebrate assemblage from the Río Yuca Formation (Venezuela) and

the northernmost record of typical Miocene mammals of high latitude (Patagonian)

affinities in South America. Geobios 49, 395–405. doi:10.1016/j.geobios.2016.06.005

Rivier, F., 1985. Sección geológica del Pacífico al Atlántico a través de Costa Rica. Revista

Geológica de América Central 2, 22–32. doi:10.15517/rgac.v0i02.10480

Robertson, J.S., 1976. Latest Pliocene mammals from Haile XV A, Alachua county,

Florida. Bulletin of the Florida State Museum, Biological Sciences 20, 111–186.

Salas-Gismondi, R., Flynn, J.J., Baby, P., Tejada-Lara, J.V., Wesselingh, F.P., Antoine, P.

O., 2015. A Miocene hyperdiverse crocodylian community reveals peculiar trophic

dynamics in proto-Amazonian mega-wetlands. Proceedings of the Royal Society B:

Biological Sciences 282, 20142490. doi: 10.1098/rspb.2014.2490

Sánchez-Villagra, M.R., 2012. Embryos in Deep Time: The Rock Record of Biological

Development. University of California Press, Berkeley.

Schmidt, D.N., 2007. The closure history of the Panama Isthmus: Evidence from isotopes

and fossils to models and molecules, in: Williams, M., Haywood, A.M., Gregory J.F.,

Schmidt, D.N. (Eds.), Deep time perspectives on climate change – marrying the

signal from computer models and biological proxies. Geological Society of London,

London, pp. 427–442.

Scillato-Yané, G.J., Carlini, A.A., Tonni, E.P., Noriega, J.I., 2005. Paleobiogeography of

the late Pleistocene pampatheres of South America. Journal of South American Earth

Sciences 20, 131–138. doi:10.1016/j.jsames.2005.06.012

Scillato-Yané, G.J., Góis, F., Zurita, A.E., Carlini A.A., González Ruiz, L.R., Krmpotic,

C.M., Oliva, C., Zamorano, M. 2013. Los Cingulata (Mammalia, Xenarthra) del

'Conglomerado Osífero' (Mioceno tardío) de la Formación Ituzaingó de Entre Ríos,

Argentina, in: Brandoni, D., Noriega, J.I. (Eds.), El Neógeno de la Mesopotamia

argentina. Asociación Paleontológica Argentina, Publicación Especial 14, Buenos

Aires, pp. 118–134.

Sigovini, M., Keppel, E., Tagliapietra, D., 2016. Open Nomenclature in the biodiversity

1348   era. Methods in Ecology and Evolution 7, 1217–1225. doi:10.1111/2041-210X.12594

1349   Simpson, G.G., 1930. *Holmesina septentrionalis*, extinct giant armadillo of Florida.

1350   American Museum Novitates 442, 1–10.

1351   Silva, J.C., Pardo, A., Cardona, A., Borrero, C., Flores, A., Navarette, R., Mejía, A., Ochoa,

1352   D., Osorio, J.A., Rosero, S., Arenas, A., 2012. Multi-Chronological Proxies to

1353   Timing Early Oligocene and Middle Miocene Deformation Events Along the Lower

1354   Magdalena Basin, NW Colombia. Resúmenes del XI Simposio Bolivariano-

1355   Exploración Petrolera en las Cuencas Subandinas, Asociación Colombiana de

1356   Geólogos y Geofísicos del Petróleo. Available in:

1357   [http://www.earthdoc.org/publication/publicationdetails/?publication=66175]

1358   Stirton, R.A., 1953. Vertebrate paleontology and continental stratigraphy in Colombia.

1359   Geological Society of America Bulletin 64, 603–622.

1360   Swofford, D.L., 2015. PAUP*: Phylogenetic analysis using parsimony (and other methods)

1361   (version 4.0a142).

1362   Swofford, D.L., Bell, C.D., 2017. PAUP* manual. Available at

1363   http://phylosolutions.com/paup-documentation/paupmanual.pdf.

1364   Tito, G., 2008. New remains of *Eremotherium laurillardi* (Lund, 1842) (Megatheriidae,

1365   Xenarthra) from the coastal region of Ecuador. Journal of South American Earth

1366   Sciences 26, 424–434. doi:10.1016/j.jsames.2008.05.001

Villarroel, C., Clavijo, J., 2005. Los mamíferos fósiles y las edades de las sedimentitas

continentales del Neógeno de la Costa Caribe Colombiana. Revista de la Academia

Colombiana de Ciencias 29, 345–356.

Vizcaíno, S.F., De Iuliis, G., Bargo, M.S., 1998. Skull shape, masticatory apparatus, and

diet of *Vassallia* and *Holmesina* (Mammalia: Xenarthra: Pampatheriidae): When

anatomy constrains destiny. Journal of Mammalian Evolution 5, 291–322. doi:

10.1023/A:1020500127041

Webb, S.D., 2006. The Great American Biotic Interchange: Patterns and Processes. Annals

of the Missouri Botanical Garden 93, 245–258. doi: 10.3417/0026-

6493(2006)93[245:TGABIP]2.0.CO;2

Woodburne, M.O., 2010. The Great American Biotic Interchange: dispersals, tectonics,

climate, sea level and holding pens. Journal of Mammalian Evolution 17, 245–264.

doi:10.1007/s10914-010-9144-8

Woodburne, M.O., Swisher, C.C. III., 1995. Land mammal high-resolution geochronology,

intercontinental overland dispersals, sea level, climate and vicariance, in: Berggren,

W.A., Kent, D.W., Aubry, M.-P., Hardenbol, J. (Eds.), Geochronology, Time Scales

and Global Stratigraphic Correlation. SEPM Special Publication 54, pp. 335–364.

Woodburne, M.O., Cione, A.L., Tonni, E.P., 2006. Central American provincialism and the

Great American Biotic Interchange. Universidad Nacional Autónoma de México,

Instituto de Geología y Centro de Geociencias, Publicación Especial 4, 73–101.

Yuan, P.B. 1984. Stratigraphy, sedimentology and geologic evolution of eastern Terraba

Trough, southwestern Costa Rica. PhD thesis, Louisiana State University (unpub.).

Zurita, A.E., Scillato-Yané, G.J., Ciancio, M., Zamorano, M., González-Ruiz, L.R., 2016.

Los Glyptodontidae (Mammalia, Xenarthra): Historia biogeográfica y evolutiva de un

grupo particular de mamíferos acorazados. Contribuciones del MACN 6, 249–262.

Figure captions

Figure 1. Geographic and stratigraphic provenance of the newly described material of pampatheriids from the Neogene of Colombia. In the left upper corner, a map of northwesternmost South America and the location of the regions of Colombia where there are outcrops of the formations with pampatheriid specimens for this study. In the right upper corner, photos of characteristic outcrops of these formations. Below in the center, a general chronostratigraphic scheme with the position of each formation within the Neogene and two important tectonic/palaeogeographic events in northwestern South America, i.e., a major, underwater uplift of the Isthmus of Panama (Schmidt 2007) and the definitive emergence of the Panamanian Land Bridge (O'dea et al. 2016). The photo of an outcrop of the Castilletes Formation was taken by Edwin Cadena.

Figure 2. Phylogenetic results. A, strict consensus tree of the parsimony with equal weights. B, strict consensus tree of the parsimony analysis with implied weights. The numbers below nodes are bootstrap resampling frequencies. Note the difference in the phylogenetic position of *Holmesina floridanus* in the two strict consensus trees. Explanation of this difference in the *Discussion* section.

Figure 3. Fixed and (semi) mobile osteoderms of *Scirrotherium hondaensis* from the La Victoria and Villavieja Formations, Municipality of Villavieja, Department of Huila, Colombia. A–B', fixed osteoderms; C'–K', (semi) mobile osteoderms. The osteoderms G, J, K, L, W, X, Y, Z, A', B', G', I', J' and K' are associated with the catalog number VPPLT 348. The osteoderms H, U and V are associated with the catalog number VPPLT 004. The osteoderms T and D' are associated with the catalog number VPPLT 701. All the former osteoderms come from the lower and middle La Victoria Formation. The osteoderms B, C,

F, I, O, P, S, C' and F' are associated with the catalog number VPPLT 1683 - MT 18 and come from the top of the La Victoria Formation. The osteoderms A, D, E, M, N, Q, R, E' and H' are associated with the catalog number VPPLT 1683 - MT 18 and come from the lower Villavieja Formation. Scale bar equal to 20 mm.

Figure 4. Photos and anatomical line drawings of the skull VPPLT 706 of *Scirrotherium hondaensis* from the middle La Victoria Formation, Municipality of Villavieja, Department of Huila, Colombia. A–B, dorsal views; C–D, ventral views; E–F, right lateral views; D–H, left lateral views. *Abbreviations*: aof, antorbital fossa; fr, frontal; iof, infraorbital foramen; la, lacrimal; Mf1, first upper molariform; Mf9, ninth upper molariform; mx, maxilla; mxf, maxillary foramen; na, nasal; pal, palatine; pm, premaxilla. Scale bar equal to 50 mm.

Figure 5. Photos and anatomical line drawings of the left femur and right ulna VPPLT 706 of *Scirrotherium hondaensis* from the middle La Victoria Formation, Municipality of Villavieja, Department of Huila, Colombia. The epiphyses of this femoral diaphysis have been reconstructed from those with catalog number UCMP 39846. A–B, anterior views of the femur; C–D, posterior views of the femur. E–F, medial views of the ulna; G–H, lateral views of the ulna. *Abbreviations*: anc, fossa for the anconeus muscle; cp, coronoid process; fh, femoral head; gt, greater trochanter; le, lateral epicondyle; me, medial epicondyle; op, olecranon process; tn, trochlear notch; tt, third trochanter. Scale bar equal to 50 mm.

Figure 6. Photos and anatomical line drawings of a thoracic vertebra (A–B) and several anterior caudal vertebrae (C–F) VPPLT 706 of *Scirrotherium hondaensis* from the middle La Victoria Formation, Municipality of Villavieja, Department of Huila, Colombia. A–B, posterior views of the thoracic vertebra. C–D, lateral views of caudal vertebrae; E–F, dorsal views of caudal vertebrae. *Abbreviations*: az, anterior zygapophyses; mp, metapophyses;

ns, neural spine; tp, transverse processes; vb, vertebral body; vla, ventrolateral apophyses. Scale bar equal to 30 mm.

Figure 7. Photos and anatomical line drawings of the astragalus (A–D) and calcaneum (E – F) UCMP 39846 of *Scirrotherium hondaensis* from the lower (?) Villavieja Formation, Municipality of Villavieja, Department of Huila, Colombia. A–B, astragalus in plantar views; C–D, astragalus in dorsal views. E–F, calcaneum in dorsal views. *Abbreviations*: ct, calcaneal tuber; ef, ectal facet; h, head of the astragalus; lt, lateral trochlea; mt, medial trochlea; sf, sustentacular facet; st, sulcus tali. Scale bar equal to 20 mm.

Figure 8. Fixed osteoderm MUN STRI 36880 of *Scirrotherium antelucanus* from the upper Sincelejo Formation, Department of Sucre, Colombia. Scale bar equal to 20 mm.

Figure 9. Pampatheriid osteoderms from the Department of La Guajira, Colombia, referred to as aff. *Scirrotherium* (**A**, MUN STRI 16718; **E**, MUN STRI 38064; and **G,** MUN STRI 16719; all these specimens are from the Castilletes Formation and they are fixed osteoderms except the latter, which consist of an anterior fragment of a mobile osteoderm); *Scirrotherium* cf. *hondaensis* (**C**, MUN STRI 36814, a fixed osteoderm from the Castilletes Formation); and *Scirrotherium* sp. (**B**, MUN STRI 36801; **D**, MUN STRI 16158; and **F**, MUN STRI 34373; all these fixed osteoderms are from the Castilletes Formation, except the latter, which comes from the Ware Formation). Note the two well-developed rows of anterior foramina in the osteoderms MUN STRI 16718 and 38064. Scale bar equal to 20 mm.

Figure 10. Osteoderms of *Kraglievichia carinatum* comb. nov. from the Ituzaingó Formation, Province of Entre Ríos, Argentina. The holotype of this species is marked with

one single asterisk (*) and paratypes with double asterisk (**). A–J, fixed osteoderms; K–R, (semi) mobile osteoderms. **A**, MLP 69-IX-8-13AC**; **B**, MLP 70-XII-29-1**; **C**, MLP 41-XII-13-905; **D**, MLP 69-IX-8-13AF; **E**, MLP 69-IX-8-13AG; **F**, MLP 41-XII-13-414A; **G**, MLP 69-IX-8-13AN; **H**, unknown catalog number; **I**, MLP 69-IX-8-13AK; **J**, MLP 41-XII-13-414B; **K**, MLP 69-IX-8-13AS; **L**, MLP 69-IX-8-13AE**; **M**, MLP 52-X-1-36; **N**, MLP 69-IX-8-13AB*; **O**, MLP 41-XII-13-909; **P**, MLP 69-IX-8-13AW; **Q**, MLP 69-IX-8-13AQ; **R,** MLP 69-IX-8-13AY. Scale bar equal to 20 mm.

Figure 11. Hypothetical time-calibrated phylogeny of the clade *Scirrotherium + Kraglievichia + Holmesina* based on the strict consensus tree under implied weights (Fig. 2(B)). Polytomies were resolved by (1) forcing the monophyly of *S. hondaensis* and *S. antelucanus* and (2) placing the species *H. septentrionalis* and *H. occidentalis* as successively basal to the largest South American *Holmesina* species, i.e., *H. paulacoutoi* and *H. major*. Note the diversification events of the clade *Scirrotherium + Kraglievichia + Holmesina* are mainly concentrated during the Burdigalian (late Early Miocene) and Plio-Pleistocene. Likewise, note the relative long ghost lineage of *Holmesina*. Images of the pampatheriids are from *PhyloPic* (all available under public domain): top, *Pampatherium humboldtii* (http://phylopic.org/name/670230e9-4775-493c-b3ab-31718fb570a3); below, *Holmesina floridanus* (http://phylopic.org/name/73635941-ed8a-4518-aae8-70e824dbee97).

Figure 12. Geographic distributions of *Scirrotherium*, *Kraglievichia* and *Holmesina* during the Neogene and Pleistocene. The symbols (i.e., circles and triangles) should not necessarily be interpreted as single localities but as approximate areas of occurrence. This

is especially true for the Pliocene and Pleistocene epochs. Further details on the

biogeography of these genera in the *Discussion* section. Scale bar equal to 2000 Km.

TABLES

Table 1. Fixed (scapular and pelvic) osteoderm measurements for taxa of interest in this study.

Taxon/Measurement	Length	Width	Thickness	References
S. hondaensis	16-35.2	17.5-27.9	3.7-6.9	This work; Góis et al. 2013
S. antelucanus	28.6-40.9	22-32.4	4.9-7.1	This work; Laurito and Valerio 2013
K. carinatum comb. nov.	20.9-33.5	17-26.1	4.1-5.9	This work; Góis et al. 2013
K. paranensis	30-45	22.5-28.3	6-11	Góis et al. 2013
H. floridanus	24.4-36.7	18.9-32.1	6-9.7	This work; Edmund 1987

Table 2. Mobile and semi-mobile osteoderm measurements for taxa of interest in this study.

Taxon/Measurement	Length	Width	Thickness	References
S. hondaensis	29.4-60	17.9-27.4	4.9-7.3	This work; Góis et al. 2013
S. antelucanus	38.2-64.6	19.4-28.9	-	Laurito and Valerio 2013
K. carinatum comb. nov.	32-54.5	17-28.9	3.9-6	This work; Góis et al. 2013
K. paranensis	60.5-70.5	25-29	7-9	Góis et al. 2013
H. floridanus	61.8-71	17.6-28.5	4.7-6.3	This work; Edmund 1987

Table 3. Selected cranial measurements for VPPLT 706 of *Scirrotherium hondaensis* and related taxa whose skulls are known.

Taxon/Measurement	GSL	NL	FL	PAL	LUR	PL	References
S. hondaensis	117.3*	~52.8	~55	-	84.1	94.3	This work
K. cf. *paranensis*	194	58	62	74	-	159	This work
*H. floridanus***	249	106.3	75	58.6	133.6	185	This work
H. septentrionalis	290	-	-	-	165	220	Góis et al. 2012

*Incomplete

**Specimen UF 191448

Table 4. Femoral measurements for *Scirrotherium hondaensis* and related taxa whose femur is known.

Measurement/Taxon	GFL	TTW	DW	References
S. hondaensis	162*	27.6	32.5	This work
K. cf. *paranensis*	164	33.7	38	This work
*H. floridanus***	195	41	47	This work
H. septentrionalis	290	70	86	Góis 2013

*Estimated from the specimens VPPLT 706 and UCMP 39846

**Specimen UF 24918

Figure 1

A. Early-to-Middle Miocene Castilletes Formation

B. Middle Miocene La Victoria Formation

C. Middle Miocene Villavieja Formation

D. Late Miocene-to-Early Pliocene Sincelejo Formation

E. Late Pliocene Waré Formation

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11

Figure 12