SYSTEMATIC REVISION AND REDEFINITION OF THE GENUS *SCIRROTHERIUM*

EDMUND & THEODOR, 1997 (CINGULATA, PAMPATHERIIDAE): IMPLICATIONS

FOR THE ORIGIN OF PAMPATHERIIDS AND THE EVOLUTION OF THE SOUTH

AMERICAN LINEAGE INCLUDING *HOLMESINA*

By KEVIN JIMÉNEZ-LARA[1,2]

[1] División de Paleontología de Vertebrados, Museo de La Plata, Facultad de Ciencias

Naturales y Museo, Universidad Nacional de La Plata, Paseo del Bosque s/n, B1900FWA

La Plata, Argentina.

[2] CONICET, Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina.

e-mail: kjimenezlara@fcnym.unlp.edu.ar

Abstract: The intrageneric relationships of the pampatheriid genus *Scirrotherium* and its

affinities with supposedly related genera, i.e. *Kraglievichia* and *Holmesina*, are revised

through parsimony phylogenetic analyses and new comparative morphological

descriptions. For this work was analyzed unpublished material of pampatheriids (numerous

osteoderms, one partial skull and a few postcranial bones) from Neogene formations of

Colombia. The results show that *Scirrotherium* is paraphyletic if we include all its referred

species, i.e. *Scirrotherium hondaensis*, *S. carinatum* and *S. antelucanus*. The species *S.*

carinatum is closer to *Kraglievichia paranensis* than to *S. hondaensis* or *S. antelucanus*,

then it is proposed the new name *K. carinatum* comb. nov. The relationships between *S.*

hondaensis and *S. antelucanus* could not be resolved, so these species should be designated

in aphyly. In spite of failing to recover *S. hondaensis* and *S. antelucanus* as one single

clade, here is preferred to maintain the generic name *Scirrotherium* in both species from

diagnostic evidence. New emended diagnoses for *Scirrotherium*, *S. hondaensis* and

Kraglievichia are provided. The genus *Holmesina* was found monophyletic and located as

the sister clade of *Scirrotherium + Kraglievichia*. The evolutionary and biogeographical

implications of the new phylogeny and taxonomical re-arrangements are discussed. It is

claimed a possible geographical origin of the family Pampatheriidae and *Scirrotherium* in

low latitudes of South America as early as Early Miocene (Burdigalian) times. The South

American ancestor or sister taxon of *Holmesina* is predicted as morphologically more

similar to *Scirrotherium* than to *Kraglievichia*.

Key words: Pampatheriidae, *Scirrotherium*, *Kraglievichia*, *Holmesina*, Great American

Biotic Interchange, Neogene.

The pampatheriids (Pampatheriidae) are a morphologically conservative extinct clade of

glyptodontoid cingulates or highly armored xenartrans (Glyptodontoidea *sensu* McKenna &

Bell 1997) with medium-to-large body sizes (Edmund 1985; Góis et al. 2013). They were

distributed from the Neogene to the Early Holocene in numerous localities of South

America (their native range), Central America, Mexico and the United States (Edmund

1985; Vizcaíno et al. 1998; Rincón et al. 2014; Góis et al. 2015 and references there). As

the modern armadillos (Dasypodidae), pampatheriids have a flexible carapace by the

presence of three transverse bands of imbricated osteoderms which conform a kind of

"articulation" between the scapular and pelvic shields (Edmund 1985). The pampatheriids

also have multiple features, especially in their skull and mandible, which, collectively,

define them as the sister group of glyptodontids –Glyptodontidae (Gaudin 2004; Gaudin &

Wible 2006; Billet et al. 2011; Delsuc et al. 2012), namely deep horizontal mandibular

ramus, laterally-directed zygomatic root, transversely-wide glenoid fossa, rough pterygoids,

among others (Gaudin & Wible, 2006).

The fossil record of Pampatheriidae is mainly represented by isolated specimens, of which

most of them are osteoderms and, in lesser extent, skulls, mandibles and postcranial bones;

fairly complete and articulated skeletons are uncommon (Edmund 1985; Góis 2013). Due

the above, the systematics of this xenartran group has historically been based on

osteodermal characters (Edmund 1985, 1987; Góis et al. 2013), as has been the case with

other cingulate clades. Overall, nearly two tens of pampatheriid species and seven genera

are known (Góis 2013). The latter conform two possible subfamilial lineages: (1) that

including to the genera *Plaina* and *Pampatherium*; and (2) that comprising the genera

Scirrotherium, *Kraglievichia* and *Holmesina* (Edmund 1985). However, it does not exist in

the scientific literature a published phylogenetic analysis on the relationships between the

different pampatheriid genera. Only Góis (2013) performed a phylogenetic analysis for

these taxa, but his results have not been published. In Góis's consensus tree, it was

corroborated the hypothesis by Edmund (1985) on the two subfamilial lineages.

The genus *Scirrotherium* is the oldest known pampatheriid in the fossil record (Góis et al.

2013; Rincón et al. 2014) and one of the four Miocene genera (the other ones are

Kraglievichia Castellanos 1927; *Vassallia* Castellanos, 1927; and *Plaina* Castellanos,

1937). This taxon was originally described by Edmund & Theodor (1997) from

craniomandibular, postcranial and osteodermal specimens collected in the Middle Miocene

(Serravalian) sedimentary sequence of the La Venta area, southwestern Colombia. These

authors interpreted that the only known species in that time and type species, *Scirrotherium*

hondaensis, has plesiomorphic traits in its osteological morphology which are expected for

its antiquity. Additionally, they highlighted morphological similarity of *S. hondaensis* with

the species *Vassallia minuta* (Late Miocene of southern and central South America; De

Iullis & Edmund 2002), more than with any other pampatheriid.

Later, Góis et al. (2013) described a second species for *Scirrotherium*, *S. carinatum*, from

the Late Miocene (Tortonian) of northeastern and southern of Argentina-northwestern

Brazil. In northeastern Argentina (Entre Ríos Province), *S. carinatum* shares basal

stratigraphic levels ("Conglomerado Osífero", literally 'bone-bearing conglomerate') in the

Ituzaingó Formation with the middle-sized pampatheriid *Kraglievichia paranensis* (Góis et

al. 2013; Scillato-Yané et al. 2013), a taxon different but not distantly related to

Scirrotherium as previously indicated. The species *S. carinatum*, based exclusively on

osteoderms of different regions of the armored carapace, has a body size comparable or

slightly smaller than that of *S. hondaensis* (Góis et al. 2013).

The phylogenetic analysis conducted by Góis (2013) recovered in polytomy to *S. carinatum*

and *S. hondaensis*, being one of these species or both of them the sister taxon/taxa of the

clade *Kraglievichia* + *Holmesina* (except *H. floridanus*). It is notorious the non-basal

position of the *Scirrotherium* species within the general topology of the cladogram despite

they conform the oldest known genus. Instead, these species are closely placed to widely

recognized terminal taxa, i.e. *Holmesina* spp. (Edmund 1985, 1987; Gaudin & Lyon 2017).

If this result is correct, it would indicate a significantly long ghost lineage at the base of the

evolutionary tree of Pampatheriidae.

Góis found phylogenetic support for *Scirrotherium* through one single synapomorphy, i.e.

presence of deep longitudinal depressions (LDs) in osteoderms. This is a distinctive feature

of *S. carinatum* but, in contrast, the LDs in *S. hondaensis* are relatively shallow, rather than

deep. Interestingly, this putative synapomorphy is actually shared by *K. paranensis*. Góis

explained the insufficient phylogenetic resolution in his analysis for *Scirrotherium* using

the argument of fragmentary character of the fossil specimens of *S. hondaensis*, although in

the case of *S. carinatum*, unlike the Colombian species, the skull, mandible and any

postcranial bone are unknown (Góis et al. 2013).

Almost simultaneously to Góis's works, Laurito and Valerio (2013) reported and studied

new pampatheriid material from the Late Miocene (Tortonian to Messinian) of Costa Rica,

which was assigned to a new species, *S. antelucanus*. This species, the largest referred to

Scirrotherium so far (body size comparable or slightly smaller than that of *K. paranensis*;

Laurito & Valerio 2013), is based on osteoderms and some postcranial bones (femoral

fragments and metatarsals). The occurrence of *S. antelucanus* in the Late Miocene of

southern Central America suggests the genus *Scirrotherium* take part earlier than any other

pampatheriid (i.e. *Plaina*, *Pampatherium*, *Holmesina*; Woodburne 2010) in the late

Cenozoic biotic interchanges of the Americas with its invasion to tropical North America

("North America" is defined here as all the continental territories north of the ancient

location of the main geographical barrier between the Americas during the early Neogene,

i.e. the Central American Seaway in northwestern Colombia), before the definitive closing

of the Panama Land Bridge (PLB) ca. 3 mya (Schmidt 2007; Coates & Stallard 2013;

O'dea et al. 2016; Jaramillo 2018).

Recently, in several contributions on fossil vertebrate assemblages from the Neogene of

Venezuela and Peru has been reported the occurrence of isolated osteoderms referred to

Scirrotherium. In basis to these discoveries, the geographical and chronological distribution

of the genus has been expanded in such a way that this taxon is now known for the Early

and Late Miocene (Burdigalian and Tortonian) of northwestern Venezuela (Rincón et al.

2014; Carrillo-Briceño et al. 2018) and Late Miocene (Tortonian) of northeastern Peru

(Antoine et al. 2016).

Assuming all the previous taxonomical assignments are correct, the latitudinal range of

Scirrotherium, from southern Central America to Patagonia (southern Argentina), is the

widest latitudinal range of a Miocene pampatheriid, only comparable with those of the Plio-

Pleistocene forms *Pampatherium* and *Holmesina* (Scillato-Yané et al. 2005). This

biogeographical inference provides support to the hypothesis that *Scirrotherium* inhabited

varied environments within its latitudinal range, and, consequently, it probably had a

relatively high ecological flexibility (Góis et al. 2013).

Despite the progress in the systematic and biogeographical research of *Scirrotherium*, it is

necessary at present a new reevaluation of several fundamental hypotheses about this taxon,

including its taxonomical definition, monophyly and evolutionary relationships with other

pampatheriid genera. Using parsimony phylogenetic analyses and morphological

comparative descriptions of new pampatheriid remains from the Neogene of Colombia, this

contribution reevaluates the taxonomic status of *Scirrotherium* and its relationships with

supposedly allied genera, i.e. *Kraglievichia* and *Holmesina*. Accordingly, I suggest a new

taxonomical and nomenclatural reorganization with emended diagnoses for *Scirrotherium*

and, complementarily, for the genus *Kraglievichia*.

Finally, considering the systematic reanalysis, I depict a model of biogeographical

evolution for the lineage *Scirrotherium-Kraglievichia-Holmesina*. From this model, I draw

out new hypotheses on the geographical origin of Pampatheriidae and the late Cenozoic

dispersal events of pampatheriids to/from North America, including a possible re-entry

event to South America of the species *S. antelucanus*.

MATERIAL AND METHODS

Selection of OTUs

I studied 12 species of pampatheriids attributed to six different genera. These species, in

alphabetic order, are: *Holmesina floridanus* Robertson, 1976; *H. major* Lund, 1842; *H.*

occidentalis Hoffstetter, 1952; *H. paulacoutoi* Cartelle & Bohórquez, 1985; *H.*

septentrionalis Leidy, 1889; *Kraglievichia paranensis* Ameghino, 1888; *Pampatherium*

humboldtii Lund, 1839; *Plaina intermedia* Ameghino, 1888; *Scirrotherium antelucanus*

Laurito & Valerio, 2013; *S. carinatum* Góis, Scillato-Yané, Carlini and Guilherme, 2013; *S.*

hondaensis Edmund & Theodor, 1997; and *Vassallia minuta* Moreno & Mercerat, 1891. It

was also included in this selection unidentified pampatheriid material (MUN STRI 16718

and 38064; see the section *Institutional abbreviations*) from the Castilletes Formation in

Colombia (see below), which is referred as "Castilletes specimens".

Among the former nominal species, I follow to Góis (2013) in considering *Vassallia*

maxima as a junior synonym of *Pl. intermedia*. The only one species of *Holmesina* not

included in this study was *H. rondoniensis* Góis, Scillato-Yané, Carlini & Ubilla, 2012.

This decision is based on a preliminary phylogenetic analysis in which *H. rondoniensis* was

identified as a "wildcard" taxon obscuring the phylogenetic resolution. It was also not

included in this analysis the species *Tonnicinctus mirus* Góis, González Ruiz, Scillato-Yané

& Soibelzon, 2015, which is considered a potentially problematic and late diverging taxon

without any apparent substantial interest with respect to the systematic issues here

addressed.

Morphological description of the specimens

The osteological morphology of the selected species was revised from direct observations

on specimens previously studied in other works, as well as through published/unpublished

descriptions (Simpson 1930, Castellanos 1937; Edmund 1985, 1987; Edmund & Theodor

1997; Góis 2013; Góis et al. 2013; Laurito & Valerio 2013; Scillato-Yané et al. 2013; Góis

et al. 2015; Gaudin & Lyon 2017). Naturally, according to the objectives of this research,

during the revision of material I focused on the species *S. antelucanus*, *S. carinatum* and *S.*

hondaensis, and, additionally, species of genera considered closely allied to *Scirrotherium*,

i.e. *K. paranensis* and *Holmesina* spp. (particularly *H. floridanus*; Appendix S1 of the

Supplementary Material).

On other hand, new undescribed cranial, postcranial and osteodermal specimens were also

used to reexamine the morphological variability of *Scirrotherium*. This material comes

from five Neogene geological units of Colombia (Fig. 1): (1) Castilletes Formation (Early

to Middle Miocene, late Burdigalian-Langhian), Municipality of Uribia, Department of La

Guajira; (2) La Victoria Formation (late Middle Miocene, Serravalian), Municipality of

Villavieja, Department of Huila; (3) Villavieja Formation (late Middle Miocene,

Serravalian), Municipality of Villavieja, Department of Huila; (4) Sincelejo Formation

(Late Miocene-Early Pliocene, Messinian-Zanclean), Municipality of Los Palmitos,

Department of Sucre; (5) Ware Formation (Late Pliocene, Piacenzian), Municipality of

Uribia, Department of La Guajira. For detailed lithological descriptions and

chronostratigraphic inferences on these formations, the reader is referred to the following

references: Moreno et al. 2015 for the Castilletes and Ware Formations; Guerrero 1997,

Flynn et al. 1997 and Anderson et al. 2016 for the La Victoria and Villavieja formations;

and Flinch 2003, Villarroel & Clavijo 2005, Bermúdez et al. 2009 and Alfaro & Holz 2014

for the Sincelejo Formation. The new fossils are deposited at the Paleontological Collection

of the Museo Mapuka de la Universidad del Norte, Barranquilla, Colombia, except those

collected in the La Victoria and Villavieja Formations. The latter are housed at the Museo

de Historia Natural La Tatacoa, La Victoria Town, Municipality of Villavieja, Department

of Huila, Colombia.

Cranial measurements, all taken on the midline of the skull (dorsally or ventrally), follow

Góis (2013). The anatomical terminology for osteoderms is based on the proposal of Góis

et al. (2013). All the measurements were taken with a digital caliper with a precision of

0.01 mm.

Selection and codification of characters

I exclusively selected cranial, dental and osteodermal characters given that the postcranial

bones of most species of Pampatheriidae is poorly known (Góis 2013). This selection was

based on personal observations and previous quantitative and qualitative analyses of the

interspecific, intergeneric and familial morphological variability of pampatheriids (e.g.

Edmund 1985, 1987; Góis 2013; Góis et al. 2013; Laurito & Valerio 2013). Overall, a

matrix of 27 characters (Appendix S2 of the Supplementary Material) was built and

managed on Mesquite version 2.75 (Maddison & Maddison 2010). In this character list, 20

characters are parsimony-informative and 7 are parsimony-uninformative; 5 are

osteodermal characters and 22 are cranial or dental. The parsimony-uninformative

characters allow to define potential autapomorphies of the studied taxa.

Cladistic analyses

Parsimony analyses under schemes of equal weights and implied weights (characters

reweighted *a posteriori*; see below) were performed in PAUP* version 4.0a142 (Swofford

2015). In both weighting schemes, the species *P. humboldtii*, *Pl. intermedia* and *V. minuta*

were defined as outgroup. The monophyly of the outgroup and rooting of trees from it was

constrained. The selection of the outgroup is based on the hypothesis about subfamilial

relationships of Pampatheriidae by Edmund (1985). The characters were treated as

unordered. The criterion for character optimization was DELTRAN (see Gaudin 2004 for

justification of this configuration). The analyses consisted of heuristic searches with

random addition sequences of 1000 replicas. For reordering of branches, it was selected the

TBR algorithm. The topological results of most parsimonious trees were summarized

through strict consensus trees.

The methodology of implied weights is intended to mitigate potential biases by limited

number of characters (especially osteodermal characters, as consequence of the

evolutionary trend in Pampatheriidae towards a simplification of the ornamentation in

comparison with that in other cingulate clades, e.g. Glyptodontidae) and the effect of

homoplastic characters (Goloboff et al. 2008; Goloboff 2014). Characters were reweighted

using the rescaled consistency index (mean value) of the equally-weighted parsimony

analysis (see Ausich et al. 2015 and references there for justification of the use of rescaled

consistency index for implied-weights parsimony analyses). A default concavity value ($k =$

3) was selected (Goloboff et al. 2018). Three successive rounds of character reweighting

were needed until identical set of strict consensus trees were found in two consecutive

searches (Swofford & Bell 2017).

Node support for the strict consensus tree resulting from the equally-weighted analysis was

evaluated from absolute Bremer support values, while the node stability for the strict

consensus tree obtained from the implied weighting analysis was evaluated using a

bootstrap resampling procedure. The software FigTree v1.4.3

(http://tree.bio.ed.ac.uk/software/figtree/) was used as graphical viewer and editor of

cladograms.

Taxonomical and nomenclatural criteria

I applied a taxonomical and nomenclatural criterion reasonably, but not strictly, constrained

by the phylogeny. This implies looking for a natural classification (i.e. based on

monophyletic groups) without ignoring possible limitations of the phylogenetic inference

related to the available information in the fossil record and major morphological gaps.

Additionally, it was used open nomenclature to indicate taxonomical uncertainty when

necessary, following general recommendations of Bengston (1988) and updated definitions

by Sigovini et al. (2016) for the qualifiers of this semantic tool of taxonomy.

Institutional abbreviations

CFM, Museo Nacional de Costa Rica, Colección de fósiles de la sección de Geología, San

José, Costa Rica; FMNH, Field Museum Natural History, Chicago, Illinois, USA; Museo

Argentino de Ciencias Naturales "Bernardino Rivadavia", Colección de Paleovertebrados,

Ciudad Autónoma de Buenos Aires, Argentina; MLP, Museo de La Plata, La Plata,

Argentina; MUN STRI, Museo Mapuka de la Universidad del Norte, Colección de

paleontología, Barranquilla, Colombia; UCMP, University of California Museum of

Paleontology, Berkeley, California, USA; UF, Florida Museum of Natural History,

Gainesville, Florida, USA; VPPLT, Museo de Historia Natural La Tatacoa, Colección de

paleontología, La Victoria Town, Huila, Colombia.

Anatomical abbreviations

AM, anterior margin; FL, frontal bone length; GFL, greatest femoral length; GSL, greatest

skull length; LCE, longitudinal central elevation; LD, longitudinal depression; LM, lateral

margin; LUR, length of the upper teeth row; ME, marginal elevation; Mf, upper

molariform; mf, lower molariform; NL, nasal bone length; PAL, parietal bone length; PL,

hard palate length; TTW, maximum width at the third trochanter of the femur; DW,

maximum width of the femoral distal epiphysis.

RESULTS

Cladistic analyses

The parsimony analysis with equal weights obtained 107 most parsimonious trees (MPTs),

each one of these with a tree length of 44 steps (consistency index = 0.909; retention index

= 0.907; rescaled consistency index = 0.825). The strict consensus tree from these MPTs

(Fig. 2A; tree length = 52; consistency index = 0.769; retention index = 0.721; rescaled

consistency index = 0.555) is not fully resolved because it has two polytomies. One of these

polytomies involves the species *S. hondaensis*, *S. antelucanus* and *H. floridanus*, while the

other one is formed by *H. septentrionalis*, *H. major*, *H. paulacoutoi* and *H. occidentalis*.

Three clades were recovered (excluding that of the entire ingroup): (1) All the ingroup taxa

except "Castilletes specimens"; (2) *S. carinatum* + *K. paranensis*; and (3) *Holmesina* spp.

except *H. floridanus*. On other hand, the parsimony analysis with implied weights yielded

30 most parsimonious trees (MPTs), each one of those with a tree length of 109 weighted

steps (consistency index = 0.982; retention index = 0.982; rescaled consistency index =

0.964). The strict consensus tree from the MPTs (Fig. 2B; tree length = 91; consistency

index = 0.978; retention index = 0.980; rescaled consistency index = 0.959), like that

produced by the equally weighted approach, is not fully resolved. Again, two polytomies,

but in this case the polytomy including *S. hondaensis*, *S. antelucanus* and *H. floridanus* was

altered. The latter taxon is placed as the basal-most *Holmesina* species. The polytomy

formed by *H. septentrionalis*, *H. major*, *H. paulacoutoi* and *H. occidentalis* was

unmodified. As consequence of the relocation of *H. floridanus* within the topology, four

clades were recovered: (1) All the ingroup taxa except "Castilletes specimens"; (2) *S.*

carinatum + *K. paranensis*; (3) *Holmesina* spp.; and (4) *H. septentrionalis*, *H. major* + *H.*

paulacoutoi + *H. occidentalis*. According to the two schemes of weighting for the

parsimony analyses, the genus *Scirrotherium* is paraphyletic if it is composed by *S.*

antelucanus, *S. hondaensis* and *S. carinatum*. *S. carinatum* is closer to *K. paranensis* than

to *S. hondaensis* or *S. antelucanus*. The relationships among *S. hondaensis* and *S.*

antelucanus is not resolved in either of the two strict consensus trees.

In the strict consensus tree from the equally-weighted analysis there is moderate node

support for the clade *S. carinatum* + *K. paranensis* (Bremer support value = 3). The clades

(1) *Scirrotherium* spp. + *K. paranensis* + *Holmesina* spp.; and (2) *Holmesina* spp. are

weakly supported (Bremer support values equal to 1 in both clades). The node stability

analysis under implied weights (Fig. 2C) shows high resampling frequencies for the clades

(1) *Scirrotherium* spp. + *K. paranensis* + *Holmesina* spp. (bootstrap value = 100); (2) *S.*

carinatum + *K. paranensis* (bootstrap value = 100); (3) *Holmesina* spp. (bootstrap value =

99); and (4) *H. septentrionalis* + *H. major* + *H. paulacoutoi* + *H. occidentalis* (bootstrap

value = 100). Low resampling frequencies are related to the clades (1) *Scirrotherium* spp. +

K. paranensis (bootstrap value = 59); and (2) *S. hondaensis* + *S. carinatum* + *K. paranensis*

(bootstrap value = 60).

SYSTEMATIC PALAEONTOLOGY

XENARTHRA Cope, 1889

CINGULATA Illiger, 1811

GLYPTODONTOIDEA Gray, 1869

Family PAMPATHERIIDAE Paula Couto, 1954

Genus *Scirrotherium* Edmund & Theodor, 1997

LSID. urn:lsid:zoobank.org:act:313358B5-3B1F-4902-8C2E-BB07CFCBEE18

Type species: *Scirrotherium hondaensis* Edmund & Theodor, 1997 by original designation.

Included species: In addition to the type species, *S. antelucanus* Laurito & Valerio 2013.

Emended diagnosis: A pampatheriid of small-to-middle body size that can be distinguished

from other pampatheriids by the following combination of features: thin non-marginal fixed

osteoderms (~3.5–7 mm in thickness); slightly to moderately rough external surface of

osteoderms; external surface of osteoderms with a sharp and uniformly narrow LCE; LCE

from superficial to well-elevated; superficial to shallow LDs with gently slope towards the

MEs; frequently one single, transversely elongated row of large foramina in the AM of

fixed osteoderms; between 6 and 11 anterior foramina as maximum number of foramina per

row.

Discussion: The taxonomical status of *Scirrotherium* is saved from invalidity by paraphyly

by exclusion of the species 'S.' *carinatum* from the genus (see below). However, according

to the preferred phylogenetic hypothesis presented here, i.e. the strict consensus tree from

the parsimony analysis under implied weights (Fig. 2B), the other two referred species of

Scirrotherium (*S. antelucanus* and *S. hondaensis*) should be designated in aphyly because

they do not have resolved relationships between them (see Ebach & Williams 2010 for

details about the phylogenetic concept of aphyly). Until new evidence available,

maintenance of the taxonomical validity of *Scirrotherium*, as defined here, is based on the

emended diagnosis of this taxon, which is partially built from ambiguous synapomorphies,

as well as from qualitative and quantitative morphological differences with other generic

taxa.

Stratigraphic and geographical distribution: [Tentatively, by badly preserved material]

Castillo Formation, upper Lower Miocene, upper Burdigalian, Lara State, Venezuela

(Rincón et al. 2014); Castilletes Formation, upper Lower to lower Middle Miocene, upper

Burdigalian to Langhian; Department of La Guajira, Colombia; La Victoria Formation and

Villavieja Formation, upper Middle Miocene, Serravalian, Department of Huila, Colombia

(Edmund & Theodor 1997); Caujarao Formation, lower Upper Miocene, Tortonian, Falcon

State, Venezuela (Carrillo-Briceño et al. 2018); Curré Formation, Upper Miocene,

Puntarenas Province, Costa Rica (Laurito & Valerio 2013); Sincelejo Formation, Upper

Miocene to Lower Pliocene, Messinian-Zanclean, Department of Sucre, Colombia; Ware

Formation, Upper Pliocene, Piacenzian, Department of La Guajira, Colombia.

Scirrotherium hondaensis Edmund & Theodor, 1997

LSID. urn:lsid:zoobank.org:act:E3B83181-91D6-44C8-90C0-BBAACEC2CDEE

Holotype: UCMP 40201, incomplete skull and left hemimandible.

Type locality and horizon: Municipality of Villavieja, Department of Huila, Colombia. La

Victoria Formation, upper Middle Miocene, Serravalian.

Referred material: VPPLT 004, several fixed osteoderms; VPPLT 264, several fixed

osteoderms and one semi-mobile osteoderm; VPPLT 348, tens of fixed and (semi) mobile

osteoderms; VPPLT 701, several fixed osteoderms; VPPLT 706, one anterior skull, one

femoral diaphysis, one ulna without distal epiphysis, several vertebrae and numerous fixed

and (semi) mobile osteoderms; VPPLT 1683 - MT 18, several fixed and (semi) mobile

osteoderms; UCMP 39846, one proximal femoral epiphysis, one left calcaneum and one

left astragalus. All the osteoderms referred to *S. hondaensis* are illustrated in the Fig. 3.

Other important specimens are illustrated in the Figs. 4–7.

Stratigraphic and geographical provenance: The samples VPPLT 004, 264, 701, 706 and

(partially) 1683 - MT 18 were collected in the La Victoria Formation, upper Middle

Miocene (Serravalian; see the Figs. 3–6 for more details on the stratigraphic provenance of

individual specimens), while the sample UCMP 39846 and part of VPPLT 1683 - MT18

comes from the Villavieja Formation, upper Middle Miocene (Serravalian).

Emended differential diagnosis: Pampatheriid of small body size that differs from other

pampatheriids on this unique combination of characters: external surface of osteoderms

with ornamentation (especially the LCE and MEs), in general terms, more protuberant than

in *S. antelucanus*, but less than in *Kraglievichia*; size range of fixed osteoderms smaller

than in *S. antelucanus* and similar to that in *Kraglievichia carinatum* comb. nov. (= '*S.*'

carinatum; Góis et al. 2013; see below); fixed osteoderms generally thicker than in *K.*

carinatum comb. nov. but less than in *K. paranensis*, similar to *S. antelucanus*; anterior

foramina smaller than in *S. antelucanus*; anterior foramina in fixed osteoderms usually

aligned in one individual row, although infrequently these osteoderms show an extra, short

or reduced row of anterior foramina; two rows of anterior foramina in mobile osteoderms,

similar to *Vassallia* (Góis 2013); last lower molariform (mf9) incipiently bilobed; frontals

prominently convex in lateral view, with this convexity in a posterior position to the

insertion of the anterior root of the zygomatic arch; anterior root of the zygomatic arch

posterolaterally projected with respect to the main body of maxilla.

Comparative description

For the original and detailed description of this species, including its osteoderms, see

Edmund & Theodor (1997). See the Tables 1 and 2 for an updated compilation of

osteodermal measurements of referred *Scirrotherium* species and comparisons with those of

related taxa. Below there are descriptions of osteological structures and traits incompletely

known or unknown for *S. hondaensis* so far.

Skull: The holotype of *S. hondaensis* UCMP 40201 includes a very fragmentary skull. This

specimen does not preserve the anterior end of the rostrum, the most of the bone

architecture at the orbit level (both dorsally and ventrally), part of the upper dental series,

ear region, braincase nor occipital region. Comparatively, the skull of the sample VPPLT

706 (Fig. 4), here originally described, is more complete, despite it also has some missing

structures. This new, small skull (see Table 3 for morphometric comparisons) is relatively

well preserved from the orbit level to the most anterior end of rostrum, except for the

anterior zygomatic arch and nasals dorsally. It also has a less deformed rostrum than that of

the holotypic skull of this species. The general aspect of the new skull is similar to those of

all known skulls of Pampatheriidae. In lateral view, this is markedly depressed towards its

anterior end. In dorsal view, it is tapered also towards its anterior rostrum, where it ends

abruptly. Proportionally, the rostrum is shorter than in *K. paranensis* and even more than in

H. floridanus. In lateral view, the facial process of premaxilla is less defined than in *H.*

floridanus and the premaxilla-maxilla suture has a convex form, like the former species

(Gaudin & Lyon, 2017). The anterorbital fossa is arranged more vertically than in *K.*

paranensis and *H. floridanus*. The lacrimal is, proportionally, the largest among

pampatheriids. This bone severely restricts the frontomaxillary contact in lateral view,

similarly to *K. paranensis*. The dorsal contribution of the lacrimal to the orbit is,

proportionally, greater than in *H. floridanus* and similar to that in *K. paranensis*. The

lacrimal foramen is anteriorly located and close to the anterior border of the orbit. The

anterior root of zygomatic arch is projected posterolaterally, unlike other pampatheriids

whose skull is known (lateral projection). The frontals show a conspicuous convexity in a

posterior position to the insertion of the anterior root of zygomatic arch, in such a way that

the posterior section of frontals is placed in a very different plane with respect to that of the

anterior section of the same bones. Dorsally, the frontals are more anteroposteriorly

elongated and more laterally expanded that in *K. paranensis*, similar to *Holmesina* spp.

Ventrally, the hard palate has a wide aspect since the rostrum is shortened in comparison

with other pampatheriids. Only two anterior molariforms are preserved (one Mf1 and one

Mf2 in different rows) and inferences about upper dentition are made from the alveoli. The

upper dental series, as in all the members of the family, is composed by nine molariforms.

Of these teeth, the last five (Mf5-Mf9) are bilobed. The anterior molariforms (Mf1-Mf4)

converge anteriorly between them, but they do not imbricate. The former teeth are rounded

to elliptical, similarly to the condition observed in *H. floridanus*. They also are less

mesiodistally elongated than in *K. paranensis*. The molariforms with greatest occlusion

area are the fifth and sixth (Mf5 and Mf6). The area of occlusion of the upper molariforms

decrease distally from the fifth and sixth molariforms to the ninth, as in all the

pampatheriids. The last upper molariform (Mf9) is the smallest of lobed molariforms and

has the lesser degree of lobulation (elliptical shape for the material described by Edmund &

Theodor 1997). In ventral view, the specimen VPPLT 706 is characterized by a gradual

lateral widening of the maxilla from the level of the anterior border of the fifth upper

molariform (Mf5). It is preserved the anterior section of palatines as far as a level slightly

posterior to the last molariform (Mf9). The maxilla-palatine suture is not recognizable.

Femur: This bone in *S. hondaensis* was unknown so far, despite the existence of a pair of

epiphyses (proximal and distal) from a left femur at the UCMP collections (UCMP 39846).

Within the sample VPPLT 706 there is a left femur (Fig. 5A–D) without epiphyses

(apparently it is not the same bone from which come the previously referred epiphyses).

Thus, the complementary description of all these anatomical elements allows to reconstruct

the overall femoral appearance and features. The estimated proximo-distal length of this

appendicular bone is ca. 162 mm and its transverse width at third trochanter is 27.6 mm.

These morphometric values are the smallest ones for known femora of Pampatheriidae

(Table 4). They are only comparable to those of the specimen MLP 69-IX-8-13A which

was referred to *K.* cf. *paranensis* (Góis 2013; Scillato-Yané et al. 2013). The femoral head

is hemispheric and the greater trochanter is less high than in *K.* cf. *paranensis*, similar to

the condition observed in *H. floridanus*. However, the greater trochanter has a more tapered

proximal end than in the latter species. In *S. hondaensis*, the lesser trochanter is less

mediolaterally expanded than in *K.* cf. *paranensis*. The femoral diaphysis is less curved

than in *K.* cf. *paranensis*, similar to that in *H. floridanus*. The border located laterally and

distally with respect to the third trochanter is more curved than in *H. floridanus*, similar to

that of *K*. cf. *paranensis*. The third trochanter is, proportionally, larger than in *K*. cf.

paranensis, but less than in *H. floridanus*. The former bone projection is poorly tapered in

comparison with the same structure in *P. humboldtii*. The patellar facets are less defined or

delimited than in *K*. cf. *paranensis*. In *S. hondaensis* these facets are oriented toward the

center of the anterior surface of the distal epiphysis, rather than laterally like in *K*. cf.

paranensis and *H. floridanus*.

Ulna: This bone is also described here for the first time. In this case, a right ulna (Fig. 5E–

H) without part of the diaphysis and the distal epyphysis. The (incomplete) proximo-distal

length is 89.5 mm. The olecranon is elongated and protuberant. In internal view, it is less

proximally tapered than in *H. floridanus*. The lateral entrance to the trochlear notch is more

restricted than in *Pampatherium*, similarly to *Holmesina*. Likewise, it is less proximo-

distally elongated than in *H. floridanus*, more similar to that in *H. paulacoutoi*. Proximally,

at the level of the trochlear notch, the posterior border is uniformly convex, not slightly

concave, like in *H. floridanus*. The depression for the insertion of the anconeus muscle is

deep and more proximally located than in *Pampatherium*, similar to *Holmesina*.

Vertebrae: Some vertebrae are also reported (Fig. 6). One of these is a thoracic vertebra and

five are caudal vertebrae, which four are articulated in two pairs and one is an isolated

distal caudal vertebra. The body of the thoracic vertebra is anteriorly eroded, as well as the

anterior zygapophyses. Posteriorly, the vertebral body has an outline similar to that of other

pampatheriids. The vertebral body is proportionally higher than in *Tonnicintus mirus*.

Notably, two ventrolateral apophyses are projected from the vertebral body, like some

thoracic vertebrae in *T. mirus* (Góis et al. 2015). Although fragmented, the neural spine of

the same vertebra is inferred as proportionally shorter than in *H. floridanus* and *T. mirus*.

On other hand, the anterior caudal vertebrae have a posteriorly oriented and tall neural

spine. The transverse processes are relatively little extended to the sides.

Astragalus: In Edmund & Theodor (1997) was mentioned the existence of numerous

undetermined postcranial elements whose description was postponed to include it in other

publication. However, that description was never published. This postcranial material

comprises, within the specimens recovered from the UCMP collections, a left astragalus

(UCMP 39846; Fig. 7A–D). In dorsal view, it is observed a lateral trochlea considerably

larger than the medial trochlea. The astragalar head is bulging, spherical, almost uniformly

convex. There is a shallow concavity in the dorsal margin of the astragalar head whose

function has been not determined, but it could be for tendinous insertion and attachment.

The astragalar neck is well-differentiated, similar to *Holmesina* and in contrast with that

observed in *Pampatherium*. In ventral view, the facets of articulation with the calcaneum,

i.e. ectal and sustentacular, are widely separated between them as one would be expect

from the observations on their counterparts in the calcaneum (Edmund 1987). According to

this, the ectal facet is noticeably larger than the sustentacular facet, unlike *H. floridanus*.

The ectal facet is kidney-shaped and the sustentacular facet has a sub-oval shape. Both of

them are concave, especially the ectal facet which is very deep. The sustentacular facet is

located in a central position within the astragalar neck.

Calcaneum: This foot bone is other postcranial element not described by Edmund and

Theodor for *S. hondaensis*. A well-preserved left calcaneum is associated with the catalog

number UCMP 39846 (Fig. 7E–F). This specimen has a proximo-distal length of 54.12 mm

and a width at the level of facets (ectal and sustentacular) of ~10.2 mm. These values are

the smallest for calcanei referred to Pampatheriidae. The only one species whose known

calcaneum is comparable in size to that of *S. hondaensis* is *H. floridanus*. The calcaneum of

the latter species is slightly more proximo-distally elongated than in *S. hondaensis*, but it is

around twice wider at the level of the ectal and sustentacular facets. This means that the

calcaneum of *H. floridanus* is more robust than that of *S. hondaensis*. The calcaneal head is

anteroposteriorly elongated, like in *H. floridanus* and unlike the proportionally short

calcaneal head of *H. septentrionalis*. The anterior end of the calcaneal head is less truncated

that in *Holmesina*. The calcaneum of *S. hondaensis* shows there is no contact between the

borders of the ectal and sustentacular facets, similar to species of *Holmesina* other than *H.*

floridanus and *Pampatherium* (Góis 2013). These facets are slightly convex and they are

separated by a moderately deep and very width groove, i.e. the *sulcus tali* (see below). Like

H. floridanus, the same facets are highly asymmetrical but in *S. hondaensis* this condition is

even extreme as the ectal facet is much larger than the sustentacular facet. Additionally, the

shape of these facets is disparate between them, i.e. kidney-shaped ectal facet and sub-oval

sustentacular facet. The ectal facet is located at an oblique angle with respect to the long

axis of the tuber calcanei, unlike *H. floridanus* and similar to *Dasypus*. Like other

pampatheriids, the sustentacular facet of the calcaneum of *S. hondaensis* is located

anteriorly to the anterior border of the ectal facet. However, this facet is even more

anteriorly placed than in other species as consequence of the wide *sulcus tali* that separates

the ectal facet with respect to the sustentacular one. Posteriorly, the calcaneal tuber is not

massive in comparison with late diverging species of *Holmesina* (e.g. *H. septentrionalis*

and *H. paulacoutoi*), but rather it is slender because a mediolateral compression,

particularly towards its dorsal side.

Stratigraphic and geographical distribution: Possibly the Castilletes Formation, upper

Lower to lower Middle Miocene, upper Burdigalian to Langhian, Department of La

Guajira, Colombia (Fig. 9C); La Victoria and Villavieja Formation, upper Middle Miocene,

Serravalian, Department of Huila, Colombia (Edmund & Theodor 1997).

Scirrotherium antelucanus Laurito & Valerio, 2013

LSID. urn:lsid:zoobank.org:act:225CD304-3B63-4B55-B8B8-33B46C90A194

Holotype: CFM-2867, mobile osteoderm.

Type locality and horizon: San Gerardo de Limoncito, Coto Brus county, Puntarenas

Province, Costa Rica. Upper Curré Formation, Upper Miocene. For further information

about the stratigraphic position of the Curré Formation, see these references: Lowery 1982;

Yuan 1984; Rivier 1985; Kolarsky et al. 1995; Alvarado et al. 2009; Aguilar et al. 2010;

Obando 2011. There are no absolute ages for this geological unit.

Referred material: MUN STRI 36880, an isolated fixed osteoderm (Fig. 8).

Stratigraphic and geographical provenance: Upper Sincelejo Formation, Upper Miocene

to Pliocene (Messinian to Zanclean). El Coley Town, Municipality of Los Palmitos,

Department of Sucre, Colombia. For further information about the stratigraphic position of

the Sincelejo Formation, see these references: Flinch 2003; Villarroel & Clavijo 2005;

Bermúdez et al. 2009; and Alfaro & Holz 2014. There are no absolute ages for this

geological unit.

Diagnosis: Unmodified (see Laurito & Valerio, 2013; p. 47).

Comparative description: The fixed osteoderm of the pelvic shield MUN STRI 36880 is

assigned to the species *S. antelucanus* on the basis of the following observations: (I) the

area and thickness of this osteoderm (linear measurements in millimetres: anteroposterior

length = 34.91; transverse width = 24; thickness = 4.45; approximate area = 837.8 mm^2) are

within the range of variability for comparable osteoderms of *S. antelucanus* and exceed the

known values of area for most of the same kind of osteoderms for *S. hondaensis*; (II) the

external surface is relatively smooth; (III) the AM is wide; (IV) the anterior foramina are

larger (2–3 millimetres of diameter) than in *S. hondaensis*, like *S. antelucanus* from Costa

Rica; (V) the number of anterior foramina (9) is within the range of variability for *S.*

antelucanus (7–10 for quadrangular osteoderms, as the specimen here described), greater

than in *S. hondaensis*; (VI) poorly elevated LCE, even superficial, like in some osteoderms

of *S. antelucanus* (generally LCE more elevated in *S. hondaensis*; see Laurito & Valerio

2013).

Stratigraphic and geographical distribution: Curré Formation, Upper Miocene, Puntarenas

Province, Costa Rica (Laurito & Valerio 2013); Sincelejo Formation, Upper Miocene to

Lower Pliocene, Messinian to Zanclean, Department of Sucre, Colombia.

aff. *Scirrotherium*

Referred material: MUN STRI 16718 (Fig. 9A), fixed osteoderm of the scapular shield;

MUN STRI 38064 (Fig. 9E), undetermined fixed osteoderm; MUN STRI 16719 (Fig. 9G),

mobile osteoderm fragmented in its AM.

Stratigraphic and geographical provenance: Castilletes Formation, upper Lower Miocene

to lower Middle Miocene, upper Burdigalian to Langhian). Localities of Makaraipao,

Kaitamana and Patajau Valley (localities with numbers 390093, 430202 and 390094 in

Moreno et al. 2015, respectively), Municipality of Uribia, Department of La Guajira,

Colombia.

Comparative description: The fixed osteoderm of the scapular shield MUN STRI 16718

(Fig. 9A) is relatively large and has a pentagonal outline. Its linear measurements in

millimetres are: anteroposterior length = 45.02; transverse width = 33.41; thickness = 6.66.

These values imply that this osteoderm has greater area than any other known area size for

osteoderms referred to *Scirrotherium* (Table 1), including those of the osteoderms of the

larger *Scirrotherium* species, i.e. *S. antelucanus* (see Appendix 1 in Laurito & Valerio

2013). Rather, this osteoderm size is similar to those reported for *H. floridanus* and *P.*

humboldtii. The external surface of the osteoderm MUN STRI 16718 is punctuated by

numerous diminutive pits, like *S. hondaensis* and *S. antelucanus*. In this surface it is not

possible to differentiate a LCE nor LDs, in such a way the osteoderm has a flattened

appearance, similar to the case of several osteoderms of *S. antelucanus* (Laurito & Valerio

2013). In contrast, the MEs are clearly recognizable. These ridges are relatively low and

narrow. There are foramina with a nearly homogeneous large size in the AM. They are

aligned in two well defined rows. The most anterior row has five foramina and the posterior

to this one has six. Collectively, the two foramina rows are equivalent to ~25% of the

anteroposterior length of the osteoderm. In *Scirrotherium*, the foramina rows of fixed

osteoderm, when present, this percentage is less than 20%.

MUN STRI 38064 (Fig. 9E): This osteoderm has a trapezoidal outline and the following

measurements in millimetres: anteroposterior length = 39.08; transverse width = 39.55;

thickness: 5.98. These values are within the range of variability of *S. antelucanus*. This

osteoderm has two long rows of anterior foramina of which the posterior row seems to

extend partially over the anterior LMs, unlike the anterior foramina row(s) in *S. hondaensis*

and *S. antelucanus*. The most anterior foramina row is formed by eight foramina and the

posterior to this one has 11 foramina. In both of these rows, foramina have similar size

between them, although a few ones are comparatively tiny. The foramina rows diverge on

the left LM and within the resultant space is located a large and isolated foramen, i.e. not

clearly aligned with any row. This osteoderm does not have recognizable LCE nor LDs, i.e.

it is flattened. Its MEs are narrow and poorly elevated. The foramina of LMs are smaller

than most of anterior foramina. As consequence of preservation factors, expected pits on

the external surface are not present.

MUN STRI 16720 (Fig. 9G): This partial mobile osteoderm has an elongated rectangular

shape. Its linear measurements in millimetres are: anteroposterior length (incomplete by

fragmentation) = 45.68; transverse width = 30.69; thickness = 6.96. The external surface is

convex and without LCE nor LDs. The AM shows an apparently unordered foramina set.

Discussion: With current evidence, the former material should not be confidently assigned

to *Scirrotherium* and even less so to create a monospecific genus. This taxonomical

decision is supported by several arguments. First, morphologically, the osteoderms here

referred to aff. *Scirrotherium* are more similar to those of *S. hondaensis* and *S. antelucanus*

than to any other osteoderms of known pampatheriids. The osteoderms of aff.

Scirrotherium fundamentally differ with respect to the osteoderms of *S. hondaensis* and *S.*

antelucanus in three characteristics: (I) greater number of anterior foramina and/or greater

development of two rows from these foramina; (II) LCE possibly absent, i.e. flattened

external surface; (III) larger maximum osteoderm area. Of these features, the third one (III)

is the less ambiguous, i.e. the maximum area of fixed osteoderms exceeds those of the

osteoderms of *S. hondaensis* and *S. antelucanus*. Comparatively, the first and second (I and

II) characteristics are more ambiguous considering that similar conditions were also

observed in *S. hondaensis* and *S. antelucanus*. These conditions are described as follows.

Some infrequent osteoderms of *S. hondaensis* have two anterior foramina rows, of which

the most anterior row is comparatively less developed (i.e. less foramina and smaller) than

in aff. *Scirrotherium*. Additionally, in *S. hondaensis* and, particularly in *S. antelucanus*,

some osteoderms have flattened and diffuse LCE, even apparently absent. These

observations imply limitations on the taxonomical resolution, especially considering that

the material on which is based aff. *Scirrotherium* is scarce and it does not allow

comparisons from a representative osteoderm series of the morphological variability within

the carapace of this animal.

Genus *Kraglievichia* Castellanos, 1927

LSDI. urn:lsid:zoobank.org:act:92C8B169-4F79-467E-B951-EF1DE6E327B1

Type species: *Kraglievichia paranensis* Ameghino, 1883

Included species: In addition to the type species, *Kraglievichia carinatum* comb. nov.

Emended differential diagnosis: Small-to-middle sized pampatheriid characterized by fixed

osteoderms with ornamentation (particularly the LCE) more conspicuous than in any other

pampatheriid; anteriorly wide posteriorly tapered LCE; very deep LDs; highly elevated and

frequently blunt MEs, even flattened towards their top; external surface of osteoderms

generally rougher than in *Scirrotherium* but less than in *Holmesina*.

Kraglievichia carinatum comb. nov.

Synonyms: *Scirrotherium carinatum* (Góis et al. 2013).

Holotype: MLP 69-IX-8-13-AB, a mobile osteoderm.

Type locality and horizon: Paraná River cliffs, Entre Ríos Province, Argentina. Ituzaingó

Formation, Upper Miocene, Tortonian.

Referred material: In addition to the holotype, the paratypes and part of the hypodigm of

this species (see Fig. 10 and Appendix S1 of the Supplementary Material).

Differential diagnosis: Unmodified (see Góis et al. 2013, p. 182).

Discussion: In their descriptive work on *K. carinatum* comb. nov., Góis et al. (2013) did

not explicitly justify the inclusion of this species within *Scirrotherium*. Interestingly, part of

the material assigned to the taxon they create, coming from northwestern Brazil (Solimões

Formation), was previously referred to *Kraglievichia* sp. by several researchers, including

Góis himself (Góis et al. 2004; Góis 2005; Cozzuol 2006; Latrubesse et al. 2010; Góis et al.

2013). However, Góis et al. (2013) refuted the original taxonomical assignment arguing

this was erroneous, although they did not offer any concrete support for their decision. In

absence of a phylogenetic analysis in Góis et al. (2013), we could assume by default that

these authors included to *K. carinatum* comb. nov. within *Scirrotherium* because they

considered osteodermal features of this species are at least compatible with the generic

diagnosis proposed by Edmund & Theodor (1997). Additionally, based on morphological

similarity, Góis and colleagues could have hypothesized closer affinities between *K.*

carinatum comb. nov. and *S. hondaensis* than those between *K. carinatum* comb. nov. and

K. paranensis.

Let do us analyse in detail each of the osteodermal features of *K. carinatum* comb. nov. in

relation to the original diagnosis of *Scirrotherium*. According to Edmund & Theodor

(1997), the fixed osteoderms of *Scirrotherium* have a small (not specified) number of large

piliferous foramina on the AM. These foramina are well spaced and interconnected

between them by a distinct channel. This is observed both in *K. carinatum* comb. nov. and

S. hondaensis. Likewise, the presence of continuous MEs, posteriorly confluent with the

LCE, is a trait also shared by the compared species. Finally, the relative osteoderm size of

K. carinatum comb. nov. is small among pampatheriids, which is in line with the original

diagnosis for *Scirrotherium.*

In consequence, the osteodermal features of *K. carinatum* comb. nov. are compatible with

those mentioned in the diagnosis for *Scirrotherium* by Edmund & Theodor (1997).

However, this does not necessarily imply that the taxonomical allocation of *K. carinatum*

comb. nov. in the genus *Scirrotherium* is correct. In fact, there are several reasons to

consider this is not reliable. Initially, the diagnosis of Edmund & Theodor (1997) contains

only three allegedly diagnostic features on osteoderms, including the relative osteodermal

size. Furthermore, and more importantly, these "diagnostic features" do not allow to

discriminate between *Scirrotherium* and any other genus of pampatheriids. Indeed, in their

analysis, Góis et al. (2013) accept that, for instance, the species *Vassallia minuta* also

shares the condition about fixed osteoderms with a small number of large anterior foramina,

which are well spaced and connected through a distinguishable canal. Independently,

Laurito & Valerio (2013) also highlighted the non-diagnostic nature for *Scirrotherium* of

the former osteodermal trait. The other osteodermal features under consideration, i.e. the

posterior confluence of the MEs with the LCE and the small osteodermal size, are also

ambiguous for positive identification of *Scirrotherium*. For instance, the confluence of MEs

and LCE is also found in *K. paranensis*, a pampatheriid clearly different from

Scirrotherium. And although apparently informative on body-size trends of some individual

pampatheriid lineages (e.g. *Holmesina* spp.) and useful as discriminant factor between

species (Góis et al. 2013; Laurito & Valerio 2013), the relative osteodermal size is not

necessarily insightful in itself to make taxonomical decisions on grouping species in one

single genus (see below). In this sense, it is noteworthy the potentially conflictive

taxonomical conclusions based predominantly on osteoderm-inferred relative body size in

relation with interpretations from non-osteodermal evidence. Thus, for example, the femur

MLP 69-IX-8-13A (an adult specimen), which comes from the Ituzaingó Formation and

was assigned to *K.* cf. *paranensis* by Scillato-Yané et al. (2013), is comparable in size to

that of *S. hondaensis*, i.e. a small pampatheriid. Then, keeping in mind the medium-to-large

body size of *K. paranensis*, is probable that the referred femur does not belong to the

species *K. paranensis*, although it is reasonable to include it in the genus *Kraglievichia* (as

the authors decided). However, Scillato-Yané et al. (2013) did not discuss the possibility

that the material assigned to *K.* cf. *paranensis*, particularly the specimen MLP 69-IX-8-

13A, has any relationship with the (partially) sympatric species of *K. paranensis*, i.e. *K.*

carinatum comb. nov., a pampatheriid whose small body size is fully compatible with the

small size of that femur. In other words, like Góis et al. (2013), they did not seriously

consider the hypothesis of *K. carinatum* comb. nov. as a small species of *Kraglievichia*,

rather than a species belonging to *Scirrotherium*.

Again analysing the original diagnosis for *Scirrotherium* by Edmund & Theodor (1997), it

should be regarded as ambiguous and hardly useful to differentiate this genus from other

genera in Pampatheriidae, at least with respect to osteodermal traits. Probably these

supposedly diagnostic features are actually symplesiomorphies for genera of the family or,

at most, a hypothetical subfamilial lineage. This means that Góis et al. (2013) did not have

a minimally strong taxonomical background from the original diagnosis of *Scirrotherium* to

assign *K. carinatum* comb. nov. to this genus. Alternatively, they may have noticed

morphological similarity between osteoderms of *K. carinatum* comb. nov. and *S.*

hondaensis from features not included in the diagnosis by Edmund and Theodor.

Nevertheless, on their publication, these authors only listed multiple morphological

differences between the former species and virtually did not mention any similarity for

them, except for potentially equivocal resemblance as that indicated by relative osteodermal

size (i.e. small osteodermal sizes in comparison with those of *K. paranensis* and *Plaina*).

The lack of usefulness of the relative osteodermal size for generic assignation is further

supported by the osteodermal morphometric analysis in Góis et al. (2013, p. 185), whose

resulting PCA and CCA plots show that, despite the similarity in relative osteodermal size,

K. carinatum comb. nov. is located far from *S. hondaensis* (which is closer to *V. minuta*, a

taxon apparently related to other main lineage within Pampatheriidae, i.e. *Plaina-*

Pampatherium) and *K. paranensis* in morphospace.

Summarizing, there is no solid, possible justification by Góis et al. (2013) on their

taxonomical decision to including *K. carinatum* comb. nov. within *Scirrotherium*. The

observation of a common general morphological pattern between *K. carinatum* comb. nov.

and *S. hondaensis* does not necessarily imply the grouping of these species under the same

generic taxon. Furthermore, we should note that Góis and colleagues, in their work on *K.*

carinatum comb. nov., did not make morphological comparisons including to *S.*

antelucanus, a species more similar in osteodermal features to *S. hondaensis* (i.e. the type

species of *Scirrotherium*). The species *S. antelucanus* was described on a scientific article

(Laurito & Valerio 2013) published nearly simultaneously, but later, to that of *K. carinatum*

comb. nov. Therefore, Góis et al. (2013) did not know about the existence of *S. antelucanus*

when they performed their systematic analysis ("Until the present study, *S. hondaensis* was

the only known species of this genus"; Góis et al. 2013, p. 177), so that their taxonomical

assignment of *K. carinatum* comb. nov. to *Scirrotherium* was biased by limited notions on

the morphological variability and diversity of *Scirrotherium* in northern South America and

southern Central America.

In this work I decide to assign *K. carinatum* comb. nov. to the genus *Kraglievichia* based

on results of a phylogenetic analysis that I designed considering the hypothesis of Edmund

(1985) on the probable subfamilial relationships within Pampatheriidae, which implicitly

sustains that the creation of supraspecific taxa from osteodermal evidence should be

determined –with the prerequisite of morphological similarity- by the degree of

development of the ornamentation. Understanding that *K. carinatum* comb. nov. has

morphologically similar osteoderms to those of *K. paranensis* (apart from relative

osteoderm size) and has one of the more conspicuous, protuberant osteodermal

ornamentations among pampatheriids, along with *K. paranensis*, as acknowledged by Góis

et al. (2013) themselves, this means that *K. carinatum* comb. should be considered closely

related to *K. paranensis* and therefore they both should also be included in the same genus,

i.e. *Kraglievichia.*

Stratigraphic and geographical distribution: Ituzaingó Formation, Upper Miocene,

Tortonian, Entre Ríos Province, Argentina; Puerto Madryn Formation, Upper Miocene,

Tortonian, Chubut Province, Argentina; Solimões Formation, Upper Miocene, Tortonian,

Acre State, Brazil (Góis et al. 2013).

Discussion

This systematic analysis is the first attempt to test the intergeneric relationships and internal

structure of the genus *Scirrotherium* with its three previously referred species, i.e. *S.*

hondaensis (type species), *'S.' carinatum* (= *K. carinatum* comb. nov.) and *S. antelucanus*.

The two strict consensus trees from the distinct character weighting schemes (equal and

implied weights) show very similar results. However, the preferred general phylogenetic

hypothesis is that supported by the implied weights analysis. According to Goloboff et al.

(2018), the parsimony analysis under implied weights outperforms equal weighting and the

model-based methods. This performance inference for the implied weights method is

empirically supported here. Beyond this preference for a particular hypothesis (further

supported below), both resultant trees agree that all the species referred to *Scirrotherium*

are not monophyletic and, consistently, from a diagnostic point of view only *S. hondaensis*

and *S. antelucanus* appears as those actually referable to *Scirrotherium*. The relationships

between *S. hondaensis* and *S. antelucanus* could not be confidently resolved, despite the

inclusion of new osteodermal characters (the only ones comparable between these species

so far) in these parsimony analyses. According to the majority-rule consensus tree obtained

from the bootstrap resampling, *S. antelucanus* is not the sister taxon of *S. hondaensis*, but it

is basal to the clade *S. hondaensis* + '*S*' *carinatum* (= *K. carinatum* comb. nov) + *K.*

paranensis. However, this clade has low resampling frequency. The basal position of *S.*

antelucanus within the majority-rule consensus tree is explained to a great extent by a large

number of missing entries for *S. antelucanus* in the character matrix. Nevertheless, a non-

monophyletic relationship among *S. hondaensis* and *S. antelucanus* should not be rule out.

In conjunction, these results suggest the need of information on craniomandibular or dental

characters for *S. antelucanus* to test its relationship with respect to *S. hondaensis*. Until new

phylogenetic evidence, the genus *Scirrotherium* is maintained as taxonomically valid using

a new, emended diagnosis which focus on its lesser degree of development of the

osteodermal ornamentation in comparison with those in *Holmesina* and, particularly,

Kraglievichia. This new diagnosis replaces the original and now inadequate diagnosis of

Edmund & Theodor (1997). Composed of *S. hondaensis* and *S. antelucanus*, *Scirrotherium*

is considered a coherent taxonomical generic unit according to the criteria of morphological

homogeneity and similarity in geographical and chronostratigraphic distributions of its

species.

Unlike the unpublished phylogeny of Góis (2013), the phylogenetic position of '*S.*'

carinatum was resolved here, i.e. this species is the sister taxon of *K. paranensis*.

Therefore, it is proposed the new name *K. carinatum* comb. nov. Despite Góis (2013) did

not recover as a clade to *S. hondaensis* and *K. carinatum* comb. nov., as expected if both

these species were assigned to *Scirrotherium*, he presented one supposed synapomorphy

that join them, i.e. very deep LDs, "in particular in *S. carinatum*" (Góis 2013, p. 215). This

feature is not a confident support for grouping *S. hondaensis* and *K. carinatum* comb. nov.

because the deepest LDs in this main pampatheriid lineage are found in *K. carinatum* comb.

nov. and *K. paranensis*, not in *S. hondaensis*. The new emended differential diagnosis for

Kraglievichia acknowledges the highly sculpted external osteoderm surface documented on

this taxon, which is more protuberant than in the Plio-Pleistocene genus *Holmesina*. This

diagnosis provides an updated and concise description of useful osteoderm features to

distinguish *Kraglievichia* from other genera within Pampatheriidae. It is important to note

that the species *K. paranensis* has several autapomorphies (see Appendix S4 of the

Supplementary Material) which, in addition to the relative osteoderm size, need to be

compared in the future with homologous, unknown (cranial) traits in *K. carinatum* comb.

nov. in order to test the phylogenetic closeness of these two species as inferred from

osteodermal traits. Provisionally, the difference in relative osteodermal size (consequently

also in relative body size) and some morphological differences between *K. carinatum*

comb. nov. and *K. paranensis*, as those noted by Góis et al. (2013), may be linked to

distinct ontogenetic growth trajectories in these species, being plesiomorphic in nature that

of *K. carinatum* comb. nov. (i.e. small body size; see Sánchez-Villagra 2012 for a

discussion on the implications for taxonomy of the ontogenetic growth in extinct species).

This hypothesis could be extrapolated in some way to the species *S. hondaensis* and *S.*

antelucanus.

An important difference here with respect to the phylogeny of Góis is that *K. paranensis*

was not recovered as the one single sister taxon of *Holmesina* spp. (except *H. floridanus*).

Instead, it is part of a group additionally formed by *S. hondaensis*, *S. antelucanus* and *K.*

carinatum comb. nov. Together, these taxa are possibly the sister clade of *Holmesina*. This

means that we do not have evidence of direct ancestral forms to *Holmesina* yet. However,

despite some striking differences, it is remarkable the greater morphological similarity

between the osteodermal ornamentation and cranial features of *Holmesina* (especially *H.*

floridanus) and *Scirrotherium*, rather than with those of *Kraglievichia*. The recognizable

similarities between *H. floridanus* and *S. hondaensis*, and at the same time differences with

K. paranensis, include a less protuberant osteodermal ornamentation; occurrence of

uniformly narrower LCE in some osteoderms; more robust skull; more expanded frontals;

frontals in a clearly different plane with respect to the parietals; less anteroposteriorly

elongated upper teeth; first two upper molariforms less obliquely oriented with respect to

the midline of the hard palate; among others.

Apart from that, Góis (2013) found that *Holmesina* is non-monophyletic by the

phylogenetic position of *H. floridanus* with respect to the clade groups the other *Holmesina*

species. This result coincides with that recovered here from the parsimony analysis with

equal weights. The species *H. floridanus* has several plesiomorphic features in comparison

with the remaining *Holmesina* species, e.g. less protuberant ornamentation, less rough

external surface of the osteoderms and less dorsal basicranium. However, the arrangement

of the calcaneal facets of the astragalus in *H. floridanus* (confluent facets) suggests that this

species is not directly related to any known South American pampatheriid (Edmund 1987).

Likewise, Gaudin & Lyon (2017) have recently found support for the monophyly of

Holmesina from craniomandibular specimens. Therefore, the position of *H. floridanus* in

polytomy with *S. hondaensis* and *S. antelucanus* within the topology of the resultant tree

with equal weights is explained from the lack of enough resolution of the analysis

(homoplastic noise), not as product of a "real" distant relationship with the remaining

Holmesina species. Conversely, the genus *Holmesina* is recovered as monophyletic by the

implied weights analysis. As expected, in this topology *H. floridanus* is the most basal

species of *Holmesina*. In both strict consensus trees, *H. septentrionalis*, the later diverging

North American species, is grouped together with all the South American *Holmesina*

species (except *H. rondoniensis*, not included in this study).

I abstained from revise the diagnosis of the genus *Holmesina* because this is considered

beyond the intended objectives of this work. However, six putative synapomorphies

(unambiguous and ambiguous) are proposed (or further supported) for this genus: (1)

Anterior and lateral margins with elongated, strong bone projections as radii directed from

the external border of the central figure towards the osteodermal borders in non-marginal

fixed osteoderms; (2) anteriorly convergent, nearly in contact medial processes of

premaxillae; (3) length of nasals greater than 30% of the maximum anteroposterior length

of the skull; (4) conspicuous and anteroposteriorly elongated maxillary ridge (Gaudin &

Lyon 2017); (5) reduced lacrimal; and (6) bilobed posterior upper molariforms (Mf5-Mf9)

with incipient trilobulation.

Evolutionary and biogeographical implications

Scirrotherium is a pampatheriid genus from the Early Miocene-Late Pliocene of northern

South America and southern Central America. This taxon, along with *Kraglievichia*,

conforms the sister evolutionary line of that gave origin to *Holmesina* (Fig. 12), a

pampatheriid probably originated in tropical southern North America (Mexico? see

Woodburne 2010). However, based on the osteological comparisons presented above,

which are expanded with respect to those of Edmund (1987), the hypothetical South

American ancestor or sister taxon of *Holmesina* probably was morphologically generalized,

more similar to *Scirrotherium* or aff. *Scirrotherium* than to *Kraglievichia*. This

interpretation is in line with that of Edmund (1987), according to which the calcaneo-

astragalar articulation of *S. hondaensis* precludes that this pampatheriid is ancestral to *H.*

floridanus, but the ornamentation pattern of the osteoderms suggests "at least some degree

of relationship" with the latter species. The genus *Kraglievichia* should be considered a

highly, anatomically divergent taxon, especially to taking into account its Miocene age. In

Edmund's words, "the osteoderms [of *H. floridanus*] are quite dissimilar to those of

Kraglievichia" (Edmund 1987, p. 16). This interpretation is partially in conflict with that of

Scillato-Yané et al. (2005), according to which *Holmesina* originated from a hypothetical

(i.e. not recorded yet) South American basal form of *Holmesina* or *Kraglievichia*. It also is

opposed to Simpson (1930) and to the phylogeny of Góis (2013) in which *Kraglievichia* is

the sister taxon of *Holmesina*.

The earliest record of *Scirrotherium*, here treated as tentative by scarce and bad-preserved

material, comes from the Early Miocene (late Burdigalian) of northwestern Venezuela

(Rincón et al. 2014; see below). Independently from the validity of occurrence of

Scirrotherium in an Early Miocene locality of northern South America, this record

represents the oldest pampatheriid reported in the scientific literature so far. This

improvement in the fossil record of Pampatheriidae makes it more congruent with the

expected time of origination of this family, i.e. Late Oligocene-Early Miocene, according to

the very few available time-calibrated phylogenies including representatives of

Pampatheriidae and its sister group, i.e. Glyptodontidae (e.g. Fernícola 2008; Billet et al.

2011). Apparently, the Early Miocene Venezuelan pampatheriid indicates an origin of these

xenartrans in low latitudes in South America. However, this hypothesis could be challenged

by a possible Late Eocene pampatheriid of Argentina, which has been not formally

described and published yet (Góis 2013).

Beyond the geographical origin of Pampatheriidae, northern South America seems to have

been a critical area for the early diversification of, at least, a main lineage of pampatheriids,

i.e. that including *Scirrotherium*. This genus probably differentiated at least as early as the

late Early Miocene-early Middle Miocene (late Burdigalian-Langhian) in northernmost

South America. The former evolutionary inference is consistent with the late Early

Miocene record referred to *Scirrotherium* from Venezuela (Rincón et al. 2014).

Collectively, *Scirrotherium* and *Kraglievichia* occupied a large area in South America

during the Neogene (Fig. 13). The geographical range of *Scirrotherium* was more restricted

than that of *Kraglievichia*, comprising only tropical low latitudes, instead of a wide

latitudinal range, as suggested by Góis et al. (2013). The revaluated distributional pattern of

Scirrotherium is comparable to that of the glyptodontid *Boreostemma*, which is recorded

from the Middle Miocene to the Late Pliocene of Colombia and Venezuela (Carlini et al.

2008; Zurita et al. 2016). In contrast, the distributional range of *Kraglievichia* is similar to

that of other Miocene xenartran taxa at the generic and specific level which occurred in

southern South America and northwestern Brazil, but not in the northern or northwestern

end of South America (see Ribeiro et al. 2014).

Overall, this evidence indicating biogeographical divergence of northwesternmost South

America as an independent faunal province, from the late Early Miocene to Middle

Miocene and possibly until the Late Miocene to Pliocene, is consistent with the results of

multiple analyses of the South American terrestrial mammal fossil record for the Neogene

(Patterson & Pascual 1968; Cozzuol 2006; Ortiz-Jaureguizar & Cladera 2006; Croft 2007;

Carrillo et al. 2015; Rincón et al. 2016; Kerber et al. 2017; Brandoni et al. 2019).

Apparently, the existence of one or several strong geographical and/or ecoclimatic barriers

(e.g. the Pebas Mega-Wetland System, whose expansion climax coincides with the Middle

Miocene) in northern South America would explain that regional endemism pattern

(MacFadden 2006; Croft 2007; Salas-Gismondi et al. 2015; Jaramillo et al. 2017). At the

same time, the development of a late Early-to-Middle Miocene biogeographical divergence

between northwesternmost South America and the rest of this continent may account for

the evolutionary (morphological) divergence of the genera *Scirrotherium* and

Kraglievichia.

In the Late Miocene, without a completely formed PLB (O'dea et al. 2016), *Scirrotherium*

expanded its geographical range to southern Central America (Fig. 13), suggesting a

hypothetical ephemeral land connection (i.e. land span) or, more likely, overwater dispersal

between South America and Central America (maybe via rafting mechanism; efficient

active swimming of a pampatheriid in a marine channel seems highly improbable). This is

the earliest dispersal event of a pampatheriid to North America (see below). The Central

American species of *Scirrotherium*, *S. antelucanus*, is larger than *S. hondaensis*, but

comparable or even smaller than aff. *Scirrotherium*. From available evidence, it is not

possible to determinate the most probable area of evolutionary differentiation of *S.*

antelucanus, but now there is support for occurrence of this species in the late Neogene of

northwestern South America, specifically in the Department of Sucre, Colombia.

The South American record of *S. antelucanus* is probably several million years younger (3–

5 my) than the Central American record. However, given the lack of absolute dating for the

fossil-bearing stratigraphic levels and the occurrence of Late Miocene strata in the same

geological unit (Sincelejo Formation) where comes the material here assigned to *S.*

antelucanus in Colombia, it should be recognized a significant age uncertainty for the new

South American record of this species. In any case, this age is considered may be Early

Pliocene or, alternatively, Latest Miocene from the stratigraphic position of the fossil-

bearing horizons (Villarroel & Clavijo, 2005; Bermúdez et al. 2009; Alfaro & Holz 2014;

Bernal-Olaya et al. 2015; Córtes et al. 2018), as well as from associated palynomorphs

(Silva et al. 2012; B. Fernandes & C. Jaramillo, pers. comm. 2014).

The biogeographical correlation across the Isthmus of Panama using *S. antelucanus* has

insightful implications for the understanding of the late Cenozoic intercontinental migratory

dynamics in the Americas, including the Great American Biotic Interchange (GABI) (Webb

2006; Woodburne et al. 2006; Woodburne 2010; Cione et al. 2015; Bloch et al. 2016).

Noteworthy, this is the first transisthmian biogeographical correlation for a Neogene

terrestrial mammal at the level of species; furthermore, it is the first short-distance

intercontinental correlation (i.e. adjacent to the Central American Seaway) with high

taxonomical resolution for Neogene land mammals of the Americas; and, finally, it

constitutes the first evidence of a distributional pattern congruent with a re-entry event to

South America by a pre-Pleistocene xenartran.

We subsequently analyse in more detail these implications. So far, we know a few

biogeographical correlations across the Isthmus of Panama which are based on records at

generic level of Neogene and Pleistocene land mammals, as well as a very few records at

species level of the latter epoch. The Neogene biogeographical correlations include the

pampatheriid genera *Plaina*, in Mexico and central-southwestern South America;

Pampatherium, in Mexico and southeastern South America; and *Holmesina*, in United

States, Mexico and El Salvador, as well as in northwestern and southeastern South America

(Woodburne 2010). At the level of species, for instance, the Pleistocene megatheriine

Eremotherium laurillardi has occurrence in both sides of the Isthmus of Panama in North-

and South America (Cartelle & De Iuliis 1995, 2006; Tito 2008; McDonald & Lundelius,

E. L. Jr. 2009; Martinelli et al. 2012; Cartelle et al. 2015).

The record in South America of *S. antelucanus* increase the taxonomical resolution of

transisthmian biogeographical correlations of Neogene land mammals, opening the

possibility of new correlations of this kind and their biostratigraphic application in circum-

Caribbean basins, in a similar way as envisioned by the renowned American

palaeontologist Ruben A. Stirton from his revision of the fossil mammal remains of "La

Peñata fauna" (Stirton 1953), the vertebrate fossil association where comes the new record

of *S. antelucanus*. This translates into direct correlation of Land Mammal Ages (in this

case, SALMA and NALMA) from migrant mammals which are shared at species level by

both South and North America. Using to *S. antelucanus*, this would mean exists a support

for faunal, not necessarily chronological, correlation of the early Hemphillian and

Montehermosan mammal (xenartran) assemblages in North America and South America,

respectively (see Laurito & Valerio, 2013). Nevertheless, naturally, any solid

intercontinental faunal correlation implies more than only one taxonomical element for

support. The direct intercontinental faunal correlations from Cenozoic land mammals

between South and North America are still underdeveloped in comparison with those

between other continents (e.g. North America and Europe or North America and Asia;

Woodburne & Swisher, C. C. III 1995; Beard & Dawson 1999; Bowen et al. 2002)

Additionally, the transisthmian correlation of *S. antelucanus* allows to increase the

geographical resolution in detection of intercontinental migrations of late Cenozoic land

mammals, which are restricted mainly to large and middle distance correlations for the

Neogene record (e.g. Mexico-southern South America; Woodburne 2010). This pattern has

prevented the exploration of possible early or intermediate phases of

anagenetic/cladogenetic events in late Cenozoic Interamerican migrant taxa, which in turn

it is reflected in the fact that we are detecting "suddenly" well-differentiated terminal taxa

(e.g. *Holmesina*) in marginal, distant areas with respect to the Central American Seaway

and adjacent terrains (Cione et al. 2015 and references therein).

On other hand, the new transisthmian correlation here presented suggests a possible

Neogene re-entry event by a xenartran to South America after its evolutionary

differentiation in North America (Fig. 13). The confirmation of this depends on a confident

determination of the differentiation area for *S. antelucanus*, i.e. if this species originated in

South America, the new record is explained more parsimoniously by population

maintenance in the ancestral area. Conversely, if this species originated in Central America

from a South American species of *Scirrotherium* as *S. hondaensis*, then we are considering

a re-entry event to South America. However, as mentioned above, it is not possible to

constrain much more than that at this moment. In any case, the possibility of a Neogene re-

entry event to South America by a xenartran is compatible with the fact that we know

several of these events during the Pleistocene. Among these Pleistocene events, there are

several involving xenartrans, including the pampatheriids *Holmesina* and *Pampatherium*,

the glyptodontid *Glyptotherium*, the pachyarmatheriid *Pachyarmatherium*, the dasypodid

Dasypus and the megatheriine *Eremotherium* (Woodburne et al. 2006; Woodburne 2010

and references therein).

Now let's discuss the evolutionary implications of this systematic analysis for the genus

Holmesina and the multiple Interamerican dispersal events of pampatheriids, including that

of *Scirrotherium* (discounting the non-confirmed re-entry event to South America). The

genus *Holmesina* has its oldest record (*Holmesina* sp.) in sedimentary rocks deposited

around the Pliocene-Pleistocene boundary (~2.4 mya) in La Florida, United States

(Edmund 1987; Woodburne 2010 and references there; Gaudin & Lyon 2017). This

northward dispersal event is part of the earliest phase of the GABI (GABI 1), in which

additionally participated other xenartrans as *Dasypus*, *Pachyarmatherium* and

Eremotherium (Woodburne 2010). Typically, *H. floridanus* has been considered the most

basal among the *Holmesina* species (Edmund 1987), as it is supported here. The time-

calibrated phylogeny introduced in this work for Pampatheriidae (Fig. 12) suggests that

exist a long ghost lineage leading to *Holmesina*, from the Early Miocene (Burdigalian) until

the Late Pliocene. The improvement of the fossil record in northern South America, Central

America and Mexico will allow to advance in the recognition of probable direct ancestral

forms for *Holmesina*.

From the above analysis, a probable model of biogeographical evolution of *Holmesina* is as

follows (Fig. 13). A hypothetical pampatheriid close to *Holmesina* or even a hypothetical

Holmesina species basal with respect to *H. floridanus* dispersed to Central America,

Mexico and United States during the Pliocene (Early Pliocene according the time-calibrated

phylogeny). Once it was established the genus *Holmesina* in North America with *H.*

floridanus, the larger species *H. septentrionalis* diverged and differentiated in the Early

Pleistocene of southern United States. Later, *H. septentrionalis* expanded southward to

Mexico and Central America during the Early-Middle Pleistocene (Aguilar & Laurito

2009). In the Middle or early Late Pleistocene, possibly *H. septentrionalis* colonized South

America, where took place an important diversification, which was likely influenced by the

Late Pleistocene climatic changes (Scillato-Yané et al. 2005). This diversification gave

origin to the species *H. occidentalis*, *H. rondoniensis*, *H. major* and to the most robust

pampatheriid, *H.paulacoutoi* (Scillato-Yané et al. 2005).

As inferred from the phylogeny and derived interpretations here presented, the dispersal

events of *S. antelucanus* and *H. floridanus* to North America are independent of each other.

This means that the number of northward intercontinental dispersal events of pampatheriids

during the late Cenozoic actually is at least three, which in chronological order are: (1)

genus *Scirrotherium* (Late Miocene); (2) lineage *Plaina-Pampatherium* (Early Pliocene);

(3) genus *Holmesina* (undetermined Pliocene). From these events, only the latter, based on

the fossil record of *H. floridanus*, is included in the GABI. The remaining two events are

classified as part of the macroevolutionary invasion "wastebasket" called "Pre-GABI"

(literally, 'before the GABI'; Woodburne et al. 2006; Woodburne 2010; also named by

Cione et al. 2015 as "ProtoGABI"). In the lineage *Plaina-Pampatherium*, it was

differentiated one genus, *Pampatherium*, and at least three species (*P. mexicanum*, *P. typum*

and *P. humboldtii*, being the two latter recorded in South America). Meanwhile, the

northward dispersal event of *Scirrotherium* seems to give no origin to any other species

different to *S. antelucanus*. Only a confirmed southward intercontinental dispersal event of

the *Scirrotherium-Kraglievichia-Holmesina* clade has been well-established, i.e. that of

Holmesina to South America in the Middle or early Late Pleistocene (Aguilar & Laurito

2009). This event probably is not part of any of the GABI phases of Woodburne (2010) but

it is chronologically located between the GABI 2 and 3.

As it has been shown, the study of more abundant and complete pampatheriid material

preserved in Neogene geological units of northern South America, in particular, and the

current intertropical region of the Americas, in general, has the potential of provide us more

complex and interesting scenarios on the evolution of this glyptodontoid family and,

specifically, the genera *Scirrotherium* and *Holmesina*.

1041

CONCLUSION

The monophyly of *Scirrotherium* has been tested through parsimony phylogenetic analyses.

This taxon is recovered as paraphyletic if we include all the referred species, i.e. *S.*

hondaensis, *S. antelucanus* and '*S.*' *carinatum*. The latter species is closer to *Kraglievichia*

paranensis and, therefore, here it is proposed the new name *K. carinatum* comb. nov. The

remaining referred species to *Scirrotherium*, *S. hondaensis* and *S. antelucanus*, are designed

in aphyly. The taxonomic validity of *Scirrotherium*, as defined here, is maintained from

diagnostic evidence. The genus *Scirrotherium* is probably the sister taxon of *Kraglievichia*,

and these two genera form the sister clade of *Holmesina*. The genus *Scirrotherium* has

occurrence from the late Early Miocene to Late Pliocene of northwestern South America

(Colombia, Venezuela and northern Peru) and the Late Miocene of southern Central

America (Costa Rica). A geographical origin of Pampatheriidae in northernmost South

America is suggested from the fossil record of *Scirrotherium* and a new time-calibrated

phylogeny. The genus *Scirrotherium* also represents the earliest member of Pampatheriidae

which participated in a dispersal event to North America, specifically to the ancient Central

American peninsula. This dispersal event happened when the PLB was not fully formed

yet. The species *S. antelucanus* lived in Central America and now also in northern

Colombia during the late Neogene. This is the first Interamerican biogeographical

correlation of a Neogene land mammal with high taxonomical resolution, i.e. at the species

level. The record of *S. antelucanus* in both sides of the ancient Central American Seaway is

compatible with a possible re-entry event of this pampatheriid to South America. In

addition, *Scirrotherium* is not probably the South American ancestor of the originally-

endemic North American genus *Holmesina*. In contrast with a previous hypothesis which

argues that *Holmesina* may have evolved from *Kraglievichia*, here it is suggested that there

is no evidence of direct ancestral forms of *Holmesina*, although the unknown South

American ancestor of *Holmesina* may be morphologically more similar to *Scirrotherium*.

1068

Acknowledgements

I am greatly indebted to all curators and managers of the collections visited for revision of

specimens during this work, especially to: Andrés Vanegas (Museo de Historia Natural La

Tatacoa), Aldo Rincón (Museo Mapuka, Universidad del Norte), Jorge Moreno-Bernal

(Museo Mapuka, Universidad del Norte), Carlos De Gracia (Museo Mapuka, Universidad

del Norte), Richard C. Hulbert Jr. (Florida Museum of Natural History), Kenneth

Angielczyk (Field Museum of Natural History), Bill Simpson (Field Museum of Natural

History, Patricia Holroyd (University of California Museum of Paleontology) and Marcelo

Reguero (Museo de La Plata). I am also grateful to Carlos Jaramillo (Smithsonian Tropical

Research Institute) and his field work team from allowing access to fossils they collected in

the Deparment of La Guajira (Colombia). Likewise, special thanks to the professor Antonio

Tovar (University of Sucre) by inspiration to explore the Sincelejo Formation; to my field

assistants during this exploration in the Department of Sucre (Colombia), Oscar Melendrez,

Angel Cruz and Juan Pacheco. I would like to thank professor César Laurito (Instituto

Nacional de Aprendizaje de Costa Rica) and Edwin Cadena (Universidad del Rosario) for

providing photos of osteoderms of the species *S. antelucanus* and one photo of outcrops of

the Castilletes Formation, respectively. This work would not have been possible without

logistic support and encourage by Alba Lara, Carolay Jiménez and Mónica Montalvo. An

anonymous reviewer provided comments that made it possible to improve the manuscript.

Financial assistance for this project was provided by several institutions/agencies:

CONICET (Internal Doctoral Fellowship); Field Museum of Natural History (Science

Visiting Scholarship); Florida Museum of Natural History (International Travel Grant);

University of California Museum of Paleontology (Welles Fund); and The Paleontological

Society (PalSIRP-Sepkoski Grant).

1093

DATA ARCHIVING STATEMENT

Data for this study are avalaible in the Dryad Digital Repository: [Intentionally blank]

The nomenclatural acts contained in this work are registered in ZooBank:

LSID. urn:lsid:zoobank.org:act:313358B5-3B1F-4902-8C2E-BB07CFCBEE18

LSID. urn:lsid:zoobank.org:act:E3B83181-91D6-44C8-90C0-BBAACEC2CDEE

LSID. urn:lsid:zoobank.org:act:225CD304-3B63-4B55-B8B8-33B46C90A194

LSID. urn:lsid:zoobank.org:act:92C8B169-4F79-467E-B951-EF1DE6E327B1

1101

REFERENCES

AGUILAR, D. H. and LAURITO, C. A. 2009. El armadillo gigante (Mammalia, Xenarthra,

Pampatheriidae) del río Tomayate, Blancano tardío-Irvingtoniano temprano, El

Salvador, América Central. *Revista Geológica de América Central*, **41**, 25–36.

AGUILAR, T., ACEVEDO, B. and ULLOA, A. 2010. Paleontología de una sección del río

Corredores, Formación Curré, Mioceno, Costa Rica. *Revista Geológica de América*

Central, **42**: 43–75.

ALFARO, E. and HOLZ, M. 2014. Review of the chronostratigraphic charts in the Sinú-

San Jacinto Basin based on new seismic stratigraphic interpretations. *Journal of South*

American Earth Sciences, **56**, 139–169.

ALVARO, G. E., BARQUERO, R., TAYLOR, W., LÓPEZ, A. CERDAS, A. and

MURILLO, J. 2009. Geología de la hoja general, Costa Rica. *Revista Geológica de*

América Central, **40**: 97–107.

AMEGHINO, F. 1888. Lista de las especies de mamíferos fósiles del Mioceno superior de

Monte Hermoso hasta ahora conocidas. Obras Completas y Correspondencia Científica,

5, 481–496.

ANDERSON, V. J., HORTON, B. K., SAYLOR, J. E., MORA, A., TESÓN, E.,

BREECKER, D. O. and KETCHAM, R. A. 2016. Andean topographic growth and

basement uplift in southern Colombia: Implications for the evolution of the Magdalena,

Orinoco, and Amazon river systems. *Geosphere*, **12** (4), 1235–1256.

ANTOINE, P. O., ABELLO, M. A., ADNET, S., SIERRA, A. J., BABY, P., BILLET, G.,

BOIVIN, M., CALDERON, Y., CANDELA, M. A., CHABAIN, J., CORFU, F.,

CROFT, D. D., GANERØD, M., JARAMILLO, C., KLAUS, S., MARIVAUX, L.,

NAVARRETE, R. E., ORLIAC, M. J. and PARRA, F. 2016. A 60-million-year

Cenozoic history of western Amazonian ecosystems in Contamana, eastern Peru.

Gondwana Research, **31**, 30–59.

AUSICH, W. I., KAMMER, T. W., RHENBERG, E. C. and WRIGHT, D. F. 2015. Early

phylogeny of crinoids within the pelmatozoan clade. *Palaeontology*, **58** (6): 937–952.

BEARD, K. C. and DAWSON, M. R. 1999. Intercontinental dispersal of Holarctic land

mammals near the Paleocene/Eocene boundary; paleogeographical, paleoclimatic and

biostratigraphic implications. *Bulletin de la Société géologique de France*, **170** (5), 697–

706.

BENGTSON, P. 1988. Open nomenclature. *Palaeontology*, **31** (1), 223–227.

BERMÚDEZ, H. D., ALVARÁN, M., GRAJALES, J. A., RESTREPO, L. C., ROSERO, J.

S., GUZMÁN, C., RUÍZ, E. C., NAVARRETE, R. E., JARAMILLO, C. and OSORNO,

J. F., 2009. Estratigrafía y evolución geológica de la secuencia sedimentaria del

Cinturón Plegado de San Jacinto. *Memorias XII Congreso Colombiano de Geología*, pp.

1–27.

BERNAL-OLAYA, R., MANN, P. and VARGAS, C. A. 2015. Earthquake, tomographic,

seismic reflection, and gravity evidence for a shallowly dipping subduction zone beneath

the Caribbean Margin of Northwestern Colombia. *In* BARTOLINI, C. and MANN, P.

(eds.) *Petroleum Geology and Potential of the Colombian Caribbean Margin*. *AAPG*

Memoir, **118**, 247–270.

BILLET, G., HAUTIER, L., DE MUIZON, C. and VALENTIN, X. 2011. Oldest cingulate

skulls provide congruence between morphological and molecular scenarios of armadillo

evolution. *Proceedings of the Royal Society of London B*, *Biological Sciences*, **278**,

2791–2797.

BLOCH, J. I., WOODRUFF, E. D., WOOD, A. R., RINCON, A. F., HARRINGTON, A.

R., MORGAN, G. S., FOSTER, D. A., MONTES, C., JARAMILLO, C. A., JUD, N. A.,

JONES, D. S. and MACFADDEN, B. J. 2016. First North American fossil monkey and

early Miocene tropical biotic interchange. *Nature*, **533** (7602), 243–258.

BOWEN, G. J., CLYDE, W. C., KOCH, P. L., TING, S., ALROY, J., TSUBAMOTO, T.,

WANG, Y. and WANG, Y. 2002. Mammalian dispersal at the Paleocene/Eocene

boundary. *Science*, **295** (5562), 2062–2065.

BRANDONI, D., RUIZ, L. G. and BUCHER, J. 2019. Evolutive Implications of

Megatericulus patagonicus (Xenarthra, Megatheriinae) from the Miocene of Patagonia

Argentina. *Journal of Mammalian Evolution*, 1–16.

CARLINI, A. A., ZURITA, A. E., SCILLATO-YANÉ, G. J., SÁNCHEZ, R. and

AGUILERA, O. A. 2008. New Glyptodont from the Codore Formation (Pliocene),

Falcón State, Venezuela, its relationship with the *Asterostemma* problem, and the

paleobiogeography of the Glyptodontinae. *Paläontologische Zeitschrift*, **82** (2), 139–

152.

CARRILLO, J. D., FORASIEPI, A., JARAMILLO, C. and SÁNCHEZ-VILLAGRA, M. R.
2015. Neotropical mammal diversity and the Great American Biotic Interchange: spatial
and temporal variation in South America's fossil record. *Frontiers in Genetics*, **5**, 451,
1–11

CARRILLO-BRICEÑO, J. D., REYES-CESPEDES, A. E., SALAS-GISMONDI, R. and
SÁNCHEZ, R. 2018. A new vertebrate continental assemblage from the Tortonian of
Venezuela. *Swiss Journal of Palaeontology*, 1–12.

CARTELLE, C. and BOHÓRQUEZ, G.A. 1985. *Pampatherium paulacoutoi*, uma nova
espécie de tatu gigante da Bahia, Brasil (Edentata, Dasypodidae). *Revista Brasileira de
Zoologia*, **2**: 229-254.

— and DE IULIIS, G. 1995. *Eremotherium laurillardi*: The Panamerican late Pleistocene
megatheriid sloth. *Journal of Vertebrate Paleontology*, **15** (4), 830–841.

— — 2006. *Eremotherium laurillardi* (Lund) (Xenarthra, Megatheriidae), the Panamerican
giant ground sloth: Taxonomic aspects of the ontogeny of skull and dentition. *Journal of
Systematic Palaeontology*, **4** (2), 199–209.

— — and PUJOS, F. 2015. *Eremotherium laurillardi* (Lund, 1842) (Xenarthra,
Megatheriinae) is the only valid megatheriine sloth species in the Pleistocene of
intertropical Brazil: A response to Faure et al., 2014. *Comptes Rendus Palevol*, **14** (1),
15–23.

CASTELLANOS, A. 1937. Anotaciones sobre la línea filogenética de clamiterios. *Serie*

Técnico-Científica de la Facultad de Ciencias Matemáticas, Físico-Químicas y

Naturales, **8**:1–35. Rosario, Argentina.

CIONE, A., GASPARINI, G., SOIBELZON, E., SOIBELZON, L. and TONNI, E. 2015.

The Great American Biotic Interchange. A South American perspective. Springer Briefs

in Earth System Sciences. South America and the Southern Hemisphere, Amsterdam,

Netherlands, 97 pp.

COATES, A. G. and STALLARD, R. F. 2013. How old is the Isthmus of Panama? *Bulletin*

of Marine Science, **89** (4), 801–813.

CORTES, J. E., AGUILERA, R., WILCHES, O., OSORNO, J. F. and CORTES, S. I. 2018.

Organic geochemical insights from oil seeps, tars, rocks, and mud volcanoes on the

petroleum systems of the Sinú-San Jacinto basin, Northwestern Colombia. *Journal of*

South American Earth Sciences, **86**, 318–341.

COZZUOL, M. A., 2006. The Acre vertebrate fauna: age, diversity, and geography.

Journal of South American Earth Sciences, **21**, 185–203.

CROFT, D. A. 2007. The Middle Miocene (Laventan) Quebrada Honda fauna, southern

Bolivia and a description of its notoungulates. *Palaeontology*, **50** (1), 277–303.

DE IULIIS, G. and EDMUND, A. G. 2002. *Vassallia maxima* Castellanos, 1946

(Mammalia: Xenarthra: Pampatheriidae), from Puerta del Corral Quemado (late

Miocene to early Pliocene), Catamarca Province, Argentina. *Smithsonian Contributions*

to Paleobiology, **93**, 49–64.

DELSUC, F., SUPERINA, M., TILAK, M. K., DOUZERY, E. J. and HASSANIN, A. 2012. Molecular phylogenetics unveils the ancient evolutionary origins of the enigmatic fairy armadillos. *Molecular Phylogenetics and Evolution*, **62**, 673–680.

EBACH, M. C. and Williams, D. M. 2010. Aphyly: a systematic designation for a taxonomic problem. *Evolutionary biology*, **37** (2-3), 123–127.

EDMUND, A. G. 1985. The Armor of fossil giant Armadillos (Pampatheriidae, Xenarthra, Mammalia). Texas Memorial Museum, *Pearce-Sellards-Series*, **40**, 1–20.

— 1987. Evolution of the Genus *Holmesina* (Pampatheriidae, Mammalia) in Florida, with Remarks on Taxonomy and Distribution. Texas Memorial Museum, *Pearce-Sellards-Series*, **45**, 1–20.

— and THEODOR, J. 1997. A new giant Armadillo. *In* KAY, R. F., CIFELLI, R. L., FLYNN, J. J., and MADDEN, R. (eds.), *Vertebrate Paleontology of the Miocene Fauna of La Venta, Colombia*. Smithsonian Institution Press, Washington, pp. 227–232

FERNÍCOLA, J. C. 2008. Nuevos aportes para la sistemática de los Glyptodontia Ameghino 1889 (Mammalia, Xenarthra, Cingulata). *Ameghiniana*, **45** (3), 553–574.

FLINCH, J. F. 2003. Structural evolution of the Sinú-Lower Magdalena area (northern Colombia). *In* BARTOLINI, C., BUFFLER, R. T. and BLICKWEDE, J. (eds.), *The Circum-Gulf of Mexico and the Caribbean: Hydrocarbon Habitats, Basin Formation, and Plate Tectonics*, *AAPG Memoir*, **79**, 776–796.

FLYNN J. J., GUERRERO J. and SWISHER, C. C. III. 1997. Geochronology of the Honda Group. *In* KAY, R. F., CIFELLI, R. L., FLYNN, J. J., and MADDEN, R. (eds.),

Vertebrate Paleontology of the Miocene Fauna of La Venta, Colombia. Smithsonian Institution Press, Washington, pp. 44–59.

GAUDIN, T. J. 2004. Phylogenetic relationships among sloths (Mammalia, Xenarthra, Tardigrada): the craniodental evidence. *Zoological Journal of the Linnean Society*, **140**, 255–305.

— and WIBLE, J. R. 2006. The phylogeny of living and extinct armadillos (Mammalia, Xenarthra, Cingulata): a craniodental analysis. *In* CARRANO, M. T., GAUDIN, T. J., BLOB, R. W. and WIBLE, J. R. (eds.), *Amniote Paleobiology: Perspectives on the Evolution of Mammals, Birds and Reptiles*, University of Chicago Press, Chicago, pp. 153–198.

— and LYON, L. M. 2017. Cranial osteology of the pampathere *Holmesina floridanus* (Xenarthra: Cingulata; Blancan NALMA), including a description of an isolated petrosal bone. *PeerJ*, **5**, e4022.

GÓIS, F. 2005. Estudo descritivo e geométrico dos Cingulata (Mammalia, Xenarthra) do Neógeno e Quaternário da Amazônia Sul-Ocidental. Unpublished bachelor thesis, Universidade Federal de Rondônia, 58 pp.

— 2013. Análisis morfológico y afinidades de los Pampatheriidae (Mammalia, Xenarthra). Unpublished PhD thesis, Universidad Nacional de La Plata, La Plata, Argentina, 330 pp.

— SCILLATO-YANÉ, G. J., CARLINI, A. A. and UBILLA, M. 2012. Una nueva especie de *Holmesina* Simpson (Xenarthra, Cingulata, Pampatheriidae) del Pleistoceno de Rondônia, sudoeste de la amazonia, Brasil. *Revista Brasileira de Paleontologia*, **15** (2), 211–227.

— SCILLATO-YANÉ, G. J., CARLINI, A. A. and GUILHERME, E. 2013. A new species

of *Scirrotherium* Edmund & Theodor, 1997 (Xenarthra, Cingulata, Pampatheriidae)

from the late Miocene of South America. *Alcheringa: An Australasian Journal of*

Palaeontology, **37** (2), 177–188.

— RUIZ, L. R., SCILLATO-YANÉ, G. J. and SOIBELZON, E. 2015. A peculiar new

Pampatheriidae (Mammalia: Xenarthra: Cingulata) from the Pleistocene of Argentina

and comments on Pampatheriidae diversity. *PloS One*, **10** (6), e0128296.

— NASCIMENTO, E. R., PORTO, A. S., HOLANDA, E. C. and COZZUOL, M. A. 2004.

Ocorrências de Cingulata dos gêneros *Kraglievichia* e *Holmesina* do Terciário e

Quaternário da Amazônia Sul-Ocidental. *Ameghiniana*, **49**, 41.

GUERRERO, J. 1997. Stratigraphy and sedimentary environments of the Honda Group in

the La Venta area. Miocene uplift of the Colombian Andes. *In* KAY, R. F., CIFELLI, R.

L., FLYNN, J. J., and MADDEN, R. (eds.), *Vertebrate Paleontology of the Miocene*

Fauna of La Venta, Colombia. Smithsonian Institution Press, Washington, pp. 15–43.

GOLOBOFF, P. A. 2014. Extended implied weighting. *Cladistics*, **30** (3): 260–272.

— TORRES, A. & ARIAS, J. S. 2018. Weighted parsimony outperforms other methods of

phylogenetic inference under models appropriate for morphology. *Cladistics*, **34** (4):

407–437.

— CARPENTER, J. M., SALVADOR ARIAS, J. and MIRANDA ESQUIVEL, D. R.

2008. Weighting against homoplasy improves phylogenetic analysis of morphological

data sets. *Cladistics*, **24** (5): 758–773.

HOFFSTETTER, R. 1952. Les mammifères pleistocenes de la République de l'Équateur. *Mémoires Societé Géologique de France*, **66**: 1–391.

JARAMILLO, C. 2018. Evolution of the Isthmus of Panama: biological, paleoceanographic, and paleoclimatological implications. *In* HOORN, C. and ANTONELLI, A. (eds.), *Mountains, climate and biodiversity*. John Wiley & Sons, Oxford, pp. 323–338

— ROMERO, I., D'APOLITO, C., BAYONA, G., DUARTE, E., LOUWYE, S., ESCOBAR, J., LUQUE, J., CARRILLO-BRICEÑO, J. D., ZAPATA, V., MORA, A., SCHOUTEN, S., ZAVADA, M., HARRINGTON, G. and WESSELING, F. P. 2017. Miocene flooding events of western Amazonia. *Science advances*, **3** (5), e1601693.

KERBER, L., NEGRI, F. R., RIBEIRO, A. M., NASIF, N., SOUZA-FILHO, J. P. and FERIGOLO, J. 2017. Tropical fossil caviomorph rodents from the southwestern Brazilian Amazonia in the context of the South American faunas: systematics, biochronology, and paleobiogeography. *Journal of Mammalian Evolution*, **24** (1), 57–70.

KOLARSKY, R. A., MANN, P. and MONTERO, W. 1995. Island arc response to shallow subduction of the Cocos Ridge, Costa Rica. *Special papers of the Geological Society of America*, **295**: 235–262.

LATRUBESSE, E. M., COZZUOL, M. A., SILVA-CAMINHA, S. A., RIGSBY, C. A., ABSY, M. L. and JARAMILLO, C., 2010. The late Miocene paleogeography of the Amazon Basin and the evolution of the Amazon River system. *Earth-Science Reviews*, **99**, 99–124.

LAURITO, C. A. and VALERIO, A. L. 2013. *Scirrotherium antelucanus*, una nueva

especie de Pampatheriidae (Mammalia, Xenarthra, Cingulata) del Mioceno Superior de

Costa Rica, América Central. *Revista Geológica de América Central*, **49**, 45–62.

LEIDY, J. 1889. Fossil Vertebrates from Florida. *Proceedings of the Academy of Natural*

Sciences of Philadelphia, **41**: 96-97.

LOWERY, B. J. 1982. Sedimentology and tectonic implications of the Middle to Upper

Miocene Curré Formation. Unpublished M.S. thesis, Louisiana State University, Baton

Rouge, 100 pp.

LUND, P. 1839. Blik paa Braziliens Dyreverden for Sidste Jordomvaeltning. Anden

Afhandling: Pattedyrene. *Det Kongelige Danske Videnskabernes*

SelskbasNaturvidenskabelige og Mathematiske Afhandlinger, **8**: 61-144.

— 1842. Blik paa Braziliens Dyreverden for Sidste Jordomvaeltning. TredieAfhandling:

Forsaettelse af Pattedyrene. *Det Kongelige Danske Videnskabernes*

SelskbasNaturvidenskabelige og Mathematiske Afhandlinger, **8**: 217–272.

MACFADDEN, B. J. 2006. Extinct mammalian biodiversity of the ancient New World

tropics. *Trends in Ecology & Evolution*, **21** (3), 157–165.

MADDISON, W. P. and MADDISON, D. R. 2010. Mesquite: a modular system for

evolutionary analysis. Version 2.73. [http://mesquiteproject.org]

MARTINELLI, A. G., FERRAZ, P. F., CUNHA, G. C., CUNHA, I. C., DE SOUZA

CARVALHO, I., RIBEIRO, L. C. B., MACEDO NETO, F., LOURENCINI

CAVELLANI, C., ANTUNES TEIXEIRA, V. P. and DA FONSECA FERRAZ, M. L.

2012. First record of *Eremotherium laurillardi* (Lund, 1842) (Mammalia, Xenarthra,

Megatheriidae) in the Quaternary of Uberaba, Triângulo Mineiro (Minas Gerais State),

Brazil. *Journal of South American Earth Sciences*, **37**, 202–207.

MCDONALD, H. G. and LUNDELIUS, E. L., Jr. 2009. The giant ground sloth

Eremotherium laurillardi (Xenarthra. Megatheriidae) in Texas. *In* ALBRIGHT, L.B., III

(ed.), Museum of Northern Arizona, *Papers on geology, vertebrate paleontology, and*

biostratigraphy in honor of Michael O. Woodburne, Bulletin **65**, 407–421.

MCKENNA, M.C. and BELL, S.K. 1997. *Classification of Mammals Above the Species*

Level. Columbia University Press, New York, 640 pp.

MORENO, E. P. and MERCERAT, A. 1891. Exploración arqueológica de la provincia de

Catamarca: Paleontología. *Revista del Museo de La Plata*, **1**: 222-236.

MORENO, F., HENDY, A. J. W., QUIROZ, L., HOYOS, N., JONES, D. S., ZAPATA, V.,

ZAPATA, S., BALLEN, G. A., CADENA, E., CÁRDENAS, A. L., CARRILLO-

BRICEÑO, J. D., CARRILLO, J. D., DELGADO-SIERRA, D., ESCOBAR, J.,

MARTÍNEZ, J. I., MARTÍNEZ, C., MONTES, C., MORENO, J., PÉREZ, N.,

SÁNCHEZ, R., SUÁREZ, C., VALLEJO-PAREJA, M. C. and JARAMILLO, C. 2015.

Revised stratigraphy of Neogene strata in the Cocinetas basin, La Guajira, Colombia.

Swiss Journal of Palaeontology, **134** (1), 5–43.

OBANDO, L. G. 2011. Stratigraphic and tectonic of northeast part of Dota quadrangle (1:

50,000), Costa Rica. *Revista Geológica de América Central*, **44**: 71–82.

O'DEA, A., LESSIOS, H. A., COATES, A. G., EYTAN, R. I., RESTREPO-MORENO, S.

A., CIONE, A. L., COLLINS, L. S., DE QUEIROZ, A., FARRIS, D. W., NORRIS, R.

D., STALLARD, R. F., WOODBURNE, M. O., AGUILERA, O., AUBRY, M-P.,

BERGGREN, W. A., BUDD, A. F., COZZUOL, M. A., COPPARD, S. E., DUQUE-

CARO, H., FINNEGAN, S., GASPARINI, G. M., GROSSMAN, E. L., JHONSON K.

G., LLOYD, D. K, KNOWLTON N., LEIGH E. G., LEONARD-PINGEL J. S.,

MARKO, P. B., PYENSON, N. D., RACHELLO-DOLMEN, P. G., SOIBELZON, E.,

SOIBELZON, L., TODD, J. A., VERMEIJ, G. J. and JACKSON, J. B. C. 2016.

Formation of the Isthmus of Panama. *Science advances*, **2** (8), e1600883.

ORTIZ-JAUREGUIZAR, E. and CLADERA, G. A. 2006. Paleoenvironmental evolution of

southern South America during the Cenozoic. *Journal of Arid Environments*, **66** (3),

498–532.

PATTERSON, B. and PASCUAL, R. 1968. The fossil mammal fauna of South America.

The Quarterly Review of Biology, **43** (4), 409–451.

RIBEIRO, A. M., MADDEN, R. H., NEGRI, F. R., KERBER, L., HSIOU, A. S. and

RODRIGUES, K. A. 2013. Mamíferos fósiles y biocronología en el suroeste de la

Amazonia, Brasil. *In* BRANDONI, D. and NORIEGA, J.I. (eds.), *El Neógeno de la*

Mesopotamia argentina, Asociación Paleontológica Argentina, Publicación Especial **14**,

207–221.

RINCÓN, A. D., SOLÓRZANO, A., BENAMMI, M., VIGNAUD, P. and MCDONALD,

H. G. 2014. Chronology and geology of an Early Miocene mammalian assemblage in

North of South America, from Cerro La Cruz (Castillo Formation), Lara State,

Venezuela: Implications in the changing course of Orinoco River hypothesis. *Andean*

geology, **41** (3), 507–528.

— — MACSOTAY, O., MCDONALD, H. G. and NÚÑEZ-FLORES, M. 2016. A new

Miocene vertebrate assemblage from the Río Yuca Formation (Venezuela) and the

northernmost record of typical Miocene mammals of high latitude (Patagonian) affinities

in South America. *GeoBios*, **49** (5), 395–405.

RIVIER, F. 1985. Sección geológica del Pacífico al Atlántico a través de Costa Rica.

Revista Geológica de América Central, **2**: 22–32.

ROBERTSON, J. S. 1976. Latest Pliocene mammals from Haile XV A, Alachua county,

Florida. *Bulletin of the Florida State Museum, Biological Sciences*, **20**: 111–186.

SALAS-GISMONDI, R., FLYNN, J. J., BABY, P., TEJADA-LARA, J. V.,

WESSELINGH, F. P. and ANTOINE, P. O. 2015. A Miocene hyperdiverse crocodylian

community reveals peculiar trophic dynamics in proto-Amazonian mega-wetlands.

Proceedings of the Royal Society B: Biological Sciences, **282** (1804), 20142490.

SÁNCHEZ-VILLAGRA, M. R. 2012. Embryos in Deep Time: The Rock Record

of Biological Development. University of California Press, Berkeley, California, 256 pp.

SCHMIDT, D. N. 2007. The closure history of the Panama Isthmus: evidence from

isotopes and fossils to models and molecules. *In* WILLIAMS, M., HAYWOOD, A. M.,

GREGORY J., F. and SCHMIDT, D. N. (eds), *Deep time perspectives on climate*

change – marrying the signal from computer models and biological proxies. Geological

Society of London, London, pp. 427–442.

SCILLATO-YANÉ, G. J., CARLINI, A. A., TONNI, E. P. and NORIEGA, J. I. 2005.

Paleobiogeography of the late Pleistocene pampatheres of South America. *Journal of*

South American Earth Sciences, **20** (1–2), 131–138.

— GÓIS, F., ZURITA, A. E., CARLINI A. A., GONZÁLEZ RUIZ, L. R., KRMPOTIC, C. M., OLIVA, C. and ZAMORANO, M. 2013. Los Cingulata (Mammalia, Xenarthra) del "Conglomerado Osífero" (Mioceno tardío) de la Formación Ituzaingó de Entre Ríos, Argentina. *In* BRANDONI, D. and NORIEGA, J.I. (eds.), *El Neógeno de la Mesopotamia argentina*, Asociación Paleontológica Argentina, Publicación Especial **14**, 118–134.

SIGOVINI, M., KEPPEL, E. and TAGLIAPIETRA, D. 2016. Open Nomenclature in the biodiversity era. *Methods in Ecology and Evolution*, **7** (10), 1217–1225.

SIMPSON, G. G. 1930. *Holmesina septentrionalis*, extinct giant armadillo of Florida. *American Museum Novitates*, **442**: 1–10.

SILVA, J. C., PARDO, A., CARDONA, A., BORRERO, C., FLORES, A., NAVARETTE, R., MEJÍA, A., OCHOA, D., OSORIO, J. A., ROSERO, S. and ARENAS, A. 2012. Multi-Chronological Proxies to Timing Early Oligocene and Middle Miocene Deformation Events Along the Lower Magdalena Basin, NW Colombia. *Resúmenes del XI Simposio Bolivariano-Exploración Petrolera en las Cuencas Subandina*s, Asociación Colombiana de Geólogos y Geofísicos del Petróleo. [http://www.earthdoc.org/publication/publicationdetails/?publication=66175]

STIRTON, R. A. 1953. Vertebrate paleontology and continental stratigraphy in Colombia. *Geological Society of America Bulletin*, **64** (6), 603–622.

SWOFFORD, D. L. 2015. PAUP*: Phylogenetic analysis using parsimony (and other methods) (version 4.0a142).

⸺ and BELL, C. D. 2017. PAUP* manual. Available at http://phylosolutions.com/paup-

documentation/paupmanual.pdf.

TITO, G. 2008. New remains of *Eremotherium laurillardi* (Lund, 1842) (Megatheriidae,

Xenarthra) from the coastal region of Ecuador. *Journal of South American Earth*

Sciences, **26** (4), 424–434.

VILLARROEL, C. and CLAVIJO, J. 2005. Los mamíferos fósiles y las edades de las

sedimentitas continentales del Neógeno de la Costa Caribe Colombiana. *Revista de la*

Academia Colombiana de Ciencias, **29** (112), 345–356.

VIZCAÍNO, S. F., DE IULIIS, G. and BARGO, M. S. 1998. Skull shape, masticatory

apparatus, and diet of *Vassallia* and *Holmesina* (Mammalia: Xenarthra:

Pampatheriidae): when anatomy constrains destiny. *Journal of Mammalian Evolution*, **5**

(4), 291–322.

WEBB, S. D. 2006. The Great American Biotic Interchange: Patterns and Processes.

Annals of the Missouri Botanical Garden, **93** (2), 245–258.

WOODBURNE, M. O. 2010. The Great American Biotic Interchange: dispersals, tectonics,

climate, sea level and holding pens. *Journal of Mammalian Evolution*, **17** (4), 245–264.

⸺ and SWISHER, C. C. III. 1995. Land mammal high-resolution geochronology,

intercontinental overland dispersals, sea level, climate and vicariance. *In* BERGGREN,

W. A., KENT, D. W., AUBRY, M.-P. and HARDENBOL, J. (eds.), *Geochronology,*

Time Scales and Global Stratigraphic Correlation. *SEPM Special Publication*, **54**, 335-

364.

— CIONE, A. L. and TONNI, E. P. 2006. Central American provincialism and the Great

American Biotic Interchange. Universidad Nacional Autónoma de México, Instituto de

Geología y Centro de Geociencias, *Publicación Especial*, **4**, 73–101.

YUAN, P. B. 1984. Stratigraphy, sedimentology and geologic evolution of eastern Terraba

Trough, southwestern Costa Rica. Unpublished PhD thesis, Louisiana State University,

Baton Rouge, 110 pp.

ZURITA, A. E., SCILLATO-YANÉ, G. J., CIANCIO, M., ZAMORANO, M. and

GONZÁLEZ-RUIZ, L. R. 2016. Los Glyptodontidae (Mammalia, Xenarthra): Historia

biogeográfica y evolutiva de un grupo particular de mamíferos acorazados.

Contribuciones del MACN, **6**, 249-262.

1429

FIGURE CAPTIONS

Figure 1. Geographical and stratigraphic provenance of the newly described material of

pampatheriids from the Neogene of Colombia. In the left upper corner, a map of

northwesternmost South America and the location of the regions of Colombia where there

are outcrops of the formations with pampatheriid specimens for this study. In the right

upper corner, photos of characteristic outcrops of these formations. Below in the center, a

general chronostratigraphic scheme with the position of each formation within the Neogene

and two important tectonic/palaeogeographical events in northwestern South America, i.e. a

major, underwater uplift of the Isthmus of Panama (Schmidt 2007) and the definitive

emergence of the Panama Land Bridge (O'dea et al. 2016). The photo of outcrops of the

Castilletes Formation was taken by Edwin Cadena.

Figure 2. Phylogenetic results. A, strict consensus tree of the parsimony with equal

weights. Numbers below nodes are absolute Bremer support values. B, strict consensus tree

of the parsimony analysis with implied weights. C, majority-rule consensus tree of the

bootstrap resampling with implied weights. The numbers below nodes are bootstrap

support values. Note the difference in the phylogenetic position of *Holmesina floridanus* in

the two strict consensus trees. Explanation of this difference in the main text.

Figure 3. Fixed and (semi) mobile osteoderms of *Scirrotherium hondaensis* from the La

Victoria and Villavieja Formations, Municipality of Villavieja, Department of Huila,

Colombia. A–B', fixed osteoderms; C'–K', (semi) mobile osteoderms. The osteoderms G,

J, K, L, W, X, Y, Z, A', B', G', I', J' and K' are associated with the catalog number VPPLT

348. The osteoderms H, U and V are associated with the catalog number VPPLT 004. The

osteoderms T and D' are associated with the catalog number VPPLT 701. All the former

osteoderms come from the lower and middle La Victoria Formation. The osteoderms B, C,

F, I, O, P, S, C' and F' are associated with the catalog number VPPLT 1683 - MT 18 and

come from the top of the La Victoria Formation. The osteoderms A, D, E, M, N, Q, R, E'

and H' are associated with the catalog number VPPLT 1683 - MT 18 and come from the

lower Villavieja Formation. Scale bar equal to 20 mm.

Figure 4. Photos and anatomical line drawings of the skull VPPLT 706 of *Scirrotherium*

hondaensis from the middle La Victoria Formation, Municipality of Villavieja, Department

of Huila, Colombia. A–B, dorsal views; C–D, ventral views; E–F, right lateral views; D–H,

left lateral views. *Abbreviations*: aof, antorbital fossa; fr, frontals; iof, infraorbital foramen;

j, jugal; la, lacrimal; Mf1, first upper molariform; Mf9, ninth upper molariform; mx,

maxilla; mxf, maxillary foramen; na, nasals; pal, palatines; pm, premaxilla. Scale bar equal

to 50 mm.

Figure 5. Photos and anatomical line drawings of the left femur and right ulna VPPLT 706

of *Scirrotherium hondaensis* from the middle La Victoria Formation, Municipality of

Villavieja, Department of Huila, Colombia. The epiphyses of this femoral diaphysis have

been reconstructed from those with catalog number UCMP 39846. A–B, anterior views of

the femur; C–D, posterior views of the femur. E–F, medial views of the ulna; G–H, lateral

views of the ulna. *Abbreviations*: anc, fossa for the anconeus muscle; cp, coronoid process;

fh, femoral head; gt, greater trochanter; le, lateral epicondyle; me, medial epicondyle; op,

olecranon process; tn, trochlear notch; tt, third trochanter. Scale bar equal to 50 mm.

Figure 6. Photos and anatomical line drawings of a thoracic vertebra (A–B) and several

anterior caudal vertebrae (C–F) VPPLT 706 of *Scirrotherium hondaensis* from the middle

La Victoria Formation, Municipality of Villavieja, Department of Huila, Colombia. A–B,

posterior views of the thoracic vertebra. C–D, lateral views of caudal vertebrae; E–F, dorsal

views of caudal vertebrae. *Abbreviations*: az, anterior zygapophyses; mp, metapophyses;

ns, neural spine; tp, transverse processes; vb, vertebral body; vla, ventrolateral apophyses.

Scale bar equal to 30 mm.

Figure 7. Photos and anatomical line drawings of the astragalus (A–D) and calcaneum (E –

F) UCMP 39846 of *Scirrotherium hondaensis* from the lower (?) Villavieja Formation,

Municipality of Villavieja, Department of Huila, Colombia. A–B, astragalus in plantar

views; C–D, astragalus in dorsal views. E–F, calcaneum in dorsal views. *Abbreviations*: ct,

calcaneal tuber; ef, ectal facet; h, head of the astragalus; lt, lateral trochlea; mf, medial

trochlea; sf, sustentacular facet; st, *sulcus tali*. Scale bar equal to 20 mm.

Figure 8. Fixed osteoderm MUN STRI 36880 of *Scirrotherium antelucanus* from the upper

Sincelejo Formation, Municipality of Los Palmitos, Department of Sucre, Colombia. Scale

bar equal to 20 mm.

Figure 9. Osteoderms referred to aff. *Scirrotherium* (**A**, MUN STRI 16718; **E**, MUN STRI

38064; and **G,** MUN STRI 16719; all these specimens are from the Castilletes Formation

and they are fixed osteoderms except the latter, which consist of an anterior fragment of a

mobile osteoderm); *Scirrotherium* cf. *hondaensis* (**C**, MUN STRI 36814, a fixed osteoderm

from the Castilletes Formation); and *Scirrotherium* sp. (**B**, MUN STRI 36801; **D**, MUN

STRI 16158; and **F**, MUN STRI 34373; all these osteoderms are from the Castilletes

Formation, except the latter, which comes from the Ware Formation); Municipality of

Uribia, Department of La Guajira, Colombia. Note the two well-developed rows of anterior

foramina in the osteoderms MUN STRI 16718 and 38064. Scale bar equal to 20 mm.

Figure 10. Osteoderms of *Kraglievichia carinatum* comb. nov. from the Ituzaingó

Formation, Entre Ríos Province, Argentina. The holotype of this species is marked with

one single asterisk (*) and paratypes with double asterisk (**). A–J, fixed osteoderms; K–

R, (semi) mobile osteoderms. **A**, MLP 69-IX-8-13AC**; **B**, MLP 70-XII-29-1**; **C**, MLP

41-XII-13-905; **D**, MLP 69-IX-8-13AF; **E**, MLP 69-IX-8-13AG; **F**, MLP 41-XII-13-414A;

G, MLP 69-IX-8-13AN; **H**, unknown catalog number; **I**, MLP 69-IX-8-13AK; **J**, MLP 41-

XII-13-414B; **K**, MLP 69-IX-8-13AS; **L**, MLP 69-IX-8-13AE**; **M**, MLP 52-X-1-36; **N**,

MLP 69-IX-8-13AB*; **O**, MLP 41-XII-13-909; **P**, MLP 69-IX-8-13AW; **Q**, MLP 69-IX-8-

13AQ; **R,** MLP 69-IX-8-13AY. Scale bar equal to 20 mm.

Figure 11. One outstanding morphological difference between typical osteoderms of

Scirrotherium and *Kraglievichia*: Uniformly narrow LCE in fixed osteoderms of

Scirrotherium (left; *S. hondaensis*); and anteriorly wide, posteriorly tapered LCE in the

same osteoderms of *Kraglievichia* (right; *K. carinatum* comb. nov.). Not to scale.

Figure 12. Time-calibrated phylogeny of the clade *Scirrotherium + Kraglievichia +*

Holmesina based on the strict consensus cladogram from this work (Fig. 2). Polytomies

from the strict consensus tree (Fig. 2B) were resolved by (1) forcing the monophyly of *S.*

hondaensis and *S. antelucanus* and (2) placing the species *H. septentrionalis* and *H.*

occidentalis as successively basal to the largest South American *Holmesina* species, i.e. *H.*

paulacoutoi and *H. major*. Note the diversification events of the clade *Scirrotherium +*

Kraglievichia + Holmesina are mainly concentrated during the Burdigalian (late Early

Miocene) and Plio-Pleistocene. Likewise, note the relative long ghost lineage of

Holmesina. Images of the pampatheriids are from *PhyloPic* (all available under public

domain): top, *Pampatherium humboldtii* (http://phylopic.org/name/670230e9-4775-493c-

b3ab-31718fb570a3); below, *Holmesina floridanus* (http://phylopic.org/name/73635941-

ed8a-4518-aae8-70e824dbee97).

Figure 13. Geographical distributions and intercontinental dispersal events of the clade

Scirrotherium + Kraglievichia + Holmesina during the Neogene-Pleistocene. The symbols

(i.e. squares, circles and triangles) should not necessarily be interpreted as single localities

but as approximate areas of occurrence. This is especially true for the Pliocene and

Pleistocene epochs.

1528

1529

1530

TABLES

Table 1. Fixed (scapular and pelvic) osteodermal measurements for taxa of interest in this study.

Taxon/Measurement	Length	Width	Thickness	References
S. hondaensis	16□35.2	17.5□27.9	3.7□6.9	This work; Góis et al. 2013
S. antelucanus	28.6□40.9	22□32.4	4.9□7.1	This work; Laurito & Valerio 2013
K. carinatum comb. nov.	20.9□33.5	17□26.1	4.1□5.9	This work; Góis et al. 2013
K. paranensis	30□45	22.5□28.3	6□11	Góis et al. 2013
H. floridanus	24.4□36.7	18.9□32.1	6□9.7	This work; Edmund 1987

Table 2. Mobile and semi-mobile osteodermal measurements for taxa of interest in this study.

Taxon/Measurement	Length	Width	Thickness	References
S. hondaensis	29.4□60	17.9□27.4	4.9□7.3	This work; Góis et al. 2013
S. antelucanus	38.2□64.6	19.4□28.9	□	Laurito & Valerio 2013
K. carinatum comb. nov.	32□54.5	17□28.9	3.9□6	This work; Góis et al. 2013
K. paranensis	60.5□70.5	25□29	7□9	Góis et al. 2013
H. floridanus	61.8□71	17.6□28.5	4.7□6.3	This work; Edmund 1987

Table 3. Selected cranial measurements for the specimen VPPLT 706 of *Scirrotherium hondaensis* and related taxa whose skulls are known.

Taxon/Measurement	GSL	NL	FL	PAL	LUR	PL	References
S. hondaensis	117.3*	~52.8	~55	□	84.1	94.3	This work
K. cf. *paranensis*	194	58	62	74	□	159	This work
*H. floridanus***	249	106.3	75	58.6	133.6	185	This work
H. septentrionalis	290	□	□	□	165	220	Góis et al. 2012

*Incomplete

**Specimen UF 191448

Table 4. Femoral measurements for *Scirrotherium hondaensis* and related taxa whose femur is known.

Measurement/Taxon	GFL	TTW	DW	References
S. hondaensis	162*	27.6	32.5	This work
K. cf. *paranensis*	164	33.7	38	This work
*H. floridanus***	195	41	47	This work
H. septentrionalis	290	70	86	Góis 2013

*Estimated from the specimens VPPLT 706 and UCMP 39846

**Specimen UF 24918

A. Early-to-Middle Miocene Castilletes Formation

B. Middle Miocene La Victoria Formation

C. Middle Miocene Villavieja Formation

D. Late Miocene-to-Early Pliocene Sincelejo Formation

E. Late Pliocene Waré Formation

1. La Venta area, La Victoria and Villavieja Formations (Middle Miocene)

2. Sincelejo Formation (Late Miocene to Pliocene)

3. Castilletes (Early to Middle Miocene) and Waré (Late Pliocene) Formations

— — — Eastern boundary of the ancient Central American Seaway

North America

South America

Central American Seaway

?

Legend:

- ● Early Miocene - *Scirrotherium*
- ● Middle Miocene - *Scirrotherium*
- ● Late Miocene - *Scirrotherium*
- ■ Late Miocene - *Kraglievichia*
- ● Pliocene - *Scirrotherium*
- ▲ Pliocene - *Holmesina*
- ▲ Pleistocene - *Holmesina*

- ⤵ Late Miocene dispersal event
- ⤵ Pliocene dispersal event
- ⤵ Pleistocene dispersal event

0   750 1500 Km