An integrated analysis of myeloid cells identifies gaps in *in vitro* models of *in vivo* biology

Nadia Rajab[1,2], Paul W Angel[1], Yidi Deng[1,3], Jennifer Gu[1], Vanta Jameson[4], Mariola Kurowska-Stolarska[5], Simon Milling[5], Chris M Pacheco[1], Matt Rutar[1], Andrew L Laslett[6,7], Kim Anh Lê Cao[3], Jarny Choi[1] and Christine A Wells*[1].

AFFILIATIONS:
1. The Centre for Stem Cell Systems, Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, 30 Royal Parade Parkville, Victoria 3010, Australia.
2. CSIRO Synthetic Biology Future Science Platform
3. Melbourne Integrative Genomics, School of Mathematics and Statistics, Faculty of Science, The University of Melbourne, 30 Royal Parade Parkville, Victoria 3010, Australia
4. Melbourne Cytometry Platform (MBC node), Department of Anatomy and Neuroscience, University of Melbourne
5. The Institute of Infection, Immunity and Inflammation, Research into Inflammatory Arthritis Centre *'Versus Arthritis'* (RACE), University of Glasgow, Glasgow, United Kingdom.
6. CSIRO Manufacturing, Clayton, Victoria 3168, Australia
7. Australian Regenerative Medicine Institute, Monash University, Victoria 3800, Australia.

*CORRESPONDING AUTHOR email wells.c@unimelb.edu.au

Summary:

The Stemformatics myeloid atlas is an integrated transcriptome atlas of human macrophages and dendritic cells that systematically compares freshly isolated tissue-resident, cultured, and stem-cell derived myeloid cell types. We identified two broad classes of tissue-resident macrophages with lung, gut and tumour-associated macrophages most similar to monocytes. Microglia, Kupffer cells and synovial macrophages shared similar profiles with each other, and with cultured macrophages. Pluripotent stem cell-derived macrophages were not reminiscent of fetal-derived cells. Instead, they were characterized by atypical expression of collagen and a highly efferocytotic phenotype. Likewise, Flt3L-derived cord blood dendritic cells were distinct from conventional dendritic cell subsets isolated from primary tissues and lacked expression of key pattern recognition receptors. Myeloid subsets were reproducible across different experimental series, showing the resource is a robust reference for new data. External users can annotate and benchmark their own samples, including annotation of myeloid single cell data at www.stemformatics.org/atlas/myeloid/.

Introduction
Macrophages are innate immune cells that are found resident in every tissue, with roles in tissue
homeostasis, and response to infection or injury. The distinct functional roles of tissue
macrophages are reflected in their transcriptional phenotypes: atlases of tissue-resident mouse
macrophages, for example, have given great insight into their complexity and heterogeneity
(Gautier et al., 2012; Gosselin et al., 2017). Individual transcriptome studies have played an
essential role in unravelling the importance of the environment on macrophage phenotype and
function (reviewed by (Huang and Wells, 2014)). These data evidence the shared molecular
pathways of myeloid cells responding to pathogenic challenge, and the receptiveness of
macrophages to environmental cues.
11
Much of our understanding of macrophage biology, including many of the molecular mechanisms
of innate immune signaling, have arisen from mouse gene knock-out studies. However, cross-
species comparisons of immune cells highlight differences between mouse and man. These
include the glycolytic switch associated with metabolic reprogramming in activated mouse
macrophages (Vijayan et al., 2019); divergent patterns of pathogen receptor expression (Vijayan
et al., 2012)and transcriptional responses to innate immune stimuli (Schroder et al., 2012). Cross-
species comparisons are further hampered by the absence of population level immune-activation
maps, with most mouse studies in macrophage biology conducted on a limited number of inbred
lines.
21
The need for improved molecular models of primary human cells is evident from the rising
popularity of single cell transcriptomic atlases, exemplified by the human cell atlas consortium
(Hay et al., 2018; Regev et al., 2017). However, unbiased profiling of cells also requires
computational predictions of cell identity, raising further questions about how best to accurately
identify immune cell populations resident in tissues, and discriminate these from circulating or
infiltrating peripheral blood cells. The isolation, and identification of tissue-resident myeloid cells
can be particularly fraught if populations are rare or hard to isolate using enzymatic or other
dissociation methods. These procedures can alter myeloid transcriptomes (Gosselin et al., 2017),
resulting in underrepresentation or phenotypic ambiguity of resident macrophages in single cell
maps of a tissue. It might be argued that human macrophages suffer from an identity crisis, relying
on equivalency to laboratory models of human macrophage biology such as *ex-vivo* culture of
monocyte-derived macrophages, which may not be appropriate as a benchmark for specialized
tissue functions.
35
The potential to model tissue residency, disease phenotypes and activation status of human
macrophages using pluripotent stem cells (PSC) is a growing area of interest (Reviewed (Lee et
al., 2018; Rajab et al., 2018)). However, the anatomical context or developmental ontogeny of
these cells is still not well understood, nor their capacity to model specialized behaviors of myeloid
cells including roles within a tissue niche. While databases such as BloodSpot (Bagger et al.,
2016) and Haematlas (Watkins et al., 2009) provide a useful snapshot of gene expression of
blood types, these lack depth with regard to tissue representation, common laboratory models or
activating stimuli. Consequently, comparisons of new models of human macrophage biology rely
on ad hoc comparisons that do not adequately benchmark their similarity to the diversity of

possible macrophage phenotypes. Our motivation, therefore, was to construct a reference atlas
of human myeloid biology that draws on important studies already in the public domain, and that
can be added to by the research community as new cell models or new profiling platforms become
available. Here, we describe an integrated myeloid transcriptome atlas to identify, benchmark and
analyze human myeloid subpopulations from *ex vivo*, *in vivo* and *in vitro* sources. The atlas is
made available as an interactive online resource https://www.stemformatics.org/atlas/myeloid.
51

A reference atlas for human myeloid biology.

We first compiled a reference transcriptional atlas (Figure 1A and Table S1) from public and
proprietary transcriptomic data from 44 studies and ~900 samples representing peripheral blood
monocytes, tissue-resident, *ex vivo* and *in vitro*-derived macrophages and dendritic cells.
Samples were curated with respect to phenotype, source and isolation method. Datasets were
processed through the Stemformatics pipeline which includes a stringent set of quality control
requirements for hosting on the Stemformatics.org portal (Choi et al., 2019a). We constructed the
atlas by implementing a two-step process (Angel et al., 2020). Firstly, transformation of
expression values from the original studies to percentile values to facilitate the comparison of
different experimental series was performed. Secondly, using a univariate estimation of their
platform dependence, genes whose expression values were significantly impacted by the way
that they were measured were removed from the atlas. This approach led to reproducible
clustering of distinct myeloid classes on a PCA (Figure 1, Figure S1A). The reproducibility of
myeloid subsets including dendritic cells, monocytes and neutrophils was validated by projecting
an independent RNAseq dataset of well annotated blood cell types from Haemopedia (Choi et al.,
2019b) (Figure 1A). Variables such as progenitor source (hematopoietic progenitor cell-, myeloid-
, or pluripotent stem cell (PSC)-derived); gene expression (Figure 1B) or culture status (Figure
1C) contributed to clustering of samples. Patterns of gene expression allow users to assess
markers of myeloid subsets, such as the expression of *TREM1* on monocytes, monocyte-derived
macrophages (MDM), alveolar macrophages and CD1c+ dendritic cells (DC2) (Figure 1B and
Table 1).
73

Two broad classes of tissue resident macrophage span either side of a monocyte border

Isolation of primary tissue-resident macrophages is particularly difficult as this can result in
alterations in phenotype (Gosselin et al., 2017). The difficulty of isolating tissue-resident
populations from healthy human tissue is evident from the spread of tissue resident macrophages
in comparison to tissue-resident dendritic cells (Figure 1), noting that several of the macrophage
datasets were obtained through surgical biopsies from patients with inflammatory disease, and
indeed mapped across the inflammatory axis. Tissue resident macrophages, including Kupffer
cells, microglia, alveolar macrophages, gut and synovial macrophages occupy a broad niche on
the atlas between dendritic cells, peripheral blood monocytes, and cultured monocytes (Figure 1
and Figure S1). Although there was no discrete partitioning of macrophages from individual
tissues, distinct classes of tissue resident macrophages were observed in the atlas – (the alveolar,
colon and macrophages isolated from tumour ascites (TAM) (Figure S1C) grouped together
between cultured monocytes and CD1c+ dendritic cells, and a second spread containing synovial
macrophages, microglia and Kupffer cells aligned with the iPSC-derived cells.
88

DC differentiated from cord blood lack an *in vivo* equivalent.
Circulating and tissue resident dendritic cells (DCs) occupy a distinct transcriptional niche from
monocytes or macrophages. *In vitro*-differentiated dendritic cells, expanded from cord blood
progenitors with FLT3L, do not resemble *in vivo* conventional DCs (DC1 or DC2; Figure 1B, Figure
2A), but are part of an activation axis shared by DC and macrophages. The hallmark of *in vitro*-
DCs are low expression of receptors such as *CX3CR1*, *IL18R1* and *TLR7* (Figure 2B, 2C and
Table S2). Other molecules, such as the cell-fusion protein *DC-STAMP* are gained in culture
(Figure 2C and Table S2). These cells are also closely associated with MDMs, which may
contribute to some of the confusion in the literature about their ontogeny. It is unlikely that this is
due to the cord blood origins of the majority of these datasets, as CD45+ cells isolated from cord
blood engrafted into humanized mice are also included in the atlas (Minoda et al., 2017) and these
recapitulate *in vivo* DC1 and DC2 transcriptional phenotypes.
101
The cDCs form distinct clusters along the activation axis of the atlas. This includes blood samples
taken from donors after vaccination (Banchereau et al., 2014) or cells isolated from inflammatory
fluids (Segura et al., 2013) and we speculate that these represent distinct and reproducible
activation phenotypes. To further evaluate DC subtypes we projected two single cell RNAseq
datasets describing blood monocytes and dendritic cells (Villani et al., 2017) and (Dutertre et al.,
2019) using the myeloid atlas as the reference (Figure 2D, Figure S2). We confirmed the original
observation of a new *AXL+ SIGLEC6+* (AS-DC) subset that shared classical cell surface markers
with plasmacytoid DCs (pDCs) and DC1s (Figure 2E). AS-DCs were suggested to be the subset
contaminating traditional pDC isolation strategies, responsible for observations that pDC can
stimulate T-cells (Villani et al., 2017). There has been some dispute between the Villani and
Dutertre studies on the identity of DC2 subsets (Dutertre et al., 2019). An advantage of projecting
both datasets to the same reference is that the subtypes can be evaluated against a broader set
of reference cell types. Here, the Dutertre DC2 subsets aligned closely with atlas DC2 cells, while
the spread of Villani clusters was much greater, and clearly aligning to respective DC and
monocyte groups on the atlas. Annotation with a Capybara similarity estimate (Figure 2E)
predicted that in addition to the 2 distinct plasmacytoid DC (pDC) subsets in the Villani data, three
distinct DC2 (CD1c+) subsets were present, including an intermediate DC2 subset that sat
between the classical monocyte and DC2 areas (Figure 2E, 2F and Figure S2). Given the number
of additional DC1 and DC2 subsets observed on the myeloid atlas, we predict that further
functional characterization of activated DC phenotypes is warranted.
122
Monocytes rapidly adapt to culture
Monocytes are post-mitotic blood cells derived from bone marrow progenitors that are short lived
in circulation and can repopulate macrophages in some tissue niches. The largest population of
circulating monocytes is marked by high levels of expression of the LPS co-receptor CD14, which
is typically used to isolate monocytes from blood. Intermediate and nonclassical subsets are
marked by acquisition of the type III FcRɣ, CD16 (Schmidl et al., 2014) and are included in the
atlas. Cultured monocytes have been previously described as 'activated', but while we observe a
distinct culture phenotype (Figure 3), the transcriptome of cultured cells mimics many of the
features of a monocyte after extravasation into tissue. Figure 3B shows the grouping of peripheral
blood monocytes, in a distinct cluster to monocytes that have been exposed to tissue culture

plastic and culture media. This culture phenotype is typified here by a decrease in endothelial-
adhesion proteins including the selectin *SELL* (Figure 3D). Regulators of RAS/RAF signaling
including *SPRED2* (Wakioka et al., 2001) have an elevated expression in cultured monocytes
(Figure 3E), consistent with spreading and migration across tissue culture plastics. The classical
MDM requires *in vitro* differentiation of monocytes using several days exposure to growth factors
such as macrophage-colony stimulating factor (M-CSF; CSF-1) or granulocyte-macrophage
colony-stimulating factor (GM-CSF). These group distinctly from the cultured monocyte cluster,
spreading further upwards along the culture axis (Figure 3F).

The culture phenotype acquired by monocytes appears to be a prelude to activation, which can
be observed along an adjacent axis in Figure 3F and is exemplified by the expression of *IL6*.
Pathogen-activated phenotypes of cultured monocytes are typified by high expression of this and
other cytokines. We observe that cultured monocytes express higher levels of chemokines, such
as *CCL2*, than peripheral monocytes (Figure 3G). Furthermore, we find that culture induces the
expression of *SLAMF1* (Table S3), which has shown to be necessary for TLR4 activation in
human macrophages (Yurchenko et al., 2018). Moreover, this culture-phenotype is further
exemplified by higher levels of *ITGB8* than circulating monocytes (Table S3), which is necessary
for activating latent TGF-β (Kelly et al., 2018)

Pluripotent stem cell-derived macrophages do not recapitulate hematopoietic ontogenies.
Many PSC-derived systems recapitulate fetal, rather than adult phenotypes, so it is no surprise
that others have argued that PSC-derivation protocols mimic primitive rather than definitive
myeloid biology. This is largely based on *MYB* expression, which is associated with definitive
hematopoiesis and has high expression in hematopoietic progenitor cells isolated from bone
marrow. It is clear that *MYB* is not required for PSC-derived myelopoiesis as macrophages can
be grown from *MYB* knock-out embryonic stem cells (Buchrieser et al., 2017). Nevertheless, *MYB*
is highly and ubiquitously expressed in PSC-myeloid progenitors, including common myeloid
progenitors and hemogenic endothelium differentiated from pluripotent stem cell, and indeed,
MYB is retained at high levels in some PSC-derived microglia (Figure 4A).

PSC-macrophage studies that were overlaid onto the atlas grouped broadly with cultured
monocytes and tissue-resident macrophages. Despite arguments on recapitulation of fetal origin,
PSC-macrophages, including PSC-microglia and PSC-Kupffer cells, form an extended group
associated with high expression of the human homologue of the *F4/80* antigen, *ADGRE1*, as well
as high expression of lipid-scavenging receptors such as *SCARB1* (Table S4). MAF expression
is indistinguishable in macrophages of different origin (Figure 4B). However, *LIN28B* is highly
expressed in *in vitro*-derived macrophages compared to *in vivo* and *ex vivo* cells (Figure 4C),
which may point to incomplete silencing of the Let7 microRNA pathway and maintenance of a
fetal state (Grant Rowe et al., 2016; Zhou et al., 2017), but which is also characteristic of myeloid
leukemias.

When pluripotent stem cells are projected onto the atlas, we can see a differentiation trajectory
that is orthogonal to the hematopoietic multipotent progenitor trajectory (Figure 4D). The
differentiation trajectory from pluripotent stem cell to macrophage aligns more closely with an

endothelial progenitor than the hematopoietic intermediates observed along PC1 of the atlas.
Projection of primitive hematopoietic progenitors from Carnegie staged embryos at CS11 to CS23
show a high degree of similarity between definitive and primitive differentiation trajectories, with
a predominant erythroid overlap in early yolk sac progenitors, followed by rapid emergence of
myeloid progenitors, and primitive macrophages. These do not overlap with the PSC-axis, but do
clearly sit with similar late fetal microglia (20 week old, (Thion et al., 2018)) and adult bone-marrow
equivalents (Watkins et al., 2009) (Figure 4E and Figure S3).

Some other phenotypes previously attributed to ontogeny in PSC-derived cells may rather reflect
a more general culture context. For example, *ADGRE1* (*F4/80*) expression has been attributed to
yolk-sac derived myeloid cells in mouse (Schulz et al., 2012). While high on PSC-derived
cells, *ADGRE1* is also clearly upregulated in *ex vivo* cultured macrophages. This is exemplified
by primary human microglia, which have low expression of *ADGRE1* in comparison to *ex vivo*
culture or PSC-derived cells (Table S4 and Table S5).

Pluripotent stem cell-derived macrophages share features with tissue-resident
macrophages despite poor maturation
Macrophages derived from human PSCs offer new opportunities to model *in vivo* macrophage
biology. When reviewing the studies contributing to this atlas, we noted that PSC-derived
macrophages are typically benchmarked against monocyte-derived macrophages (MDMs), or
cultured primary cells, using a suite of phenotyping techniques. Each experiment includes a small
number of samples for transcriptional profiling, with a few notable exceptions (Alasoo et al., 2018).
We argue that, given the spectrum of possible resident tissue macrophage phenotypes, it would
be more useful to compare PSC-derived cells against an atlas of possible macrophage
phenotypes. Whilst several groups reuse publicly available tissue macrophage data, the
opportunity to carry out large-scale comparisons to different primary myeloid cells has been
limited by the availability of relevant data on a compatible platform.

Primary microglia included in the atlas include both *in vivo* isolated fetal and cultured *ex vivo* fetal
and adult microglia. The profiles of *in vivo* isolated fetal microglia cluster apart from the spread of
ex vivo cultured adult and fetal microglia (Figure S1B). Projection of single cell data from human
fetal yolk sac and head (Bian et al., 2020) show high concordance with the atlas fetal microglia
data, suggesting that age and culture environment override an ontogeny signature (Figure 4 and
Figure S3). Cultured tissue macrophages including *ex vivo* primary microglia or Kupffer cells
shared a broad transcriptional signature with MDMs, and pluripotent-stem cell derived myeloid
cells. Microglia represent just over a third of PSC-directed myeloid differentiation studies in the
atlas. These do not resolve into a unique cluster but share transcriptional phenotypes with PSC-
derived and tissue-resident macrophages

There are exceptions which include some 'cytokine-matured' PSC-derived microglia samples
from (Abud et al., 2017). These are close to the *in vivo* fetal microglia samples of the PSC-
microglia but are also closely associated with other primary tissue resident macrophages from
lung, joint and gut. The atlas does provide an opportunity to review the expression of markers
thought to distinguish microglia from other primary macrophages. *TMEM119*, for example, is

largely restricted to primary or PSC-derived microglia, although some PSC-microglia samples
have low expression of this marker (Figure S1B). *P2RY12* is variably expressed across all
microglial samples, but its expression is also evident in different tissue-resident samples including
those derived from gut and synovial tissues (Figure S1C and Table S4 and Table S5).

The majority of PSC-derived macrophages have low expression of HLA relevant genes including
CIITA, a known master regulator of MHCII gene expression, which suggests poor maturation and
limited capacity to present antigen to lymphocytes. Nevertheless, some *in vitro*-derived
macrophages cultured with stimulating factors (Figure S1D) do show inducible *CIITA* expression,
demonstrating that they have the capacity to express antigen-presenting machinery. It is also
worth noting additional culture conditions that result in high *CIITA* expression without interferon-
stimulation (Figure S1D). This may be the result of long-term culture conditions for microglia
samples (Abud et al., 2017), or reflect prior conditioning of myeloid progenitors (Honda-Ozaki et
al., 2018).

Pluripotent stem cell-derived macrophages display transcriptional hallmarks of
efferocytosis
Efferocytosis, or apoptotic cell clearance, has broad immunomodulatory effects (Reviewed by
(Elliott et al., 2017)) and active engulfment and clearance of cells by PSC-macrophages is clearly
observed in the absence of any inflammatory activation (Supplementary Video). Efferocytosis
(Figure 5A) modulates macrophages from a pro-inflammatory phenotype to one with resolving
qualities (Yamaguchi et al., 2014), consistent with the patterns of gene expression observed in
cultured macrophages. A study (Cao et al., 2019) demonstrated higher lipid uptake in PSC-
macrophages compared to peripheral blood MDMs, concordant with higher expression of
efferocytosis-related genes including *S1PR1* and *MERTK*. We confirm that *MERTK* is generally
highly expressed in PSC-derived macrophages, but that there is also a tissue-resident distribution
of *MERTK* expression, with very low levels observed in primary alveolar macrophages, and
highest levels observed in human fetal microglia (Figure 5B). Tissue-resident macrophages are
known first-responders to tissue damage and are key in orchestrating inflammation and its
subsequent resolution. This appears to be a phenotype that is selected for in cultured
macrophages and may be an inevitable consequence of derivation that strives for high cell yield.

Lipid homeostasis is an important role for resident tissue macrophages. A high proportion of
genes differentially expressed between *in vitro*-derived macrophages/microglia/kupffer cells and
tissue resident cells are involved in lipid transport, catabolism and in buffering the cells from
concomitant stresses associated with lipid turn-over. For example, reduced expression of
ABCA6 is consistent with high efflux of cholesterol from these macrophages (Figure 5C). Higher
levels of mitochondrial acyl-CoA dehydrogenase *ACADM* (Figure 5D) and phosphatidate
phosphatase *LPIN3* (Figure 5E) suggests high lipid turnover.

There has been growing interest in the importance of metabolic reprogramming in macrophage
responses, so we asked whether media supplementation could explain the spread of PSC-derived
macrophages on the atlas. All PSC-derivation protocols supplement media with fatty- or amino-
acids, including L-Glutamine, non-essential amino acids (NEAA), Linoleic and Linolenic acids.

Some methods add fetal bovine or calf serum, but there was no obvious correlation between
serum addition and without. Overall, factors are so ubiquitously used that a particular supplement
alone could not explain the differences between PSC and cultured primary macrophages.
268

Pluripotent stem cell-derived macrophages express high levels of collagen

Collagen production and deposition alongside extracellular matrix remodeling are processes
involved in wound healing and scarring. Macrophages are instrumental in instructing tissue repair,
particularly through the production of growth factors such as TGF-β, IGF1 and PDGF (Shook et
al., 2018). Secreted growth factors drive fibroblasts and endothelial cells to produce extracellular
matrix components, promoting keloid formation as well as angiogenesis. This model has
macrophages influencing collagen deposition by neighboring stromal cells, however, in some
instances are capable of contributing to collagen deposition as demonstrated in mouse and
zebrafish injury models (Simões et al., 2020). Gene-set enrichment analysis of the genes that are
most correlated with *in vitro*-derived macrophages moving away (Figure 6A) from the tissue
resident populations revealed that the most significant pathways in these cells involved collagen
synthesis and production (Table S6). A STRING protein-protein interaction network (Figure 6B)
shows that this phenotype is significantly enriched for highly connected matrix remodeling,
collagen deposition and cadherin-mediated cell-cell and cell-matrix interactions.
283

Perhaps pluripotent stem cell-derived cells are being driven to adopt a pro-fibrotic phenotype
through the derivation or culture contents. Initial observations on analysis of myeloid-, pluripotent
stem cell- and hematopoietic progenitor-derived cells, highlight higher expression of collagen
genes in pluripotent stem cell-derived cells (Figure 6C). To investigate whether this phenotype is
due to culture impact, we isolated peripheral blood monocytes and derived human PSC-
progenitors and cultured these cells with the same culture media in the presence of M-CSF for 5
days to drive macrophage differentiation. On day 5, cells were either stimulated with LPS for 2
hours before extraction or extracted as control samples for sequencing analysis. Analysis of
sequencing samples showed the same enriched collagen and cadherin networks with high
expression of these genes observed in PSC-macrophages, with little or no expression in
monocyte-derived macrophages regardless of whether cells were stimulated or not (Figure 6D).
To determine this at a protein level, we carried out intracellular flow cytometry analysis of type I
collagen in PSC- and monocyte-derived cells (Figure 6E). Compared with staining controls, we
observed higher levels of type I collagen in PSC-macrophages. Collectively, our results suggest
that although culture media can impact cell phenotype, it is not the main driver for this pro-fibrotic
phenotype in PSC-macrophages in the final stages of differentiation.
300

Discussion

Human macrophage biology is integral to the development of homeostasis and disease
mechanisms in every tissue in the body, but our understanding of human myeloid biology is limited
by the quality by the models available to us. Here we describe an integrated myeloid transcriptome
atlas as a novel resource to identify myeloid cells in single cell datasets and to benchmark *in vitro*
models of *in vivo* biology. Implementation in Stemformatics enables users to upload their own
data to benchmark cell types against the atlas for rapid and intuitive cell-classification. The

resource is scalable and will grow as the availability of new tissue resident samples and myeloid
models become available.
310
Transcriptional benchmarking of myeloid subsets, and particularly those benchmarking new
derivation methods, typically draw on a small number of reference samples. Standard analysis
workflows include normalisation methods that remove technical batch effects, and harmonise the
behaviour of samples assigned to the same class. When combining data that confounds technical
batch with the biology of interest, this process inevitably overstates similarities within groups and
over emphases differences between groups. The result is a somewhat self-fulfilling analysis, with
the outcome predicted by the data processing approach. Here, we offer an alternative for
benchmarking new models of myeloid biology against a reference atlas constructed from a large
number of well phenotyped and curated published data. We demonstrate reproducible
classification of major myeloid cell classes, including the influence of culture or derivation method
on the phenotype of those cells. Projection of external data into the atlas further demonstrates
that these phenotypic patterns are reproducible, even at the resolution of the single cell.
323
Our analyses highlight that there is room for improvement in the development of *in vitro* model
systems that attempt to mimic *in vivo* counterparts. We demonstrate that cord blood-derived
dendritic cells differentiated ex vivo from monocytes or CD34 progenitors don't adequately
capture key aspects of *in vivo* myeloid biology. Likewise, by benchmarking curated public data of
PSC-macrophages and their precursor cells against the atlas, it is apparent that these represent
neither definitive nor primitive myelopoiesis, or rather, that they imperfectly recapitulate aspects
of both. PSC-conditions clearly do not mimic the developmental time-frame nor tissue niche of
yolk-sac, fetal liver or bone marrow. PSC-macrophages do recapitulate many aspects of ex-vivo
cultured tissue macrophages, but there is little evidence for cultured microglia being distinct from
any other cultured macrophage. In our hands, PSC-macrophages display transcriptional
hallmarks of efferocytosis and surprisingly collagen production, which may suggest that the
derivation process of these cells are driving a pro-fibrotic phenotype. These observations offer
new opportunities to co-opt additional transcriptional networks that may arise from the tissue
niche, to improve stem cell derived myeloid models.
338

Acknowledgements
The authors thank Tyrone Chen and Othmar Korn for assistance with data processing and Isha
Nagpal for website development to support the Stemformatics interactive viewer. The authors
thank Ramaciotti Centre for Genomics (University of New South Wales; Sydney) and University
of Glasgow Polyomics facility for mRNA sequencing, and the Melbourne Cytometry Platform for
flow cytometry assistance. This work was funded by Stem Cells Australia, an Australian
Research Council Special Research Initiative [SRI110001002] to CAW; NHMRC Synergy
[APP1186371] to CAW; Wellcome Trust catalyst funding WELLCOME (097821/Z/11/B) to CAW,
SM, MKS; SM work was supported by the Research into Inflammatory Arthritis Centre Versus
Arthritis (RACE) [#20298]; MKS is funded by Versus Arthritis UK [#20298 & #22072]. NR is
funded by the Centre for Stem Cell Systems and the CSIRO Synthetic Biology Future Science
Platform. CAW is funded by a Future Fellowship from the Australian Research Council
[FT150100330]. JC is funded by the JEM Research Foundation to the Stem Cell Atlas.

Author Contributions

Conception NR, JC, CAW; Experimental Investigation and Interpretation NR, VJ, JG, CAW; Experimental Resources ALL, CAW; Methodology PWA, JC, YD; Data provider NR, MKS, SM; Curation NR, MR, CMP, CAW; Statistical analysis YD, KALC, PWA; Writing – original draft NR, CAW; Writing - review and editing NR, PWA, SM, ALL, KALC, JC, CAW; Supervision CAW, ALL, KAC; Project Funding - CAW.

Declaration of Interests

None

References

Abud, E.M., Ramirez, R.N., Martinez, E.S., Healy, L.M., Nguyen, C.H.H., Newman, S.A., Yeromin, A. V., Scarfone, V.M., Marsh, S.E., Fimbres, C., et al. (2017). iPSC-Derived Human Microglia-like Cells to Study Neurological Diseases. Neuron *94*, 278-293.e9.

Alasoo, K., Rodrigues, J., Mukhopadhyay, S., Knights, A.J., Mann, A.L., Kundu, K., Hale, C., Dougan, G., and Gaffney, D.J. (2018). Shared genetic effects on chromatin and gene expression indicate a role for enhancer priming in immune response. Nat. Genet. *50*, 424–431.

Angel, P.W., Rajab, N., Deng, Y., Pacheco, C.M., Chen, T., Le Cao, K.-A., Choi, J., and Wells, C. (2020). A simple, scalable approach to building a cross-platform transcriptome atlas. BioRxiv 2020.03.09.984468.

Bagger, F.O., Sasivarevic, D., Sohi, S.H., Laursen, L.G., Pundhir, S., Sønderby, C.K., Winther, O., Rapin, N., and Porse, B.T. (2016). BloodSpot: A database of gene expression profiles and transcriptional programs for healthy and malignant haematopoiesis. Nucleic Acids Res. *44*, D917–D924.

Buchrieser, J., James, W., and Moore, M.D. (2017). Human Induced Pluripotent Stem Cell-Derived Macrophages Share Ontogeny with MYB-Independent Tissue-Resident Macrophages. Stem Cell Reports *8*, 334–345.

Cao, X., Yakala, G.K., van den Hil, F.E., Cochrane, A., Mummery, C.L., and Orlova, V. V. (2019). Differentiation and Functional Comparison of Monocytes and Macrophages from hiPSCs with Peripheral Blood Derivatives. Stem Cell Reports *12*, 1282–1297.

Choi, J., Pacheco, C.M., Mosbergen, R., Korn, O., Chen, T., Nagpal, I., Englart, S., Angel, P.W., and Wells, C.A. (2019a). Stemformatics: visualize and download curated stem cell data. Nucleic Acids Res. *47*, D841–D846.

Choi, J., Baldwin, T.M., Wong, M., Bolden, J.E., Fairfax, K.A., Lucas, E.C., Cole, R., Biben, C., Morgan, C., Ramsay, K.A., et al. (2019b). Haemopedia RNA-seq: a database of gene expression during haematopoiesis in mice and humans. Nucleic Acids Res. *47*, D780–D785.

Dutertre, C.A., Becht, E., Irac, S.E., Khalilnezhad, A., Narang, V., Khalilnezhad, S., Ng, P.Y., van den Hoogen, L.L., Leong, J.Y., Lee, B., et al. (2019). Single-Cell Analysis of Human Mononuclear Phagocytes Reveals Subset-Defining Markers and Identifies Circulating Inflammatory Dendritic Cells. Immunity *51*, 573-589.e8.

Elliott, M.R., Koster, K.M., and Murphy, P.S. (2017). Efferocytosis Signaling in the Regulation of Macrophage Inflammatory Responses. J. Immunol. *198*, 1387–1394.

Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P., Haw, R.,

Jassal, B., Korninger, F., May, B., et al. (2018). The Reactome Pathway Knowledgebase. Nucleic Acids Res. *46*, D649–D655.

Gautier, E.L., Shay, T., Miller, J., Greter, M., Jakubzick, C., Ivanov, S., Helft, J., Chow, A., Elpek, K.G., Gordonov, S., et al. (2012). Gene-expression profiles and transcriptional regulatory pathways that underlie the identity and diversity of mouse tissue macrophages. Nat. Immunol. *13*, 1118–1128.

Gosselin, D., Skola, D., Coufal, N.G., Holtman, I.R., Schlachetzki, J.C.M., Sajti, E., Jaeger, B.N., O'Connor, C., Fitzpatrick, C., Pasillas, M.P., et al. (2017). An environment-dependent transcriptional network specifies human microglia identity. Science (80-.). *356*, 1248–1259.

Grant Rowe, R., Wang, L.D., Coma, S., Han, A., Mathieu, R., Pearson, D.S., Ross, S., Sousa, P., Nguyen, P.T., Rodriguez, A., et al. (2016). Developmental regulation of myeloerythroid progenitor function by the Lin28b-let-7-Hmga2 axis. J. Exp. Med. *213*, 1497–1512.

Hay, S.B., Ferchen, K., Chetal, K., Grimes, H.L., and Salomonis, N. (2018). The Human Cell Atlas bone marrow single-cell interactive web portal. Exp. Hematol. *68*, 51–61.

Honda-Ozaki, F., Terashima, M., Niwa, A., Saiki, N., Kawasaki, Y., Ito, H., Hotta, A., Nagahashi, A., Igura, K., Asaka, I., et al. (2018). Pluripotent Stem Cell Model of Nakajo-Nishimura Syndrome Untangles Proinflammatory Pathways Mediated by Oxidative Stress. Stem Cell Reports *10*, 1835–1850.

Huang, E., and Wells, C.A. (2014). Environmental Influences Interface of Genetic, Epigenetic, and Responsiveness Is Determined at the The Ground State of Innate Immune Downloaded from. J Immunol J. Immunol. J. Immunol. *193*, 13–19.

Jones, J.C., Sabatini, K., Liao, X., Tran, H.T., Lynch, C.L., Morey, R.E., Glenn-Pratola, V., Boscolo, F.S., Yang, Q., Parast, M.M., et al. (2013). Melanocytes derived from transgene-free human induced pluripotent stem cells. J. Invest. Dermatol. *133*, 2104–2108.

Joshi, K., Elso, C., Motazedian, A., Labonne, T., Schiesser, J. V., Cameron, F., Mannering, S.I., Elefanty, A.G., and Stanley, E.G. (2019). Induced pluripotent stem cell macrophages present antigen to proinsulin-specific T cell receptors from donor-matched islet-infiltrating T cells in type 1 diabetes. Diabetologia *62*, 2245–2251.

Kelly, A., Gunaltay, S., McEntee, C.P., Shuttleworth, E.E., Smedley, C., Houston, S.A., Fenton, T.M., Levison, S., Mann, E.R., and Travis, M.A. (2018). Human monocytes and macrophages regulate immune tolerance via integrin αvβ8-mediated TGFβ activation. J. Exp. Med. *215*, 2725–2736.

Kong, W., Fu, Y.C., and Morris, S.A. (2020). Capybara: A computational tool to measure cell identity and fate transitions. BioRxiv 2020.02.17.947390.

Lee, C.Z.W., Kozaki, T., and Ginhoux, F. (2018). Studying tissue macrophages in vitro: are iPSC-derived cells the answer? Nat. Rev. Immunol. *18*, 716–725.

Ng, E.S., Davis, R., Stanley, E.G., and Elefanty, A.G. (2008). A protocol describing the use of a recombinant protein-based, animal product-free medium (APEL) for human embryonic stem cell differentiation as spin embryoid bodies. Nat. Protoc. *3*, 768–776.

Rajab, N., Rutar, M., Laslett, A.L., and Wells, C.A. (2018). Designer macrophages: Pitfalls and opportunities for modelling macrophage phenotypes from pluripotent stem cells. Differentiation *104*, 42–49.

Regev, A., Teichmann, S.A., Lander, E.S., Amit, I., Benoist, C., Birney, E., Bodenmiller, B., Campbell, P., Carninci, P., Clatworthy, M., et al. (2017). The Human Cell Atlas. Elife *6*.

Schmidl, C., Renner, K., Peter, K., Eder, R., Lassmann, T., Balwierz, P.J., Itoh, M., Nagao-Sato,
S., Kawaji, H., Carninci, P., et al. (2014). Transcription and enhancer profiling in human
monocyte subsets. Blood *123*, e90–e99.

Schroder, K., Irvine, K.M., Taylor, M.S., Bokil, N.J., Le Cao, K.-A., Masterman, K.-A., Labzin,
L.I., Semple, C.A., Kapetanovic, R., Fairbairn, L., et al. (2012). Conservation and divergence in
Toll-like receptor 4-regulated gene expression in primary human versus mouse macrophages.
Proc. Natl. Acad. Sci. U. S. A. *109*, E944-53.

Shook, B.A., Wasko, R.R., Rivera-Gonzalez, G.C., Salazar-Gatzimas, E., López-Giráldez, F.,
Dash, B.C., Muñoz-Rojas, A.R., Aultman, K.D., Zwick, R.K., Lei, V., et al. (2018). Myofibroblast
proliferation and heterogeneity are supported by macrophages during skin repair. Science (80-.
). *362*, eaar2971.

Simões, F.C., Cahill, T.J., Kenyon, A., Gavriouchkina, D., Vieira, J.M., Sun, X., Pezzolla, D.,
Ravaud, C., Masmanian, E., Weinberger, M., et al. (2020). Macrophages directly contribute
collagen to scar formation during zebrafish heart regeneration and mouse heart repair. Nat.
Commun. *11*, 1–17.

Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M.,
Doncheva, N.T., Morris, J.H., Bork, P., et al. (2019). STRING v11: protein–protein association
networks with increased coverage, supporting functional discovery in genome-wide
experimental datasets. Nucleic Acids Res. *47*, D607–D613.

Turlach, B.A., and Weingessel, A. (2019). quadprog: Functions to Solve Quadratic
Programming Problems. R package version 1.5-8.

Vijayan, D., Radford, K.J., Beckhouse, A.G., Ashman, R.B., and Wells, C.A. (2012). Mincle
polarizes human monocyte and neutrophil responses to *Candida albicans*. Immunol. Cell Biol.
90, 889–895.

Vijayan, V., Pradhan, P., Braud, L., Fuchs, H.R., Gueler, F., Motterlini, R., Foresti, R., and
Immenschuh, S. (2019). Human and murine macrophages exhibit differential metabolic
responses to lipopolysaccharide - A divergent role for glycolysis. Redox Biol. *22*, 101147.

Villani, A.C., Satija, R., Reynolds, G., Sarkizova, S., Shekhar, K., Fletcher, J., Griesbeck, M.,
Butler, A., Zheng, S., Lazo, S., et al. (2017). Single-cell RNA-seq reveals new types of human
blood dendritic cells, monocytes, and progenitors. Science (80-.). *356*.

Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski,
E., Peterson, P., Weckesser, W., Bright, J., et al. (2019). SciPy 1.0--Fundamental Algorithms for
Scientific Computing in Python. Nat. Methods *17*, 261–272.

Vlahos, K., Sourris, K., Mayberry, R., McDonald, P., Bruveris, F.F., Schiesser, J. V., Bozaoglu,
K., Lockhart, P.J., Stanley, E.G., and Elefanty, A.G. (2019). Generation of iPSC lines from
peripheral blood mononuclear cells from 5 healthy adults. Stem Cell Res. *34*, 101380.

Wakioka, T., Sasaki, A., Kato, R., Shouda, T., Matsumoto, A., Miyoshi, K., Tsuneoka, M.,
Komiya, S., Baron, R., and Yoshimura, A. (2001). Spred is a Sprouty-related suppressor of Ras
signalling. Nature *412*, 647–651.

Watkins, N.A., Gusnanto, A., de Bono, B., De, S., Miranda-Saavedra, D., Hardie, D.L.,
Angenent, W.G.J., Attwood, A.P., Ellis, P.D., Erber, W., et al. (2009). A HaemAtlas:
characterizing gene expression in differentiated human blood cells. Blood *113*, e1-9.

Yamaguchi, H., Maruyama, T., Urade, Y., and Nagata, S. (2014). Immunosuppression via
adenosine receptor activation by adenosine monophosphate released from apoptotic cells. Elife
3.

Yurchenko, M., Skjesol, A., Ryan, L., Richard, G.M., Kandasamy, R.K., Wang, N., Terhorst, C.,
Husebye, H., and Espevik, T. (2018). SLAMF1 is required for TLR4-mediated TRAM-TRIF–
dependent signaling in human macrophages. J. Cell Biol. *217*, 1411–1429.

Zhou, J., Chan, Z.L., Bi, C., Lu, X., Chong, P.S.Y., Chooi, J.Y., Cheong, L.L., Liu, S.C., Ching,
Y.Q., Zhou, Y., et al. (2017). LIN28B activation by PRL-3 promotes leukemogenesis and a stem
cell-like transcriptional program in AML. Mol. Cancer Res. *15*, 294–303.

Figure Titles and Legends

Figure 1: A reference atlas for human myeloid biology
(A) Stemformatics myeloid atlas with samples coloured by cell type. Navy blue - monocytes,
blue - macrophages, aqua -Dendritic cells, dark green - CD141+ DC, light green - CD1c+ DC,
yellow - pDC, brown – granulocytes, pink stem and progenitor cells, hemogenic endothelium.
Validation with Haemopedia RNAseq myeloid samples: diamond shape – monocytes, circle
granulocytes, cross DC. (B) Atlas coloured by ranked expression of TMEM1. (Scale bar: high
ranked expression (dark red) to low ranked expression (grey) indicating axis of activation) (C)
Atlas coloured by culture status (sample source). See also Table 1, Figure S1, and Table S1.

Figure 2: Cultured and *in vitro*-derived dendritic cells do not capture aspects *in vivo*
myeloid biology
(A) DC subsets displayed in the atlas – aqua *in vitro* (cord blood derived) DC, dark green DC1
(CD141+), light green DC2 (CD1c+), yellow - pDC, brown neutrophils (B) atlas coloured by ranked
expression of TLR7. (Scale bar: high ranked expression (dark red) to low ranked expression
(grey)). (C) Ranked expression (Y-axis) of receptor CX3CR1 and cell-fusion protein DC-STAMP
in vivo dendritic cells (n=145), *ex vivo* dendritic cells (n=17) and *in vitro*-derived dendritic cells
(n=57). Grey stripe indicates variance attributable to platform. P-value: Mann-Whitney-Wilcoxon
rank sum test. (D) Single cell projections of Villani *et al.* (2017) and Dutertre *et al.* (2019) samples
onto atlas (F) Heatmap derived from Capybara analysis of Villani *et al.* (2017) and Dutertre *et al.*
(2019) samples compared to myeloid cell types. Colour gradients reflect similarity of single cell
clusters to atlas cell types (dark least similar, to light most similar). See also Figure S2 and Table
S2.

Figure 3: Monocytes acquire a culture phenotype
(A) Schematic of rolling monocytes, highlighting cultured cells mimic many of the features of a
monocyte after extravasation (B) Cultured monocytes form a distinct cluster away from *in vivo*
monocytes along PC3 (C) STRING-DB network of top-ranked genes differentially expressed
between peripheral blood (*in vivo*, n=107) and cultured monocytes (*ex vivo*, n=171) indicating
upregulation of cytoskeletal proteins and down regulation of endothelial-adhesion proteins (D)
Ranked expression (Y-axis) of gene involved in endothelial adhesion, SELL, comparing cultured
monocytes (n=171) with monocytes directly profiled from blood (*in vivo*, n=107). Grey stripe
indicates variance attributable to platform. P-value: Mann-Whitney-Wilcoxon rank sum test
(E)Ranked expression (Y-axis) of gene involved in the regulation of RAS/RAF signaling
comparing cultured monocytes (n=171) with monocytes directly profiled from blood (*in vivo*,
n=107). Grey stripe indicates variance attributable to platform. P-value: Mann-Whitney-Wilcoxon

rank sum test (F) atlas coloured by ranked expression of IL-6 (Scale bar: high ranked
expression (dark red) to low ranked expression (grey) indicating axis of activation) (G) Ranked
expression (Y-axis) of Chemokine CCL2 comparing cultured monocytes (n=171) with
monocytes directly profiled from blood (*in vivo*, n=107). Grey stripe indicates variance
attributable to platform. P-value: Mann-Whitney-Wilcoxon rank sum test. See also Table S3.
534
Figure 4: Pluripotent stem cell-derived macrophages do not recapitulate hematopoietic
ontogenies
(A) atlas coloured by ranked expression of MYB (Scale bar: high ranked expression (dark red) to
low ranked expression (grey). (B) Ranked expression (Y-axis) of MAF expression from *in vivo*
(n=61), *ex vivo* (n=26) and *in vitro-* (n=96) derived macrophages (gut, synovial, kupffer, microglia,
macrophage). Grey stripe indicates variance attributable to platform. (C) Ranked expression (Y-
axis) of LIN28B expression from *in vivo* (n=61), *ex vivo* (n=26) and *in vitro-* (n=96) derived
macrophages (gut, synovial, kupffer, microglia, macrophage). Grey stripe indicates variance
attributable to platform. P-value MannWhitney-Wilcoxon rank sum test. (D) atlas with samples
coloured by cell type with projection of iPSC samples highlighting their position in comparison to
in vitro-derived macrophages, hematopoietic multipotent progenitors and hemangioblast. (E)
Single cell projection of (Bian et al., 2020) human fetal yolk sac cell clusters onto the atlas
following an 8 cell aggregation. See also Figure S3 and Table S4 and Table S5.
548
Figure 5: Pluripotent stem cell-derived macrophages display transcriptional hallmarks of
efferocytosis
(A)Schematic of impact of efferocytosis on cell metabolic reprogramming and function (B) atlas
coloured by ranked expression of MERTK (scale bar: high ranked expression (dark red) to low
ranked expression (grey). (C,D,E) Ranked expression (Y-axis) of genes comparing *in vivo* (n=61),
ex vivo (n=26) and *in vitro*-derived (n=96) macrophages (gut, synovial, kupffer, microglia,
macrophage) for (C) cholesterol efflux (D) mitochondrial acyl-CoA dehydrogenase (E)
phosphatidate phosphatase. See also Table S4 and Table S5.
557
Figure 6: Pluripotent stem cell-derived macrophages express high levels of collagen
(A) atlas coloured by cell source to highlight *in vitro*-derived macrophages used for regression
testing (B) STRING_DB Protein-Protein network of *in vitro*-derived macrophages highlights
enrichment of collagen, growth factor and cadherin networks. Line color indicates the type of
interaction evidence. Light blue solid lines indicate known interactions from curated databases,
solid pink line indicates known interactions that have been experimentally determined, bright
green lines indicate gene neighborhood predicted interactions, red lines indicate gene fusions
predicted interactions, dark blue lines indicate gene co-occurrence predicted interactions,
yellow/green lines indicate textmining, black lines indicate co-expression, and light purple lines
indicate protein homology. (C) Violin plots of myeloid- (n=584), pluripotent stem cell- (n=116) and
hematopoietic progenitor- (n=201) derived cells comparing expression of collagen genes
(COL1A1 and COL4A2)(D) mRNA-seq gene expression from human peripheral blood monocyte-
derived macrophages (HMDM) (n=3) and human pluripotent stem cell derived macrophages
(PSCM) (n=2) samples (C= control, S = stimulated) (E) Intracellular flow cytometry analysis of
human peripheral blood monocyte-derived macrophages (HMDM) and human pluripotent stem

cell derived macrophages (PSCM), representative of 2 experimental repeats (n=2 HMDM, n=2
PSCM). Red = no primary antibody control, Black = isotype control, Purple = Type I Collagen
stain. See also Table S6.
576
577   **Supplemental Figures**
578
Figure S1: A reference atlas and resource for human myeloid biology. Related to Figure
1.
(A) Pre-gene filtering (left) and Post-gene filtering (right) atlas coloured by platform: red various
microarray platforms, black RNAseq platforms. (B) atlas coloured by ranked expression of
TMEM119 (Scale bar: high ranked expression (dark red) to low ranked expression (grey) (C) atlas
coloured by ranked expression of P2RY12 (Scale bar: high ranked expression (dark red) to low
ranked expression (grey) (D) Ranked expression (Y-axis) of Class II transactivator (CIITA) *in vivo*
versus *in vitro*-derived macrophages (gut, synovial, kupffer, microglia, macrophage). Red –
activated, yellow – (Abud et al., 2017) microglia samples, khaki- Honda-Ozaki et al. (2018)
macrophage samples.
589
Figure S2: Single cell aggregation and projection. Related to Figure 2.
(A) Kolmogorov–Smirnov (KS) statistics (y-axis) to assess the difference in gene expression
distribution between pseudo-bulk single cells DC6 and bulk sample plasmacytoid dendritic cells
from the atlas, with respect to the number of single cells that are aggregated (x-axis). Each line
indicates one of thirty random sub-samplings with replacement trial. KS statistics are calculated
on each gene and averaged across all genes. A minimum KS statistic is obtained when
aggregating 8 cells. (B) atlas cell types before single cell projection (C) Single cell projection of
(Dutertre et al., 2019) and (Villani et al., 2017) samples onto the atlas where a 8 cells are
aggregated based on (A).
599
600   **Figure S3: Fetal Ontogeny. Related to Figure 4.**
601   atlas coloured by cell type with (Bian et al., 2020) projection of single cell data from human fetal
602   yolk sac and head.
603
604   **Tables with Titles and Legends**
605
606   **Table 1: Atlas inclusion of tissue-resident monocytes and macrophages**

Tissue	Tier 1 (sample #)	Tier 2	Cell Type
Blood	*in vivo* (64) *ex vivo* (162)	myeloid	monocyte
Cord Blood	*ex vivo* (51)	myeloid	monocyte
Gut	*in vivo (15)*	myeloid	macrophage
Synovium	*in vivo (18)*	myeloid	macrophage
Brain	*in vivo* (10) *ex vivo* (21)	myeloid	microglia
Lung	*in vivo (14)*	myeloid	macrophage
Liver	*ex vivo* (5)	myeloid	Kupffer cell
Ovarian Tumour	*in vivo* (4)	myeloid	macrophage

607
608 **Supplemental Tables**
609
Table S1: Datasets and samples to compile atlas and single cell projection. Related to
Figure 1.
Tissue resident macrophages and dendritic cells from peripheral blood, spleen, thymus, joint,
lung, gut, brain and liver. Samples also included monocytes from peripheral and cord blood, as
well as *in vitro* differentiated DCs from cord blood progenitors or monocyte-derived
macrophages. Columns include dataset accession ID, platform, Stemformatics Dataset ID,
number of samples, tier categorization, cell type and relevant tissue/organism part.
617
618 **Table S2: *in vivo* vs. *ex vivo* vs. *in vitro* dendritic cells. Related to Figure 2.**
619 Comparison of gene expression of *in vivo* (n=145), *ex vivo* (n=17) and *in vitro*- (n=57) derived
620 dendritic cells. Columns refer to gene symbols, P-values re-calculated by Mann-
621 WhitneyWilcoxon rank-sum test, mean and standard deviation.
622
623 **Table S3: *in vivo* vs. *ex vivo* monocytes. Related to Figure 3.**
624 Comparison of gene expression of *ex vivo* (n=171) and *in vivo* (n=107) monocytes. Columns
625 refer to gene symbols, P-values re-calculated by Mann-Whitney-Wilcoxon rank-sum test, mean
626 and standard deviation.
627
628 **Table S4: *in vivo* vs. *ex vivo* vs. *in vitro* macrophages. Related to Figure 4 and Figure 5.**
629 Comparison of gene expression of *in vivo* (n=61), *ex vivo* (n=26), *in vitro*- (n=96) derived
630 macrophages (gut, synovial, kupffer, microglia, macrophage). Columns refer to gene symbols,
631 P-values recalculated by Mann-Whitney-Wilcoxon rank-sum test, mean and standard deviation.
632
633 **Table S5: *in vivo* vs. *ex vivo* vs. *in vitro* microglia. Related to Figure 4 and Figure 5.**
634 Comparison of gene expression of in vivo (n=10), ex vivo (n=21) and in vitro- (n=43) derived
635 microglia. Columns refer to gene symbols, P-values re-calculated by Mann-Whitney-Wilcoxon
636 rank-sum test, mean and standard deviation.
637
638 **Table S6: Gene-Set Enrichment Analysis. Related to Figure 6.**
639 Table of the top 10 Reactome pathways enriched in genes highly correlated with *in vitro*-derived
640 macrophage spread. Enrichment: number of genes in the list/number of genes in that pathway
641 (False Discovery Rate-value). Genes: multiple entries assigned to the same gene indicated by
642 underlining of gene symbol with UniProt accessions in brackets.
643
644 **Supplemental Video**
645 Active engulfment and clearance of cells by pluripotent stem cell-derived macrophages.
646
Methods
Atlas formation was developed as described in (Angel et al., 2020) and is comprised of 44
datasets, 901 samples and 3757 genes. Mapping, and analysis of microarray and RNA
sequencing datasets were conducted in the Stemformatics platform. Scripts are available for

download from the Stemformatics BitBucket (Choi et al., 2019a)). All datasets and relevant
samples (Supplementary Table 6) passed stringent quality control checks required for hosting
on the stemformatics platform. Quality control checks include evaluation of library quality, and
inclusion of replicates associated with experimental design. These datasets were either already
hosted on stemformatics, or were downloaded from public depositories and processed through
the stemformatics pipeline for inclusion.
657
Platform Effect Analysis and Gene Selection for PCA
This method assesses each gene independently for a dependence upon experimental platform.
The initial step is to transform expression values from RNA Sequencing and Microarray into
percentile values. The second step uses a univariate estimate of gene platform dependence
and then selects genes with a small ratio of platform dependent variance to total variance.
These genes are used to run the PCA (3757 genes passed this cut). The threshold for gene
selection is empirically determined by using the Kruskal Wallis H Test to assess the difference
in platform expression distribution for each principal component. A platform variance fraction of
0.2 is found to remove the platform effect for the first three principal component. For further
details, please refer to (Angel et al., 2020).
668
669 Quantification and Statistical Analysis
670 P-values were re-calculated using the Mann-Whitney-Wilcoxon rank-sum test (two-sided). This
671 was implemented via the python (version 3.7.5) SciPy package (version 1.3.1)(Virtanen et al.,
672 2019)Multiple testing over the set of genes was accounted for with Bonferroni correction
673 implemented in the statsmodels package (Seabold and Perktold, 2010)
674
Pseudo-bulk samples from Villani's Single Cell Data
Single cells were aggregated to form pseudo-bulk samples to mitigate library size differences
between single cell and bulk data, and to project samples onto the atlas. Each group of (known)
cell type in the single cell data was randomly sampled for k single cells with replacement.
Aggregation consisted in summing up their expression profiles. k was determined by the
number of sampling (i.e. how many pseud-bulk samples for each cell type) as half of the cell
type's population size. The optimum aggregation size k was investigated by evaluating the
similarity in distribution between the atlas' plasmacytoid dendritic cells and the aggregated
pseudo-bulk DC6 samples using Kolmogorov–Smirnov (KS) statistic D averaged across all
genes. KS statistic measured the difference between the empirical cumulative distribution
functions of two groups of samples; the smaller the value, the closer the aggregated DC6
resembled the reference transcriptional profiles from the pDC in the bulk atlas. A minimum D
value was obtained for $k = 8$ across 30 iterations (Supplementary Figure 1). Similar results were
obtained for other cell types. Thus, every pseudo-bulk samples were aggregated from 8 single
cells.
690
Capybara Cell Score
Capybara (Kong et al., 2020)cell scores was used to measure cell identities continuum of the
pseudo-bulk samples using the atlas as the reference. Capybara cell scores were calculated by
performing restricted linear regression of reference samples on each of the pseudo-bulk

samples' expression profiles. Denote y_i the expression profile of the i^{th} pseudo-bulk sample of
length G, where G represents the total number of genes in the data, and X a (GXT) the
reference matrix, where T represents the number of known cell types of interest. We considered
5 cell types: Dendritic cell, Monocyte, CD141+ dendritic cell, CD1c+ dendritic cell and
Plasmacytoid dendritic cells. X is obtained by averaging the expression profiles of the
Stemformatics myeloid samples according to their cell types. Capybara solves the optimization
problem

$$argmin_\beta \ (y_i - X\beta_i)^T(y_i - X\beta_i)$$

under the constraint that

$$\beta_{it} > 0 \ \ \forall t \in \{1, 2, ..., T\}, \ \sum_t^T \beta_{it} < 1$$

where β_{it} is a regression coefficient, or cell score, for each pseudo-bulk sample i and each atlas
cell type t. The cell score is obtained using quadratic programming implemented with R (version
3.6.2) package *quadprog (version 1.5-8)*(Turlach and Weingessel, 2019).

<u>Enrichment analysis and Protein-Protein Network</u>
An enrichment analysis was conducted on the top 92 genes ranked by Pearson correlation
(≥0.7) along the upward axis including in vitro-derived cells. Enriched pathways were identified
using these genes at Reactome (Fabregat et al., 2018) and significance ranked by p-value/false
discovery rate. A protein-protein network was generated using the top 92 genes on STRING-DB
(Szklarczyk et al., 2019). Disconnected nodes not shown.

<u>Pluripotent Stem Cell Differentiation</u>
Stem cell work was carried out in accordance with The University of Melbourne ethics
committee HREC (approval 1851831). Stem cell lines used were PB001.1 (Vlahos et al., 2019),
a kind gift from the Stem Cell Core Facility at the Murdoch Children's Research Institute, and
HDF51((Jones et al., 2013); RRID:CVCL_UF42) was kindly provided to ALL by Prof. Jeanne
Loring (The Scripps Research Institute, CA, USA). Human pluripotent stem cells were
differentiated into macrophages based on protocol described by ((Joshi et al., 2019; Ng et al.,
2008)), with modifications. Modifications were as follows: embryoid bodies were kept in
rotational cultures without transference to matrigel plates for adherence, and the collection of
progenitors from week 2 were immediately re-suspended in RPMI-1640 containing L-
Glutamine (Life Technologies) and 10% Fetal Bovine Serum for macrophage differentiation (see
macrophage differentiation).

<u>Monocyte isolation from peripheral blood</u>
Buffy Coat was obtained from the Australian Red Cross Blood Service in accordance with The
University of Melbourne ethics committee HREC (approval 1646608). The blood was diluted with
PBS at a 1:3 dilution and underlayed with Ficoll-Hypaque. The underlayed blood samples were
centrifuged at 350g for 30 minutes at 24°C with no brake. Peripheral blood mononuclear cells
were isolated from the interphase and washed twice by using MACs buffer (DPBS, 0.5% heat
inactivated Fetal Bovine Serum, 2mM EDTA) and centrifuging at 400g for 5 minutes at 4°C. Cells
were centrifuged at 400g for 5 minutes at 4°C and resuspended in 40µl MACs buffer per 107 cells.
Monocytes were positively selected by a magnetic field using Human
CD14 MicroBeads (MACS Miltenyi Biotec) and LS Columns (Miltenybiotec). These cells were
plated for macrophage differentiation (see Macrophage differentiation

Macrophage differentiation

Monocytes/progenitor cells were cultured in tissue-culture treated 6 well plates. Cells were
cultured in RPMI-1640 medium containing L-Glutamine (Life Technologies) with 10% Fetal
Bovine Serum and 100ng/ml recombinant Human M-CSF (R&D Systems; 216-MC) for 5 days.
Media changes were carried out on day 4.

Flow Cytometry

HMDM and PSCM were collected and centrifuged at 400g for 5 minutes. Supernatant was
aspirated and 5µl mouse serum was added to 'dry' pellets for 5 minutes on ice. Cells were
resuspended in FACS Buffer (Hanks Balanced Salt Solution, 0.5% Human Serum Albumin) and
stained with CD14 or matched isotype control antibodies on ice for 20 minutes then washed twice
(3ml FACS Buffer, spun at 400g, 5 minutes). Resuspended cells were fixed with 4%
paraformaldehyde (PFA) for 15 minutes at room temperature. PFA was washed out and cells
rinsed twice in FACS buffer before resuspending in PBS and stored overnight at 4°C. Fixed cells
were permeabilized with 0.1% Triton X-100 (in 1XPBS) for 10 minutes at room temperature and
washed twice. Blocking buffer (0.3M glycine buffer, 10% Goat Serum, 1XPBS) was added to 'dry'
pellets on ice for 1 hour. Cells were stained with antibodies to Type I Collagen or matched isotype
control on ice for 20 minutes. Cells were washed twice then incubated with secondary antibody
for 20 minutes on ice in the dark. Cells were then washed twice and resuspended for analysis on
a CytoFLEX S flow cytometer (Beckman Coulter, Brea, CA) using CytExpert acquisition software.
Post-acquisition analysis was performed with FCS Express 7 flow cytometry software.

Flow Cytometry antibodies	
CD14	Brilliant Violet 421™ anti-Human CD14 (BioLegend: Cat. No. 325628; RRID:AB_2563296)
CD14 Isotype	Brilliant Violet 421™ Mouse IgG1 (BioLegend: Cat. No. 400157; RRID:AB_10897939)
Type I Collagen Primary antibody	Rabbit Anti-Collagen I antibody (Abcam: Cat. No. ab264074)
Secondary Antibody	Goat Anti-Rabbit IgG H&L (Alexa Fluor®488) (Abcam: Cat. No. ab150077; RRID:AB_2630356)
Type I Collagen Isotype	Rabbit IgG monoclonal Isotype Control (Abcam: Cat. No. Ab172730; RRID:AB_2687931)

Stimulation Assay

On day 5 of differentiation, one well containing peripheral blood monocyte- or human pluripotent
stem cell-derived macrophages were stimulated with 10ng/ml Lipopolysaccharide (LPS) (Sigma-
Aldrich; *Salmonella enterica* serotype minnesota) for 2 hours. After stimulation period, media was
aspirated, and the wells were washed twice with PBS (Ca2+Mg2+ free). before cell lysis using 2-
mercaptoethanol (Sigma-Aldrich) and RNeasy Plus Lysis Buffer (Qiagen). Samples were placed
into Eppendorf's and stored at -80°C before RNA extraction.

RNA extraction

Total RNA was isolated using the RNeasyⓇ Plus Mini Kit (Qiagen) according to manufacturer's
instructions. In summary: for the removal of genomic DNA, samples were placed into gDNA
columns and centrifuged for 30 seconds at 8000g. Ethanol (70%) was mixed with the flow through
and samples were transferred to RNeasy spin columns. The columns were centrifuged for 15

seconds at 8000g. Buffer RW1 was then added to the columns and columns were centrifuged for
15 seconds at 8000g. Buffer RPE was added to the columns and columns were centrifuged for
15 seconds. Buffer RPE was again added to the columns with centrifugation at 8000g for 2
minutes. Columns were then placed into new collection tubes and centrifuged at full speed for 1
minute to dry the membrane. RNase-free water was then added directly onto the column
membrane and columns placed into Eppendorf's and centrifuged at 8000g for 1 minute to collect
RNA. RNA quality and quantity were determined using a Tapestation (Agilent Technologies
2200). Samples were stored at -80°C. Zymo Research RNA Clean & Concentrator-25 Kit was
used to pool replicates (from the same donor) together and elute in elute into smaller volume with
maximum concentration.
787
788   RNA sequencing
789   RNA samples were processed by the Ramaciotti Centre for Genomics (University of New South
790   Wales; Sydney). Illumina Novaseq_6000 was used for mRNA-sequencing.
791
792   Data and Code Availability
793   mRNA-sequencing data is available through accession GSE150893. All public accessions are
794   listed in Supplementary Table 6. Stemformatics code is publicly available
795   at bitbucket.org/stemformatics. Atlas code is available
796   at bitbucket.org/stemformatics/s4m_pyramid/src/master/scripts/atlas.py.
797
798   Graphing software and Illustration
799   Graphs for mRNA-seq gene expression were generated using Graphpad Prism. Violin plots
800   were generated through the www.stemformatics.org platform. Schematic Figure illustrations
801   created with BioRender.com

(A) Coloured by cell type

PC3

Atlas samples
- HSC Progenitors
- Endothelium
- Macrophages
- Monocytes
- Neutrophils
- Dendritic Cells

Projected samples
- ◇ Monocyte
- O Neutrophil
- + Dendritic Cell

PC2

PC1

(B) Coloured by gene expression

PC3

alveolar macrophages

14 week fetal microglia

cultured adult microglia

ex vivo monocytes

gut macrophages

TREM1
0.9
0.8
0.7
0.6
0.5
0.4
0.3
0.2
0.1

PC2

PC1

(C) Coloured by culture status

Sample source
- *in vivo*
- *ex vivo*
- *in vitro*

PC3

MDM

TRM
ex vivo monocytes
TRM

in vivo monocytes

PC2

PC1

Figure 1: A reference atlas for human myeloid biology

Figure 2: Cultured and *in vitro*-derived dendritic cells do not capture aspects of *in vivo* myeloid biology

Figure 3: Monocytes rapidly adapt to culture

Figure 4: Pluripotent stem cell-derived macrophages do not recapitulate hematopoietic ontogenies

(A)

apoptotic
cells

efferocytosis

e.g.
S1PR1
MerTK

uptake of
apoptotic bodies

metabolic reprogramming

lipid metabolism
fatty acid synthesis
efflux of cholesterol
lipid turnover

functional impact

inflammatory   resolving

(B)

fetal microglia

PSC-
macrophages

MerTK

0.9
0.8
0.7
0.6
0.5
0.4
0.3
0.2
0.1

Kupffer
cells

alveolar
macrophages

gut macrophages

(C)

ABCA6

P=2.22E-16

P=0.58

P=3.83E-05

in vivo   *ex vivo*   *in vitro*

(D)

ACADM

P=5.21E-18

P=1.20E-07

in vivo   *ex vivo*   *in vitro*

(E)

LPIN3

P=4.13E-13

P=2.38E-05

in vivo   *ex vivo*   *in vitro*

Figure 5: Pluripotent stem cell-derived macrophages display transcriptional hallmarks of efferocytosis

Figure 6: Pluripotent stem cell-derived macrophages express high levels of collagen

Figure S1: A reference atlas and resource for human myeloid biology. Related to Figure 1.

(A) Pre-gene filtering (left) and Post-gene filtering (right) iMAC atlas coloured by platform: red various microarray platforms, black RNAseq platforms. (B) iMAC atlas coloured by ranked expression of TMEM119 (Scale bar: high ranked expression (dark red) to low ranked expression (grey) (C) iMAC atlas coloured by ranked expression of P2RY12 (Scale bar: high ranked expression (dark red) to low ranked expression (grey) (D) Ranked expression (Y-axis) of Class II transactivator (CIITA) *in vivo* versus *in vitro*-derived macrophages (gut, synovial, kupffer, microglia, macrophage). Red – activated, yellow – (Abud et al., 2017) microglia samples, khaki- Honda-Ozaki *et al.* (2018) macrophage samples.

Figure S2: Single cell aggregation and projection. Related to Figure 2.

(A) Kolmogorov–Smirnov (KS) statistics (y-axis) to assess the difference in gene expression distribution between pseudo-bulk single cells DC6 and bulk sample plasmacytoid dendritic cells from the atlas, with respect to the number of single cells that are aggregated (x-axis). Each line indicates one of thirty random sub-samplings with replacement trial. KS statistics are calculated on each gene and averaged across all genes. A minimum KS statistic is obtained when aggregating 8 cells. (B) iMAC atlas cell types before single cell projection (C) Single cell projection of (Dutertre et al., 2019) and (Villani et al., 2017) samples onto the iMAC atlas where a 8 cells are aggregated based on (A).

Figure S3: Fetal Ontogeny. Related to Figure 4.

iMAC atlas coloured by cell type with (Bian et al., 2020) projection of single cell data from human fetal yolk sac and head.

Supplemental Tables

Table S1: Datasets and samples to compile iMAC atlas and single cell projection. Related to Figure 1.
Tissue resident macrophages and dendritic cells from peripheral blood, spleen, thymus, joint, lung, gut, brain and liver. Samples also included monocytes from peripheral and cord blood, as well as *in vitro* differentiated DCs from cord blood progenitors or monocyte-derived macrophages. Columns include dataset accession ID, platform, Stemformatics Dataset ID, number of samples, tier categorization, cell type and relevant tissue/organism part.

Table S2: dendritic cells. Related to Figure 2.
Comparison of gene expression of *in vivo* (n=145), *ex vivo* (n=17) and *in vitro*- (n=57) derived dendritic cells. Columns refer to gene symbols, P-values re-calculated by Mann-WhitneyWilcoxon rank-sum test, mean and standard deviation.

Table S3: *in vivo* vs. *ex vivo* monocytes. Related to Figure 3.
Comparison of gene expression of *ex vivo* (n=171) and *in vivo* (n=107) monocytes. Columns refer to gene symbols, P-values re-calculated by Mann-Whitney-Wilcoxon rank-sum test, mean and standard deviation.

Table S4: *in vivo* vs. *ex vivo* vs. *in vitro* macrophages. Related to Figure 4 and Figure 5.
Comparison of gene expression of *in vivo* (n=61), *ex vivo* (n=26), *in vitro*- (n=96) derived macrophages (gut, synovial, kupffer, microglia, macrophage). Columns refer to gene symbols, P-values recalculated by Mann-Whitney-Wilcoxon rank-sum test, mean and standard deviation.

Table S5: *in vivo* vs. *ex vivo* vs. *in vitro* microglia. Related to Figure 4 and Figure 5.
Comparison of gene expression of in vivo (n=10), ex vivo (n=21) and in vitro- (n=43) derived microglia. Columns refer to gene symbols, P-values re-calculated by Mann-Whitney-Wilcoxon rank-sum test, mean and standard deviation.

Table S6: Gene-Set Enrichment Analysis. Related to Figure 6.
Table of the top 10 Reactome pathways enriched in genes highly correlated with *in vitro*-derived macrophage spread. Enrichment: number of genes in the list/number of genes in that pathway (False Discovery Rate-value). Genes: multiple entries assigned to the same gene indicated by underlining of gene symbol with UniProt accessions in brackets.

Supplemental Video
Active engulfment and clearance of cells by pluripotent stem cell-derived macrophages.

Table S6: Gene-Set Enrichment Analysis

<table>
<tr><td colspan="3" align="center">Table 1</td></tr>
<tr><td align="center">Pathway</td><td align="center">Enrichment</td><td align="center">Genes</td></tr>
<tr>
<td>Collagen biosynthesis and modifying enzymes</td>
<td>11/76 (3.76e-09)</td>
<td>ADAMTS3, COL1A2, COL4A2, SERPINH1, COL14A1, COL3A1, COL4A5, COL1A1, <u>COL4A1</u> <i>(P02462, Q03692)</i>, COL5A1</td>
</tr>
<tr>
<td>Extracellular matrix organization</td>
<td>18/329 (4.48e-09)</td>
<td>ADAMTS1, COL1A1, <u>COL4A1</u><i>(P02462, Q03692)</i>, COL5A1, LTBP1, PTPRS, ADAMTS3, COL1A2, COL4A2, KDR, LUM, SERPINH1, COL14A1, COL3A1, COL4A5, LAMB1, MFAP4</td>
</tr>
<tr>
<td>Collagen chain trimerization</td>
<td>9/44 (4.81e09)</td>
<td>COL14A1, COL3A1, COL4A5, COL1A1, <u>COL4A1</u><i>(P02462, Q03692)</i>, COL5A1, COL1A2, COL4A2</td>
</tr>
<tr>
<td>Collagen formation</td>
<td>11/104 (2.39e-08)</td>
<td>ADAMTS3, COL1A2, COL4A2, SERPINH1, COL14A1, COL3A1, COL4A5, COL1A1, <u>COL4A1</u><i>(P02462, Q03692)</i>, COL5A1</td>
</tr>
<tr>
<td>ECM proteoglycans</td>
<td>10/79 (2.39e-08)</td>
<td>COL1A1, COL4A1, COL5A1, PTPRS, COL1A2, COL4A2, LAMB1, COL3A1, COL4A5, LUM</td>
</tr>
<tr>
<td>Non-integrin membrane-ECM interactions</td>
<td>9/61 (3.83e-08)</td>
<td>COL1A1, <u>COL4A1</u><i>(P02462, Q03692)</i>, COL5A1, COL1A2, COL4A2, LAMB1, COL3A1, COL4A5</td>
</tr>
<tr>
<td>Integrin cell surface interactions</td>
<td>10/86 (3.83e-08)</td>
<td>COL1A1, <u>COL4A1</u><i>(P02462, Q03692)</i>, COL5A1, COL1A2, COL4A2, KDR, COL3A1, COL4A5, LUM</td>
</tr>
<tr>
<td>Assembly of collagen fibrils and other multimeric structures</td>
<td>9/67 (6.95e-08)</td>
<td>COL14A1, COL3A1, COL4A5, COL1A1, <u>COL4A1</u><i>(P02462, Q03692)</i>, COL5A1, COL1A2, COL4A2</td>
</tr>
<tr>
<td>Collagen degradation</td>
<td>9/69 (7.93e-08)</td>
<td>COL14A1, COL3A1, COL4A5, COL1A1, <u>COL4A1</u><i>(P02462, Q03692)</i>, COL5A1, COL1A2, COL4A2</td>
</tr>
<tr>
<td>Degradation of the extracellular matrix</td>
<td>11/148 (3.73e-07)</td>
<td>ADAMTS1, COL1A2, COL4A2, LAMB1, COL14A1, COL3A1, COL4A5, COL1A1, <u>COL4A1</u><i>(P02462, Q03692)</i>, COL5A1</td>
</tr>
</table>

Table of the top 10 Reactome pathways enriched in genes highly correlated with *in vitro*-derived macrophage spread. Enrichment: number of genes in the list/number of genes in that pathway (False Discovery Rate-value). Genes: multiple entries assigned to the same gene indicated by underlining of gene symbol with UniProt accessions in brackets.