Pseudomonas **can survive bacteriocin-mediated killing via a persistence-like mechanism**

Kandel PP[1], David A. Baltrus[2], Kevin L. Hockett*[1,3]

[1]Department of Plant Pathology and Environmental Microbiology, Pennsylvania State

University, University Park, PA, 16802, USA

[2]School of Plant Sciences, University of Arizona, Tucson, AZ 85721, USA

[3]The Huck Institutes of the Life Sciences, The Pennsylvania State University, University Park,

PA 16802, USA

*klh450@psu.edu

Abstract

Phage tail-like bacteriocins (tailocins) are bacterially-produced protein toxins that can mediate

competitive interactions between co-colonizing bacteria. Both theoretical and empirical research

has shown there are intransitive interactions between bacteriocin-producing, bacteriocin-

sensitive, and bacteriocin-resistant populations, whereby producers outcompete sensitive,

sensitive outcompete resistant, and resistant outcompete producers. These so-called 'rock-paper-

scissor' dynamics explain how all three populations can be maintained in the same environment,

without one genotype driving the others extinct. Using *Pseudomonas syringae* as a model

system, we demonstrate that otherwise sensitive bacterial cells have the ability to survive

bacteriocin exposure through a physiological mechanism. This mechanism is similar to the

persister phenotype that allows cells to survive antibiotic exposure, without acquiring antibiotic

resistance. We show that a significant fraction of the target cells that survive a lethal dose of

tailocin did not exhibit any detectable increase in survival in subsequent exposure (i.e. they

survived through a persistence-like mechanism). Tailocin persister cells were more prevelant in

stationary rather than log phase cultures. Of the fraction of cells that gained detectable tailocin

resistance, there was a range of resistance from complete (insensitive) to incomplete (partially

sensitive). By genomic sequencing and genetic engineering we showed that a mutation in a

hypothetical gene containing 8-10 transmembrane domains causes tailocin high-persistence and

genes of various glycosyl transferases cause incomplete and complete tailocin resistance.

Importantly, of the several classes of mutations, only those causing complete tailocin resistance

compromised host fitness. This result, combined with previous research, indicates that bacteria

likely utilize persistence as a means to survive bacteriocin-mediated killing without suffering the

costs associated with resistance. This research provides important insight into how bacteria can

escape the trap of fitness trade-offs associated with gaining *de novo* tailocin resistance, and

expands our understanding of how sensistive bacterial populations can persist in the presence of

lethal competitors.

Introduction

Diverse microbes inhabit shared environments and compete for limited resources. Competition

for these resources can be mediated by secretion of toxins such as antibiotics, type VI effectors,

and bacteriocins that enable the producing cells to maintain their dominance (1, 2). Bacteriocins

are bacterially produced protein toxins that are lethal toward strains related to the producer (3, 4).

These antibacterial toxins have been proposed as antibiotic alternatives to treat or prevent

infection spread in both humans and plants (5, 6), in addition to being used in food preservation

for several decades (7). Given their ubiquitous nature, where sequenced bacteria are commonly

predicted to encode at least one bacteriocin (8-10) and bacteria isolated from diverse

environments often produce detectable bacteriocins [eg. (8, 11, 12), it is reasonable to predict

that bacteriocin producing populations exert a selective force on co-colonizing sensitive

populations to either gain resistance, or avoid killing through a different mechanism. Resistance

evolution, often involving a heritable mutation in either the toxin receptor or membrane

translocator genes, is a common mechanism to avoid being killed (13, 14). Resistant mutants,

however, are likely to suffer fitness costs associated with the mutation, which reduces their

ability to proliferate in the environment. Therefore, resistant mutants are only competitive in

environments where there is substantial or sustained bacteriocin exposure, otherwise they are

competitively inferior to their sensitive progenitors. Conversely, bacteriocin producing

populations are dominant over sensitive populations, but are competitively inferior to resistant

populations, as the result of the resources wasted on ineffective toxins. These non-transitive

dynamics underly a rock-paper-scissor model for microbial competition (13-15), where a

community composed of all three genotypes is maintained as a result of negative frequency-

dependent selection (14). These dynamics, however, which are built primarily on modelling and

laboratory culture-based experiments, are dependent on a small number of qualitative states. It is

unknown whether or how quantitative resistance or physiological tolerance in otherwise sensitive

populations will affect community competitive dynamics. The ability of bacteriocin sensitive

cells to persist under toxin pressure has important implications for the ecology of microbes

generally, and for the potential to use bacteriocins as therapeutics specifically. For instance,

previous studies have reported that, although sensitive cells may not co-exist with the producing

strain in a well-mixed environment (16-18), they still prevail in the competitive *in vivo*

environments (13, 14, 19). Little is known, however, regarding the mechanism(s) that allow

maintainance of a sensitive population despite sustained bacteriocin exposure.

Bacteriocins are classified into different groups based on their structure, composition, and mode

of action. Tailocins are bacteriocins that resemble bacteriophage tails and are grouped into R-

type (with a retractile core tube) and F-type (flexible) (20, 21). In the opportunistic human

pathogen *P. aeruginosa* and other environmental Pseudomonads and *Burkholderia*, phage-

derieved tailocin bacteriocins were shown to antagonize competitors including pathogenic strains

(22-28). In fact, a recent study by Principe et al. suggested the effectiveness of foliar sprays of

tailocins produced by *P. fluorescens* in reducing the severity and incidence of bacterial-spot

disease in tomato caused by *Xanthomonas vesicatoria* (28). Other studies have also indicated the

potential use of engineered R-tailocins in suppressing foodborne pathogens using *in vivo* models

(29, 30). Our group has previously characterized a R-type tailocin from a plant pathogenic

bacterium *P. syringae* pv. *syringae* (*Psy*) B728a (31). This R-type tailocin showed antagonistic

potential against several pathovars of *P. syringae* that cause serious diseases and substantial

losses in economically important crops such as common bean (pv. *phasiolicola*), soybean (pv.

glycinea), chestnut (pv. *aesculi*), and kiwifruit (pv. *actinidae*) (31). A broad spectrum of tailocin

mediated antagonistic interaction in *P. syringae* has been described recently (11). Yet, we have a

very limited understanding regarding the defense responses against tailocin by the target

pathogen, a key information to design tailocins as therapeutic agents.

Tailocins are considered to be potent killers as a single tailocin particle is predicted to kill a

sensitive cell and an induced producer cell can release as many as 200 particles (32, 33). R-

tailocins, once bound to the cell surface receptors of the target cells, puncture through the cell

membrane and cause cell death by membrane dissipation (3, 32). Specific lipopolysaccharide

(LPS) components of the target cells are known to serve as surface receptors of tailocins (34-36).

LPS is composed of three components: the lipid A, core oligosaccharide, and the O-

polysaccharide (O-antigen) (37, 38). Although the lipid A and core region are mostly conserved

within a species, the O-antigen region varies extensively in its chain length and composition of

sugars (39). Biosynthesis and transport of LPS to the outer membrane as well the modification of

O-antigen involves complex processes involving a number of highly diverse genes (38). Little is

known about the extent of LPS modification required for tailocin resistance and persistence and

the consequence of these modifications in host fitness and virulence in the target pathogen .

This study aimed to examine the phenomena that addresses both of the crucial questions related

to bacteriocins: 1) how sensitivie cells survive lethal bacteriocin exposure and 2) can

bacteriocins be used as stand alone control measures? We demonstrate that a sensitive population

can employ multiple strategies to survive toxin exposure without acquiring an otherwise fitness-

reducing mutation. One strategy was physiological persistence, a mechanism that enables a sub-

population of sensitive cells to transiently survive lethal doses of the bacteriocin without

undergoing genetic changes. The second strategy relies on acquiring subtle genetic changes

(incomplete resistance) that do not impose a detectable fitness cost, while still allowing the

mutants to better survive bacteriocin exposure. We found pronounced differences between the

frequencies of persistent cells depending on growth phase of the target cells. Moreover, we

identified ten unique mutant alleles with likely roles in the lipopolysaccharide (LPS) O-antigen

biosynthesis leading to various degrees of tailocin resistance. We also recovered a mutation in an

open reading frame, located in an LPS biosynthesis operon, that results in increased bacteriocin

persistence. In addition to increasing the basal persistence frequency, this mutation results in a

loss of growth phase-dependent difference in tailocin peristence. Finally, we demonstrated that

the complete mutants suffered a fitness cost within a susceptible host plant, whereas both high

persisteter, as well as incomplete resistant mutants were equally fit compared to the wild-type.

This work suggests that bacterial cells can employ mechanisms to survive antagonistic toxins but

still preserve their host colonization potential. This work has important implications for how

bacteria can potentially avoid rock-paper-scissor dynamics widely understood to be important in

mediating interactions between toxin producing, sensitive, and resistant genotypes and in using

bacteriocins as potential therapeutic agents.

Results

A sub-population of *Pph* cells survived tailocin by persistence that increased in the stationary state

The relative activity of the purified tailocin was determined to be 10^3-10^4 activity units (AU) and 1.25×10^7- 4.25×10^9 lethal killing units/ml. The minimum inhibitory concentration was estimated to be 100 AU when exposed to ~10^6 viable target cells at their logarithmic growth. No loss of tailocin activity was observed for a period of over six months in the buffer (10 mM Tris PH 7.0, and 10 mM MgSO4) at 4◦C.

Purified tailocin was used to test its killing effects on stationary and log phase cultures of the *Pph* target cells in a broth environment. After an hour of 100 AU tailocin treatment, a consistent reduction (3.59 ± 0.12 log) in the viable population occurred for logarithmic cultures, while a significantly lower reduction (1.38 ± 0.14 log) occurred for the stationary cultures. Further analysis showed that, upon treatment of equivalent number of viable cells, stationary cells consistently survived 10 to 100-fold more than the logarithmic cells (**Fig 1** and **Fig S1**). Surviving colonies, especially those from the stationary phase, were predominantly sensitive upon tailocin re-exposure suggesting survival by persistence mechanism (see below).

The persistent sub-population was maintained under prolonged exposure time and increased concentration of tailocin

Tailocin treatments were applied to both stationary and log cultures of *Pph* for up to 24 hours with enumeration of surviving population before and after 1, 4, 8, and 24 hours of tailocin treatment to generate a tailocin death curve. After a steep reduction in the population within the

first hour of treatment, further killing of the cells that survived the first hour treatment, did not

occur in either culture (**Fig 2A**). Twenty four hours post-treatment, although the overall

population increased (**Fig 2A**), individual treatments showed different results: for some replicate

treatments, the population remained constant suggesting maintenance of the persistent state,

while for some other replicates, population increased due to replication of cells that acquired

tailocin resistance (see **Fig S2**).

Upon tailocin re-treatment, >90% of stationary and >60% of log cells that survived the first hour

treatment, were as sensitive as the wild type (i.e. persistent) as in (**Fig 2B**). The proportion of

persistent survivors was higher in the stationary cultures than in the log cultures at all time points

(**Fig 2B**). Tailocin persistent cells were recovered from both cultures even after 24 hours of

tailocin treatment, although the proportion decreased over time (**Fig 2B**). Tailocin activity was

detected in the supernatants recovered from the treated samples that contained persistent cells

(**Fig S3**), confirming saturation of tailocin in the treatment. Although a slight reduction of

activity was observed when the tailocin preparation was mixed with undiluted stationary

supernatant compared to log supertanant, no difference was detected upon diluting the supertants

up to 1,000-20,000-fold before mixing with tailocin (similar to how the cultures were diluted for

tailocin treatment) (**Fig S4**). This suggested that the increased tailocin persistence in the

stationary phase is not related to inhibition of tailocin activity by an extracellular component.

Upon treating the cells with a concentrated tailocin (900 AU) the surviving population decreased

such that no difference in survival between the stationary and log cultures was detected (**Fig 3A**).

However, even with this higher level of tailocin applied, the proportion of tailocin persistent cells

remained higher for stationary phase survivors than that for the log phase survivors (**Fig 3B**).

175 **Tailocin exposure selected for heritable mutants showing increased persistence and**

176 **heterogenous resistance**

177 In addition to the recovery of tailocin persistent sub-population, we recovered an unique mutant,

178 refered here as high persistent-like (HPL), which survived significantly greater than the wild type

179 under liquid-broth treatment (**Fig. 4A**). However, under a long-term exposure of overlay

180 condition, it showed similar sensitivity to the wild type (**Fig. 4B**). Furthermore, the HPL

181 phenotype did not differ in survival between the stationary and log phases even at a higher

182 concentration of tailocin applied (**Fig 5**). Next category of mutants recovered were conditionally-

183 sensitive and are referred here as incomplete resistant (IR) mutants (see **Fig 2B and 3B** for

184 proportion). These mutants lost sensitivity in the broth even at high tailocin concentration (**Fig.**

185 **4A)**, but displayed some sensitivity in the overlay (**Fig. 4B**). Lastly, complete tailocin resistant

186 (R) mutants that were insensitive to tailocin under both treatment conditions were also recovered.

187 Of the four complete resistant mutants we selected, two (R1 and R4) showed an unique rough

188 colony morphotype.

189

190 **Mutations involved various genes likely associated to LPS biogenesis and modification**

191 Genome sequencing and variant identification of the high-persistent like (n=1), incomplete

192 resistant (n=9), and resistant mutants (n=4) was performed by mapping the Illumina reads with

193 the parental reference sequence. Mutants isolated at different experiments showed mutation in a

194 different locus. A specific region (**Fig 6**) was identified in the *Pph* genome that showed the most

195 prominent role in tailocin activity. The HPL mutant contained a 16 bp deletion in an ORF that

196 caused frameshift near the C-terminal of a hypothetical protein that is co-transcribed with the

197 LPS genes. No functional evidence could be found for the hypothetical protein by *in silico*

analysis except that it was predicted to contain 8-10 transmembrane domains. Majority of genes

identified for complete and incomplete resistance were glycosyl transferases and related proteins

that are likely involved in LPS biogenesis (**Table 1 and Fig 6**). For example, for the gene

PSPPH_0957 that encodes a glycosyl transferase family 1 protein, a frame-shift and a missense

mutation in the middle of the gene caused complete resistance while insertion of few bases at the

3'end of the gene caused incomplete resistance. Moreover, one of the incomplete resistant

mutant class (IR6) showed mobilization of the 100 bp MITE sequence present in the *Pph*

genome as described by Bardaji et al (40). The insertion of MITE inactivated a gene that is

annotated to encode a FAD-dependent oxidoreductase.

The mutant phenotypes detected by genome sequencing were further confirmed by sanger

sequencing of the target region as well as by swapping the mutant alleles to the wild type

background and vice-versa of the selected mutants (HPL, IR4, R1, R3, and R4). In all cases, the

allele-swapped strains showed the phenotypes as expected (**Table 1**). This confirmed that the

mutation identified by genome sequencing were responsible for the tailocin high persistent-like

and resistant phenotypes.

LPS analysis of the mutants and the wild type *Pph* showed that the complete resistant mutants

lacked fully-formed O-antigen region, whereas the high persistent-like and majority of

incomplete resistant mutants still possessed the O-antigen with minor to undetactable changes.

One of the incomplete resistant mutants (IR4), however, showed a very different and faint O-

antigen band (**Fig S5**).

In planta fitness was compromised only for complete resistant mutants devoid of LPS O-antigen

As indicated by the population levels of each mutant lines in the green bean plants, while the high persistent-like and incomplete resistant mutants did not suffer any, the complete resistant mutants suffered a significant fitness cost as earley as 24 hours post-inoculation (**Fig. 7 and Fig. S6**). Up to a 100-fold reduction in the population was seen for the complete resistant mutants (equivalent to the reduction in the type III secretion system mutant). At 48 hpi, although the complete resistant mutants population increased compared to the type III mutant, it was still significantly lower than wild type, HPL and IR mutants (**Fig. 7 and Fig. S6**). These results suggest that persistence and incomplete resistance are mechanisms to survive attack by the competitor strains while keeping the host fitness and virulence potential intact.

Discussion

There have been renewed research interests in alternative treatment strategies for bacterial pathogens due mainly to the growing threats of antibiotic resistant infections (5, 7). Bacteriocins including tailocins have long been proposed as effective and more specific alternatives to broad spectrum antibiotics (5, 6). However, a critical question that remains unaddressed in designing bacteriocins as pathogen control agents is; how does the sensitive pathogen population respond to application of lethal doses of bacteriocins? Moreover, although it is predicted in the rock-paper-scissors model of bacteriocin mediated interaction, that sensitive cells co-exist with the producer through ecological mechanisms (13, 14, 19), there is no description of the possibility of maintaining a bacteriocin sensitive population through a physiological mechanism.

In this study, we addressed these questions using a phage-tail like bacteriocin (i.e. tailocin)

produced by *P. syringae* pv. *syringae* strain B728a in killing target cells of *P. syringae* pv.

phaseolicola 1448A. We showed that, upon exposure of a lethal dose of tailocin, a sub-

population of sensitive cells survives without undergoing genetic changes. The fraction of this

sub-population, termed here as tailocin persistent sub-population, increased significantly in the

stationary phase than in the logarithmic phase of growth. By repeated exposures of this sub-

population to same or higher doses of tailocin, we showed that they have not gained any

heritable resistance, and physiological persistence is the only mechanism for their survival.

Moreover, a prolonged tailocin exposure generated a killing pattern similar to that reported for

persistent sub-population upon antibiotic treatment (41, 42). Persistence was maintained for at

least 24 hours with tailocin exposure, a phenomenon that was more evident in some treatment

replicates in which resistant evolution did not occur (see Fig. S2). Although increasing the

tailocin concentration killed some of the persistent survivors, and the difference in survival

between the two growth phases was no longer seen, stationary phase-derived cells still exhibited

higher persistence than the log phase cells upon re-exposure. This indicated that the stationary

cells may require multiple hits by tailocin particles as opposed to the one-hit-one-kill mechanism

of killing described for tailocins (32, 33), or that the probabilty of a successful hit in stationary

phase is lower than in log phase. Since tailocins are thought to be target cell specific, and are not

known to have off-target effects, higher concentration of tailocin could be used to achieve a more

effective pathogen control. However, although at a low level, persistence was still maintained

even with high-dose tailocin treatment and inherent emergence of either complete or incomplete

resistance was frequently observed. As such, although a significant reduction in pathogen

population and disease pressure can be obtained with tailocins, a stand-alone tailocin treatment

might not be enough to achieve a sustainable pathogen control.

The use of the term 'persistence' in relation to antimicrobial survival is disputed to some extent

and is sometimes used interchangably with 'tolerence' and 'viable but not culturable state'. In

this paper, we used 'persistence' as this phenotype was only seen in a sub-population, resulted in

a bi-phasic death curve, cells resuscitated almost immediately upon tailocin removal, and were

equally sensitive to the wild type cells upon re-exposure. This definition of persistence has been

suggested previously (42). Persistence to antimicrobials is being increasingly recognized for its

role in antimicrobial treatment failures with bacterial infections (43). Various mechanisms are

implicated in the maintainance of persistence (44, 45), although persistence responses can be

different based on the stresses involved and their mode of action (46). Of these mechanisms,

toxin-antitoxin (TA) systems are the most studied ones for formation of persister-subpopulation

(47-49). TA systems were shown to be induced when cells were starved for certain sugars and

amino acids or by exposure to osmotic stresses that altered ATP levels in the cell (46, 50).

However, It was also shown that activation of TA system does not always induce persister

formation (50). Additionally, recent findings have indicated a mechanism mediated by the

guanosine penta- or -tetraphosphate (ppGpp) for persister formation that is not dependent on a

TA system (51). A strong stationary state effect, that likely involved starvation response, was

shown to increase persistence by 100-1,000 fold in *Staphylococcus aureus* with ciprofloxacin

treatment (50). Whether similar mechanisms of TA and independent ppGpp systems regulate

tailocin persistence or a specific mechanism for tailocin and/or related bacteriophage exists,

remains to be determined. Nevertheless, our data of the difference in tailocin persistence between

the stationary and log cultures suggests that metabolic inactivity and starvation-induced stress

could be a strong factor in tailocin persistence.

Few previous studies have demonstrated growth-phase dependent differences in LPS O-antigen

chain length and composition or their regulatory pathways (52, 53). In a previous study with *P.*

fluorescens, exponentially growing cells had a significantly higher rate of cell lysis than

stationary or decline phase cells with bacteriophage PhiS1 (54). Moreover, a recent study

showed that sensitivity to colicin, a bacteriocin from *Eschericia coli*, could be altered by growth

condition dependent LPS O-antigen changes (55). Growth phase dependent LPS modification,

although has not been reported so far in *P. syringae*, could be contributing to the difference in

tailocin persistence. We also observed that undiluted stationary phase supernatant inhibited

tailocin activity to some extent compared to the log phase supernatant. However, whether this is

linked to differences in the secreted LPS between the two supernatants or other cellular factors

needs further assessment. Moreover, we also demonstrated that mutation in one of the

hypothetical proteins containing a signal peptide and several trans-membrane domains caused

increased tailocin persistence. Since the hypothetical protein occurs in the same operon as other

LPS biogenesis genes, it is likely that it plays a role in O-antigen biogenesis and/or modification,

thereby reducing tailocin interaction with the cells.

Phase variation is another mechanism that is known to cause increased survival to surface active

antimicrobials (eg. phages and host immune defenses) (56). Phase variation is a gene regulation

system that induces heterogenous expression of specific genes in a clonal population (56-58).

Although phase variation is heritable, the 'ON' 'OFF' switch from variant to wild type

phenotype occurs randomly amounting to 10_{-4} to 10_{-1} per generation, significantly greater than

what is expected by mutational events (59). Phase variation has been shown to modify LPS

operons in *S. enterica* spp (60), *P. aeruginosa* (61)*,* and *P. fluorescence* (62)*.* In *S. enterica*

serovar Typhimurium phase variable glucosylation of the O-antigen was reported to cause

temporal development of phage resistance (63). Other reports of the role of phase variation in

defense against bacteriophages, that also target LPS, are available (64, 65), including in the

closely related *P. aeruginosa* (61). Therefore, phase variation in tailocin survival can not be

ruled out since LPS is the receptor involved. Since the rate of switch from the tailocin surviving

variant to the sensitive wild type upon tailocin removal will have to be highly greater than

reported (59), the tailocin persistent phenotype is unlikely to be the result of phase variation. It

remains to be determined, however, if the incomplete resistance could be caused by phase

variable gene switching. In one of the incomplete resistant mutants, the target gene was modified

by movement and integration of an internal mobile genetic element (MGE). MGEs have been

shown to cause phase variation in other bacterial systems (66, 67). A high-throughput sensitivity

screening of the mutant progeny colonies will be required to confirm if the tailocin resistant

phenomena involves phase variable gene switch.

To our knowledge, the persistent as well as incomplete resistant phenotypes observed here have

not been described before with bacteriocins. Here, using a *Pseudomonas syringae* tailocin, we

showed that, in addition to the persistent sub-population, resistant lines with various degrees of

sensitivity are selceted by exposure of target cells to tailocins. LPS analysis and *in planta* fitness

assessement showed that, to gain complete tailocin resistance, cells have to loose their LPS O-

antigen. This comes with a fitness trade-off. On the contrary, by maintaining a persistent sub-

population and/or by undergoing subtle genetic changes, the LPS integrity was preserved for the

most part and fitness within the host was maintained. Similar results of reduced virulence and

plant fitness was shown in phage resistant mutants devoid of O-antigen (68). Moreover, two of

the complete resistant mutants (R1 and R4) in our study showed a typical rough-colony

morphotype. Rough-colony morphotypes lacking LPS O-antigen were resistant to phages and

had reduced *in planta* virulence as described previously in *P. syringae* pv. *morsprunorum*

isolates from plum and cherry (62). In other plant pathogenic bacteria such *as Xanthomonas*

oryzae pv. *oryzae* and *Xylella fastidiosa,* loss of O-antigen, respectively, reduced type III

secretion into plants (69), and increased recognition of the pathogen by the host immune

response (70). In both cases, plant virulence of the mutants was significantly reduced (69, 70).

Also, the incomplete resistant mutants, although contained mutations in the LPS biogenesis

genes and mostly had their O-antigen region intact, did not show any fitness trade-offs in our

growth chamber experiments. However, under a natural environment that involves more severe

environmental stresses, it could be expected that their fitness would be altered. Under these

circumstances, the persister sub-population, that does not involve any genetic changes, would

enable the target pathogen to withstand the host defence while surviving the competition and

ensure that its lineage is maintained.

Taken together, both previous reports and our results suggest that full resistance to tailocin incurs

significant fitness cost. Our work demonstrates that persister and incomplete resistant sub-

populations of the sensitive strain preserve their host fitness despite suffering bacteriocin

exposure. These results have important implications for the mechanisms and ecological

processes that can promote the co-existence of both sensitive and bacteriocin producing

populations. In **Fig 8**, we present a model of the three cell types (resistant, incomplete resistant,

and persister) that can be detected following bacteriocin-mediated selection. In particular, we

view the persister sub-population as one that can survive bacteriocin exposure, without paying

any long term fitness cost. In the short term, however, we believe that persister cells have a

fitness of zero because they are likely not replicating (see Figure S2). Thus, bacteriocin persisters

may best be thought of as a means to switch at high frequency between being phenotypically

resistant and phenotypically sensitive.

Materials and Method

Bacterial strains, media, and culture conditions

All bacterial strains, plasmids, and mutants used in this study are listed in Table 1 and 2. *P.*

syringae pv. *syringae* (*Psy*) wild type (WT) strain B728a and its tailocin defective mutant ΔRrbp

were used to prepare the treatment supernatants. *P. syringae* pv. *phaseolicola* (*Pph*) 1448A was

used as the target strain. Tailocin high-persistent like, resistant, and selected complementated

strains of *Pph* generated in this study are described in Table 1. King's medium B (KB) was used

to culture the strains. Liquid cultures were prepared by inoculating individual colonies from a

two-day old KB agar plate into 2 ml of liquid medium at 28◦C with shaking at 200 rpm.

Antibiotics Kanamacyin (Km), Chloramphenicol (Cm), Tetracycline (Tet), Gentamycin (Gm),

Rifampcin (Rif), Nitrofurantoin (NFT), and Nalidixic Acid (NA) were used at 50, 25, 10, 50, 50,

50, 50 µg/ml final concentration, respectively.

Tailocin induction and purification

Purified tailocin and control treatments were prepared from logarithmic (log) cultures of *Psy*

B728a and ΔRrbp, respectively using a polyethylene glycol (PEG) precipitation protocol as

previously described (31, 71). Briefly, 100-fold diluted overnight B728a cultures were sub-

cultured for 4-5 hours in KB before inducing with 0.5 µg ml$_{-1}$ final concentration of mitomycin C

(GoldBio). Induced cultures were incubated for 24 hours with shaking at 28∘C. Next, cells were

pelleted by centrifugation and the supernatants were mixed with 10% (w/v) PEG 8000

(FisherScientific) and 1M NaCl. Supernatants were then incubated either in ice for 1 hour and

centrifuged at 16,000 g for 30 min at 4∘C or incubated overnight at 4∘C and centrifuged at 7000g

for one hour at 4∘C. Pellets were resuspended (1/10- 1/20th of the original volume of the

supernatant) in a buffer (10 mM Tris PH 7.0, and 10 mM MgSO4). Two extractions with equal

volume of chloroform were performed to remove residual PEG. Tailocin activity was confirmed

by spotting dilutions of 3-5 µl of both the tailocin and control supernatants onto soft agar

overlays of *Pph*. The relative activity of tailocin was expressed as arbitrary units (AU) as

obtained from the reciprocal of the highest dilution that exhibited visible tailocin killing in an

overlay seeded with ~10_8 CFUs of *Pph* log cells. Lethal killing units of the purified tailocin were

determined using a Poisson distribution of the number of surviving colonies after treatment with

different dilutions of the tailocin as described previously (27, 72).

Tailocin treatment and survival assessment for stationary and log cultures

To assess tailocin activity against the stationary and log phases of *Pph*, individual colonies

growing on KB agar plates for ~2 days were inoculated into 2 ml of liquid KB medium.

Following incubation at 28∘C with shaking at 200 rpm overnight, the cultures were back diluted

1000-fold into fresh KB. The back diluted cultures were either incubated for 28-30 hours to

prepare stationary cultures, or back-diluted 100-fold at 24 hours and cultured for another 4-6

hours to prepare log cultures (**see Fig S7** for a growth curve of *Pph*).

Stationary cultures were diluted 20,000-fold and logarithmic cultures were diluted 1,000-fold

[~10_5 -10_6 CFUs/ml for both cultures **see Fig 1**] in fresh KB before tailocin treatment. Treatment

was applied by mixing 10 µl of diluted cultures in 90 µl of purified tailocin diluted in KB. After

treatment, samples were incubated for ~ 1 hour at 28∘C and washed twice to remove residual

tailocin particles. Washing was performed by mixing the treated culture in 900 µl of fresh KB

followed by centrifugation at 12,000 g for 2 min. The top 900 µl fraction was discarded and the

bottom 100 µl fraction was serially diluted and either spread- or spot-plated to enumerate

surviving population. Plates were incubated at 28∘C for 1-3 days before enumeration. Serial

dilutions of both stationary and log cultures were spotted onto KB agar to enumerate the

untreated population. Experiments were performed with various tailocin concentrations (i.e. 100

AU, 500 AU, and 900 AU).

Tailocin re-treatment to differentiate persistence and resistance

Surviving colonies were treated again with tailocin to differentiate them into persistent or

resistant colonies. Re-treatment was performed by an overlay method as described previously

(71), or by broth treatment as discussed above. Overlay method was used to determine the AU of

the tailocin preparation with the selected mutant lines. Broth exposure was used to calculate

reduction in the population of log cultures after treatment. Surviving colonies were differentiated

into various phenotypes as follows: persistent (sensitive to tailocin to the wild-type level in both

the overlay and broth method), high -persistent like (completely sensitive in the overlay but

survived significantly more than wild-type under broth condition), incomplete resistant [showed

conditional sensitivity (i.e. some sensitivity in overlay but no significant sensitivity in the broth),

and complete resistant (were insensitive under both conditions).

Time dependent death curve with tailocin treatment

Prolonged tailocin exposure was performed with both the stationary and log-phase cultures of

Pph to generate death curves. Treatment was applied in a 96-well plate as before. Surviving

populations were enumerated at 1, 4, 8, and 24 hours following treatment and randomly selected

surviving colonies were re-exposed to tailocin to differentiate them into persistent or resistant

phenotypes.

Tailocin recovery from the treated samples and activity testing

Stationary and log cultures treated with purified tailocin for 1-24 hours as described above were

centrifuged and the supernatant was collected, and filter sterilized using a 0.22 µm syringe filter.

Supernatants were diluted 5-, 10-, 50-, and 100-fold in KB and spotted on *Pph* overlay. Purified

tailocin particles diluted in KB was also included as control treatment.

Determining the effect of stationary and log supernatant on tailocin activity

Stationary and log phase cultures were prepared as described above by culturing *Pph* cells in KB

broth for either 28-30 hours or for 4-6 hours, respectively. Cultures were centrifuged for 2 min at

12,000 g and the supernatant was filter sterilized using a 0.2 µm syringe filter. Stationary and log

supernatants were diluted 1,000-20,000-fold in KB (according to how the cultures were diluted

for tailocin treatment). Various dilutions (10-, 50-, 100-, and 1000-folds) of purified tailocin

were prepared in the stationary and log supernatants. Dilutions were spotted on a *Pph* lawn using

the overlay method.

Genome sequencing and analysis of the tailocin high-persistent and resistant mutants

The tailocin high-persistent like (HPL) and complete and incomplete resistant mutants recovered

from tailocin treatment of *Pph* wild type cells (and confirmed by re-treatment) were selected for

genome sequencing. DNA was extracted from the overnight cultures using Promega Wizard

Genomic DNA purification Kit using manufacturer's protocol. DNA quantity and quality were

assessed with Qubit 3 Fluorometer using Qubit™ dsDNA HS Assay Kit (Thermo Fisher

Scientific) and Nanodrop 2000 (Thermo Scientific). A uniquely indexed library from each

mutant line was prepared using Illumina DNA Flex kit (Illumina). An approximately equimolar

pool of libraries was generated and 150bp paired-end reads were sequenced on an Illumina

MiSeq.

Resulting forward and reverse read files were paired and mapped to the *Pph* 1448A

chromosomal and plasmid sequences using Geneious R10.2 with default parameters for medium

sensitivity. Next, genetic variants were identified using 'Find variations/SNPs' program within

Geneious using default settings. Regions supported by a minimum of 10 reads and >90% variant

frequency were selected. Moreover, variants shared among all the mutants that were generated in

independent experiments were discarded as misalignments. Variants identified by Geneious were

confirmed in the contigs generated by De Novo assembly of the paired reads using SPAdes

3.11.0 using K-mer sizes of 21, 33, 55, 77, 99, 127 with careful mode selected to minimize

mismatches and short indels. Indels were also detected and visualized in the contigs by Harvest

suit tools (73). Sequencing generated 1,986,400 -2,857,324 paired reads per genome with 48-

69X genome coverage. *De novo* assembly generated 308-338 contigs per genome with the N50s

of 75,222-81,220bp. The total assembly size was 5.97 Mbp with a GC content of 57.96%, values

similar to *Pph* reference genome (74). The read files of all the genomes generated in this study

have been deposited to the NCBI Short Read Archive (SRA) database under a Bioproject

PRJNA608702 and biosamples SAMN14206700- SAMN14206710. The SRA accession

numbers are SRR11179092- SRR11179102.

Presence absence and similarity searches of the selected genomic regions and genes implicated

in tailocin persistence and resistance were performed with NCBI and IMG-JGI databases using

BLAST algorithm using the *Pph* sequences as query. InterProScan (75) and Phobious program

within the Geneious plugin was used to predict functional domains in the amino acid sequences.

Confirmation of mutant phenotype by allele swap

The mutations were further confirmed with sanger sequencing and by swapping the mutant allele

to the wild type background and vice-versa. Allele swap experiments of the selected incomplete

resistant (IR4) and complete resistant mutants (R1, R3, and R4) were performed as previously

described in Hockett et al (31). Briefly, a ~1kb fragment containing the mutant or a WT allele

was amplified using a Phusion High-Fidelity DNA Polymerase (New England Biolabs) using

standard protocols. Primers used for generating the PCR fragments and mutant confirmation are

listed in Table 3. The PCR fragment contained gateway cloning sites added through the primer

extension. Purified PCR fragment was cloned into pDONR207 and further recombined into

pMTN1907 using BP and LR clonase enzymes, respectively. pMTN1907 containing a desired

clone was transformed to S17-1 and conjugated to the *Pph* wild type or mutant background by

bi-parental mating. Tet$_R$ merodiploids of *Pph* selected on Tet, Rif, and NFT plates were counter-

selected in KB supplemented with 10% Sucrose, followed by PCR confirmation of the desired

allele-swapped strains. Allele swap of the high persistent-like (HPL) mutant was performed

similarly, but using a one-step gateway vector pDONR1K18ms and the merodiploids were

selected in Km, Rif, NFT plates.

LPS extraction and visualization

LPS extraction was performed as described by Davis and Goldberg (76) from pellets of one

milliliter of overnight cultures (OD_{600} 0.5). After extraction, 10 µl samples were separated by

SDS-PAGE. Samples were labeled using the Pro-Q Emerald 300 glycoprotein stain kit

(ThermoFisher Scientific, #P21857) according to manufacturer's instructions. For LPS size

determination, CandyCane™ glycoprotein molecular weight standards (ThermoFisher Scientific,

#C21852) were included. Gels were visualized using Molecular Imager Gel-Doc XR+ (Bio-Rad)

with Image Lab Software.

In planta fitness test of the high-persistent, incomplete resistant, and resistant mutants

The high persistent-like (HPL), selected incomplete resistant, and complete resistant mutant of

Pph including a type III secretion mutant Δ*hrpL::Pph* (77) were tested for their *in planta*

fitness. *In planta* experiment was performed in a growth chamber (Conviron) maintained at 24◦C,

75% RH and 16 and 8 hours of day/night cycles. Plants of Dwarf French Bean (*Phaseolus*

vulgaris) variety 'Canadian Wonder' were grown in a Dillen 6.0 Standard pots (Onliant) in

Sunshine Mix 4 Aggregate Plus Professional Growing Mix (Growerhouse). Plants were irrigated

daily. Nine days post seeding, plants were infiltrated with suspension of the bacteria. For

inoculant preparation, overnight cultures of WT *Pph*, Δ*hrpL::Pph*, and the tailocin persistent and

resistant lines were pelleted by centrifugation, washed twice, and resuspended in equal volume

of 10 mM MgCl2 buffer. Optical density (OD_{600}) was adjusted to 0.1 using a Spectronic 200

Spectrophometer (Thermo Scientific) and diluted 50 times. ~200 µl diluted cultures were

infiltrated onto designated areas of the two primary leaves using 1 ml BD syringes and infiltrated

areas were marked. Infiltrated areas were harvested using a 1cm cork borer in a 2 ml tube

containing 200 µl 10 mM MgCl2 and 2 of the 3 mm glass beads (VWR) at 0, 24, and 48 hours

post infiltration. Harvested leaf discs were homogenized in a FastPrep-24 instrument (MP

Biomedicals) for 20 sec. Homogenate (5 µl) after serial dilution were spotted on KB agar plates

supplemented with 50 µg/ml of Nalidixic Acid and CFUs were counted after two days. *In planta*

experiments were repeated at least twice with 8 replications per time.

Statistical analysis

Means of total and surviving population between treatments were compared using the Glimmix

protocol in SAS 9.4 with experimental repeat used as a random factor. Whenever required, post

hoc analysis was performed with Tukey's Honest Significant Difference (Tukey HSD) test at 5%

significance level (P=0.05). *In planta* enumeration data were analyzed in JMP Pro 14 (SAS Inc.)

using Fit Y by X model and One way ANOVA and Tukey HSD at P=0.05 after confirming that

data were normally distributed and had equal variances.

Acknowledgements

This research was supported by the following sources for KLH: the USDA National Institute of

Food and Federal Appropriations under project PEN04648 and Accession 1015871, the USDA

NIFA Foundational Program [2019-67013-29353], and the Lloyd Huck Early Career

Professorship.

References

1. Granato ET, Meiller-Legrand TA, Foster KR. The evolution and ecology of bacterial warfare. Curr Biol. 2019;29(11):R521-r37.

2. Stubbendieck RM, Straight PD. Multifaceted interfaces of bacterial competition. J Bacteriol. 2016;198(16):2145-55.

3. Ghequire MGK, De Mot R. The tailocin tale: peeling off phage tails. Trends Microbiol. 2015;23(10):587-90.

4. Baba T, Schneewind O. Instruments of microbial warfare: bacteriocin synthesis, toxicity and immunity. Trends Microbiol. 1998;6(2):66-71.

5. Cotter PD, Ross RP, Hill C. Bacteriocins - a viable alternative to antibiotics? Nat Rev Microbiol. 2013;11(2):95-105.

6. Vidaver AK. Prospects for control of phytopathogenic bacteria by bacteriophages and bacteriocins. Annu Rev Phytopathol. 1976;14(1):451-65.

7. Cotter PD, Hill C, Ross RP. Bacteriocins: developing innate immunity for food. Nat Rev Microbiol. 2005;3:777.

8. Gordon DM, O'Brien CL. Bacteriocin diversity and the frequency of multiple bacteriocin production in *Escherichia coli*. Microbiology. 2006;152(Pt 11):3239-44.

9. Riley MA, Wertz JE. Bacteriocin diversity: ecological and evolutionary perspectives. Biochimie. 2002;84(5-6):357-64.

10. Ghequire MG, De Mot R. Ribosomally encoded antibacterial proteins and peptides from *Pseudomonas*. FEMS Microbiol Rev. 2014;38(4):523-68.

11. Baltrus DA, Clark M, Smith C, Hockett KL. Localized recombination drives diversification of killing spectra for phage-derived syringacins. ISME J. 2019;13(2):237-49.

12. Gordon DM, Oliver E, Littlefield-Wyer J. The diversity of bacteriocins in Gram-negative bacteria. In: Riley MA, Chavan MA, editors. Bacteriocins: Ecology and Evolution. Berlin, Heidelberg: Springer Berlin Heidelberg; 2007. p. 5-18.

13. Kerr B, Riley MA, Feldman MW, Bohannan BJ. Local dispersal promotes biodiversity in a
real-life game of rock-paper-scissors. Nature. 2002;418(6894):171-4.

14. Czaran TL, Hoekstra RF, Pagie L. Chemical warfare between microbes promotes
biodiversity. Proc Natl Acad Sci U S A. 2002;99(2):786-90.

15. Kirkup BC, Riley MA. Antibiotic-mediated antagonism leads to a bacterial game of rock-
paper-scissors in vivo. Nature. 2004;428(6981):412-4.

16. Chao L, Levin BR. Structured habitats and the evolution of anticompetitor toxins in
bacteria. Proc Natl Acad Sci U S A. 1981;78(10):6324-8.

17. Durrett R, Levin S. Allelopathy in spatially distributed populations. J Theor Biol.
1997;185(2):165-71.

18. Gordon DM, Riley MA. A theoretical and empirical investigation of the invasion dynamics
of colicinogeny. Microbiology. 1999;145 (Pt 3):655-61.

19. Riley MA, Wertz JE, Goldstone C. The ecology and evolution of microbial defense systems
in *Escherichia coli*. EcoSal Plus. 2004;1(1).

20. Nakayama K, Takashima K, Ishihara H, Shinomiya T, Kageyama M, Kanaya S, et al. The
R-type pyocin of *Pseudomonas aeruginosa* is related to P2 phage, and the F-type is related
to lambda phage. Mol Microbiol. 2000;38(2):213-31.

21. Michel-Briand Y, Baysse C. The pyocins of *Pseudomonas aeruginosa*. Biochimie.
2002;84(5-6):499-510.

22. Smidt ML, Vidaver AK. Isolation and characterization of syringacin W-1, a bacteriocin
produced by *Pseudomonas syringae* pv. *syringae*. Can J Microbiol. 1986;32(3):231-6.

23. Dorosky RJ, Yu JM, Pierson LS, Pierson EA. *Pseudomonas chlororaphis* produces two
distinct R-tailocins that contribute to bacterial competition in biofilms and on roots. Appl
Environ Microbiol. 2017;83(15):e00706-17.

24. Ghequire MGK, Dillen Y, Lambrichts I, Proost P, Wattiez R, De Mot R. Different
ancestries of R tailocins in rhizospheric *Pseudomonas* isolates. Genome Biol Evol.
2015;7(10):2810-28.

25. Yao GW, Duarte I, Le TT, Carmody L, LiPuma JJ, Young R, et al. A broad-host-range
tailocin from *Burkholderia cenocepacia*. Appl Environ Microbiol. 2017;83(10):e03414-16.

26. Vidaver AK, Mathys ML, Thomas ME, Schuster ML. Bacteriocins of the phytopathogens
Pseudomonas syringae, *P. glycinea*, and *P. phaseolicola*. Can J Microbiol. 1972;18(6):705-
13.

27. Haag WL, Vidaver AK. Purification and characterization of syringacin 4-A, a bacteriocin
from *Pseudomonas syringae* 4-A. Antimicrob Agents Chemother. 1974;6(1):76-83.

28. Principe A, Fernandez M, Torasso M, Godino A, Fischer S. Effectiveness of tailocins
produced by *Pseudomonas fluorescens* SF4c in controlling the bacterial-spot disease in
tomatoes caused by *Xanthomonas vesicatoria*. Microbiol Res. 2018;212-213:94-102.

29. Ritchie JM, Greenwich JL, Davis BM, Bronson RT, Gebhart D, Williams SR, et al. An
Escherichia coli O157-specific engineered pyocin prevents and ameliorates infection by *E.*

coli O157:H7 in an animal model of diarrheal disease. Antimicrob Agents Chemother.
2011;55(12):5469-74.

30. Scholl D, Cooley M, Williams SR, Gebhart D, Martin D, Bates A, et al. An engineered R-
type pyocin is a highly specific and sensitive bactericidal agent for the food-borne pathogen
Escherichia coli O157:H7. Antimicrob Agents Chemother. 2009;53(7):3074-80.

31. Hockett KL, Renner T, Baltrus DA. Independent co-option of a tailed bacteriophage into a
killing complex in *Pseudomonas*. mBio. 2015;6(4):e00452.

32. Scholl D. Phage tail-like bacteriocins. Annu Rev Virol. 2017;4(1):453-67.

33. Matsui H, Sano Y, Ishihara H, Shinomiya T. Regulation of pyocin genes in *Pseudomonas*
aeruginosa by positive (prtN) and negative (prtR) regulatory genes. J Bacteriol.
1993;175(5):1257-63.

34. Filiatrault MJ, Munson RS, Campagnari AA. Genetic analysis of a pyocin-resistant
lipooligosaccharide (LOS) mutant of *Haemophilus ducreyi*: restoration of full-length LOS
restores pyocin sensitivity. J Bacteriol. 2001;183(19):5756.

35. Hockett KL, Clark M, Scott S, Baltrus DA. Conditionally redundant bacteriocin targeting by
Pseudomonas syringae. bioRxiv. 2017:167593.

36. Kohler T, Donner V, van Delden C. Lipopolysaccharide as shield and receptor for R-
pyocin-mediated killing in *Pseudomonas aeruginosa*. J Bacteriol. 2010;192(7):1921-8.

37. Maldonado RF, Sa-Correia I, Valvano MA. Lipopolysaccharide modification in Gram-
negative bacteria during chronic infection. FEMS Microbiol Rev. 2016;40(4):480-93.

38. Whitfield C, Trent MS. Biosynthesis and export of bacterial lipopolysaccharides. Annu Rev
Biochem. 2014;83(1):99-128.

39. Lerouge I, Vanderleyden J. O-antigen structural variation: mechanisms and possible roles in
animal/plant-microbe interactions. FEMS Microbiol Rev. 2002;26(1):17-47.

40. Bardaji L, Anorga M, Jackson RW, Martinez-Bilbao A, Yanguas-Casas N, Murillo J.
Miniature transposable sequences are frequently mobilized in the bacterial plant pathogen
Pseudomonas syringae pv. *phaseolicola*. PLoS One. 2011;6(10):e25773.

41. Kaldalu N, Hauryliuk V, Tenson T. Persisters-as elusive as ever. Appl Microbiol
Biotechnol. 2016;100(15):6545-53.

42. Brauner A, Fridman O, Gefen O, Balaban NQ. Distinguishing between resistance, tolerance
and persistence to antibiotic treatment. Nat Rev Microbiol. 2016;14(5):320-30.

43. Lewis K. Persister cells. Annu Rev Microbiol. 2010;64:357-72.

44. Behiels E, Wen Y, Devreese B. Toxin–Antitoxin systems: their role in persistence, biofilm
formation, and pathogenicity. Pathog Dis. 2014;70(3):240-9.

45. Wang X, Wood TK. Toxin-antitoxin systems influence biofilm and persister cell formation
and the general stress response. Appl Environ Microbiol. 2011;77(16):5577.

46. Shan Y, Brown Gandt A, Rowe SE, Deisinger JP, Conlon BP, Lewis K. ATP-dependent
persister formation in *Escherichia coli*. mBio. 2017;8(1):e02267-16.

47. Grant SS, Hung DT. Persistent bacterial infections, antibiotic tolerance, and the oxidative
stress response Virulence. 2013;4(4):273-83.

48. Martins PMM, Merfa MV, Takita MA, De Souza AA. Persistence in phytopathogenic
bacteria: do we know enough? Front Microbiol. 2018;9:1099-.

49. Shidore T, Triplett LR. Toxin-antitoxin systems: implications for plant disease. Annu Rev
Phytopathol. 2017;55:161-79.

50. Conlon BP, Rowe SE, Gandt AB, Nuxoll AS, Donegan NP, Zalis EA, et al. Persister
formation in *Staphylococcus aureus* is associated with ATP depletion. Nat Microbiol.
2016;1:16051.

51. Song S, Wood TK. ppGpp ribosome dimerization model for bacterial persister formation
and resuscitation. bioRxiv. 2019:663658.

52. Bravo D, Silva C, Carter JA, Hoare A, Alvarez SA, Blondel CJ, et al. Growth-phase
regulation of lipopolysaccharide O-antigen chain length influences serum resistance in
serovars of *Salmonella*. J Med Microbiol. 2008;57(Pt 8):938-46.

53. Lahtinen P, Brzezinska A, Skurnik M. Temperature and growth phase regulate the
transcription of the o-antigen gene cluster of *Yersinia enterocolitica* O:3. In: Skurnik M,
Bengoechea JA, Granfors K, editors. The Genus Yersinia: Entering the Functional Genomic
Era. Boston, MA: Springer US; 2003. p. 289-92.

54. Sillankorva S, Oliveira R, Vieira MJ, Sutherland I, Azeredo J. *Pseudomonas fluorescens*
infection by bacteriophage PhiS1: the influence of temperature, host growth phase and
media. FEMS Microbiol Lett. 2004;241(1):13-20.

55. Sharp C, Boinett C, Cain A, Housden NG, Kumar S, Turner K, et al. O-antigen dependent
colicin insensitivity of uropathogenic *Escherichia coli*. J Bacteriol. 2019;201(4).

56. Bayliss CD. Determinants of phase variation rate and the fitness implications of differing
rates for bacterial pathogens and commensals. FEMS Microbiol Rev. 2009;33(3):504-20.

57. Henderson IR, Owen P, Nataro JP. Molecular switches--the ON and OFF of bacterial phase
variation. Mol Microbiol. 1999;33(5):919-32.

58. van der Woude MW. Phase variation: how to create and coordinate population diversity.
Curr Opin Microbiol. 2011;14(2):205-11.

59. van der Woude MW. Re-examining the role and random nature of phase variation. FEMS
Microbiol Lett. 2006;254(2):190-7.

60. Broadbent SE, Davies MR, van der Woude MW. Phase variation controls expression of
Salmonella lipopolysaccharide modification genes by a DNA methylation-dependent
mechanism. Mol Microbiol. 2010;77(2):337-53.

61. Webb JS, Lau M, Kjelleberg S. Bacteriophage and phenotypic variation in *Pseudomonas
aeruginosa* biofilm development. J Bacteriol. 2004;186(23):8066.

62. Sánchez-Contreras M, Martín M, Villacieros M, O'Gara F, Bonilla I, Rivilla R. Phenotypic
selection and phase variation occur during alfalfa root colonization by *Pseudomonas
fluorescens* F113. J Bacteriol. 2002;184(6):1587-96.

63. Kim M, Ryu S. Spontaneous and transient defence against bacteriophage by phase-variable
682   glucosylation of O-antigen in *Salmonella enterica* serovar Typhimurium. Mol Microbiol.
683   2012;86(2):411-25.

684   64. Seed KD, Faruque SM, Mekalanos JJ, Calderwood SB, Qadri F, Camilli A. Phase variable
685   O antigen biosynthetic genes control expression of the major protective antigen and
686   bacteriophage receptor in *Vibrio cholerae* O1. PLoS Pathog. 2012;8(9):e1002917-e.

687   65. Cota I, Sánchez-Romero MA, Hernández SB, Pucciarelli MG, García-del Portillo F,
688   Casadesús J. Epigenetic control of *Salmonella enterica* o-antigen chain length: a tradeoff
689   between virulence and bacteriophage resistance. PLOS Genet. 2015;11(11):e1005667.

690   66. Szekely E, Simon M. DNA sequence adjacent to flagellar genes and evolution of flagellar-
691   phase variation. J Bacteriol. 1983;155(1):74.

692   67. Ziebuhr W, Krimmer V, Rachid S, Lossner I, Gotz F, Hacker J. A novel mechanism of
693   phase variation of virulence in *Staphylococcus epidermidis*: evidence for control of the
694   polysaccharide intercellular adhesin synthesis by alternating insertion and excision of the
695   insertion sequence element IS256. Mol Microbiol. 1999;32(2):345-56.

696   68. León M, Bastías R. Virulence reduction in bacteriophage resistant bacteria. Front Microbiol.
697   2015;6:343-.

698   69. Wang L, Vinogradov EV, Bogdanove AJ. Requirement of the lipopolysaccharide O-chain
699   biosynthesis gene *wxocB* for type III secretion and virulence of *Xanthomonas oryzae* pv.
700   *Oryzicola*. J Bacteriol. 2013;195(9):1959-69.

701   70. Rapicavoli JN, Blanco-Ulate B, Muszynski A, Figueroa-Balderas R, Morales-Cruz A, Azadi
702   P, et al. Lipopolysaccharide O-antigen delays plant innate immune recognition of *Xylella*
703   *fastidiosa*. Nat Commun. 2018;9(1):390.

704   71. Hockett KL, Baltrus DA. Use of the soft-agar overlay technique to screen for bacterially
705   produced inhibitory compounds. J Vis Exp. 2017(119).

706   72. Mayr-Harting A, Hedges AJ, Berkeley RCW. Methods for studying bacteriocins. In: Norris
707   JR, Ribbons DW, editors. Methods in Microbiology. 7: Academic Press; 1972. p. 315-422.

708   73. Treangen TJ, Ondov BD, Koren S, Phillippy AM. The Harvest suite for rapid core-genome
709   alignment and visualization of thousands of intraspecific microbial genomes. Genome Biol.
710   2014;15(11):524.

711   74. Joardar V, Lindeberg M, Jackson RW, Selengut J, Dodson R, Brinkac LM, et al. Whole-
712   genome sequence analysis of *Pseudomonas syringae* pv. *phaseolicola* 1448A reveals
713   divergence among pathovars in genes involved in virulence and transposition. J Bacteriol.
714   2005;187(18):6488-98.

715   75. Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, et al. InterProScan:
716   protein domains identifier. Nucleic Acids Res. 2005;33(Web Server issue):W116-20.

717   76. Davis MR, Jr., Goldberg JB. Purification and visualization of lipopolysaccharide from
718   Gram-negative bacteria by hot aqueous-phenol extraction. J Vis Exp. 2012(63).

77. Baltrus DA, Nishimura MT, Dougherty KM, Biswas S, Mukhtar MS, Vicente J, et al. The
molecular basis of host specialization in bean pathovars of *Pseudomonas syringae*.
Molecular Plant-Microbe Interactions. 2012;25(7):877-88.

78. Feil H, Feil WS, Chain P, Larimer F, DiBartolo G, Copeland A, et al. Comparison of the
complete genome sequences of *Pseudomonas syringae* pv. *syringae* B728a and pv. *tomato*
DC3000. Proc Natl Acad Sci USA. 2005;102(31):11064-9.

Table 1. Tailocin sensitivity, predicted loci of mutation and effect on target gene of tailocin
high-persistent and resistant mutants and selected allele-swapped strains

Strain/mutant	Population reduction after 900 AU tailocin (log CFU/ml)	Mutated locus	Gene involved	Change
Pph	5.61±0.28	WT	none	none
HPL	1.66±0.12	PSPPH_0964	Hypothetical protein	16 bp deletion, frameshift, truncation
HPL in WT*	1.56±0.11	Same as in HPL		
WT in HPL*	5.32±0.27	Same as WT		
IR1	0.48±0.08	PSPPH_3226	Glycosyl transferase	T>P
IR2	0.78±0.28	PSPPH_2810	4-amino-4-deoxy-L-arabinose transferase	S>I
IR3	0.33±0.14	PSPPH_0960	Glycosyl transferase family 2	(N>I)
IR4	0.12±0.05	PSPPH_0957	Glycosyl transferase 1	T insertion, extra AA added at the end
IR4 in WT*	0.04±0.09	Same as in IR4		
WT in IR4*	5.37±0.15	Same as WT		
IR5	-0.1±0.07	PSPPH_0520	Glycosyl transferase	H>R
IR6	-0.260±0.06	PSPPH_0963	FAD- dependent oxidoreductase	100 bp insertion, truncated protein (mite)
R1	-0.26±0.15	PSPPH_0957	Glycosyl transferase 1	C>A substitution, truncated protein
R1 in WT*	-0.48±0.16	Same as in R1		
WT in R1*	5.26±0.19	Same as WT		
R2	-0.17±0.08	PSPPH_0957	Glycosyl transferase 1	E>K
R3	-0.02±0.05	PSPPH_0952	Nucleoside-diphosphate-sugar epimerase	R>W
R3 in WT*	-0.01±0.04	Same as in R3		
WT in R3*	5.31±0.24	Same as WT		
R4	-0.45±0.11	PSPPH_0983	Glycosyl transferase	Frameshift
R4 in WT*	-0.48±0.05	Same as in R4		
WT in R4*	5.62±0.19	Same as WT		

WT=wild type, HPL= high persister-like, IR= incomplete resistant, R= resistant, * indicates allele-swapped strains to confirm
mutant phenotype.

733   **Table 2.** Bacterial strains and plasmids used in this study

Strain	Background	Characteristics	Source
B728a	*P. syringae* pv. *syringae* (*Psy*)	Tailocin producing WT strain	(78)
B728a:ΔRrbp	*Psy*	A tailocin deficient B728a mutant	(31)
1448a	*P. syringae* pv. *syringae* (*Pph*)	A tailocin sensitive WT strain	(74)
Δ*hrpL:Pph*	*Pph*	Type III secretion mutant of *Pph*	(77)
S17-1	*Eschericia coli*	Strain that can conjugate *P. syringae*	
pDONR207	plasmid	Gate way entry clone (Gm$_R$)	Invitrogen
pMTN1907	plasmid	Gateway compatible destination vector (Km$_R$,Tet$_R$, Sucs)	(77)
pDONR1K18ms	plasmid	Gateway vector (Km$_R$,Cm$_R$, Sucs)	Unpublished Gifted by Dr. Brian Kvitko, University of Georgia

744   **Table 3.** Primers used in this study

Primer	Sequence (5'-3')*	Description
HPL_upF	GGGGACAAGTTTGTACAAAAAAGCAGGCTCC GAGCCGCTGATCAACTC	Amplify the mutant and WT allele of HPL region
HPL_DnR	GGGGACCACTTTGTACAAGAAAGCTGGGTG ACCGACGTAGAAAAAGCA	
HPL_outF	ATCAAGGACTGGAAGACAATC	Target outside of the construct region and used for mutant confirmation by sequencing
HPL_outR	CAGCTACCACACTGACTATCG	
HPL_seqF	GTCTTCAACTCCAAAGTCACC	Target within the construct region and used for mutant confirmation
HPL_seqR	GCATTGTTGTAGACGTCAAAG	
HPL_testF	ATCGTTACCCATCTCTTTGTC	Diagnostic primers to selectively amplify the WT allele of HPL
HPL_testR	CGAAAAAGTGACAACCAATC	
M13F	TGTAAAACGACGGCCAGT	Confirmation of successful cloning in pDONR1k18ms
PK18M13R	AACAGCTATGACATGA	
BRM_11_15_16_5'S	GGGGACAAGTTTGTACAAAAAAGCAGGCTCCGA GCAGCTACAGCAGG	Amplify the mutant and WT allele of IR4 and R1 region
BRM_11_15_16_3'AS	GGGGACCACTTTGTACAAGAAAGCTGGGTGCGA GATAACCGTGCAC	
BRM_11_5'S	TGCGAGATCACAAAATT	Diagnostic primers to selectively amplify the WT allele of IR4
BRM_11_3'AS	CGCCTTCCACATTAA	
BRM_16_5'S	CCATTTCCGACTCCTC	Diagnostic primers to selectively amplify the WT allele of R1
BRM_16_3'AS	CTTCCTGGCGCG	
Pph_BRM_258_5'S	GGGGACAAGTTTGTACAAAAAAGCAGGCTCCAG ACGCCAGATTGAAGTTCTC	Amplify the mutant and WT allele of R3 region
Pph_BRM_258_3'AS	GGGGACCACTTTGTACAAGAAAGCTGGGTGCCC AACCGAAGTCGAGC	
Pph_BRM_258_5'S_C	GGTGTAATTGAACGGCCT	Diagnostic primers to selectively amplify the WT allele of R3
Pph_BRM_258_3'AS_C	GCCCATCGTCATTGCCA	
Pph_BRM_328_5'S	GGGGACAAGTTTGTACAAAAAAGCAGGCTCCCG GCAGTCGAGAAAGC	Amplify the mutant and WT allele of R4 region
Pph_BRM_328_3'AS	GGGGACCACTTTGTACAAGAAAGCTGGGTGGGC ATCGGCATCTATTCC	
Pph_BRM_328_5'S_C	CCAGGTTTGAGGCCG	Diagnostic primers to selectively amplify the WT allele of R3
Pph_BRM_328_3'AS_C	GAGTGCTGGTTACACACG	

*Gateway recombination sites are underlined

Fig 1. Difference in tailocin survival between stationary and log phase cultures of _Pph_.

Cultures were treated with a lethal dose of tailocin (100 AU) and viable cells pre- and post-

treatment were enumerated. Experiments were repeated at least five times with 3-6 biological

replicates per time. Mean and standard error of mean are graphed. _P_<0.05 indicate significant

differences within grouped bars as analyzed in SAS 9.4 with proc Glimmix.

Fig 2. Time dependent dynamics of tailocin survial. **A)** Stationary and log cultures of _Pph_

were treated with tailocin (100 AU) and the viable population was enumerated before and 1, 4, 8,

and 24 hours post-treatment. Three independent experiments were performed with 3-6 biological

replicates per time. Mean and standard error of mean are graphed. Different letters represent

significant differences as analyzed in SAS 9.4 with proc Glimmix at _P_=0.05. **B)** Percentage of

surviving colony phenotye upon tailocin re-treatment. Randomly selected surviving colonies

(n=12-44 for each growth phase and hours of treatment) from the initial tailocin treatment were

sub-cultured and re-treated with tailocin, and percentage of the surviving phenotype was

calculated. Colonies recovered from three independent experiments were used for the re-

treatment to calculate this percentage.

Fig 3. **Dynamics of tailocin survival with concentrated tailocin treatment**. **A)** Cultures were

treated with high dose of tailocin (900 AU) and viable populations pre- and post-treatment were

determined. Experiments were repeated at least three times with 3-6 biological replicates per

time. Mean and standard error of mean are graphed. _P_<0.05 indicates significant difference

within grouped bars as analyzed in SAS 9.4 with proc Glimmix. **B)** Percentage of surviving

colony phenotype after treatment with concentrated (900 AU) tailocin for one hour. Although

most of the surviving colonies were either incomplete resistant or resistant, persistent cells were

maintained even at high tailocin concentration. Surviving colonies were tested during three

independently repeated experiments for both cultures. (Refer **Fig. S3** for visual details that leads

upto this dynamics.)

Fig 4. **Treatment response of tailocin persistent and resistant lines. A)** Reduction in the

population of tailocin persistent and resistant mutant lines upon re-treatment with tailocin. Log

cultures of each lines were treated with 900 AU of tailocin and change in the population was

calculated after an hour of tailocin treatment. At least three separate colonies of each lines were

tested and experiments were repeated a minimum of three times. Means of the difference in log

transformed viable population pre- and post treatment are graphed. Error bars indicate stantard

error of mean. **B)** Assessement of response of mutant lines to tailocin under overlay condition.

Dilutions of tailocins (shown on the left most column) were spotted over the culture lawn of each

of the lines. Yellow line indicates the dilution upto wchich visible killing was observed. HPL;

high persistent-like, IR; incomplet resistant, R; resistant.

Fig 5. Dynamics of tailocin survival for the high persistent-like (HPL) mutant. Cultures

were treated with various tailocin doses and viable cells pre- and post-treatment were

enumerated. Means and standard error from three independently repeated experiments with at

least three biological replicates per experiment are reported. More killing occurred at high

tailocin concentration and no difference in stationary and log phage was observed. *P*>0.05

indicate no significant differences within grouped bars as analyzed in SAS 9.4 with proc

Glimmix.

Fig 6. **A genomic region (PSPPH_0957- PSPPH_0964) of *Pph* that showed prominent effect**

on tailocin sensitivity and resistance. Mutations of this region gave rise to multiple phenotypes

that differed in tailocin persistence and resistance. Refer Table 2 for a complete list of genes at

this and other genomic locations.

Fig. 7. *In planta* characterization of tailocin high persistent, selected incomplete resistant,

and resistant mutants. Strains and mutants were syringe infiltrated into green bean leaves and

population change was monitered by harvesting infiltrated leaf discs and enumeration using

dilution plating. Experiments were repeated atleast twice with 8 replications per strain. A

representative experiment is presented. Error bars indicate standard error. Different letters

indicate significant difference ($P<0.05$) for a given time point.

Fig. 8. A visual representation of the various tailocin surviving sub-populations and their

phenotypes upon exposure to tailocin.

Supporting Information

Fig S1. Visual representation of the difference in tailocin survival between the stationary and log

phase upon 100 AU of tailocin treatment for one hour. St, Stationary phase culture, Log-

logarithmic phase culture.

Fig S2. Visual representation of the dynamics of persistent and resistant cells after 100 AU of

tailocin treatment for 1 and 24 hours. For stationary 1 (St 1), surviving cells did not grow upto 24

hours (indicated by no change in the viable cells 1 and 24 hours post treatment) and were

sensitive on re-exposure (i.e. persistent) . For stationary 2, although the first hour survivors were

sensitive, at 24 hours, population increased due to growth of resistant cells as indicated by their

insensitivity to tailocin upon re-exposure (lower panel plates).

Fig S3. Testing tailocin activity of supernatants recovered after treatment. Stationary (S) and Log

(L) cultures were treated as described above with tailocin for specified number of hours. The

treatments were centrifuged, and the supernatant was collected and filter-sterilized. Dilution of

the supernatant were tested with *Pph* overlay. Active tailocin particles were recovered from both

stationary and log phage treatments at all time points tested (stationary treatments from this

experiment gave rise to a lot of tolerant colonies, log had few). Two independent experiments

were performed with similar results.

Fig S4. Test of tailocin activity after mixing purified tailocin in dilutions of filter-sterilized

stationary- and log- phase culture supernatants. Although a slight inhibition of tailocin activity

was observed in undiluted stationary supernatants (left), no inhibition was observed after diluting

the supernatant (100-1000 fold) as was done for tailocin treatment of cells. After mixing with

tailocin, supernatants were incubated for one hour. Dilutions (as shown in the left panel) of

tailocin and supernatant mixtures were spotted on a *Pph* overlay. Experiment was repeated twice

with three biological replicates and two technical replicates per time (n=12 in total).

Fig S5. SDS-Page separation of LPS extracted from wild-type *(Pph),* tailocin high-persistent like

mutant of *Pph* (HPL), incomplete resistant (IR) and resistant (R) mutants of *Pph*. IR 7 was one

of the incomplete resistant mutants not included in other analyses. Band sizes indicate molecular

weight in kilodalton.

Fig. S6. *In planta* characterization of selected tailocin resistant mutants together with the

complemented strains (mutant allele was swapped to WT and vice-versa). Results from two

separate experiments with 8 replication per experiment are presented. All complete mutant

phenotypes at 24 and 48 hpi had significantly lower population levels compared to WT and

incomplete mutant.

Fig S7. Growth curve of *Pph* in KB. Cells were inoculated in in a 96-well culture plate with KB

and optical density was measured at 600nm every two hours until 24 hourrs. Growth curve

experiment was performed twice. One representive experiment is shown.

A)

Population change after treatment (log CFU/ml)

900 AU tailocin applied to log cultures for 1 hour

B)

R1 R2 IR4 IR3 IR6 HPL

PSPPH_0957   PSPPH_0960   PSPPH_0963   PSPPH_0964
Glycosyltransferase 1   Glycosyltransferase 2   FAD-dependent   hypothetical
                                            oxidoreductase   protein

→ Resistance (R)   → Incomplete Resistance (IR),   → High Persistent-like (HPL)

Sensitive Incomplete resistant Resistant Persister

Tailocin particle Genetic transition Physiological transition

Tailocin-mediated selection

Immune

Transition is difficult to reverse; reduced fitness within host

Decreased sensitivity

Transition is difficult to reverse; no fitness effects in host

Transition is easy to reverse; no fitness effects in host; can not replicate

Immune