The negative adipogenesis regulator DLK1 is transcriptionally regulated by TIS7 (IFRD1) and translationally by its orthologue SKMc15 (IFRD2)

[1]Ilja Vietor[*], [1,13]Domagoj Cikes, [1,14]Kati Piironen, [1]Theodora Vasakou, [2]David Heimdörfer [1,15]Ronald Gstir, [2]Matthias David Erlacher, [3]Ivan Tancevski, [3]Philipp Eller, [3]Egon Demetz, [4]Michael Hess, [5]Volker Kuhn (†), [6]Gerald Degenhart, [7,8]Jan Rozman, [9,10,11]Martin Klingenspor, [7,8,12]Martin Hrabe de Angelis, [1]Taras Valovka, [1, 14]Lukas A. Huber

[1]Institute of Cell Biology, Biocenter, [2]Division of Genomics and RNomics, Biocenter, [3]Department of Internal Medicine II, [4]Division of Histology and Embryology, [5]Department Trauma Surgery, [6]Department Radiology, Medical University Innsbruck, Austria, [7]German Mouse Clinic, Institute of Experimental Genetics, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany, [8]German Center for Diabetes Research (DZD), Neuherberg, Germany, [9]Chair of Molecular Nutritional Medicine, Technical University of Munich, School of Life Sciences, Weihenstephan, Freising, Germany, [10]EKFZ - Else Kröner Fresenius Center for Nutritional Medicine, Technical University of Munich, Freising, Germany, [11]ZIEL - Institute for Food & Health, Technical University of Munich, Freising, Germany, [12]Chair of Experimental Genetics, Technical University of Munich, School of Life Sciences, Weihenstephan, Freising, Germany, [13]IMBA, Institute of Molecular Biotechnology of the Austrian Academy of Sciences, Vienna, Austria, [14]Division of Pharmaceutical Chemistry and Technology, Faculty of Pharmacy, University of Helsinki, Finland, [15]ADSI – Austrian Drug Screening Institute GmbH, Innsbruck, Austria

* Address correspondence to: Ilja Vietor, Institute of Cell Biology, Biocenter, Innsbruck Medical University, Innrain 80-82, A-6020 Innsbruck, Austria. Phone: +43 512 9003 70175 E-mail: ilja.vietor@i-med.ac.at

† Deceased May 2019

Keywords: adipocyte differentiation, diet-induced obesity, lean phenotype, translational inhibition

Running title: Identification of novel regulators of adipogenesis

Conflict of interest: The authors declare that they have no conflict of interest.

SYNOPSIS

This study uncovered that IFRD1 (TIS7) and its orthologue IFRD2 (SKMc15) are two essential regulators of adipogenesis. These proteins are highly similar on the sequence level, yet they regulate adipocyte differentiation using different but complementary mechanisms. Our main findings are:

- IFRD1 (TIS7) and IFRD2 (SKMc15) knockout mice are resistant against diet-induced obesity
- IFRD1 (TIS7) and IFRD2 (SKMc15) are critical for proper nutritional fat uptake and adipogenesis
- IFRD1 (TIS7) controls adipogenesis via Wnt/β-catenin-dependent transcriptional regulation of adipocyte-specific genes
- IFRD2 (SKMc15) regulates adipocyte-specific genes acting as a novel general translational inhibitor

Abstract

Delta-like homolog 1 (DLK1), an inhibitor of adipogenesis, controls the cell fate of adipocyte

progenitors. Here we identify two independent regulatory mechanisms, transcriptional and

translational, by which TIS7 (IFRD1) and its orthologue SKMc15 (IFRD2) regulate DLK1

levels. Mice deficient in both TIS7 and SKMc15 (dKO) had severely reduced adipose tissue and

were resistant to high fat diet-induced obesity. Wnt signaling, a negative regulator of adipocyte

differentiation was significantly up regulated in dKO mice. Elevated levels of the Wnt/β-catenin

target protein Dlk-1 inhibited the expression of adipogenesis regulators PPARγ and C/EBPα, and

fatty acid transporter CD36. Although both, TIS7 and SKMc15, contributed to this phenotype,

they utilized two different mechanisms. TIS7 acted by controlling Wnt signaling and thereby

transcriptional regulation of Dlk-1. On the other hand, here we provide distinctive experimental

evidence that SKMc15 acts as a general translational inhibitor significantly affecting DLK-1

protein levels. Our study provides data describing novel mechanisms of DLK1 regulation in

adipocyte differentiation involving TIS7 and SKMc15.

46   *Nonstandard abbreviations used:* CD36, cluster of differentiation 36; HFD, high fat diet; IFRD,

47   interferon related developmental regulator; KO, knockout; dKO, double knockout; RD, regular

48   diet; SKMc15, skeletal muscle cDNA library clone15; SVF, stromal vascular fraction; TIS7,

49   TPA Induced Sequence 7

Introduction

Adipogenesis is a complex process in which multipotent stem cells are converted into preadipocytes before terminal differentiation into adipocytes (Sarantopoulos, Banyard et al., 2018). These mechanisms involve protein factor regulators, epigenetic factors, and miRNAs. TPA Induced Sequence 7 (TIS7) protein has been shown to be involved in the mainly transcriptional regulation of differentiation processes in various cell types, e.g. neurons (Iacopetti, Barsacchi et al., 1996), enterocytes (Wang, Iordanov et al., 2005), myocytes (Vadivelu, Kurzbauer et al., 2004) and also adipocytes (Nakamura, Hinoi et al., 2013).

Multiple lines of evidence link the regulation of Wnt/β-catenin signaling to the physiological function of TIS7, known in human as Interferon related developmental regulator 1 (IFRD1) (Iezaki, Onishi et al., 2016, Vietor, Kurzbauer et al., 2005). Our data based on experiments with TIS7 knockout mice show a negative effect of TIS7 on Wnt signaling and a positive on adipocyte differentiation (Vietor et al., 2005, Yu, 2010 #3955). TIS7 deficiency leads to a significant up regulation of Wnt/β-catenin transcriptional activity in both primary osteoblasts and in mouse embryonic fibroblasts (MEFs) derived from TIS7 knockout (KO) mice. It was shown that TIS7 is also involved in the control of adipocytes differentiation in mice and was up regulated in both visceral (vWAT) and subcutaneous (sWAT) white adipose tissues (WAT) of genetically obese ob/ob mice (Nakamura et al., 2013). TIS7 transgenic mice have increased total body adiposity and decreased lean mass compared with wild type (WT) littermates (Wang et al., 2005). On HFD, TIS7 transgenic mice exhibit a more rapid and proportionately greater gain in body weight with persistently elevated total body adiposity. Enhanced triglyceride (TG) absorption in the gut of TIS7 transgenic mice (Wang et al., 2005) indicated that TIS7 expressed in the gut epithelium has direct effects on fat absorption in enterocytes. We have shown that as a result of impaired intestinal lipid absorption TIS7 KO mice displayed lower body adiposity (Yu, Jiang et al., 2010). Compared with WT littermates, TIS7 KO mice do not gain weight when

chronically fed an HFD and TIS7 deletion results in delayed lipid absorption and altered
intestinal and hepatic lipid trafficking, with reduced intestinal TG, cholesterol, and free fatty acid
mucosal levels in the jejunum (Garcia, Wakeman et al., 2014). TIS7 protein functions as a
transcriptional co-regulator (Micheli, Leonardi et al., 2005) due to its interaction with protein
complexes containing either histone deacetylases (HDAC) (Park, Horie et al., 2017, Vadivelu et
al., 2004, Vietor, Vadivelu et al., 2002, Wick, Schleiffer et al., 2004) or protein methyl
transferases, in particular PRMT5 (Lammirato, Patsch et al., 2016). The analysis of adipocyte
differentiation in preadipocytic 3T3-L1 cells suggested an involvement of TIS7 in the regulation
of adipogenesis in the Wnt/β-catenin signaling context (Nakamura et al., 2013).

SKMc15, also known as Interferon related developmental regulator 2 (IFRD2), a second member
of TIS7 gene family, is highly conserved in different species (Latif, Duh et al., 1997). Mouse
TIS7 and SKMc15 are highly homologous, with a remarkable identity at both the cDNA and
amino acid levels (58% and 88%, respectively). However, there was so far no information about
the physiological function and mechanisms of action of SKMc15 and its possible involvement in
differentiation of various tissues. Recently, cryo-electronmicroscopy (cryo-EM) analyses of
inactive ribosomes identified IFRD2 (SKMc15) as a novel ribosome-binding protein inhibiting
translation to regulate gene expression (Brown, Baird et al., 2018). The physiological function of
SKMc15 matching the mechanism based on the above mentioned cryo-EM data was so far not
shown and we postulate it could be involved in adipogenesis, since a significant reduction of
whole protein synthesis was previously shown as a major regulatory event during early
adipogenic differentiation (Marcon, Holetz et al., 2017).

Adipogenesis occurs late in embryonic development and in postnatal periods. Adipogenic
transcription factors CCAAT/enhancer binding protein α (C/EBPα) and peroxisome proliferator
activated receptor γ (PPARγ) play critical roles in adipogenesis and in the induction of adipocyte
markers (Farmer, 2006). PPARγ is the major downstream target of Delta-like protein 1 (DLK-1).

It is inactivated by the induction of the MEK/ERK pathway leading to its phosphorylation and

proteolytic degradation (Wang & Sul, 2009). DLK-1, also known as Pref-1 (preadipocyte factor

1) activates the MEK/ERK pathway to inhibit adipocyte differentiation (Kim, Kim et al., 2007).

C/EBPα is highly expressed in mature adipocytes and can bind DNA together with PPARγ to a

variety of respective target genes (Lefterova, Zhang et al., 2008). Besides that, PPARγ binding to

C/EBPα gene induces its transcription, thereby creating a positive feedback loop (Lowell, 1999).

Both proteins have synergistic effects on the differentiation of adipocytes that requires a

balanced expression of both C/EBPα and PPARγ.

Wnt/β-catenin signaling is one of the extracellular signaling pathways specifically affecting

adipogenesis (Li, Luo et al., 2008, Ross, Hemati et al., 2000, van Tienen, Laeremans et al., 2009)

by maintaining preadipocytes in an undifferentiated state through inhibition of C/EBPα and

PPARγ (Tontonoz & Spiegelman, 2008). PPAR-γ and Wnt/β-catenin pathways are regarded as

master mediators of adipogenesis (Xu, Wang et al., 2016). Wnt signaling is a progenitor fate

determinator and negatively regulates preadipocyte proliferation through DLK-1 (Mortensen,

Jensen et al., 2012). Mice overexpressing DLK-1 are resistant to HFD-induced obesity, whereas

DLK-1 KO mice have accelerated adiposity (Moon, Smas et al., 2002). DLK-1 transgenic mice

show reduced expression of genes controlling lipid import (CD36) and synthesis (Srebp1c,

Pparγ) (Barclay, Nelson et al., 2011). DLK-1 expression coincides with altered recruitment of

PRMT5 and β-catenin to the DLK-1 promoter (Paul, Sardet et al., 2015). PRMT5 acts as a co-

activator of adipogenic gene expression and differentiation (LeBlanc, Konda et al., 2012). SRY

(sex determining region Y)-box 9 (Sox9), a transcription factor expressed in preadipocytes, is

down regulated preceding adipocyte differentiation. DLK-1 prevents down regulation of Sox9 by

activating ERK, resulting in inhibition of adipogenesis (Sul, 2009). The PRMT5- and histone-

associated protein COPR5, affects PRMT5 functions related to cell differentiation (Paul, Sardet

et al., 2012). Adipogenic conversion is delayed in MEFs derived from COPR5 KO mice and

WAT of COPR5 KO mice is reduced when compared to control mice. DLK-1 expression is up

regulated in COPR5 KO cells (Paul et al., 2015).

Here we show the involvement of TIS7 and SKMc15 in the process of adipocyte differentiation.

dKO mice had strongly decreased amounts of the body fat when fed with even regular, chow diet

and were resistant against the high fat diet induced obesity. We found two independent

molecular mechanisms through which TIS7 and SKMc15 fulfill this function. The fact that these

two genes use independent mechanisms of action supported by the observation that whole body

deficiency of both genes led to a stronger phenotype when compared to single knockouts of TIS7

or SKMc15. TIS7 regulates the Wnt signaling pathway activity and restricts DLK-1 protein

levels, thereby allowing adipocyte differentiation. In contrast, SKMc15 KO did not affect Wnt

signaling, but as we show here, cells lacking SKMc15 have significantly up regulated

translational activity. In addition, we identified strongly enriched Dlk-1 mRNA concentrations

specifically in polyribosomes isolated from SKMc15 knockout MEFs when compared to the WT

MEFs. This was true also for dKO, but not for the single TIS7 knockout cells. The ablation of

both TIS7 and SKMc15 genes significantly affected the expression of genes essential for

adipocyte differentiation and function. Since dKO mice render a substantially leaner phenotype

on chow diet, even without any challenge by HFD-induction, we propose that TIS7 and SKMc15

represent novel players in the process of physiological adipocyte differentiation.

143
Results

Mice lacking TIS7 and SKMc15 genes have lower body mass, less fat and are resistant against

HFD-induced obesity.

In order to clarify whether both, TIS7 and SKMc15 are involved in the regulation of the

adipocyte differentiation and if they act through same or different mechanisms, we have

generated mice lacking both genes by crossing TIS7 with SKMc15 single KO mice. dKO pups

were viable, and adult male and female mice were fertile. At birth, body weights of dKO and

WT mice were similar. Nevertheless, already during weaning, both the male and female dKO

mice failed to gain weight when compared to their WT littermates and this persisted in the

following weeks when the mice were fed regular diet (RD; chow diet) containing 11 % kcal of

fat. At 10 weeks of age, dKO mice displayed 30 % and later up to 44.9 % lower body weight

compared with WT mice (Fig 1A).

Based on Dual Energy X-ray Absorptiometry (DEXA) measurement, 6-month-old WT mice had

substantially higher amounts of fat than their dKO littermates (Fig 1B, left panel). The effect of

TIS7 and SKmc15 double knockout was even more pronounced when the total fat and lean mass

values were normalized to the body weight since the dKO mice were smaller than their WT

counterparts were. The percentage of fat was in the WT mice 37.7 ± 4 % vs. 6 ± 3 % of the total

body mass in dKO animals. Furthermore, the percentage of lean tissue mass in WT animals was

lower than in dKO animals (60 ± 4.6 % vs. 92 ± 3.14 %; Fig 1B, right panel). Nevertheless, the

dKO mice were not significantly smaller, since there was only a minor difference in body length,

including the tail differed between WT and dKO mice (Fig EV1A). Next, we analyzed the

contribution of TIS7 and SKMc15 to the whole-body fat content of mice. Three-dimensional

reconstruction of images based on micro-computed tomography (micro-CT) of sex/age-matched

adult mice disclosed that both TIS7 and SKMc15 single KO mice already had less abdominal fat

than WT controls and that the dKO of TIS7 and SKMc15 genes caused the strongest decrease in

abdominal fat content and size (Fig 1C). Quantitative analyses of micro-CT measurements

showed that TIS7 deficiency caused a substantial lack of the abdominal fat tissues (P=0.002

when compared to WT mice, Fig 1D). Whereas TIS7 KO mice had less fat mass but were not

significantly lighter than their WT littermates (Fig 1E), SKMc15 KO were lighter (P=0.007) and

leaner (P=0.0001) and dKO mice were both significantly lighter (P = 0.0001) and had

significantly less abdominal fat (P=0.006) than the WT mice (Fig 1D,E).

The indirect calorimetry trial with sex- and age-matched WT and dKO animals where did not

identify any significant difference in respiratory exchange ratios (RER = VCO2/VO2) of WT

and dKO mice (Fig EV1B). However, dKO mice showed substantially reduced body weight,

mainly because of lacking fat, despite identical food intake, activity and no major differences in

several metabolic parameters. To investigate potential links between TIS7, SKMc15 and obesity,

we studied the response of dKO mice to HFD. At 2 months of age, male mice were housed

individually and fed with HFD for 21 days. Food intake was measured every second day and

body weight was measured every fourth day. Feces were collected every second day to analyze

the composition of excreted lipids and blood samples were collected after the third week of HFD

feeding to measure the concentrations of hepatic and lipoprotein lipases, respectively. Already

within the first week of HFD feeding, WT mice gained more weight than the dKO mice (Fig 1F),

although there were no obvious differences in food consumption (Suppl fig 1A) or in levels of

lipolytic enzymes (Suppl fig 1B,C). These differences in body weight gain continued to increase

during the second and third week, at which time the body weight of WT mice increased

additionally for 30% (30.0 ± 2.3%) when compared with the beginning of the HFD feeding

period. In contrast, the weight of dKO animals increased only slightly (7.6 ± 1.6%) (Fig 1F).

Both genes, TIS7 and SKMc15 contributed to this phenotype and we could see a stronger effect

following their deletion (Fig 1F).

Adipocyte differentiation in dKO mice is inhibited due to up regulated DLK-1 levels.

A possible explanation of the lean phenotype of dKO mice was that TIS7 and SKMc15 regulate

adipocyte differentiation. Primary MEFs derived from totipotent cells of early mouse

mammalian embryos are capable of differentiating into adipocytes and they are versatile models

to study adipogenesis as well as mechanisms related to obesity such as genes, transcription

factors and signaling pathways implicated in the adipogenesis process (Ruiz-Ojeda, Ruperez et

al., 2016). To test whether TIS7 and SKMc15 are required for adipogenesis we treated MEFs

derived from WT, SKMc15 and TIS7 single and dKO mice using an established adipocyte

differentiation protocol (Wang, Sun et al., 2015). Expression levels of both TIS7 and SKMc15

mRNA increased during the adipocyte differentiation of WT MEFs. TIS7 expression reached

maximum levels representing 5.7-fold increase compared to proliferating WT MEFs on day 3

(Suppl fig 1D) and SKMc15 reached on day 5 the maximum of 2.5-fold expression levels of

proliferating MEFs (Suppl fig 1E). These data suggested that both proteins play a regulatory role

in adipogenesis, however differ in their mechanisms and timing. Eight days after initiation of

adipocyte differentiation a remarkable reduction of adipocyte differentiation potential in

SKMc15, TIS7 KO MEFs and dKO stromal vascular fraction (SVF) cells isolated from inguinal

WAT was observed, as characterized by the formation of lipid droplets stained by oil red O (Fig

2A). Quantification of this staining revealed that fat vacuole formation in cells derived from

SKMc15, TIS7 and dKO mice represented 23 %, 48 % and 12 % of the WT cells, respectively

(Fig 2B). Stable ectopic expression of SKMc15 significantly increased adipocyte differentiation

in both, single and double TIS7 and SKMc15 knockout MEF cell lines (Fig EV1C,D and

EV2A). Ectopic expression of TIS7 significantly induced the adipocyte differentiation in TIS7

single knockout MEFs (Fig EV1C). These data indicated that both, TIS7 and SKMc15 were

critical for adipocyte differentiation and that the defect in adipogenesis could be responsible for

the resistance of dKO mice to HFD-induced obesity.

Wnt/β-catenin signaling is an important regulatory pathway for adipocyte differentiation

(Prestwich & Macdougald, 2007). Consistently, others and we have shown that TIS7 deficiency

causes up regulation of Wnt/β-catenin transcriptional activity. Therefore, we have assessed Wnt

signaling activity in MEFs by measuring transcriptional activity using TOPflash, TCF-binding

luciferase reporter assays. As shown in Fig 2C, Wnt signaling activity, when compared to the

WT MEFs, was not regulated in SKMc15 KO, but highly significantly (P<0.0001) up regulated

in TIS7 KO and also in dKO MEFs. Supporting these findings, Western blot analyses also

identified increased β-catenin protein levels in inguinal WAT of TIS7 KO and in dKO mice

(P<0.0001), but not in SKMc15 KO mice (Fig 2D, left top panel) suggesting the involvement of

TIS7 but not SKMc15 in the Wnt signaling pathway regulation.

Next, we analyzed the expression levels of DLK-1, a negative regulator of adipogenesis and at

the same time known target of Wnt signaling (Paul et al., 2015). DLK-1 protein was significantly

up regulated in inguinal WAT isolated from TIS7 and SKMc15 KO as well as from dKO mice

(Fig 2D, right top panel). A significant (P<0.001) up regulation of DLK-1 mRNA and protein

levels in dKO inguinal WAT samples revealed the qPCR and confirmed Western blot analyses

(Fig 2E). Up regulation of DLK-1 in dKO mice both on RNA and protein results were confirmed

in undifferentiated SVF cells isolated from inguinal fat (Fig 2F,G). DLK-1 mRNA levels were

even stronger up regulated following the 8 days differentiation protocol of SVF cells (Fig 2H).

Moreover, whereas in WT MEFs the expression of DLK-1 was strongly up regulated only during

the first day of adipocyte differentiation and then over the next days declined to basal levels, in

dKO MEFs DLK-1 was up regulated (P<0.001) throughout the entire 8 days of differentiation

(Fig 3A). This result complemented protein analyses of lysates from 8 day differentiated

adipocytes (Fig EV2B middle panel). A rescue experiment confirmed that DLK-1 expression

was TIS7- and SKMc15-dependent. DLK-1 mRNA levels were analyzed by RT-qPCR in dKO

MEFs stably expressing TIS7 and/or SKMc15. Ectopic expression of TIS7, SKMc15 and mainly

their combination significantly (P<0.001) down regulated DLK-1 RNA and protein levels (Fig

3B). Accordingly, these experiments documented that TIS7 and/or SKMc15 were involved in the

regulation of DLK-1 expression, but the molecular mechanism remained unclear.

The adipocyte differentiation deficiency of dKO inguinal SVF cells suggested that C/EBPα and

PPARγ might also be regulated through TIS7 and/or SKMc15. It was previously shown that

elevated levels of the cleaved ectodomain of DLK1 have been correlated with reduced

expression of PPARγ (Lee, Villena et al., 2003). Therefore, we analyzed the differences in

PPARγ expression between WT and dKO inguinal SVF cells. While PPARγ and C/EBPα

mRNA levels were strongly induced in undifferentiated WT, these were barely detectable in

dKO SVF cells (Fig 3C,D). Similarly, we identified significantly decreased PPARγ and C/EBPα

mRNA levels in TIS7, SKMc15 and dKO inguinal SVF cells following 8 days adipocyte

differentiation protocol (Fig 3E,F).

Our earlier study revealed that TIS7 binds directly to DNA and via interaction with PRMT5

regulates gene expression (Lammirato et al., 2016). Therefore, we performed chromatin

immunoprecipitation (ChIP) experiments where we studied binding of TIS7 and SKMc15 as

well as of the transcription factor β-catenin and symmetrically dimethylated histone H4 at

arginine residue 3 (H4R3me2s) (Paul et al., 2015) to the regulatory elements of the DLK-1 gene.

In dKO MEFs treated 8 days with the adipocyte differentiation cocktail we found increased β-

catenin binding to the β-catenin/TCF binding site 2 of the Dlk-1 regulatory element when

compared to WT MEFs (Fig 3G). On the other hand, binding of H4R3me2s to the same Dlk-1

regulatory element was significantly reduced (P<0.001). We could not identify any direct

binding of TIS7 or SKMc15 proteins to two different DLK-1 regulatory elements (Dlk-1 region

A and β-cat/TCFbs2), neither in WT nor in dKO MEFs, suggesting rather an epigenetic

regulation than *via* their direct binding to DLK-1 regulatory elements. We concluded that TIS7

and SKMc15 are required to restrain the DLK-1 levels, through the Wnt/β-catenin signaling

pathway and yet another so far unknown mechanism.

After finding significantly increased DLK-1 protein levels in inguinal WAT of SKMc15 single

knockout mice (Fig 2D) we measured DLK-1 expression in MEFs generated from these mice.

These were significantly increased (>70-fold) when compared to WT MEFs (Fig 3H). In a search

for a SKMc15-specific regulatory mechanism of Dlk-1 levels we focused on the translational

regulation. It was previously shown that general reduction of protein synthesis and

downregulation of the expression and translational efficiency of ribosomal proteins are events

crucial for the regulation of adipocyte differentiation (Marcon et al., 2017). Initially, we have

measured specifically polyribosome-bound DLK-1 RNA in SKMc15 knockout MEFs. Our

analysis identified significantly (P<8,26104 E-07) higher DLK-1 mRNA levels in polyribosome

RNA fraction of SKMc15 KO when compared to wild type MEFs (Fig 3I). Interestingly, IFRD2

(SKMc15) was recently identified as a novel specific factor capable of translationally

inactivating ribosomes (Brown et al., 2018). Because SKMc15 may play a crucial role in the

adipogenesis regulation we tested in our following experiment the effect of SKMc15 knockout

on the general translational efficiency of WT, TIS7, SKMc15 single and double KO MEFs. Cells

were incubated 30 minutes in the absence of methionine and cysteine, followed by one hour in

the presence of ^{35}S-methionine. As shown in Fig 4A, there was a significant increase in the

general translational activity of MEFs lacking SKMc15 alone or both TIS7 and SKMc15, but not

in TIS7 single knockout cells. Therefore, we concluded that SKMc15 alone, but not TIS7

inhibits the general translational activity necessary for the induction of adipogenic

differentiation, also via the translational regulation of Dlk-1. This finding supported the result

shown in Fig 2D where the DLK-1 protein levels were significantly induced in inguinal WAT

samples isolated from SKMc15 single knockout mice.

TIS7 and SKMc15 regulate adipocyte differentiation through DLK-1, MEK/ERK pathway,

PPARγ and C/EBPα.

DLK-1 protein carries a protease cleavage site in its extracellular domain (Lee, Helman et al.,

1995) and is secreted. The extracellular domain of DLK-1 is cleaved by ADAM17, TNF-α

Converting Enzyme to generate the biologically active soluble DLK-1 (Wang & Sul, 2009).

DLK-1 mRNA and protein levels are high in preadipocytes but DLK-1 expression is absent in

mature adipocytes. Hence, adding soluble DLK-1 to the medium inhibits adipogenesis (Garces,

Ruiz-Hidalgo et al., 1999). To test if the dKO MEFs secreted DLK-1, cell culture media from

MEFs treated 8 days with the adipocyte differentiation cocktail were collected and analyzed by

Western blotting. As shown in Fig 4B, dKO cells secreted DLK-1 protein, but in an identical

volume of the cell culture medium from WT MEFs no DLK-1 could be detected. In contrary to

that, an unrelated secreted protein, namely Collagen I, was found in media of both, WT and dKO

MEFs in similar, in the medium of WT MEFs even slightly higher amounts. To prove that

secreted DLK-1 could inhibit adipocyte differentiation of dKO MEFs, we cultured WT MEFs

with conditioned medium from dKO MEFs. Adipocyte differentiation of WT MEFs was strongly

inhibited by the dKO MEFs-conditioned medium when compared to the control WT cells (Fig

4C, quantified in Fig 4D). Another indication that dKO inhibited adipocyte differentiation via

DLK-1 protein up regulation delivered the experiment where we specifically knocked down

DLK-1. We targeted either all Dlk1 mRNA splice variants (oligo shDLK-1 391) or only Dlk1

mRNA splice variants containing coding sequences for the protease site for extracellular

cleavage (oligo shDLK-1 393) as previously published in (Mortensen et al., 2012). Both DLK-1

knockdown constructs stably expressed in dKO MEFs significantly increased (P<0.001)

adipocyte differentiation as documented in Fig EV2C. Hes1 levels, together with DLK-1 are

continuously down regulated during the process of adipogenesis while PPARγ are rising (Huang,

Yang et al., 2010). In our experimental setup, down regulation of DLK-1 levels was paralleled

by significant (P<0.001) decrease in Hes1 mRNA levels (Fig EV3A). Opposite, treatment with a

recombinant DLK-1 protein or stable DLK-1 ectopic expression documented by qRT-PCR (Fig

EV3D) significantly (P<0.001) inhibited adipocyte differentiation of WT MEFs as shown in Fig

EV3B,C. This was accompanied by a significant (P<0.001) decrease in C/EBPα mRNA levels

(Fig EV3E). Furthermore, DLK-1 mRNA quantification (Fig 4E) documented that dKO MEF

cell lysates contained significant amounts of DLK-1 mRNA when compared to WT control or

WT cells treated with dKO MEFs-conditioned medium. In parallel, WT cells treated with dKO

MEFs-conditioned medium expressed significantly lower amounts of PPARγ mRNA when

compared to WT cells incubated with control medium (Fig 4F). In addition, ectopic expression

of SKMc15 and co-expression with TIS7 in dKO MEFs rescued almost up to WT levels the

adipocyte differentiation potential of these cells (Fig EV2A). Ectopic expression of both, TIS7

and SKMc15 significantly (P<0.001) down regulated DLK-1 mRNA expression in dKO MEFs

(Fig EV2A). Moreover, conditioned medium from dKO MEFs expressing DLK-1 shRNA

knockdown constructs significantly (P<0.001) lost the ability to inhibit adipocyte differentiation

of WT MEFs as the medium from dKO MEFs did (Fig EV3F). Based on these results we

concluded that cells derived from dKO mice express increased DLK-1 levels and secreted,

soluble DLK-1 may inhibit adipocyte differentiation *in vivo*.

Previous studies showed that soluble DLK-1 activates MEK/ERK signaling, which is required

for inhibition of adipogenesis (Kim et al., 2007). As DLK-1 was strongly up regulated in dKO

SVF cells during adipocyte differentiation we subsequently analyzed possible activation of the

MEK/ERK pathway in gonadal WAT samples of WT and dKO mice (Fig 4G). The

phosphorylation of p44 and of p42 was up regulated 1.8-fold (P<0.001) in the dKO when

compared to the WT G WAT samples (Fig 4G). Next, we addressed the question of expression

levels of PPARγ and C/EBPα in WAT depots. Both adipocyte differentiation regulators PPARγ

and C/EBPα mRNA expression levels were in gonadal WAT samples isolated from dKO mice

significantly (P<0.01) downregulated when compared to the values of WT control animals (Fig

4H). Furthermore, we analyzed the possible effect of TIS7, SKMc15 and their combined

knockout on the DLK-1 levels in brown adipose tissue (BAT). Western blot analysis identified

significant (P<0.001) increase in DLK-1 protein levels in BAT samples in knockout mice of all

three genotypes (Fig 4I). The MEK/ERK pathway was similarly as in gonadal WAT, up

regulated also in MEFs generated from dKO mice (Fig 5A). The phosphorylation of p42 and of

p44 was up regulated 3-fold and 4.3-fold, respectively (P<0.01), in the adipocyte-differentiated

dKO when compared to the WT MEFs (Fig 5B). Previously, activation of MEK/ERK by DLK-1

was shown to up regulate the expression of the transcription factor SOX9, resulting in the

inhibition of adipogenesis (Sul, 2009). Therefore, we measured SOX9 mRNA expression by RT

qPCR in 8 days adipocyte-differentiated MEFs. SOX9 expression was significantly (P<0.001) up

regulated in 8 days adipocyte-differentiated dKO when compared to WT MEFs (Fig 5C). PPARγ

and C/EBPα mRNA levels were both continuously up regulated during the 8 days differentiation

protocol of WT MEFs (Fig 5D,E). On the contrary, we did not find any significant increase in

PPARγ and C/EBPα mRNA levels in dKO MEFs (Fig 5D,E). In a rescue experiment we showed

that only the co-expression of TIS7 and SKMc15 strongly increased the expression of the

PPARγ in undifferentiated dKO MEFs (P<0.001), almost up to the levels of WT MEFs (Fig 5F).

The expression of C/EBPα was also strongly up regulated by the co-expression of TIS7 and

SKMc15 (P<0.001) (Fig 5G). Based on these results we concluded that TIS7 and SKMc15

regulate the expression of both PPARγ and C/EBPα, crucial regulators of adipocyte

differentiation.

Lipid absorption is reduced in TIS7 and SKMc15 double knockout mice.

dKO mice were despite identical food intake, activity and RER (Suppl fig 1A and Fig EV1B)

leaner than their WT littermates. Therefore, we tested the possibility that dKO mice store energy

ectopically. In the feces of dKO mice fed with high fat diet we identified significantly higher

(P<0.05) amounts of free fatty acids than in that of their WT siblings (Fig 6A). Secondly, the

energy content of dried feces from dKO mice determined by bomb calorimetry was significantly

higher (P<0.001) than that of WT mice (Fig 6B). PPARγ induces the expression of CD36, a very

long chain fatty acids (VLCFA) transporter in heart, skeletal muscle, and adipose tissues

(Coburn, Knapp et al., 2000). The regulation of CD36 by PPARγ contributes to the control of

blood lipids. Interestingly, CD36 null mice exhibit elevated circulating LCFA and triglycerides

levels consistent with the phenotype of dKO mice and CD36 deficiency partially protected from

HFD-induced insulin resistance (Wilson, Tran et al., 2016). Because of down regulated PPARγ

levels in adipose tissues of dKO mice we decided to study more in detail CD36 regulation of

these as well. In inguinal WAT from dKO mice, we found strongly reduced CD36 mRNA

expression levels (64 % of the WT values) (Fig 6C). Next, we analyzed CD36 in WT and dKO

MEFs before onset and during adipocyte differentiation. As long as CD36 mRNA expression

substantially increased in WT MEFs, there were almost undetectable transcript levels of CD36 in

dKO cells treated with the adipocyte differentiation cocktail (Fig 6D). Diacylglycerol

acyltransferase 1 (DGAT1), a protein associated with the enterocytic triglyceride absorption and

intracellular lipid processing (Nozaki, Tanaka et al., 1999) is besides CD36 another target gene

of adipogenesis master regulator PPARγ (Koliwad, Streeper et al., 2010). DGAT1 mRNA levels

are strongly up regulated during adipocyte differentiation (Cases, Smith et al., 1998), its

promoter region contains a PPARγ binding site (Ludwig, Mahley et al., 2002) and DGAT1 is

also negatively regulated by the MEK/ERK pathway (Tsai, Qiu et al., 2007). DGAT1 expression

was shown to be increased in TIS7 transgenic mice (Wang et al., 2005) and its expression was

decreased in the gut of high fat diet-fed TIS7 KO mice (Yu et al., 2010). Importantly, DGAT1

expression in adipocytes and inguinal WAT is up regulated by PPARγ activation (Koliwad et al.,

2010). Therefore, we measured DGAT1mRNA levels during the differentiation of WT and dKO

MEFs into adipocytes to analyze the role of TIS7 and SKMc15 on the regulation of this protein

involved in adipogenesis and triglyceride processing. As long as DGAT1 expression

substantially increased during the differentiation of WT MEFs, there was no difference in

DGAT1 mRNA levels in dKO cells treated with the adipocyte differentiation cocktail (Fig 6E).

We concluded that TIS7 and SKMc15 regulate expression of multiple proteins involved both, in

adipocyte differentiation and in fat uptake, thereby contributing to the lean phenotype of dKO

mice through multiple means.

Discussion

In this study, we show that simultaneous depletion of TIS7 and of its orthologue SKMc15, a

protein recently identified as a translational inhibitor, caused severe reduction of adipose tissues

and resistance against high fat-induced obesity in mice. We identified two parallel mechanisms

leading to a deficiency in adipocyte differentiation. Firstly, TIS7-regulated Wnt signaling

affected Dlk-1 transcription and secondly, our study provided the first experimental evidence for

SKMc15 acting as a translational inhibitor that among other proteins controls DLK-1 protein

levels, thereby contributing to the regulation of adipogenesis (Fig 6F).

dKO mice were phenotypically similar with both, CD36 deficient and DLK-1 transgenic mice,

namely in decreased amounts of WAT and resistance to HFD-induced obesity. Previous studies

showed that overexpression of TIS7 caused increased intestinal lipid transport resulting in

elevated body weight gain during HFD feeding (Wang et al., 2015). Knockout of TIS7 and

SKMc15 impaired absorption of free fatty acids from the lumen into enterocytes and reduced

rate of fat absorption from intestines into the circulation. Simultaneously, higher concentrations

of free fatty acids in the feces of dKO mice (Fig 6A) suggested that mice lacking TIS7 and

SKMc15 suffer from an intestinal lipid uptake deficiency, yet another contributing reason for the

lean phenotype of these mice. In dKO mice, we found inhibited PPARγ and C/EBPα levels,

induced MEK/ERK pathway, and decreased expression of CD36 and DGAT1, all hallmarks of

up regulated DLK-1, a known adipogenesis inhibitor. TIS7 and SKMc15 acted after the

commitment to the preadipocyte stage since the elevated DLK-1 and β-catenin levels found in

WAT of dKO mice were characteristic for the preadipocyte stage (Gautam, Khedgikar et al.,

2017).

It is known that disruption of Wnt signaling in embryonic fibroblasts results in spontaneous

adipocyte differentiation (Bennett, Hodge et al., 2003) and opposite, stabilized β-catenin keeps

cells in the preadipocyte stage (Ross et al., 2000). A subpopulation of the stromal vascular

fraction of adipose tissue is adipogenic and at the same time has a weaker Wnt/β-catenin signal

(Hu, Yang et al., 2015). Consistent with this knowledge we found up regulated β-catenin protein

levels and increased Wnt signaling activity in WAT of TIS7 single and dKO MEFs. It has to be

426 mentioned that our data were different to those by Nakamura *et al.* who showed that following

427 the TIS7 overexpression in adipocytes was up regulated Wnt/β-catenin signaling and inhibited

428 oil red O staining (Nakamura et al., 2013). However, one has to take into consideration the

429 difference in cell systems used in these two studies. All results presented in Nakamura's study

430 were obtained in 3T3-L1 cells fibroblasts following overexpression of TIS7. Opposite to that, we

431 have studied the role of endogenous TIS7 in cells derived from wild type or knockout mice.

432 Moreover, we have found ubiquitous TIS7 expression in all WT mouse organs without any pre-

433 treatment such as e.g. hypoxia. On the other hand, Nakamura *et al.* identified up regulated TIS7

434 expression levels in WAT of obesity model mice. This result, however, fully supports our

435 findings of lean phenotype in TIS7 and SKMc15 dKO mice.

436 We demonstrated that in dKO MEFs, induced for adipocyte differentiation, the increase in Wnt

437 signaling led to up regulated binding of β-catenin to the DLK-1 gene regulatory elements,

438 resulting in sustained DLK-1 expression. In adipogenesis DLK-1 expression is down regulated

439 through histone methylation by COPR5 and PRMT5 that prevent β-catenin binding to the DLK-

440 1 gene (Paul et al., 2015). Interestingly, in dKO MEFs we found weaker binding of dimethylated

441 H4R3 to the DLK-1 gene that was consistent with our previous findings on the negative role of

442 TIS7 in epigenetic regulation of gene expression including the PRMT5 activity (Lammirato et

443 al., 2016).

444 There was a difference in effects of ectopic expression of TIS7, SKMc15 and their co-expression

445 on DLK-1 levels (Fig 3B), confirming the hypothesis that TIS7 and SKMc15 regulate DLK-1

446 levels via two independent pathways/mechanisms. Despite the fact that SKMc15 knockout had

447 no effect on Wnt signaling it nevertheless affected DLK-1 levels, suggesting a contribution of

448 SKMc15 to its regulation. Since SKMc15 was identified as a novel factor translationally

449 inactivating ribosomes (Brown et al., 2018) and given that down regulation of the protein

450 synthesis machinery is an essential regulatory event during early adipogenic differentiation

(Marcon et al., 2017), we proposed that SKMc15 regulates DLK-1 levels through translational

regulation. Experiments presented here confirmed specific regulation of DLK-1 through this

mechanism. Elevated DLK-1 levels in dKO MEFs activated the MEK/ERK pathway thereby

decreasing PPARγ and C/EBPα levels important for adipogenic differentiation. Besides that, the

expression of Sox9 and Hes1 were up regulated, suggesting that the dKO MEFs keep their

proliferative state and cannot enter differentiation into mature adipocytes (Kim et al., 2007).

Down regulation of PPARγ and C/EBPα implied changes in expression of downstream

adipogenesis-related genes. Among them, decreased CD36 and DGAT1 levels were a plausible

explanation of the dKO mice lean phenotype since it is known that both proteins play a

functional role in differentiation of murine adipocytes and their deficiency impairs fat pad

formation independent of lipid uptake (Christiaens, Van Hul et al., 2012). Our findings implicate

that TIS7 and SKMc15 play a up to now unknown role in the regulation of CD36 and therefore

possibly contribute to CD36 related pathogenesis of human metabolic diseases, such as

hypoglycemia (Nagasaka, Yorifuji et al., 2011), hypertriglyceridemia (Kashiwagi, Tomiyama et

al., 2001), and disordered fatty acid metabolism (Glazier, Scott et al., 2002, Tanaka, Nakata et

al., 2001).

We and others have previously shown that TIS7 is involved under both, chow and HFD

conditions in intestinal triglyceride absorption. Here we show for the first time the role of TIS7

and its orthologue SKMc15 in adipocyte differentiation. Moreover, our data presented here

confirm the physiological function of SKMc15 as a translational inhibitor. To our surprise,

although TIS7 and SKMc15 share high sequence homology and a functional role in the

adipogenesis, they use two independent regulatory mechanisms.

Methods

Generation of animal models. All animal experiments were performed in accordance with

Austrian legislation BGB1 Nr. 501/1988 i.d.F. 162/2005).

Mice lacking TIS7 were previously described (Vadivelu et al., 2004). SKMc15 KO mice

generation: targeting construct contained SKMc15 gene locus exons. A loxP site and a neomycin

resistance gene inserted at position 105515 (AY162905). The neomycin cassette was flanked by

two frt sites. A second loxP site was inserted at position 102082 (AY162905). Further

downstream, 15 additional nucleotides, part of intron 1, exon 2 (splice site: donor and acceptor)

and the CDS of hrGFP from the Vitality hrGFP mammalian expression vector pIRES-hrGFP-2a

(Stratagene) were added. The targeted construct was electroporated into Sv129 mouse ES cells.

After the selection with G418 single cell clones were screened by PCR and confirmed by

Southern blot analysis. After Cre recombinase treatment cell clones were screened by PCR.

Clones with floxed gene deleted, were used for blastocyst injection into C57Bl/6J mice. Two

male chimeras with ≥ 40 % or more agouti coat color were mated to C57Bl6 females. The

knockout mouse strain was derived from one male mouse carrying the allele of interest.

Heterozygous mice were back-crossed to C57Bl6 mice for 9 generations. dKO mice were

generated by crossing the SKMc15 KO mice with the TIS7 single KO mice. The resulting mice

were screened by PCR and double heterozygous mice used for further breeding until

homozygosity. In order to achieve maximal homogeneity of experimental groups, in all

experiments presented here we used only male mice.

SKMc15 knockout Southern blot analysis. XbaI restriction sites (position 109042; position

102079; position 92253) located in the SKMc15 locus (AY162905). Fragment detected by the

Southern probe: 9.6 kb wt (Suppl fig 1F, insert, band V); 15 kb in deleted locus (band IV). The

566 bp long probe for hybridization from genomic DNA (96605 – 97171; AY162905). PCR

primer sequences: RK 150 Fwd: 5'- GGTCCTGCCACTAATGCACTG-3'; RK 151 Rev: 5'-

GCAGACAGATGCCAGGAAGAC-3'.

SKMc15 knockout PCR genotyping analysis. hrGFP insert in the 3´ UTR detected by primer

GFP2 5´-AGCCATACCACATTTGTAGAG-3´ and RK101 3´ UTR (5´-

TGATGATAGCTTCAAAGAGAA-3´; 100617 – 100591 of the SKMc15 locus (AY162905).

PCR product 1700 bp. SKMc15 detection: RG1; 5´-TGTGGCCTTTATCCTGAGTC-3´;

102286–102266) and RG2; 5´-TGGCTTCATTTACACTACTCCTT-3´; 101860 – 101882

primers (Suppl fig 1F). WT allele PCR product 426 bp and the targeted allele PCR product 1772

bp (Suppl fig 1G). TIS7 genotype tested as explained previously (Vadivelu et al., 2004) (Suppl

fig 1H).

Growth monitoring and body composition measurement. Mice weaned at 3 weeks, regular chow

diet, weighed weekly. Dual energy X-ray absorptiometry was measured with Norland scanner

(Fisher Biomedical). Micro-CT experiments performed using vivaCT 40 (Scanco Medical AG).

The scans were performed using 250 projections with 1024 samples, resulting in a 38μm

isotropic resolution. Tube settings: 45kV voltage, 177μA current, integration time 300ms per

projection. Image matrix 1024*1024 voxels and a grayscale depth of 16 bit. The length of the

image stack was individually dependent, starting from the cranial end of the fist lumbar

vertebrae, to the caudal end of the fifth lumbar vertebrae. The image reconstruction and post

processing were performed using the Scanco Medical system software V6.6. For the adipose

tissue evaluation an IPL (image processing language) script by Judex et al., provided by Scanco

medical AG was modified to the scanner individual parameters leading to 2 values lower

threshold than in the original script for the adipose tissue filters 76. The script calculated the total

abdominal volume without potential air in the cavities. A separation of subcutaneous and

visceral fat mass was used only for visualization. For the quantitative fat mass analysis we used

56 male 5 to 12 month old mice (mean age 9.35 ± 2.03 month) reflecting an adult to mid aged

cohort defined by Flurkey et al. 2007 (Neeland, Ayers et al., 2013). The development of adipose

tissue in healthy mice is stable between 4 and 12 months (Lemonnier, 1972). Therefore, sex and

age could have only very limited influence on the experimental results. For the quantitative

comparison, the percent contribution of the abdominal fat to the body weight was calculated

using a mean weight of 0.9196 g/ml for adipose tissue (Neeland et al., 2013). Statistics were

performed using an ANCOVA with a Bonferroni corrected Post Hoc testing on the µCT fat data

and the body mass data.

Metabolic measurements. The indirect calorimetry trial monitoring gas exchange, activity, and

food intake was conducted over 21 hours (PhenoMaster TSE Systems). Body mass and rectal

body temperature before and after the trial were measured. The genotype effects were

statistically analyzed using 1-way ANOVA. Food intake and energy expenditure were analyzed

using a linear model including body mass as a co-variate.

HFD feeding. Age-matched (7-week-old) male TIS7-/- SKMc15-/- and WT mice were caged

individually and maintained up to 3 weeks on a synthetic, high saturated fat (HFD) diet

(TD.88137; Ssniff). Animals were weighted every 4th day between 08:00 and 10:00. Small

intestines were harvested for oil red O staining to detect lipid accumulation. Intestines, liver,

muscles and adipose tissue were collected for total RNA and protein isolation. Unfixed intestines

were flushed with PBS using a syringe, embedded in Tissue-Tek (Sakura, 4583) and frozen in

liquid nitrogen for immunohistochemical analysis.

Quantitative food consumption and fecal fat determination. Adult mice were acclimatized to

individual caging and to the HFD for a week, monitored for weight gain and their food intake

daily. The daily food intake data were pooled for the following 7 days and the food intake was

estimated (g/day). Feces were collected daily and weighted for 7 days after second week of the

HFD consumption. 50 mg of dried feces were boiled in 1 ml alkaline methanol (1M

NaOH/Methanol, 1 3 v/v) at 80°C for 2 h after addition of 50 nmol 5α-Cholestane (Sigma,

C8003) as internal standard for neutral sterol analysis. After cooling down to room temperature,

neutral sterols were extracted using three times 3 ml of petroleum ether, boiling range 60–80°C.

The residual sample was diluted 1:9 with distilled water. 100 µl of the solution were subjected to

an enzymatic total bile acid measurement. The extracted neutral sterols were converted to

trimethylsilyl derivatives. Neutral sterol composition of prepared feces samples was determined

by capillary gas chromatography on an Agilent gas chromatograph (HP 6890) equipped with a

25 m×0.25 mm CP-Sil-19 fused silica column (Varian) and a Flame Ionization Detector. The

working conditions were the following: Injector temperature 280°C; pressure 16.0 psi; column

flow constant at 0.8 ml/min; oven temperature program: 240°C (4 min), 10°C/min to 280°C (27

min); detector temperature 300°C (Harchaoui, Franssen et al., 2009). Energy content of dried

egested feces samples (~ 1 g per sample) was determined by bomb calorimetry (IKA C 7000,

IKA, Staufen, Germany) (Pfluger, Kabra et al., 2015).

Antibodies, viral and cDNA constructs. Antibodies: anti-CD36 AbD Serotec (MCA2748),

Abcam (ab36977), p44/42 MAPK (Erk1/2) and Phospho-p44/42 MAPK (Erk1/2)

(Thr202/Tyr204) Cell Signaling Technology (9102, 9101), β-catenin antibody Sigma (C2206),

anti-DLK Abcam (ab119930), anti-histone H4R3me2s antibody Active Motif (61187). For ChIP

experiments were used anti-TIS7 (Vietor et al., 2002) and anti-SKMc15 (Vadivelu et al., 2004)

rabbit polyclonal antibodies previously proven for ChIP suitability in (Lammirato et al., 2016).

pTOPflash reporter construct was a gift from H. Clevers (University of Utrecht, Holland). TIS7

construct was described previously (Vietor et al., 2002). Partial cds of mSKMc15 was amplified

by PCR and cloned into pcDNA3.1(-)/MycHis6 (Invitrogen).

Cell culture and adipocyte differentiation. MEFs were generated from 16-day old embryos. After

dissection of head for genotyping, removal of limbs, liver and visceral organs embryos were

minced and incubated in 1 mg/mL collagenase (Sigma-Aldrich, C2674) 30 mins at 37°C.

Embryonic fibroblasts were maintained in growth medium containing DMEM high glucose (4.5

g/l); sodium pyruvate, L-glutamine, 10% FCS (Invitrogen, 41966029), 10%

penicillin/streptomycin at 37°C in 5% CO_2. Adipocyte differentiation treatment was medium

with 0.5 mM 3-isobutyl-1-methylxanthine (I5879), 1 µM dexamethasone (D4902), 5 µg/ml

insulin (I2643) and 1 µM Rosiglitazone (Sigma-Aldrich, R2408). After 3 days growth medium

containing 1 µg/ml insulin cells were differentiated for 8 days. To visualize lipid accumulation

adipocytes were washed with PBS, fixed with 6 % formaldehyde overnight, incubated with 60 %

isopropanol, air dried and then incubated with oil red O. Microscopic analysis was followed by

isopropanol elution and absorbance measurement at 490 nm.

Construction of expression plasmids and generation of stable cell lines

The pRRL CMV GFP Sin-18 plasmid (Zufferey et al., 1998) was used to generate TIS7,

SKMC15 and Dlk1 expression lentiviral constructs. For this, the corresponding cDNAs were

cloned into the BamHI and SalI sites of the pRRL CMV GFP Sin-18 plasmid. A cap-

independent translation enhancer (CITE) fused to the puromycin resistance pac gene and the

woodchuck hepatitis virus post-transcriptional regulatory element (WPRE) were introduced

downstream of the TIS7, SKMC15 and Dlk1 coding sequences. All DNA constructs were

verified by sequencing.

To generate the Dlk1 shRNA lentiviral vectors, oligonucleotides targeting either all Dlk1 mRNA

splice variants (Dlk1Total: 5'-GATCCCCAGATCGTAGCCGCAACCAATTCAAGAGATTGG

TTGCGGCTACGATCTTTTTTGGAAA-3') or only Dlk1 mRNA splice variants containing the

extracellular cleavage sequence (Dlk1PS: 5'-

GATCCCCTCCTGAAGGTGTCCATGAATTCAA

GAGATTCATGGACACCTTCAGGATTTTTGGAAA-3') (Mortensen et al., 2012) were fused

with the H1 promoter and cloned into the pRDI292 vector as reported (Reintjes et al., 2016). The

GFP-targeting pRDI-shRNA-GFP plasmid was used as a control (Reintjes et al., 2016).

The viral supernatants were obtained as described previously (Leitner, Jakschitz et al., 2022),

concentrated with Retro-X Concentrator (Clontech, Takara Bio) and used to infect WT, TIS7 and

SKMc15 single and dKO MEFs. The selection was carried out for 2 weeks in DMEM

supplemented with 10% (v/v) FBS, 100 U mL^{-1} penicillin and 100 µg mL^{-1} streptomycin, and 2

µg mL^{-1} puromycin.

Polysome profiling

Polysome profiling was performed as described in (Savant-Bhonsale & Cleveland, 1992), with

modifications. The day before harvesting the cells, continuous 15-45 % (w/v) sucrose gradients

were prepared in SW41 tubes (Beckman) in Polysome Gradient Buffer (10 mM Hepes-KOH, pH

7.6; 100 mM KCl; 5 mM MgCl2) employing the Gradient Master ip (Biocomp) and stored

overnight at 4 °C. All steps of the protein extraction were performed on ice. Exponentially

growing cells were washed twice with ice cold DPBS (Gibco) supplemented with 100 µg/mL f.

c. cycloheximide, scraped in 300 µL Polysome Lysis Buffer (10 mM Hepes-KOH, pH 7.6; 100

mM KCl; 5 mM MgCl2; 0.5 % IGEPAL® CA-630; 100 µg/mL cycloheximide) supplemented

with 0.1 U/µL murine RNase Inhibitor (NEB), and passed through a G25 needle 25 times. Nuclei

were pelleted at 16,000 g for 6 min at 4 °C and the supernatants were carefully layered onto the

sucrose gradients. Samples were centrifuged at 35,000 rpm for 2 h at 4 °C (with brakes switched

off) using an SW 41 Ti rotor (Beckmann). 20 fractions of 0.6 mL were collected by use of a

peristaltic Pump P1 (Amersham Biosciences) and polysome profiles were generated by optical

density measurement at 254 nm using optical unit UV-1 (Amershan Biosciences) and chart

recorder Rec 111 (Amersham Biosciences).

Analysis of translation by metabolic labelling. Pulse labelling of mitochondrial proteins was

performed as described before (Popow, Alleaume et al., 2015), with the following changes: 1x

10^6 wild-type and dKO MEFs cells per plate were seeded and cultivated for 24 h. Cells were

washed twice with PBS followed by one wash with labelling medium (ESC medium without

cysteine and methionine) and a 30 min incubation in labelling medium. To label newly translated

products, 200 µCi of [^{35}S]-methionine (10 mCi/ml; Hartman Analytic) were added and the cells

were incubated for 1 h. Cells were washed and incubated another 10 min in standard medium at

37 °C. Cells were then harvested by trypsinizing, washed once with ice-cold PBS, extracted with

ice-cold RIPA buffer containing protease-inhibitors and briefly sonicated. Proteins were

fractionated by gel electrophoresis in 16% Tricine gels (Invitrogen) and stained with Coomassie

brilliant blue, and radioactive signals were visualized by phosphorimaging. Signal intensities

were quantified using the Image Studio Lite (v5.2) software.

RT-PCR. Tissues from animals fed for 3 weeks with HFD were snap frozen and stored at –80°C.

Total RNA was isolated using the TRIzol reagent (Invitrogen, 15596026). RNA was then

chloroform-extracted and precipitated with isopropanol. The yield and purity of RNA was

determined by spectroscopic analysis; RNA was stored at –80°C until use.

Quantitative RT-PCR and statistics. Total RNA were treated with DNAse1 and reverse

transcribed to cDNA by Revert Aid First Strand cDNA Synthesis Kit (Thermo Scientific,

K1622) with oligo dT primers. Quantitative RT-PCR was performed using Taqman probes and

primer sets (Applied Biosystems) specific for CD36 (assay ID Mm00432398_m1), Dgat1

(Mm00515643_m1), Pparγ (Mm00440940_m1), C/EBPα (Mm00514283_s1), DLK-1

(Mm00494477_m1) and Sox9 (Mm00448840_m1). Ribosomal protein 20 (assay ID

Mm02342828_g1) was used as normalization control for quantification by the ddCt method.

PCR reactions were performed using 10 µl cDNA in PikoReal 96 real-time PCR system (Thermo

Scientific). Quantification data were analyzed by 2-tailed, Homoscedastic t-tests are based on the

assumption that variances between two sample data ranges are equal type 2 Student's t test.

Transient transfections and luciferase assay. pGL2-Basic (Promega, E1641) or pGLCD36

(Shore, Dietrich et al., 2002) were used as reporter constructs. Expression constructs or empty

vector DNA as a control were co-transfected. pCMV-β-Gal plasmid was used to normalize for

transfection efficiency. For luciferase reporter assays, 1.5x105 cells were seeded into 24-well

plates and transfected after 24 hours with indicated plasmid combinations using Lipofectamine

Plus™ Reagent (Invitrogen, 15338030). The total amount of transfected DNA (2 µg DNA per

well) was equalized by addition of empty vector DNA. Cells were harvested 48 h post-

transfection in 0.25 M Tris, pH 7.5, 1% Triton X-100 buffer and assayed for both luciferase and

β-galactosidase activities. Luciferase activity and β-galactosidase activity were assayed in

parallel by using the Lucy 2® detection system (Anthos). Transfections were performed in

triplicates and all experiments were repeated several times.

Chromatin immunoprecipitation (ChIP). Chromatin was isolated from TIS7 WT and dKO

formaldehyde-treated, 8 days adipocyte differentiated MEFs using the EpiSeeker Chromatin

Extraction Kit (Abcam, ab117152). ChIP analyses were carried out as described previously

(Reintjes et al., 2016). Sequence of the oligonucleotides for two regions of the Dlk-1 promoter,

encompassing TCF and β-catenin binding sites were as defined in (Paul et al., 2015). Sonicated

chromatin was centrifuged at 15.000x g for 10 min at 4°C, and the supernatant (65 µg of sheared

DNA per each IP) was diluted ten-fold with cold ChIP dilution buffer containing 16.7 mM Tris-

HCl pH 8.1, 167 mM NaCl, 0.01% (w/v) SDS, 1.1% (w/v) Triton X-100 and 1.2 mM EDTA

with protease inhibitors. Samples were pre-cleared for 1 h with protein A Sepharose CL-4B

(Sigma-Aldrich, 17-0780-01) beads blocked with 0.2 µg/µl sonicated herring sperm DNA

(Thermo Fisher, 15634017) and 0.5 µg/µl BSA (NEB, B9000 S). Immunoprecipitations were

performed at 4°C overnight. Immune complexes were collected with protein A Sepharose for 1h

at 4°C followed by centrifugation at 1000 rpm and 4°C for 5 min. Beads were washed with 1ml

low salt wash buffer (20 mM Tris-HCl pH 8.1, 150 mM NaCl, 0.1%(w/v) SDS, 1%(w/v) Triton

X-100 (Merck), 2 mM EDTA), high salt wash buffer (20 mM Tris-HCl pH 8.1, 500 mM NaCl,

0.1% (w/v) SDS, 1% (w/v) Triton X-100, 2 mM EDTA), LiCl wash buffer (10 mM Tris-HCl pH

8.1, 250 mM LiCl, 1% (w/v) sodium deoxycholate, 1% (w/v) IGEPAL-CA630, 1 mM EDTA)

for 5 min at 4°C on a rotating wheel, and twice with 1 ml TE buffer (10 mM Tris-HCl pH 8.0, 1

mM EDTA). Protein-DNA complexes were eluted from antibodies by adding a freshly prepared

elution buffer containing 1% SDS and 0.1 M $NaHCO_3$. The eluate was reverse cross linked by

adding NaCl to a final concentration of 0.2 M and incubating at 65 °C for four hours. Afterwards

the eluate was treated with Proteinase K at 45 °C for 1 hour. The immunoprecipitated DNA was

then isolated by phenol/chloroform precipitation and used as a template for real-time-quantitative

PCR. The primer pairs specific for regulatory regions of the Dlk-1 gene were selected as

described before (Paul et al., 2015). Reactions with rabbit IgG or with 1.23% of total chromatin

(input) were used as controls. For real-time-quantitative PCR a PikoReal System was used.

Signals were normalized to input chromatin and shown as % input. The raw cycle threshold (Ct)

values of the input were adjusted to 100% by calculating raw Ct – log2(100/input). To calculate

the % input of the immunoprecipitations, the equation $100 \times 2[Ct$ (adjusted input to 100%) – Ct

(IP)] was applied.

Statistical analyses

Statistical analyses were performed with ANOVA, Student's unpaired t-test using GraphPad

Prism Ver. 9.2 (GraphPad La Jolla,CA) software, or as indicated in the legends. P-value is

indicated by asterisks in the figures:*$P \leq 0.05$,**$P < 0.01$,***$P < 0.001$,****$P < 0.0001$. Data from

SVF cells were analyzed using ordinary one-way ANOVA with Holm-Šidák's multiple

comparisons test.

Acknowledgements

The authors thank Robert Kurzbauer for generation of dKO mice, Stephan Geley, Laura M. de

Smalen, Laura De Gaetano and Karin Schluifer for the technical assistance, Christiane Heim for

serum analyses and free fatty acid uptake measurements, Frans Stellaard for the analysis of fatty

acids content in feces, Mayra Eduardoff for RNA processing for Affymetrix chip analysis,

Alexander Magnutzki for advice with the statistical analyses of data, David Teis and Zlatko

Trajanoski for critical reading of the manuscript. Furthermore, we would like to thank Dr. Paul

Shore for providing us with the pGLCD36 construct. We are indebted to the staff at the Animal

Facility of Innsbruck Medical University for their care of our mice. This work was supported by

P18531-B12 and P22350-B12 grants from the Austrian FWF grant agency to Ilja Vietor and by

the German Federal Ministry of Education and Research (Infrafrontier grant 01KX1012) to

Martin Hrabe de Angelis.

Author contributions

I.V., D.C., M.E., I.T., G.D., J.R., T.V. and L.A.H. designed the study.

Experiments were performed by: Fig.1A DC, B VK+DC, C VK+GD, D,E GD, F DC; Fig.2A,B

ThV, C IV, D,E,F,G ThV, H IV+ThV+KP; Fig.3A KP, B IV+ThV, C,D,E,F ThV+KP, G

IV+TV, H,I IV+DH+ME+TV; Fig.4A IV+ME, B IV, C,D,E,F,G ThV, H ThV+KP, I ThV; Fig.5

KP; Fig.6 A,B IT+FS+DC, C,D,E KP; Fig.EV1A IV, B JR, C,D TV+ThV; Fig.EV2 TV+ThV;

Fig.EV3A TV+ThV, B,C, D, E IV+TV; F TV+ThV; Suppl.Fig.1 A JR, B,C IT+PE+ED, D KP,

E,F RG.

I.V., D.C., T.V. and L.A.H. wrote the manuscript.

Figure legends

Figure 1. dKO mice display postnatal growth retardation, less body fat and are resistant

against HFD-induced obesity

A) Growth curves of WT and dKO mice on chow diet (n fluctuate depending on the time

point of measurement).

B) DEXA measurements of WT and dKO mice. Left panel depicts the absolute values of fat

and lean mass per animal and the right panel represents values normalized to the total

body weight of animals (n=15).

C) Micro-computed tomography measurements identified a lack of abdominal fat in single

and dKO mice. 3-dimensional reconstitution of images of the abdominal fat mass

distribution in WT and KO mice. Yellow color represents subcutaneous and red visceral

fat mass. Black compartments are shadows resulting as part of the lightning model for the

3D volume rendering.

D) Mass contribution [%] of the abdominal fat in correlation to the total body weight [g]. A

linear regression is overlaid for each group individually. The regression results for the

WT group are y=0,342x-11,13 with a R^2 of 0,74; for the TIS7 KO group y=0,143x-3,91

with a R^2 of 0,82; for the SKMc15 KO y=0,005x+0,48 with a R^2 of 0,04 and for the dKO

y=0,011x+0,01 with a R^2 of 0,1. ANCOVA analysis for the fat mass as a percentage of

the body weigth as covariant was performed.

E) Effect of single SKMc15 or TIS7 knockout and TIS7 SKMc15 double knockout on total

body weight in [g] and body fat amount normalized to the body weight in [%].

F) TIS7 and SKMc15 knockout reduced the gain of body weight of mice fed with HFD. 7

weeks old male WT and dKO mice were caged individually and maintained up to 21 days

on HFD. Data shown are mean ± STD, n≥9 per genotype. Data were analyzed applying

one-way ANOVA with Holm-Šidák's multiple comparisons test. *P<0.05, **P<0.01,

**** P<0.0001.

Figure 2. Adipocyte differentiation, DLK-1 levels and Wnt signaling pathway are

significantly affected in dKO mice.

A) Representative oil red O staining of inguinal SVF cells following 8 days of the adipocyte

differentiation protocol. Scale bar is 20µm.

B) Quantification of oil red O staining from three independent adipocyte differentiation

experiments shown in panel A. Ordinary one-way ANOVA **** P<0.0001.

C) Wnt signaling activity was measured in MEFs transiently co-transfected with the Tcf

reporter plasmid pTOPflash and β-galactosidase expression vectors. Transfection

efficiency was normalized using the β-galactosidase values. The experiment was repeated

three times. Data shown are mean ± STD. Ordinary one-way ANOVA Holm-Šídák's

multiple comparisons test **** P<0.0001.

D) Left top panel: Western blot analysis of β-catenin amounts in inguinal WAT samples.

Representative Western blot from 3 biological repetitions is shown. Right top panel:

DLK-1 is significantly up regulated both in TIS7, SKMc15 single KO and in dKO mice.

Bottom panels: actin Western blots were used for normalization of sample loading.

E) DLK-1 mRNA expression measured by qPCR in inguinal WAT tissue samples (top).

Values were normalized on GAPDH, n=3. Error bars indicate standard deviations. A

representative Western blot image detecting DLK-1 and GAPDH proteins (bottom).

F) Dlk-1 mRNA expression detected in undifferentiated cells isolated from inguinal WAT

SVF cells. Normalized on GAPDH. Error bars indicate standard deviations, ***P<0.001.

G) DLK-1 protein Western blot analysis in undifferentiated cells isolated from inguinal

WAT SVF cells. Mean of 3 biological repeats; insert is one representative Western blot.

***P<0.001.

H) Dlk-1 mRNA expression detected in 8 days adipocyte differentiated cells isolated from

inguinal WAT SVF cells. Normalized on GAPDH. Error bars indicate standard

deviations, ***P<0.001.

Figure 3. TIS7 and SKMc15 regulate Dlk-1 and thereby affect adipocyte regulators

expression.

A) DLK-1 mRNA expression measured in MEFs treated with the adipocyte differentiation

cocktail for given times. WT MEFs values were set as 1. *P<0.05, ***P<0.001.

B) DLK-1 mRNA levels (top) and protein levels (bottom) in dKO MEFs were down

regulated following ectopic co-expression of TIS7 and SKMc15. mRNA expression

levels were analyzed by RT qPCR in stably transfected cells following 8 days adipocyte

protocol differentiation. WT MEFs values were set as 1, n=3; Error bars indicate standard

deviations. *** P<0.001

C) PPARγ mRNA expression detected in undifferentiated cells isolated from inguinal WAT

SVF cells. Normalized on GAPDH. Error bars indicate standard deviations, ***P<0.001.

D) C/EBPα mRNA expression detected in undifferentiated cells isolated from inguinal

WAT SVF cells. Normalized on GAPDH. Error bars indicate standard deviations,

***P<0.001.

E) PPARγ mRNA expression detected in 8 days adipocyte differentiated cells isolated from

inguinal WAT SVF cells. Normalized on GAPDH. Error bars indicate standard

deviations, ***P<0.001.

F) C/EBPα mRNA expression detected in 8 days adipocyte differentiated cells isolated from

inguinal WAT SVF cells. Normalized on GAPDH. Error bars indicate standard

deviations, ***P<0.001.

G) Recruitment of indicated proteins to regulatory regions of the Dlk-1 promoter in WT and

KO MEFs was analyzed by ChIP at day 8 of the adipocyte differentiation. Values are

expressed as the percentage of immunoprecipitated chromatin relative to input and are

the mean of triplicates. ChIP analysis identified increased specific β-catenin binding to its

DLK-1 regulatory element in dKO samples. Pre-immune serum and IgG were used as

background controls. n=3, Data shown are mean ± STD, Student's t-test ***P<0.001.

H) Dlk-1 mRNA expression detected in undifferentiated MEF cells. Normalized on

GAPDH. Error bars indicate standard deviations, ***P<0.001.

I) Real-time qPCR detection of Dlk-1 RNA in polyribosomes. Normalized on GAPDH.

Error bars indicate standard deviations, ***P<0.001.

Figure 4. SKMc15 inhibits DLK-1 translation. dKO-secreted DLK-1 inhibits adipocyte

differentiation through MEK/ERK signaling.

A) Translational analysis – *in vivo* metabolic staining of MEFs with ^{35}S-methionine. Equal

numbers of cells were seeded and 24 hours later treated as explained in Methods section.

Identical volumes of cell lysates were separated by SDS-PAGE, gel was dried and

analyzed by a phosphorimager. Equal loading was documented by the Coomassie blue

staining of the gel.

B) dKO MEFs secrete DLK-1 protein into the cell culture medium. Identical volumes of

media from WT and dKO MEF cells 8 days treated with the adipocyte differentiation

cocktail were analyzed by Western blot. Collagen I was present in both samples in

similar amounts. Equal loading of samples was evaluated by amido black staining of the

membrane.

C) dKO MEFs-conditioned medium inhibited adipocyte differentiation in WT MEFs. MEFs

were treated 8 days with the adipocyte differentiation cocktail. WT MEFs treated with

the dKO-conditioned medium (replaced 3 times every 2 days) showed reduced

differentiation. Images representing three biological repeats were cropped and space bars

represent always 20μm.

D) Quantification of oil red O staining from three independent adipocyte differentiation

experiments. Data shown are mean ± STD, ordinary one-way ANOVA P=0.0016.

Student's t-test, **P<0.01, *** P<0.001.

E) DLK1 mRNA levels in WT and dKO MEFs treated 8 days with the adipocyte

differentiation cocktail or in WT MEFs treated with the dKO-conditioned medium, ***

P<0.001.

F) PPARγ mRNA expression levels in same cells as shown in panel D and E, *** P<0.001.

G) Representative Western blots of phospho-p44, phospho-p42, p44 and p42 in gonadal

WAT of WT and dKO mice. Normalization on p44 and p42; n=3. Error bars indicate

standard deviations. WT values were set as 1, *** P<0.001.

H) PPARγ and C/EBPα mRNA expression in gonadal WAT. Normalization on GAPDH;

n=3. Error bars indicate standard deviations. WT values were set as 1, ** P<0.01.

I) DLK-1 protein is up regulated in brown adipose tissues (BAT) of SKMc15, TIS7 single

knockout and in dKO mice. Western blot analysis was performed on 5 samples of each

genotype. Normalization on GAPDH. Error bars indicate standard deviations,

***P<0.001.

Figure 5. MAPK signaling and Sox9 expression are induced while adipocyte differentiation

regulatory genes are down regulated in dKO MEFs.

A) Representative Western blots of phospho-p44, phospho-p42, p44 and p42 from 8 days

adipocyte differentiation cocktail-treated MEFs.

B) Quantitative analysis of Western blot data. Normalization on p44 and p42; n=3. Error

bars indicate standard deviations. WT values were set as 1; **P<0.01.

C) SOX9 expression was measured by qPCR in 8 days adipocyte-differentiated WT and

dKO MEFs. SOX9 values were normalized on GADPH expression. WT MEFs values

were set as 1, n=3. Data shown are mean ± STD. Student's t-test ***P<0.001.

D) PPARγ and

E) C/EBPα mRNA levels were down regulated in dKO adipocyte-differentiated MEFs.

Gene expression was measured by qPCR during the treatment with the adipocyte

differentiation cocktail. Values were normalized on GADPH expression. WT MEFs

values were set as 1, n=3. Data shown are mean ± STD, *** P<0.001.

F) Ectopic co-expression of TIS7 and SKMc15 in undifferentiated MEFs significantly

increased levels of adipogenic genes PPARγ and

G) C/EBPα. Data shown are mean ± STD, Student's t-test ***P<0.001.

Figure 6. FFA uptake is inhibited in dKO mice.

A) Free fatty acids concentrations in feces of WT and dKO mice. Feces were collected every

second day and the composition of excreted lipids was determined by capillary gas

chromatography. Data shown are mean ± STD, *P<0.05.

B) Energy content of dried egested feces samples was determined by bomb calorimetry.

Data shown are mean ± STD, **P<0.01.

C) qPCR analysis of CD36 mRNA levels in inguinal WAT. Data shown are mean ± STD,

*P<0.05.

D) qPCR analysis of CD36 in MEFs differentiating into adipocytes. Relative expression

levels were normalized on GAPDH expression. WT MEFs values were set as 1. Data

shown are mean ± STD, ***P<0.001.

E) DGAT1 mRNA levels were down regulated in dKO adipocyte-differentiated MEFs. Data

shown are mean ± STD, ***P<0.001.

F) Proposed model of TIS7 and SKMc15 molecular mechanisms of action during adipocyte

differentiation. Two parallel mechanisms leading to a deficiency in adipocyte

differentiation: TIS7-regulated Wnt signaling affected Dlk-1 transcription and SKMc15

acting as a translational inhibitor also contributed to the regulation of adipogenesis.

Figure EV1. Body length and respiratory exchange of dKO do not differ from WT mice.

TIS7 and SKMc15 positively regulate adipocyte differentiation of MEF cells.

A) Measurement of whole body length, including the tale of age- and sex-matched mice.

n=14 and 17, respectively. Data shown are mean ± STD, *P<0.01

B) Indirect calorimetry measurements of WT and dKO animals. No significant differences

were found in respiratory exchange ratios (RER) of WT and dKO mice.

C) Oil red O staining of MEFs following 8 days adipocyte differentiation protocol. Ectopic

expression of TIS7 and/or SKMc15 induced adipocyte differentiation of TIS7 KO MEFs,

size bar 20µm. Quantification of oil red O staining, n=3. Data shown are mean ± STD,

***P<0.001.

D) Ectopic expression of SKMc15 induced adipocyte differentiation of SKMc15 KO MEFs,

size bar 20µm. Quantification of oil red O staining, n=3. Data shown are mean ± STD,

***P<0.001.

900   Figure EV2. TIS7 and SKMc15 positively regulate adipocyte differentiation of MEF cells

901   and inhibit DLK-1 expression. DLK-1 inhibits adipocyte differentiation of dKO MEFs.

A) Ectopic expression of TIS7 and/or SKMc15 induced adipocyte differentiation of dKO

MEFs, size bar 20µm. Quantification of oil red O staining, n=3, *** P<0.001. Ectopic

expression of TIS7 and/or SKMc15 inhibited DLK-1 mRNA expression detected in dKO

MEF cells. Normalized on GAPDH. Error bars indicate standard deviations, WT MEFs

values were set as 1. Data shown are mean ± STD, ***P<0.001.

B) Cells from which the conditioned medium was transferred during the differentiation. Oil

red O staining (panels on the left) after 8 days of adipocyte differentiation. Western blot

detection of DLK-1 protein in differentiated cells.

C) Oil red O staining (left panel) of dKO control and shDLK-1 MEFs following 8 days

adipocyte differentiation protocol. DLK-1 knockdown induced adipocyte differentiation

of dKO MEFs, size bar 20µm. Western blot documenting (middle panel) the knockdown

efficiency of two independent shRNA constructs targeting DLK-1. Quantification of oil

red O staining, n=3. Data shown are mean ± STD, *** P<0.001.

Figure EV3. DLK-1 regulates Hes1 expression, adipocyte differentiation of dKO MEF

cells, and C/EBPα expression. dKO MEFs-secreted DLK-1 inhibits adipocyte

differentiation of WT MEF cells.

A) Hes1 mRNA expression detected in 8 days differentiated MEF cells. Normalized on

GAPDH. Error bars indicate standard deviations, WT MEFs values were set as 1. Data

shown are mean ± STD, ***P<0.001.

B) Oil red O staining (left panel) of 8 days adipocyte-differentiated WT control MEFs, WT

MEFs treated with a recombinant DLK-1 or expressing ectopic DLK-1.

C) Quantification of oil red O staining shown in panel E, n=3. Data shown are mean ± STD,

***P<0.001.

D) DLK-1 mRNA expression detected in 8 days differentiated MEF cells.

E) C/EBPα mRNA expression detected in 8 days differentiated MEF cells. Normalized on

GAPDH. Error bars indicate standard deviations, WT MEFs values were set as 1. Data

shown are mean ± STD, ***P<0.001.

F) Oil red O staining of WT control MEFs (left image) following 8 days adipocyte

differentiation protocol. WT cells differentiated in conditioned medium from dKO MEFs

(middle image) and WT MEFs treated with conditioned medium from dKO MEFs

expressing DLK-1 shRNA construct (right image).   Size bar 20μm. Western blot

documenting (middle panel) the knockdown efficiency of shRNA constructs targeting

DLK-1. Quantification of oil red O staining, n=3. Data shown are mean ± STD,

***P<0.001.

Supplementary figure 1. No difference between WT and dKO mice in food consumption.

Expression patterns of TIS7 and SKMc15 in WT MEFs during adipocyte differentiation.

SKMc15 knockout strategy.

A) No significant genotype effect on food intake; when adjusted to the body mass, non-

significantly increased in dKO mice (4.0 g WT and 3.4 g dKO mice). Data shown are

mean ± STD.

B) Hepatic lipase serum concentrations of WT (n=4) and dKO (n=4) 11 weeks old male

mice after 3 weeks of HFD. Data shown are mean ± STD.

C) Lipoprotein lipase serum concentrations of WT (n=4) and dKO (n=4) 8 weeks old male

mice following 3 weeks of HFD. Data shown are mean ± STD.

D) TIS7 expression reaches its maximum on day 3 of MEF adipocyte differentiation. Data

shown are mean ± STD. Student's t-test P<0.01, n=3.

E) SKMc15 reaches its maximum expression on day 5 of MEF adipocyte differentiation.

Data shown are mean ± STD. Student's t-test P<0.05, n=3. All Ct values were

normalized to GAPDH.

F) Modification of the SKMc15 gene locus and characterization of the targeted and null

genotype. Homologous recombination with the targeting vector inserts one loxP site and

a frt site flanked neomycin resistance gene (PGK–Neo) into intron 1 at position 105515

(AY162905). A second loxP site inserted behind the poly A tail into the 3´ UTR of the

SKMc15 gene at position 102082 (AY162905). Further downstream were added 15

additional nucleotides, parts of intron 1, exon 2 (splice site: donor and acceptor) and the

CDS of hrGFP (out of Vitality hrGFP Mammalian Expression Vector pIRES-hrGFP-2a;

Stratagene). Arrows indicated the orientation of the hrGFP gene and the loxP sites.

Southern blot analysis. The fragment detected by the Southern probe, was 9.6 kb in the

WT (insert, band V) and 15 kb in the deleted locus (band IV).

G) PCR genotype analysis of SKMc15 targeted and deleted alleles. I: 426 bp WT SKMc15

PCR fragment. II: 1772 bp SKMc15 targeted allele PCR product. III: 1700 bp SKMc15

GFP screening PCR product can be amplified from the targeted (F) and the deleted allele

(-/-), but not from the WT (+/+) locus.

H) TIS7 PCR detection in dKO mice. Left panel: detection of 1800 bp TIS7 WT allele; right

panel: detection of 2100 bp TIS7 KO allele.

Conflict of interest.

The authors declare that they have no conflict of interest.

References

Barclay JL, Nelson CN, Ishikawa M, Murray LA, Kerr LM, McPhee TR, Powell EE, Waters MJ
(2011) GH-dependent STAT5 signaling plays an important role in hepatic lipid metabolism.
Endocrinology 152: 181-92

Bennett CN, Hodge CL, MacDougald OA, Schwartz J (2003) Role of Wnt10b and C/EBPalpha
in spontaneous adipogenesis of 243 cells. *Biochem Biophys Res Commun* 302: 12-6

Brown A, Baird MR, Yip MC, Murray J, Shao S (2018) Structures of translationally inactive
mammalian ribosomes. *Elife* 7

Cases S, Smith SJ, Zheng YW, Myers HM, Lear SR, Sande E, Novak S, Collins C, Welch CB,
Lusis AJ, Erickson SK, Farese RV, Jr. (1998) Identification of a gene encoding an acyl
CoA:diacylglycerol acyltransferase, a key enzyme in triacylglycerol synthesis. *Proc Natl Acad*
Sci U S A 95: 13018-23

Christiaens V, Van Hul M, Lijnen HR, Scroyen I (2012) CD36 promotes adipocyte
differentiation and adipogenesis. *Biochim Biophys Acta* 1820: 949-56

Coburn CT, Knapp FF, Jr., Febbraio M, Beets AL, Silverstein RL, Abumrad NA (2000)
Defective uptake and utilization of long chain fatty acids in muscle and adipose tissues of CD36
knockout mice. *J Biol Chem* 275: 32523-9

Farmer SR (2006) Transcriptional control of adipocyte formation. *Cell Metab* 4: 263-73

Garces C, Ruiz-Hidalgo MJ, Bonvini E, Goldstein J, Laborda J (1999) Adipocyte differentiation
is modulated by secreted delta-like (dlk) variants and requires the expression of membrane-
associated dlk. *Differentiation* 64: 103-14

Garcia AM, Wakeman D, Lu J, Rowley C, Geisman T, Butler C, Bala S, Swietlicki EA, Warner
BW, Levin MS, Rubin DC (2014) Tis7 deletion reduces survival and induces intestinal
anastomotic inflammation and obstruction in high-fat diet-fed mice with short bowel syndrome.
Am J Physiol Gastrointest Liver Physiol 307: G642-54

Gautam J, Khedgikar V, Kushwaha P, Choudhary D, Nagar GK, Dev K, Dixit P, Singh D,
Maurya R, Trivedi R (2017) Formononetin, an isoflavone, activates AMP-activated protein
kinase/beta-catenin signalling to inhibit adipogenesis and rescues C57BL/6 mice from high-fat
diet-induced obesity and bone loss. *Br J Nutr* 117: 645-661

Glazier AM, Scott J, Aitman TJ (2002) Molecular basis of the Cd36 chromosomal deletion
underlying SHR defects in insulin action and fatty acid metabolism. *Mamm Genome* 13: 108-13

Harchaoui KE, Franssen R, Hovingh GK, Bisoendial RJ, Stellaard F, Kuipers F, Kastelein JJ,
Kuivenhoven JA, Stroes ES, Groen AK (2009) Reduced fecal sterol excretion in subjects with
familial hypoalphalipoproteinemia. *Atherosclerosis* 207: 614-6

Hu L, Yang G, Hagg D, Sun G, Ahn JM, Jiang N, Ricupero CL, Wu J, Rodhe CH, Ascherman
JA, Chen L, Mao JJ (2015) IGF1 Promotes Adipogenesis by a Lineage Bias of Endogenous
Adipose Stem/Progenitor Cells. *Stem Cells* 33: 2483-95

Huang Y, Yang X, Wu Y, Jing W, Cai X, Tang W, Liu L, Liu Y, Grottkau BE, Lin Y (2010)
gamma-secretase inhibitor induces adipogenesis of adipose-derived stem cells by regulation of
Notch and PPAR-gamma. *Cell Prolif* 43: 147-56

Iacopetti P, Barsacchi G, Tirone F, Cremisi F (1996) Expression of the PC4 gene in the
developing rat nervous system. *Brain Res* 707: 293-7.

Iezaki T, Onishi Y, Ozaki K, Fukasawa K, Takahata Y, Nakamura Y, Fujikawa K, Takarada T,
Yoneda Y, Yamashita Y, Shioi G, Hinoi E (2016) The Transcriptional Modulator Interferon-
Related Developmental Regulator 1 in Osteoblasts Suppresses Bone Formation and Promotes
Bone Resorption. *J Bone Miner Res* 31: 573-84

Kashiwagi H, Tomiyama Y, Nozaki S, Kiyoi T, Tadokoro S, Matsumoto K, Honda S, Kosugi S,
Kurata Y, Matsuzawa Y (2001) Analyses of genetic abnormalities in type I CD36 deficiency in
Japan: identification and cell biological characterization of two novel mutations that cause CD36
deficiency in man. *Hum Genet* 108: 459-66

Kim KA, Kim JH, Wang Y, Sul HS (2007) Pref-1 (preadipocyte factor 1) activates the
MEK/extracellular signal-regulated kinase pathway to inhibit adipocyte differentiation. *Mol Cell
Biol* 27: 2294-308

Koliwad SK, Streeper RS, Monetti M, Cornelissen I, Chan L, Terayama K, Naylor S, Rao M,
Hubbard B, Farese RV, Jr. (2010) DGAT1-dependent triacylglycerol storage by macrophages
protects mice from diet-induced insulin resistance and inflammation. *J Clin Invest* 120: 756-67

Lammirato A, Patsch K, Feiereisen F, Maly K, Nofziger C, Paulmichl M, Hackl H, Trajanoski Z,
Valovka T, Huber LA, Vietor I (2016) TIS7 induces transcriptional cascade of methylosome
components required for muscle differentiation. *BMC Biol* 14: 95

Latif F, Duh FM, Bader S, Sekido Y, Li H, Geil L, Zbar B, Minna JD, Lerman MI (1997) The
human homolog of the rodent immediate early response genes, PC4 and TIS7, resides in the lung
cancer tumor suppressor gene region on chromosome 3p21. *Hum Genet* 99: 334-41.

LeBlanc SE, Konda S, Wu Q, Hu YJ, Oslowski CM, Sif S, Imbalzano AN (2012) Protein
arginine methyltransferase 5 (Prmt5) promotes gene expression of peroxisome proliferator-
activated receptor gamma2 (PPARgamma2) and its target genes during adipogenesis. *Mol
Endocrinol* 26: 583-97

Lee K, Villena JA, Moon YS, Kim KH, Lee S, Kang C, Sul HS (2003) Inhibition of
adipogenesis and development of glucose intolerance by soluble preadipocyte factor-1 (Pref-1). *J*
Clin Invest 111: 453-61

Lee YL, Helman L, Hoffman T, Laborda J (1995) dlk, pG2 and Pref-1 mRNAs encode similar
proteins belonging to the EGF-like superfamily. Identification of polymorphic variants of this
RNA. *Biochim Biophys Acta* 1261: 223-32

Lefterova MI, Zhang Y, Steger DJ, Schupp M, Schug J, Cristancho A, Feng D, Zhuo D,
Stoeckert CJ, Jr., Liu XS, Lazar MA (2008) PPARgamma and C/EBP factors orchestrate
adipocyte biology via adjacent binding on a genome-wide scale. *Genes Dev* 22: 2941-52

Leitner PD, Jakschitz T, Gstir R, Stuppner S, Perkams S, Kruus M, Trockenbacher A, Griesbeck
C, Bonn GK, Huber LA, Valovka T (2022) Anti-Inflammatory Extract from Soil Algae
Chromochloris zofingiensis Targeting TNFR/NF-kappaB Signaling at Different Levels. *Cells* 11

Lemonnier D (1972) Effect of age, sex, and sites on the cellularity of the adipose tissue in mice
and rats rendered obese by a high-fat diet. *J Clin Invest* 51: 2907-15

Li HX, Luo X, Liu RX, Yang YJ, Yang GS (2008) Roles of Wnt/beta-catenin signaling in
adipogenic differentiation potential of adipose-derived mesenchymal stem cells. *Mol Cell*
Endocrinol 291: 116-24

Lowell BB (1999) PPARgamma: an essential regulator of adipogenesis and modulator of fat cell
function. *Cell* 99: 239-42

Ludwig EH, Mahley RW, Palaoglu E, Ozbayrakci S, Balestra ME, Borecki IB, Innerarity TL,
Farese RV, Jr. (2002) DGAT1 promoter polymorphism associated with alterations in body mass
index, high density lipoprotein levels and blood pressure in Turkish women. *Clin Genet* 62: 68-
73

Marcon BH, Holetz FB, Eastman G, Origa-Alves AC, Amoros MA, de Aguiar AM, Rebelatto
CK, Brofman PRS, Sotelo-Silveira J, Dallagiovanna B (2017) Downregulation of the protein
synthesis machinery is a major regulatory event during early adipogenic differentiation of human
adipose-derived stromal cells. *Stem Cell Res* 25: 191-201

Micheli L, Leonardi L, Conti F, Buanne P, Canu N, Caruso M, Tirone F (2005) PC4 Coactivates
MyoD by Relieving the Histone Deacetylase 4-Mediated Inhibition of Myocyte Enhancer Factor
2C. *Mol Cell Biol* 25: 2242-2259

Moon YS, Smas CM, Lee K, Villena JA, Kim KH, Yun EJ, Sul HS (2002) Mice lacking
paternally expressed Pref-1/Dlk1 display growth retardation and accelerated adiposity. *Mol Cell*
Biol 22: 5585-92

Mortensen SB, Jensen CH, Schneider M, Thomassen M, Kruse TA, Laborda J, Sheikh SP,
Andersen DC (2012) Membrane-tethered delta-like 1 homolog (DLK1) restricts adipose tissue
size by inhibiting preadipocyte proliferation. *Diabetes* 61: 2814-22

Nagasaka H, Yorifuji T, Takatani T, Okano Y, Tsukahara H, Yanai H, Hirano K, Hui SP,
Hirayama S, Ito T, Chiba H, Miida T (2011) CD36 deficiency predisposing young children to
fasting hypoglycemia. *Metabolism* 60: 881-7

Nakamura Y, Hinoi E, Iezaki T, Takada S, Hashizume S, Takahata Y, Tsuruta E, Takahashi S,
Yoneda Y (2013) Repression of adipogenesis through promotion of Wnt/beta-catenin signaling
by TIS7 up-regulated in adipocytes under hypoxia. *Biochim Biophys Acta* 1832: 1117-28

Neeland IJ, Ayers CR, Rohatgi AK, Turer AT, Berry JD, Das SR, Vega GL, Khera A, McGuire
DK, Grundy SM, de Lemos JA (2013) Associations of visceral and abdominal subcutaneous
adipose tissue with markers of cardiac and metabolic risk in obese adults. *Obesity (Silver Spring)*
21: E439-47

Nozaki S, Tanaka T, Yamashita S, Sohmiya K, Yoshizumi T, Okamoto F, Kitaura Y, Kotake C,
Nishida H, Nakata A, Nakagawa T, Matsumoto K, Kameda-Takemura K, Tadokoro S, Kurata Y,
Tomiyama Y, Kawamura K, Matsuzawa Y (1999) CD36 mediates long-chain fatty acid transport
in human myocardium: complete myocardial accumulation defect of radiolabeled long-chain
fatty acid analog in subjects with CD36 deficiency. *Mol Cell Biochem* 192: 129-35

Park G, Horie T, Kanayama T, Fukasawa K, Iezaki T, Onishi Y, Ozaki K, Nakamura Y, Yoneda
Y, Takarada T, Hinoi E (2017) The transcriptional modulator Ifrd1 controls PGC-1alpha
expression under short-term adrenergic stimulation in brown adipocytes. *The FEBS journal* 284:
784-795

Paul C, Sardet C, Fabbrizio E (2012) The histone- and PRMT5-associated protein COPR5 is
required for myogenic differentiation. *Cell Death Differ* 19: 900-8

Paul C, Sardet C, Fabbrizio E (2015) The Wnt-target gene Dlk-1 is regulated by the Prmt5-
associated factor Copr5 during adipogenic conversion. *Biol Open* 4: 312-6

Pfluger PT, Kabra DG, Aichler M, Schriever SC, Pfuhlmann K, Garcia VC, Lehti M, Weber J,
Kutschke M, Rozman J, Elrod JW, Hevener AL, Feuchtinger A, Hrabe de Angelis M, Walch A,
Rollmann SM, Aronow BJ, Muller TD, Perez-Tilve D, Jastroch M et al. (2015) Calcineurin
Links Mitochondrial Elongation with Energy Metabolism. *Cell Metab* 22: 838-50

Popow J, Alleaume AM, Curk T, Schwarzl T, Sauer S, Hentze MW (2015) FASTKD2 is an
RNA-binding protein required for mitochondrial RNA processing and translation. *RNA* 21:
1873-84

Prestwich TC, Macdougald OA (2007) Wnt/beta-catenin signaling in adipogenesis and
metabolism. *Curr Opin Cell Biol* 19: 612-7

Reintjes A, Fuchs JE, Kremser L, Lindner HH, Liedl KR, Huber LA, Valovka T (2016)
Asymmetric arginine dimethylation of RelA provides a repressive mark to modulate
TNFalpha/NF-kappaB response. *Proc Natl Acad Sci U S A* 113: 4326-31

Ross SE, Hemati N, Longo KA, Bennett CN, Lucas PC, Erickson RL, MacDougald OA (2000)
Inhibition of adipogenesis by Wnt signaling. *Science* 289: 950-3

Ruiz-Ojeda FJ, Ruperez AI, Gomez-Llorente C, Gil A, Aguilera CM (2016) Cell Models and
Their Application for Studying Adipogenic Differentiation in Relation to Obesity: A Review. *Int*
J Mol Sci 17

Sarantopoulos CN, Banyard DA, Ziegler ME, Sun B, Shaterian A, Widgerow AD (2018)
Elucidating the Preadipocyte and Its Role in Adipocyte Formation: a Comprehensive Review.
Stem Cell Rev 14: 27-42

Savant-Bhonsale S, Cleveland DW (1992) Evidence for instability of mRNAs containing
AUUUA motifs mediated through translation-dependent assembly of a > 20S degradation
complex. *Genes Dev* 6: 1927-39

Shore P, Dietrich W, Corcoran LM (2002) Oct-2 regulates CD36 gene expression via a
consensus octamer, which excludes the co-activator OBF-1. *Nucleic Acids Res* 30: 1767-73

Sul HS (2009) Minireview: Pref-1: role in adipogenesis and mesenchymal cell fate. *Mol*
Endocrinol 23: 1717-25

Tanaka T, Nakata T, Oka T, Ogawa T, Okamoto F, Kusaka Y, Sohmiya K, Shimamoto K,
Itakura K (2001) Defect in human myocardial long-chain fatty acid uptake is caused by
FAT/CD36 mutations. *J Lipid Res* 42: 751-9

Tontonoz P, Spiegelman BM (2008) Fat and beyond: the diverse biology of PPARgamma. *Annu*
Rev Biochem 77: 289-312

Tsai J, Qiu W, Kohen-Avramoglu R, Adeli K (2007) MEK-ERK inhibition corrects the defect in
VLDL assembly in HepG2 cells: potential role of ERK in VLDL-ApoB100 particle assembly.
Arterioscler Thromb Vasc Biol 27: 211-8

Vadivelu SK, Kurzbauer R, Dieplinger B, Zweyer M, Schafer R, Wernig A, Vietor I, Huber LA
(2004) Muscle regeneration and myogenic differentiation defects in mice lacking TIS7. *Mol Cell*
Biol 24: 3514-25

van Tienen FH, Laeremans H, van der Kallen CJ, Smeets HJ (2009) Wnt5b stimulates
adipogenesis by activating PPARgamma, and inhibiting the beta-catenin dependent Wnt
signaling pathway together with Wnt5a. *Biochem Biophys Res Commun* 387: 207-11

Vietor I, Kurzbauer R, Brosch G, Huber LA (2005) TIS7 regulation of the beta-catenin/Tcf-4
target gene osteopontin (OPN) is histone deacetylase-dependent. *J Biol Chem* 280: 39795-801

Vietor I, Vadivelu SK, Wick N, Hoffman R, Cotten M, Seiser C, Fialka I, Wunderlich W, Haase
A, Korinkova G, Brosch G, Huber LA (2002) TIS7 interacts with the mammalian SIN3 histone
deacetylase complex in epithelial cells. *Embo J* 21: 4621-31

Wang S, Sun Z, Zhang X, Li Z, Wu M, Zhao W, Wang H, Chen T, Yan H, Zhu J (2015) Wnt1
positively regulates CD36 expression via TCF4 and PPAR-gamma in macrophages. *Cell Physiol*
Biochem 35: 1289-302

Wang Y, Iordanov H, Swietlicki EA, Wang L, Fritsch C, Coleman T, Semenkovich CF, Levin
MS, Rubin DC (2005) Targeted intestinal overexpression of the immediate early gene tis7 in
transgenic mice increases triglyceride absorption and adiposity. *J Biol Chem* 280: 34764-75

Wang Y, Sul HS (2009) Pref-1 regulates mesenchymal cell commitment and differentiation
through Sox9. *Cell Metab* 9: 287-302

Wick N, Schleiffer A, Huber LA, Vietor I (2004) Inhibitory effect of TIS7 on Sp1-C/EBPalpha
transcription factor module activity. *J Mol Biol* 336: 589-95

Wilson CG, Tran JL, Erion DM, Vera NB, Febbraio M, Weiss EJ (2016) Hepatocyte-Specific
Disruption of CD36 Attenuates Fatty Liver and Improves Insulin Sensitivity in HFD-Fed Mice.
Endocrinology 157: 570-85

Xu C, Wang J, Zhu T, Shen Y, Tang X, Fang L, Xu Y (2016) Cross-Talking Between PPAR and
WNT Signaling and its Regulation in Mesenchymal Stem Cell Differentiation. *Curr Stem Cell*
Res Ther 11: 247-54

Yu C, Jiang S, Lu J, Coughlin CC, Wang Y, Swietlicki EA, Wang L, Vietor I, Huber LA, Cikes
D, Coleman T, Xie Y, Semenkovich CF, Davidson NO, Levin MS, Rubin DC (2010) Deletion of
Tis7 protects mice from high-fat diet-induced weight gain and blunts the intestinal adaptive
response postresection. *J Nutr* 140: 1907-14

Zufferey R, Dull T, Mandel RJ, Bukovsky A, Quiroz D, Naldini L, Trono D (1998) Self-
inactivating lentivirus vector for safe and efficient in vivo gene delivery. *J Virol* 72: 9873-80

1167

A 8 days differentiated SVF cells from inguinal WAT

WT

SKMc15 KO

TIS7 KO

dKO

oil red O staining

B SVF

[% of WT]

WT SKMc15 KO TIS7 KO dKO

oil red O staining

C MEFs

[fold-increase]

N.S.

WT SKMc15 KO TIS7 KO dKO

TOPflash activity

D inguinal WAT

β-catenin — 95 kDa
actin — 50 kDa

DLK-1 — 50 kDa
actin — 50 kDa

E inguinal WAT

DLK-1 mRNA

WT dKO

DLK-1 — 50 kDa
GAPDH — 35 kDa

F inguinal WAT undifferentiated cells

DLK-1 mRNA

WT dKO

G

DLK-1 [fold change]

WT dKO

DLK-1 — 50 kDa
GAPDH — 35 kDa

H inguinal WAT 8 days differentiated cells

DLK-1 mRNA

WT dKO

A

^{35}S-Methionine coomassie blue

B

DLK-1 - 50 kDa

Collagen1 - 40 kDa
- 35 kDa

amido black - 60 kDa
- 40 kDa

C

8 days differentiated MEFs

Exp. 1 Exp. 2 Exp. 3

WT +WT medium

WT +dKO medium

dKO

oil red O staining

D

E

F

G

G WAT

mouse: 1 2 1 2

40 - Pp42/44

40 - p42/44

H

G WAT

I

BAT

WT dKO SKMc15 TIS7
KO KO

mouse: 1 2 1 2 1 2 1 2

DLK-1 - 50 kDa

GAPDH - 35 kDa

A

B

8 days differentiated MEFs

C

D

adipocyte differentiation

E

F

G

A

B

C

inguinal WAT

D

adipocyte differentiation

E

adipocyte differentiation

F