The negative adipogenesis regulator DLK1 is transcriptionally regulated by TIS7 (IFRD1) and translationally by its orthologue SKMc15 (IFRD2)

[1]Ilja Vietor*, [1,12]Domagoj Cikes, [1,13]Kati Piironen, [1,14]Ronald Gstir, [2]Matthias David Erlacher, [3]Michael Hess, [4]Volker Kuhn (†), [5]Gerald Degenhart, [6,7]Jan Rozman, [8,9,10]Martin Klingenspor, [6,7,11]Martin Hrabe de Angelis, [1]Taras Valovka, [1, 14]Lukas A. Huber

[1]Institute of Cell Biology, Biocenter, [2]Division of Genomics and RNomics, Biocenter, [3]Division of Histology and Embryology, [4]Department Trauma Surgery, [5]Department Radiology, Medical University Innsbruck, Austria, [6]German Mouse Clinic, Institute of Experimental Genetics, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany, [7]German Center for Diabetes Research (DZD), Neuherberg, Germany, [8]Chair of Molecular Nutritional Medicine, Technical University of Munich, School of Life Sciences, Weihenstephan, Freising, Germany, [9]EKFZ - Else Kröner Fresenius Center for Nutritional Medicine, Technical University of Munich, Freising, Germany, [10]ZIEL - Institute for Food & Health, Technical University of Munich, Freising, Germany, [11]Chair of Experimental Genetics, Technical University of Munich, School of Life Sciences, Weihenstephan, Freising, Germany, [12]IMBA, Institute of Molecular Biotechnology of the Austrian Academy of Sciences, Vienna, Austria, [13]Division of Pharmaceutical Chemistry and Technology, Faculty of Pharmacy, University of Helsinki, Finland, [14]ADSI – Austrian Drug Screening Institute GmbH, Innsbruck, Austria

* Address correspondence to: Ilja Vietor, Institute of Cell Biology, Biocenter, Innsbruck Medical University, Innrain 80-82, A-6020 Innsbruck, Austria. Phone: +43 512 9003 70175 E-mail: ilja.vietor@i-med.ac.at

† Deceased May 2019

Running title: Identification of novel regulators of adipogenesis

Conflict of interest: The authors declare that they have no conflict of interest.

Abstract

Delta-like homolog 1 (DLK1), an inhibitor of adipogenesis, controls the cell fate of adipocyte progenitors.

Understanding regulatory mechanisms affecting DLK1 levels is therefore of high interest. Here we identify

two independent mechanisms, transcriptional and translational, by which TIS7 (IFRD1) and its orthologue

SKMc15 (IFRD2) regulate DLK1 levels. Mice deficient in both TIS7 and SKMc15 (dKO) had severely reduced

adipose tissue and were resistant to high fat diet-induced obesity. Wnt signaling, a negative regulator of

adipocyte differentiation was significantly up regulated in dKO mice. Elevated levels of the Wnt/β-catenin

target protein Dlk-1 inhibited the expression of adipogenesis regulators PPARγ and C/EBPα, and fatty acid

transporter CD36. Although both, TIS7 and SKMc15, contributed to this phenotype, they utilized two different

mechanisms. TIS7 acted by controlling Wnt signaling and thereby transcriptional regulation of Dlk-1. On the

other hand, here we provide experimental evidence that SKMc15 acts as a general translational inhibitor

affecting DLK-1 protein levels. Our study provides data describing novel mechanisms of DLK1 regulation in

adipocyte differentiation involving TIS7 and SKMc15.

Nonstandard abbreviations used: CD36, cluster of differentiation 36; HFD, high fat diet; IFRD, interferon

related developmental regulator; KO, knockout; dKO, double knockout; RD, regular diet; SKMc15, skeletal

muscle cDNA library clone15; TIS7, TPA Induced Sequence 7

Introduction

Adipogenesis is a complex process in which multipotent stem cells are converted into preadipocytes before

terminal differentiation into adipocytes [1]. These mechanisms involve protein factor regulators, epigenetic

factors, and miRNAs. TPA Induced Sequence 7 (TIS7) protein has been shown to be involved in the mainly

transcriptional regulation of differentiation processes in various cell types, e.g. neurons [2], enterocytes [3],

myocytes [4] and also adipocytes [5].

Multiple lines of evidence link the regulation of Wnt/β-catenin signaling to the physiological function of TIS7,

known in human as Interferon related developmental regulator 1 (IFRD1) [6, 7]. Our data based on experiments

with TIS7 knockout mice show a negative effect of TIS7 on Wnt signaling and a positive on adipocyte

differentiation [6, 8]. TIS7 deficiency leads to a significant up regulation of Wnt/β-catenin transcriptional activity

in both primary osteoblasts and in mouse embryonic fibroblasts (MEFs) derived from TIS7 knockout (KO)

mice. Recent data show that TIS7 is also involved in the control of adipocytes differentiation in mice and was

up regulated in both visceral (vWAT) and subcutaneous (sWAT) white adipose tissues (WAT) of genetically

obese ob/ob mice [5]. TIS7 transgenic mice have increased total body adiposity and decreased lean mass

compared with wild type (WT) littermates [3]. On HFD, TIS7 transgenic mice exhibit a more rapid and

proportionately greater gain in body weight with persistently elevated total body adiposity. Enhanced

triglyceride (TG) absorption in the gut of TIS7 transgenic mice [3] indicated that TIS7 expressed in the gut

epithelium has direct effects on fat absorption in enterocytes. We have shown that as a result of impaired

intestinal lipid absorption TIS7 KO mice displayed lower body adiposity [8]. Compared with WT littermates,

TIS7 KO mice do not gain weight when chronically fed an HFD and TIS7 deletion results in delayed lipid

absorption and altered intestinal and hepatic lipid trafficking, with reduced intestinal TG, cholesterol, and free

fatty acid mucosal levels in the jejunum [9]. TIS7 protein functions as a transcriptional co-regulator [10] due to its

interaction with protein complexes containing either histone deacetylases (HDAC) [4, 11, 12, 13] or protein methyl

transferases, in particular PRMT5 [14]. The analysis of adipocyte differentiation in preadipocytic 3T3-L1 cells

suggested an involvement of TIS7 in the regulation of adipogenesis in the Wnt/β-catenin signaling context [5].

SKMc15, also known as Interferon related developmental regulator 2 (IFRD2), a second member of TIS7

gene family, is highly conserved in different species [15]. Mouse TIS7 and SKMc15 are highly homologous,

with a remarkable identity at both the cDNA and amino acid levels (58% and 88%, respectively). However,

there was so far no information about the physiological function and mechanisms of action of SKMc15 and

its possible involvement in differentiation of various tissues. Recently, cryo-electronmicroscopy (cryo-EM)

analyses of inactive ribosomes identified IFRD2 (SKMc15) as a novel ribosome-binding protein inhibiting

translation to regulate gene expression [16]. The physiological function of SKMc15 matching the mechanism

based on the above mentioned cryo-EM data was so far not shown and we postulate it could be involved in

adipogenesis, since a significant reduction of whole protein synthesis was previously shown as a major

regulatory event during early adipogenic differentiation [17].

Adipogenesis occurs late in embryonic development and in postnatal periods. Adipogenic transcription

factors CCAAT/enhancer binding protein α (C/EBPα) and peroxisome proliferator activated receptor γ

(PPARγ) play critical roles in adipogenesis and in the induction of adipocyte markers [18]. PPARγ is the major

downstream target of Delta-like protein 1 (DLK-1). It is inactivated by the induction of the MEK/ERK pathway

leading to its phosphorylation and proteolytic degradation [19]. DLK-1, also known as Pref-1 (preadipocyte

factor 1) activates the MEK/ERK pathway to inhibit adipocyte differentiation [20]. C/EBPα is highly expressed

in mature adipocytes and can bind DNA together with PPARγ to a variety of respective target genes [21].

Besides that, PPARγ binding to C/EBPα gene induces its transcription, thereby creating a positive feedback

loop [22]. Both proteins have synergistic effects on the differentiation of adipocytes that requires a balanced

expression of both C/EBPα and PPARγ.

Wnt/β-catenin signaling is one of the extracellular signaling pathways specifically affecting adipogenesis [23,

24, 25] by maintaining preadipocytes in an undifferentiated state through inhibition of C/EBPα and PPARγ [26].

PPAR-γ and Wnt/β-catenin pathways are regarded as master mediators of adipogenesis [27]. Wnt signaling is

a progenitor fate determinator and negatively regulates preadipocyte proliferation through DLK-1 [28]. Mice

over-expressing DLK-1 are resistant to HFD-induced obesity, whereas DLK-1 KO mice have accelerated

adiposity [29]. DLK-1 transgenic mice show reduced expression of genes controlling lipid import (CD36) and

synthesis (Srebp1c, Pparγ) [30]. DLK-1 expression coincides with altered recruitment of PRMT5 and β-catenin

to the DLK-1 promoter [31]. PRMT5 acts as a co-activator of adipogenic gene expression and differentiation [32].

SRY (sex determining region Y)-box 9 (Sox9), a transcription factor expressed in preadipocytes, is down

regulated preceding adipocyte differentiation. DLK-1 prevents down regulation of Sox9 by activating ERK,

resulting in inhibition of adipogenesis [33]. The PRMT5- and histone-associated protein COPR5, affects PRMT5

functions related to cell differentiation [34]. Adipogenic conversion is delayed in MEFs derived from COPR5

KO mice and WAT of COPR5 KO mice is reduced when compared to control mice. DLK-1 expression is up

regulated in COPR5 KO cells [31].

Here we show the involvement of TIS7 and SKMc15 in the process of adipocyte differentiation. dKO mice

had strongly decreased amounts of the body fat when fed with even regular, chow diet and were resistant

against the high fat diet induced obesity. We found two independent molecular mechanisms through which

TIS7 and SKMc15 fulfill this function in a synergistic manner. TIS7 regulates the Wnt signaling pathway

activity and restricts DLK-1 protein levels, thereby allowing adipocyte differentiation. In contrast, SKMc15 KO

did not affect Wnt signaling, but as we show here, cells lacking SKMc15 have significantly up regulated

translational activity. This was true also for dKO, but not for the single TIS7 knockout cells. The ablation of

both TIS7 and SKMc15 genes significantly affected the expression of genes essential for adipocyte

differentiation and function. Since dKO mice render a substantially leaner phenotype on chow diet, even

without any challenge by HFD-induction, we propose that TIS7 and SKMc15 represent novel players in the

process of physiological adipocyte differentiation.

118
Results

Mice lacking TIS7 and SKMc15 genes have lower body mass, less fat and are resistant against HFD-induced

obesity.

In order to clarify whether both, TIS7 and SKMc15 are involved in the regulation of the adipocyte

differentiation and if they act through same or different mechanisms, we have generated mice lacking both

genes by crossing TIS7 with SKMc15 single KO mice. dKO pups were viable, and adult male and female

mice were fertile. At birth, body weights of dKO and WT mice were similar. Nevertheless, already during

weaning, both the male and female dKO mice failed to gain weight when compared to their WT littermates

and this persisted in the following weeks when the mice were fed regular diet (RD; chow diet) containing 11

% kcal of fat. At 10 weeks of age, dKO mice displayed 30 % and later up to 44.9 % lower body weight

compared with WT mice (Figure 1a).

Based on Dual Energy X-ray Absorptiometry (DEXA) measurement, 6-month-old WT mice had substantially

higher amounts of fat than their dKO littermates (Figure 1b, left panel). The effect of TIS7 and SKmc15 double

knockout was even more pronounced when the total fat and lean mass values were normalized to the body

weight since the dKO mice were smaller than their WT counterparts. The percentage of fat was in the WT

mice 37.7 ± 4 % vs. 6 ± 3 % of the total body mass in dKO animals. Furthermore, the percentage of lean

tissue mass in WT animals was lower than in dKO animals (60 ± 4.6 % vs. 92 ± 3.14 %; Figure 1b, right

panel). Nevertheless, the dKO mice were not overall smaller, since their body length, including the tail did

not significantly differ between WT and dKO mice at different age (n=20; data not shown). Next, we analyzed

the contribution of TIS7 and SKMc15 to the whole-body fat content of mice. Three-dimensional reconstruction

of images based on micro-computed tomography (micro-CT) of sex/age-matched adult mice disclosed that

both TIS7 and SKMc15 single KO mice already had less abdominal fat than WT controls and that the dKO

of TIS7 and SKMc15 genes caused the strongest decrease in abdominal fat content and size (Figure 1c).

Quantitative analyses of micro-CT measurements showed that TIS7 deficiency caused a substantial lack of

the abdominal fat tissues (P=0.002 when compared to WT mice, Figure 2a). Whereas TIS7 KO mice had

less fat mass but were not significantly lighter than their WT littermates (Figure 2b), SKMc15 KO were lighter

(P=0.007) and leaner (P=0.0001) and dKO mice were both significantly lighter (P = 0.0001) and had

significantly less abdominal fat (P=0.006) than the WT mice (Figures 2a, b).

We conducted an indirect calorimetry trial with sex- and age-matched WT and dKO animals (Supplementary

figure 1). We did not identify any significant difference in respiratory exchange ratios (RER = VCO2/VO2) of

WT and dKO mice (Supplementary figure 1a). However, dKO mice showed substantially reduced body

weight, mainly because of lacking fat, despite identical food intake, activity and no major differences in several

metabolic parameters. To investigate potential links between TIS7, SKMc15 and obesity, we studied the

response of dKO mice to HFD. At 2 months of age, male mice were housed individually and fed with HFD for

21 days. Food intake was measured every second day and body weight was measured every fourth day.

Feces were collected every second day to analyze the composition of excreted lipids and blood samples

were collected after the third week of HFD feeding to measure the concentrations of hepatic and lipoprotein

lipases, respectively. Already within the first week of HFD feeding, WT mice gained more weight than the

dKO mice (Figure 2c), although there were no obvious differences in food consumption (Supplementary

figure 1b) or in levels of lipolytic enzymes (Supplementary figures 1c, d). These differences in body weight

gain continued to increase during the second and third week, at which time the body weight of WT mice

increased additionally for 30% (30.0 ± 2.3%) when compared with the beginning of the HFD feeding period.

In contrast, the weight of dKO animals increased only slightly (7.6 ± 1.6%) (Figure 2c). Both genes, TIS7 and

SKMc15 contributed to this phenotype and we could see an additive effect following their deletion (Figure

2c).

Adipocyte differentiation in dKO mice is inhibited due to up regulated DLK-1 levels.

A possible explanation of the lean phenotype of dKO mice was that TIS7 and SKMc15 regulate adipocyte

differentiation. Primary MEFs derived from totipotent cells of early mouse mammalian embryos are capable

of differentiating into adipocytes and they are versatile models to study adipogenesis as well as mechanisms

related to obesity such as genes, transcription factors and signaling pathways implicated in the adipogenesis

process [35]. To test whether TIS7 and SKMc15 are required for adipogenesis we treated MEFs derived from

WT, SKMc15 and TIS7 single and dKO mice using an established adipocyte differentiation protocol [36].

Expression levels of both TIS7 and SKMc15 mRNA increased during the adipocyte differentiation of WT

MEFs. TIS7 expression reached maximum levels representing 5.7-fold increase compared to proliferating

WT MEFs on day 3 and SKMc15 reached on day 5 the maximum of 2.5-fold expression levels of proliferating

MEFs (Supplementary figure 2). These data suggested that both proteins play a regulatory role in

adipogenesis, however differ in their mechanisms and timing. Eight days after initiation of adipocyte

differentiation a remarkable reduction of adipocyte differentiation potential in SKMc15, TIS7 KO MEFs and

dKO MEFs was observed, as characterized by the formation of lipid droplets stained by oil red O (Figure 3a).

Quantification of this staining revealed that fat vacuole formation in cells derived from SKMc15, TIS7 and

dKO mice represented 65 %, 35 % and 24 % of the WT cells, respectively (Figure 3b). These data indicated

that both, TIS7 and SKMc15 were critical for adipocyte differentiation and that the defect in adipogenesis

could be responsible for the resistance of dKO mice to HFD-induced obesity.

Wnt/β-catenin signaling is an important regulatory pathway for adipocyte differentiation [37]. Consistently,

others and we have shown that TIS7 deficiency causes up regulation of Wnt/β-catenin transcriptional activity.

Therefore, we have assessed Wnt signaling activity in MEFs by measuring transcriptional activity using

TOPflash, TCF-binding luciferase reporter assays. As shown in Figure 3c, Wnt signaling activity, when

compared to the WT MEFs, was not regulated in SKMc15 KO, but significantly up regulated in TIS7 KO

($P<0.0001$) and also in dKO MEFs ($P<0.01$). Supporting these findings, Western blot analyses of WAT

samples also identified increased β-catenin protein levels in WAT of TIS7 KO and in dKO mice ($P<0.0001$),

but not in SKMc15 KO mice (Figure 3d, top panel) suggesting the involvement of TIS7 but not SKMc15 in

the Wnt signaling pathway regulation.

Next, we analyzed the expression levels of DLK-1, a negative regulator of adipogenesis and at the same

time known target of Wnt signaling [31]. DLK-1 protein was significantly upregulated in WAT isolated from TIS7

and SKMc15 KO as well as from dKO mice (Figure 3d, bottom panel). The qPCR analysis revealed a

significant up regulation of DLK-1 mRNA levels in dKO WAT samples ($P<0.01$) and similarly in dKO MEFs

($P<0.05$) when compared to the WT controls (Figure 4a). Moreover, whereas in the WT MEFs the expression

of DLK-1 was strongly up regulated only during the first day of adipocyte differentiation and then over the

next days declined to basal levels, in dKO MEFs DLK-1 was up regulated ($P<0.001$) throughout the entire 8

days of differentiation (Figure 4b). This result was confirmed on the protein level by Western blot analyses of

lysates from 8 day differentiated adipocytes (Figure 4c). A rescue experiment was conducted to confirm that

DLK-1 expression was TIS7- and SKMc15-dependent. dKO MEFs were transiently co-transfected with TIS7

and SKMc15 vectors and 48 hours later the mRNA expressions were analyzed by RT-qPCR. Ectopic co-

expression of TIS7 and SKMc15 significantly ($P < 0.05$) down regulated the DLK-1 expression (Figure 4d).

Accordingly, TIS7 and/or SKMc15 are involved in the regulation of DLK-1 expression, but the molecular

mechanism remained unclear.

Our earlier study revealed that TIS7 binds directly to DNA and via interaction with PRMT5 regulates gene

expression [14]. Therefore, we performed chromatin immunoprecipitation (ChIP) experiments where we

studied the binding of TIS7 and SKMc15 as well as known transcription factors (β-catenin and H4R3me2s

(symmetrically dimethylated histone H4 at arginine residue 3)) [31] to the regulatory elements of the DLK-1

gene. In dKO MEFs treated 8 days with the adipocyte differentiation cocktail we found increased β-catenin

binding to the β-catenin/TCF binding site 2 of the Dlk-1 regulatory element when compared to WT MEFs

(Figure 4e). On the other hand, binding of H4R3me2s to the same Dlk-1 regulatory element was significantly

reduced ($P < 0.0001$). We could not identify any direct binding of TIS7 or SKMc15 proteins to two different

DLK-1 regulatory elements (Dlk-1 region A and β-cat/TCFbs2), neither in WT nor in dKO MEFs, suggesting

rather an epigenetic regulation than *via* their direct binding to DLK-1 regulatory elements. We concluded that

TIS7 and SKMc15 are required to restrain the DLK-1 levels, through the Wnt/β-catenin signaling pathway

and yet another so far unknown mechanism.

In a search for a SKMc15-specific regulatory mechanism of Dlk-1 levels we focused on the translational

regulation since previously was shown that general reduction of protein synthesis and downregulation of the

expression and translational efficiency of ribosomal proteins are events crucial for the regulation of adipocyte

differentiation [17]. Interestingly, IFRD2 (SKMc15) was recently identified as a novel specific factor capable of

translationally inactivating ribosomes [16]. Because SKMc15 may play a crucial role in the adipogenesis

regulation we tested in our following experiment the effect of SKMc15 knockout on the general translational

efficiency of WT, TIS7, SKMc15 single and double KO MEFs. Cells were incubated 30 minutes in the absence

of methionine and cysteine, followed by one hour in the presence of [35]S-methionine. As shown in Figure 5a,

there was a significant increase in the general translational activity of MEFs lacking SKMc15 alone or both

TIS7 and SKMc15, but not in TIS7 single knockout cells. Therefore, we concluded that SKMc15 alone, but

not TIS7 inhibits the general translational activity necessary for the induction of adipogenic differentiation,

also via the translational regulation of Dlk-1. This finding supported the result shown in Figure 3d where the

DLK-1 protein levels were significantly induced in WAT samples isolated from SKMc15 single knockout mice.

TIS7 and SKMc15 regulate adipocyte differentiation through DLK-1, MEK/ERK pathway, PPARγ and

C/EBPα.

DLK-1 protein carries a protease cleavage site in its extracellular domain [38] and is secreted. The extracellular

domain of DLK-1 is cleaved by ADAM17, TNF-α Converting Enzyme to generate the biologically active

soluble DLK-1 [19]. DLK-1 mRNA and protein levels are high in preadipocytes but DLK-1 expression is absent

in mature adipocytes. Hence, adding soluble DLK-1 to the medium inhibits adipogenesis [39]. To test if the dKO

MEFs secreted DLK-1, cell culture media from MEFs treated 8 days with the adipocyte differentiation cocktail

were collected and analyzed by Western blotting. As shown in Figure 5b, dKO cells secreted DLK-1 protein,

but in an identical volume of the cell culture medium from WT MEFs no DLK-1 could be detected. To prove

that secreted DLK-1 could inhibit adipocyte differentiation of dKO MEFs, we cultured WT MEFs with

conditioned medium from dKO MEFs. Adipocyte differentiation of WT MEFs was strongly inhibited by the

dKO MEFs-conditioned medium when compared to the control WT cells (Figure 5c, quantified in Figure 5d).

Based on these results we concluded that cells derived from dKO mice express increased DLK-1 levels and

secreted, soluble DLK-1 may inhibit adipocyte differentiation *in vivo*.

Previous studies showed that soluble DLK-1 activates MEK/ERK signaling, which is required for inhibition of

adipogenesis [20]. As DLK-1 was strongly up regulated in dKO MEFs during adipocyte differentiation we

subsequently analyzed possible activation of the MEK/ERK pathway (Figure 6a). The phosphorylation of p44

and of p42 was up regulated 5-fold and 3-fold (P<0.05), respectively, in the adipocyte-differentiated dKO

when compared to the WT MEFs (Figure 6b). Previously, activation of MEK/ERK by DLK-1 was shown to up regulate the expression of the transcription factor SOX9, resulting in the inhibition of adipogenesis [33]. Therefore, we measured SOX9 mRNA expression by RT qPCR in 8 days adipocyte-differentiated MEFs. SOX9 expression was significantly (P<0.001) up regulated in 8 days adipocyte-differentiated dKO when compared to WT MEFs (Figure 6c).

The adipocyte differentiation deficiency of dKO MEFs suggested that C/EBPα and PPARγ might also be regulated through TIS7 and/or SKMc15. It was previously shown that elevated levels of the cleaved ectodomain of DLK1 have been correlated with reduced expression of PPARγ [40]. Therefore, we analyzed the differences in PPARγ expression between WT and dKO MEFs during their differentiation into adipocytes. While PPARγ expression was strongly induced during adipocyte differentiation from WT MEFs this was barely detectable in dKO MEFs (Figure 7a) and also C/EBPα expression was never induced up to the levels of WT (Figure 7b). In a rescue experiment we showed that the co-expression of TIS7 and SKMc15 in undifferentiated MEFs strongly increased the expression of the PPARγ in dKO MEFs (P<0.001), almost up to the levels of WT MEFs (Figure 7c). The expression of C/EBPα was also strongly up regulated by the co-expression of TIS7 and SKMc15 (P<0.01) (Figure 7d). Based on these results we concluded that TIS7 and SKMc15 regulate the expression of both PPARγ and C/EBPα, both crucial for adipocyte differentiation.

TIS7 and SKMc15 regulate the expression of the lipid transporters CD36 and DGAT1.

PPARγ induces the expression of CD36, a very long chain fatty acids (VLCFA) transporter in heart, skeletal muscle, and also adipose tissues [41]. The regulation of CD36 by PPARγ contributes to the control of blood lipids. Interestingly, CD36 null mice exhibit elevated circulating LCFA and triglycerides levels consistent with the phenotype of dKO mice and CD36 deficiency partially protected from HFD-induced insulin resistance [42]. Because of down regulated PPARγ levels in adipose tissues of dKO mice we decided to study in more detail the regulation of CD36 in these as well. We have found strongly reduced CD36 mRNA expression levels in WAT from dKO mice (64 % of the WT values) (Figure 8a). Next, we analyzed CD36 in WT and dKO MEFs

before onset and during adipocyte differentiation. As long as CD36 mRNA expression substantially increased

in WT MEFs, there were almost undetectable transcript levels of CD36 in dKO cells treated with the adipocyte

differentiation cocktail (Figure 8b). Diacylglycerol acyltransferase 1 (DGAT1), a protein associated with the

enterocytic triglyceride absorption and intracellular lipid processing [83] is besides CD36 another target gene

of adipogenesis master regulator PPARγ [43]. DGAT1 mRNA levels are strongly up regulated during adipocyte

differentiation [44], its promoter region contains a PPARγ binding site [45] and DGAT1 is also negatively regulated

by the MEK/ERK pathway [46] . DGAT1 expression was shown to be increased in TIS7 transgenic mice [3] and

its expression was decreased in the gut of high fat diet-fed TIS7 KO mice [8]. Importantly, DGAT1 expression

in adipocytes and WAT is up regulated by PPARγ activation [43]. Therefore, we measured DGAT1mRNA levels

during the differentiation of WT and dKO MEFs into adipocytes to analyze the role of TIS7 and SKMc15 on

the regulation of this protein involved in adipogenesis and triglyceride processing. As long as DGAT1

expression substantially increased during the differentiation of WT MEFs, there was no difference in DGAT1

mRNA levels in dKO cells treated with the adipocyte differentiation cocktail (Figure 8c). Therefore, we

concluded that both, TIS7 and SKMc15 regulate expression of multiple proteins involved in adipocyte

differentiation and function, thereby contributing to the lean phenotype of dKO mice.

Discussion

In this study we show that simultaneous depletion of TIS7 and of its orthologue SKMc15, a protein recently

identified as a translational inhibitor, caused severe reduction of adipose tissues and resistance against high

fat-induced obesity in mice. We identified two parallel mechanisms leading to a deficiency in adipocyte

differentiation. Firstly, TIS7-regulated Wnt signaling affected Dlk-1 transcription and secondly, our study

provided the first experimental evidence for SKMc15 acting as a translational inhibitor that among other

proteins might control DLK-1 protein levels, thereby also contributing to the regulation of adipogenesis (Figure

9).

dKO mice were phenotypically similar with both, CD36 deficient and DLK-1 transgenic mice, namely in

decreased amounts of WAT and resistance to HFD-induced obesity. Previous studies showed that over-

expression of TIS7 caused increased intestinal lipid transport resulting in elevated body weight gain during

HFD feeding [36]. Knockout of TIS7 and SKMc15 impaired absorption of free fatty acids from the lumen into

enterocytes and reduced rate of fat absorption from intestines into the circulation. Simultaneously, higher

concentrations of free fatty acids in the feces of dKO mice suggested that mice lacking TIS7 and SKMc15

suffer from an intestinal lipid uptake deficiency, yet another contributing reason for the lean phenotype of

these mice (data not shown). In dKO mice we found inhibited PPARγ and C/EBPα levels, induced MEK/ERK

pathway, and decreased expression of CD36 and DGAT1, all hallmarks of up regulated DLK-1, a known

adipogenesis inhibitor. TIS7 and SKMc15 acted after the commitment to the preadipocyte stage since the

elevated DLK-1 and β-catenin levels found in WAT of dKO mice were characteristic for the preadipocyte

stage [47].

It is known that disruption of Wnt signaling in embryonic fibroblasts results in spontaneous adipocyte

differentiation [48] and opposite, stabilized β-catenin keeps cells in the preadipocyte stage [23]. Consistent with

this knowledge we found up regulated β-catenin protein levels in WAT of dKO mice and increased Wnt

signaling activity in dKO MEFs. It has to be mentioned that our data were different to those by Nakamura *et*

al. who showed that following the TIS7 overexpression in adipocytes was up regulated Wnt/β-catenin

signaling and inhibited oil red O staining [5]. However, one has to take into consideration the difference in cell

systems used in these two studies. All results presented in Nakamura's study were obtained in 3T3-L1 cells

fibroblasts following overexpression of TIS7. Opposite to that, we have studied the role of endogenous TIS7

in cells derived from wild type or knockout mice. Moreover, we have found ubiquitous TIS7 expression in all

WT mouse organs without any pre-treatment such as e.g. hypoxia. On the other hand, Nakamura *et al.*

identified up regulated TIS7 expression levels in WAT of obesity model mice. This result, however, fully

supports our findings of lean phenotype in TIS7 and SKMc15 dKO mice.

We demonstrated that in dKO MEFs, induced for adipocyte differentiation, the increase in Wnt signaling led

to up regulated binding of β-catenin to the DLK-1 gene regulatory elements, resulting in sustained DLK-1

expression. In adipogenesis DLK-1 expression is down regulated through histone methylation by COPR5

and PRMT5 that prevent β-catenin binding to the DLK-1 gene [31]. Interestingly, in dKO MEFs we found weaker

binding of dimethylated H4R3 to the DLK-1 gene that was consistent with our previous findings on the

negative role of TIS7 in epigenetic regulation of gene expression including the PRMT5 activity [14].

There was a difference in effects of ectopic expression of TIS7, SKMc15 and their co-expression on DLK-1

levels (data not shown), confirming the hypothesis TIS7 and SKMc15 regulate DLK-1 levels via two

independent pathways/mechanisms. Despite the fact that SKMc15 knockout had no effect on Wnt signaling

it nevertheless affected DLK-1 levels, suggesting a contribution of SKMc15 to its regulation. Since SKMc15

was identified as a novel factor translationally inactivating ribosomes [16] and since down regulation of the

protein synthesis machinery is an essential regulatory event during early adipogenic differentiation [17], we

hypothesize that SKMc15 regulates DLK-1 levels through translational regulation, however additional

investigations will be necessary to confirm this mechanism. Elevated DLK-1 levels in dKO MEFs activated

the MEK/ERK pathway thereby decreasing PPARγ and C/EBPα levels important for adipogenic

differentiation. Besides that, the expression of SOX9 was up regulated, suggesting that the dKO MEFs keep

their proliferative state and cannot enter differentiation into mature adipocytes [20].

Down regulation of PPARγ and C/EBPα implied changes in expression of downstream adipogenesis-related

genes. Among them, decreased CD36 and DGAT1 levels were a plausible explanation of the dKO mice lean

phenotype since it is known that both proteins play a functional role in differentiation of murine adipocytes

and their deficiency impairs fat pad formation independent of lipid uptake [49]. Our findings implicate that TIS7

and SKMc15 play a up to now unknown role in the regulation of CD36 and therefore possibly contribute to

CD36 related pathogenesis of human metabolic diseases, such as hypoglycemia [50], hypertriglyceridemia [51],

and disordered fatty acid metabolism [52, 53].

We and others have previously shown that TIS7 is involved under both, chow and HFD conditions in intestinal

triglyceride absorption. Here we show for the first time the role of TIS7 and its orthologue SKMc15 in

adipocyte differentiation. Moreover, our data presented here document the physiological function of SKMc15

as a translational inhibitor. To our surprise, although TIS7 and SKMc15 share high sequence homology and

a functional role in the adipogenesis, they use two independent regulatory mechanisms.

Methods

Generation of animal models. All animal experiments were performed in accordance with Austrian legislation

BGB1 Nr. 501/1988 i.d.F. 162/2005).

Mice lacking TIS7 were previously described [4]. SKMc15 KO mice generation: targeting construct contained

SKMc15 gene locus exons. A loxP site and a neomycin resistance gene inserted at position 105515

(AY162905). The neomycin cassette was flanked by two frt sites. A second loxP site was inserted at position

102082 (AY162905). Further downstream, 15 additional nucleotides, part of intron 1, exon 2 (splice site:

donor and acceptor) and the CDS of hrGFP from the Vitality hrGFP mammalian expression vector pIRES-

hrGFP-2a (Stratagene) were added. The targeted construct was electroporated into Sv129 mouse ES cells.

After the selection with G418 single cell clones were screened by PCR and confirmed by Southern blot

analysis. After Cre recombinase treatment cell clones were screened by PCR. Clones with floxed gene

deleted, were used for blastocyst injection into C57Bl/6J mice. Two male chimeras with ≥ 40 % or more

agouti coat color were mated to C57Bl6 females. The knockout mouse strain was derived from one male

mouse carrying the allele of interest. Heterozygous mice were back-crossed to C57Bl6 mice for 9

generations. dKO mice were generated by crossing the SKMc15 KO mice with the TIS7 single KO mice. The

resulting mice were screened by PCR and double heterozygous mice used for further breeding until

homozygosity. In order to achieve maximal homogeneity of experimental groups, in all experiments

presented here we used only male mice.

SKMc15 knockout Southern blot analysis. XbaI restriction sites (position 109042; position 102079; position

92253) located in the SKMc15 locus (AY162905). Fragment detected by the Southern probe: 9.6 kb wt

(Supplementary figure 3a, insert, band V); 15 kb in deleted locus (band IV). The 566 bp long probe for

hybridization from genomic DNA (96605 – 97171; AY162905). PCR primer sequences: RK 150 Fwd: 5'-

GGTCCTGCCACTAATGCACTG-3'; RK 151 Rev: 5'- GCAGACAGATGCCAGGAAGAC-3'.

SKMc15 knockout PCR genotyping analysis. hrGFP insert in the 3´ UTR detected by primer GFP2 5´-

AGCCATACCACATTTGTAGAG-3´ and RK101 3´ UTR (5´-TGATGATAGCTTCAAAGAGAA-3´; 100617 –

100591 of the SKMc15 locus (AY162905). PCR product 1700 bp. SKMc15 detection: RG1; 5´-

TGTGGCCTTTATCCTGAGTC-3´; 102286–102266) and RG2; 5´-TGGCTTCATTTACACTACTCCTT-3´;

101860 – 101882 primers (Supplementary figure 3a). WT allele PCR product 426 bp and the targeted allele

PCR product 1772 bp (Supplementary figure 3b). TIS7 genotype tested as explained previously [4]

(Supplementary figure 3c).

Growth monitoring and body composition measurement. Mice weaned at 3 weeks, regular chow diet,

weighed weekly. Dual energy X-ray absorptiometry was measured with Norland scanner (Fisher Biomedical).

Micro-CT experiments performed using vivaCT 40 (Scanco Medical AG). The scans were performed using

250 projections with 1024 samples, resulting in a 38µm isotropic resolution. Tube settings: 45kV voltage,

177µA current, integration time 300ms per projection. Image matrix 1024*1024 voxels and a grayscale depth

of 16 bit. The length of the image stack was individually dependent, starting from the cranial end of the fist

lumbar vertebrae, to the caudal end of the fifth lumbar vertebrae. The image reconstruction and post

processing were performed using the Scanco Medical system software V6.6. For the adipose tissue

evaluation an IPL (image processing language) script by Judex et al., provided by Scanco medical AG was

modified to the scanner individual parameters leading to 2 values lower threshold than in the original script

for the adipose tissue filters 76. The script calculated the total abdominal volume without potential air in the

cavities. A separation of subcutaneous and visceral fat mass was used only for visualization. For the

quantitative fat mass analysis we used 56 male 5 to 12 month old mice (mean age 9.35 ± 2.03 month)

reflecting an adult to mid aged cohort defined by Flurkey et al. 2007 [54]. The development of adipose tissue

in healthy mice is stable between 4 and 12 months [55]. Therefore, sex and age could have only very limited

influence on the experimental results. For the quantitative comparison, the percent contribution of the

abdominal fat to the body weight was calculated using a mean weight of 0.9196 g/ml for adipose tissue [54].

Statistics were performed using an ANCOVA with a Bonferroni corrected Post Hoc testing on the µCT fat

data and the body mass data.

Metabolic measurements. The indirect calorimetry trial monitoring gas exchange, activity, and food intake

was conducted over 21 hours (PhenoMaster TSE Systems). Body mass and rectal body temperature before

and after the trial were measured. The genotype effects were statistically analyzed using 1-way ANOVA.

Food intake and energy expenditure were analyzed using a linear model including body mass as a co-variate.

HFD feeding. Age-matched (7-week-old) male TIS7-/- SKMc15-/- and WT mice were caged individually and

maintained up to 3 weeks on a synthetic, high saturated fat (HFD) diet (TD.88137; Ssniff). Animals were

weighted every 4th day between 08:00 and 10:00. Small intestines were harvested for oil red O staining to

detect lipid accumulation. Intestines, liver, muscles and adipose tissue were collected for total RNA and

protein isolation. Unfixed intestines were flushed with PBS using a syringe, embedded in Tissue-Tek (Sakura,

4583) and frozen in liquid nitrogen for immunohistochemical analysis.

Quantitative food consumption and fecal fat determination. Adult mice were acclimatized to individual caging

and to the HFD for a week, monitored for weight gain and their food intake daily. The daily food intake data

were pooled for the following 7 days and the food intake was estimated (g/day). Feces were collected daily

and weighted for 7 days after second week of the HFD consumption.

Antibodies, viral and cDNA constructs. Antibodies: anti-CD36 AbD Serotec (MCA2748), Abcam (ab36977),

p44/42 MAPK (Erk1/2) and Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) Cell Signaling Technology

(9102, 9101), β-catenin antibody Sigma (C2206), anti-DLK Abcam (ab119930), anti-histone H4R3me2s

antibody Active Motif (61187). For ChIP experiments were used anti-TIS7 [11] and anti-SKMc15 [4] rabbit

polyclonal antibodies previously proven for ChIP suitability in [14]. pTOPflash reporter construct was a gift from

H. Clevers (University of Utrecht, Holland). TIS7 construct was described previously [11]. Partial cds of

mSKMc15 was amplified by PCR and cloned into pcDNA3.1(-)/MycHis6 (Invitrogen).

Cell culture and adipocyte differentiation. MEFs were generated from 16-day old embryos. After dissection

of head for genotyping, removal of limbs, liver and visceral organs embryos were minced and incubated in 1

mg/mL collagenase (Sigma-Aldrich, C2674) 30 mins at 37°C. Embryonic fibroblasts were maintained in

growth medium containing DMEM high glucose (4.5 g/l); sodium pyruvate, L-glutamine, 10% FCS (Invitrogen,

41966029), 10% penicillin/streptomycin at 37°C in 5% CO_2. Adipocyte differentiation treatment was medium

with 0.5 mM 3-isobutyl-1-methylxanthine (I5879), 1 µM dexamethasone (D4902), 5 µg/ml insulin (I2643) and

1 µM Rosiglitazone (Sigma-Aldrich, R2408). After 3 days growth medium containing 1 µg/ml insulin cells

were differentiated for 8 days. To visualize lipid accumulation adipocytes were washed with PBS, fixed with

6 % formaldehyde overnight, incubated with 60 % isopropanol, air dried and then incubated with oil red O.

Microscopic analysis was followed by isopropanol elution and absorbance measurement at 490 nm.

Analysis of translation by metabolic labelling. Pulse labelling of mitochondrial proteins was performed as

described before [56], with the following changes: 1x 10^6 wild-type and dKO MEFs cells per plate were seeded

and cultivated for 24 h. Cells were washed twice with PBS followed by one wash with labelling medium (ESC

medium without cysteine and methionine) and a 30 min incubation in labelling medium. To label newly

translated products, 200 µCi of [^{35}S]-methionine (10 mCi/ml; Hartman Analytic) were added and the cells

were incubated for 1 h. Cells were washed and incubated another 10 min in standard medium at 37 °C. Cells

were then harvested by trypsinizing, washed once with ice-cold PBS, extracted with ice-cold RIPA buffer

containing protease-inhibitors and briefly sonicated. Proteins were fractionated by gel electrophoresis in 16%

Tricine gels (Invitrogen) and stained with Coomassie brilliant blue, and radioactive signals were visualized

by phosphorimaging. Signal intensities were quantified using the Image Studio Lite (v5.2) software.

RT-PCR. Tissues from animals fed for 3 weeks with HFD were snap frozen and stored at –80°C. Total RNA

was isolated using the TRIzol reagent (Invitrogen, 15596026). RNA was then chloroform-extracted and

precipitated with isopropanol. The yield and purity of RNA was determined by spectroscopic analysis; RNA

was stored at –80°C until use.

Quantitative RT-PCR and statistics. Total RNA were treated with DNAse1 and reverse transcribed to cDNA

by Revert Aid First Strand cDNA Synthesis Kit (Thermo Scientific, K1622) with oligo dT primers. Quantitative

RT-PCR was performed using Taqman probes and primer sets (Applied Biosystems) specific for CD36

(assay ID Mm00432398_m1), Dgat1 (Mm00515643_m1), Pparγ (Mm00440940_m1), C/EBPα

(Mm00514283_s1), DLK-1 (Mm00494477_m1) and Sox9 (Mm00448840_m1). Ribosomal protein 20 (assay

ID Mm02342828_g1) was used as normalization control for quantification by the ddCt method. PCR reactions

were performed using 10 µl cDNA in PikoReal 96 real-time PCR system (Thermo Scientific). Quantification

data were analyzed by 2-tailed, type 2 Student's t test.

Transient transfections and luciferase assay. pGL2-Basic (Promega, E1641) or pGLCD36 [57] were used as

reporter constructs. Expression constructs or empty vector DNA as a control were co-transfected. pCMV-β-

Gal plasmid was used to normalize for transfection efficiency. For luciferase reporter assays, 1.5x10⁵ cells

were seeded into 24-well plates and transfected after 24 hours with indicated plasmid combinations using

Lipofectamine Plus™ Reagent (Invitrogen, 15338030). The total amount of transfected DNA (2 µg DNA per

well) was equalized by addition of empty vector DNA. Cells were harvested 48 h post-transfection in 0.25 M

Tris, pH 7.5, 1% Triton X-100 buffer and assayed for both luciferase and β-galactosidase activities. Luciferase

activity and β-galactosidase activity were assayed in parallel by using the Lucy 2® detection system (Anthos).

Transfections were performed in triplicates and all experiments were repeated several times.

Chromatin immunoprecipitation (ChIP). Chromatin was isolated from TIS7 WT and dKO formaldehyde-

treated, 8 days adipocyte differentiated MEFs using the EpiSeeker Chromatin Extraction Kit (Abcam,

ab117152). ChIP analyses were carried out as described previously [58]. Sequence of the oligonucleotides for

two regions of the Dlk-1 promoter, encompassing TCF and β-catenin binding sites were as defined in [31].

Sonicated chromatin was centrifuged at 15.000x g for 10 min at 4°C, and the supernatant (65 µg of sheared

DNA per each IP) was diluted ten-fold with cold ChIP dilution buffer containing 16.7 mM Tris-HCl pH 8.1, 167

mM NaCl, 0.01% (w/v) SDS, 1.1% (w/v) Triton X-100 and 1.2 mM EDTA with protease inhibitors. Samples

were pre-cleared for 1 h with protein A Sepharose CL-4B (Sigma-Aldrich, 17-0780-01) beads blocked with

0.2 µg/µl sonicated herring sperm DNA (Thermo Fisher, 15634017) and 0.5 µg/µl BSA (NEB, B9000 S).

Immunoprecipitations were performed at 4°C overnight. Immune complexes were collected with protein A

Sepharose for 1h at 4°C followed by centrifugation at 1000 rpm and 4°C for 5 min. Beads were washed with

1ml low salt wash buffer (20 mM Tris-HCl pH 8.1, 150 mM NaCl, 0.1%(w/v) SDS, 1%(w/v) Triton X-100

(Merck), 2 mM EDTA), high salt wash buffer (20 mM Tris-HCl pH 8.1, 500 mM NaCl, 0.1% (w/v) SDS, 1%

(w/v) Triton X-100, 2 mM EDTA), LiCl wash buffer (10 mM Tris-HCl pH 8.1, 250 mM LiCl, 1% (w/v) sodium

deoxycholate, 1% (w/v) IGEPAL-CA630, 1 mM EDTA) for 5 min at 4°C on a rotating wheel, and twice with 1

ml TE buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA). Protein-DNA complexes were eluted from antibodies by

adding a freshly prepared elution buffer containing 1% SDS and 0.1 M $NaHCO_3$. The eluate was reverse

cross linked by adding NaCl to a final concentration of 0.2 M and incubating at 65 °C for four hours.

Afterwards the eluate was treated with Proteinase K at 45 °C for 1 hour. The immunoprecipitated DNA was

then isolated by phenol/chloroform precipitation and used as a template for real-time-quantitative PCR. The

primer pairs specific for regulatory regions of the Dlk-1 gene were selected as described before [31]. Reactions

with rabbit IgG or with 1.23% of total chromatin (input) were used as controls. For real-time-quantitative PCR

a PikoReal System was used. Signals were normalized to input chromatin and shown as % input. The raw

cycle threshold (Ct) values of the input were adjusted to 100% by calculating raw Ct – log2(100/input). To

calculate the % input of the immunoprecipitations, the equation 100 × 2[Ct (adjusted input to 100%) – Ct (IP)]

was applied.

Statistics. Unpaired, 2-tailed Student's t tests were used for all statistical comparisons. A P value of < 0.05

was considered significant.

Acknowledgements

The authors thank Robert Kurzbauer for generation of dKO mice, Stephan Geley, Laura M. de Smalen and

Karin Schluifer for the technical assistance, David Teis and Zlatko Trajanoski for critical reading of the

manuscript. Furthermore, the Animal Facility of Medical University Innsbruck for their care of our mice. This

work was supported by P18531-B12 and P22350-B12 grants from the Austrian FWF grant agency to Ilja

Vietor and by the German Federal Ministry of Education and Research (Infrafrontier grant 01KX1012) to

Martin Hrabe de Angelis.

Author contributions

I.V., D.C., M.D.E., V.K., J.Ro., M.K., T.V. and L.A.H. designed the study. I.V., D.C., K.P., R.G., J.Ro., M.H.,

V.K., G.D., F.S., and T.V. performed experiments. I.V., D.C., T.V. and L.A.H. wrote the manuscript.

Figure legends

Figure 1. dKO mice display postnatal growth retardation and less body fat.

a) Growth curves of WT and dKO mice on chow diet (n fluctuate depending on the time point of

measurement). b) DEXA measurements of WT and dKO mice. Left panel depicts the absolute values of fat

and lean mass per animal and the right panel represents values normalized to the total body weight of

animals (n=15). c) Micro-computed tomography measurements identified a lack of abdominal fat in single

and dKO mice. 3-dimensional reconstitution of images of the abdominal fat mass distribution in WT and KO

mice. Yellow color represents subcutaneous and red visceral fat mass. Black compartments are shadows

resulting as part of the lightning model for the 3D volume rendering.

Figure 2. dKO mice are resistant against HFD-induced obesity.

a) Mass contribution [%] of the abdominal fat in correlation to the total body weight [g]. A linear regression

is overlaid for each group individually. The regression results for the WT group are y= 0,342x-11,13 with a

R^2 of 0,74; for the TIS7 KO group y=0,143x-3,91 with a R^2 of 0,82; for the SKMc15 KO y=0,005x+0,48 with

a R^2 of 0,04 and for the dKO y=0,011+0,01 with a R^2 of 0,1. ANCOVA analysis for the fat mass as a

percentage of the body weigth as covariant was performed. b) Effect of single SKMc15 or TIS7 knockout

and TIS7 SKMc15 double knockout on total body weight in [g] and body fat amount normalized to the body

weight in [%]. c) TIS7 and SKMc15 knockout reduced the gain of body weight of mice fed with HFD. 7

weeks old male WT and dKO mice were caged individually and maintained up to 21 days on HFD. Data

shown are mean ± SEM, n≥9 per genotype, *P< 0.05, **P < 0.01, *** P < 0.001.

Figure 3. Adipocyte differentiation and Wnt signaling pathway are significantly affected in dKO

mice. a) Representative oil red O staining of MEF cells following 8 days of the adipocyte differentiation

protocol. Images were cropped and space bars represents always 20µm. b) Quantification of oil red O

staining from three independent adipocyte differentiation experiments. Student's t-test ***P<0.001. c) Wnt

signaling activity was measured in MEFs transiently co-transfected with the Tcf reporter plasmid pTOPflash

and β-galactosidase expression vectors. Transfection efficiency was normalized using the β-galactosidase

values. The experiment was repeated three times. Data shown are mean ± SEM. d) Top panel: Western

blot analysis of β-catenin amounts in WAT samples. One representative Western blot from 3 biological

repetitions is shown. Bottom panel: DLK-1 is significantly up regulated both in TIS7, SKMc15 single KO and

also in dKO mice.

Figure 4. DLK-1 levels are lower and β-catenin binding to the Dlk-1 gene is increased in dKO MEFs.

a) DLK-1 mRNA expression measured by qPCR in WAT and MEFs. Values were normalized on GAPDH,

n=3. Error bars indicate standard deviations. b) DLK-1 mRNA expression measured in MEFs treated with

the adipocyte differentiation cocktail for given times. WT MEFs values were set as 1. c) DLK-1 protein

Western blot analysis in MEFs after 8 days of adipocyte differentiation. Mean of 3 biological repeats; insert

is one representative Western blot. d) DLK-1 mRNA levels in dKO MEFs were down regulated following

ectopic co-expression of TIS7 and SKMc15. mRNA expression levels were analyzed by RT qPCR 48 hours

following the transfection. WT MEFs values were set as 1, n=3; Student's t-test *P<0.05. Error bars indicate

standard deviations. e) Recruitment of indicated proteins to regulatory regions of the Dlk-1 promoter in WT

and KO MEFs was analyzed by ChIP at day 8 of the adipocyte differentiation. Values are expressed as the

percentage of immunoprecipitated chromatin relative to input and are the mean of triplicates. ChIP analysis

identified increased specific β-catenin binding to its DLK-1 regulatory element in dKO samples. Pre-immune

serum and IgG were used as background controls. n=3, Data shown are mean ± SEM, Student's t-test ***P

< 0.001.

Figure 5. SKMc15 is a translational inhibitor. Up regulated DLK-1 in dKO MEFs inhibits adipocyte

differentiation. a) translational analysis – *in vivo* metabolic staining of MEFs with ^{35}S-methionine. Equal

numbers of cells were seeded and 24 hours later treated as explained in Methods section. Identical volumes

of cell lysates were separated by SDS-PAGE, gel was dried and analyzed by a phosphorimager. Equal

loading was documented by the Coomassie blue staining of the gel. b) dKO MEFs secrete DLK-1 protein into

the cell culture medium. Identical volumes of media from WT and dKO cells 8 days treated with the adipocyte

differentiation cocktail were analyzed by Western blot. Equal loading of samples was evaluated by amido

black staining of the membrane. c) dKO MEFs-conditioned medium inhibited adipocyte differentiation in WT

MEFs. MEFs were treated 8 days with the adipocyte differentiation cocktail. WT MEFs treated with the dKO-

conditioned medium showed reduced differentiation. Images were cropped and space bars represent always

20µm. d) Quantification of the oil red O staining of samples shown in panel C.

Figure 6. MAPK signaling and Sox9 expression are induced in dKO MEFs.

a) Representative Western blots of phospho-p44 and phospho-p42 from 8 days adipocyte differentiation

cocktail-treated MEFs. b) Normalization on p44 and p42; n=3. Error bars indicate standard deviations. WT

values were set as 100 %. c) SOX9 mRNA levels were up regulated in dKO MEFs. SOX9 expression was

measured by qPCR in 8 days adipocyte-differentiated WT and dKO MEFs. SOX9 values were normalized

on GADPH expression. WT MEFs values were set as 1, n=3. Data shown are mean ± SEM. Student's t-test

**P < 0.01.

Figure 7. Adipocyte differentiation regulatory genes are down regulated in dKO mice.

a) PPARγ and b) C/EBPα mRNA levels were down regulated in dKO adipocyte-differentiated MEFs. Gene

expression was measured by qPCR during the treatment with the adipocyte differentiation cocktail. Values

were normalized on GADPH expression. WT MEFs values were set as 1, n=3. Data shown are mean ±

SEM. c) Ectopic co-expression of TIS7 and SKMc15 in undifferentiated MEFs significantly increased levels

of adipogenic genes PPARγ and d) C/EBPα. Student's t-test **P < 0.01, ***P < 0.001.

Figure 8. Target genes of adipocyte differentiation regulatory genes are down regulated in dKO mice.

a) qPCR analysis of CD36 mRNA levels in WAT, *P< 0.05 b) qPCR analysis of CD36 in MEFs

differentiating into adipocytes. Relative expression levels were normalized on GAPDH expression. WT

MEFs values were set as 1. c) DGAT1 mRNA levels were down regulated in dKO adipocyte-differentiated

MEFs, ***P < 0.001.

Figure 9. Proposed model of TIS7 and SKMc15 molecular mechanisms of action during adipocyte

differentiation.

Two parallel mechanisms leading to a deficiency in adipocyte differentiation: TIS7-regulated Wnt signaling

affected Dlk-1 transcription and SKMc15 acting as a translational inhibitor also contributed to the regulation

of adipogenesis.

Conflict of interest.

The authors declare that they have no conflict of interest.

References

1.   Sarantopoulos CN, Banyard DA, Ziegler ME, Sun B, Shaterian A, Widgerow AD. Elucidating the
Preadipocyte and Its Role in Adipocyte Formation: a Comprehensive Review. *Stem Cell Rev* 2018,
14(1)**:** 27-42.

612
613 2.   Iacopetti P, Barsacchi G, Tirone F, Cremisi F. Expression of the PC4 gene in the developing rat
614   nervous system. *Brain Res* 1996, **707**(2)**:** 293-297.

615
616 3.   Wang Y, Iordanov H, Swietlicki EA, Wang L, Fritsch C, Coleman T*, et al.* Targeted intestinal
617   overexpression of the immediate early gene tis7 in transgenic mice increases triglyceride
618   absorption and adiposity. *J Biol Chem* 2005, **280**(41)**:** 34764-34775.

619
620 4.   Vadivelu SK, Kurzbauer R, Dieplinger B, Zweyer M, Schafer R, Wernig A*, et al.* Muscle
621   regeneration and myogenic differentiation defects in mice lacking TIS7. *Mol Cell Biol* 2004,
622   **24**(8)**:** 3514-3525.

623
624 5.   Nakamura Y, Hinoi E, Iezaki T, Takada S, Hashizume S, Takahata Y*, et al.* Repression of
625   adipogenesis through promotion of Wnt/beta-catenin signaling by TIS7 up-regulated in
626   adipocytes under hypoxia. *Biochim Biophys Acta* 2013, **1832**(8)**:** 1117-1128.

627
628 6.   Vietor I, Kurzbauer R, Brosch G, Huber LA. TIS7 regulation of the beta-catenin/Tcf-4 target gene
629   osteopontin (OPN) is histone deacetylase-dependent. *J Biol Chem* 2005, **280**(48)**:** 39795-39801.

7.   Iezaki T, Onishi Y, Ozaki K, Fukasawa K, Takahata Y, Nakamura Y*, et al.* The Transcriptional Modulator Interferon-Related Developmental Regulator 1 in Osteoblasts Suppresses Bone Formation and Promotes Bone Resorption. *J Bone Miner Res* 2016, **31**(3)**:** 573-584.

8.   Yu C, Jiang S, Lu J, Coughlin CC, Wang Y, Swietlicki EA*, et al.* Deletion of Tis7 protects mice from high-fat diet-induced weight gain and blunts the intestinal adaptive response postresection. *J Nutr* 2010, **140**(11)**:** 1907-1914.

9.   Garcia AM, Wakeman D, Lu J, Rowley C, Geisman T, Butler C*, et al.* Tis7 deletion reduces survival and induces intestinal anastomotic inflammation and obstruction in high-fat diet-fed mice with short bowel syndrome. *Am J Physiol Gastrointest Liver Physiol* 2014, **307**(6)**:** G642-654.

10. Micheli L, Leonardi L, Conti F, Buanne P, Canu N, Caruso M*, et al.* PC4 Coactivates MyoD by Relieving the Histone Deacetylase 4-Mediated Inhibition of Myocyte Enhancer Factor 2C. *Mol Cell Biol* 2005, **25**(6)**:** 2242-2259.

11. Vietor I, Vadivelu SK, Wick N, Hoffman R, Cotten M, Seiser C*, et al.* TIS7 interacts with the mammalian SIN3 histone deacetylase complex in epithelial cells. *Embo J* 2002, **21**(17)**:** 4621-4631.

12. Wick N, Schleiffer A, Huber LA, Vietor I. Inhibitory effect of TIS7 on Sp1-C/EBPalpha transcription factor module activity. *J Mol Biol* 2004, **336**(3)**:** 589-595.

13. Park G, Horie T, Kanayama T, Fukasawa K, Iezaki T, Onishi Y*, et al.* The transcriptional modulator Ifrd1 controls PGC-1alpha expression under short-term adrenergic stimulation in brown adipocytes. *The FEBS journal* 2017, **284**(5)**:** 784-795.

14. Lammirato A, Patsch K, Feiereisen F, Maly K, Nofziger C, Paulmichl M*, et al.* TIS7 induces transcriptional cascade of methylosome components required for muscle differentiation. *BMC Biol* 2016, **14**(1)**:** 95.

15. Latif F, Duh FM, Bader S, Sekido Y, Li H, Geil L*, et al.* The human homolog of the rodent immediate early response genes, PC4 and TIS7, resides in the lung cancer tumor suppressor gene region on chromosome 3p21. *Hum Genet* 1997, **99**(3)**:** 334-341.

16. Brown A, Baird MR, Yip MC, Murray J, Shao S. Structures of translationally inactive mammalian ribosomes. *Elife* 2018, **7**.

17. Marcon BH, Holetz FB, Eastman G, Origa-Alves AC, Amoros MA, de Aguiar AM*, et al.* Downregulation of the protein synthesis machinery is a major regulatory event during early adipogenic differentiation of human adipose-derived stromal cells. *Stem Cell Res* 2017, **25:** 191-201.

18. Farmer SR. Transcriptional control of adipocyte formation. *Cell Metab* 2006, **4**(4)**:** 263-273.

19. Wang Y, Sul HS. Pref-1 regulates mesenchymal cell commitment and differentiation through
Sox9. *Cell Metab* 2009, **9**(3)**:** 287-302.

20. Kim KA, Kim JH, Wang Y, Sul HS. Pref-1 (preadipocyte factor 1) activates the MEK/extracellular
signal-regulated kinase pathway to inhibit adipocyte differentiation. *Mol Cell Biol* 2007, **27**(6)**:**
2294-2308.

21. Lefterova MI, Zhang Y, Steger DJ, Schupp M, Schug J, Cristancho A*, et al.* PPARgamma and C/EBP
factors orchestrate adipocyte biology via adjacent binding on a genome-wide scale. *Genes Dev*
2008, **22**(21)**:** 2941-2952.

22. Lowell BB. PPARgamma: an essential regulator of adipogenesis and modulator of fat cell
function. *Cell* 1999, **99**(3)**:** 239-242.

23. Ross SE, Hemati N, Longo KA, Bennett CN, Lucas PC, Erickson RL*, et al.* Inhibition of adipogenesis
by Wnt signaling. *Science* 2000, **289**(5481)**:** 950-953.

24. van Tienen FH, Laeremans H, van der Kallen CJ, Smeets HJ. Wnt5b stimulates adipogenesis by
activating PPARgamma, and inhibiting the beta-catenin dependent Wnt signaling pathway
together with Wnt5a. *Biochem Biophys Res Commun* 2009, **387**(1)**:** 207-211.

25. Li HX, Luo X, Liu RX, Yang YJ, Yang GS. Roles of Wnt/beta-catenin signaling in adipogenic
differentiation potential of adipose-derived mesenchymal stem cells. *Mol Cell Endocrinol* 2008,
291(1-2)**:** 116-124.

26. Tontonoz P, Spiegelman BM. Fat and beyond: the diverse biology of PPARgamma. *Annu Rev*
Biochem 2008, **77:** 289-312.

27. Xu C, Wang J, Zhu T, Shen Y, Tang X, Fang L*, et al.* Cross-Talking Between PPAR and WNT
Signaling and its Regulation in Mesenchymal Stem Cell Differentiation. *Curr Stem Cell Res Ther*
2016, **11**(3)**:** 247-254.

28. Mortensen SB, Jensen CH, Schneider M, Thomassen M, Kruse TA, Laborda J*, et al.* Membrane-
tethered delta-like 1 homolog (DLK1) restricts adipose tissue size by inhibiting preadipocyte
proliferation. *Diabetes* 2012, **61**(11)**:** 2814-2822.

29. Moon YS, Smas CM, Lee K, Villena JA, Kim KH, Yun EJ*, et al.* Mice lacking paternally expressed
Pref-1/Dlk1 display growth retardation and accelerated adiposity. *Mol Cell Biol* 2002, **22**(15)**:**
5585-5592.

30. Barclay JL, Nelson CN, Ishikawa M, Murray LA, Kerr LM, McPhee TR*, et al.* GH-dependent STAT5
signaling plays an important role in hepatic lipid metabolism. *Endocrinology* 2011, **152**(1)**:** 181-
192.

719
720 31. Paul C, Sardet C, Fabbrizio E. The Wnt-target gene Dlk-1 is regulated by the Prmt5-associated
721   factor Copr5 during adipogenic conversion. *Biol Open* 2015, **4**(3)**:** 312-316.

722
723 32. LeBlanc SE, Konda S, Wu Q, Hu YJ, Oslowski CM, Sif S*, et al.* Protein arginine methyltransferase 5
724   (Prmt5) promotes gene expression of peroxisome proliferator-activated receptor gamma2
725   (PPARgamma2) and its target genes during adipogenesis. *Mol Endocrinol* 2012, **26**(4)**:** 583-597.

726
727 33. Sul HS. Minireview: Pref-1: role in adipogenesis and mesenchymal cell fate. *Mol Endocrinol*
728   2009, **23**(11)**:** 1717-1725.

729
730 34. Paul C, Sardet C, Fabbrizio E. The histone- and PRMT5-associated protein COPR5 is required for
731   myogenic differentiation. *Cell Death Differ* 2012, **19**(5)**:** 900-908.

732
733 35. Ruiz-Ojeda FJ, Ruperez AI, Gomez-Llorente C, Gil A, Aguilera CM. Cell Models and Their
734   Application for Studying Adipogenic Differentiation in Relation to Obesity: A Review. *Int J Mol Sci*
735   2016, **17**(7).

736
737 36. Wang S, Sun Z, Zhang X, Li Z, Wu M, Zhao W*, et al.* Wnt1 positively regulates CD36 expression
738   via TCF4 and PPAR-gamma in macrophages. *Cell Physiol Biochem* 2015, **35**(4)**:** 1289-1302.

739
740 37. Prestwich TC, Macdougald OA. Wnt/beta-catenin signaling in adipogenesis and metabolism. *Curr
741   Opin Cell Biol* 2007, **19**(6)**:** 612-617.

742
743 38. Lee YL, Helman L, Hoffman T, Laborda J. dlk, pG2 and Pref-1 mRNAs encode similar proteins
744   belonging to the EGF-like superfamily. Identification of polymorphic variants of this RNA.
745   *Biochim Biophys Acta* 1995, **1261**(2)**:** 223-232.

746
747 39. Garces C, Ruiz-Hidalgo MJ, Bonvini E, Goldstein J, Laborda J. Adipocyte differentiation is
748   modulated by secreted delta-like (dlk) variants and requires the expression of membrane-
749   associated dlk. *Differentiation* 1999, **64**(2)**:** 103-114.

750
751 40. Lee K, Villena JA, Moon YS, Kim KH, Lee S, Kang C*, et al.* Inhibition of adipogenesis and
752   development of glucose intolerance by soluble preadipocyte factor-1 (Pref-1). *J Clin Invest* 2003,
753   **111**(4)**:** 453-461.

754
755 41. Coburn CT, Knapp FF, Jr., Febbraio M, Beets AL, Silverstein RL, Abumrad NA. Defective uptake
756   and utilization of long chain fatty acids in muscle and adipose tissues of CD36 knockout mice. *J
757   Biol Chem* 2000, **275**(42)**:** 32523-32529.

759   42.   Wilson CG, Tran JL, Erion DM, Vera NB, Febbraio M, Weiss EJ. Hepatocyte-Specific Disruption of
760   CD36 Attenuates Fatty Liver and Improves Insulin Sensitivity in HFD-Fed Mice. *Endocrinology*
761   2016, **157**(2)**:** 570-585.

762
763   43.   Koliwad SK, Streeper RS, Monetti M, Cornelissen I, Chan L, Terayama K*, et al.* DGAT1-dependent
764   triacylglycerol storage by macrophages protects mice from diet-induced insulin resistance and
765   inflammation. *J Clin Invest* 2010, **120**(3)**:** 756-767.

766
767   44.   Cases S, Smith SJ, Zheng YW, Myers HM, Lear SR, Sande E*, et al.* Identification of a gene
768   encoding an acyl CoA:diacylglycerol acyltransferase, a key enzyme in triacylglycerol synthesis.
769   *Proc Natl Acad Sci U S A* 1998, **95**(22)**:** 13018-13023.

770
771   45.   Ludwig EH, Mahley RW, Palaoglu E, Ozbayrakci S, Balestra ME, Borecki IB*, et al.* DGAT1 promoter
772   polymorphism associated with alterations in body mass index, high density lipoprotein levels
773   and blood pressure in Turkish women. *Clin Genet* 2002, **62**(1)**:** 68-73.

774
775   46.   Tsai J, Qiu W, Kohen-Avramoglu R, Adeli K. MEK-ERK inhibition corrects the defect in VLDL
776   assembly in HepG2 cells: potential role of ERK in VLDL-ApoB100 particle assembly. *Arterioscler
777   Thromb Vasc Biol* 2007, **27**(1)**:** 211-218.

778
779   47.   Gautam J, Khedgikar V, Kushwaha P, Choudhary D, Nagar GK, Dev K*, et al.* Formononetin, an
780   isoflavone, activates AMP-activated protein kinase/beta-catenin signalling to inhibit
781   adipogenesis and rescues C57BL/6 mice from high-fat diet-induced obesity and bone loss. *Br J
782   Nutr* 2017, **117**(5)**:** 645-661.

783
784   48.   Bennett CN, Hodge CL, MacDougald OA, Schwartz J. Role of Wnt10b and C/EBPalpha in
785   spontaneous adipogenesis of 243 cells. *Biochem Biophys Res Commun* 2003, **302**(1)**:** 12-16.

786
787   49.   Christiaens V, Van Hul M, Lijnen HR, Scroyen I. CD36 promotes adipocyte differentiation and
788   adipogenesis. *Biochim Biophys Acta* 2012, **1820**(7)**:** 949-956.

789
790   50.   Nagasaka H, Yorifuji T, Takatani T, Okano Y, Tsukahara H, Yanai H*, et al.* CD36 deficiency
791   predisposing young children to fasting hypoglycemia. *Metabolism* 2011, **60**(6)**:** 881-887.

792
793   51.   Kashiwagi H, Tomiyama Y, Nozaki S, Kiyoi T, Tadokoro S, Matsumoto K*, et al.* Analyses of genetic
794   abnormalities in type I CD36 deficiency in Japan: identification and cell biological
795   characterization of two novel mutations that cause CD36 deficiency in man. *Hum Genet* 2001,
796   **108**(6)**:** 459-466.

797
798   52.   Glazier AM, Scott J, Aitman TJ. Molecular basis of the Cd36 chromosomal deletion underlying
799   SHR defects in insulin action and fatty acid metabolism. *Mamm Genome* 2002, **13**(2)**:** 108-113.

53. Tanaka T, Nakata T, Oka T, Ogawa T, Okamoto F, Kusaka Y*, et al.* Defect in human myocardial
long-chain fatty acid uptake is caused by FAT/CD36 mutations. *J Lipid Res* 2001, **42**(5)**:** 751-759.

54. Neeland IJ, Ayers CR, Rohatgi AK, Turer AT, Berry JD, Das SR*, et al.* Associations of visceral and
abdominal subcutaneous adipose tissue with markers of cardiac and metabolic risk in obese
adults. *Obesity (Silver Spring)* 2013, **21**(9)**:** E439-447.

55. Lemonnier D. Effect of age, sex, and sites on the cellularity of the adipose tissue in mice and rats
rendered obese by a high-fat diet. *J Clin Invest* 1972, **51**(11)**:** 2907-2915.

56. Popow J, Alleaume AM, Curk T, Schwarzl T, Sauer S, Hentze MW. FASTKD2 is an RNA-binding
protein required for mitochondrial RNA processing and translation. *RNA* 2015, **21**(11)**:** 1873-
1884.

57. Shore P, Dietrich W, Corcoran LM. Oct-2 regulates CD36 gene expression via a consensus
octamer, which excludes the co-activator OBF-1. *Nucleic Acids Res* 2002, **30**(8)**:** 1767-1773.

58. Reintjes A, Fuchs JE, Kremser L, Lindner HH, Liedl KR, Huber LA*, et al.* Asymmetric arginine
dimethylation of RelA provides a repressive mark to modulate TNFalpha/NF-kappaB response.
Proc Natl Acad Sci U S A 2016, **113**(16)**:** 4326-4331.

a

n=9-22

Vietor_Figure 1

n=7-17

b

DEXA analysis

absolute mass

relative to body weight

c

WT

TIS7
KO

visceral
fat

subcutaneous
fat

SKMc15
KO

TIS7
SKMc15
dKO

5 mm

a

b

c

a

b

c

d

a

P-p44
P-p42
- 43 kDa
p44
p42
- 43 kDa
WT   dKO

8d dif. MEFs

b

c

a

b

c

d

a

b

c