Extraction of active RhoGTPases by RhoGDI regulates spatiotemporal patterning of RhoGTPases

Adriana Golding[1†], Ilaria Visco[2†], Peter Bieling[2††], William Bement[3,4††]

[1]Graduate Program in Cell and Molecular Biology, University of Wisconsin-Madison, Madison, WI, USA

[2]Department of Systemic Cell Biology, Max Planck Institute of Molecular Physiology, Dortmund, Germany

[3]Laboratory of Cell and Molecular Biology, University of Wisconsin-Madison

[4]Department of Integrated Biology, University of Wisconsin-Madison

[†] **co-first authors**

[††]**co-corresponding authors**

Addresses for correspondence:

William Bement   Peter Bieling

Laboratory of Cell and Molecular Biology   Department of Systemic Cell Biology

University of Wisconsin-Madison   Max Planck Institute of Molecular Physiology

1525 Linden Drive   Otto-Hahn-Straße 11

Madison, WI 53706   44227 Dortmund, Germany

wmbement@wisc.edu   peter.bieling@mpi-dortmund.mpg.de

Abstract

The RhoGTPases are characterized as membrane-associated molecular switches cycling

between active, GTP-bound and inactive, GDP-bound states. However, 90-95% of RhoGTPases

are maintained in a soluble form by RhoGDI, which is generally viewed as a passive shuttle for

inactive RhoGTPases. Our current understanding of RhoGTPase:RhoGDI dynamics has been

limited by two experimental challenges: direct visualization of the RhoGTPases *in vivo* and

reconstitution of the cycle *in vitro*. We developed methods to directly image vertebrate

RhoGTPases *in vivo* or on lipid bilayers *in vitro*. Using these tools, we identified pools of active

and inactive RhoGTPase associated with the membrane, showed that RhoGDI can actively

extract both inactive and active RhoGTPases, and that the extraction of active RhoGTPase

contributes to their spatial regulation around wounds. In contrast to the textbook model of the

RhoGTPase cycle, these results indicate that RhoGDI actively contributes to spatiotemporal

patterning by removing active RhoGTPases from the plasma membrane.

Introduction

      The Rho family GTPases, including Rho, Rac and Cdc42, are essential signaling proteins that mediate morphological changes in cells by directing local cytoskeletal rearrangements (Bishop & Hall, 2000; Kimura et al., 1996). These rearrangements are generally initiated at and confined to specific subcellular regions. For example, a narrow, concentrated zone of Rho activity directs the formation of a ring of actin filaments and myosin-2 at the equatorial cortex that drives cytokinesis (Bement, Benink, and Von Dassow 2005; Yonemura, Hirao-Minakuchi, and Nishimura 2004; Yüce, Piekny, and Glotzer 2005). Similarly, Rho, Rac and Cdc42 are activated near the leading edge of crawling cells in patterns that correspond to local cycles of protrusion, adhesion and retraction (Machacek et al., 2009; Martin et al., 2016). Because tight spatiotemporal regulation of the GTPases is a fundamental feature of these cellular processes, considerable effort has been invested in studying GTPase regulation.

      The RhoGTPases are classically characterized as cycling between membrane-associated, active states and soluble, inactive states as a result of interactions with three classes of regulatory proteins: guanine nucleotide exchange factors (GEFs), which activate GTPases by promoting exchange of GDP for GTP (Rossman et al. 2005); GTPase activating proteins (GAPs), which inactivate GTPases by promoting GTP hydrolysis (Moon & Zheng, 2003); and guanine nucleotide dissociation inhibitor (GDI), which solubilizes GTPases to generate a large reservoir of heterodimeric GTPase:GDI complexes in the cytoplasm (Garcia-Mata, Boulter, & Burridge, 2011). In the canonical model of GTPase regulation, GTPase cycling is thought to proceed as follows: a GTPase is activated by a GEF at the plasma membrane following its release from GDI, is subsequently inactivated by a GAP, and is then returned to the soluble pool by GDI. Thus, in the canonical model, the lifetime of GTPase activity at the plasma membrane is thought

to be controlled entirely by GEFs and GAPs, with GDI essentially serving as a passive shuttle

that interacts exclusively with inactive GTPases.

A limitation of this traditional view is that the function and the biochemical activities of

RhoGDI, in contrast to GEFs and GAPs, are not well understood. Presently, no consensus exists

as to i) whether GDI actively extracts GTPases from membranes or merely solubilizes them by

sequestration (Johnson, Erickson, and Cerione 2009; Zhang et al. 2014), ii) whether it interacts

with GTPases in a nucleotide-specific manner (Nomanbhoy & Cerione, 1996; Tnimov et al.,

2012), or iii) how its activity is coordinated with GEFs or GAPs (Garcia-Mata et al., 2011). The

notion that GDI works as a passive shuttle rests largely on two findings. First, when

GTPase:GDI complexes are purified from cell lysates, the great majority of GTPase within the

complex is in the inactive, GDP-bound form (Abo, Webb, Grogan, & Segal, 1994), as expected

if the GDI solubilizes GTPases after inactivation by a GAP. Second, binding of GTPases by GDI

strongly suppresses GTP hydrolysis (Hart et al., 1992), indicating that hydrolysis must precede

the extraction from the membrane. However, other results raise a serious, albeit contentious,

challenge to this idea: multiple studies indicate that GDI binds both inactive and active GTPase

with relatively high affinity *in vitro* (Hancock & Hall, 1993; Hart et al., 1992; Nomanbhoy &

Cerione, 1996; Tnimov et al., 2012), implying that GDI may have the potential to interact with

active as well as inactive GTPase *in vivo*. As such, GDI might have the potential to exert a more

direct role in the regulation of GTPase activity than currently appreciated.

Understanding the pattern forming ability of RhoGTPases requires the mechanistic

dissection of the interactions between GTPases and GDI. Unfortunately, our ability to study

GTPase:GDI dynamics has been hampered due to two major experimental limitations: first,

visualization of the GTPases in living cells is limited by the fact that labeling with fluorescent

protein on the amino terminus impairs GTPase regulation and function, while carboxy-terminal

labeling prevents GTPase prenylation (Howell et al., 2012; Yonemura et al., 2004). In the

absence of direct visualization, GTPase dynamics must be inferred from activity probes. Second,

with a few important exceptions (Johnson et al., 2009; Nomanbhoy, Erickson, & Cerione, 1999),

in vitro studies of GTPase:GDI dynamics have either utilized unprenylated GTPases, omitted

membranes, or both. Additionally, nearly all reconstitution experimentation focused on the effect

of GDI on the distribution between membrane-associated and soluble forms of GTPases at

thermodynamic equilibrium (Zhang et al., 2014). Thus, we do not currently understand how GDI

affects the transitions between membrane and soluble GTPase states kinetically, especially under

conditions which mimic the cellular environment which is far from equilibrium due to the

constant dissipation of energy.

To overcome these limitations, we developed two distinct methods to directly visualize

vertebrate RhoGTPases on membranes *in vivo* and or on supported lipid bilayers *in vitro*. Using

these tools, we identify co-existing pools of active and inactive GTPases associated with the

plasma membrane. We also demonstrate that GDI can actively extract GTPases from the

membrane and, unexpectedly, that GDI can extract both inactive and active GTPases. Finally, we

show that the extraction of active GTPase also occurs *in vivo* and that this contributes to the

spatial regulation of GTPase activity. Collectively, these data indicate that the textbook model of

the GTPase cycling must be reassessed because GDI can directly mediate the spatiotemporal

regulation of GTPase activity.

Results

Visualization of RhoGTPases around cell wounds

Traditional fusion of fluorescent protein with the amino- or carboxyl-termini of the

RhoGTPases impairs GTPase localization and function (SuppFig1; Yonemura, Hirao-Minakuchi,

and Nishimura 2004). To overcome this problem, we first adapted an approach described by

Bendezú et al. (2015) for labeling of yeast Cdc42. Specifically, we inserted green fluorescent

protein into a solvent-exposed external loop of the *Xenopus* GTPases (see Methods). To test the

internally-tagged (IT) GTPases *in vivo,* we exploited the cell wound repair model in *Xenopus*

laevis oocytes where wounding elicits a robust accumulation of active Rho and Cdc42 in

discrete, concentric zones at the cortex (Fig1A; Benink & Bement, 2005). IT-GTPases were co-

expressed with wild-type (WT) GDI to avoid GTPase aggregation (Boulter et al., 2010). Both

IT-Rho and IT-Cdc42 were recruited to concentric rings around the wound (Fig1B,C).

Comparison of IT-Rho to a Rho activity reporter (mRFP-2xrGBD; Davenport et al. 2016)

revealed that IT-Rho spatially overlapped with the Rho activity zone. Comparison of IT-Cdc42

to a Cdc42 activity reporter (mRFP-wGBD; Benink and Bement 2005) revealed that IT-Cdc42

localized throughout the active Cdc42 zone, but extended slightly beyond it towards the wound

center (Fig1B,C; see also below). We also tested the behavior of IT-Rac and found that it

concentrated around wounds in the same region as IT-Cdc42 as expected from previous

experiments (SuppFig2; Abreu-Blanco, Verboon, & Parkhurst, 2014; Benink & Bement, 2001).

These results indicate that the IT and Cy3-tagged GTPase variants can interact with diverse

regulators required to achieve their normal enrichment at wounds.

As an alternative approach, and as a means to obtain labeled RhoGTPases that could be

used both *in vivo* and *in vitro*, purified recombinant Rho and Cdc42 were prenylated and coupled

to Cy3 via a short N-terminal peptide via sortase-mediated ligation (see Methods). Microinjected

Cy3-Rho and Cy3-Cdc42 localized to wounds (Fig1D,E), in a manner indistinguishable from

their IT counterparts expressed in the oocyte (Fig1F,G). As observed with IT-Rho and IT-Cdc42,

Cy3-Rho completely overlapped with the zone of Rho activity while Cy3-Cdc42 localized

throughout and slightly interior of the active Cdc42 zone.

Visualization of RhoGTPases in other cellular contexts

To further test the behavior of the IT- and Cy3-labeled RhoGTPases, we sought to

determine if they localize to the plasma membrane in other cellular processes. This is important

because these processes likely depend on different regulators from those that operate during cell

wound repair. IT-Rho and Cy3-Rho localized to the cytokinetic apparatus and epithelial

junctions in *Xenopus* embryos, consistent with previous results obtained with a Rho activity

reporter (Fig2A-C; Bement, Benink, & Von Dassow 2005). Similarly, IT-Cdc42 localized to

exocytosing cortical granules (Fig2D), consistent with previous results obtained with a Cdc42

activity reporter (Yu and Bement 2007). IT-Cdc42 was also recruited to cell-cell junctions and

enriched there upon wounding (Fig2E), a behavior previously revealed using a Cdc42 activity

reporter (Clark et al., 2009).

Next, we wanted to determine whether IT-RhoGTPases can functionally substitute for the

endogenous GTPases. The *Xenopus laevis* oocyte system is not conducive to traditional

knockdown approaches due to its large stores of maternal protein and relatively slow protein

turnover. Therefore, we employed C3-exotransferase, a Rho-specific toxin, to inhibit endogenous

Rho activity, and expressed an IT-Rho in which the C3 ribosylation site (N41) is mutated to a

residue that cannot be ribosylated (N41V; Sekine, Fujiwara, and Narumiyas 1989). In control

oocytes expressing the probe for active Rho, Rho activity around wounds was suppressed by C3

(Fig 2F,G). In contrast, cells expressing IT-Rho-N41V generated a spatially defined zone of Rho

activity around the wound that closed over similar timescales compared to the control.

Collectively, the above results indicate that both IT- and Cy3-labeled GTPases are faithful

reporters of the distribution of GTPases and show that IT-Rho can functionally substitute for its

endogenous counterpart.

Pools of active and inactive RhoGTPase accumulate in the plasma membrane

The observation that Cdc42 extended slightly beyond its zone of activity suggests that

there may be a pool of inactive, membrane-bound Cdc42 at this location. This notion is

consistent with the previous observation that Abr, a Cdc42 GAP thought to regulate Cdc42

activity at wounds, also localizes interior to the Cdc42 zone (Vaughan et al. 2011). To

understand the relationship between activity and membrane-association of Cdc42, we

overexpressed Abr, a manipulation previously shown to decrease Cdc42 activity around wounds

(Vaughan et al., 2011). Remarkably, this resulted in a dose-dependent loss of active Cdc42 at

wounds while having far less effect on Cy3-Cdc42 (Fig3A,B). These results further demonstrate

that the IT- and Cy3-GTPases are functional. More importantly, they demonstrate that substantial

pools of both active and inactive GTPases can be dynamically maintained at the plasma

membrane.

RhoGDI is recruited to concentrated areas of RhoGTPase activity

Efforts to visualize RhoGDI at the plasma membrane have generally failed (Ngo et al.,

2017), likely because GDI only transiently interacts with GTPases upon release into or extraction

from the membrane. However, we reasoned it might be possible to detect GDI at wound sites due

to the high local concentration of Rho and Cdc42. Indeed, we found that 3xGFP-GDI is enriched

at wounds and overlaps with both the Rho and Cdc42 zones (Fig4A). To confirm that

endogenous GDI also localizes to wounds, antibodies were raised against *X. laevis* GDI

(SuppFig3A) and used to immunolabel wounded oocytes. Consistent with the results obtained

with 3XGFP-GDI, endogenous GDI accumulated at wounds (Fig4B). These results demonstrate

that GDI accumulation occurs at discrete regions of the plasma membrane that are enriched with

its GTPase clients.

RhoGDI differentially regulates Rho and Cdc42

The localization of RhoGDI around wounds suggests that it might play an active role in

delivery to or extraction of RhoGTPases from the membrane and thus their spatiotemporal

patterning. As an initial test of this possibility, we overexpressed GDI via mRNA microinjection.

This manipulation potently suppressed both Rho and Cdc42 activity, as well as Cy3-Rho and

Cy3-Cdc42 localization, suggesting that GDI exerts its effects via extraction of the GTPases

(Fig5A). To obtain a more quantitative understanding of the relationship between GDI and

GTPase activity, we microinjected purified GDI (SuppFig3B) at increasing concentrations prior

to wounding. High concentrations of microinjected GDI suppressed both Rho and Cdc42 activity

at wounds (Fig5B,C), consistent with the results obtained from GDI via mRNA-mediated

overexpression. However, more modest increases revealed differential effects on Rho and

Cdc42. Specifically, increases between 18-190% in GDI levels (based on proteomic data from

Wühr et al. 2014) resulted in a greater reduction of Cdc42 activity compared to Rho (Fig5C). We

found the same to be true for bovine GDI (SuppFig4). These results show that GDI differentially

impacts Rho and Cdc42 activity *in vivo* and that this effect does not require gross

overexpression.

RhoGDI extracts active and inactive RhoGTPase in vitro

To directly probe the mechanism by which RhoGDI inhibits Rho and Cdc42 activity *in*

vivo, we established a real-time GTPase dissociation assay on supported lipid-bilayers (SLBs)

(Fig6A; SuppFig5; see Methods). The addition of Cy3-Cdc42 to SLBs resulted in their

membrane binding as detected by total internal reflection microscopy (TIRF), which was

dependent on its C-terminal prenyl moiety, as expected (Fig6B). We then studied the time course

of Cdc42 release from SLBs under buffer flow which continuously flushed out unbound proteins

from solution. Spontaneous release of inactive, GDP-bound Cdc42 from the membrane was

rather slow (t1/2=37.24±3.05 sec), however the addition of excess GDI lead to a dramatic

acceleration of dissociation (ca.20-fold; Fig6C; SuppFig3C). To determine whether this was the

result of either simple sequestration in solution or, alternatively, active catalytic extraction of

Cdc42 from membranes, we performed assays in the presence of an alternative solubilizer

(RabGGTase beta) that sequesters the GTPase prenyl moiety (Fig6C). Sequestration alone only

marginally affected the rate of dissociation, demonstrating that GDI actively extracts GTPases

from membranes (Fig6D,E).

To characterize membrane extraction more quantitatively, we carried out experiments

over a wide range of RhoGDI concentrations using either inactive (GDP-bound) or active (GTP-

bound, constitutively active) forms of Cdc42 or Rho. We observed that WT RhoGTPases

hydrolyze even GTP analogs such as GTPγS over the long time period (hours) required for

performing a full titration in our SLB assay (data not shown), which lead to them accumulating

in the inactive, GDP-bound form during the course of the experiment. We therefore turned to

two constitutively active GTPase variants -Q61L and G12V- and found that the former was the

most suitable for our in vitro assay because of its extremely low rate of spontaneous GTP

hydrolysis. Nonetheless, G12V produced qualitatively similar results in this assay (see SuppFig5

and Supplemental Discussion). Remarkably, GDI was able to extract both inactive (GDP-bound)

and active (GTP-bound, Q61L/Q63L) Cdc42 and RhoA in a concentration-dependent manner

(Fig7A,B). Although GDI extracted GDP-bound GTPase more efficiently than GTP-bound

GTPases (3-5 fold difference), it was still able to effectively facilitate the dissociation of the

latter (Figure 7C-F). The affinities of GDI for the active and inactive GTPases on membranes,

determined by hyperbolic fits to the extraction rates, were surprisingly similar (Cdc42:GDP

$3.41\pm0.56\mu M$, Cdc42Q61L:GTP $14.40\pm2.50\mu M$, RhoA:GDP $5.77\pm0.87\mu M$, RhoAQ63L:GTP

$19.58\pm2.45\mu M$). On the other hand, the maximal rates of extraction were not, indicating that the

rate-limiting step of membrane extraction depends on the activity state of GTPases. To

investigate whether extraction of active and inactive GTPases is a conserved ability among GDI

proteins, we also studied mammalian GDI. Similar to its *Xenopus* ortholog, bovine GDI1 was

able to extract both GDP- as well as GTP-bound Cdc42 and RhoA (SuppFig6). These data

clearly demonstrate that GDIs can catalytically extract both inactive and active RhoGTPase from

membranes *in vitro*.

Identification of an extraction-deficient RhoGDI

The canonical RhoGTPase cycle assumes that GDI does not extract GTPase without its

prior inactivation by a GAP (*reviewed by* Garcia-Mata, Boulter, & Burridge 2011). However, the

above results suggest that GDI might directly attenuate GTPase activity at the plasma membrane

via extraction of active GTPase. If this hypothesis is correct, then expression of an extraction-

deficient GDI would influence the spatiotemporal patterning of GTPase activity. We therefore

sought to generate an extraction-deficient GDI that could still initiate contact with GTPases but

fail to extract them from the plasma membrane. Mutants were screened by quantifying their

recruitment to wounds relative to WT GDI, based on the rationale that mutants capable of

binding but not extracting should remain at the membrane longer and thus accumulate at wounds

more than WT GDI.

Using this screen, we first tested RhoGDI mutants that were previously reported to be

deficient in extraction (Dransart, Morin, Cherfils, & Olofsson, 2005; Ueyama et al., 2013). None

of these extraction-deficient mutants were recruited to wounds more strongly than WT GDI,

suggesting that they were impaired in binding to GTPases at wounds rather than extraction

(SuppFig7). We therefore designed three novel mutants: (1) the isolated regulatory arm of GDI

that initiates contact with the GTPase but lacks a binding pocket for the hydrophobic prenyl

group (Δ51-199) and (2,3) mutation of residues E158/9 previously hypothesized to be

responsible for GTPase extraction (Hoffman, Nassar, & Cerione, 2000). GDI mutants Δ51-199,

E158/9A and E158/9Q were halo-tagged, expressed in oocytes and their recruitment to wounds

was quantified relative to WT GDI. GDI Δ51-199 showed minimal recruitment to wounds,

however both GDI E158/9A and E158/9Q showed a significant increase in recruitment to

wounds relative to WT GDI, with GDI E158/9Q (GDI-QQ) having the greatest increase

(Fig8A,B).

To directly test whether RhoGDI E158/9Q (QQ) is deficient in extraction, its functional

capabilities were tested *in vitro* in the SLB assay. WT GDI was able to extract both GDP-bound

and GTP-bound Cdc42 Q61L from the supported lipid bilayers (Fig8C). In contrast, GDI-QQ

retained most of its ability (less than two-fold reduction) to extract inactive Cdc42 but was

completely deficient in extracting active Cdc42 (Fig8C). Quantitatively similar results were

obtained using Cdc42 G12V (data not shown). The same was found to be true for Rho (Fig8C)

and to be conserved for bovine GDI1 (SuppFig8). These results confirm that the GDI-QQ mutant

is indeed extraction deficient: modestly so for inactive GTPases and completely so for active

GTPases.

RhoGDI extracts active RhoGTPase in vivo

We sought to directly test whether RhoGDI can extract active GTPases *in vivo* by

employing GDI-QQ. First, we compared the effects of WT vs. QQ GDI overexpression on

wounded oocytes expressing constitutively-active Cdc42 (G12V). (Q61L could not be used as it

failed to elevate Cdc42 levels around wounds; see SuppFig11 and Supplemental Discussion).

While WT GDI significantly reduced the amount of constitutively-active Cdc42 around wounds,

GDI-QQ did not (Fig9A,B). Second, we compared the effects of WT vs. QQ GDI overexpression

on wounded oocytes microinjected with Cy3-Cdc42 bound to GTPɣS. WT GDI significantly

reduced the amount of Cy3-Cdc42(GTPɣS) around wounds while GDI-QQ did not (Fig9C,D).

Collectively, these data suggest that GDI can extract active GTPase from the plasma membrane

in vivo.

The above results imply that the extraction of active RhoGTPase by GDI might

contribute to its spatiotemporal patterning *in vivo*. To test this hypothesis, we expressed GDI-QQ

and monitored the consequences on Rho and Cdc42 activity following wounding. Strikingly,

Cdc42 activity around wounds was significantly elevated, in contrast to Rho which was

unaffected (Fig9E,F). To assess whether the increase in activity was due to an increase of total

Cdc42 around wounds as opposed to competition between GDI-QQ and the Cdc42 activity

probe, we repeated the experiment with Cy3-Cdc42. Similar to the results obtained with the

activity reporter, expression of GDI-QQ elevated Cy3-Cdc42 levels around wounds relative to

controls (Fig9G,H). Cumulatively, these data suggest that GDI directly extracts active Cdc42

throughout the Cdc42 zone, and that extraction of active Cdc42 is necessary for its regulation

around wounds.

Discussion

Direct visualization of the RhoGTPases in living cells is essential for the understanding

of their complex spatiotemporal dynamics. We have established two methods to fluorescently

label vertebrate GTPases such that they are functional: internal tagging with a fluorescent protein

or by sortase-mediated labeling with a fluorescent dye. This now provides us with widely

applicable reagents to analyze GTPase function. These probes faithfully mimic the distribution

of the endogenous GTPases based on their comparison to activity reporters in several processes:

cell wound repair, cytokinesis, junctional integrity and epithelial wound repair. Further, the

successful rescue of Rho function at wounds in the presence of C3 by a C3-insensitive mutant of

IT-Rho indicates that these proteins are capable of replacing their endogenous counterparts. It

will be important to assess the ability of IT- or Cy3-labeled GTPases substitute for their

endogenous counterparts in other vertebrate cellular processes in the future, although we note

that IT-Cdc42 has been shown to be functional in fission yeast (Bendezú et al., 2015).

Nonetheless, the combination of the two labeling approaches is powerful as it permits side-by-

side comparison of results obtained *in vivo* and *in vitro,* as demonstrated here.

Visualization of functional, labeled RhoGTPases in combination with activity reporters in

living cells led to an unexpected observation: pools of inactive Cdc42 at the plasma membrane.

To the best of our knowledge, this is the first time that inactive GTPases have been detected at

membranes under conditions other than gross GTPase overexpression. Notably, the pool of

inactive Cdc42 spatially coincides with a local Cdc42-GAP, Abr (Vaughan et al. 2011). The pool

of inactive Cdc42 expands with overexpression of Abr, further demonstrating that locally-

inactivated Cdc42 can remain associated with the plasma membrane. This finding has important

mechanistic implications for the regulation of GTPase activity. It suggests that GTP hydrolysis

and extraction of GTPases, while are likely indirectly linked, are not necessarily tightly coupled.

This raises the possibility that GTPases might cycle through multiple rounds of activation and

inactivation while remaining associated with the membrane.

Remarkably, in addition to the RhoGTPases themselves, GDI also localized to the plasma

membrane in proximity to wounds. The localization of GDI in the same place where the

GTPases are especially abundant implies that its accumulation reflects interaction with is

GTPase clients. Consistent with this idea, GDI mutants deficient in GTPase binding fail to

recruit to wounds (data not shown). It will be important to investigate the detailed mechanism of

GDI localization and the control of its turnover at sites of high GTPase activity in the future.

The most significant result of this study is that RhoGDI extracts active GTPase,

particularly active Cdc42, during cell wound repair. This finding is based on both *in vitro* assays

showing that WT, but not GDI-QQ, extracts active GTPase from supported lipid bilayers and the

in vivo demonstration that WT, but not GDI-QQ, extracts constitutively-active and GTPɣS-

loaded Cdc42 from the plasma membrane. We thus conclude that GDI has the capacity to extract

active GTPases. Moreover, this ability is harnessed to limit the level of Cdc42 activity during

cell repair. While this finding may seem heretical, it has the virtue of explaining previous results

in the oocyte wound repair system. That is, based on an indirect approach involving

photoactivatable Rho and Cdc42 activity reporters, it was found that Cdc42 activity is lost

throughout its zone, while Rho activity is preferentially lost at the trailing edge of its zone

(Burkel, Benink, Vaughan, von Dassow, & Bement, 2012). The results presented here suggest

that GDI is responsible for the removal of active Cdc42 throughout the Cdc42 zone while Rho is

inactivated by a trailing edge GAP prior to extraction. This would also explain why a mild

overexpression of GDI significantly reduced Cdc42 activity but had no effect on Rho activity:

loss of active Cdc42 can be controlled at the level of GDI while Rho inactivation is controlled at

the level of a GAP (SuppFig9).

The broader implications of RhoGDI's ability to extract active GTPase are two-fold.

First, the canonical GTPase cycle needs a new branch in which active GTPase can be directly

extracted from the plasma membrane (Fig10). Further, because GDI binding strongly inhibits

GTP hydrolysis and nucleotide exchange (Hart et al., 1992; Ueda, Kikuchi, Ohga, Yamamoto, &

Takai, 2001), active GTPase may exist in its soluble form in complex with GDI. However,

complementary evidence from biochemical and biological studies suggest that active GTPases

are less stably bound to GDI compared to their inactive form (Hodgson et al., 2016; Slaughter,

Das, Schwartz, Rubinstein, & Li, 2009; Tnimov et al., 2012). As such, the secondary extraction

branch may actually represent a loop through which active GTPases are not only removed from

cellular membranes, but rapidly returned to them (Fig10). Such a mechanism might enhance the

spatial reach of GTPase activity within the plasma membrane or even mediate its spreading

between different membrane compartments (Palamidessi et al., 2008).

Second, RhoGDI's ability to extract active GTPase forces us to reassess its role in

GTPase regulation in different cellular processes. While the field primarily studies local GTPase

regulation at the level of GEFs and GAPs, we should reconsider GDI's role in regulation, as well

as the regulation of GDI itself.

Materials and Methods

Plasmids

The active RhoGTPase probes, mRFP-wGBD, eGFP-wGBD, eGFP-2xrGBD, BFP-

2xrGBD and mRFP-2xrGBD in pCS2+ were generated as previously described (Sokac et al.

2003; Benink and Bement 2005; Davenport et al. 2016). mCh-Rho, mCh-Rac, mCh-Cdc42

(Benink 2005), untagged Rho, Rac, and Cdc42 (wild-type (WT) and constitutively-active

(G12V)) in pCS2+ were made as previously described (Benink and Bement 2005). For

expression and purification in *E. coli*, codon-optimized Cdc42 and RhoA (Eurofins Genomics

Germany GmbH, Ebersberg, Deutschland) were subcloned into a pETMz2 vector via Gibson

assembly cloning (Gibson et al., 2009), with primers GTPase(GeneStrand)fwd and -rev and

pETfwd and -rev (all primer sequences in SuppTable1). A pentaglycine for sortase-mediated

labeling was added onto the 5' of the GTPases. The constitutively-active Cdc42 G12V and RhoA

G14V mutants were generated by Quickchange mutagenesis with primers Cdc42(G12V)fwd and

-rev and RhoA(G14V)fwd and -rev, respectively.

X. laevis IT-Cdc42 in pCS2+ was generated according to Bendezú et al. (2015): a linker -

SGGSACSGPPG- was cloned into Cdc42 after Q134. The linker encodes for *BamH1* and *Asc1*

restriction sites for digestion and insertion of GFP into the linker region. The 5' end of Cdc42

was amplified with primers Cdc42(1) and Cdc42(2); the 3' end was amplified separately with

primers Cdc42(3) and Cdc42(4). The two products were joined by PCR stitching with primers

Cdc42(1) and Cdc42(4). The single product was digested with *EcoR1* and *Xho1* and ligated into

pCS2+. The resulting construct was mutated by Quickchange with primers pCS2+-Cdc42(1) and

pCS2+-Cdc42(2) to remove the *BamH1* restriction site upstream of the insertion in the multiple

cloning site. eGFP was amplified from eGFP-wGBD with primers eGFP(1) and eGFP(2). Both

the Quickchanged construct and eGFP were digested with *BamH1* and *Asc1*, and eGFP was

ligated into the linker region internal to the Cdc42 coding sequence. *X. laevis* Rho and Rac were

similarly tagged internally after residues Q136 and L134, respectively.

X. laevis RhoGDI Clone ID:7010361 (GE-Healthcare Dharmacon, Lafayette, CO) was

subcloned into pCS2+ with *Cla1* and *Xho1*, and into N'3xGFP and N'Halo (Promega, Madison,

WI) pCS2+ with *BspE1* and *Xho1*. A FLAG-tag was added by PCR onto the 5' of RhoGDI, and

the product was subcloned into pFast-Bac1 with *Cla1* and *Not1*. The following mutations were

made by Quickchange mutagenesis to untagged and N'3xGFP RhoGDI in pCS2+: E158/9A,

E158/9Q, D40A, D40N, D180A and D180N (Dransart et al., 2005). Mutant RhoGDI 8(A) had

the first 8 charged amino acids to mutated to alanine (D3/5, E11-13, E15-17A) by sequential

PCR (Ueyama et al., 2013). The 3' end of RhoGDI was amplified with primers 8(A)F1 and R1.

The product was amplified and added to at its 5' end with primers 8(A)F2 and R1, and for a third

time with 8(A)F3 and R1. The final PCR product was subcloned into pCS2+ by Infusion PCR

(Takara Bio, Kusatsu, Japan). For subcloning into N'3xGFP-pCS2+, the third PCR from above

was repeated with 8(A)F4 and R2. Mutant RhoGDI helix replacement (HR) had alpha helix D39-

Q48 replaced with a glycine linker GGGGSGGGGS. This was done by sequential PCR as

described above with four rounds of PCR: HR1 and R1, HR2 and R1, HR3 and R1, then either

HR4 and R1 for subcloning into pCS2+ or HR5 and R2 for subcloning into N'3xGFP-pCS2+.

RhoGDI mutant Δ51-199 was generated by adding a stop codon after L50 by Quickchange

mutagenesis. To make RhoGDI Δ1-19, the 3' end of RhoGDI was amplified with primers (-)20

F1 and R for subcloning by infusion into pCS2+, and primers (-)20 F2 and R for N'3xGFP-

pCS2+. Primers (-55) F1, F2 and R were used to generate RhoGDI Δ1-54 as described above

(Hoffman et al., 2000; Ueyama et al., 2013). For expression and purification in *E. coli*, *X. laevis*

RhoGDI WT and E158/9Q were subcloned into a pGEX-6P-2 vector via Gibson assembly

cloning (Gibson et al., 2009) using XlRhoGDIfwd and -rev. A cysteine for labeling was added

by PCR onto the 5' of RhoGDI. Bovine RhoGDI1 in pGEX-6P was a gift of Dr. Tomotaka

Komori. E163/4Q mutation was made by Quickchange mutagenesis with the E163/4Qfwd and -

rev primers. Mutant bovine RhoGDI Δ1-22 and Δ1-59 were subcloned with BamHI and NotI

into a pGEX-6P-2 with Δ1-22fwd and -rev and Δ1-59fwd and -rev, respectively. Mutant bovine

RhoGDI HR was generated via Gibson assembly cloning (Gibson et al., 2009), with primers

HRfwd and -rev and pGEXHRfwd and -rev, for amplification of RhoGDI and the

pFASTBacH10 vector, respectively. RabGGTase 2 beta in a pGATEV vector was kindly

provided by Dr. Konstantin Gavriljuk.

Expression and purification of recombinant protein from E. coli for sortase-labeling

Rosetta(DE3) chemically competent *E. coli* cells were transformed with WT or mutant

RhoGDIs, induced with 250µM IPTG and incubated at 18°C ON. Bacteria cells were harvested,

centrifuged at 4000xg for 20min, and pellets flash frozen in liquid nitrogen and stored at -80°C.

Frozen pellets were resuspended in a 3x volume of lysis buffer (50mM KPi pH 8, 400mM KCl,

1mM EDTA, 5mM βME, 1mM PMSF, 1mM benzamidine) and lysed with a high pressure

homogenizer at 4°C. Lysate was clarified by centrifugation at 100,000xg for 1hr and applied to a

glutathione sepharose 4 fast flow column bed (GE-Healthcare, Chicago, IL) equilibrated with

wash buffer (50mM KPi pH 8, 400mM KCl, 1mM EDTA, 5mM BME, 1mM benzamidine). The

column was washed with wash buffer, and protein was eluted with elution buffer (50mM KPi pH

8, 400mM KCl, 1mM EDTA, 5mM BME, 1mM benzamidine, 10mM reduced L-glutathione).

Peak fractions were pooled, protein concentration was estimated with Bradford assay (Bio-Rad

Laboratories, Inc., Hercules, CA), and PreScission protease was added at 1:30. After ON

incubation on ice, the sample was concentrated using 5,000 MWCO Vivaspin15R centrifugal

concentrators (Sartorius AG, Göttingen, Germany), buffer exchanged in wash buffer on a HiPrep

26/10 desalting column (GE-Healthcare) and recirculated on the same glutathione sepharose 4

fast flow column bed re-equilibrated in wash buffer. Flow through was collected, concentrated,

spun down and gel filtered on a HiLoad Superdex 75 pg column (GE-Healthcare) in storage

buffer (20mM HEPES pH 7.5, 150mM KCl, 0.5mM TCEP, 20% Glycerol). Peak fractions were

pooled, concentrated, flash frozen in liquid nitrogen and stored at -80°C. Protein purification and

purity were determined by Coomassie stain of 12% SDS-PAGE, protein concentration measuring

absorbance at 280nm. WT and mutant GTPases were expressed and purified similarly to GDIs

with the following differences: L21(DE3) chemically competent *E. coli* cells were used and

proteins were expressed with 1 mM IPTG at 37°C for 4hrs. The affinity step was performed on

HiTrap Chelating HP columns loaded with cobalt and equilibrated in 50mM HEPES pH 7.5,

50mM NaCl, 5mM MgCl2, 0.5mM BME, 100µM ATP and 100µM GDP/GTP. RabGTTase Beta

was expressed and purified as described before (Gavriljuk, Itzen, Goody, Gerwert, & Kötting,

2013). After gel filtration, nucleotide bound to Cdc42 G12V was exchange to GTPγS, incubating

the protein with 10-fold excess EDTA and GTPγS for 30 minutes on ice. The new nucleotide

state was stabilized by adding 20-fold excess MgCl$_2$. Before freezing, protein buffer was

exchanged with a NAP-5 column (GE-Healthcare) in storage buffer (50mM HEPES pH 7.5,

50mM NaCl, 2mM MgCl2, 2mM DTT, 20% Glycerol).

Cy3-labeling and in vitro prenylation of RhoGTPases

RhoGTPases were labeled at the N-t with Cy3 using a sortase-mediated reaction and *in*

vitro prenylated as previously described (Gavriljuk et al., 2013; Popp, Antos, Grotenbreg,

Spooner, & Ploegh, 2007). In brief, RhoGTPases were incubated with sortase and Cy3 N-t

labeled LPETGG peptide at 3:1:15 ratio in labeling buffer (Tris pH 8.0, 150mM KCl, 6µM

CaCl2, 0.5mM TCEP) and incubated ON at 16°C. The entire reaction was mixed with

geranylgeranyltransferase type 1 and geranylgeranyl diphosphate at 10:1:30 ratio in prenylation

buffer (50mM HEPES pH 7.5, 50mM NaCl, 2mM $MgCl_2$, 2mM DTT, 30µM GDP/GTP, 2%

CHAPS), and incubated ON on a rotating mixer at 4°C. The sample was spun in a TLA-100

rotor (Beckman Coulter, Brea, CA) at 80,000 rpm for 30 minutes at 4°C and gel filtered on a

HiLoad Superdex 75pg column (GE-Healthcare) equilibrated with prenylation buffer with 0.5%

CHAPS. Peak fractions were pooled, concentrated using 5,000 MWCO Vivaspin4 centrifugal

concentrators (Sartorius AG) and buffer exchanged in prenylation buffer without CHAPS on a

NAP-5 column (GE-Healthcare). Residual detergent was removed by Pierce™ Detergent

Removal Spin Column (Thermo Fisher). After sortase-mediated labeling with Cy3, unprenylated

proteins were directly spun down and gel filtered in absence of CHAPS.

Oocyte collection and preparation

Ovarian tissue was harvested from adult *X. laevis* via surgical procedures approved by the

University of Wisconsin-Madison Institutional Animal Care and Use Committee. Oocytes were

stored in 1x modified Barth's solution (88mM NaCl, 1mM KCl, 2.4mM $NaHCO_3$, 0.82mM

$MgSO_4$, 0.33mM $NaNO_3$, 0.41mM $CaCl_2$, 10mM HEPES, pH 7.4) with 100µg/mL gentamicin

sulfate, 6µg/mL tetracycline and 25µg/mL ampicillin at 16°C. Prior to manual defolliculation

with forceps, oocytes were treated with 8mg/mL type I collagenase (Life Technologies, Grand

Island, NY) in 1x modified Barth's solution for 1hr at 16°C on an orbital shaker.

mRNA preparation

mRNA was generated *in vitro* using the mMessage mMachine SP6 transcription kit

(Thermo Fisher, Carlsbad, CA) and purified using the RNeasy Mini Kit (Qiagen, Hilden,

Germany). Transcript size was verified on a 1% agarose/formaldehyde denaturing gel relative to

the Millennium Marker (Life Technologies) RNA molecular weight standard.

Oocyte microinjection

Oocytes were microinjected with a 40nL injection volume using a p-100 microinjector

(Harvard Apparatus, Holliston, MA). mRNA encoding probes for active Rho (2xrGBD) and

active Cdc42 (wGBD) were injected at a final needle concentration of 30µg/mL and 100µg/mL,

respectively. IT-Rho, Rac and Cdc42 were each injected at a final needle concentration of

125µg/mL, with 63µg/mL of WT RhoGDI to stabilize the exogenous GTPase and maintain

stoichiometric ratio of GTPase:GDI (Boulter et al., 2010). mRNA encoding Abr was injected at a

final needle concentration of 25-500µg/mL. 3xGFP-WT GDI and mutants at 333µg/mL,

untagged RhoGDI at 300µg/mL, Halo-WT GDI and mutants at 200µg/mL, Cdc42 G12V at

28µg/mL, and untagged GDI E158/9Q at 1.5mg/mL. For purified protein, Cy3-Rho and Cy3-

Cdc42, *in vitro* prenylated and complexed with RhoGDI, were injected at a final needle

concentration of 4.56µM. C3 exotransferase was injected at a final needle concentration of

497 1.1µg/mL in 1mM DTT and WT GDI at 3.5-114µM for the standard curve. For wounding

498 experiments, all mRNA was injected 20-24hrs before imaging, and purified protein was injected

499 at least 2hrs before imaging, except for C3 which was injected 30min prior to imaging. For

500 imaging cortical granule exocytosis, oocytes were injected 16hrs before imaging and matured

501 ON in progesterone. Two-cell embryos were microinjected with a 5nL injection volume at a final

502 needle concentration of 167µg/mL for IT-Rho and IT-Cdc42 mRNA, and 18.24µM Cy3-Rho and

503 Cy3-Cdc42.

505 *Purification of recombinant protein from E. coli for antibody purification*

506  BL21 pLysS cells (Thermo Fisher) were transformed with GST-RhoGDI in pGEX6p.1.

507 A positive clone was used to inoculate 12mL of lysogeny broth (LB) supplemented with

508 25µg/mL ampicillin and cultured ON. The 12mL culture was added to 1L of LB with ampicillin

509 and shaken at 37℃ until OD600~0.6. The culture was induced by adding a final concentration of

510 0.1mM IPTG and shaken at 37℃ for 2hrs. BL21 pLysS cells were pelleted at 5300rpm for 10min

511 at 4℃, and the pellet resuspended in Buffer A (50mM Tris-HCL, pH 7.6; 50mM NaCl with 1mM

512 DTT in PBS). Pellets were stored at -80℃. Pellets were thawed at room temperature to promote

513 cell lysis. Triton X-100 was added to a final concentration of 0.6%, PMSF at 500uM, lysoszyme

514 at 1mM in 10mM Tris pH 8.0, 400µM Peflabloc, 1µg/mL aprotinin, 1µg/mL leupeptin.

515 Solubilate was incubated at RT for 30min, DNAse1 was added to a final concentration of

516 10ug/mL, incubated again for at RT for 30min, and centrifuged at 16,000xg for 10min at 4℃. The

517 supernatant was collected and exposed to a column containing glutathione-sepharaose 4B

518 (MilliporeSigma). The column was washed 5x with Buffer A and the protein eluted with 20mM

519 Tris, pH 8.0, 20mM glutathione, 400µM Peflabloc, 1.25µg/mL aprotinin, 14.25µg/mL leupeptin,

520   0.25mM E-64, 0.5mM PMSF. Protein concentration was determined by Coomassie stain of a

521   12% SDS-PAGE alongside a BSA standard curve.

Antibody generation and purification

FLAG-RhoGDI purified from Sf9 cells was used as an antigen for antibody production in

rabbits (Covance, Princeton, NJ). The serum was heat-inactivated at 56°C for 30min, diluted 1:1

in 20mM Tris, pH 7.5, and filtered through a 0.22µm syringe. The diluted, filtered serum was

loaded onto a column containing GST-GDI coupled to Affi-Gel 15, to minimize antibody cross-

reactivity to the FLAG-tag on the antigen. The column was washed 20x with 20mM Tris, pH 7.5

and 20x with 20mM Tris, pH 7.5, 500mM NaCl. Antibody was first eluted with 100mM glycine,

pH 2.5 into 1M Tris, pH 8.8 for neutralization. The column was washed 20x with 20mM Tris,

pH 8.8. Antibody remaining on the column was eluted with 100mM Triethylamine, pH 11.5 into

concentrated HCl and 1M Tris, pH 7.5 for neutralization. The concentration of each fraction was

determined by A_{280}. The peak antibody fractions were pooled, dialyzed against PBS (2x2L) ON,

and concentrated using a 100K MW Amicon Ultra-15 Centrifugal filter (MilliporeSigma).

Antibody specificity was determined by western blotting of purified protein, *X. laevis* oocyte

whole cell lysate (WCL), WCL of oocytes overexpressing GDI and WCL of oocytes expressing

3xGFP-GDI.

Expression and purification of recombinant protein from insect cells

DH10Bac-competent *E. coli* (Thermo Fisher) were transformed with FLAG-WT RhoGDI

or E158/9Q in pFast-Bac1 and positive clones were identified by blue/white screening. Bacmid

was purified and transfected into Sf9 cells using Cellfectin II reagent (Thermo Fisher). High-

expressing clones were identified and baculovirus was generated for two additional passages. Sf9

cells, 22×10^6 per 15cm plate, were infected with high-titer baculovirus and incubated 27°C for

72hrs. Sf9 cells were harvested, centrifuged at 500xg for 5min, and pellets were stored at -80°C.

Frozen pellets were resuspended in a 5x volume of solubilization buffer (1xPBS pH 7.5,

1% Triton X-100, 0.5µg/mL leupeptin, 0.5µg/mL aprotinin, 0.5µg/mL Pepstatin A, 40µg/mL

PMSF, 100µg/mL benzamidine, 0.5µg/mL E64) and incubated at 4°C with end-over-mixing for

1hr. Lysate was clarified by centrifugation at 21,000xg for 15min and applied to an anti-FLAG

M2 agarose column bed (MilliporeSigma, Burlington, MA). The column was washed 3x with

wash buffer (1xPBS, 0.5µg/mL leupeptin, 0.5µg/mL aprotinin, 0.5µg/mL Pepstatin A, 40µg/mL

PMSF, 100µg/mL benzamidine, 0.5µg/mL E64). A buffer exchange was performed with 1:1

wash buffer:HEPES (25mM HEPES pH 7.5, 100mM KCl), and the column washed 2x with

HEPES. Protein was eluted with 1M Arginine pH 4.4 into an equal volume of collection buffer

(50mM HEPES pH 7.5, 200mM KCl). Fractions were analyzed by coomassie stain of a 12%

SDS-PAGE. Peak fractions were pooled and concentrated using a 10MW Amicon Ultra-15

Centrifugal filter (MilliporeSigma). A buffer exchange was performed during concentrating with

HEPES such that the final Arginine concentration was less than 1mM. Protein purification and

purity was determined by comparison to a BSA standard curve by Coomassie stain of a 12%

SDS-PAGE.

Fixing and staining of wounded oocytes

Oocytes were wounded, allowed to heal for 2-3min and fixed for 2hrs in 10mM EGTA,

100mM KCl, 3mM MgCl$_2$, 10mM HEPES, 150mM sucrose (pH 7.6), 4% PFA, 0.1%

glutaraldehyde, 0.1% Triton X-100. Fixed oocytes were washed 5x in TBSN/BSA (5mg/mL

BSA in 1xTBS containing 0.1% NP-40). Oocytes were bisected and blocked in TBSN/BSA for

4hrs at 4°C. Oocytes were stained with rabbit α-RhoGDI at 1:1000 in TBSN/BSA for 12hrs,

washed 5x in TBSN/BSA over 12hrs, stained with chicken α-rabbit Alexa Fluor 647 (Invitrogen,

Carlesbad, CA) at 1:10,000 for 12hrs in BSN/BSA at 4°C, and washed 5x in TBSN/BSA over

12hrs.

Image acquisition, wounding and data analysis

Laser scanning confocal microscopy was performed using a Nikon Eclipse Ti inverted

microscope with a Prairie Point Scanner confocal system (Bruker, Middleton, WI). The

microscope was fitted with a 440-nm dye laser pumped by a MicroPoint 337-nm nitrogen laser

(Andor, South Windsor, CT) for wounding. Brightest-point projections, measurements of

fluorescence intensities, area and distances were made in FIJI (Schindelin et al., 2012). Bio-

Formats Importer and De-Flicker plugins were used. Ring intensity corrected for background

was calculated by quantifying the mean intensity of the ring and subtracting the mean intensity

of the background. Total activity was calculated by multiplying the mean intensity of the zone

(corrected for background) by the area of the zone, normalized for wound width. GraphPad

Prism was used to plot quantifications and perform statistical analyses. An unpaired student's T-

test with a 2-tailed distribution and unequal variance was used to compare two conditions, one-

way ANOVA with a Tukey post hoc analysis was used to analyze more than two conditions.

*$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

Supported lipid bilayer assay

Small unilamellar vesicles and supported lipid bilayers were prepared with 100% 1,2-

dioleoyl-sn-glycero-3-phosphocholine (18:1 DOPC; Avanti Polar Lipids, Inc., Alabaster, AL) as

described before (Hansen et al., 2019). 250µL Cy3 labeled GTPases were incubated on SLBs at

200nM final concentration until equilibrium was reached. A 0.5mm silicone tubing was attached

drop-to-drop to the chamber with the equilibrated sample via a male luer connector (ibidi GmbH,

Gräfelfing, Germany). After acquisition of few frames in absence of flow, the chamber was

flushed at 10µL/sec with imaging buffer (20mM HEPES pH 7.0, 150mM KCl, 1.5mM $MgCl_2$,

0.5mM EGTA, 100µM GDP/GTP) alone or in presence of a GTPase solubilizer until baseline

was reached. Oxygen scavenger system (1.25mg/mL glucose oxidase, 0.2 mg/mL catalase, 400

mg/mL glucose) was added fresh to each sample and buffer before imaging. TIRF was

performed on a Nikon Eclipse Ti inverted microscope with a VisiScope TIRF-FRAP Cell

Explorer system (Visitron Systems GmbH, Puchheim, Germany) using a 60× Apo TIRF oil-

immersion objective (1.49 N.A.). Cy3-labeled proteins were excited with a 561-nm laser line,

excitation light was passed through a ET-561nm Laser Bandpass Set (Chroma Technology

Corporation, Bellows Falls, VT) before illuminating the sample. Fluorescence emission was

detected on a Evolve 512 Delta EMCCD camera (Teledyne Photometrics, Tucson, AZ).

Measurements of fluorescence intensities were made in FIJI (Schindelin et al., 2012). Plot Z-axis

profile tool and Bio-Formats Importer plugins were used. Fluorescence intensity was corrected

for background. To display multiple curves on the same graph, data from different experiments

were aligned using the overshoot signal occurring after the flow was started and normalized

dividing by the maximum intensity. Data from wash off experiments were fitted with a one

component exponential decay function ($y = y_0 + A\,e^{-\lambda x}$; y_0= y offset, A=amplitude,

λ=exponential decay constant), choosing a fitting range that did not include the initial overshoot.

This was possible because a monoexponential function can be fitted to a range of the data set

without affecting the λ value obtained. K_{off} titration curves were fitted with a hyperbolic function

($y = y_0 + \frac{\lambda_{max}\,x}{K_d + x}$). Origin Pro (OriginLab Corporation, Northampton, MA) was used to analyze

data, plot quantifications and perform statistical analyses. An unpaired student's T-test with a 2-

tailed distribution and equal variance was used to compare two conditions. **$p<0.01$,

***$p<0.001$, ****$p<0.0001$.

Acknowledgements

This work was supported by National Institutes of Health Grant GM52932 to W.B., a Dr. Stanley and Dr. Eva Lurie Weinreb Fellowship to A.G. and HSFP CDA00070/2017-2 to P.B. I.V is supported by the MaxSynBio Consortium, which is jointly funded by the Federal Ministry of Education and Research of Germany and the Max Planck Society. We acknowledge National Institutes Health Grant R44 MH065724 to LOCI at UW-Madison. We are also grateful to both our labs for their continued input.

References

Abo, A., Webb, M., Grogan, A., & Segal, A. (1994). Activation of NADPH oxidase involves the
dissociation of p21rac from its inhibitory GDP/GTP exchange protein (rhoGDI) followed
by its translocation to the plasma membrane. *The Biochemical Journal*, *298 Pt 3*(3), 585–
591. https://doi.org/10.1042/bj2980585

Abreu-Blanco, M. T., Verboon, J. M., & Parkhurst, S. M. (2014). Coordination of Rho Family
GTPase Activities to Orchestrate Cytoskeleton Responses during Cell Wound Repair.
Current Biology, *24*(2), 144–155. https://doi.org/10.1016/j.cub.2013.11.048

Bement, W. M., Benink, H. A., & Von Dassow, G. (2005). A microtubule-dependent zone of
active RhoA during cleavage plane specification. *Journal of Cell Biology*, *170*(1), 91–101.
https://doi.org/10.1083/jcb.200501131

Bendezú, F. O., Vincenzetti, V., Vavylonis, D., Wyss, R., Vogel, H., & Martin, S. G. (2015).
Spontaneous Cdc42 Polarization Independent of GDI-Mediated Extraction and Actin-Based
Trafficking. *PLoS Biology*, *13*(4), 1–30. https://doi.org/10.1371/journal.pbio.1002097

Benink, H. (2005). *Characterization of Rho GTPases during wound healing - Catalog - UW-*
Madison Libraries. Retrieved from
https://search.library.wisc.edu/catalog/9910091280802121

Benink, H. A., & Bement, W. M. (2005). Concentric zones of active RhoA and Cdc42 around
single cell wounds. *Journal of Cell Biology*, *168*(3), 429–439.
https://doi.org/10.1083/jcb.200411109

Benink, H., & Bement, W. (2001). Concentric rings of Rho and Rac/Cdc42 activity encircle
oocyte wounds - UW Madison. *Molecular Biology of the Cell*, *12*, 396A. Retrieved from
http://uwi-primoalma-
prod.hosted.exlibrisgroup.com/primo_library/libweb/action/openurl?sid=google&auinit=H
A&aulast=Benink&atitle=Concentric+rings+of+Rho+and+Rac/Cdc42+activity+encircle+o
ocyte+wounds&title=Molecular+biology+of+the+cell+/&volume=12&date=200

Bishop, A. L., & Hall, A. (2000). *Rho GTPases and their effector proteins*. *Biochem. J* (Vol.
348). Retrieved from
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1221060/pdf/10816416.pdf

Boulter, E., Garcia-Mata, R., Guilluy, C., Dubash, A., Rossi, G., Brennwald, P. J., & Burridge,
K. (2010). Regulation of Rho GTPase crosstalk, degradation and activity by RhoGDI1.
Nature Cell Biology, *12*(5), 477–483. https://doi.org/10.1038/ncb2049

Burkel, B. M., Benink, H. A., Vaughan, E. M., von Dassow, G., & Bement, W. M. (2012). A
Rho GTPase Signal Treadmill Backs a Contractile Array. *Developmental Cell*, *23*(2), 384–
396. https://doi.org/10.1016/j.devcel.2012.05.025

Clark, A. G., Miller, A. L., Vaughan, E., Yu, H. Y. E., Penkert, R., & Bement, W. M. (2009).
Integration of Single and Multicellular Wound Responses. *Current Biology*, *19*(16), 1389–
1395. https://doi.org/10.1016/j.cub.2009.06.044

Davenport, N. R. (2016). *Molecular and Cellular Determinants of Pattern Formation During*

Wound Repair in Xenopus laevis. University of Wisconsin-Madison.

Davenport, N. R., Sonnemann, K. J., Eliceiri, K. W., & Bement, W. M. (2016). Membrane
dynamics during cellular wound repair. *Molecular Biology of the Cell*, *27*(14), 2272–2285.
https://doi.org/10.1091/mbc.E16-04-0223

Dransart, E., Morin, A., Cherfils, J., & Olofsson, B. (2005). Uncoupling of inhibitory and
shuttling functions of Rho GDP dissociation inhibitors. *Journal of Biological Chemistry*,
280(6), 4674–4683. https://doi.org/10.1074/jbc.M409741200

Garcia-Mata, R., Boulter, E., & Burridge, K. (2011). The "invisible hand": regulation of RHO
GTPases by RHOGDIs. *Nature Reviews Molecular Cell Biology*, *12*(8), 493–504.
https://doi.org/10.1038/nrm3153

Gavriljuk, K., Itzen, A., Goody, R. S., Gerwert, K., & Kötting, C. (2013). Membrane extraction
of Rab proteins by GDP dissociation inhibitor characterized using attenuated total reflection
infrared spectroscopy. *Proceedings of the National Academy of Sciences of the United
States of America*, *110*(33), 13380–13385. https://doi.org/10.1073/pnas.1307655110

Gibson, D. G., Young, L., Chuang, R.-Y., Venter, J. C., Hutchison, C. A., & Smith, H. O.
(2009). Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nature
Methods*, *6*(5), 343–345. https://doi.org/10.1038/nmeth.1318

Hancock, J. F., & Hall, A. (1993). A novel role for RhoGDI as an inhibitor of GAP proteins. *The
EMBO Journal*, *1*(1), 91–95.

Hansen, S. D., Huang, W. Y. C., Lee, Y. K., Bieling, P., Christensen, S. M., & Groves, J. T.
(2019). Stochastic geometry sensing and polarization in a lipid kinase-phosphatase
competitive reaction. *Proceedings of the National Academy of Sciences of the United States
of America*, 201901744. https://doi.org/10.1073/pnas.1901744116

Hart, M. J., Maru, Y., Leonard, D., Witte, O. N., Evans, T., & Cerione, R. A. (1992). A GDP
dissociation inhibitor that serves as a GTPase inhibitor for the Ras-like protein CDC42Hs.
Science, *258*(5083), 812–815. https://doi.org/10.1126/science.1439791

Hodgson, L., Spiering, D., Sabouri-Ghomi, M., Dagliyan, O., DerMardirossian, C., Danuser, G.,
& Hahn, K. M. (2016). FRET binding antenna reports spatiotemporal dynamics of GDI-
Cdc42 GTPase interactions. *Nature Chemical Biology*, *12*(10), 802–809.
https://doi.org/10.1038/nchembio.2145

Hoffman, G. R., Nassar, N., & Cerione, R. A. (2000). Structure of the Rho family GTP-binding
protein Cdc42 in complex with the multifunctional regulator RhoGDI. *Cell*, *100*(3), 345–
356. https://doi.org/10.1016/S0092-8674(00)80670-4

Howell, A. S., Jin, M., Wu, C.-F., Zyla, T. R., Elston, T. C., & Lew, D. J. (2012). Negative
feedback enhances robustness in the yeast polarity establishment circuit. *Cell*, *149*(2), 322–
333. https://doi.org/10.1016/j.cell.2012.03.012

Johnson, J. L., Erickson, J. W., & Cerione, R. A. (2009). New Insights into How the Rho
Guanine Nucleotide Dissociation Inhibitor Regulates the Interaction of Cdc42 with
Membranes. *Journal of Biological Chemistry*, *284*(35), 23860–23871.
https://doi.org/10.1074/jbc.M109.031815

Kimura, K., Ito, M., Amano, M., Chihara, K., Fukata, Y., Nakafuku, M., … Kaibuchi, K. (1996). Regulation of myosin phosphatase by Rho and Rho-associated kinase (Rho-kinase). *Science (New York, N.Y.)*, *273*(5272), 245–248. https://doi.org/10.1126/SCIENCE.273.5272.245

Machacek, M., Hodgson, L., Welch, C., Elliott, H., Pertz, O., Nalbant, P., … Danuser, G. (2009). Coordination of Rho GTPase activities during cell protrusion. *Nature*, *461*(7260), 99–103. https://doi.org/10.1038/nature08242

Martin, K., Reimann, A., Fritz, R. D., Ryu, H., Jeon, N. L., & Pertz, O. (2016). Spatio-temporal co-ordination of RhoA, Rac1 and Cdc42 activation during prototypical edge protrusion and retraction dynamics. *Scientific Reports*, *6*(1), 21901. https://doi.org/10.1038/srep21901

Moon, S. Y., & Zheng, Y. (2003). Rho GTPase-activating proteins in cell regulation. *Trends in Cell Biology*, *13*(1), 13–22. https://doi.org/10.1016/S0962-8924(02)00004-1

Ngo, A. T. P., Thierheimer, M. L. D., Babur, Ö., Rocheleau, A. D., Huang, T., Pang, J., … Aslan, J. E. (2017). Assessment of roles for the Rho-specific guanine nucleotide dissociation inhibitor Ly-GDI in platelet function: a spatial systems approach. *American Journal of Physiology - Cell Physiology*, *312*(4), C527–C536. https://doi.org/10.1152/ajpcell.00274.2016

Nomanbhoy, T., Erickson, J., & Cerione, R. (1999). Kinetics of Cdc42 Membrane Extraction by Rho-GDI Monitored by Real-Time Fluorescence Resonance Energy Transfer. *Biochemistry*, *38*(6), 1744–1750. https://doi.org/10.1021/BI982198U

Nomanbhoy, T. K., & Cerione, R. A. (1996). *Characterization of the Interaction between RhoGDI and Cdc42Hs Using Fluorescence Spectroscopy**. Retrieved from http://www.jbc.org/

Palamidessi, A., Frittoli, E., Garré, M., Faretta, M., Mione, M., Testa, I., … Di Fiore, P. P. (2008). Endocytic Trafficking of Rac Is Required for the Spatial Restriction of Signaling in Cell Migration. *Cell*, *134*(1), 135–147. https://doi.org/10.1016/J.CELL.2008.05.034

Popp, M. W., Antos, J. M., Grotenbreg, G. M., Spooner, E., & Ploegh, H. L. (2007). Sortagging: a versatile method for protein labeling. *Nature Chemical Biology*, *3*(11), 707–708. https://doi.org/10.1038/nchembio.2007.31

Rossman, K., Der, C., & Sondek, J. (2005). GEF means go: turning on RHO GTPases with guanine nucleotide-exchange factors. https://doi.org/10.1038/nrm1587

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., … Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Sekine, A., Fujiwara, M., & Narumiyas, S. (1989). Asparagine Residue in the rho Gene Product Is the Modification Site for Botulinum ADP-ribosyltransferase. *The Journal of Biological Chemistry*, *264*(15), 8602–8605. Retrieved from http://www.jbc.org/content/264/15/8602.full.pdf

Slaughter, B. D., Das, A., Schwartz, J. W., Rubinstein, B., & Li, R. (2009). Dual Modes of Cdc42 Recycling Fine-Tune Polarized Morphogenesis. *Developmental Cell*, *17*(6), 823–835. https://doi.org/10.1016/J.DEVCEL.2009.10.022

Smith, M. J., Neel, B. G., & Ikura, M. (2013). NMR-based functional profiling of RASopathies
and oncogenic RAS mutations. *Proceedings of the National Academy of Sciences*, *110*(12),
4574–4579. https://doi.org/10.1073/pnas.1218173110

Sokac, A. M., Co, C., Taunton, J., & Bement, W. (2003). Cdc42-dependent actin polymerization
during compensatory endocytosis in Xenopus eggs. *Nature Cell Biology*, *5*(8), 727–732.
https://doi.org/10.1038/ncb1025

Tnimov, Z., Guo, Z., Gambin, Y., Nguyen, U. T. T. T., Wu, Y.-W. W., Abankwa, D., …
Alexandrov, K. (2012). Quantitative analysis of prenylated RhoA interaction with its
chaperone, RhoGDI. *Journal of Biological Chemistry*, *287*(32), 26549–26562.
https://doi.org/10.1074/jbc.M112.371294

Ueda, T., Kikuchi, A., Ohga, N., Yamamoto, J., & Takai, Y. (2001). Purification and
Characterization from Bovine Brain Cytosol of a Novel Regulatory Protein Inhibiting the
Dissociation of GDP from and the Subsequent Binding of GTP to rhoB ~20, a ras p214ike
GTP- binding Protein*. *The Journal of Biological Chemistry*, *265*(16), 1–9. Retrieved from
http://www.jbc.org/content/265/16/9373.full.pdf%0Apapers3://publication/uuid/E9334306-
D989-40B8-A264-39306D63F2D4

Ueyama, T., Son, J., Kobayashi, T., Hamada, T., Nakamura, T., Sakaguchi, H., … Saito, N.
(2013). Negative Charges in the Flexible N-Terminal Domain of Rho GDP-Dissociation
Inhibitors (RhoGDIs) Regulate the Targeting of the RhoGDI-Rac1 Complex to Membranes.
The Journal of Immunology, *191*(5), 2560–2569. https://doi.org/10.4049/jimmunol.1300209

Vaughan, E. M., Miller, A. L., Yu, H. Y. E., & Bement, W. M. (2011). Control of local Rho
GTPase crosstalk by Abr. *Current Biology*, *21*(4), 270–277.
https://doi.org/10.1016/j.cub.2011.01.014

Wühr, M., Freeman, R. M., Presler, M., Horb, M. E., Peshkin, L., Gygi, S. P., & Kirschner, M.
W. (2014). Deep Proteomics of the Xenopus laevis Egg using an mRNA-Derived Reference
Database. *Current Biology*, *24*(13), 1467–1475. https://doi.org/10.1016/J.CUB.2014.05.044

Yonemura, S., Hirao-Minakuchi, K., & Nishimura, Y. (2004). Rho localization in cells and
tissues. *Experimental Cell Research*, *295*(2), 300–314.
https://doi.org/10.1016/J.YEXCR.2004.01.005

Yu, H. Y. E., & Bement, W. M. (2007). Control of local actin assembly by membrane fusion-
dependent compartment mixing. *Nature Cell Biology*, *9*(2), 149–159.
https://doi.org/10.1038/ncb1527

Yüce, Ö., Piekny, A., & Glotzer, M. (2005). An ECT2-centralspindlin complex regulates the
localization and function of RhoA. *The Journal of Cell Biology*, *170*(4), 571–582.
https://doi.org/10.1083/jcb.200501097

Zhang, S.-C., Gremer, L., Heise, H., Janning, P., Shymanets, A., Cirstea, I. C., … Ahmadian, M.
R. (2014). Liposome Reconstitution and Modulation of Recombinant Prenylated Human
Rac1 by GEFs, GDI1 and Pak1. *PLoS ONE*, *9*(7), e102425.
https://doi.org/10.1371/journal.pone.0102425

Figure 1: Direct visualization of Rho and Cdc42 during cell wound repair. A) Left: image of

active Cdc42 (magenta) and active Rho (green) around a single-cell wound; right: schematic

diagram indicating zone regions; B) Wound in oocyte microinjected with rGBD (magenta) and

IT-Rho (green); B') Line scan of normalized fluorescence intensity from (B); C) As in B but

with wGBD (magenta) and IT-Cdc42 (green); D,D') As in B but with Cy3-Rho (magenta) and

rGBD (green); E,E') As in B but with Cy3-Cdc42 (magenta) and wGBD (green); F,F') As in B

but with Cy3-Rho (magenta) and IT-Rho (green); G,G') As in B but with Cy3-Cdc42 (magenta)

and IT-Cdc42 (green) and line scan. Scale bar 10μm, time min:sec.

Figure 2: Directly-labeled Rho and Cdc42 during cytokinesis, cortical granule exocytosis,

epithelial wound repair and at junctions. A) Cytokinesis in *X. laevis* embryo microinjected

with IT-Rho; IT-Rho accumulates at nascent cleavage furrow (arrowhead); B) Cytokinesis in *X.*

laevis embryo microinjected with Cy3-Rho; Cy3-Rho accumulates at nascent cleavage furrow

(arrowhead); C) *X. laevis* embryo microinjected with IT-Rho (left) and Cy3-Rho (right); both are

enriched at cell-cell junctions; D) Meiotically mature *Xenopus* egg microinjected with IT-Cdc42;

IT-Cdc42 is recruited to exocytosing cortical granules (arrowheads) following egg activation

(0:42); E) *X. laevis* embryo microinjected with IT-Cdc42; IT-Cdc42 concentrates at cell-cell

junctions (arrowhead; 0:00) and, following damage, is recruited to the wound and becomes

enriched at junctions (arrow); F) C3-insensitive IT-Rho rescues Rho activation in presence of

C3. Control: cell microinjected with rGBD shows normal Rho activity accumulation and wound

closure; C3: cell microinjected with rGBD fails to activate Rho in presence of C3; IT-Rho-

N41V: cell microinjected with rGBD and C3-insensitive IT-Rho normally activates Rho; IT-

Rho-N41V+C3: cell microinjected with rGBD and C3-insensitive IT-Rho rescues Rho activity in

presence of C3.Scale bar 10μm, time min:sec. G) Quantification of Rho activity, corrected for

background (n=8-12). One-way ANOVA with Tukey post-test statistical analysis. *p<0.05,

***p<0.001, ****p<0.0001.

Figure 3: Pools of inactive and active Cdc42 at the plasma membrane. A) Oocytes

microinjected with wGBD (green), Cy3-Cdc42 (magenta) and indicated concentrations of

mRNA encoding the Cdc42-GAP Abr. Scale bar 10μm, time min:sec.; B) Quantification of Cy3-

Cdc42, active Cdc42 and ratio of active Cdc42 to Cy3-Cdc42 for each condition. n=8-24. One-

way ANOVA with Tukey post-test statistical analysis. *p<0.05, **p<0.01, ***p<0.001,

***p<0.0001.

Figure 4: RhoGDI is recruited to single-cell wounds enriched in Rho and Cdc42 activity. A)

Oocytes microinjected with wGBD (magenta), rGBD (blue) and GDI (green); B) Wounded

oocytes fixed and stained with anti-*X. laevis* RhoGDI. Scale bar 10μm, time min:sec.

Figure 5: RhoGDI overexpression differentially regulates Rho and Cdc42 activity. A) Top 2

rows: oocytes microinjected with Cy3-Cdc42 (magenta) and wGBD (green) alone or with

RhoGDI. Bottom 2 rows: Oocytes microinjected with Cy3-Rho (magenta) and rGBD (green)

alone or with GDI; B) Oocytes microinjected with wGBD (magenta), rGBD (green) and

increasing concentrations of RhoGDI protein; C) Standard curve of decrease in Rho and Cdc42

activity with increasing concentrations of RhoGDI (n=10-23 for each concentration). Scale bar

10μm, time min:sec.

Figure 6: RhoGDI actively extracts RhoGTPases from membranes *in vitro*. A) Experimental

setup of *in vitro* experiments: prenylated RhoGTPases were reconstituted on supported lipid

bilayers (SLBs) in flow chambers and imaged by TIRF. Wash off experiments were designed to

avoid RhoGTPase rebinding to membranes and performed controlling the flow rate via a syringe

pump; B) TIRF imaging allows for selective imaging of RhoGTPases at the membrane.

Prenylated and unprenylated Cdc42 were imaged in the same conditions and TIRF signal at

membranes was quantified (n=4 for each condition); C) Wash off experiments: prenylated Cdc42

reconstituted on SLBs were washed with imaging buffer only (control), in presence of 5μM

RhoGDI or RabGTTase Beta. Time lapse images at selected time points and quantification of the

full experiments are shown. Decay curves were fitted with a monoexponential function; D)

Comparison of the K_{off} values obtained by fitting the decay curves (n=3 for control and

RabGTTase Beta, n=2 for RhoGDI); E) Schematic representation of the proposed mode of action

of the two RhoGTPases solubilizers. RabGGTase Beta sequesters RhoGTPases in solution,

whereas RhoGDI actively extracts RhoGTPases from the membranes. Scale bar 10μm. Unpaired

student's t-test, 2-tailed distribution, equal variance statistical analysis. **p<0.01, ***p<0.001,

****p<0.0001.

Figure 7: RhoGDI actively extracts both inactive and active RhoGTPases from membranes

***in vitro*.** A) Wash off experiments: prenylated Cdc42 in both inactive (Cdc42:GDP) and

constitutively active (Cdc42Q61L:GTP) states were reconstituted on SLBs and washed in

presence of 5 μM RhoGDI. Time lapse images at selected time points are shown; B)

Quantification of wash off experiments in which the concentration of RhoGDI was titrated

between 0 and 20μM; C) K_{off} values obtained for inactive and constitutively active Cdc42Q61L

fitting the decay curves with a monoexponential decay function are plotted against RhoGDI

concentration. Extraction rates were fitted with a hyperbolic function; D) Ratio of K_{off} obtained

for inactive and constitutively active Cdc42Q61L at the same RhoGDI concentration; F) Same as

in C-D for inactive (RhoA:GDP) and constitutively active (RhoAQ63L:GTP) RhoA. Scale bar

10μm.

Figure 8: Identification of mutant RhoGDI deficient in extraction of active RhoGTPase. A)

Oocytes microinjected with halo-tagged WT, Δ51-199, E158/9A and E158/9Q GDI mutants.

Scale bar 10μm, time min:sec; B) Quantification of RhoGDI intensity at wounds (n=7-13).

Unpaired student's t-test, 2-tailed distribution, unequal variance statistical analysis to WT.

****p<0.0001. C) Comparison of K_{off} values obtained for inactive (Cdc42:GDP, RhoA:GDP)

and constitutively-active (Cdc42Q61L:GTP, Cdc42Q63L:GTP) Cdc42 and RhoA from wash off

experiments in presence of either WT (black) or E158/9Q (QQ) (red) RhoGDI. Extraction rates

were fitted with a hyperbolic function.

Figure 9: RhoGDI extracts active Cdc42 *in vivo*. A) Oocytes microinjected with wGBD alone

or with constitutively-active Cdc42 (G12V), WT or QQ GDI; B) Quantification of total Cdc42

activity for (A), (n=12); C) oocytes microinjected with Cy3-Cdc42 bound to GTPγS alone or

with WT or QQ GDI; D) Quantification of intensity for (C), (n=18). Scale bar 10μm, time

min:sec. One-way ANOVA with Tukey post-test statistical analysis; E) Oocytes microinjected

with wGBD (magenta), rGBD (green) alone or with QQ GDI; F) Quantification of total Cdc42

(magenta) and Rho (green) activity from (E) (n=9); G) Cy3-Cdc42 or Cy3-Rho alone or with QQ

GDI; H) Quantification of total recruitment of Cy3-Cdc42 (magenta) and Cy3-Rho (green) (n=6-

11). Scale bar 10μm, time min:sec. Unpaired student's t-test, 2-tailed distribution, unequal

variance statistical analysis. *p<0.05, **p<0.01, ***p<0.001, ***p<0.0001.

Figure 10: Schematic of proposed update to RhoGTPase cycle. We propose that in addition

to the canonical GTPase cycle, GDI can extract active GTPase from the plasma membrane.

Based on evidence that GTPase:GDI binding prevents GTP hydrolysis and nucleotide exchange

(Hart et al., 1992; Ueda et al., 2001), active GTPase extracted by GDI would still be active upon

its release back into the plasma membrane.

Supplemental Figure 1: Amino-terminally tagged RhoGTPases do not localize properly to

wounds. Oocytes injected with A) mCh-Cdc42 (magenta) and wGBD (green), B) mCh-Rho

(magenta) and rGBD (green) or C) mCh-Rac (magenta) and wGBD (green) with A'-C')

Corresponding line scans. Scale bar 10μm, time min:sec.

Supplemental Figure 2: Internally-tagged Rac localizes to wounds. A) Oocyte injected with

wGBD (magenta) and IT-Rac (green); A') Corresponding line scan. Scale bar 10μm, time

min:sec.

Supplemental Figure 3*: X. laevis* **RhoGDI antibody specificity and purified RhoGDI**

protein. A) Western blot stained with αGDI antibody to determine specificity; lane 1: X. laevis

oocyte whole cell lysate (WCL), lane 2: WCL of oocytes overexpressing GDI, lane 3: WCL of

oocytes expressing 3xGFP-RhoGDI, lane 4: purified FLAG-RhoGDI; B) Coomassie stain of

12% SDS-PAGE to assess purity of FLAG-RhoGDI; lane 1: start, lane 2: void, lanes 3-10:

elution fractions. C) Coomassie stain of 12% SDS-PAGE to assess purity of RhoGDI purified

from bacteria; lane 1: start, lane 2: flow through, lane 3: TEV cleavage, lane 4: tag removal, lane

5: end product after gel filtration.

Supplemental Figure 4: Bovine RhoGDI decreases Rho and Cdc42 activity in a dose-

dependent manner *in vivo*. Standard curve of decrease in Rho and Cdc42 activity with

increasing concentrations of bovine RhoGDI (n=2-30 for each concentration).

Supplemental Figure 5: *In vitro* data analysis. A) Raw data. Wash off experiments were

started after the RhoGTPase signal at the membrane was stable. After starting the syringe pump,

a signal overshoot occurred; B) The overshoot signal was used as a reference point to align

different experiments and cut off for further analysis; C) Data were background corrected and

normalized to display multiple curve on the same graph; D) Data were fitted with a

monoexponential decay curve to obtain dissociation constants (K_{off}) of the GTPases from the

membrane.

Supplemental Figure 6: Comparison of bovine and *Xenopus* RhoGDI in their ability to

actively extract both inactive and active RhoGTPases from synthetic membranes. A) K_{off}

values obtained for inactive and constitutively active Cdc42 at different bovine RhoGDI

concentrations. Extraction rates were fitted with a hyperbolic function; A') Ratio of K_{off} obtained

for inactive and constitutively active Cdc42 at the same bovine RhoGDI concentration; B-B')

same as in A-A' for inactive (RhoA:GDP) and constitutively active (RhoAQ63L:GTP) RhoA.

Supplemental Figure 7: **Analysis of previously-described extraction-deficient mutants.** A-I)

Oocytes microinjected with WT or mutant RhoGDI and quantification of localization relative to

WT RhoGDI; Unpaired student's t-test, 2-tailed distribution, unequal variance statistical

analysis; J) Average K_{off} values obtained for inactive RhoA and Cdc42 in absence (control) and

presence of 5 μM of either bovine RhoGDI WT or bovine RhoGDI mutants.

Supplemental Figure 8: Mutant E163/4Q bovine RhoGDI is deficient in extraction of active

Cdc42 and RhoA *in vitro*. Comparison of K_{off} values obtained for inactive (Cdc42:GDP,

RhoA:GDP) and constitutively-active (Cdc42Q61L:GTP, Cdc42Q63L:GTP) Cdc42 and RhoA

from wash off experiments in presence of either WT (black) or E163/4Q (QQ) (red) bovine

RhoGDI. Extraction rates were fitted with a hyperbolic function.

Supplemental Figure 9: Schematic of RhoGDI's role in RhoGTPase zone definition around

wounds. Active Rho (green) and active Cdc42 (red) are activated in discrete, concentric zones

that close inward as the wound heals. Cdc42 inactivation is variable throughout its zone, thus its

zone is shaped by activation (Burkel et al., 2012). Conversely, the Rho zone is shaped by

inactivation as it is subject to RhoGAPs 1/8 at the trailing edge of its zone (Davenport, 2016).

Thus, we hypothesize that GDI extracts active Cdc42 throughout the Cdc42 zone and inactive

Rho from the trialing edge of the Rho zone.

Supplemental Figure 10: Comparison between G12V and Q61L constitutively-active

RhoGTPases. K_{off} values obtained for constitutively-active Cdc42 G12V fitting the decay

curves with a biexponential decay function are plotted against RhoGDI concentration. The fast

rate (black) corresponds to the GDP-bound state (dotted black line), the slow rate (red) to the

active GTP-state as inferred from the Q61L mutant (dotted red line).

Supplemental Figure 11: Cdc42 Q61L does not behave like constitutively-active Cdc42 *in*

***vivo*.** Oocytes injected with wGBD alone or with Cdc42 Q61L. Scale bar 10μm, time min:sec.

Supplemental Discussion

Two naturally occurring, oncogenic mutations originally discovered in Ras (Q61L and

G12V) are commonly used as constitutively-active variants of Ras-like GTPases such as Cdc42

and RhoA. While often used interchangeably, it is important to note that the biochemical

properties of these two single site mutations are not equivalent (Smith, Neel, & Ikura, 2013). The

Q61L substitution directly affects a catalytic residue within the RhoGTPase active site and

therefore directly and strongly impairs spontaneous nucleotide hydrolysis. This leads to the

Q61L mutant being dominantly GTP-bound in the absence of external regulators such as GEFs

and GAPs. In other words, GTPases affected by the Q61L substitution do not require GEF

activity to adopt an active, GTP-bound state.

The G12V mutation, on the other hand, targets an auxiliary site important for GAP-

mediated hydrolysis. GTPases affected by the G12V mutation are still capable of hydrolyzing

GTP intrinsically and can thus accumulate in the inactive, GDP-bound state in the absence of

external regulators. Their "constitutive activity" rather originates from a complete deficiency in

inactivation via GAP-induced GTP hydrolysis. In line with this, we found that these two

mutants, when purified recombinantly, accumulate in different nucleotide states as determined

by HPLC analysis: Q61L mutant proteins were GTP-bound, whereas G12V ones were GDP

bound after purification. We also found that the two produced different effects *in vivo* with

G12V behaving as expected for a constitutively-active (ie its expression resulted in an excess of

Cdc42 activity around the wound) while Q61L does not (ie its expression modestly or

significantly reduced Cdc42 activity around wounds; SuppFig11).

With respect to the results obtained with these mutants *in vitro*, Q61L mutant GTPases

(constitutively bound to GTP) showed monophasic membrane dissociation kinetics upon

addition of GDI with rates that were clearly slower than those obtained for inactive GTPases

(Fig7). This shows that the Q61L mutant uniformly adopts an active, GTP-like state distinct from

the GDP form.

Since the G12V mutant was bound to GDP after purification, we first exchanged its

nucleotide to GTPγS. GTPases prepared in this manner were however extracted by RhoGDI with

biphasic kinetics with two characteristic rates, indicating the presence of two biochemically

distinct GTPase species. Interestingly, the fast rate corresponded to the GDP-bound state,

whereas the slow rate was nearly identical to the active GTP-state as inferred from the Q61L

mutant (SuppFig10). From these observations, we drew the following conclusions: i) the G12V

mutant likely hydrolyzes a considerable fraction of their associated GTPγS during the long time

needed to prepare our extraction assays (often multiple hours, see Methods), ii) the "active

states" adopted by Q61L (GTP, complete) and G12V (GTPγS, partial) are identical concerning

GDI-mediated extraction from membranes. Because of these two reasons combined, we chose to

work with the Q61L mutants as a proxy for the active GTPase state for all our *in vitro* extraction

assays.

1019

1020

1021

1022

1023

1024

1025 **Figure 1**

1026

1027

1028 **Figure 2**

1029

1036   **Figure 4**

1037

1038

1039 **Figure 5**

1040

1041   **Figure 6**

1042

1043   **Figure 7**

1044

1045

1046

1047 **Figure 8**

1048

1049

1050

1051

1052

1053

1054   **Figure 9**

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067   **Figure 10**

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079   **Supplemental Table 1: Primers.**

Primer Name	Primer sequence (5'-3')
Cdc42(1)	cccgggcccgggaattcatgcagacaattaaatggtgtagtcgttgg
Cdc42(2)	cgcgccagagcatgcggatccgccagactgtttgtttttttgccag
Cdc42(3)	gatccgcatgctctggcgcgccgggcaaaccaatc
Cdc42(4)	cccgcgcggccctcgagttatagcagcatacacttgcgtttcttc
Cdc42(Q61L)fwd	ctttttgatactgcagggctagaggattatgacagattacg
Cdc42(Q61L)rev	cgtaatctgtcataatcctctagccctgcagtatcaaaaag
Cdc42(G12V)fwd	gtgttgttgtgggcgatgttgctgttggtaaaacatg
Cdc42(G12V)rev	catgttttaccaacagcaacatcgcccacaacaacac
GTPase(GeneStrand)fwd	gagaatctttattttcagggc
GTPase(GeneStrand)rev	ggtggtggtgctcgagtgc
Cdc42(Sf9)fwd	ccactactgagaatctttattttcaggggtggtggtggtggtatgcagacaattaagtgtgttgttg
Cdc42(Sf9)rev	gcaggctctagattcgaaagcgttattatagcagcacacacctgcgg
pETfwd	ggccgcactcgagcaccacc
pETrev	cgccctgaaaataaagattctc
pFASTBacH10fwd	cgctttcgaatctagagcctgc
pFASTBacH10rev	ctgaaaataaagattctcagtagtgg
pCS2+-Cdc42(1)	cttgttcttttgcaggaaaccatcgattcgaattcatgcagac
pCS2+-Cdc42(2)	tcgaatcgatggtttcctgcaaaaagaacaagtagcttgtattc
Rac(1)	cccgggcccggaattcatgcaggccattaaatgtgtgg
Rac(2)	cgcgccagagcatgcggatccgccagagagtttcttttctttcagcttctcaatagtgtc
Rac(3)	gatccgcatgctctggcgcgccgggcacccc
Rac(4)	cccgcgcggccctcgagttacaacagccgacatcttc
pCS2+-Rac(1)	ctttttgcaggcccccatcgatatgcagccattaaatgtg
pCS2+-Rac(2)	gcctgcatatcgatggggggcctgcaaaaagaacaagtagc
Rho(1)	cccgggcccggaattcatggcagccattcgtaagaagctcg
Rho(2)	cgcgccagagcatgcggatccgccagactgtttcattttggtgagctccct
Rho(3)	gatccgcatgctctggcgcgccgggcgagcctgtgaag
Rho(4)	cccgcgcggccctcgagttagatgagaaggcacgtgg
RhoA(Q63L)fwd	gggacacagctgggctggaagattatgatcgc
RhoA(Q63L)rev	gcgatcataatcttccagcccagctgtgtccc
RhoA(G14V)fwd	atgtctttccacaggctacatcaccaacaatcacc
RhoA(G14V)rev	ggtgattgttggtgatgtagcctgtggaaagacat
RhoA(Sf9)fwd	ccactactgagaatctttattttcaggggtggtggtggtggtatggctgccatccggaagaaac
RhoA(Sf9)rev	gcaggctctagattcgaaagcgttattacaagacaaggcaaccagattttttc
pCS2+-Rho(1)	caagctacttgttcttttgcaggatggcatcgattcgaattc
pCS2+-Rho(2)	ggctgccatgaattcgaatcgatgccatcctgcaaaaagaac
eGFP(1)	ctggcggatccatggtgagcaaggg
eGFP(2)	gcattggcgcgcgcccttgtacagctc
8(A)F1	ggcgctgcagctgtggcagcagcagtcgaactgaac
8(A)F2	ggccgcaaaggcaggcatcaagcatggcgctgcagctg
8(A)F3	gatcccatcgattcgaattcatggccgcaaaggcag

8(A)F4	cgagctgtacaagtccggaatggccgcaaaggcag
R1	cactatagttctagaggctcgagttaatctttccactctttc
R2	gaattcgaagcttgagctcgacgttaatctttccactc
HR F1	ggacaagggaggtggtggaagtggtggaggaggttctgcccaagtggatcc
HR F2	tgaactacaaggccccggagatgaaatctctgcaggaaatccaagagttggacaagggag
HR F3	catcaagcatggcgaggaggaggtggaagaagaagtcgaactgaactacaag
HR F4	gatcccatcgattcgaattcatggccgacaaggatggcatcaagcatgg
HR F5	cgagctgtacaagtccggaatggccgacaaggatggcatcaagcatgg
(-)20 F1	ggatcccatcgattcgaattcatgaactacaaggccccg
(-)20 F2	gctgtacaagtccggatgaactacaaggcccc
(-)55 F1	gatcccatcgattcgaattcatggcccaagtggatcctaaccttc
(-)55 F2	cgagctgtacaagtccggaatggcccaagtggatcctaaccttc
XlRhoGDIfwd	ctgttccaggggcccctgggatcctgtatggccgacaaggatggc
XlRhoGDIrev	gatcgtcagtcagtcacgatgcggccgcttatcaatctttccactctttctttatgg
E163/4Qfwd	gtatgagttcctgaccccccatgcagcaggcgcccaagggcatgc
E163/4Qrev	gcatgcccttgggcgcctgctgcatggggggtcaggaactcatac
Δ1-22fwd	ctgttccaggggcccctgggatcctgtcactcagtcaactataagc
Δ1-22rev	cgtcagtcagtcacgatgcggccgctcagtccttcc
Δ1-59fwd	ctgttccaggggcccctgggatcctgtgctgtgtctgctgacccc
Δ1-59rev	cgtcagtcagtcacgatgcggccgctcagtccttcc
HRfwd	gtggtggaagtggtggaggaggttctgctgtgtctgctgacccc
HRrev	ctcctccaccacttccaccacctcccttatagttgactgagtgctcg
pGEXHRfwd	cgagcactcagtcaactataagggaggtggtggaagtggtggaggaggttctgctgtgtctgctgacccc
pGEXHRrev	cgagcactcagtcaactataagggaggtggtggaagtggtggaggag

1080

1081   **Supplemental Figure 1**

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092 **Supplemental Figure 2**

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108 **Supplemental Figure 3**

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122 **Supplemental Figure 4**

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136 **Supplemental Figure 5**

1137

1138

1139

1140

1141

1142

1143

1144

1145 **Supplemental Figure 6**

1146

1147

1148

1149

1150

1151

1152

1153

1154   **Supplemental Figure 7**

1155

1156   **Supplemental Figure 8**

1157

1158

1159

1160

1161

1162   **Supplemental Figure 9**

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176 **Supplemental Figure 10**

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189 **Supplemental Figure 11**

1190

1191

1192

1193

1194

1195