Title: Effects of the copper IUD on composition of the vaginal microbiota in the olive

baboon

Authors: Eastman AJ[a], Sack D[a], Chai D[b], Bassis CM[c], Young VB[c,d], Bergin IL[e] and Bell

JD[a].

Affiliations:

[a]University of Michigan Department of Obstetrics and Gynecology, 1500 East Medical

Center Drive, L4100, Ann Arbor, MI 48109, USA

[b]Institute of Primate Research, National Museum of Kenya, End of Karen Road, P.O.

Box 24481 Karen 00502, Nairobi, Kenya

[c]University of Michigan Department of Internal Medicine Division of Infectious

Diseases, 3101 Taubman Center, SPC 5368, 1500 East Medical Center Drive, Ann

Arbor, MI 48109, USA

[d]University of Michigan Department of Microbiology and Immunology, 1150 West

Medical Center Drive , 5641 Medical Science II Ann Arbor , Michigan 48109-5620

[e]University of Michigan Unit for Laboratory Animal Medicine, 2800 Plymouth Rd, Ann

Arbor, MI 48109, USA

Corresponding Author:

Jason Bell

University of Michigan

Department of Obstetrics & Gynecology

1500 East Medical Center Drive, L4100

Ann Arbor, MI, USA 48109

(p) (734) 232-3897

(f) (734)647-9727

jasonbel@umich.edu

Abstract word count: 100

Manuscript word count: 996

Abstract:

Objectives: Determine the feasibility of copper intrauterine devices (Cu-IUD) in baboons

and impact on vaginal microbiota.

Study design: Vaginal swabs were taken before insertion (pre-IUD), during IUD use

(IUD), and after removal (post-IUD) and microbiota assessed by 16S rRNA-encoding

gene sequence analysis.

Results: No animals showed physical changes or discomfort during pre-IUD, IUD, or

post-IUD phases. There were significant vaginal bacterial community differences in the

post-IUD phase.

Conclusions: The baboon is a viable model for Cu-IUD investigation, with Cu-IUD

removal, but not insertion, altering vaginal microbiota.

Implications: A baboon model of Cu-IUD allows investigation into the intersection of

Cu-IUDs and human reproductive tract disorders and pathogens.

Introduction

A non-human primate (NHP) model for copper intrauterine devices (Cu-IUD) that mimics all aspects of human Cu-IUD usage, including both the device and transcervical insertion, has been elusive. Macaques have a tortuous cervix, which prevents transcervical insertion, and their small size may make device tolerance challenging. The olive baboon (*Papio anubis*) is larger and has a straight cervical canal, making it an ideal animal model for intrauterine contraception. A NHP model for Cu-IUDs will permit studies regarding the impact of Cu-IUDs on the biology of the female reproductive tract and the relationship between intrauterine contraception and infectious agents (*e.g.* *Chlamydia trachomatis*, HIV) that may be ethically impossible in humans.

Our laboratory has previously developed a NHP model of levonorgestrel-releasing intrauterine system (LNG-IUS) with olive baboons, which assessed changes to the vaginal microbiome in baboons with LNG-IUS. We found inter-individual differences, but no consistent device-associated shift in microbial communities [1]. Animals with LNG-IUSs had a moderate shift towards more stable microbial communities over time [1]. In this study, we assessed physiologic changes in a cohort of 5 baboons with Cu-IUD and analyzed populations of the vaginal microbiota over 24 weeks.

Methods

Regulatory approval

The Institutional Review Committee at the Institute of Primate Research (IPR) in Nairobi, Kenya (NIH Office of Laboratory Animal Welfare foreign assurance A5796)

approved this study. This study received an off-site exemption from the University of

Michigan University Institutional Animal Care and Use Committee. An export permit

was obtained from the Kenyan Wildlife Service and import permits were obtained from

the US CDC [1].

Study site, population, design, sampling

This study was conducted at the IPR (for IPR description, see [1]). Five wild-caught

sexually mature female olive baboons (*P. anubis*) were utilized in this study, quarantined,

examined, housed individually, and vaginally sampled as previously described [1]. Study

timecourse was as follows: pre-IUD phase weeks 0-4, weekly sampling; Cu-IUDs

inserted according to package insert after week 4 sampling; Cu-IUD phase weeks 5-20,

sampling every 4 weeks; Cu-IUDs removed after week 20 sampling; post-IUD phase

weeks 21-24, weekly sampling.

Microbiome DNA preparation, sequencing, and analysis

DNA extraction from vaginal swabs was performed with the PowerMag Soil DNA

Isolation Kit (Mo Bio, Carlsbad, CA, USA), and amplification of the V4 region of the

bacterial 16S rRNA gene, sequencing (Illumina MiSeq), sequence processing and

analysis were performed as previously described [1].

Results

Clinical signs

All baboons retained the Cu-IUD for the planned duration (from week 5-20). No animals

showed any physical changes or discomfort (*e.g.* changes in weight, temperature, pulse,

respiration, and frequency of discharge) during the pre-IUD (weeks 0-5), IUD (weeks 6-

20), or post-IUD (weeks 21-24) phases of the study, nor did the IUD appear to disrupt the

menstrual cycle (data not shown).

Week-to-week vaginal microbial community structure variation

16S rRNA gene sequence analysis of vaginal swabs was used to compare baboon

bacterial communities, both between individual animals and within an individual animal

over time. We first analyzed community structure within and between baboons by θ_{YC}

dissimilarity coefficient, a measure of both the presence of specific operational

taxonomic units (OTUs) and their relative abundance [2]. Examination of the average θ_{YC}

distance showed high within-animal week-to-week variation in community structure (Fig.

S1). The greatest shift in bacterial community composition, indicated by the highest θ_{YC}

distances, occurred between week 0 and week 1, when the animals were moved from

group housing to single cages. Community stability, measured by week-to-week θ_{YC}

distances, did not change significantly between pre-IUD, IUD, and post-IUD timepoints

(Fig. 1).

Microbial community composition and clustering

The most frequent bacterial phyla across all baboons were Bacteroidetes, Fusobacteria,

Proteobacteria, and Firmicutes (Fig. 1). When we compared the pre-IUD, IUD, and post-

IUD timepoints by analysis of molecular variance (AMOVA) [3] on θ_{YC} distances, there

was a significant difference between post-IUD communities and both pre-IUD

communities (p-value: 0.003) and IUD communities (p-value: <0.001), but not a

significant difference between pre-IUD and IUD communities (p-value: 0.126) (Fig. 2).

Differentially abundant OTUs identified by LEfSe [4] included OTU1 (Leptotrichiaceae),

which was higher in the post-IUD timepoints relative to both pre-IUD and IUD (Fig. S2,

Table S1).

When we followed community structure over time, communities within an

individual were more similar then those found in different animals (Fig. S3).

Discussion

This study was designed to test the feasibility of Cu-IUD placement in baboons

and assess vaginal microbiota changes due to Cu-IUD placement. We found no clinical

changes to baboons (*e.g.* temperature, weight, menstrual cycle, discharge) and a slight but

statistically significant alteration of the vaginal microbiota after Cu-IUD removal. As we

have demonstrated with the LNG-IUS in the baboon, we conclude that the olive baboon

is a feasible model for Cu-IUD studies.

The mechanism behind the change in vaginal microbiota is unclear, as is the

biological significance. Further investigation is necessary to determine why removal of

the Cu-IUD altered community structure while insertion of the Cu-IUD did not. The lack

of change between the pre-IUD phase and IUD phase is consistent with recently

published work in humans, although the human study did not extend to after IUD

removal [5].

Animal models of Cu-IUD are limited. In mice, insertion of a copper wire into the

uterine horn [6] is limited by the reliance on surgery, the vastly different estrus cycle and

reproductive tract physiology from humans, and the more limited range of naturally-

occurring sexually-transmitted pathogens relative to humans and NHP. NHP studies of

Cu-IUD have been successfully performed in the pigtail macaque [7], but surgery is

necessary for device insertion, which carries significant risks to the animal. While the

baboon is not the primary NHP model for HIV, it is a viable model [8], and studies

wishing to evaluate the interaction of IUDs with HIV are feasible. Moreover, baboons are

used for models of endometriosis [9] and their high vaginal pH and diverse microbial

communities may model aspects of bacterial vaginosis [10]. Our study opens a new

avenue of research into the intersection of the Cu-IUD and human reproductive tract

disorders and pathogens.

Acknowledgements

The authors would like to acknowledge the Human Microbiome Initiative.

References

[1] Liechty ER, Bergin IL, Bassis CM, et al. The levonorgestrel-releasing intrauterine system is associated with delayed endocervical clearance of Chlamydia trachomatis without alterations in vaginal microbiota. Pathogens and disease. 2015;73:ftv070.

[2] Yue JC, Clayton MK. A Similarity Measure Based on Species Proportions. Communications in Statistics - Theory and Methods. 2005;34:2123-31.

[3] Anderson MJ. A new method for non-parametric multivariate analysis of variance. Austral Ecology. 2001;26:32-46.

[4] Segata N, Izard J, Waldron L, et al. Metagenomic biomarker discovery and explanation. Genome biology. 2011;12:R60.

[5] Bassis CM, Allsworth JE, Wahl HN, Sack DE, Young VB, Bell JD. Effects of intrauterine contraception on the vaginal microbiota. Contraception. 2017;96:189-95.

[6] Chou CH, Chen SU, Shun CT, Tsao PN, Yang YS, Yang JH. Divergent endometrial inflammatory cytokine expression at peri-implantation period and after the stimulation by copper intrauterine device. Scientific reports. 2015;5:15157.

[7] Engel RM, Morris M, Henning T, et al. Evaluation of pigtail macaques as a model for the effects of copper intrauterine devices on HIV infection. Journal of medical primatology. 2014;43:349-59.

[8] Locher CP, Witt SA, Herndier BG, Tenner-Racz K, Racz P, Levy JA. Baboons as an animal model for human immunodeficiency virus pathogenesis and vaccine development. Immunological reviews. 2001;183:127-40.

[9] Braundmeier AG, Fazleabas AT. The non-human primate model of endometriosis: research and implications for fecundity. Molecular human reproduction. 2009;15:577-86.

[10] Rivera AJ, Frank JA, Stumpf R, et al. Differences between the normal vaginal bacterial community of baboons and that of humans. American Journal of Primatology. 2011;73:119-26.

Figure Legends:

Figure Legends:

Figure 1. Bacterial communities exhibit inter-animal variation, but no pattern in change

between pre-IUD, IUD, and post-IUD periods. Week-to-week change in θYC values in

each animal is matched to the bar chart of microbial communities at each timepoint.

Microbial communities are color coded as indicated, and are grouped by phylum

indicated. LEfSe analysis performed on each individual animal by grouping communities

found in the pre-IUD, IUD, and post-IUD timepoints found: animal 3903 had significant

difference in relative abundance between the pre-IUD and IUD phases; animal 3919 had

a significant difference in relative abundance between the IUD and post-IUD timepoints;

animals 3919 and 4047 had shifts in relative abundance of microbial communities

between the pre-IUD and post-IUD periods.

Figure 2. Principal Coordinates Analysis (PCoA) shows no distinct clustering of OTUs

with regard to IUD insertion but changes after IUD removal. Baboons at each timepoint

are represented by a circle, colored to indicate pre-IUD, IUD, and post-IUD periods.

Axes indicate percentage of variation by the plotted principal coordinates. Operational

Taxonomic Units (OTUs) of related bacteria are overlaid on the data. Significant

differences (in OTU1 Leptotrichaceae) were seen by AMOVA when post-IUD timepoints

were compared to both pre-IUD and IUD timepoints.

Supplemental Figure Legends:

Figure S1. Averaged week-to-week θYC values do not change significantly between pre-

IUD, IUD, and post-IUD periods.

Figure S2. Principal Coordinates Analysis (PCoA) show animals are most similar to

themselves and change little over 24 weeks. Baboons at each timepoint are represented

by a circle, colored to indicate individual animals. Axes indicate percentage of variation

by the plotted principal coordinates. Operational Taxonomic Units (OTUs) of related

bacteria are overlaid on the data. No significant differences were seen by AMOVA.

Figure S3. Significant differences by LEfSe in 10 OTUs with abundance greater than

0.001 in any group. OTU number and p-value (by LEfSe) are indicated on the Y-axis and

the relative abundance is indicated on the X-axis. LEfSe analysis was a 3-way

comparison between timepoints pre-IUD, IUD, and post-IUD. When further analyzed by

2-way ANOVA with multiple comparisons: OTU0001 was significantly different from

pre-IUD and IUD; OTU0004 was significantly different in post-IUD relative to IUD; and

OTU0007 was significantly different in post-IUD relative to IUD. Our cutoff for

inclusion was OTUs with a relative abundance greater than 0.001 in any group.

Table S1. OTU identities for groups with significant differences in abundance between

timepoints determined by LEfSE.

Figure 1

Figure 2

Figure S1

Figure S2

Table S1

OTU	Size	Taxonomy
Otu0001	271779	Bacteria(100);Fusobacteria(100);Fusobacteria(100);Fusobacteriales(100);Leptotrichiaceae(100);unclassified(100);
Otu0004	89043	Bacteria(100);Bacteroidetes(100);Bacteroidia(100);Bacteroidales(100);Bacteroidaceae(100);Bacteroides(100);
Otu0007	71274	Bacteria(100);Bacteroidetes(100);Bacteroidia(100);Bacteroidales(100);Prevotellaceae(100);Prevotella(100);
Otu0015	31833	Bacteria(100);Firmicutes(100);Clostridia(100);Clostridiales(100);unclassified(100);unclassified(100);
Otu0020	23142	Bacteria(100);Actinobacteria(100);Actinobacteria(100);Actinomycetales(100);Actinomycetaceae(100);Mobiluncus(100);
Otu0021	22837	Bacteria(100);Bacteroidetes(100);Bacteroidia(100);Bacteroidales(100);Porphyromonadaceae(100);Porphyromonas(100);
Otu0031	9988	Bacteria(100);Firmicutes(100);Negativicutes(100);Selenomonadales(100);Veillonellaceae(100);unclassified(100);
Otu0036	7269	Bacteria(100);Firmicutes(100);Clostridia(100);Clostridiales(100);Clostridiales_Incertae_Sedis_XI(100);Anaerococcus(100);
Otu0042	6190	Bacteria(100);Firmicutes(100);Clostridia(100);Clostridiales(100);Peptostreptococcaceae(100);unclassified(100);
Otu0048	5547	Bacteria(100);Firmicutes(100);Clostridia(100);Clostridiales(100);Lachnospiraceae(100);Catonella(100);

Figure S3