# Neural and behavioral correlates of declarative memory are associated with temporal discounting in older adults

Karolina M. Lempert[a], Dawn J. Mechanic-Hamilton[b,c], Long Xie[d,e], Laura E.M. Wisse[d,e], Robin de Flores[d,e], Jieqiong Wang[d,e], Sandhitsu R. Das[d,e], Paul A. Yushkevich[d,e], David A. Wolk[b,c], Joseph W. Kable[a]

[a]Department of Psychology, University of Pennsylvania, Philadelphia, PA 19104, USA.

[b]Penn Memory Center, University of Pennsylvania, Philadelphia, PA 19104, USA.

[c]Department of Neurology, University of Pennsylvania, Philadelphia, PA 19104, USA.

[d]Penn Image Computing and Science Laboratory (PICSL), University of Pennsylvania, Philadelphia, PA 19104, USA.

[e]Department of Radiology, University of Pennsylvania, Philadelphia, PA 19104, USA.

Correspondence: Joseph W. Kable, kable@psych.upenn.edu

The authors declare no competing financial interests.

**Abstract**

When facing intertemporal choices, or decisions involving trade-offs between smaller, sooner and larger, delayed rewards, people tend to discount the value of future rewards. There are substantial individual differences in this tendency toward temporal discounting, however. The neurocognitive mechanisms underlying these individual differences remain unclear. One possibility is that individual differences in declarative memory ability underlie individual differences in temporal discounting. Here we tested this hypothesis in older adults (both cognitively normal and with mild cognitive impairment), given that declarative memory declines with age, at rates that vary across individuals. We related temporal discounting to performance on neuropsychological measures of declarative memory, as well as to structural measures of medial temporal lobe regions that are critical for memory. Performance on episodic and semantic memory retrieval tasks was associated with temporal discounting, such that people with better memory discounted delayed rewards less. This relationship held even when controlling for executive function abilities that might influence memory retrieval. Underlining the specificity of this association, performance on measures of executive function (Trail Making Test, Digit Span Backwards, lexical fluency) was unrelated to temporal discounting, and declarative memory ability was unrelated to risk preferences. Finally, cortical thickness in the entorhinal cortex was associated with reduced temporal discounting, with episodic memory performance partially mediating this association. These findings suggest that individual differences in temporal discounting are primarily driven by memory ability, and not executive function, and that a decline in medial temporal lobe structural integrity with aging may impact time preferences.

**Keywords:** temporal discounting, intertemporal choice, declarative memory, executive function, mild cognitive impairment, aging

**1. Introduction**

People often have to decide between smaller/sooner and larger/later rewards. In these intertemporal choices (Strotz, 1956), individuals tend to devalue, or discount, future outcomes. This leads them to prefer immediate rewards to even larger ones received after a delay (Mazur, 1987), a tendency known as temporal discounting. While most individuals exhibit some degree of temporal discounting, people vary widely in the rate at which they discount delayed rewards (Peters & Büchel, 2011). Steep temporal discounting (i.e., overvaluing the present) has been linked with problematic behaviors, such as smoking (Bickel, Yi, Kowal, & Gatchalian, 2008; Yi & Landes, 2012), alcohol abuse (Vuchinich & Simpson, 1998), gambling (Reynolds, 2006), drug addiction (MacKillop et al., 2011), and excessive credit card borrowing (Meier & Sprenger, 2010). Although there is a substantial literature examining neural correlates of value at the time of intertemporal choice (Frost & McNaughton, 2017; Kable & Glimcher, 2007), the neurocognitive mechanisms underlying individual differences in temporal discounting remain unclear. Here we test whether individual differences in temporal discounting are linked to variability in declarative memory abilities, and to structural measures of critical neural substrates for declarative memory in the medial temporal lobe (MTL). To do this, we leverage variability in cognitive abilities among older adults (both cognitively normal and with mild cognitive impairment, MCI), given that declarative memory declines as people age (Buckner, 2004; Park et al., 1996) at rates that vary across individuals (Rubin et al., 1998).

Better functioning of the declarative memory system, which includes both semantic and episodic memory, may support more future-oriented intertemporal choice by allowing for more vivid imagination of future outcomes (Addis et al., 2007). Consistent with this idea, imagining positive future events and retrieving positive autobiographical memories has been shown to

decrease temporal discounting in young adults (Benoit, Gilbert, & Burgess, 2011; Lempert, Speer, Delgado, & Phelps, 2017; Peters & Büchel, 2010). The critical memory processes may not be specific to *episodic* memory, since individuals with semantic dementia, a disorder characterized by a loss of semantic memory, show increased temporal discounting (Chiong et al., 2016). However, a clear association between declarative memory abilities and temporal discounting across individuals has not yet been established. There is some evidence that individuals in at-risk states for dementia, such as subjective cognitive impairment (Hu et al., 2017) and MCI (Lindbergh, Puente, Gray, Mackillop, & Miller, 2014) have higher temporal discounting rates, but this research is inconsistent (Chiong et al., 2016; Coelho et al., 2016). Two studies using a correlational design in older adults found null results when examining the association between episodic memory and temporal discounting (Boyle et al., 2012; Seinstra, Grzymek, & Kalenscher, 2015), but these studies included only cognitively normal individuals, who might have limited variance in episodic memory ability. They also used measures that are less susceptible to age-related decline (e.g., associative recognition; Howard, Bessette-Symons, Zhang, & Hoyer, 2006). Here we leveraged a large and well-characterized older adult sample with substantial variability in episodic memory retrieval ability, including individuals with MCI, and used measures that emphasized delayed free recall to relate performance on memory measures to temporal discounting.

In younger adults, MTL gray matter volume (Owens et al., 2017; Pehlivanova et al., 2018) and white matter integrity (Olson et al., 2009; Yu, 2012) significantly predict temporal discounting rates. However, this association has not been examined in older adults. Since the MTL is critical for both episodic and semantic memory (Henry & Crawford, 2004; Moscovitch et al., 2005; Pihlajamäki et al., 2000), and is especially vulnerable to structural changes with

aging (Jernigan et al., 1991; Wolk et al., 2016), we additionally examined the relationship between cortical thickness in MTL subregions and temporal discounting.

An alternative hypothesis links individual differences in temporal discounting to executive functions (e.g., working memory and cognitive flexibility), which depend on the dorsolateral prefrontal cortex (McClure & Bickel, 2014; McClure, Laibson, Loewenstein, & Cohen, 2004; Wesley & Bickel, 2014). Therefore, we also examined the relationship between temporal discounting and a composite measure of executive function, including the Trail Making Test and the Digit Span test. This allowed us to determine whether any potential relationship with temporal discounting was specific to declarative memory, rather than driven by a relationship between temporal discounting and cognitive abilities more globally. We also controlled for executive function when examining the relationship between declarative memory and temporal discounting to better isolate declarative memory from executive control processes.

To our knowledge, no study has directly compared whether frontal lobe-mediated executive function or temporal lobe-mediated declarative memory can better explain individual variability in temporal discounting. The neuropsychological tasks used to measure these two cognitive functions differ in a number of ways, however. For example, the episodic retrieval tasks rely more on verbal abilities, which also decline with aging. Therefore, here we performed a secondary analysis examining the association between measures of verbal fluency – lexical and semantic fluency – and temporal discounting. These fluency tasks share many features, including that they both require speeded responding and speech production: lexical fluency probes the ability to generate words beginning with a certain letter (e.g., "F"), whereas semantic fluency measures the ability to retrieve as many exemplars within a category (e.g., animals) as possible. Lexical fluency, however, has been shown to depend on the frontal lobe (Mummery, Patterson,

Hodges, & Wise, 1996; Tupak et al., 2012), since it involves keeping rules in mind (e.g., no proper nouns, no number words) and rapidly switching between categories of words. Good performance on semantic fluency, in contrast, requires an intact temporal lobe (Gourovitch et al., 2000; Henry & Crawford, 2004; Troyer, Moscovitch, Winocur, Alexander, & Stuss, 1998), including MTL structures (Greenberg, Keane, Ryan, & Verfaellie, 2009; Pihlajamäki et al., 2000), since this task depends on retrieving semantic knowledge. Lexical fluency may even require inhibiting associations between words that are based on meaning (Perret, 1974). Thus, if semantic fluency is associated with temporal discounting even when controlling for lexical fluency, this would provide further evidence for the involvement of memory and temporal lobe function in intertemporal preferences. It would also suggest that any relationship between declarative memory and temporal discounting is not limited to episodic memory.

Finally, given previous evidence that cognitive abilities are linked to preferences in both the time and risk domains, we assessed risk tolerance with a risky choice task. This enabled us to determine whether any potential association between cognitive abilities and temporal discounting was specific to temporal discounting, or if it extended to economic preferences in general.

**2. Materials and methods**

2.1. Participants

100 older adults (age: 72.01±6.82 years, range: 58-93 years; male/female: 42/58, White/Black/Multi-racial: 62/36/2) completed the study. To ensure sufficient variability in our cognitive measures, we included individuals with MCI in our sample. MCI is a syndromic label often conceptualized as an intermediate stage of cognitive decline between normal cognitive aging and mild dementia. While about 50% of MCI patients likely have underlying Alzheimer's

Disease pathology (Vos et al., 2015), the category is heterogeneous and not indicative of a specific pathological process. All subjects are part of the Clinical Core cohort of the University of Pennsylvania Alzheimer's Disease Core Center. Given the constraint that we would only recruit from this well-characterized cohort, we selected 100 as our target sample size, in order to detect a modest correlation ($r = 0.3$) with 90% power. This study was approved by the Institutional Review Board of the University of Pennsylvania, and all participants provided informed consent. Choice task data were collected from 1/5/2017 to 2/14/2018, and all participants completed the National Alzheimer's Coordinating Center Uniform Data Set neuropsychological test battery (Morris et al., 2006; https://www.alz.washington.edu/WEB/data_descript.html) within one year of completing the choice tasks (87.76±70.62 days;  range: 0 – 315 days). All subjects were deemed cognitively normal ($n = 74$; average Mini-Mental State Examination (MMSE) score: 29.06±1.10) or as having amnestic or multi-domain MCI ($n = 26$; MMSE: 27.04±2.20) based on consensus conference diagnosis by Alzheimer's Disease clinical experts. Individuals with MCI either had a single domain of impairment, memory ($n = 6$) or were impaired in memory and at least one other domain (e.g., language or executive function; $n = 20$). As of this point (10/24/2019) based on annual evaluations, seven of the MCI participants have converted to dementia, and one of the cognitively normal participants has converted to MCI. These changes in diagnosis were recorded 335±179 days (range: 131 – 671 days) after choice task data were collected.

2.2. Procedure

Participants completed choice tasks assessing temporal discounting and risk tolerance (details below). The order of the tasks was counterbalanced across subjects. Both tasks were

computerized (programmed in E-Prime 2.0, Psychology Software Tools, Sharpsburg, PA). Subjects were given extensive instructions as well as practice trials to confirm that they understood the tasks fully. They were also instructed that their choices were incentive-compatible. That is, at the end of the session, one choice from either the intertemporal choice or risky choice task was randomly selected to determine a bonus. Since participants did not know which choice would count, their best strategy was to treat each one as if it were the one that counted. The bonus was paid using a pre-paid debit card (Greenphire Clincard) on the day the payment was due. Because all payments were made this way, we introduced no differences in transaction costs for different types of payments (risky choice task payment, intertemporal choice immediate payment or intertemporal choice delayed payment). For delayed payments, subjects received payment on their Clincard on the date corresponding to the delay for the chosen option. The procedure lasted approximately 15 minutes. Both choice tasks were self-paced, and participants had up to 20 s to respond on each trial.

*2.2.1. Intertemporal choice task*

On each of the 51 trials in this task (Pehlivanova et al., 2018; Senecal, Wang, Thompson, & Kable, 2012; Yu et al., 2017), participants chose between a small amount of money available immediately, and a larger amount of money available at a specified delay. The delayed outcome was always one of three amounts ($25, $30, $35). Immediate reward amounts varied from $10 - $34, and delays ranged from 1-180 days. After they selected their choice, a checkmark appeared on the screen indicating which side they had pressed. Immediate and delayed rewards switched sides of the screen at random from trial to trial.

*2.2.2. Risky choice task*

On each trial of this task (60 choices), participants chose between a small amount of money ($1-$68) available for certain, and a larger amount of money ($10-$100) available with some risk. All risky options entailed a 50% chance of the larger amount and a 50% chance of $0. Probabilities were displayed graphically, using pie charts. The gamble and the safe option alternated sides of the screen randomly. If a participant chose the risky option on the randomly selected trial, the experimenter flipped a coin to determine if they would receive payment or $0.

2.3. Episodic memory measures

Scores on the following episodic memory tasks (Word List Memory Delayed Recall; Craft Story Delayed Recall; Benson Complex Figure Delayed Recall) were transformed to *z*-scores and averaged to form an episodic memory composite score.

*2.3.1. Word List Memory Delayed Recall*

In this test (Morris et al., 1989), participants were presented with a list of 10 high-frequency words one at a time and were asked to read the words aloud (2 second presentation). The word list was presented 3 consecutive times, in randomized order. After every presentation, participants were asked to recall the words (Immediate Recall). After a short delay of approximately 5 minutes, the participant was asked to recall as many of the ten words as they could. We included this Delayed Recall score as one of our measures of memory performance. Finally, participants were asked to identify the target words from the list of 10 presented words and 10 distractor words. Because performance on this recognition task was at ceiling across our sample (maximum score = 20; mean score = 19.38), we did not include it in our composite score.

*2.3.2. Craft Story Delayed Recall*

The Craft Story 21 (Craft et al., 1996) is a paragraph story learning and recall test, similar to the Logical Memory subtest of the Wechsler Memory Scales (Wechsler, 1987). The examiner read a story aloud once, then asked the participant to repeat the details of the story in the same words read by the examiner or in their own words. Points for verbatim (exact content words) and paraphrase recall (similar contextual story units) were summed individually. After ~15 minutes (14.52±2.30 min), the participant was asked to recall the story again. Once again, points for verbatim and paraphrase recall were summed individually. If the subject recalled no items from the Craft Story after the delay, the examiner provided a cue ("It was a story about a boy"). For this study, only the delayed paraphrase recall score (range: 1 to 25) was included in the composite score.

*2.3.3. Benson Complex Figure Delayed Recall*

In this assessment of visuospatial memory (Possin, Laluz, Alcantar, Miller, & Kramer, 2011), participants are first asked to copy a complex figure (a simplified version of the Rey-Osterrieth complex figure), and then to draw it from memory approximately 10-15 minutes later. Their recall score is based on the accuracy and placement of reproduced elements present in the figure drawing. We used their recall score (range: 0 to 17) as our third measure of episodic memory.

2.4. Semantic fluency measures

Semantic fluency (Morris et al., 1989) was measured by having participants name as many animals as they could in 60 seconds and as many vegetables as they could in 60 seconds. The total number of correct and unique animal and vegetable words were tallied, converted to *z*-scores, and then averaged.

2.5. Executive function measures

Scores on the Digit Span Backwards and Trails B-A (described below) were transformed to *z*-scores and averaged to form an executive function composite score.

*2.5.1. Digit Span Task (Backwards condition)*

The Digit Span Task (Wechsler, 1997), part of the Wechsler Adult Intelligence Scale, is among the most widely used neuropsychological tests. It assesses auditory attention and the maintenance and manipulation of information in short-term memory. In the Forwards condition, participants are asked to repeat, in the same order, a series of digits that are read aloud to them. In the Backwards condition, participants are asked to repeat the digits in the reverse order. Because the Backwards condition is a more sensitive measure of executive function (Groeger, Field, & Hammond, 1999), and the dorsolateral prefrontal cortex is engaged during performance of the Backwards condition (Gerton et al., 2004), the number of correct trials in this condition was used as one measure of executive function.

*2.5.2. Trail Making Test (Trails B-A)*

This test (Reitan, 1992) is given in two parts, A and B. Part A involves drawing a line connecting consecutive numbers from 1 to 25 (the numbers are scattered randomly on a page) as

quickly as possible. Part B involves drawing a line, connecting alternating numbers and letters in sequence as quickly as possible (i.e., 1-A-2-B, etc.). The time to complete each "trail" is recorded. The Trails B portion of the Trail Making Test is a widely used neuropsychological measure of frontal executive function (Davidson, Gao, Mason, Winocur, & Anderson, 2008; Stuss et al., 2001), involving attention, cognitive flexibility (Kortte, Horner, & Windham, 2002), and set-shifting. The difference between Part B time and Part A time is considered a measure of executive function, since performance on Part A accounts for any motor or processing speed differences between subjects. Because the distribution of time is skewed, "Trails B-A" scores were natural log-transformed before they were $z$-scored. They were also negated before being averaged with the other measure in the executive function composite, since higher Trails B-A scores are indicative of worse performance.

## 2.6. Lexical fluency measures

Lexical fluency (Benton & Hamsher, 1976) was measured by having the participant list as many words beginning with the letter "F" as they could in 60 seconds and as many "L" words as they could in 60 seconds. The total number of correct and unique "F" and "L" words were counted. Then the scores from these two trials were converted to $z$-scores and averaged.

## 2.7. Data analysis

Participants' individual choice data for the intertemporal and risky choice tasks were fit with the following logistic function using maximum likelihood estimation:

$$P_1 = \frac{1}{1 + e^{-\beta(SV1 - SV2)}}, P_2 = 1 - P_1$$

where $P_1$ refers to the probability that the participant chose option 1, and $P_2$ refers to the probability that the participant chose option 2. $SV_1$ and $SV_2$ refer to the participant's estimated subjective value of option 1 and option 2 respectively. $\beta$ was used as a scaling factor and was fitted for each individual task.

In the risky choice task, $P_1$ was the probability of choosing the risky option. $SV_1$ and $SV_2$ (for the risky option and safe option, respectively) were calculated using a power utility function:

$$SV = p * A^\alpha$$

Here $p = .5$ for the risky option, $p = 1$ for the certain option, and $\alpha$ is a risk tolerance parameter that varies across subjects. Higher $\alpha$ indicates greater risk tolerance (less risk aversion).

In the intertemporal choice task, $P_1$ was the probability of choosing the delayed option, and the subjective values of the options were estimated using a hyperbolic function, assuming a linear utility function (Green & Myerson, 2004; Mazur, 1987):

$$SV = \frac{A}{1 + kD}$$

Here $A$ is the amount of the option, $D$ is the delay until the receipt of the outcome (for immediate rewards, $D = 0$), and $k$ is a discount rate parameter that varies across subjects. Higher $k$ indicates higher discounting (less tolerance of delay). Since $k$ and $\alpha$ were not normally distributed, these values were natural log-transformed before conducting statistical analyses.

As an additional analysis to ensure that our temporal discounting results were not driven by differences in risk tolerance, we also estimated discount rates using a utility-transformed hyperbolic function (Lopez-Guzman, Konova, Louie, & Glimcher, 2018):

$$SV = \frac{A^\alpha}{1 + kD}$$

Here $\alpha$ is the risk tolerance parameter derived from the risky choice task. Fitting this additional model also ensures that our results do not depend on our choice of the linear-utility hyperbolic model.

Two multiple linear regressions were performed, for the temporal discount rate $k$ and the risk tolerance parameter $\alpha$. The independent variables of interest were the episodic memory composite score and the executive function composite score. In each regression, age, gender (0 = male; 1 = female), and years of education were entered as covariates of no interest. Partial Pearson correlation coefficients are reported. Given that we were testing four hypotheses (the relationships between two cognitive variables and two choice variables), we Bonferroni-corrected our significance threshold to 0.0125. In these regression analyses, and any analyses that included the executive function composite measure, four participants were excluded for being outliers on the Trail Making Test: one did not complete Trail Making Test Part B in the allotted time, and three had Trail Making Test Part B completion times that were more than 3 SD > mean (times of 280 s, 300 s, and 300 s).

For our secondary analysis on verbal fluency, we again conducted two multiple linear regressions, for the temporal discount rate $k$ and the risk tolerance parameter $\alpha$. The independent variables of interest were the semantic fluency score and the lexical fluency score. In these regressions too, age, gender, and years of education were entered as covariates of no interest. Partial Pearson correlation coefficients are reported.

In an exploratory analysis, we also conducted a one-way ANOVA to test for the effects of diagnosis (MCI vs. cognitively normal) on temporal discounting and risk tolerance.

2.8. Structural MRI data acquisition and analysis

Ninety-two participants in the sample also underwent MRI scanning. Most ($n = 53$) completed the choice tasks on the same day as their structural MRI scanning session, but a subset completed the tasks at a different time. For those 39 participants, MRI scans were acquired on average 345.79±167.76 days (range: 23 – 637 days) from the choice task session.

MRI data were obtained on a Siemens Prisma 3T MRI with a 64-channel head coil. T1-weighted high-resolution magnetization-prepared rapid-acquisition gradient echo (MPRAGE; 0.8 x 0.8 x 0.8 mm$^3$ voxels; TR/TE/TI=1600/3.87/950 ms; flip angle=15°) anatomical scans were collected. The medial temporal lobe subregions were segmented using an automatic pipeline, ASHS–T1 (Xie et al., 2019)[1]. This technique uses a multi-atlas label fusion approach (Wang et al., 2013) together with a tailored segmentation protocol to take into account anatomical variability in MTL cortex. It also explicitly labels dura, which has similar appearance to gray matter in T1-weighted MRI, resulting in more accurate segmentation of MTL cortex compared to other T1-MRI segmentation pipelines, such as FreeSurfer (Xie et al., 2019). In addition to a volume measure of the hippocampus, we obtained measures of mean cortical thickness in the following regions-of-interest (ROIs): entorhinal cortex, perirhinal cortex subregions Brodmann areas 35 and 36 (BA35 and BA36), and parahippocampal cortex, using a graph-based multi-template thickness analysis pipeline taking the ASHS-T1 automatic segmentation as input (Xie et al., 2017). We conducted five regression analyses, one for each ROI, examining the relationship between its thickness/volume and temporal discounting rate. Measures were averaged across hemispheres. Age, gender, and years of education were entered as covariates of no interest. For

---

[1] ASHS-T1 is a follow-up project to ASHS (Automatic Segmentation of Hippocampal Subfields) which is used to label hippocampal subfields and MTL cortical subregions in T2-MRI. It was recently extended to T1-MRI and modified to label MTL cortical subregions and partition hippocampus into anterior and posterior regions, since hippocampal subfields are not generally distinguishable in T1-MRI.

the hippocampal volume analyses, we included the (square root-transformed) intracranial volume as an additional covariate, since volume measures are more biased by total intracranial volume than mean cortical thickness measures are (Barnes et al., 2010). In the regressions examining BA35 and BA36, one participant was excluded because segmentation in these regions did not pass quality control in either hemisphere. In the few cases ($n = 7$) where segmentation could be performed on one side but image quality was inadequate on the other, the mean was replaced with the thickness measure from the available side. All of the above analyses were done with the temporal discounting rate $k$ and the risk tolerance parameter $\alpha$ as dependent variables. In addition, we examined the relationship between our chosen MTL ROIs and the executive function composite and episodic memory composite measures.

If we found that structural measures of any of the MTL ROIs were significantly associated with discount rate, we planned to conduct a mediation analysis (Baron & Kenny, 1986), to see if that association was mediated by episodic memory ability. We used a Sobel test (Sobel, 1982) to test for the presence of an indirect effect (specifically, that MTL structural integrity influences discount rate via episodic memory ability), and we estimated the proportion of the total effect that could be explained through the indirect effect, along with a 95% confidence interval (Hicks & Tingley, 2011).

## 3. Results

3.1. Episodic memory, but not executive function, is associated with temporal discounting

100 older adults completed the National Alzheimer's Coordinating Center Uniform Data Set neuropsychological testing battery, as well as an intertemporal choice task and a risky choice

task. See Table 1 for sample characteristics. In all analyses reported below, age, gender, and years of education were entered as covariates, and partial correlation coefficients are reported.

Episodic memory, but not executive function, was associated with temporal discounting rate. When entering both the episodic memory composite score and executive function composite score as independent variables in a regression predicting temporal discounting rate, better performance on episodic memory tasks was associated with reduced temporal discounting ($r$ = -0.32; $p$ = 0.002; $n$ = 96; Fig. 1A), while the association between executive function and temporal discounting was not significant, and in the positive direction ($r$ = 0.08; $p$ = 0.461; Fig. 1B). That is, individuals with better episodic memory tended to discount delayed rewards less, even after controlling for variance that could be explained by executive function. When conducting separate regressions for episodic memory and executive function (still controlling for age, gender, and years of education), there was a significant association between episodic memory and temporal discounting ($r$ = -0.30; $p$ = 0.003; $n$ = 100), but not between executive function and temporal discounting ($r$ = -0.04; $p$ = 0.693; $n$ = 96).

The association between episodic memory and temporal discounting was not driven by any specific test included in the composite score. The episodic memory composite score included the Word List Memory Delayed Recall, Craft Story Delayed Recall, and Benson Complex Figure Delayed Recall tests. All memory test scores (Word List Memory Delayed Recall: $r$ = -0.32; $p$ = 0.001; Craft Story Delayed Recall: $r$ = -0.21; $p$ = 0.040; Benson Complex Figure Delayed Recall: $r$ = -0.26; $p$ = 0.009) predicted temporal discounting to a comparable degree. Neither of the individual executive function measures was associated with temporal discounting (Digit Span Backwards: $r$ = -0.02; $p$ = 0.813; Trails B − A: $r$ = 0.05; $p$ = 0.653).

The association between episodic memory and temporal discounting did not depend on assumptions about the utility function for money. We also examined the hyperbolic temporal discount rate estimated assuming a non-linear (rather than linear) utility function (see *Materials and methods*) and obtained similar results. Specifically, when entered into the same regression, the episodic memory composite ($r = -0.30$, $p = 0.004$), but not the executive function composite ($r = 0.13$; $p = 0.217$), was significantly correlated with temporal discounting. We obtained similar results with separate regressions for episodic memory ($r = -0.26$, $p = 0.010$) and executive function ($r = 0.02$; $p = 0.840$).

Finally, the association between episodic memory measures and temporal discounting was specific to discounting, and did not extend to decision tendencies in the risk domain. From our risky choice task, we derived a measure of an individual's risk tolerance by assuming a power function for utility and estimating a risk tolerance parameter $\alpha$. There was no relationship between the episodic memory composite score and risk tolerance or between executive function and risk tolerance, whether both composite scores were entered as predictors in the regression together (episodic memory $r = 0.07$, $p = 0.486$; executive function $r = 0.08$; $p = 0.466$) or separately (episodic memory $r = 0.10$, $p = 0.350$; executive function $r = 0.11$; $p = 0.297$).

**Table 1.**

Characteristics of participants ($n$ = 100)

| Characteristic | Mean, SD, Range (or %) |
|---|---|
| Age | 72.01 (6.82, 58-93) |
| Gender | 58% Female, 42% Male |
| Race | 62% White, 36% Black, 2% Multi-racial |
| Years of education | 15.96 (2.82, 9-20) |
| Diagnosis | 74% Cognitively normal; 26% MCI |
| Cognitive measures: Raw scores | Mean, SD, Range |
| Word List Memory Delayed Recall | 7.5 (2.61, 0-10) |
| Craft Story Delayed Recall | 13.63 (5.90, 0-23) |
| Benson Complex Figure Delayed Recall | 9.93 (4.43, 0-16) |
| Semantic fluency (Animals) | 20.89 (6.03, 9-41) |
| Semantic fluency (Vegetables) | 13.88 (4.71, 0-30) |
| Lexical fluency (F-words) | 15.08 (4.66, 2-26) |
| Lexical fluency (L-words) | 13.78 (4.15, 5-25) |
| Digit Span Task (Backwards) number correct | 6.73 (2.09, 2-13) |
| Trail Making Test (Part B minus Part A) RT | 46.61 (25.48, 10-153)* |

Note: MCI = mild cognitive impairment; RT = reaction time. *$n$ = 4 participants excluded for not completing Trail Making Test Part B in the allotted time ($n$ = 1) or for having a reaction time on Trail Making Test Part B that was more than 3 SD > mean ($n$ = 3; 280 s, 300 s, and 300 s).

**Fig. 1.** Associations between cognitive measures and temporal discounting. The episodic memory composite measure (A) is significantly correlated with temporal discounting rate: people with better memory are more patient for future rewards. The residual discount rate after adjusting for age, gender, years of education, and executive function is shown. Performance on the executive function composite measure (B) is not significantly correlated with discount rate. The residual discount rate after adjusting for age, gender, years of education, and episodic memory is shown. MCI = Mild Cognitive Impairment. Linear fits to the data along with 95% confidence intervals are overlaid on the plots.

3.2. Semantic, but not lexical, fluency is associated with temporal discounting

As a secondary test of the roles of temporal lobe-mediated vs. frontal lobe-mediated cognitive processes in temporal discounting, we examined the association between verbal fluency measures and temporal discounting. Semantic fluency, but not lexical fluency, was associated with temporal discounting rate. When entering both semantic fluency and lexical fluency as independent variables in a regression predicting temporal discounting rate, better performance on semantic fluency was associated with reduced temporal discounting ($r = -0.23$; $p = 0.026$; $n = 100$; Fig. 2A), while the association between lexical fluency and temporal discounting was not significant ($r = 0.09$; $p = 0.411$; Fig. 2B). Thus, even when controlling for the search and retrieval processes involved in verbal fluency tasks in general, people with better semantic fluency, which requires accessing semantic knowledge, were more likely to choose larger, delayed rewards. When conducting separate regressions for semantic fluency and lexical fluency (still controlling for age, gender, and years of education), there was a significant association between semantic fluency and temporal discounting ($r = -0.22$; $p = 0.032$), but not between lexical fluency and temporal discounting ($r = -0.06$; $p = 0.584$).

These results remained qualitatively similar when assuming a non-linear utility function for money (in the same regression, semantic fluency: $r = -0.31$; $p = 0.002$; lexical fluency: $r = 0.18$; $p = 0.077$; in separate regressions, semantic fluency: $r = -0.25$; $p = 0.013$; lexical fluency: $r = 0.0004$; $p = 0.997$). Moreover, neither fluency measure was associated with risk tolerance (in the same regression, semantic fluency: $r = -0.15$; $p = 0.137$; lexical fluency: $r = 0.17$; $p = 0.094$; in separate regressions, semantic fluency: $r = -0.07$; $p = 0.521$; lexical fluency: $r = 0.10$; $p = 0.314$).

**Fig. 2.** Associations between verbal fluency measures and temporal discounting. Semantic

fluency (A) is significantly correlated with temporal discounting rate: people with better

semantic fluency are more patient for future rewards. The residual discount rate after adjusting

for age, gender, years of education, and lexical fluency is shown. Performance on lexical fluency

(B) is not significantly correlated with discount rate. The residual discount rate after adjusting for

age, gender, years of education, and semantic fluency is shown. MCI = Mild Cognitive

Impairment. Linear fits to the data along with 95% confidence intervals are overlaid on the plots.

3.3. Temporal discounting is increased in mild cognitive impairment

Perhaps not surprisingly given the relationship between declarative memory and temporal discounting, there was a significant effect of diagnosis (mild cognitive impairment vs. cognitively normal) on discount rate ($F_{(1,98)} = 5.01$; $p = 0.028$), with MCI participants displaying increased temporal discounting overall. This provides additional support for the finding that declarative memory and temporal discounting are associated, since MCI participants performed significantly worse on all episodic memory measures (Word List Memory Delayed Recall: $F_{(1,98)} = 105.21$, $p < 0.001$; Craft Story Delayed Recall: $F_{(1,98)} = 90.37$, $p < 0.001$; Benson Complex Figure Delayed Recall: $F_{(1,98)} = 86.03$, $p < 0.001$) and semantic fluency measures (Animal fluency: $F_{(1,98)} = 23.18$; $p < 0.001$; Vegetable fluency: $F_{(1,98)} = 39.19$; $p < 0.001$) compared to cognitively normal individuals.

Within each diagnosis subgroup of the sample, there was no significant association between episodic memory and discount rate (controlling for executive function: cognitively normal, $n = 74$: $r = -0.07$, $p = 0.553$; MCI, $n = 22$: $r = -0.45$; $p = 0.064$), but the correlation coefficients suggest that the relationship was stronger in the MCI group. There was also no significant association between semantic fluency and discount rate (controlling for lexical fluency: cognitively normal, $n = 74$: $r = -0.13$, $p = 0.270$; MCI, $n = 26$: $r = -0.06$; $p = 0.779$), but here the effect sizes were comparable between the two groups. With the exception of Digit Span Backwards ($F_{(1,98)} = 3.09$, $p = 0.082$), MCI participants performed significantly worse on executive function (Trails B-A: $F_{(1,94)} = 7.09$, $p = 0.009$), and lexical fluency (F-word fluency: $F_{(1,98)} = 10.05$, $p = 0.002$; L-word fluency: $F_{(1,98)} = 7.82$, $p = 0.006$) as well. MCI participants were not more risk-seeking or risk-averse overall ($F_{(1,98)} = 0.20$, $p = 0.652$).

3.4. Entorhinal cortical thickness is associated with temporal discounting

First, we confirmed that structural integrity of the medial temporal lobe was associated

with episodic memory, but not executive function, in our dataset. A large subset of participants

($n = 92$) had structural MRI data. We examined associations with structural measures from five

medial temporal lobe subregions: cortical thickness in entorhinal cortex, BA35, BA36, and

parahippocampal cortex, and hippocampal volume. Structural measures from all of these medial

temporal lobe regions were associated with episodic memory (entorhinal cortex $r = 0.54$, $p <$

$0.001$; BA35 $r = 0.41$, $p < 0.001$; BA36 $r = 0.28$; $p = 0.009$; parahippocampal cortex $r = 0.24$, $p$

$= 0.024$; hippocampus $r = 0.57$; $p < 0.001$). In contrast, none of the medial temporal lobe

structural measures were associated with executive function (entorhinal cortex $r = 0.07$, $p =$

$0.543$; BA35 $r = 0.10$, $p = 0.385$; BA36 $r = 0.08$; $p = 0.466$; parahippocampal cortex $r = 0.06$, $p$

$= 0.586$; hippocampus $r = 0.10$; $p = 0.367$).

Given the association between behavioral measures of episodic memory and temporal

discounting, next we conducted an exploratory analysis relating structural integrity in the medial

temporal lobe to temporal discounting. Temporal discounting was associated with mean cortical

thickness in the entorhinal cortex ($r = -0.28$; $p = 0.008$; $n = 92$; Fig. 3), but the association did

not reach significance with cortical thickness in BA35 ($r = -0.09$; $p = 0.382$; $n = 91$), BA36 ($r = -$

$0.18$; $p = 0.094$; $n = 91$), or parahippocampal cortex ($r = -0.17$; $p = 0.116$; $n = 92$), or with the

volume of the hippocampus ($r = -0.06$, $p = 0.603$; $n = 92$). The association between entorhinal

cortex and discount rate did not depend on our choice of utility function, as it remained

significant when estimating the discount function assuming a non-linear utility function ($r = -$

$0.24$; $p = 0.022$). Cortical thickness in entorhinal cortex was also related specifically to

discounting, and not to risk tolerance ($r = 0.05$, $p = 0.642$; see Table 2 for full list of partial

correlation coefficients). In addition, this relationship does not depend on our inclusion of MCI participants, as the association between entorhinal cortical thickness and discount rate remained significant even when including diagnosis (MCI vs. cognitively normal) as an additional covariate ($r = -0.21$; $p = 0.044$). However, when examining the two subsamples separately, the association was stronger in the MCI group (MCI, $n = 24$: $r = -0.44$; $p = 0.047$; cognitively normal, $n = 68$: $r = -0.08$; $p = 0.506$). This pattern resembles what was observed in the behavioral data, and most likely reflects increased variance in MTL atrophy in the MCI subgroup.

Episodic memory ability partially mediated the relationship between entorhinal cortical thickness and temporal discounting. When entorhinal cortical thickness and episodic memory were entered in the same regression to predict discount rate (with age, gender, and years of education as covariates), we found that entorhinal cortical thickness no longer predicted discounting ($r = -0.14$; $p = 0.249$). A Sobel test revealed that there was a significant indirect effect ($t = 2.07$; $p = 0.039$), whereby entorhinal cortical thickness influences temporal discounting through its influence on episodic memory. Approximately 49% (bootstrapped 95% CI: [28.45%, 179.11%]) of the total effect of entorhinal cortical thickness on temporal discounting could be explained through this indirect effect.

Finally, to investigate if entorhinal cortical thickness was associated with temporal discounting above and beyond the other medial temporal lobe thickness/volume measures, we entered all five structural measures into the same regression, with intracranial volume, age, gender, and years of education as regressors of no interest. Even when controlling for cortical thickness in parahippocampal cortex, BA35, BA36, and the volume of the hippocampus, entorhinal cortical thickness still predicted discount rate ($r = -0.26$; $p = 0.018$), and none of the

other measures did (parahippocampal cortex $r = -0.13$; $p = 0.249$; BA35 $r = 0.12$; $p = 0.271$;

BA36 $r = -0.14$; $p = 0.209$; hippocampus $r = 0.10$; $p = 0.351$).

**Fig. 3.** Association between entorhinal cortical thickness (in millimeters) and temporal discounting rate (*n* = 92). The entorhinal cortex was the only one of our medial temporal lobe ROIs that was significantly and robustly associated with temporal discounting rate: people with more entorhinal cortical thickness were more patient for future rewards. The residual plot after adjusting for age, gender, and years of education is shown. MCI = Mild Cognitive Impairment. Linear fits to the data along with 95% confidence intervals are overlaid on the plots.

**Table 2.**
Associations between cortical thickness in regions of interest and choice measures.

| Region | Discount rate | Discount rate (utility-transformed) | Risk tolerance |
|---|---|---|---|
| Entorhinal cortex | **-0.28\*\*** | **-0.24\*** | 0.05 |
| BA36 | -0.18 | **-0.23\*** | -0.13 |
| BA35 | -0.09 | -0.09 | -0.02 |
| Parahippocampal cortex | -0.17 | -0.15 | 0.03 |
| Hippocampus (volume) | -0.06 | -0.03 | 0.11 |

Note: Partial correlation coefficients, controlling for age, gender, years of education, and intracranial volume (for hippocampus only) are shown. *p<0.05, **p<0.01 (uncorrected for multiple comparisons).

**4. Discussion**

Here we found in a diverse group of cognitively normal and MCI older adults that better declarative memory ability was associated with reduced temporal discounting, or a relatively greater preference for larger, delayed rewards. This relationship held even when controlling for executive function abilities that may impact memory retrieval. This finding was not limited to *episodic* memory, since semantic fluency was also correlated with temporal discounting, even when adjusting for lexical fluency, another task with similar executive demands. The specificity of these associations is underlined by the findings that executive function and lexical fluency were unrelated to temporal discounting, and declarative memory was unrelated to risk preferences. In an exploratory analysis relating cortical thickness in the MTL to temporal discounting, we found that thickness in the entorhinal cortex was associated with temporal discounting, and that the relationship between entorhinal cortical thickness and discount rate was partially mediated by episodic memory ability.

Declarative memory is one of the first cognitive abilities to decline as individuals age (Buckner, 2004; Nyberg, 2016; Nyberg, Lövdén, Riklund, Lindenberger, & Bäckman, 2012), and this decline is associated with degeneration in the MTL. Here we found that behavioral markers of MTL function (episodic memory retrieval and semantic fluency) and cortical thickness in the entorhinal cortex were associated with discounting in older adults. To our knowledge, this is the first study to link declarative memory abilities with temporal discounting. Previous studies of intertemporal choice in individuals with MCI and Alzheimer's Disease have yielded inconsistent results (Chiong et al., 2016; Coelho et al., 2016; Lebreton et al., 2013; Lindbergh et al., 2014). Even in "normal" aging, however, there are large individual differences in the extent and rate of cognitive decline, suggesting that a correlational design might be more

appropriate in this population. One previous correlational study conducted with healthy older adults found that global cognition was associated with temporal discounting, but episodic and semantic memory composite scores were not (Boyle et al., 2012). Another study that investigated the relationship between episodic memory ability and temporal discounting in older adults also yielded null results (Seinstra et al., 2015). These two studies, however, included only cognitively normal older adults. Furthermore, one of these studies (Seinstra et al., 2015) focused primarily on associative recognition and autobiographical memory recall, which may be less impacted by aging compared to free recall and episodic retrieval, respectively (Howard et al., 2006; Schroots, Van Dijkum, & Assink, 2004). In the other study (Boyle et al., 2012), the episodic memory measure did not distinguish between immediate and delayed recall. Here, by leveraging a large and well-characterized sample with substantial variability in episodic and semantic memory retrieval ability, including individuals with MCI, and using measures that emphasized delayed free recall and semantic memory, we detected a significant association between temporal discounting and memory function.

Another contribution of the current study is linking neural correlates of memory with temporal discounting. Structural integrity in the MTL has previously been associated with temporal discounting in adolescents (Pehlivanova et al., 2018) and young and middle-aged adults (Owens et al., 2017). Here we found in older adults, who are more likely to have atrophy in MTL, that entorhinal cortical thickness was associated with temporal discounting. Perhaps surprisingly, hippocampal volume was not associated with temporal discounting, despite the importance of this region for the formation of context-rich episodic memories. The entorhinal cortex, however, was the region that was associated with temporal discounting in one previous study (Owens et al., 2017). It is also one of the earliest deposition sites for the neurofibrillary

tangle pathology associated with aging and Alzheimer's Disease (Braak & Braak, 1991). The exact mechanism by which the entorhinal cortex supports future-oriented choice warrants further study. One possibility is that given its role as a "relay station" between hippocampus and prefrontal cortex (Agster & Burwell, 2009; Apergis-Schoute, Pinto, & Paré, 2006), the entorhinal cortex facilitates the modulation of valuation by memory retrieval at the time of choice. Another possibility is that its involvement in memory for timing or duration (Lositsky et al., 2016; Montchal, Reagh, & Yassa, 2019) influences prospective duration estimates. While our analysis of MTL subregions was exploratory and preliminary, our findings are strengthened by our use of state-of-the-art methods for segmentation (which included rigorous quality checks) in a well-characterized group of older adults with variability in the structural integrity of these regions.

The inclusion of MCI participants was critical here, as there was no significant correlation between declarative memory measures and temporal discounting in the participants who were classified as cognitively normal. This reflects a limited range of impairment in the cognitively normal group, and may explain why this association has not been observed in the literature before. While we cannot definitively rule out the possibility that MCI participants have higher discount rates for a reason other than their decline in episodic memory (e.g., their diagnosis), a few results argue against this possibility. First, MCI participants were also impaired on executive function tasks, but it was declarative memory specifically that correlated with temporal discounting, not global cognition. Second, within the MCI group, the partial correlation coefficient between episodic memory and temporal discount rate was high ($r = -0.45$); this continuous association between memory ability and temporal discounting would be unlikely if it was the diagnosis itself that changed preferences. Finally, the relationship between entorhinal cortical thickness and discount rate was significant even when controlling for diagnosis. Future

studies using episodic and semantic memory measures that show sufficient variance in young and cognitively normal older adults will shed additional light on the generalizability of this relationship. Nevertheless, the finding that MCI participants, a group that is *defined* by their memory deficits, show increased temporal discounting is evidence for the hypothesis that episodic memory drives individual differences in intertemporal preferences.

Previous studies have proposed that it is *episodic* memory and episodic future thinking specifically that underlie individual differences in temporal discounting (Peters & Büchel, 2010), since a richer picture of one's personal future might lead that person to view the future as more concrete, certain, and closer in time (Fujita, Trope, Liberman, & Levin-Sagi, 2006). However, here we found that a measure of *semantic* memory – semantic fluency – also correlated with discount rate. While semantic fluency taps into a few cognitive processes on its own, it is differentiated from lexical fluency by its reliance on the retrieval of semantic knowledge and its dependence on intact temporal lobe function (Gourovitch et al., 2000; Henry & Crawford, 2004). Semantic fluency is also more likely than lexical fluency to engage a strategy of "clustering" items into categories rather than "switching" between categories (Troyer et al., 1998). We were unable to analyze trial-level data to examine whether the association between semantic fluency and discounting is driven by the use of the "clustering" strategy or by the use of semantic knowledge itself. Intriguingly, however, prior studies have found evidence of links between semantic memory and temporal discounting. Individuals with semantic dementia show increased temporal discounting, even relative to individuals with Alzheimer's Disease (Chiong et al., 2016). They also show impaired episodic future thinking despite having intact autobiographical memory retrieval (Irish, Addis, Hodges, & Piguet, 2012; Irish, Addis, Hodges, Piguet, et al., 2012), and this impairment is mediated by reduction in gray matter volume in semantic memory

regions (e.g., temporal pole, inferior temporal gyrus; Irish et al., 2012a). In addition, individuals with hippocampal damage, who have impaired episodic memory but intact semantic memory (Giovagnoli, Erbetta, & Bugiani, 2001; Schmolck, Kensinger, Corkin, & Squire, 2002), display temporal discounting rates that are no different from those of normal controls (Kwan et al., 2012), suggesting that semantic knowledge may be sufficient for deciding about the future. Since we saw a relationship between discounting and both semantic and episodic memory tasks here, however, another possibility is that both of these tasks tap into a shared process that is integral to making patient intertemporal choices.

In contrast, performance on standard measures of executive function (Digit Span Backwards, Trails B, and lexical fluency) was not associated with temporal discounting. In fact, when controlling for episodic memory, the relationship between executive function and temporal discounting was in the positive direction (with better executive function being associated with higher discounting). Thus, our findings provide key evidence that declarative memory processes are a more important contributor to future-directed decision-making than executive function. Although there is a well-documented association between temporal discounting and fluid intelligence (Burks, Carpenter, Goette, & Rustichini, 2009; Kable et al., 2017; Shamosh et al., 2008), in principle, it could be that episodic memory underpins this association. Recent research has shown that the strong correlation between working memory and general intelligence (Ackerman, Beier, & Boyle, 2005) may be driven by individual differences in episodic memory processes, such as search and retrieval (Healey, Crutchley, & Kahana, 2014; Mogle, Lovett, Stawski, & Sliwinski, 2008; Unsworth, Brewer, & Spillers, 2009). Furthermore, the most successful manipulations of temporal discounting to date involve activating episodic memory circuitry by encouraging people to imagine future events (Bulley, Henry, & Suddendorf, 2016).

On the other hand, taxing frontal executive processes does not necessarily increase impulsive choice, but rather, might just decrease choice consistency (i.e., the ability to maintain consistent preferences across trials; Franco-Watkins et al., 2010; Olschewski et al., 2018). Therefore, executive function may play some role in intertemporal decision-making, such as helping individuals to integrate costs and benefits or maintain consistency, but individual differences in temporal discounting rate are more likely to be explained by declarative memory.

We found no association between either executive function or declarative memory and risk tolerance. Previous research has linked cognitive abilities with risk preferences, including in older adults (Boyle, Yu, Buchman, Laibson, & Bennett, 2011). Specifically, people who are more educated (Donkers et al., 1999), have higher intelligence (Burks et al., 2009), and have better global cognition (Boyle et al., 2011; Frederick, 2005) are more risk-seeking, especially when risky choices have higher expected value. It is possible that other cognitive or affective abilities not assessed here are preferentially involved in decisions under risk.

The results of the current study elucidate the inconsistent findings related to aging and intertemporal decision-making. We did not find a significant relationship between age and temporal discounting rate, whether looking at our sample overall ($r = 0.06$, $p = 0.56$), or just within the cognitively normal  ($r = 0.03$, $p = 0.79$) or MCI ($r = 0.17$, $p = 0.41$) groups. This is consistent with previous studies showing that sound economic decision-making remains intact in older adults (Li, Baldassi, Johnson, & Weber, 2013; Li et al., 2015), and that changes in economic decision-making with aging are more likely to be underpinned by changes in cognitive abilities (Henninger, Madden, & Huettel, 2010). Our results suggest specifically that temporal discounting may increase with age to the extent that declarative memory declines. However, we cannot draw this conclusion from our cross-sectional investigation. Longitudinal work has shown

that changes in cognitive function in older adults are associated with concomitant changes in temporal discounting (James, Boyle, Yu, Han, & Bennett, 2015). Future research, perhaps with the same cohort used here, will reveal whether episodic and/or semantic memory decline has a causal influence on intertemporal decision-making.

In sum, the current study sheds light on the cognitive and neural mechanisms underlying individual differences in temporal discounting. It also contributes to our understanding of decision-making in the context of aging. These findings may aid in the development of interventions to promote more economic patience, especially as cognition declines.

**Acknowledgements**

We would like to thank Laura Saad, Arun Pilania, Joseph Harrison, and Jacqueline Lane for assistance with data collection.

**Funding**

KML was supported by a National Research Service Award (F32-AG-054032-02) from the National Institute on Aging. This project received funding from the TIAA Institute and Wharton School's Pension Research Council/Boettner Center.

**References**

Ackerman, P. L., Beier, M. E., & Boyle, M. O. (2005). Working Memory and Intelligence: The Same or Different Constructs? *Psychological Bulletin*, *131*(1), 30–60. https://doi.org/10.1037/0033-2909.131.1.30

Addis, D. R., Wong, A. T., & Schacter, D. L. (2007). Remembering the past and imagining the

future: Common and distinct neural substrates during event construction and elaboration. *Neuropsychologia*, *45*(7), 1363–1377. https://doi.org/10.1016/j.neuropsychologia.2006.10.016

Agster, K. L., & Burwell, R. D. (2009). Cortical efferents of the perirhinal, postrhinal, and entorhinal cortices of the rat. *Hippocampus*, *19*(12), 1159–1186. https://doi.org/10.1002/hipo.20578

Apergis-Schoute, J., Pinto, A., & Paré, D. (2006). Ultrastructural organization of medial prefrontal inputs to the rhinal cortices. *European Journal of Neuroscience*, *24*(1), 135–144. https://doi.org/10.1111/j.1460-9568.2006.04894.x

Barnes, J., Ridgway, G. R., Bartlett, J., Henley, S. M. D., Lehmann, M., Hobbs, N., … Fox, N. C. (2010). Head size, age and gender adjustment in MRI studies: a necessary nuisance? *NeuroImage*, *53*(4), 1244–1255. https://doi.org/10.1016/j.neuroimage.2010.06.025

Baron, R. M., & Kenny, D. A. (1986). The moderator–mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*, *51*(6), 1173–1182. https://doi.org/10.1037/0022-3514.51.6.1173

Benoit, R. G., Gilbert, S. J., & Burgess, P. W. (2011). A neural mechanism mediating the impact of episodic prospection on farsighted decisions. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *31*(18), 6771–6779. https://doi.org/10.1523/JNEUROSCI.6559-10.2011

Benton, A. L., & Hamsher, K. (1976). *Multilingual Aphasia Examination manual*. Iowa City, IA: University of Iowa.

Bickel, W. K., Yi, R., Kowal, B. P., & Gatchalian, K. M. (2008). Cigarette smokers discount past

and future rewards symmetrically and more than controls: is discounting a measure of

impulsivity? *Drug and Alcohol Dependence*, *96*(3), 256–262.

https://doi.org/10.1016/j.drugalcdep.2008.03.009

Boyle, P. A., Yu, L., Buchman, A. S., Laibson, D. I., & Bennett, D. A. (2011). Cognitive

function is associated with risk aversion in community-based older persons. *BMC

Geriatrics*, *11*(1), 53. https://doi.org/10.1186/1471-2318-11-53

Boyle, P. A., Yu, L., Segawa, E., Wilson, R. S., Buchman, A. S., Laibson, D. I., & Bennett, D.

A. (2012). Association of cognition with temporal discounting in community based older

persons. *BMC Geriatrics*, *12*(1), 48. https://doi.org/10.1186/1471-2318-12-48

Braak, H., & Braak, E. (1991). Neuropathological stageing of Alzheimer-related changes. *Acta

Neuropathologica*, *82*(4), 239–259. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/1759558

Buckner, R. L. (2004). Memory and Executive Function in Aging and AD. *Neuron*, *44*(1), 195–

208. https://doi.org/10.1016/j.neuron.2004.09.006

Bulley, A., Henry, J., & Suddendorf, T. (2016). Prospection and the present moment: The role of

episodic foresight in intertemporal choices between immediate and delayed rewards. *Review

of General Psychology*, *20*(1), 29–47. https://doi.org/10.1037/gpr0000061

Burks, S. V, Carpenter, J. P., Goette, L., & Rustichini, A. (2009). Cognitive skills affect

economic preferences, strategic behavior, and job attachment. *Proceedings of the National

Academy of Sciences of the United States of America*, *106*(19), 7745–7750.

https://doi.org/10.1073/pnas.0812360106

Chiong, W., Wood, K. A., Beagle, A. J., Hsu, M., Kayser, A. S., Miller, B. L., & Kramer, J. H.

(2016). Neuroeconomic dissociation of semantic dementia and behavioural variant

frontotemporal dementia. *Brain : A Journal of Neurology*, *139*(Pt 2), 578–587.

https://doi.org/10.1093/brain/awv344

Coelho, S., Guerreiro, M., Chester, C., Silva, D., Maroco, J., Paglieri, F., & de Mendonça, A.

(2016). Delay discounting in mild cognitive impairment. *Journal of Clinical and*

*Experimental Neuropsychology*, 1–11. https://doi.org/10.1080/13803395.2016.1226269

Craft, S., Newcomer, J., Kanne, S., Dagogo-Jack, S., Cryer, P., Sheline, Y., … Alderson, A.

(n.d.). Memory improvement following induced hyperinsulinemia in Alzheimer's disease.

*Neurobiology of Aging*, *17*(1), 123–130. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/8786794

Davidson, P. S. R., Gao, F. Q., Mason, W. P., Winocur, G., & Anderson, N. D. (2008). Verbal

fluency, Trail Making, and Wisconsin Card Sorting Test performance following right

frontal lobe tumor resection. *Journal of Clinical and Experimental Neuropsychology*, *30*(1),

18–32. https://doi.org/10.1080/13803390601161166

Donkers, B., Melenberg, B., van Soest, A., Donkers, B., Melenberg, B., & van Soest, A. (1999).

Estimating Risk Attitudes Using Lotteries; A Large Sample Approach. Retrieved from

https://econpapers.repec.org/paper/tiutiucen/94cd10d5-d110-4d60-87f2-086013112b71.htm

Franco-Watkins, A. M., Rickard, T. C., & Pashler, H. (2010). Taxing Executive Processes Does

Not Necessarily Increase Impulsive Decision Making. *Experimental Psychology*, *57*(3),

193–201. https://doi.org/10.1027/1618-3169/a000024

Frederick, S. (2005). Cognitive Reflection and Decision Making. *Journal of Economic*

*Perspectives*, *19*(4), 25–42. https://doi.org/10.1257/089533005775196732

Frost, R., & McNaughton, N. (2017). The neural basis of delay discounting: A review and

preliminary model. *Neuroscience & Biobehavioral Reviews*, *79*, 48–65.

https://doi.org/10.1016/J.NEUBIOREV.2017.04.022

Fujita, K., Trope, Y., Liberman, N., & Levin-Sagi, M. (2006). Construal levels and self-control. *Journal of Personality and Social Psychology*, *90*(3), 351–367. Retrieved from http://psycnet.apa.orgjournals/psp/90/3/351

Gerton, B. K., Brown, T. T., Meyer-Lindenberg, A., Kohn, P., Holt, J. L., Olsen, R. K., & Berman, K. F. (2004). Shared and distinct neurophysiological components of the digits forward and backward tasks as revealed by functional neuroimaging. *Neuropsychologia*, *42*(13), 1781–1787. https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2004.04.023

Giovagnoli, A. R., Erbetta, A., & Bugiani, O. (2001). Preserved Semantic Access in Global Amnesia and Hippocampal Damage. *The Clinical Neuropsychologist*, *15*(4), 508–515. https://doi.org/10.1076/clin.15.4.508.1889

Gourovitch, M. L., Kirkby, B. S., Goldberg, T. E., Weinberger, D. R., Gold, J. M., Esposito, G., … Berman, K. F. (2000). A comparison of rCBF patterns during letter and semantic fluency. *Neuropsychology*, *14*(3), 353–360. https://doi.org/10.1037/0894-4105.14.3.353

Green, L., & Myerson, J. (2004). A discounting framework for choice with delayed and probabilistic rewards. *Psychological Bulletin*, *130*(5), 769–792. https://doi.org/10.1037/0033-2909.130.5.769

Greenberg, D. L., Keane, M. M., Ryan, L., & Verfaellie, M. (2009). Impaired category fluency in medial temporal lobe amnesia: the role of episodic memory. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *29*(35), 10900–10908. https://doi.org/10.1523/JNEUROSCI.1202-09.2009

Groeger, J. A., Field, D., & Hammond, S. M. (1999). Measuring Memory Span. *International Journal of Psychology*, *34*(5–6), 359–363. https://doi.org/10.1080/002075999399693

Healey, M. K., Crutchley, P., & Kahana, M. J. (2014). Individual differences in memory search and their relation to intelligence. *Journal of Experimental Psychology: General*, *143*(4), 1553–1569. https://doi.org/10.1037/a0036306

Henninger, D. E., Madden, D. J., & Huettel, S. A. (2010). Processing speed and memory mediate age-related differences in decision making. *Psychology and Aging*, *25*(2), 262–270. https://doi.org/10.1037/a0019096

Henry, J. D., & Crawford, J. R. (2004). A Meta-Analytic Review of Verbal Fluency Performance Following Focal Cortical Lesions. *Neuropsychology*, *18*(2), 284–295. https://doi.org/10.1037/0894-4105.18.2.284

Hicks, R., & Tingley, D. (2011). Causal Mediation Analysis. *The Stata Journal: Promoting Communications on Statistics and Stata*, *11*(4), 605–619. https://doi.org/10.1177/1536867X1201100407

Howard, M. W., Bessette-Symons, B., Zhang, Y., & Hoyer, W. J. (2006). Aging selectively impairs recollection in recognition memory for pictures: Evidence from modeling and receiver operating characteristic curves. *Psychology and Aging*, *21*(1), 96–106. https://doi.org/10.1037/0882-7974.21.1.96

Hu, X., Uhle, F., Fliessbach, K., Wagner, M., Han, Y., Weber, B., & Jessen, F. (2017). Reduced future-oriented decision making in individuals with subjective cognitive decline: A functional MRI study. *Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring*, *6*, 222–231. https://doi.org/10.1016/j.dadm.2017.02.005

Irish, M., Addis, D. R., Hodges, J. R., & Piguet, O. (2012). Exploring the content and quality of episodic future simulations in semantic dementia. *Neuropsychologia*, *50*(14), 3488–3495. https://doi.org/10.1016/j.neuropsychologia.2012.09.012

Irish, M., Addis, D. R., Hodges, J. R., Piguet, O., Addis, D., Cheng, T., … Smith, S. (2012). Considering the role of semantic memory in episodic future thinking: evidence from semantic dementia. *Brain : A Journal of Neurology*, *135*(Pt 7), 2178–2191. https://doi.org/10.1093/brain/aws119

James, B. D., Boyle, P. A., Yu, L., Han, S. D., & Bennett, D. A. (2015). Cognitive decline is associated with risk aversion and temporal discounting in older adults without dementia. *PloS One*, *10*(4), e0121900. https://doi.org/10.1371/journal.pone.0121900

Jernigan, T. L., Archibald, S. L., Berhow, M. T., Sowell, E. R., Foster, D. S., & Hesselink, J. R. (1991). Cerebral structure on MRI, Part I: Localization of age-related changes. *Biological Psychiatry*, *29*(1), 55–67. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/2001446

Kable, J. W., Caulfield, M. K., Falcone, M., McConnell, M., Bernardo, L., Parthasarathi, T., … Lerman, C. (2017). No Effect of Commercial Cognitive Training on Brain Activity, Choice Behavior, or Cognitive Performance. *The Journal of Neuroscience*, *37*(31), 7390–7402. https://doi.org/10.1523/JNEUROSCI.2832-16.2017

Kable, J. W., & Glimcher, P. W. (2007). The neural correlates of subjective value during intertemporal choice. *Nature Neuroscience*, *10*(12), 1625–1633. https://doi.org/10.1038/nn2007

Kortte, K. B., Horner, M. D., & Windham, W. K. (2002). The Trail Making Test, Part B: Cognitive Flexibility or Ability to Maintain Set? *Applied Neuropsychology*, *9*(2), 106–109. https://doi.org/10.1207/S15324826AN0902_5

Kwan, D., Craver, C. F., Green, L., Myerson, J., Boyer, P., & Rosenbaum, R. S. (2012). Future decision-making without episodic mental time travel. *Hippocampus*, *22*(6), 1215–1219. https://doi.org/10.1002/hipo.20981

Lebreton, M., Bertoux, M., Boutet, C., Lehericy, S., Dubois, B., Fossati, P., & Pessiglione, M. (2013). A Critical Role for the Hippocampus in the Valuation of Imagined Outcomes. *PLoS Biology*, *11*(10), e1001684. https://doi.org/10.1371/journal.pbio.1001684

Lempert, K. M., Speer, M. E., Delgado, M. R., & Phelps, E. A. (2017). Positive autobiographical memory retrieval reduces temporal discounting. *Social Cognitive and Affective Neuroscience*. https://doi.org/10.1093/scan/nsx086

Li, Y., Baldassi, M., Johnson, E. J., & Weber, E. U. (2013). Complementary cognitive capabilities, economic decision making, and aging. *Psychology and Aging*, *28*(3), 595–613. https://doi.org/10.1037/a0034172

Li, Y., Gao, J., Enkavi, A. Z., Zaval, L., Weber, E. U., & Johnson, E. J. (2015). Sound credit scores and financial decisions despite cognitive aging. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(1), 65–69. https://doi.org/10.1073/pnas.1413570112

Lindbergh, C. A., Puente, A. N., Gray, J. C., Mackillop, J., & Miller, L. S. (2014). Delay and Probability Discounting as Candidate Markers for Dementia: An Initial Investigation. *Archives of Clinical Neuropsychology*, *29*(7), 651–662. https://doi.org/10.1093/arclin/acu043

Lopez-Guzman, S., Konova, A. B., Louie, K., & Glimcher, P. W. (2018). Risk preferences impose a hidden distortion on measures of choice impulsivity. *PLOS ONE*, *13*(1), e0191357. https://doi.org/10.1371/journal.pone.0191357

Lositsky, O., Chen, J., Toker, D., Honey, C. J., Shvartsman, M., Poppenk, J. L., … Norman, K. A. (2016). Neural pattern change during encoding of a narrative predicts retrospective duration estimates. *ELife*, *5*. https://doi.org/10.7554/eLife.16070

MacKillop, J., Amlung, M. T., Few, L. R., Ray, L. A., Sweet, L. H., & Munafò, M. R. (2011). Delayed reward discounting and addictive behavior: A meta-analysis. *Psychopharmacology*, *216*(3), 305–321. https://doi.org/10.1007/s00213-011-2229-0

Mazur, J. (1987). An adjusting procedure for studying delayed reinforcement. In M. L. Commons, J. E. Mazur, & J. A. Nevin (Eds.), *Quantitative Analyses of Behavior: Vol. 5: The Effect of Delay and of Intervening Events on Reinforcement Value* (pp. 55–73). Hillsdale, NJ: Erlbaum.

McClure, S. M., & Bickel, W. K. (2014). A dual-systems perspective on addiction: contributions from neuroimaging and cognitive training. *Annals of the New York Academy of Sciences*, *1327*, 62–78. https://doi.org/10.1111/nyas.12561

McClure, S. M., Laibson, D. I., Loewenstein, G., & Cohen, J. D. (2004). Separate neural systems value immediate and delayed monetary rewards. *Science*, *306*(5695), 503–507. https://doi.org/10.1126/science.1100907

Meier, S., & Sprenger, C. (2010). Present-Biased Preferences and Credit Card Borrowing. *American Economic Journal: Applied Economics*, *2*(1), 193–210. https://doi.org/10.1257/app.2.1.193

Mogle, J. A., Lovett, B. J., Stawski, R. S., & Sliwinski, M. J. (2008). What's So Special About Working Memory? *Psychological Science*, *19*(11), 1071–1077. https://doi.org/10.1111/j.1467-9280.2008.02202.x

Montchal, M. E., Reagh, Z. M., & Yassa, M. A. (2019). Precise temporal memories are supported by the lateral entorhinal cortex in humans. *Nature Neuroscience*, *22*(2), 284–288. https://doi.org/10.1038/s41593-018-0303-1

Morris, J C, Heyman, A., Mohs, R. C., Hughes, J. P., van Belle, G., Fillenbaum, G., … Clark, C.

(1989). The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part I.
Clinical and neuropsychological assessment of Alzheimer's disease. *Neurology*, *39*(9),
1159–1165. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/2771064

Morris, John C., Weintraub, S., Chui, H. C., Cummings, J., DeCarli, C., Ferris, S., … Kukull, W.
A. (2006). The Uniform Data Set (UDS): Clinical and Cognitive Variables and Descriptive
Data From Alzheimer Disease Centers. *Alzheimer Disease & Associated Disorders*, *20*(4),
210–216. https://doi.org/10.1097/01.wad.0000213865.09806.92

Moscovitch, M., Rosenbaum, R. S., Gilboa, A., Addis, D. R., Westmacott, R., Grady, C., …
Nadel, L. (2005). Functional neuroanatomy of remote episodic, semantic and spatial
memory: a unified account based on multiple trace theory. *Journal of Anatomy*, *207*(1), 35–
66. https://doi.org/10.1111/j.1469-7580.2005.00421.x

Mummery, C. J., Patterson, K., Hodges, J. R., & Wise, R. J. (1996). Generating "tiger" as an
animal name or a word beginning with T: differences in brain activation. *Proceedings.
Biological Sciences*, *263*(1373), 989–995. https://doi.org/10.1098/rspb.1996.0146

Nyberg, L. (2016). Functional brain imaging of episodic memory decline in ageing. *Journal of
Internal Medicine*. https://doi.org/10.1111/joim.12533

Nyberg, Lars, Lövdén, M., Riklund, K., Lindenberger, U., & Bäckman, L. (2012). Memory
aging and brain maintenance. *Trends in Cognitive Sciences*, *16*(5), 292–305.
https://doi.org/10.1016/j.tics.2012.04.005

Olschewski, S., Rieskamp, J., & Scheibehenne, B. (2018). Taxing cognitive capacities reduces
choice consistency rather than preference: A model-based test. *Journal of Experimental
Psychology: General*, *147*(4), 462–484. https://doi.org/10.1037/xge0000403

Olson, E. A., Collins, P. F., Hooper, C. J., Muetzel, R., Lim, K. O., & Luciana, M. (2009). White

Matter Integrity Predicts Delay Discounting Behavior in 9- to 23-Year-Olds: A Diffusion Tensor Imaging Study. *Journal of Cognitive Neuroscience*, *21*(7), 1406–1421. https://doi.org/10.1162/jocn.2009.21107

Owens, M. M., Gray, J. C., Amlung, M. T., Oshri, A., Sweet, L. H., & MacKillop, J. (2017). Neuroanatomical foundations of delayed reward discounting decision making. *NeuroImage*, *161*, 261–270. https://doi.org/10.1016/j.neuroimage.2017.08.045

Park, D. C., Smith, A. D., Lautenschlager, G., Earles, J. L., Frieske, D., Zwahr, M., & Gaines, C. L. (1996). Mediators of long-term memory performance across the life span. *Psychology and Aging*, *11*(4), 621–637. https://doi.org/10.1037/0882-7974.11.4.621

Pehlivanova, M., Wolf, D. H., Sotiras, A., Kaczkurkin, A., Moore, T. M., Ciric, R., … Satterthwaite, T. D. (2018). Diminished Cortical Thickness is Associated with Impulsive Choice in Adolescence. *The Journal of Neuroscience*, 2200–2217. https://doi.org/10.1523/JNEUROSCI.2200-17.2018

Perret, E. (1974). The left frontal lobe of man and the suppression of habitual responses in verbal categorical behaviour. *Neuropsychologia*, *12*(3), 323–330. https://doi.org/10.1016/0028-3932(74)90047-5

Peters, J., & Büchel, C. (2010). Episodic Future Thinking Reduces Reward Delay Discounting through an Enhancement of Prefrontal-Mediotemporal Interactions. *Neuron*, *66*(1), 138–148. https://doi.org/10.1016/j.neuron.2010.03.026

Peters, J., & Büchel, C. (2011). The neural mechanisms of inter-temporal decision-making: Understanding variability. *Trends in Cognitive Sciences*. https://doi.org/10.1016/j.tics.2011.03.002

Pihlajamäki, M., Tanila, H., Hänninen, T., Könönen, M., Laakso, M., Partanen, K., … Aronen,

H. J. (2000). Verbal fluency activates the left medial temporal lobe: a functional magnetic resonance imaging study. *Annals of Neurology*, *47*(4), 470–476. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/10762158

Possin, K. L., Laluz, V. R., Alcantar, O. Z., Miller, B. L., & Kramer, J. H. (2011). Distinct neuroanatomical substrates and cognitive mechanisms of figure copy performance in Alzheimer's disease and behavioral variant frontotemporal dementia. *Neuropsychologia*, *49*(1), 43–48. https://doi.org/10.1016/j.neuropsychologia.2010.10.026

Reitan, R. M. (1992). *Trail Making Test. Manual for administration and scoring.* Tucson, AZ: Reitan Neuropsychology Laboratory.

Reynolds, B. (2006). A review of delay-discounting research with humans: relations to drug use and gambling. *Behavioural Pharmacology*, *17*(8), 651–667. https://doi.org/10.1097/FBP.0b013e3280115f99

Rubin, E. H., Storandt, M., Miller, J. P., Kinscherf, D. A., Grant, E. A., Morris, J. C., & Berg, L. (1998). A Prospective Study of Cognitive Function and Onset of Dementia in Cognitively Healthy Elders. *Archives of Neurology*, *55*(3), 395. https://doi.org/10.1001/archneur.55.3.395

Schmolck, H., Kensinger, E. A., Corkin, S., & Squire, L. R. (2002). Semantic knowledge in patient H.M. and other patients with bilateral medial and lateral temporal lobe lesions. *Hippocampus*, *12*(4), 520–533. https://doi.org/10.1002/hipo.10039

Schroots, J. J. F., Van Dijkum, C., & Assink, M. H. J. (2004). Autobiographical Memory from a Life Span Perspective. *The International Journal of Aging and Human Development*, *58*(1), 69–85. https://doi.org/10.2190/7A1A-8HCE-0FD9-7CTX

Seinstra, M., Grzymek, K., & Kalenscher, T. (2015). Gender-Specific Differences in the

Relationship between Autobiographical Memory and Intertemporal Choice in Older Adults. *PLOS ONE*, *10*(9), e0137061. https://doi.org/10.1371/journal.pone.0137061

Senecal, N., Wang, T., Thompson, E., & Kable, J. W. (2012). Normative arguments from experts and peers reduce delay discounting. *Judgment and Decision Making*, *7*(5), 568–589. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/23596504

Shamosh, N. A., Deyoung, C. G., Green, A. E., Reis, D. L., Johnson, M. R., Conway, A. R. A., … Gray, J. R. (2008). Individual differences in delay discounting: relation to intelligence, working memory, and anterior prefrontal cortex. *Psychological Science*, *19*(9), 904–911. https://doi.org/10.1111/j.1467-9280.2008.02175.x

Sobel, M. E. (1982). *Asymptotic Confidence Intervals for Indirect Effects in Structural Equation Models*. *Source: Sociological Methodology* (Vol. 13). Retrieved from http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.452.5935&rep=rep1&type=pdf

Strotz, R. H. (1956). Myopia and Inconsistency in Dynamic Utility Maximization. *Review of Economic Studies*, *23*(3), 165–180. Retrieved from http://www.jstor.org/stable/2295722?seq=1#page_scan_tab_contents

Stuss, D. T., Bisschop, S. M., Alexander, M. P., Levine, B., Katz, D., & Izukawa, D. (2001). The Trail Making Test: a study in focal lesion patients. *Psychological Assessment*, *13*(2), 230–239. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/11433797

Troyer, A. K., Moscovitch, M., Winocur, G., Alexander, M. P., & Stuss, D. (1998). Clustering and switching on verbal fluency: the effects of focal frontal- and temporal-lobe lesions. *Neuropsychologia*, *36*(6), 499–504. https://doi.org/10.1016/S0028-3932(97)00152-8

Tupak, S. V., Badewien, M., Dresler, T., Hahn, T., Ernst, L. H., Herrmann, M. J., … Ehlis, A.-C. (2012). Differential prefrontal and frontotemporal oxygenation patterns during phonemic

and semantic verbal fluency. *Neuropsychologia*, *50*(7), 1565–1569.

https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2012.03.009

Unsworth, N., Brewer, G. A., & Spillers, G. J. (2009). There's more to the working memory

capacity—fluid intelligence relationship than just secondary memory. *Psychonomic Bulletin*

*& Review*, *16*(5), 931–937. https://doi.org/10.3758/PBR.16.5.931

Vos, S. J. B., Verhey, F., Frölich, L., Kornhuber, J., Wiltfang, J., Maier, W., … Visser, P. J.

(2015). Prevalence and prognosis of Alzheimer's disease at the mild cognitive impairment

stage. *Brain*, *138*(5), 1327–1338. https://doi.org/10.1093/brain/awv029

Vuchinich, R. E., & Simpson, C. A. (1998). Hyperbolic temporal discounting in social drinkers

and problem drinkers. *Experimental and Clinical Psychopharmacology*, *6*(3), 292–305.

Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/9725113

Wang, H., Suh, J. W., Das, S. R., Pluta, J. B., Craige, C., & Yushkevich, P. A. (2013). Multi-

Atlas Segmentation with Joint Label Fusion. *IEEE Transactions on Pattern Analysis and*

*Machine Intelligence*, *35*(3), 611–623. https://doi.org/10.1109/TPAMI.2012.143

Wechsler, D. (1987). *Wechsler Memory Scale-Revised Manual.* San Antonio, TX: The

Psychological Corporation.

Wechsler, D. (1997). *WAIS--III: Wechsler Adult Intelligence Scale (3rd ed.) Administration and*

*scoring manual. The Psychological Corporation, San Antonio, TX.*

Wesley, M. J., & Bickel, W. K. (2014). Remember the future II: meta-analyses and functional

overlap of working memory and delay discounting. *Biological Psychiatry*, *75*(6), 435–448.

https://doi.org/10.1016/j.biopsych.2013.08.008

Wolk, D. A., Das, S. R., Mueller, S. G., Weiner, M. W., Yushkevich, P. A., Apostolova, L. G.,

… Bookheimer, S. Y. (2016). Medial temporal lobe subregional morphometry using high

resolution MRI in Alzheimer's disease. *Neurobiology of Aging*, *0*(0), 2867–2873. https://doi.org/10.1016/j.neurobiolaging.2016.09.011

Xie, L., Pluta, J. B., Das, S. R., Wisse, L. E. M., Wang, H., Mancuso, L., … Yushkevich, P. A. (2017). Multi-template analysis of human perirhinal cortex in brain MRI: Explicitly accounting for anatomical variability. *NeuroImage*, *144*, 183–202. https://doi.org/10.1016/J.NEUROIMAGE.2016.09.070

Xie, L., Wisse, L. E. M., Pluta, J., de Flores, R., Piskin, V., Manjón, J. V., … Yushkevich, P. A. (2019). Automated segmentation of medial temporal lobe subregions on in vivo T1-weighted MRI in early stages of Alzheimer's disease. *Human Brain Mapping*. https://doi.org/10.1002/hbm.24607

Yi, R., & Landes, R. D. (2012). Temporal and probability discounting by cigarette smokers following acute smoking abstinence. *Nicotine & Tobacco Research : Official Journal of the Society for Research on Nicotine and Tobacco*, *14*(5), 547–558. https://doi.org/10.1093/ntr/ntr252

Yu, L. Q., Lee, S., Katchmar, N., Satterthwaite, T. D., Kable, J. W., & Wolf, D. H. (2017). Steeper discounting of delayed rewards in schizophrenia but not first-degree relatives. *Psychiatry Research*, *252*, 303–309. https://doi.org/10.1016/j.psychres.2017.02.062

Yu, R. (2012). Regional white matter volumes correlate with delay discounting. *PloS One*, *7*(2), e32595. https://doi.org/10.1371/journal.pone.0032595