Title: Biosurfactant production maintains viability in anoxic conditions by depolarizing the membrane in *Bacillus subtilis*

Authors

Heidi A. Arjes[1], Lam Vo[1], Caroline Marie Dunn[2], Lisa Willis[1], Christopher A. DeRosa[3], Cassandra L. Fraser[3], Daniel B. Kearns[2,*], Kerwyn Casey Huang[1,4,5,*]

Affiliations

[1]Department of Bioengineering, Stanford University School of Medicine, Stanford, CA 94305, USA

[2]Department of Biology, Indiana University, Bloomington, IN 47405, USA

[3]Department of Chemistry, University of Virginia, Charlottesville, VA 22904, USA

[4]Department of Microbiology & Immunology, Stanford University School of Medicine, Stanford, CA 94305, USA

[5]Chan Zuckerberg Biohub, San Francisco, CA 94158

*Corresponding authors: kchuang@stanford.edu and dbkearns@indiana.edu

Lead author: Kerwyn Casey Huang (kchuang@stanford.edu)

Keywords: surfactin, biosurfactant, oxygen depletion, cell lysis, protoplast, respiration, membrane potential, membrane energetics, LytC

Summary

The presence or absence of oxygen in the environment is a strong effector of cellular metabolism and physiology. Like many eukaryotes and some bacteria, *Bacillus subtilis* is an obligate aerobe that primarily utilizes oxygen during respiration to generate ATP. Despite the importance of oxygen for *B. subtilis* survival, we know little about how oxygen is consumed during growth and how populations respond to shifts in oxygen availability. Here, we find that when oxygen was depleted from stationary phase cultures ~90% of *B. subtilis* 3610 cells died and lysed due to autolysin activity; the remaining cells maintained colony-forming ability. Interestingly, the domesticated 168 strain maintained a higher optical density than 3610 during oxygen depletion due to the formation of cell-wall-less protoplasts, but the remaining, rod-shaped cells were >100-fold less viable than 3610. We discovered that the higher viability in 3610 was due to its ability to produce the antibacterial compound surfactin, as surfactin addition rescued 168 viability and also increased yield in aerobic growth. We further demonstrate that surfactin strongly depolarizes the *B. subtilis* membrane, and that other known membrane-potential disruptors restore viability to 168. These findings highlight the importance of surfactin for survival during oxygen-depleted conditions and demonstrate that antimicrobials normally considered harmful can instead benefit cells in stressful conditions when the terminal electron acceptor in respiration is limiting.

Introduction

Many species across all domains of life use oxygen as the terminal electron acceptor during aerobic respiration, which produces approximately ten-fold more ATP per glucose molecule than via glycolysis and fermentation in the absence of respiration. In most human cells, oxygen depletion causes exhaustion of ATP and eventual death, either through lysis caused by osmotic stress due to the inability to regulate osmolyte levels [1, 2] or through activation of signaling cascades that lead to apoptosis [3]. Thus, maintaining concentration gradients of ions across cell membranes is of paramount importance when oxygen is lacking. Like human cells, certain microbes such as the pathogen *Mycobacterium tuberculosis* as well as most fungi (with the exception of yeasts) [4] are considered strict aerobes due to their inability to make ATP in the absence of oxygen. Under conditions of rapidly depleted oxygen, *M. tuberculosis* cells show a loss in viability [5, 6]. The related, soil-dwelling species *Mycobacterium smegmatis* also loses viability upon oxygen depletion, and the remaining viable cells maintain the redox balance using hydrogen fermentation and activate stress response genes critical for survival [7, 8]. Despite the clinical importance of *M. tuberculosis*, surprisingly few studies have interrogated how these and other strict aerobes respond to oxygen depletion and the genes responsible for survival.

The Gram-positive model bacterium *Bacillus subtilis* is considered a strict aerobe when grown in the absence of nitrate or nitrite [9]. *B. subtilis* is naturally found in the soil and is used as an additive to help prevent infections and promote growth in plants [10, 11]. In the soil, *B. subtilis* undergoes constant shifts in oxygen concentration, as oxygen is readily available in dry soils but diffuses less and becomes depleted in wet or flooded soils

following a rain [12]. Early observations of *B. subtilis* culture lysis upon a shift to anoxic

environments have yet to be further characterized [13], and it remains a mystery whether

B. subtilis has strategies to cope with oxygen limitation.

B. subtilis has long been domesticated in the lab, leading to a multitude of genetic

tools, strain libraries, and online databases and resources [14-16]. The high level of genetic

relatedness between biofilm-forming "wild" strains and derivative non-biofilm-forming

laboratory strains has been exploited to identify genetic differences that underlie biofilm

community behaviors [17]. For instance, the commonly studied laboratory strain 168,

which is derived from the biofilm-forming strain 3610, lacks an extrachromosomal plasmid

and harbors several point mutations that reduce or abolish social behaviors such as matrix

production [18]. Notably, 3610 produces the small molecule surfactin, a powerful

surfactant that has previously been implicated in swarming motility [19-21]. Surfactin has

also been shown to kill fungi [22] and some bacteria *in vitro* [22-26]. However, fitness

benefits of surfactin production in the context of planktonic cultures have yet to be

identified, although some surfactants can accelerate oxygen diffusion through the air-water

interface [27].

Here, we characterize the interplay between oxygen availability and surfactin

production during the growth and death of *B. subtilis* cultures. We show that oxygen

becomes limiting in the culture during the transition to stationary phase and that surfactin

secretion improves growth yield in stationary phase by increasing oxygen availability.

During a shift to anoxic conditions, we demonstrate that the majority of *B. subtilis* cells die

and lyse due to the activity of the LytC autolysin and surfactin. Finally, we discover that

surfactin maintains the viability of the remaining cells by causing membrane

depolarization that allows these cells to survive until oxygen is restored.

Results

Oxygen depletion leads to rapid death and lysis of most *B. subtilis* cells

When culturing the biofilm-forming *B. subtilis* strain 3610 in LB, we noticed that once a test tube containing a late exponential culture was shifted from a shaking incubator to a stationary rack on the bench, the opacity of the tube decreased continuously over a period of 10 hours, suggesting that cells were dying and lysing (Fig. 1A). Consistent with lysis, microscopic observation of cultures left to sit on the bench for 10 hours revealed substantial phase-gray cell remnants in addition to phase-dark rod-shaped cells (Fig. 1A). We conclude that *B. subtilis* 3610 cultures exhibit death and lysis during static incubation after cessation of rapid growth.

The static culture would be limited in oxygen diffusion, and as *B. subtilis* is a strict aerobe that relies on its use of oxygen as a terminal electron acceptor during respiration, we hypothesized that the death in the culture was due to oxygen limitation. To measure oxygen levels during oxygen growth and depletion, we added phosphorescent oxygen-sensitive nanoparticles (Methods) to cell cultures in microtiter plates and measured optical density (OD) and light emission over time. To establish a controlled environment, we grew cultures in sealed 96-well microtiter plates with a hole poked in the seal to allow oxygen exchange, or limited oxygen by completely sealing the wells (Fig. 1B, top); we utilized linear shaking to prevent an oxygen gradient in the culture. In media without cells, oxygen initially diffused into the media due to shaking, and then remained at an approximately constant value (Fig. 1B, bottom). In media with cells, oxygen levels initially increased (again due to shaking) but then began to decrease when the culture reached an $OD_{600} \sim 0.05$,

indicating that the cells were consuming the oxygen faster than it dissolved into the media

(Fig. 1B). When cultures reached an $OD_{600}\sim0.5$, oxygen levels were undetectable but the

culture continued to grow (Fig. 1B), presumably because the cells rapidly consumed any

oxygen that dissolved into the media.

When the cultures reached an $OD_{600}\sim0.8$, we sealed the wells to abolish oxygen

exchange in the headspace, and returned the cultures to a shaking environment. As

expected, the measured oxygen levels in the cultures remained low (Fig. 1B, lower right).

Within 2 hours, the OD_{600} began to decrease, reminiscent of our observations of OD loss in

standing test tubes (Fig. 1A), and plateaued at ~0.1 after 48 h (Fig. 1B, upper right).

Consistent with the OD drop upon sealing the well, cell viability (defined here as the ability

of a cell to form a colony on an agar plate) dropped approximately ten-fold after 48 h (Fig.

1C). These observations suggest that the loss of OD in standing tubes and in agitated sealed

plates was due to cell lysis during oxygen depletion.

To further explore the correlation between cell lysis and oxygen depletion, we

investigated the effect of cell density on hypoxia-induced cell lysis by growing cultures

aerobically to different optical densities prior to sealing the wells. Regardless of starting

culture density (OD_{600} from 0.4 to 0.8), the OD_{600} either stayed approximately constant or

increased slightly for ~1 hour after sealing, and then the cultures exhibited large decreases

in OD over time (Fig. 1D, S1A). The maximum lysis rate (the absolute value of the most

negative slope of the $\ln(OD_{600})$ curve) increased with increasing starting OD, such that the

cultures that began at the highest density ($OD_{600}\sim0.8$) had a lysis rate of 40% decrease per

hour (Fig. 1D). Since the lysis behavior varied with initial OD, we standardized all further

experiments by growing 3610 cells to an OD_{600} of ~0.9-1.1 before cutting off oxygen

exchange. Transfer of OD~1 cultures into an anaerobic chamber resulted in an even higher

maximum lysis rate of 86% decrease per hour (Fig. S1B). From these data, we infer that

populations at high density deplete the remaining oxygen more rapidly, and that rapid

oxygen depletion more readily triggers cell lysis.

A cell wall autolytic enzyme is necessary for lysis of non-viable cells upon oxygen

depletion

In addition to losing viability upon oxygen depletion, the non-viable population also lysed

and was removed from the population of intact cells, as indicated by the substantial

decrease in optical density. Given the importance of the cell wall for maintaining cellular

integrity in bacteria, we hypothesized that this lysis involves the activity of cell wall

autolytic enzymes. Cell wall growth requires both insertion of new material and cleavage of

the existing peptidoglycan by autolysins [28]. In *B. subtilis*, the major autolysin LytC is

under control of the vegetative sigma factor σ^A and the alternative sigma factor σ^D [29, 30],

preventing the unchecked breakdown of the cell wall when insertion is disrupted [31]. We

measured the OD of a $\Delta lytC$ 3610 mutant and found that the mutant exhibited reduced lysis

upon oxygen depletion (Fig. 1C). As a specificity control, deletion of another σ^D-dependent

autolysin LytD phenocopied wild-type behavior (Fig. S1C), suggesting that LytC is the

primary autolysin activated in oxygen-depleted conditions. Despite the higher biomass in

the $\Delta lytC$ cultures after oxygen depletion, a similar proportion of cells (~10%) retained

viability as in wild-type cultures (Fig. 1C). Single-cell imaging of the oxygen-depleted

cultures revealed mixed populations of rod-shaped cells, some of which were phase-gray

"ghosts" and others that were phase dark (Fig. 1C). Time-lapse microscopy demonstrated

that only a portion of the phase-dark rod-shaped Δ*lytC* cells were capable of resuming

growth (Fig. S1D, movie S1, S2), consistent with our plating efficiency data (Fig. 1C). These

data indicate that lysis and cell viability are genetically separable phenotypes, since many

Δ*lytC* mutant cells remain intact yet still lose viability. Thus, since LytC degrades the cell

wall but does not impact the viability of 3610 cultures upon oxygen depletion, we conclude

that lysis is downstream of viability loss during oxygen depletion.

Laboratory-domesticated *B. subtilis* strains exhibit more viability loss despite

reduced lysis upon oxygen depletion

B. subtilis strain 168, a genetic derivative of 3610, is commonly used in research

laboratories as it was selected for dispersed growth in culture and improved genetic

tractability [17]. In microtiter plate assays, we found that upon oxygen depletion the OD of

168 cultures decreased at a lower rate and remained at a higher level throughout 48 h of

monitoring (Fig. 2A). Imaging of oxygen-depleted 168 cultures revealed the presence of

round cells in addition to rod-shaped cells (Fig. 2B). Some round cells were intact and did

not stain with propidium iodide (PI), while others had compromised membranes and thus

stained brightly with PI (Fig. 2B). In cell cultures that were first stained with fluorescent D-

amino acids (FDAAs) to label cell walls, the vast majority of the rounded cells did not retain

FDAA staining, indicating that they were protoplasts without cell walls (Fig. S2A). Indeed, a

168 Δ*lytC* strain did not form round cells and exhibited less lysis (Fig. S2B), indicating that

the protoplasts released in 168 are more fragile than their walled counterparts. Thus, we

conclude that LytC activity during oxygen depletion degrades the cell wall in 168, releasing

membrane-bound protoplasts that account for the higher OD in 168 cultures compared

with 3610 cultures.

Despite having a higher OD_{600} than 3610 cultures, the colony-forming units (CFU) in

168 cultures depleted of oxygen for 24 h decreased ~100-fold relative to 3610 (Fig. 2C).

Indeed, time-lapse imaging on fresh medium with oxygen showed that despite having

intact cell envelopes, only ~1% of the rod-shaped 168 cells were able to grow and divide,

whereas ~95% of the 3610 cells exhibited growth (Fig. S2C, Movie S3,S4). We never

observed growth either in increased cell number or increased cell size during 12 h time-

lapse imaging of any of the protoplasts on fresh LB or on filtered spent medium, suggesting

that the protoplasts formed in strain 168 are not viable. We conclude that despite reduced

cell lysis, 168 cultures experience a much more drastic loss of viability upon oxygen

depletion.

Surfactin increases lysis and restores viability to the domesticated strain 168

Strains 3610 and 168 differ in a number of chromosomal loci, and 168 is defective in

phenotypes related to secretion of extracellular products that support multicellular

behaviors such as biofilm formation and swarming motility [18]. To determine whether

extracellular products were responsible for the oxygen-related phenotypic differences

between the two strains, we subjected a 1:1 volumetric mixture of the strains to oxygen

depletion. The viability of 168 (which has a colony morphology visually distinct from that

of 3610) increased dramatically in the co-culture (Fig. 4C). A candidate compound that

could restore viability to 168 is surfactin, which is a strong surfactant that creates K^+-

permeable pores and solubilizes lipid bilayer vesicles *in vitro* [32]. Surfactin is produced by

3610 but not 168, due to a mutation in *sfp*, which encodes an enzyme necessary to activate

the surfactin biosynthesis complex [19]. We found that exogenous surfactin addition at

either the time of inoculation of the culture or at the initiation of oxygen depletion

increased the lysis rate of 168 and eliminated protoplasts (Fig. 2D,E), consistent with

surfactin-induced protoplast lysis.

Remarkably, exogenous surfactin addition restored the viability of oxygen-depleted

168 cultures to similar levels as 3610 (Fig. 2F). Moreover, a 168 strain with *sfp* genetically

complemented to restore surfactin production phenocopied the surfactin-treated 168

cultures in terms of lysis, cell morphology, and viability (Fig. 2F, S2). Thus, we conclude

that surfactin production both promotes protoplast lysis and maintains viability in oxygen-

depleted *B. subtilis* cultures.

Surfactin improves growth yield of the domesticated strain 168 by increasing oxygen

diffusion

One intriguing distinction between the aerobic growth of 3610 and 168 was the divergence

of the growth curves around OD_{600}~0.3; after this point, 3610 cultures completed another

~1.5 mass doublings to reach OD_{600}~0.9, while 168 cultures underwent only 1 mass

doubling to reach OD_{600}~0.6 over the same period (Fig. 3A). We noted that a surfactin-

supplemented culture experienced more growth after the OD reached ~0.3 than untreated

cultures, resulting in a higher overall yield similar to that of 3610 (Fig. 3B). When surfactin

was added just prior to the divergence in growth curves, the yield was similarly restored

(Fig. 3B). Thus, we inferred that surfactin production was responsible for the increase in

yield of 3610 relative to 168. The detergent Tween 80 similarly increased growth yield in

168 cultures (Fig. 3C), indicating that the detergent properties of surfactin were likely

responsible for the increased yield.

Certain compounds have the ability to increase oxygen diffusion in liquid [33], and it

has been proposed that oxygen diffusivity is rate-limiting for the function of some

biological systems [34]. To test whether surfactin increased oxygen in the media, we used

our oxygen nanoprobe to compare oxygen levels in 168 cultures with and without added

surfactin. We found that exogenous surfactin increased the oxygen levels in cultures during

late-exponential phase, when oxygen would normally be depleted lower than our limit of

detection (Fig. 3D). A similar increase occurred due to addition of Tween 80 (Fig. 3D). A

biophysical model incorporating diffusion and cellular consumption of oxygen predicted

that the observed ~1.2-fold increase in peak oxygen level is consistent with a ~1.4-fold

increase in diffusion rate (Methods). Thus, we infer that the presence of surfactin results in

higher growth yield due to the enhanced availability of oxygen during the transition to

stationary phase when oxygen would otherwise be limiting for growth.

A transposon screen supports surfactin production as the main determinant of

survival upon oxygen depletion

Given the large decreases in viability after 24 h of oxygen depletion, we performed a

genetic screen to attempt to identify any mutations that would increase the survival of 168

and 3610 without oxygen. We made 20-30 independent transposon libraries of 5000-

10,000 individual transposon mutants per library in each background and subjected these

libraries to oxygen depletion, hypothesizing that any mutants with enhanced survival

would be enriched (Fig. S3A). We could not identify any such 168 mutants, suggesting that

there is not an easily obtainable loss-of-function mutant in a surfactin-independent

pathway to increase viability upon oxygen depletion.

250   We identified two categories of mutants in the 3610 background with decreased

lysis upon oxygen depletion: surfactin production (*comA*, *comP*, *srfAA,* and a hit upstream of

rghR (*rghR*$_{us}$)) and flagella-related (*fliI, fliJ*, and *fliF*) (Fig. S3B, Table S1). *srfAA* encodes the

surfactin synthetase subunit A, which is part of the enzyme that synthesizes surfactin (Fig.

4A) [19, 35]. ComA and ComP form a two-component system that activates surfactin

production at high cell density [36]. RghR regulates RapG and RapH, two repressors of the

srfA operon (Fig 4A) [37]. The flagella-related mutants disrupted *fliI* and *fliJ*, which encode

accessories of the flagellar type III secretion apparatus, and *fliF*, which encodes the flagellar

basal ring [38-41]. In-frame markerless deletions of these mutants had ~4-fold lower

viability than the parent after 24 h of oxygen depletion (Fig. S3C), which we hypothesized

was due to the surfactin in the pooled library cultures increasing their survival advantage

in the mixed population but not in isolation. Indeed, upon exogenous surfactin addition, all

mutants responded with increased lysis and removal of protoplasts/cell debris (Fig. 4B,C,

S3D). Taken together, the fact that we obtained no hits that increased viability in a 168

background and that all hits in a 3610 background are directly related to surfactin

production or respond to surfactin points to surfactin as the primary determinant of

viability in the absence of oxygen.

Surfactin restores viability by depolarizing the membrane

In vitro, surfactin creates potassium ion-permeable pores in lipid bilayers [32]. If this

behavior occurs *in vivo*, such pores would reduce the strength of the potassium ion

gradient across the cell membrane and alter membrane potential. Thus, we tested the

energetic state of cells after various chemical treatments using the membrane potential-

sensitive dyes $DiSC_3(5)$ and ThT [42, 43]. $DiSC_3(5)$ is taken up by cells and the fluorescence

signal is initially quenched. Agents that depolarize cells release $DiSC_3(5)$ into the medium,

resulting in an increased signal [42]. By contrast, ThT enters cells with polarized

membranes and fluoresces inside the cell; upon depolarization, ThT exits the cells and the

fluorescence is reduced [43]. As expected, we found that treatment of 168 cells with

valinomycin, a known depolarizing agent that functions as a potassium-specific transporter

[44], led to an increase in $DiSC_3(5)$ and a decrease in ThT signal (Fig. 5A). CCCP, a proton

ionophore that dissipates the proton motive force [45], strongly reduced the ThT signal

and slightly increased $DiSC_3(5)$ fluorescence (Fig. S4). Surfactin treatment led to a large

increase in $DiSC_3(5)$ fluorescence and a greater decrease in ThT fluorescence than

valinomycin (Fig. 5A), demonstrating that surfactin can strongly depolarize *B. subtilis* cells.

Based on these findings, we hypothesized that de-energizing the membrane would

be sufficient to rescue the colony-forming ability of 168 cultures upon oxygen depletion.

Indeed, valinomycin and CCCP both restored viability to a similar extent as surfactin (Fig.

5B). Interestingly, valinomycin- and CCCP-treated cultures had protoplasts after 24 h of

oxygen depletion, consistent with surfactin being necessary for protoplast lysis and

indicating that protoplast formation is independent from viability maintenance. Taken

together, these data indicate that membrane potential dictates the ability of *B. subtilis* cells

to survive oxygen depletion.

Discussion

While surfactin has been recognized to promote swarming motility and complex colony architecture in *B. subtilis* communities such as biofilms [18, 46], its role in planktonic cultures has remained mysterious. Here, we demonstrate three independent roles of surfactin in planktonic cultures. During aerobic growth, the detergent properties of surfactin increase growth yield by increasing the oxygen available to cells entering stationary phase (Fig. 3). During oxygen depletion, surfactin plays two independent roles: (1) it works in tandem with LytC to remove non-viable cells from the culture (Fig. 2), and (2) it depolarizes the remaining cells and thereby maintains their viability during oxygen depletion (Fig. 5).

The increase in growth yield due to surfactin suggests that oxygen is limiting even in aerobic cultures and that increasing oxygen availability can increase yield. This finding has industrial implications as *B. subtilis* is commonly used to produce many biological compounds such as enzymes and antibiotics [47], and maintaining oxygen levels are critical to optimize production of the desired compound [48, 49]. If this oxygen-related growth yield enhancement holds across species, detergent addition may increase bioproduction in other commonly utilized organisms such as *Streptomycetes* species, which are used to isolate numerous antibiotics, and the industrial powerhouse *Escherichia coli*, which is widely used to synthesize many biopharmaceuticals such as insulin [47]. In addition to oxygen, surfactin may also facilitate growth by increasing nutrient diffusion to cells during late exponential phase when nutrients become limiting.

We also found that surfactin has at least two functions during oxygen starvation that are distinct from its role during aerobic growth. Surfactin causes lysis in cells that experience LytC-mediated cell wall degradation (Fig. S2), presumably through membrane disruption. Moreover, surfactin maintains viability of the remaining intact cells through membrane depolarization, which may allow these cells to enter a metabolically inactive state where they can ride out the stress of the oxygen depletion. While it was formally possible that cell lysis helped maintain viability of the remaining cells, the observations that $\Delta lytC$ mutants have the same viability as wild-type cultures even though the non-viable cells remain intact (Fig. 1C) and that CCCP or valinomycin treatment of 168 cultures rescues viability without removing protoplasts (Fig. 5C) demonstrate that surfactin maintains viability by acting directly upon the membrane potential of the surviving cells.

Generally, OD is used as a proxy for cell number, which holds true for our measurements of 3610 cultures undergoing oxygen depletion as the 10-fold drop in OD measured reflects cell lysis and quantitatively mirrors the decrease in viability (Fig. 1C). However, our work highlights instances during oxygen depletion when biomass measured by OD is uncoupled from viability: both $\Delta lytC$ 3610 cells and 168 cells remain intact and the cultures have a relatively high OD compared with wild-type 3610, but most cells cannot form colonies (Fig. 2C, S1C). This uncoupling between OD and viability has been observed previously in cell-cycle mutants where cells remain intact and metabolically active but cannot divide and form colonies [51], and may be more prevalent than is currently appreciated, motivating future studies that rely on OD to perform assays to verify culture viability.

Since cell depolarization can actually improve the viability of cells undergoing

oxygen depletion, under conditions with limiting terminal electron acceptors, certain

antibacterial treatments that inhibit growth may actually keep cells viable. Such a

possibility needs to be taken into consideration when treating bacterial infections and

removing bacteria in low-oxygen clinical and industrial settings, particularly those in which

the microbe primarily gains energy through aerobic respiration and hence the terminal

electron acceptor may be limiting. Moreover, it remains generally unclear which

antimicrobial treatments will disrupt membrane potential as a side effect of their primary

activity or mechanistically how antibiotic treatments affect cells in low oxygen

environments, motivating further studies of membrane energetics during growth-

inhibition.

The cell-wall breakdown of *B. subtilis* cells during oxygen depletion provides further

support for a recently discovered regulatory role of membrane potential in cell-wall

synthesis [52]. Indeed, LytC is activated upon sodium azide treatment that deprotonates

the cell wall [53, 54] and the activity of peptidoglycan synthesis enzymes in *E. coli* are

regulated by pH [55, 56]. In addition, the recent observation that actively growing and

dormant cells in a *B. subtilis* culture respond oppositely to an electrical pulse wherein they

either hyperpolarize or depolarize, respectively [57], suggests that membrane energetics

may explain the observed population heterogeneity in our oxygen-depleted 3610 cultures.

Thus, as membrane potential is of utmost importance in metabolism and cell growth, is

becoming more appreciated in regulating cell-wall remodeling, and likely feeds back into

many additional aspects of cell physiology, bacteria must employ strategies to maintain

and/or modulate membrane potential during changing environmental conditions. The

ability of surfactin to both alter membrane energetics to maintain viability during oxygen-

depletion and to enhance growth in oxygen-limited conditions has likely provided multiple

fitness advantages to *B. subtilis* in spatially structured and complex environments such as

native soil communities, and strategies for regulating membrane potential may be an

important factor for survival of other strict aerobes as well.

Methods

Media and growth conditions

All strains and their genotypes are listed in Table S2. Strains were grown in LB (Lennox broth with 10 g/L tryptone, 5 g/L NaCl, and 5 g/L yeast extract). Antibiotics for selection of mutant strains were used as follows: kanamycin (5 µg/mL), MLS (a combination of erythromycin at 0.5 µg/mL and lincomycin at 12.5 µg/mL), chloramphenicol (5 µg/mL), and spectinomycin (100 µg/mL). Surfactin was added at a final concentration of 0.05 mg/mL unless otherwise noted. Strains were cultured either in 5 mL of medium in a test tube on a roller drum or in 200 µL of medium in a 96-well plate in a Biotek Epoch2 spectrophotometer under linear shaking (shaking was set to 567 cycles per minute (cpm), 3-mm magnitude of shaking). For all experiments, the initial inoculum was from a fresh colony struck from a -80 °C freezer stock onto LB 1.5% agar plates and incubated overnight at 37 °C.

Strain construction

Strains were constructed using SPP1 phage transduction [58]. The donor strain was grown for >6 h in TY medium (LB supplemented with 0.01 M $MgSO_4$ and 0.1 mM $MnSO_4$ after autoclaving). Ten-fold dilutions of SPP1 phage were added to the culture and 3 mL TY soft (0.5%) agar was mixed with the culture/phage mixture and poured over a TY plate (1.5% agar) overnight. A plate was chosen that exhibited nearly total clearing of cells without a large number of phage-resistant mutants. Five milliliters of TY medium were added to this plate and a 1-mL filter tip was used to scrape up the soft agar. This soft agar/liquid mix was

filtered through a 0.4-μm filter. The phage was added to a stationary-phase (grown for 6-10

h) culture in TY medium of the recipient strain (10 μL undiluted phage + 100 μL recipient

cells, and optionally 900 μL TY medium) and incubated at 37 °C for 30 min, then plated

onto LB + antibiotic and 0.01 M sodium citrate (sodium citrate was omitted for MLS

selection). Plates were incubated for 24 h and transductants were struck for single colonies

to eliminate the phage.

To generate the Δ*lytC* in-frame marker-less deletion construct, the region upstream

of *lytC* was PCR-amplified using the primer pair 1427/1428 and digested with SalI and

EagI, and the region downstream of *lytC* was PCR-amplified using the primer pair

1425/1426 and digested with EagI and BamHI. The two fragments were then

simultaneously ligated into the SalI and BamHI sites of pMiniMAD, which carries a

temperature-sensitive origin of replication and an erythromycin resistance cassette [59], to

generate pDP299.

To generate the Δ*lytD* in-frame marker-less deletion construct, the region upstream

of *lytD* was PCR-amplified using the primer pair 1429/1430 and digested with SalI and

EagI, and the region downstream of *lytD* was PCR-amplified using the primer pair

1431/1432 and digested with EagI and BamHI. The two fragments were then

simultaneously ligated into the SalI and BamHI sites of pMiniMAD to generate pDP300.

Deletion plasmids were introduced into strain DK1042 (Table S2) via single cross-

over integration by transformation at the restrictive temperature for plasmid replication

(37 °C) using MLS resistance as a selection. To evict the plasmid, the strain was incubated

in 3 mL LB broth at a permissive temperature for plasmid replication (22 °C) for 14 h, and

serially diluted and plated on LB agar at 37 °C. Individual colonies were patched on LB

plates and LB plates containing MLS to identify MLS-sensitive colonies that had evicted the

plasmid. Chromosomal DNA from colonies that had evicted the plasmid was purified and

screened by PCR using primers 1427/1426 or 1430/1431 to determine isolates that had

retained the Δ*lytC* or Δ*lytD* allele, respectively [59, 60].

In-frame deletions of MLS-knockout mutants were constructed as outlined

previously [14]. Briefly, the strain of interest was transformed with pDR244 at 30 °C.

Several transformants were struck onto LB at 42 °C. Resultant colonies were patched on

MLS and spectinomycin plates to confirm that the plasmid and MLS cassette were lost.

Colonies were screened for altered oxygen phenotype.

Growth and lysis assays

Strains were struck out for single colonies on the evening prior to the experiment. Colonies

were inoculated into fresh LB and grown aerobically at 37 °C, either in 5 mL LB on a roller

drum or in 200 µL in a Greiner 96-well plate. To deplete oxygen from cultures grown in test

tubes, 200 µL were aliquoted into a 96-well plate and the plate was sealed with optical film

(Excel Scientific AeraSeal). To grow cultures in the plate reader, 200 µL of stationary-phase

inocula were aliquoted into each well of a 96-well plate, with the edge wells containing

medium only. Optical film was used to cover the plates, and one hole per well was poked

using a 20-gauge needle to allow for air exchange. 168 and 3610 cultures were then grown

until the 3610 culture reached an $OD_{600} \sim 1.0$. To deplete the oxygen, the plate was taken

out of the plate reader, the optical seal with holes was removed, and a new seal was placed

onto the plate to fully cut off oxygen exchange. We grew and depleted cultures at 37 °C

using linear shaking (567 cpm, 3-mm magnitude) and read OD_{600} every 7.5 min in a Biotek

Epoch2 spectrophotometer.

Mariner transposon library construction

The mariner transposon was used to create a library of insertion mutants. The parent

strains (HA1235 and HA1414) were struck for single colonies onto MLS plates at 30 °C. One

colony per library was grown in 3 mL LB+kanamycin at room temperature overnight.

Transposon-insertion libraries were selected by plating 10-fold dilutions of the cultures on

prewarmed LB+kanamycin plates and incubating overnight at 37 °C.

Screen to enhance for mutants that survive better without oxygen

Each library (~5000-10,000 colonies per library) was inoculated into 5 mL LB and grown

on a roller drum at 37 °C to an OD_{600}~1.0. Libraries were then aliquoted into wells of a 96-

well plate (200 μL per well) and plates were sealed to deplete oxygen. Oxygen was

depleted at 37 °C for 2-8 days for 168 libraries or at 30 °C for 4-8 days for 3610 libraries. At

various time points following the start of oxygen depletion, one well of the library was

harvested and struck out for single colonies. One or two colonies per well were tested for

either enhanced lysis upon oxygen depletion (3610 background) or enhanced colony-

forming ability (168 background). Any mutants found to have these phenotypes were back-

crossed into the parent using SPP1 phage transduction to ensure the transposon mutation

was the causative agent. To map the mutation, genomic DNA was prepped from each

mutant. Inverse PCR was carried out using Phusion polymerase with the primers IPCR1

and IPCR2. The PCR products were gel-purified and sequenced using the IPCR2 primer. The

sequences were mapped onto the *B. subtilis* genome using BLASTN.

Oxygen nanoprobe measurements

Relative oxygen levels were measured using the oxygen-sensitive nanoprobe

(BF$_2$nbm(I)PLA) that emits an oxygen-dependent phosphorescence reading and an oxygen-

independent fluorescence reading, which together can be used to calculate the relative

oxygen level of the medium [61, 62]. Briefly, the nanoprobe was added at 5% (10 μL into

190 μL) to the inoculum before growth of the cultures. In addition to OD$_{600}$ readings,

fluorescence readings (ex/em: 414/450 nm) and phosphorescence readings (ex/em:

415/560 nm, with a 2-ms delay between excitation and emission) were taken using a

Biotek Synergy H1 spectrophotometer. Readings were taken every 7.5 min, with incubation

at 37 °C and linear shaking (567 cpm, 3-mm magnitude of shaking).

PI staining and phase microscopy

Five hundred nanoliters of cultures were spotted onto LB pads made with 1.5% agar and

10 μM PI. Once dry, a coverslip was added and cells were imaged on a Nikon Ti-E inverted

microscope using a 100X objective (NA: 1.4). Phase and fluorescence (mCherry filter,

ex/em: 570/645 nm) images were acquired. Images were processed identically in Adobe

Photoshop and merged using FIJI.

Time-lapse microscopy of oxygen-depleted cultures

The bottom of a rectangular Singer PlusPlate culture plate was used to make a large pad

[63], in which 35 mL of LB+1.5% agar was pipetted onto the plate ~1 h before imaging so

that the agar could solidify completely. Once solid (after ~5-10 min), a second Singer

PlusPlate was placed on top of the agar pad to prevent contamination and drying. One

microliter of cultures was spotted in the center of the pad and allowed to dry. A large 113

by 77 mm custom-made no. 1.5 glass coverslip (Nexterion) was applied [63]. Imaging was

carried out in a heated environmental chamber with a water bubbler and several

reservoirs of water to humidify the chamber. Phase images were acquired on a Nikon Ti-E

inverted microscope every 5 min using a 40X air objective (NA: 0.95) with 1.5X

magnification. Images were compiled into movies and analyzed using Matlab or FIJI. All

rod-shaped cells were identified in the first frame and defined as able to grow if they at

least doubled in mass and divided without lysis during the experiment.

Fluorescent D-amino acid staining

HADA [64] was added to cultures at a final concentration of 500 µM during the last mass

doubling of growth. To reduce background staining, cultures were diluted 1:10 in MSgg

solution (5 mM potassium phosphate buffer + 0.05 M MOPS at pH 7) and then spotted onto

a pad of that solution made with 1.5% agar. Cells were imaged using a Nikon Ti-E inverted

microscope with a 100X oil objective (NA: 1.4). Phase and fluorescence (DAPI filter, ex/em:

375/460 nm, exposure time 2 s) images were acquired. Phase and fluorescence images

were adjusted identically in Adobe Photoshop and merged in FIJI.

Plating efficiency

Cultures were harvested and diluted 10-fold into LB. One hundred microliters of the

dilutions were plated onto an LB plate. These plates were incubated overnight at 37 °C in a

single layer (not stacked). CFU/mL values of the original culture were calculated from the

colony counts of the dilutions that had distinct colonies.

Lysis assay starting at different optical densities

Three colonies were inoculated into a single 10 mL LB culture and mixed well. Six two-fold

serial dilutions of the culture were carried out and 5 mL of each dilution were transferred

to a test tube and incubated at 37 °C on a roller drum until the undiluted culture reached an

OD_{600}~0.8. Two hundred microliters of each culture were then aliquoted into a 96-well

plate, the plate was sealed with optical film, and OD_{600} was monitored in a Biotek Epoch2

spectrophotometer.

Oxygen depletion in an anaerobic chamber

Cultures were grown in aerobic conditions (96-well plate sealed with optical film, with

holes poked through the film for each well). Cultures were then transferred into a Coy

anaerobic chamber and OD_{600} was monitored using a Biotek Epoch2 spectrophotometer as

above.

Biophysical model linking oxygen diffusivity and concentration in a growing culture

Since the addition of surfactin increased oxygen levels in late-exponential phase of a

growing culture (Fig. 3D), we sought to understand whether oxygen diffusivity could

explain the increase. In a small region of extent Δ*l* at a depth *l* below the air-liquid interface

of the culture, oxygen is depleted at a rate $\gamma \times \rho(l) \times c(l)\Delta l$, where γ is the absorption rate

per unit oxygen in close proximity to a cell, $\rho(l)$ is the cell density at depth l, and $c(l)$ is the

oxygen concentration at depth l. At steady state, the rate of oxygen uptake in the region of

extent Δl must be balanced by the rate of supply via diffusion, that is

526
$$D\frac{\partial^2 c}{\partial l^2}\Delta l = \gamma\rho c\Delta l, \qquad (1)$$

where D is the diffusivity of oxygen. The solution to Eq. 1 is $c(l) = c_0 e^{-l\sqrt{\gamma\rho/D}}$, assuming

that c_0 is the oxygen concentration at the surface, $c \to 0$ as $l \to \infty$, and $\gamma\rho/D$ is independent

of l. Hence, as D increases, oxygen reaches increased depths as $\sim \sqrt{D/\gamma\rho}$, and the total

amount of oxygen in the culture increases as $\int c(l)dl = c_0\sqrt{D/\gamma\rho}$. Thus, a 1.2-fold increase

in the total amount of oxygen corresponds to a 1.4-fold increase in D of $\sim$ 1.4-fold,

assuming that c_0 is linked to solubility and γ and ρ remain approximately constant.

Membrane potential measurements

Membrane potential was measured using a protocol modified from [42]. Cells were grown

in 5 mL LB in a roller drum to $OD_{600}\sim1$. Cells were washed in a buffer containing 10 mM

potassium phosphate, 5 mM MgSO4, and 250 mM sucrose (pH 7.0) and then resuspended

to a calculated OD_{600} of 0.085 in that same buffer (pelleting steps were at 5100 rcf for 3

min). Two hundred microliters of this mixture were added to wells in a 96-well plate. 1 μM

DiSC3(5) or 10 μM Thioflavin T (ThT) were added to the wells. Fluorescence readings

(em/ex: 620/685 for DISC and 450/482 for ThT) were taken every 12 s with 5 s of linear

shaking (567 cpm, 3-mm magnitude) on a Biotek Synergy H1 spectrophotometer. Readings

were collected for ~3 min before the drug was added and then readings were collected for

another 10 min.

Acknowledgements

The authors thank Alfred Spormann, Kyler Lugo, and the Huang lab for helpful discussions, Chao Jiang for help with strain construction, and Maya Farha and Eric Brown for help with membrane potential measurements. The authors acknowledge support from the Allen Discovery Center at Stanford on Systems Modeling of Infection (to H.A.A. and K.C.H.), National Institutes of Health (NIH) grants R35 GM131783 (to C.M.D. and D.B.K.) and R01 CA167250 (to C.A.D. and C.L.F.), and the Stanford Summer Research Amgen Scholars Program (to L.V.). K.C.H. is a Chan Zuckerberg Biohub Investigator.

Figure Legends

Figure 1: *B. subtilis* **3610 lyses due to oxygen depletion.**

(A) *B. subtilis* cultures lyse when not shaking. Wild-type strain 3610 (WT) was grown in

a test tube and then incubated at room temperature without shaking for 10 h.

Phase-contrast images were acquired at 0 and 10 h. Scale bar: 5 μm; arrowhead

points to cell debris.

(B) Oxygen is depleted during exponential growth and remains low throughout

subsequent oxygen depletion while cultures are sealed. Cells were cultured with

oxygen-sensitive nanoparticles (Methods). OD_{600} (top) and the relative oxygen

levels (bottom, oxygen level at first timepoint normalized to 1) of the cultures were

measured. Lines represent the average and shading represents one standard

deviation (SD), $n=3$.

(C) LytC is necessary for lysis. Left: cultures were grown to an $OD_{600} \sim 1$ and then oxygen

was depleted at $t=0$ h and OD_{600} was monitored. Shading represents 1 SD, $n=3$.

Despite the differences in OD, the $\Delta lytC$ 3610 mutant has similar viability to wild

type ($p=0.07$ at 24 h and 0.67 at 48 h, Student's t-test). Right: phase-contrast images

of wild-type and $\Delta lytC$ cells at 24 h post-oxygen depletion. Scale bar: 5 μm;

arrowhead points to phase-gray, lysed cell.

(D) Culture lysis is strongly correlated with initial cell density when the initial

$OD_{600} > 0.45$. Representative lysis curves of 6 cultures that vary in initial OD_{600} (see

Fig. S1 for other independent replicates). Inset: the maximum lysis rate vs. initial

OD_{600}. A linear regression analysis was performed on all data with initial

$OD_{600} > 0.45$.

Figure 2: Surfactin production is necessary to maintain viability.

(A) *B. subtilis* strain 168 lyses less upon oxygen depletion than 3610. *B. subtilis* 3610 and 168 strains were grown aerobically and then depleted for oxygen at 0 h. Lines represent the average and shading represents 1 SD, *n*=7. Inset: maximum lysis rate of 3610 culture is significantly higher than that of 168 (*: *p*=1.7×10⁻⁸, Student's t-test; Methods). Circles show individual experiment values, and error bars represent 1 SD.

(B) Many 168 cells form cell-wall-less protoplasts upon oxygen depletion. Merge of phase-contrast and fluorescence images of propidium iodide (PI)-stained cells at 0, 24, and 48 h post oxygen depletion. Red indicates membrane-compromised cells.

(C) Co-culturing 168 with 3610 rescues its viability upon oxygen depletion. 168

viability in monoculture is significantly different than 3610 (*: $p<0.005$; student's t-

test). Inset: 3610 and 168 have distinct colony morphologies when plated on LB.

Error bars represent 1 SD, n=3-5.

(D) Culturing with exogenous surfactin increases lysis of 168 cultures upon oxygen

depletion. OD curves during oxygen depletion of 3610, 168, and 168 genetically

rescued for surfactin production (168sfp+) or with 48 µM exogenous surfactin

added before growth (srf$_{inoc}$) or at depletion (srf$_{dep}$). Lines represent the average

and shading represents 1 SD, n=3-5. Inset: maximum lysis rates (*: $p <0.001$;

Student's t-test).

(E) Culturing with exogenous surfactin eliminates protoplasts from 168. Top: surfactin

molecular structure. Bottom: phase-contrast and PI fluorescence imaging at 24 h

post-oxygen depletion of 168sfp+ cells or with exogenous surfactin.

(F) Surfactin restores the viability of 168 cultures to near 3610 levels upon oxygen

depletion. Error bars represent 1 SD, n = 3-5. Surfactin-treated 168 (srf$_{inoc}$, srf$_{dep}$,

and 168sfp+) cultures are each significantly different than 168 ($p<0.001$ at 24 h,

$p<0.01$ at 48 h, Student's t-test). By contrast, the viability of surfactin-treated 168

cultures are not significantly different than that of 3610 at 24 h ($p>0.2$, Student's t-

test).

Figure 3: Surfactin restores the growth yield of 168 due to its detergent properties.

(A) 3610 aerobic cultures achieve a higher growth yield than 168. Lines represent the
average and shading represents 1 SD, $n=3$. *: time period over which 168 growth
differed significantly from that of 3610 ($p<0.05$, Student's t-test).

(B) Surfactin addition rescues growth yield. Growth curves of 168 with surfactin
restored genetically (168sfp+) or 48 µM added exogenously at inoculation
(168+srf$_{0h}$) or at t = 3 h (168+srf$_{3h}$). Lines represent the average and shading
represents 1 SD, $n=3$. The mean 3610 growth curve from (A) is shown as dotted
blue line. Inset: zoom-in to the period of growth divergence.

(C) Tween 80 addition rescues growth yield in a concentration-dependent manner.

Lines represent the average and shading represents 1 SD, *n*=3. The mean 3610

growth curve from (A) is shown as dotted blue line. Inset: zoom-in to the period of

growth divergence.

(D) Surfactin addition increases oxygen levels during late exponential phase. Relative

oxygen levels (bottom) during growth (top) of 168 with surfactin (48 µM) or Tween

80 (0.76 mM) added. *: time period over which oxygen levels of 168+surfactin

(yellow) or 168+Tween 80 (black) were significantly different from 168 cultures

(*p*<0.05, Student's t-test).

Figure 4: Transposon mutagenesis identifies genes that impact lysis during oxygen

depletion.

(A) Schematic of regulation of the surfactin synthetase gene operon (*srfAA-AD*). Known

regulators of SrfAA are shown in green. Our data suggests flagellar proteins (purple)

may also regulate surfactin.

(B) Surfactin-treated cultures of the transposon-disrupted mutants have fewer

protoplasts and cell debris. Phase-contrast images of mutants depleted of oxygen

for 24 h with and without 48 μM exogenous surfactin. Top: arrowheads show

636 protoplasts, phase-gray dead cells, and cell debris, all of which were not observed in

637 the surfactin-treated cultures.

638 (C) Transposon hits exhibit faster lysis when treated with exogenous surfactin. Black

639 curves are without surfactin, medium and light gray curves are with 24 μM and 48

640 μM surfactin, respectively. The dashed line is the parent (3610).

Figure 5: Surfactin maintains viability upon oxygen depletion by depolarizing the membrane.

(A) Surfactin depolarizes the membrane in *B. subtilis*. Membrane potential assays of 168

cells using the dyes $DiSC_3(5)$ (top) and ThT (bottom). The time of addition of

surfactin (48 µM), valinomycin (50 µM) and CCCP (5 µM) is marked by the black

arrowhead. One representative experimental replicate is shown (other replicates

are in Fig. S4).

(B) Treatment with the membrane depolarizing agents valinomycin (5 µM) and CCCP (5

µM) restore plating efficiency of *B. subtilis* 168 after oxygen depletion, similar to

surfactin (48 µM). Error bars represent 1 SD, *n*=3-5. 168 plating efficiency data are

significantly different than those of 168+surfactin, 168+valinomycin, 168+CCCP,

and 3610 (*p*<0.005, Student's t-test).

(C) Valinomycin and CCCP-treated 168 cultures exhibit protoplasts, demonstrating that

protoplast removal is not necessary for viability enhancement. Overlays of phase-

contrast and PI fluorescence (red) images at 24 h post-oxygen depletion. Scale bar:

5 μm.

Supplemental Figures

Figure S1: Characterization of depletion in an anaerobic chamber and of Δ*lytC* and Δ*lytD* mutants. (Related to Figure 1)

(A) Culture lysis depends on cell density at the time of oxygen depletion. Two sets of

independent lysis curves that vary in initial OD_{600} are shown; see Fig. 1D for the

other independent experiments.

(B) Cell lysis is more rapid under anaerobic conditions. Cultures were grown in 96-well

plates to an $OD_{600}\sim1$ and then transferred to an anaerobic chamber to rapidly

remove oxygen from the culture. Line represents the average and shading

represents 1 SD, *n*=3.

(C) Δ*lytD* cells show similar rates of lysis to wild-type (WT) 3610 cells upon oxygen

depletion, unlike Δ*lytC* cells that exhibit slower lysis. Strains were grown to an

OD600 ~1 and then oxygen was depleted at $t = 0$.

(D) After oxygen-depletion, the majority of WT cells but only a small subset of Δ*lytC*

cells resume growth. After 24 h of oxygen depletion, WT and Δ*lytC* cultures were

spotted onto an LB pad with oxygen and cell growth was monitored. Nearly all WT

cells grew but only a few Δ*lytC* cells grew (arrowheads). Images shown are frames

from Supplemental movies 1 and 2.

Figure S2: Oxygen depletion causes production of wall-less protoplasts in 168 and a

low percentage of cells that are able to resume growth. (Related to Figure 2)

(A) Cell-wall staining reveals that protoplasts have lost their cell wall. Phase-contrast

and HADA (cell wall) staining of 168 cultures at 24 h post-oxygen depletion. Cells

were stained with HADA for the last doubling before oxygen was depleted and the

dye was present during the duration of depletion. Orange arrowheads: phase-dark

spherical cells that lack cell wall staining. Scale bar: 5 μm.

(B) 168 Δ*lytC* mutants remain rod-shaped following oxygen depletion. Left: OD_{600} of

168 Δ*lytC* exhibited somewhat slower lysis than the parent. Right: overlay of phase-

contrast and propidium iodide staining of 168 Δ*lytC* cells.

(C) Almost all 3610 cells grow once oxygen is restored following 24 h of oxygen

depletion, but only ~1% of 168 cells grow in similar circumstance. Cells from 3610

and 168 cultures after 24 h of oxygen depletion were spotted onto an LB pad and

imaged in time-lapse to determine which cells were capable of growing and

dividing. For 3610, the arrowhead points to a rare rod-shaped cell that did not

grow. For 168, the arrowhead points to a rare rod-shaped cell that grew. Scale bar:

10 μm. n = 536 and 604 rod-shaped cells for 3610 and 168, respectively. Images

shown are frames from Movies S3 and S4.

Figure S3: Characterization of hits from transposon screen. (Related to Figure 4)

(A) Schematic of transposon screen design. A transposon mutant library was grown to
an OD_{600}~1, and then oxygen was depleted. We screened cells that were able to
recover after oxygen depletion for oxygen depletion phenotypes.

(B) Transposon mutants have a reduced lysis rate. OD_{600} curves during oxygen
depletion of cultures related to surfactin regulation (green) or flagella (purple).
Lines represent the average and shading represents 1 SD, n=5. Inset: maximum lysis
rates. Error bars represent 1 SD, n=5.

706   (C) Plating efficiency of clean deletions of *srfAA, fliI,* and *rghR* are lower than that of

707   3610 2 and 4 days following oxygen depletion.

708   (D) Addition of 48 μM surfactin increases the lysis rate of hits from transposon mutant

709   screen.

Figure S4: Membrane potential assays of 168 cells. (Related to Figure 5)

DiSC$_3$(5) (top) and ThT (bottom) both show that surfactin (48 μM), valinomycin (50

μM) and CCCP (5 μM) (time of addition marked by the black arrow) cause

depolarization. Two experimental replicates are shown (replicate 1 is shown in Fig. 5).

Supplemental Movie Legends

Supplemental Movie 1: The majority of oxygen-depleted cells from strain 3610 grow once

oxygen is restored. Cells from a 24 h oxygen-depleted wild-type 3610 culture were spotted

onto an LB pad and imaged every 5 min at 37 °C. This experiment was done in tandem with

the 3610 Δ*lytC* cultures in Movie S2.

Supplemental Movie 2: A small subset of oxygen-depleted cells in the 3610 Δ*lytC* cultures

grow once oxygen is restored. Cells from a 24 h oxygen-depleted Δ*lytC* culture were spotted

on an LB pad and imaged every 5 min at 37 °C. This experiment was done in tandem with

the 3610 cultures in Movie S1.

Supplemental Movie 3: The majority of oxygen-depleted cells from strain 3610 grow once

oxygen is restored. This experiment was carried out as in Movie S1, and was done in

tandem with the 168 cultures in Movie S4.

Supplemental Movie 4: A very small subset of oxygen-depleted, rod-shaped cells from

strain 168 grow once oxygen is restored. Cells from a 24 h oxygen-depleted culture were

spotted on an LB pad and imaged every 5 min at 37 °C. This experiment was done in

tandem with the 3610 cultures in Movie S3.

Supplemental Tables

Table S1: Information about hits from transposon screen.

Gene hit	# independent hits	Gene function
srfAA	1	surfactin synthetase
comA	3	*comP/comA* quorum sensing response regulator
comP	4	*comP/comA* quorum sensing sensor histidine kinase
rghRA	1	transcriptional repressor
*rghRA-*upstream	1	transcriptional repressor
fliI	1	flagella chaperone
fliF	1	flagella chaperone
fliJ	1	flagella basal ring

 Table S2: List of strains used in this study.

Strain name	Nickname	Genotype	Reference or source
HA10	3610	*trpC+, rapP+, sfp+, epsC+, swrA+, degQ+, pBS32*	Kearns lab
DK5073	3610 *ΔlytC*	3610, *ΔlytC*	This work
DK5075	3610 *ΔlytD*	3610, *ΔlytD*	This work
HA1	168	*trpC2*	Carol Gross lab
BKE35620	168 *ΔlytC*	168, *ΔlytC::MLS*	[14]
HA1417	168*sfp+*	168, *swrAfs, sfp⁰, amyE::Psfp-sfp cmR*	This work
HA1161	parent	3610, *ΔSPB, ΔPBSX, ΔpBS32*	Kearns lab
HA1235	N/A	3610, *ΔSPB, ΔPBSX, ΔpBS32, pMarA-kan*	This work
HA1414	N/A	168, *pMarA-kan*	This work
Back-crossed transposon mutagenesis strains			
HA1225	*comA*	3610, *ΔSPB, ΔPBSX, ΔpBS32, comA::pMarA-kan*	This work
HA1227	*comP*	3610, *ΔSPB, ΔPBSX, ΔpBS32, comP::pMarA-kan*	This work
HA1228	*fliJ*	3610, *ΔSPB, ΔPBSX, ΔpBS32, fliJ::pMarA-kan*	This work
HA1229	*fliI*	3610, *ΔSPB, ΔPBSX, ΔpBS32, fliI::pMarA-kan*	This work
HA1233	*fliF*	3610, *ΔSPB, ΔPBSX, ΔpBS32, fliF::pMarA-kan*	This work
HA1234	*srfAA*	3610, *ΔSPB, ΔPBSX, ΔpBS32, srfAA::pMarA-kan*	This work
HA1281	*yvaN*	3610, *ΔSPB, ΔPBSX, ΔpBS32, yvaN::pMarA-kan*	This work
HA1282	*yvaN-us (upstream)*	3610, *ΔSPB, ΔPBSX, ΔpBS32, yvaN-upstream::pMarA-kan*	This work
Clean deletions of transposon mutagenesis strains			
HA1369	*ΔsrfAA*	3610, *srfAA*	This work
HA1374	*ΔfliI*	3610, *fliI*	This work
HA1377	*ΔrghRA*	3610, *rghRA*	This work

BKECG 168 strains used to create clean deletions in 3610			
BKE03480	N/A	168, *srfAA::MLS*	[14]
BKE16240	N/A	168, *fliI::MLS*	[14]
BKE33660	N/A	168, *rghRA::MLS*	[14]

 Table S3: Primers and plasmids used in this study.

Primer number	Construct name	Sequence	Reference
1427	lytCUF SalI	AGGAGGTCGACGAATTAGTCTTGATGGAAAGCAGTAT	This work
1428	lytCUR EagI	CTCCTCGGCCGAAGCTGTTGGCACAAAAAGTATGAG	This work
1425	lytCDF EagI	AGGAGCGGCCGCATCAGATGCAAGTAAATTGAAGCA	This work
1426	lytCDR BamHI	ctcctggatcccatgatttgttttgtaatacttggcat	This work
1429	lytDUR EagI	AGGAGCGGCCGAAAAAACTTAGAAAGTTGCAAATAG GCT	This work
1430	lytDUF SalI	CTCCTGTCGACTTGATATGAAGAATAGACAGTTGGCA	This work
1431	lytDDR BamHI	AGGAGGGATCCAGTATTCCTTCAGAATCAACGGGGGA AT	This work
1432	lytDDF EagI	CTCCTCGGCCGTTAGTCTCTTTTTCATTCCTTCTCCTC TT	This work
695	IPCR1	GCTTGTAAATTCTATCATAATTG	This work
696	IPCR2	AGGGAATCATTTGAAGGTTGG	This work
Plasmids		**Genotype**	**Reference**
pDP299		*ΔlytC ori^{TsBs} mls amp*	This work
pDP300		*ΔlytD ori^{TsBs} mls amp*	This work
pMiniMAD		*ori^{TsBs} mls amp*	[59]
pDR244		*ori^{Ts} , cre constitutively expressed, amp/spc*	[14]

References

1. Boutilier, R.G. (2001). Mechanisms of cell survival in hypoxia and hypothermia. J Exp Biol *204*, 3171-3181.

2. Hochachka, P.W. (1986). Defense strategies against hypoxia and hypothermia. Science *231*, 234-241.

3. Sendoel, A., and Hengartner, M.O. (2014). Apoptotic cell death under hypoxia. Physiology (Bethesda) *29*, 168-176.

4. Prescott, L.M., Harley, J.P., and Klein, D.A. (2005). Microbiology, 6th Edition, (Dubuque, IA: McGraw-Hill Higher Education).

5. Wayne, L.G., and Hayes, L.G. (1996). An in vitro model for sequential study of shiftdown of *Mycobacterium tuberculosis* through two stages of nonreplicating persistence. Infect Immun *64*, 2062-2069.

6. Rao, S.P., Alonso, S., Rand, L., Dick, T., and Pethe, K. (2008). The protonmotive force is required for maintaining ATP homeostasis and viability of hypoxic, nonreplicating *Mycobacterium tuberculosis*. Proc Natl Acad Sci U S A *105*, 11945-11950.

7. Berney, M., Greening, C., Conrad, R., Jacobs, W.R., Jr., and Cook, G.M. (2014). An obligately aerobic soil bacterium activates fermentative hydrogen production to survive reductive stress during hypoxia. Proc Natl Acad Sci U S A *111*, 11479-11484.

8. O'Toole, R., Smeulders, M.J., Blokpoel, M.C., Kay, E.J., Lougheed, K., and Williams, H.D. (2003). A two-component regulator of universal stress protein expression and adaptation to oxygen starvation in *Mycobacterium smegmatis*. J Bacteriol *185*, 1543-1554.

9.   Nakano, M.M., Dailly, Y.P., Zuber, P., and Clark, D.P. (1997). Characterization of

anaerobic fermentative growth of *Bacillus subtilis*: identification of fermentation end

products and genes required for growth. J Bacteriol *179*, 6749-6755.

10.   Pusey, P.L., and Wilson, C.L. (1984). Postharvest biological control of stone fruit

brown rot by *Bacillus subtilis*. Plant Disease *68*, 753-756.

11.   Kumar, A.S., Lakshmanan, V., Caplan, J.L., Powell, D., Czymmek, K.J., Levia, D.F., and

Bais, H.P. (2012). Rhizobacteria *Bacillus subtilis* restricts foliar pathogen entry

through stomata. The Plant Journal *72*, 694-706.

12.   Sierra, J., and Renault, P. (1998). Temporal Pattern of Oxygen Concentration in a

Hydromorphic Soil. Soil Science Society of America Journal *62*, 1398-1405.

13.   Kaufman, W., and Bauer, K. (1958). Some studies of the mechanism of the" anaerobic

autolysis" of *Bacillus subtilis* J. Gen. Microbiol. *18*.

14.   Koo, B.M., Kritikos, G., Farelli, J.D., Todor, H., Tong, K., Kimsey, H., Wapinski, I.,

Galardini, M., Cabal, A., Peters, J.M., et al. (2017). Construction and Analysis of Two

Genome-Scale Deletion Libraries for *Bacillus subtilis*. Cell Syst *4*, 291-305 e297.

15.   Peters, J.M., Colavin, A., Shi, H., Czarny, T.L., Larson, M.H., Wong, S., Hawkins, J.S., Lu,

C.H.S., Koo, B.M., Marta, E., et al. (2016). A Comprehensive, CRISPR-based Functional

Analysis of Essential Genes in Bacteria. Cell *165*, 1493-1506.

16.   Zhu, B., and Stulke, J. (2018). SubtiWiki in 2018: from genes and proteins to

functional network annotation of the model organism *Bacillus subtilis*. Nucleic Acids

Res *46*, D743-D748.

17.   Zeigler, D.R., Pragai, Z., Rodriguez, S., Chevreux, B., Muffler, A., Albert, T., Bai, R.,

Wyss, M., and Perkins, J.B. (2008). The origins of 168, W23, and other *Bacillus*

subtilis legacy strains. J Bacteriol *190*, 6983-6995.

18.   Julkowska, D., Obuchowski, M., Holland, I.B., and Seror, S.J. (2005). Comparative

analysis of the development of swarming communities of Bacillus subtilis 168 and a

natural wild type: critical effects of surfactin and the composition of the medium. J

Bacteriol *187*, 65-76.

19.   Nakano, M.M., Corbell, N., Besson, J., and Zuber, P. (1992). Isolation and

characterization of sfp: a gene that functions in the production of the lipopeptide

biosurfactant, surfactin, in Bacillus subtilis. Mol Gen Genet *232*, 313-321.

20.   Nakano, M.M., Marahiel, M.A., and Zuber, P. (1988). Identification of a genetic locus

required for biosynthesis of the lipopeptide antibiotic surfactin in *Bacillus subtilis.* J

Bacteriol *170*, 5662-5668.

21.   Arima, K., Kakinuma, A., and Tamura, G. (1968). Surfactin, a crystalline peptidelipid

surfactant produced by *Bacillus subtilis*: isolation, characterization and its inhibition

of fibrin clot formation. Biochem Biophys Res Commun *31*, 488-494.

22.   Płaza, G.A., Turek, A., Król, E., and Szczygłowska, R. (2013). Antifungal and

antibacterial properties of surfactin isolated from *Bacillus subtilis* growing on

molasses. African Journal of Microbiology Research *7*, 3165-3170.

23.   Korenblum, E., de Araujo, L.V., Guimaraes, C.R., de Souza, L.M., Sassaki, G., Abreu, F.,

Nitschke, M., Lins, U., Freire, D.M., Barreto-Bergter, E., et al. (2012). Purification and

characterization of a surfactin-like molecule produced by *Bacillus sp. H2O-1* and its

antagonistic effect against sulfate reducing bacteria. BMC Microbiol *12*, 252.

24. Rosenberg, G., Steinberg, N., Oppenheimer-Shaanan, Y., Olender, T., Doron, S., Ben-Ari, J., Sirota-Madi, A., Bloom-Ackermann, Z., and Kolodkin-Gal, I. (2016). Not so simple, not so subtle: the interspecies competition between *Bacillus simplex* and *Bacillus subtilis* and its impact on the evolution of biofilms. NPJ Biofilms Microbiomes *2*, 15027.

25. Snook, M.E., Mitchell, T., Hinton, D.M., and Bacon, C.W. (2009). Isolation and characterization of leu7-surfactin from the endophytic bacterium *Bacillus mojavensis* RRC 101, a biocontrol agent for *Fusarium verticillioides*. J Agric Food Chem *57*, 4287-4292.

26. Zhi, Y., Wu, Q., Du, H., and Xu, Y. (2016). Biocontrol of geosmin-producing Streptomyces spp. by two *Bacillus* strains from Chinese liquor. Int J Food Microbiol *231*, 1-9.

27. Olmeda, B., Villen, L., Cruz, A., Orellana, G., and Perez-Gil, J. (2010). Pulmonary surfactant layers accelerate O(2) diffusion through the air-water interface. Biochim Biophys Acta *1798*, 1281-1284.

28. Cabeen, M.T., and Jacobs-Wagner, C. (2005). Bacterial cell shape. Nat Rev Microbiol *3*, 601-610.

29. Lazarevic, V., Margot, P., Soldo, B., and Karamata, D. (1992). Sequencing and analysis of the *Bacillus subtilis lytRABC* divergon: a regulatory unit encompassing the structural genes of the N-acetylmuramoyl-L-alanine amidase and its modifier. J Gen Microbiol *138*, 1949-1961.

832   30.   Margot, P., Mauel, C., and Karamata, D. (1994). The gene of the N-
833   acetylglucosaminidase, a *Bacillus subtilis* 168 cell wall hydrolase not involved in
834   vegetative cell autolysis. Mol Microbiol *12*, 535-545.

835   31.   Koch, A.L. (2001). Autolysis control hypotheses for tolerance to wall antibiotics.
836   Antimicrob Agents Chemother *45*, 2671-2675.

837   32.   Sheppard, J.D., Jumarie, C., Cooper, D.G., and Laprade, R. (1991). Ionic channels
838   induced by surfactin in planar lipid bilayer membranes. Biochim Biophys Acta *1064*,
839   13-23.

840   33.   Laidig, K.E., Gainer, J.L., and Daggett, V. (1998). Altering Diffusivity in Biological
841   Solutions through Modification of Solution Structure and Dynamics. J Am Chem Soc
842   *120*, 9394-9395.

843   34.   Hotez, L., Dailey, J.W., Geelhoed, G.W., and Gainer, J.L. (1977). The role of oxygen
844   diffusivity in biochemical reactions. Experientia *33*, 1424-1425.

845   35.   Cosmina, P., Rodriguez, F., de Ferra, F., Grandi, G., Perego, M., Venema, G., and van
846   Sinderen, D. (1993). Sequence and analysis of the genetic locus responsible for
847   surfactin synthesis in *Bacillus subtilis*. Mol Microbiol *8*, 821-831.

848   36.   Nakano, M.M., Xia, L.A., and Zuber, P. (1991). Transcription initiation region of the
849   *srfA* operon, which is controlled by the *comP-comA* signal transduction system in
850   *Bacillus subtilis*. J Bacteriol *173*, 5487-5493.

851   37.   Hayashi, K., Kensuke, T., Kobayashi, K., Ogasawara, N., and Ogura, M. (2006). *Bacillus*
852   *subtilis* RghR (YvaN) represses *rapG* and *rapH*, which encode inhibitors of
853   expression of the *srfA* operon. Mol Microbiol *59*, 1714-1729.

854   38.   Mukherjee, S., and Kearns, D.B. (2014). The structure and regulation of flagella in

855   *Bacillus subtilis*. Annu Rev Genet *48*, 319-340.

856   39.   Ueno, T., Oosawa, K., and Aizawa, S. (1992). M ring, S ring and proximal rod of the

857   flagellar basal body of *Salmonella typhimurium* are composed of subunits of a single

858   protein, FliF. J Mol Biol *227*, 672-677.

859   40.   Evans, L.D., Stafford, G.P., Ahmed, S., Fraser, G.M., and Hughes, C. (2006). An escort

860   mechanism for cycling of export chaperones during flagellum assembly. Proc Natl

861   Acad Sci U S A *103*, 17474-17479.

862   41.   Minamino, T., Chu, R., Yamaguchi, S., and Macnab, R.M. (2000). Role of FliJ in flagellar

863   protein export in *Salmonella*. J Bacteriol *182*, 4207-4215.

864   42.   Farha, M.A., Verschoor, C.P., Bowdish, D., and Brown, E.D. (2013). Collapsing the

865   proton motive force to identify synergistic combinations against *Staphylococcus*

866   *aureus*. Chem Biol *20*, 1168-1178.

867   43.   Prindle, A., Liu, J., Asally, M., Ly, S., Garcia-Ojalvo, J., and Suel, G.M. (2015). Ion

868   channels enable electrical communication in bacterial communities. Nature *527*, 59-

869   63.

870   44.   Varma, S., Sabo, D., and Rempe, S.B. (2008). K+/Na+ selectivity in K channels and

871   valinomycin: over-coordination versus cavity-size constraints. J Mol Biol *376*, 13-22.

872   45.   Strahl, H., Burmann, F., and Hamoen, L.W. (2014). The actin homologue MreB

873   organizes the bacterial cell membrane. Nat Commun *5*, 3442.

874   46.   Branda, S.S., Gonzalez-Pastor, J.E., Ben-Yehuda, S., Losick, R., and Kolter, R. (2001).

875   Fruiting body formation by *Bacillus subtilis*. Proc Natl Acad Sci U S A *98*, 11621-

876   11626.

877   47.   Pham, J.V., Yilma, M.A., Feliz, A., Majid, M.T., Maffetone, N., Walker, J.R., Kim, E., Cho,

878   H.J., Reynolds, J.M., Song, M.C., et al. (2019). A Review of the Microbial Production of

879   Bioactive Natural Products and Biologics. Front Microbiol *10*, 1404.

880   48.   Hu, J., Lei, P., Mohsin, A., Liu, X., Huang, M., Li, L., Hu, J., Hang, H., Zhuang, Y., and Guo,

881   M. (2017). Mixomics analysis of Bacillus subtilis: effect of oxygen availability on

882   riboflavin production. Microb Cell Fact *16*, 150.

883   49.   Garcia-Ochoa, F., and Gomez, E. (2009). Bioreactor scale-up and oxygen transfer rate

884   in microbial processes: an overview. Biotechnol Adv *27*, 153-176.

885   50.   Baeshen, N.A., Baeshen, M.N., Sheikh, A., Bora, R.S., Ahmed, M.M., Ramadan, H.A.,

886   Saini, K.S., and Redwan, E.M. (2014). Cell factories for insulin production. Microb

887   Cell Fact *13*, 141.

888   51.   Arjes, H.A., Kriel, A., Sorto, N.A., Shaw, J.T., Wang, J.D., and Levin, P.A. (2014). Failsafe

889   mechanisms couple division and DNA replication in bacteria. Curr Biol *24*, 2149-

890   2155.

891   52.   Rojas, E.R., Huang, K.C., and Theriot, J.A. (2017). Homeostatic Cell Growth Is

892   Accomplished Mechanically through Membrane Tension Inhibition of Cell-Wall

893   Synthesis. Cell Syst *5*, 578-590 e576.

894   53.   Blackman, S.A., Smith, T.J., and Foster, S.J. (1998). The role of autolysins during

895   vegetative growth of *Bacillus subtilis* 168. Microbiology *144 (Pt 1)*, 73-82.

896   54.   Calamita, H.G., Ehringer, W.D., Koch, A.L., and Doyle, R.J. (2001). Evidence that the

897   cell wall of Bacillus subtilis is protonated during respiration. Proc Natl Acad Sci U S

898   A *98*, 15260-15263.

899   55.   Mueller, E.A., Egan, A.J., Breukink, E., Vollmer, W., and Levin, P.A. (2019). Plasticity of

900   *Escherichia coli* cell wall metabolism promotes fitness and antibiotic resistance

901   across environmental conditions. Elife *8*.

902   56.   Peters, K., Kannan, S., Rao, V.A., Biboy, J., Vollmer, D., Erickson, S.W., Lewis, R.J.,

903   Young, K.D., and Vollmer, W. (2016). The Redundancy of Peptidoglycan

904   Carboxypeptidases Ensures Robust Cell Shape Maintenance in *Escherichia coli*. MBio

905   *7*.

906   57.   Stratford, J.P., Edwards, C.L.A., Ghanshyam, M.J., Malyshev, D., Delise, M.A., Hayashi,

907   Y., and Asally, M. (2019). Electrically induced bacterial membrane-potential

908   dynamics correspond to cellular proliferation capacity. Proc Natl Acad Sci U S A *116*,

909   9552-9557.

910   58.   Yasbin, R.E., and Young, F.E. (1974). Transduction in *Bacillus subtilis* by

911   bacteriophage SPP1. J Virol *14*, 1343-1348.

912   59.   Patrick, J.E., and Kearns, D.B. (2008). MinJ (YvjD) is a topological determinant of cell

913   division in *Bacillus subtilis*. Mol Microbiol *70*, 1166-1179.

914   60.   Arnaud, M., Chastanet, A., and Debarbouille, M. (2004). New vector for efficient

915   allelic replacement in naturally nontransformable, low-GC-content, gram-positive

916   bacteria. Appl Environ Microbiol *70*, 6887-6891.

917   61.   DeRosa, C.A., Samonina-Kosicka, J., Fan, Z., Hendargo, H.C., Weitzel, D.H., Palmer,

918   G.M., and Fraser, C.L. (2015). Oxygen Sensing Difluoroboron Dinaphthoylmethane

919   Polylactide. Macromolecules *48*, 2967-2977.

920   62.   DeRosa, C.A., Seaman, S.A., Mathew, A.S., Gorick, C.M., Fan, Z., Demas, J.N., Peirce,

921   S.M., and Fraser, C.L. (2016). Oxygen Sensing Difluoroboron beta-Diketonate

Polylactide Materials with Tunable Dynamic Ranges for Wound Imaging. ACS Sens *1*,

1366-1373.

63.   Shi, H., Colavin, A., Lee, T.K., and Huang, K.C. (2017). Strain Library Imaging Protocol:

high-throughput, automated single-cell microscopy for large bacterial collections

arrayed on multiwell plates. Nature Protocols.

64.   Kuru, E., Hughes, H.V., Brown, P.J., Hall, E., Tekkam, S., Cava, F., de Pedro, M.A., Brun,

Y.V., and VanNieuwenhze, M.S. (2012). In Situ probing of newly synthesized

peptidoglycan in live bacteria with fluorescent D-amino acids. Angew Chem Int Ed

Engl *51*, 12519-12523.

931