Out of the blue; Phototropins of the leaf vascular bundle sheath

mediate the blue light regulation of the leaf hydraulic conductance

Yael Grunwald[*], **Sanbon Chaka Gosa**[*], **Tanmayee Torne**, **Nava Moran**, **Menachem**

Moshelion[#]

The Robert H. Smith Institute of Plant Sciences and Genetics in Agriculture, The Robert H. Smith Faculty of

Agriculture, Food and Environment, The Hebrew University of Jerusalem, Rehovot 76100, Israel

*Both authors contributed equally to this work

[#]Correspondence: menachem.moshelion@mail.huji.ac.il

ABSTRACT

The leaf vascular bundle sheath cells (BSCs), which tightly envelop the leaf veins,

constitute a selective dynamic barrier to water and solutes radially entering the mesophyll

and play a major role in regulating the leaf radial hydraulic conductance (K_{leaf}). Recently,

we showed that the BSCs' plasma membrane H^+-ATPase, AHA2, increases K_{leaf} by

acidifying the xylem sap. Since BL reportedly increases K_{leaf} and we found the blue light

(BL) receptor genes, *PHOT1* and *PHOT2* expressed in the Arabidopsis BSCs, we

hypothesized that, similar to the guard cells (GCs) BL signal transduction pathway, the

BSCs' PHOT1 and PHOT2 activate the BSCs' H^+-ATPase and thus regulate K_{leaf}. Indeed,

under BL illumination, the K_{leaf} in the knockout mutant lines *phot1-5*, *phot2-1, phot1-*

5phot2-1 and *aha2-4* was lower than in WT. BSCs-directed complementation (using the

SCR promoter) of *phot1-5* and *aha2-4* respectively by *PHOT1* and *AHA2*, restored the

BL-induced K_{leaf} increase. BSCs-specific silencing of *PHOT1* or *PHOT2* (using the SCR

promoter) abolished the BL-induced K_{leaf} increase. Xylem-fed PHOT inhibitor, tyrphostin

9, also abolished the BL-induced K_{leaf} increase in WT. Moreover, in WT plants, white

light (WL) acidified the xylem sap compared to dark, but did not acidify the xylem sap of

the *phot1-5* mutant. BSCs-specific complementation of *phot1-5* by SCR: PHOT1,

restored the WL-induced xylem acidification. On a cellular level, BL hyperpolarized the

BSCs, which was prevented by tyrphostin 9, and alkalinized their cytosol. In addition, the

osmotic water permeability coefficient (P_f) of the BSCs was higher under WL treatment.

Our results link the blue light control of water fluxes from the xylem to the mesophyll via

the BSCs in the following model:

BL →BSCs' *PHOT*s activation →tyrosine phosphorylation→BSCs' H^{+-} ATPase

activation →BSCs hyperpolarization and cytosol alkalinization, xylem acidification →P_f

elevation → K_{leaf} increase. This independent BL signal transduction pathway that we

revealed within the bundle sheath explains – at least in part – how light increases the leaf

hydraulic conductance.

Introduction

Light, the energy source for photosynthesis, has also evolved as a signal that regulates

growth and development (Kronenberg and Kendrick, 1986; Briggs and Huala, 1999) as well

as physiological traits such as stomatal conductance [(g_S); (Zeiger and Helper, 1977;

Karlsson, 1986; Kinoshita et al., 2001; Talbott et al., 2003; Van Ieperen et al., 2012)] and

leaf hydraulic conductance [(K_{leaf}); (Voicu et al., 2008; Ben Baaziz et al., 2012; Aasamaa

and Sõber, 2012; Prado and Maurel, 2013)]. One of the most conserved and well-studied

light-sensing mechanisms is that of the blue light (BL, 390-550 nm) which invokes stomatal

opening (Grondin et al., 2015). In this signal transduction pathway, BL is perceived by the

guard cells (GCs) light-activated protein kinases PHOT1 and PHOT2 (Briggs and Christie,

2002; Kinoshita et al., 2001), eventually activating the plasma membrane H^+-ATPases

(Kinoshita and Shimazaki, 1999; Svennelid et al., 1999), which, in turn, hyperpolarizes the

GCs and acidifies their surrounding apoplast [(Ueno et al., 2005; Den Os et al., 2007;

Elmore and Coaker, 2011; Kinoshita and Shimazaki, 1999), recently reviewed by Inoue &

Kinoshita (2017)]. Consequently, the hyperpolarization-activated inward-rectifying K^+

channels (K_{in}) open, enabling potassium ions (K^+) influx driven by K^+ electrochemical

potential difference. Inward K^+ flux results in an osmolyte accumulation reducing the water

potential inside the cell, which, in turn, drives an influx of water leading to GCs swelling

and stomatal opening (Assmann, 1993; Roelfsema and Hedrich, 2005; Shimazaki et al.,

2007; Oishi et al., 2010; Yamauchi et al., 2016). Conversely, at night, in the dark, the guard

cells' H^+-ATPases are deactivated, the GCs are depolarized and their apoplast becomes

alkaline (Kinoshita et al., 2001; Kinoshita and Shimazaki, 1999).

Blue light or white light (WL) illumination has also been shown to increase the hydraulic

conductance (K_{leaf}) of the entire leaf in several plant species (Voicu et al., 2008 (bur oak);

Voicu et al., 2009; Sellin et al., 2011 (silver birch); Aasamaa and Sber, 2012 (deciduous

trees); Ben Baaziz et al., 2012 (walnut)). Interestingly, K_{leaf} showed a faster response to

light than g_S (Guyot et al., 2012). However, the molecular mechanism of K_{leaf} increase by

light quality is not yet fully understood.

In the past decade, it has been established that the bundle sheath cells (BSCs), a

parenchymatous layer which tightly enwraps the entire leaf vasculature, can act as a

dynamic and selective xylem-mesophyll barrier to water and ions, which participates in

K_{leaf} regulation (Shapira et al., 2009; Shatil-Cohen and Moshelion, 2012; Sade et al., 2014;

Wigoda et al., 2014, 2017; Grunwald et al., 2021). Moreover, we recently discovered that

the BSCs' H^+-ATPase proton pump, AHA2, participated in regulating K_{leaf} via changes in

the xylem pH and the positive correlation between AHA2-driven xylem acidification and

K_{leaf} was due to an increase in the osmotic water permeability of the BSCs membrane

(Grunwald et al., 2021). AHA2 is expressed particularly abundantly in the BSCs (at an

over-three-fold higher level as compared to the neighboring mesophyll cells; Wigoda et al.,

2017, GEO repository Experiment GSE85463), explaining how such acidification is

possible. In addition, the same BSC transcriptome analysis (ibid.) revealed a substantial

expression of the BL receptors PHOT1 and PHOT2 in the BSCs, as well as their immediate

phosphorylation substrate, BLUS1 [(Blue Light Signaling1, At4g14480; (Takemiya et al.,

2013)] and the BLUS1-interacting kinase, BHP (Blue-light-dependent H^+-ATPase

Phosphorylation, At4g18950), the elements of early BL signaling in the guard cells

(reviewed by Inoue and Kinoshita, 2017). These findings led us to hypothesize that a BL

signal transduction pathway similar to that which is active in opening stomata, is active also

in the vascular BSCs and that it has a role in the regulation of K_{leaf}. In confirmation of this

hypothesis, we describe here a few of the physiological and molecular details of the

mechanism underlying the blue light-dependent K_{leaf} regulation by BSCs.

Materials and Methods

Plant material

Plant types. We used WT (wild type) *Arabidopsis thaliana* plants ecotype Colombia, Col-

0 (aka Col), the T-DNA insertion AHA2 mutants *aha2-4* (SALK_ 082786) (Col) and *aha2-*

4 complemented with *SCR: AHA2* (line 56) (Col), and *SCR:GFP* (Col) as already described

(Grunwald et al., 2021). In addition, we used WT Arabidopsis (Col) with the Glabrous

mutation (WT Col-gl), *phot1-5 (nph1-5)* (Col-gl), *phot 2-1(npl1-1 or cav1-1)* (Col-gl) and

the double mutant *phot1-5phot2-1 (nph1-5 npl1-1)* (Col-gl) (Kagawa et al., 2001), which

were kindly provided by the Ken-Ichiro Shimazaki lab (Tokyo, Japan).

Construction of transgenic plant lines. *SCR:mirPHOT* plants**:** The premiR-PHOT1or

PHOT2 and synthetic genes were synthesized by Hylabs (Rehovot, Israel), based on a

premiR164 backbone (Alvarez et al., 2006). We used the Web-based MicroRNA Designer

(WMD, http://wmd3.weigelworld.org) to produce a premiRNA gene MIR319a as

described in WMD. After sequence verification, the premiR-PHOT1or premiR-PHOT2

were cloned into the pDONR™ 221 and the SCR promoter into pDONRP4P1r (Invitrogen)

vectors which are Gateway® compatible by BP reactions, and later cloned into the

pB7M24GW (Invitrogen) two-fragment binary vector by LR reaction according to the

manufacturer's instructions. Each binary vector was transformed into *Agrobacterium*

tumefaciens by electroporation and transformed to WT Col0 using the floral dip method

(Clough and Bent, 1998). Transformants were selected based on their BASTA (Glufosinate

ammonium, Sigma cat # 45520) resistance, grown on plates with MS (Murashige and

Skoog, Duchefa cat# M222.0050) basal medium + 1 % sucrose and 20 µg/ml BASTA.

DNA insertion was verified in selected lines by PCR targeting the junction of the premiR-

gene and the 35S terminator with the forward primer located about 1000bp from the 3' end

of premiR-gene and reverse primer on the 35S terminator (see primer list in supplemental

Table S1), PCR fragments were then sequenced and verified.

SCR: PHOT1 plants. Binary vectors were constructed with the *PHOT1* gene as described

above and then transformed into *phot1-5* (Col-gl1) plants, and successful transformation

was verified by PCR.

Plant growth conditions. Plants were grown in soil (Klasmann686

Klasmann-Deilmann, Germany) + 4g/l Osmocote® 6M in a growth chamber at 22 °C and

70% relative humidity, and 8-h light/16-h dark photoperiod. During the day, the

illumination, humidity and VPD changed in a pattern resembling natural conditions, as in

(Negin and Moshelion, 2017). The illumination intensity provided by LED lights strips

(EnerLED 24 V-5630, 24 W/m, 3000 K (50%)/6000 K (50%)) reached up to 150-200 µmol

m^{-2} sec^{-1} light at the plant level. The plants were irrigated twice a week.

Detached leaves preparation for gas exchange measurements. Fully expanded leaves

from 7-8- week-old plants were excised at the petiole base using a wet sharp blade under a

drop of water. Petioles were immediately submerged in 0.5 ml Eppendorf tubes with

artificial xylem solution (AXS; 3 mM KNO_3, 1 mM $Ca(NO_3)_2$, 1 mM $MgSO_4$, 3 mM $CaCl_2$,

0.25 mM NaH_2PO_4, 90 µM EDFS (Sigma). The leaves were excised, shortly before "lights

OFF" transition (around 5:00 PM) on the evening preceding the measurements and placed

in gas-sealed transparent 25 cm x 20 cm x 15 cm plastic boxes with water-soaked tissue

paper on the bottom to provide ~90% humidity. The transparent boxes were then inserted

into a larger light-proof plastic box overnight and kept in total darkness until the start of

light treatments in the next morning.

Light treatments of detached leaves

All gas exchange experiments were conducted in a dark room (<1 umol m^{-2} s^{-1}) at a constant

temperature of 22 °C, in the morning between 10:00 AM and 1:00 PM. The excised leaves

were taken out of the light-sealed box, and placed in one of two custom-made light

chambers (either red light (RL) or combined red and blue light (RL+BL), as specified

further down). Leaves were exposed to either RL or RL+BL for 15 minutes in ambient

vapor pressure deficit (VPD) of 1.3-1.5 kPa. A fan circulated the air in each light chamber

for uniform and constant temperature and VPD. Next, leaves were transferred to a **LI-COR**

6400XT for an additional 2-3 minutes, under the same illumination conditions with VPD

of about 1.3kPa, until stabilization of the gas exchange measurements, after which the

measurement was recorded. The total illumination intensity in both light chambers was set

to roughly 220 µmol m^{-2} s^{-1}. In the RL chamber the leaves were illuminated only with red

light (660 nm) and in the 'RL+BL' chamber they were illuminated with approximately 90%

red light and 10% blue light (450 nm). D-LED lighting with adjustable current source (DR-

SD6/12) was used as light sources (http://www.d-led.net/). Light intensity was verified

daily using a LI-250A light meter with a LI-190SA quantum sensor (LI-COR, Lincoln, NE,

USA).

Determination of gas exchange, E and g$_S$, and hydraulic conductance, K$_{leaf}$

Immediately after the light treatments, the leaves were placed in a LI-COR 6400XT for

measurements of stomatal conductance (g$_S$) and transpiration (E) similar to Grunwald et

al., 2021, with the following changes: all the experiments were conducted in the dark at 22

C^0 and the illumination in the LI-COR cuvette was adjusted to match the preceding light

treatments. Immediately thereafter, the water potential (Ψ_{leaf}) was determined in a pressure

chamber (ARIMAD-3000; MRC Israel) and K_{leaf} was calculated (as in Grunwald et al.,

2021) by:

Eq. 1: $K_{leaf} = -E / (\Psi_{leaf} - \Psi_{AXS}) \approx -E / \Psi_{leaf}$,

Additionally, g_s was measured at the same time as above using LI-COR 6400XT on intact

leaves of whole plants left under the growth-room illumination for 2-4 hours after Lights

ON. Illumination settings in the instrument's cuvette were set to the same conditions as in

the growth room.

Inhibition of light transduction in detached leaves

Tyrphostin 9 (ENZO, cat. # BML-EI21, 100 mM in DMSO (Sigma, cat # W387520), kept

in aliquots at -18 °C), was added to the AXS to a final concentration of 10 µM. AXS with

DMSO at a concentration of 100 µl/l (vol/vol) served as a control in this set of experiments.

Excised leaves were kept overnight until the measurements as described above. For surface

application, 0.05% of Tween 20 (J.T. Baker cat# X251-07) was added to the Tyrphostin 9

and control solutions and sprayed prior to overnight perfusion with unmodified AXS.

Boxes holding the sprayed leaves did not contain the damp tissue paper, and leaves were

patted dry prior to placing them in the light chamber to ensure the leaf surface is dry when

placed in LI-COR 6400xt cuvette.

Determination of xylem sap pH in detached leaves

Leaf preparation for imaging. On the eve of the experiment, leaves from 6-7 week old

plants, approximately 2.5 cm long and 1 cm wide, were excised with a sharp blade and

perfused (as described) with AXS containing 100 µM of the dual-excitation fluorescent pH

probe, FITC-D (fluorescein isothiocyanate conjugated to 10 kD dextran, (Sigma cat. #:

FD10S), added from a 10 mM stock in DDW). The excised leaves were then placed in the

"double box" setup (i.e. sealed plastic box inside a dark box) and kept in the dark until

measurements the next morning.

Dark-treated leaves: Leaves were taken out of the dark boxes and immediately prepared

for vein imaging on a microscope slide under low light conditions (<5 μmol m^{-2} sec^{-1}).

Light-treated leaves: Leaves were taken out of the dark boxes and placed in the growth

chamber inside transparent gas sealed boxes (boxes as in K_{leaf}) for 30 minutes of growth

room white light illumination (see above under "Plant growth conditions") and were

subsequently imaged.

The leaves were imaged using an inverted microscope (Olympus-IX8, as detailed by

Grunwald et al., 2021). Image capture and image analysis of the intra-xylem pH were as

already described (ibid.).

Determination of the BSCs membrane potential

BSCs protoplasts imaging. For the evaluation of protoplast membrane potential, we used a

fluorescent membrane-potential-sensitive dual-excitation ratiometric dye Di-8-ANEPPS

(Pucihar et al., 2009) and an inverted epifluorescence microscope (Eclipse Ti-S, Nikon,

Tokyo, Japan) coupled to an IXON Ultra 888 camera (Andor, UK). BSC protoplasts

isolated (Shatil-Cohen et al., 2014) from 6-8 weeks old SCR:GFP (Col) plants were kept at

room temperature in the dark. A sample with a few protoplasts was placed in a bath solution

in an experimental chamber, containing 30 μM di-8-ANEPPS and 1% pluronic acid,

without or with 10 µM tyrphostin 9 (see Solutions below), and allowed to settle for 10 min

on a glass coverslip bottom. A Nikon 40X objective (Plan Fluor 40x/ 0.75, OFN25 DIC

M/N2) was used for protoplast viewing and imaging. An individual, perfectly round BSC

(25-32.5 μM diameter) was selected for further imaging based on its GFP fluorescence

(excitation of 490/15 nm was delivered by a xenon lamp monochromator, Polychrome II

(Till Photonics, Munich, Germany), under the control of IW6.1 (Imaging Workbench 6.1)

software (Indec Biosystems, Santa Clara, CA), and the emitted fluorescence was filtered

via a 515 nm dichroic mirror and a 525/50 nm band-pass barrier filter).

Protoplasts light treatment and imaging. The selected BSC protoplast in the experimental

chamber was then darkened for 10-20 min (Control), and a first pair of images of di-8-

ANEPPS fluorescence was recorded (excitation by a pair of 50 ms pulses, 438 nm then 531

nm, 3 ms apart, was delivered by the Polychrome II; the emitted fluorescence was filtered

via a dichroic mirror of 570 nm and 585/20 nm emission band-pass filter (Chroma

Technology Corp., Bellows Falls, VT, USA)). Subsequently, the protoplast was illuminated

for 5-10 min (by a home-made illuminator of crystal clear LEDs from Tal-Mir electronics,

Israel) with either RL (660 nm, 220 µmol m^{-2} s^{-1}) or RL+BL (220 µmol m^{-2} s^{-1}, BL 450 nm,

10-15% of the total intensity) and a second pair of fluorescence images was recorded as

above. The images were processed using FIJI (Abràmoff et al., 2004; Schindelin et al.,

2012) to yield a pixel-by-pixel fluorescence ratio as described earlier (Wigoda et al., 2017).

Fluorescence ratio calibration. BSCs protoplasts incubated in di-8-ANEPPS as above were

subjected to 5 s-long voltage pulses in the range of +17 to -223 mV using a patch-clamp

pipette in a whole-cell configuration. A pair of images were recorded during the 2nd half of

each pulse, during a steady-state. The patch-clamp-imaging setup was described in detail

by Wigoda et al. (2017), except for the current substitution of the image acquisition

instruments with the abovementioned Eclipse microscope and IXON camera. This

calibration confirmed the positive correlation between the di-8-ANEPPS fluorescence ratio

and depolarization (Suppl. Fig.8).

Solutions. <u>Di-8-ANEPPS</u> (Molecular probes, Life technologies, cat. # D3167, Oregon,

USA): a 10 mM stock solution in DMSO was stored in 10 μL aliquots at -20 °C.

<u>Pluronic F-127</u> (Molecular probes, Life technologies, cat. # P6867, Oregon, USA), a 20%

stock solution in DMSO was stored in 100 uL aliquots at RT.

<u>Bath Solution (in mM)</u>: 5 KCl, 1 $CaCl_2$, 4 $MgCl_2$, 10 MES; pH 5.6; osmolality: 435 mOsm,

adjusted w/ D-sorbitol.

<u>Patch-clamp pipette solution (in mM)</u>:112 K-Gluconate, 28 KCl. 2 $MgCl_2$, 10 HEPES, 328

Sorbitol, pH 7.5, Osmolarity: 480 mOsm.

<u>Tyrphostin 9</u> was added to the bath solution from aliquots used in the detached leaves

experiments.

Determination of the BSCs cytosolic pH (pH$_{CYT}$)

The experiments were performed between 10 AM and 4 PM. BSC protoplasts were isolated

(Shatil-Cohen et al., 2014) from 6-8 weeks old SCR:GFP (Col) plants and kept in an

Eppendorf vial at room temperature (22 ± 2 °C) under dark until use. The protoplasts were

treated as already detailed (Torne et al., 2021) with the following modifications: A 10 min

'dark treatment' RL+BL or RL illumination treatments (illumination setting as described

above for membrane potential experiments) were applied and only the unmodified bath

solution was used for flushing out the external SNARF-AM during the last 1 min of the

dark/ light treatments. . Image acquisition and analysis to determine the pH$_{CYT}$ values were

performed as already described (Torne et al., 2021).

Solutions: <u>Bath solution</u> for protoplasts' pH$_{CYT}$ determination (in mM): 5 KCl, 1 $CaCl_2$, 4

$MgCl_2$, 10 MES; pH 5.6; osmolality: 435 mOsm, adjusted w/ D-sorbitol.

<u>SNARF1-AM</u>: (Molecular Probes/Thermo Fisher cat. #: C-1272); 50 µg portions were

dissolved in DMSO sequentially, as needed; a 10 mM stock solution was stored in aliquots

at -20°C.

<u>Pluronic F-127:</u> (Molecular Probes, Life Technologies, cat. # P6867, Oregon, 741 USA), a

20% stock solution in DMSO was stored in 100 µL aliquots at RT.

Leaf vein density measurements – vein density measurements were performed as

described earlier (Grunwald et al., 2021).

Osmotic water permeability coefficient (P_f) determination under different light

regimes

Protoplast isolation and P_f determination were performed on *SCR:GFP* (Col) as already

described   (Shatil-Cohen et al., 2014) with the following modifications: The P_f

determination by the hypo-osmotic challenge (150 mOsm less than in the isotonic solution)

was carried out under either light (6 V-30 W halogen lamp, PHILIPS 5761) or dark

treatment.

Light treatments: After isolation, the protoplasts were divided into two test tubes, one tube

("light treated") was kept in constant light in the growth room and the other tube ("dark

treated") was kept in a light-proof box until the measurements. The light source during the

hypo-osmotic challenge of the light-treated protoplasts was the built-in microscope bulb.

During the osmotic challenge of the dark-treated protoplasts, the room was darkened,

protoplasts were added to the experimental chamber and let sink and stick to the bottom,

and a GFP-positive BSC was quickly located. Next, the microscope light was turned off,

and the cell was kept in the chamber in total darkness for 10 minutes. During the 1 min

videotaping of the protoplast's hypoosmotic challenge, the chamber was illuminated by the

microscope lamp via a red plastic band-pass filter (~600-700nm/ 30 µmol m^{-2} sec^{-1}) to

block out the BL. Henceforth, in P_f experiments, we designate this low-level red light

illumination as "Dark". Both wash solutions (isotonic and hypotonic) were buffered to pH

5.7.

Determination of guttation drops pH

Whole intact WT (Col) plants were covered overnight with black plastic bags to increase

the relative humidity in their atmosphere and to prevent transpiration. The following

morning immediately after the growth room lights went ON), guttation droplets were

collected from the tips of the leaves. Ten to twenty guttation droplets were collected and

pooled to reach a sample volume of approximately 20 µL in a 200-µL vial, which was

immediately sealed. pH was measured within 10 min of sample collection using a micro-

pH electrode MI-410 (Microlectrode, Inc., New Hampshire, USA).

RESULTS

PHOT1 and PHOT2 light receptors are involved in K_{leaf} regulation

Following the GC model in which the photoreceptors PHOT1 and PHOT2 transduce the BL stomatal opening signal, and encouraged by the substantial expression of the blue light receptor genes *PHOT1* and *PHOT2* in the Arabidopsis BSCs transcriptome (Wigoda et al., 2017), we explored the roles of these two light receptors in the regulation of the leaf hydraulic conductance, K_{leaf}. We compared, under two light regimes, red light (RL) and red light + blue light (RL+ BL), the K_{leaf} of WT plants to K_{leaf} of knockout mutants lacking one or both of these light receptors (*phot1-5, phot 2-1 and phot1-5 phot 2-1*; Fig. 1). Under RL+BL, WT leaves' K_{leaf} was more than 3-fold higher than under RL treatment and more than 2-fold higher than K_{leaf} of the RL+BL-treated three mutants. Also, under RL+BL, *phot2-1* mutant's K_{leaf} was about 40 % higher than under the RL treatment. In contrast, BL did not seem to increase the *phot1-5* mutant's or the double mutant's K_{leaf} compared to K_{leaf} under the RL treatment. Under RL, K_{leaf} did not differ among all the mutants, but while the WT's K_{leaf} was no different than the single-mutants' K_{leaf}, it was about double that in the double mutant (Fig. 1).

The higher calculated value of WT's K_{leaf} under BL (as compared to RL), resulted from a greatly increased (less negative) leaf water potential, Ψ_{leaf}, and an appreciably higher transpiration rate, E (Eq. 1, Materials and methods). This invites the interpretation that the BL-induced a highly conductive water pathway *into* the leaf enabling higher radial water influx which, in turn, was able to compensate for the BL-enhanced E, i.e. water efflux from the leaf, via the BL-increased stomata conductance, g_S (suppl. Fig. S1).

BSCs-specific silencing of PHOT1 and PHOT2 receptors decreased K_{leaf}

To examine specifically the participation of the BSCs' light receptors in the BL-induced

K_{leaf} increase, we silenced either the *PHOT1* or the *PHOT2* gene using amiRNA (artificial

microRNA) under the BSCs-specific promoter, *SCARECROW* (*SCR*, see Materials and

Methods). Under RL+BL illumination, the K_{leaf} values of leaves of both types of transgenic

plants, *SCR:mirphot1* and *SCR:mirphot2*, were lower (by 40 -50 %) than the K_{leaf} values of

WT leaves (Fig. 2). In contrast, all the *SCR:mirphot* plants had WT'-like g_S and E, while

their Ψ_{leaf} was lower (by 50-90 %) than WT's, suggesting a reduction of the radial water

influx to the leaf (Suppl Fig. S2).

BSCs-specific complementation of the *phot1-5* mutant with PHOT1 restored K_{leaf}

We complemented the *phot1-5* (Col-gl) mutant plants with *SCR*:*PHOT1* to restore PHOT1

activity exclusively in the BSCs (Materials and Methods) and compared them to WT (Col-

gl) plants and to *phot1-5* plants. *SCR:PHOT1* plants illuminated with RL+BL had their

K_{leaf} restored from *phot1-5* values to WT values (Fig. 3). While, as expected, BL increased

the WT's g_S, the mutant's g_S and that of the complemented mutant did not change under

BL, also as expected from the main-BL-receptor-devoid guard cells (Suppl. Fig. S3).

Notably, under BL, the Ψ_{leaf} of the *SCR*:*PHOT1* complemented mutant was restored to the

WT's values, and both were higher by over 50% than the mutant's Ψ_{leaf} (Suppl. Fig. S3).

This was the major contribution to the restored high K_{leaf} of the complemented mutant,

suggesting a restoration of radial water influx to the leaf.

The kinase inhibitor, tyrphostin 9, inhibited the RL+BL-induced K_{leaf} increase

The kinase inhibitor tyrphostin 9 suppressed the blue-light-dependent H^+-ATPase

phosphorylation in guard cells (Hayashi et al., 2017). To examine further whether the

vascular PHOT receptors initiate a similarly sensitive phosphorylating event in the BL-

dependent K_{leaf} regulation pathway, we perfused detached WT leaves with AXS, without

or with 10 μM tyrphostin 9, and exposed them to RL or RL+BL treatments (Materials and

Methods). The K_{leaf} values of RL+BL-illuminated leaves perfused with tyrphostin 9 were

about 50% lower than in RL+BL-illuminated control leaves (without tyrphostin 9), and

were no different than K_{leaf} in leaves illuminated with RL only (whether or not exposed to

tyrphostin 9; Fig.4a, left panel, suppl. Fig S5). However, gs was not affected by the petiole-

delivered tyrphostin 9 (Fig.4b, left panel). In contrast to the perfused tyrphostin 9, spraying

the inhibitor on the leaf surface did not interfere with the BL-induced almost 3-fold

increase of K_{leaf} (Fig. 4a, right panel), but it averted the BL-induced gs increase (Fig. 4b,

right panel). The impaired stomatal response as a result of tyrphostin 9 application

directly on the leaf surface, is in agreement with tyrphostin 9 inhibition of BL stomatal

opening in epidermal peels (Hayashi et al., 2017). The leaf transpiration, E, behaved in a

pattern similar to gS (Suppl. Fig S4a). Petiole-fed tyrphostin 9 abolished the BL-induced

increase of Ψ_{leaf}, but tyrphostin 9 sprayed on the leaf surface did not prevent it (Suppl. Fig

S4b).

BL hyperpolarizes the BSCs and alkalinizes their cytosolic pH

To investigate further whether the BSCs plasma-membrane H^+-ATPases plays a role in the

BL signal transduction pathway, resembling GCs, we monitored the membrane potential of

BSC protoplasts using the potentiometric dual-excitation dye di-8-ANEPPS. Indeed, 5-10

min of RL+BL illumination hyperpolarized the BSCs protoplasts relative to RL

illumination or to Dark (absence of illumination), while 10 μM tyrphostin 9 added in the

bath (Materials and Methods) not only abolished the BL-induced hyperpolarization but

even depolarized the BSCs (Fig. 5, Suppl. Fig. S8), linking BL to the activity of plasma

membrane proton pumps.

Exploring further our hypothesis of H^+-ATPases activation by BL, we monitored the

cytosolic pH (pH_{CYT}) using the ratiometric, dual-emission fluorescent pH-sensitive dye,

SNARF1 (Materials and methods). pH_{CYT} of BSCs illuminated for 10 min with RL+BL

(following dark treatment), was significantly more alkaline than pH_{CYT} of dark-treated

BSCs (0.72 pH units higher) and pH_{CYT} of RL-only-treated BSCs (1.06 pH units higher),

indicating that BL initiates the activity of H^+-ATPases which extrude protons from the

BSCs into the xylem (Fig. 5b).

AHA2 plays a role in the BL-induced K_{leaf} regulation

Recently, we reported that AHA2 plays a major role in K_{leaf} regulation by acidifying the

xylem sap. To test further whether the BSC's H^+-ATPase AHA2 participates in the blue

light-initiated K_{leaf} regulation pathway, we examined the AHA2 knockout mutant *aha2-4*,

and its BSCs-complemented transgenic plant line *SCR*:*AHA2* [i.e., the *aha2-4* mutant with

AHA2 expressed only in its BSCs (Grunwald et al., 2021)] under the two light regimes (RL

and RL+BL). K_{leaf} of *aha2-4* leaves did not respond to the RL+BL illumination and was

only about 40 % of K_{leaf} of the BL-treated WT leaves, but, a similar illumination increased

K_{leaf} of the AHA2-complemented (*SCR*:*AHA2*) plants to almost 65 % of the WT's K_{leaf} (Fig.

6), indicating that the BL-induced K_{leaf} increase depended on AHA2. In contrast, under

RL+BL, E and g_S of *SCR*:*AHA2* did not differ from E and g_S of the *aha2-4* mutant, and they

remained unaffected by BL and lower than the WT's E and g_S (Suppl. Fig S5a & b).

Notably, Ψ_{leaf} of *SCR*:*AHA2* was higher by about 40 % than the mutant's and did not differ

from the WT's Ψ_{leaf}, suggesting a restoration of radial water influx to the leaf (Suppl. Fig

S5c).

Morning light after night darkness acidifies the xylem sap

To delve further into the acidification-mediated link between light and Kleaf, we explored the diurnal transition between the xylem sap pH in the night and in the morning,, we first measured the pH of guttation drops collected from the tips of WT leaves before "dawn" (i.e., before lights-ON, Materials and Methods) and found that their mean pH was 7.3. Next, we measured the xylem sap pH in detached leaves, comparing leaves before dawn to leaves after dawn (i.e., leaves at the end of an overnight dark treatment, to leaves after a 30 min WL illumination in the growth chamber; Materials and Methods). This comparison included leaves of WT, *phot1-5* and *SCR:PHOT1* plants. Illumination acidified the xylem sap in WT plants by approx. 0.6 pH units (from 6.0 to 5.4) and in the *SCR: PHOT1* plants by approx. 0.8 pH units (from 6.1 to 5.3), while the xylem pH in *phot1-5* remained unchanged, at about 5.9 (Fig. 7), suggesting that PHOT1 in the BSCs is necessary for light-activated xylem acidification, which, in turn, we have already shown to increase K_{leaf} (Grunwald et al., 2021).

Light increases the osmotic water permeability, P_f, of the BSCs

To explore the link between light and K_{leaf} at the cellular level, we tested whether light affects the osmotic water permeability (P_f) of the BSCs membrane. We determined the P_f of isolated BSC protoplasts from videotapes of their swelling during a hypotonic challenge after white light (WL) illumination or after darkening for 10 min (Materials and Methods). The illuminated BSCs had higher P_f than the dark-treated ones (Fig. 8), suggesting that the morning increase of water fluxes from the xylem to the BSCs (and then, presumably – further into the mesophyll) is mediated also by aquaporins, independently from xylem sap acidification.

403 **Leaf vein architecture is not affected by the expression of PHOT1 or PHOT2**

404 In order to resolve whether the diminished K_{leaf} of the Phot receptor mutants we observed

405 under BL were due to differences in leaf architecture and morphology, we quantified the

406 vein density in all genotypes we studied and found no differences among them (Suppl. Fig

407 S9).

DISCUSSION

BL signal transduction in the BSCs compared to the GC's BL signal transduction

We demonstrate here an autonomous BL signaling mechanism in Arabidopsis BSCs which underlies the increase of K_{leaf} by light and which resembles the "classical" short-term BL-induced signaling pathway mediated by PHOT1 and PHOT2 photoreceptors found in stomatal opening, as well as chloroplasts movement in response to BL (Zeiger et al., 1983; Sakai et al., 2001; Kinoshita et al., 2001b; and reviewed by Briggs and Christie, 2002). To explain the underlying mechanism, we built on our earlier findings that, on the one hand, the BSCs express substantial levels of *PHOT1* and *PHOT2* receptors (Wigoda et al., 2017) and, on the other hand, K_{leaf} is enhanced by xylem sap acidification by the BSCs' AHA2 (Grunwald et al., 2021). Here, we link these findings using the PHOT receptors mutants *phot1-5* and *phot2-1*, and the double mutant *phot1-5 phot2-1*, and, moreover, we focus on the roles of both receptors PHOT1 and PHOT2 specifically in the BSCs and demonstrate the independence of the BSCs' BL signaling from the GCs' BL signaling. This notion of independence is particularly important since the BSC layer constitutes a hydraulic valve between the xylem and the leaf, regulating water loss to the atmosphere *in series* with the GCs apertures.

The similarity of BSCs BL-signaling to the stomata-opening BL-signaling, is highlighted by the requirement for the presence of both active PHOT1 and PHOT2 receptors for the BL-induced respective increase of K_{leaf} and g_S; we base this conclusion on the particularly severe depression of both responses in the double mutant relative to the two single mutants' responses (13 % of the WT's K_{leaf} BL response remaining in the double mutant, vs 32 and 40 % of the WT's K_{leaf} in *phot1-5* and in *phot2-1*, respectively (Fig.1), and 35% of the WT's g_S BL response remaining in the double mutant, vs 57 % and 71% of the WT's g_S in *phot1-*

5 and in *phot2-1*, respectively (Suppl. Fig.S1a; similar BL effects on stomatal apparture in

these mutants have been reported earlier (Kinoshita et al., 2001)).

Similar between the two systems (BSCs and GCs) is also the slightly higher (than in RL)

sensitivity to BL in the *phot2-1* mutant seen both in K_{leaf} (Fig.1) and in g_S (Suppl. Fig.S1a),

the latter also already noted earlier (Kinoshita et al., 2001), and both likely due to the

presence of PHOT1. Notably, based on the complete abolishment of the BL-K_{leaf} response

(relative to RL) in the "reciprocal" *phot1-5* mutant, presumed to possess PHOT2 activity,

PHOT2 would appear completely redundant, were it not for the even more severe

impairment of the double mutant's K_{leaf} (even below the WT's K_{leaf} under RL alone; Fig.1).

Interestingly, after 2-4 hours of illumination (beyond our usual 15 minutes of illumination)

the *phot* single mutants's g_S re-acquired the WT-like BL sensitivity (Suppl. Fig. S7), similar

to an earlier report on stomatal aperture after 2-4 hours of BL illumination (Kinoshita et al.,

2001); this could perhaps reflect a gradually accumulating effect of the GCs' AHA activity,

overcoming the partially impaired PHOT-to-AHA signal transduction in the single *phot*

mutants.

That the PHOTs (and particularly PHOT1) are indispensible to BL signaling in the BSCs,

as they are in GCs, is emphasized additionally by our two genetic manipulations focused

entirely on the BSCs: (a) the severe impairment of the BL-induced K_{leaf} increase in WT

plants with either PHOT1 or PHOT2 silenced by *SCR:PHOT-mirs* (Fig. 2) and (b) the full

restoration of K_{leaf} sensitivity to BL by *phot1-5* complementation with *SCR:PHOT1* (Fig.

3).

While the results of these genetic manipulations on the BSCs PHOTs attest, importantly,

to their involvement in the BL regulation of radial water influx to the leaf, they also

emphasize the independence of the BSCs BL signal transduction from that in the GCs – as

deduced from the lack of their effect on the g_S (Suppl. Figs. S2a, S3a).

Additionally, the localized effects of the PHOT inhibitor, tyrphostin 9 demonstrated both

the similarity between the BSCs' and the GC's BL signaling pathways in WT plants and

the independence of the first from the latter; both signaling pathways were inhibited by it,

but only when it was applied in close proximity to the tissue of action: BSCs were affected

only by the petiole-fed tyrphostin 9, which inhibited the BL-induced increase of K_{leaf} and

Ψ_{leaf} without affecting g_S, while GCs were affected only by tyrphostin 9 sprayed on the leaf

surface (i.e., on GCs), which abolished the BL-induced g_S increase without affecting K_{leaf}

(Fig.4, suppl. Fig. S4).

The cellular BL signal transduction pathway of the BSCs

The BSCs hyperpolarization in response to BL that we observed (Fig. 5) resembles the GCs

hyperpolarization in response to BL (Assmann et al., 1985; Roelfsema et al., 2001; Taylor

and Assmann, 2001) and its inhibition by tyrphostine (Fig. 5) suggests the involvement of

the PHOT receptors in a similar way (Hayashi et al., 2017). Hyperpolarization is known to

be generated by the plasma membrane H^+-ATPase activity (Spanswick, 1981). Recently we

reported that the BSCs H^+-ATPase plays a major role in the acidification of the xylem sap

which in turn increases K_{leaf} (Grunwald et al., 2021). Indeed – compared to WT – the H^+-

ATPase mutant, *aha2-4* had a diminished BL-induced K_{leaf} increase (possibly due to a more

alkaline xylem pH), as well as a diminished BL-induced g_S increase, while *aha2-4* with the

SCR:AHA2 complementation limited to BSCs had its BL-induced K_{leaf} increase restored,

while its g_S remained unaffected, affirming the mediation by *AHA2* of the BL increase of

K_{leaf} (Fig. 6, Suppl. Fig. S6). Notably, this is an interesting contrast with the otherwise

similar GCs' BL-signaling pathway, which terminates upon AHA1 (Yamauchi et al., 2016).

Here, we report that, in addition to external low pH, light is also needed for the BSCs P_f

increase (Fig.8). The role of BSCs aquaporins in controlling K_{leaf} was already identified

earlier (in walnut, Ben Baaziz et al., 2012; in grapevine, Vitali et al., 2016; in poplar, Muries

et al., 2019; in Arabidopsis, Sade et al., 2014; Prado et al., 2019). In particular, Prado et al.,

(2019) demonstrated a contribution of the abundant – also in the leaf veins – aquaporin

AtPIP2;1 to the rhythmicity of the diurnal and circadian hydraulic conductance of the

Arabidopsis rosette, K_{ros}, which peaked before mid-day. Thus, in addition to the likely P_f-

promoting effect of the slightly acidic pH (6) perfusion solution (YG,NW et al., 2020) that

Prado et al. (2019) used in their K_{ros} measurements, the changes in K_{ros} reflected also the

effects of light and the circadian clock (Prado et al., (2019).

Elaborating on the correlation between K_{ros} and AtPIP2;1 phosphorylation that they found,

these authors demonstrated that frog oocytes expressing AtPIP2;1 had higher P_f induced by

co-expressing two of the Arabidopsis 14-3-3 protein isoforms which bound to the AtPIP2;1

(Prado et al., 2019). It would be tempting to speculate, on this basis, that, resembling the

PHOT-initiated BL signaling cascade involving 14-3-3 proteins which bind to and

upregulate the GCs' AHA1 activity (e.g., Emi et al., 2001) also AtPIP2;1 activity, and

perhaps that of other vascular aquaporins may be enhanced by light via similar signaling

(i.e., by phosphorylation and 14-3-3 protein binding).

An alternative tempting speculation would be that the light-induced BSCs' P_f increase is

due to BL-mediated activation of AHA2 and the BSCs' cytosol alkalinization resulting

from proton secretion. Earlier work already documented that P_f increases due to aquaporin

activation by cytosol alkalinization, mediated by a conserved histidine (His 197 in

AtPIP2;2; Tournaire-Roux et al., 2003) and a leucine (Leu206 in BvPIP2;2 Fortuna et al.,

2019) in the cytosol-facing D loop of the aquaporin protein. Our results of cytosolic

alkalisation (Fig. 5b) in the BSCs, in response to ilumintion with BL, can explain the

increase in P_f under light (Fig. 8), underlying the light-induced increase in K_{leaf}. Our results

showing increased P_f due to cytosolic alkalisation, are in accord with (Tournaire-Roux et

al., 2003) affirming that that aquaporin gating is involved in this mechanism. Moreover,

could the concurrent hyperpolarization of BSCs by BL (also via AHA2 activation) imply a

possible role of hyperpolarization in the regulation of the BSCs AQPs as suggested by in-

silico simulations (Hub et al., 2010)? This possibility needs yet to be explored

experimentally.

The model of K_{leaf} activation by BL, in a GC-like BL signal transduction pathway, that our

results support, is as follows: the BSCs' PHOT1 and PHOT2 (only partially redundant)

perceive the blue light, activate the plasma membrane H^+-ATPase (AHA2), which

hyperpolarizes the BSCs, alkalinizes the BCSs' cytosol and acidifies the xylem pH, leading

to an increase of the BSC P_f and, in turn, to a rise of K_{leaf}, or, schematically:

BL →BSCs' *PHOT*s activation →tyrosin phosphorylation* →BSCs' H^+-ATPase

(AHA2) activation →BSCs hyperpolarization,cytosol alkalization **and xylem**

acidification → BSCs P_f elevation →K_{leaf} increase.

* Notably, the abovementioned tyrosine phosphorylation step – suggested to occur, in the

guard cells, on the BHP protein upstream of the H^+-ATPase (Hayashi et al., 2017) – is yet

to be explored in the BSCs; currently, our guess is that similarly to the GC, this step occurs

upstream of the BSCs' AHA2 (Fig. 9).

The physiological relevance of K_{leaf} increase by blue light

Both GCs opening and K_{leaf} increase occur in the early morning hours (e.g., Brodribb and

Holbrook, 2004; Domec et al., 2009; Locke and Ort, 2015) when the irradiance spectrum

is relatively enriched in blue wavelengths (Chiang et al., 2019, Matthews et al., 2020, and

references therein). We suggest that the advantage of this blue light response of stomata is

in allowing the acquisition of CO_2 during sub-maximum VPD (vapor pressure deficit, a

measure of the driving force for leaf water evaporation), thus increasing water use

efficiency (WUE).

What is the advantage of such early K_{leaf} response to light?

Interestingly, not only does K_{leaf} increase in response to blue light as does stomatal aperture

but, in accord with our conclusions about the independence of the BSCs' BL signal

transduction from that of the GCs', K_{leaf} responds to light faster than g_S (Guyot et al., 2012).

One possible advantage of this almost-simultaneity, could be to match stomata opening

with increased water flux into the leaf. Were it not for this accompanying K_{leaf} increase,

BL-induced stomata opening, which peaks in the morning hours (the "Golden hour"; Gosa

et al., 2018), could lead to an imbalance of leaf water, even in the presence of a relatively

low VPD, leading to a drop in the leaf water potential (Ψ_{leaf}); this in turn, could close

stomata (Raschke, 1970; Guyot et al., 2012; Klein, 2014; Scoffoni et al., 2018) and thus

limit photosynthesis. Instead, an early morning increase of K_{leaf} would elevate Ψ_{leaf} thereby

increasing and prolonging the opening stomata (ibid). Thus, the BL-response of K_{leaf} could

support and enhance the stomata BL response. Hypothetically, under more extreme

conditions, e.g., with high VPD already in the early morning, increased K_{leaf} would prevent

a hydraulic pathway failure when stomatal conductance is at its peak (Brodribb and

Holbrook, 2004; Halperin et al., 2017; Gosa et al., 2019).

 Another possible advantage of the early morning BL-induced K_{leaf} increase may be related

to CO_2 permeability via aquaporins. A few lines of evidence converge to support this

hypothesis: (a) CO_2 can cross cell membranes through aquaporins, and, in particular, those

of the PIP1 family (Kaldenhoff and Fischer, 2006), and (b) the transcripts of PIP1 family

aquaporins in walnut were upregulated concomitantly with K_{leaf} increase by a brief (15 min)

white light illumination (Biaaziz et al., 2012), (c) in soybean, the transcript of a PIP1 family

aquaporin (PIP1;8) was more abundant in the morning than in midday in correlation with a

higher K_{leaf} in the morning than in the midday (Locke and Ort, 2015), (d) we demonstrated

that aquaporins control K_{leaf} (Shatil-Cohen and Moshelion, 2012; Sade et al., 2014); (e)

xylem-transported CO_2 was suggested to be a source for CO_2 assimilated in the bundle

sheath and mesophyll (Janacek et al., 2009; Hubeau et al., 2019). Thus, assuming the K_{leaf}

increase with the first light of day parallels the increase of CO_2 permeability of aquaporins

in the BSCs, the passage to the mesophyll of CO_2 from the xylem originating in roots

respiration will be enhanced even before full stomatal opening. The increased CO_2

availability at this time, when the photosynthetic light is already sufficient, would enhance

CO_2 assimilation – a great advantage to the plant. To strengthen this hypothesis Stutz and

Hanson, (2019) have shown that xylem transported CO2 is assimilated in plants at different

fractions throughout the day depending on the lighting and CO2 availability. Interestingly,

photosynthetic CO_2 uptake reaches a maximum within 10 minutes of blue light (Doi et al.,

2015), which is over threefold faster than the full opening of stomata (Grondin et al., 2015).

In conclusion, our data provide new evidence for the role of the BSCs' phototropins

(PHOT1 and PHOT2) in their AHA2-mediated increase of K_{leaf} by blue light.

An additional interesting outcome from our results is the corroboration – and explanation –

of the reports on alkalinized xylem sap pH in the absence of light, e.g., at night (Urrestarazu

et al., 1996 (in tomatoes); Aubrey et al., 2011 (in poplars)), or even on cloudy days (in

poplars; Aubrey et al., 2011). Similarly, in Arabidopsis, only after dawn and the activation

of the BL signal transduction pathway does the xylem sap become acidic (as in Fig.7) and

K_{leaf} increases (Fig. 3).

Our results, together with the fact that in the dark the BSC P_f is low (Fig. 8), may explain

how the mesophyll does not get flooded at night when root pressure increases. While during

day time the BSCs' higher P_f underlies the higher K_{leaf} which is important for the support

of transpiration (Sade et al., 2015; Shatil-Cohen and Moshelion, 2012; Sade and Moshelion,

2014; Grunwald et al., 2021), at night, the P_f and K_{leaf} reduction likely prevents water influx

into the leaf, and the buildup of root pressure is relieved via guttation, and if not — xylem

sap may flood the leaf (Shatil-Cohen and Moshelion, 2012). Not surprisingly, and in accord

with the dark-alkalinization of the xylem sap, guttation drops pH was 7.2 ± 0.04 ($\pm$SE, 15

biological repeats/ collected over three days).

The inactivity of the BSCs H^+-ATPases, manifested in the nightly alkalinization of the

xylem sap, could be advantageous for the plant as a means of energy conservation, as

ATPases operate at a high energetic cost (Palmgren, 2001).

These findings expand our understanding of the molecular basis of the leaf water influx

regulation. That BL increases K_{leaf} in several other species (Voicu et al., 2008; Voicu et al.,

2009; Sellin et al., 2011; Aasamaa and Sber, 2012; Ben Baaziz et al., 2012), imparts an

even more general importance to our results. Thus, a focus on the hydraulic valve in series

with the stomata should provide new directions for studying the plant water relations within

its environment.

AUTHOR CONTRIBUTIONS

598 YG and SG constructed all transgenic lines, planned and preformed the K_{leaf} and vein pH

599 experiments, and wrote the paper. YG planned and preformed Pf experiments. TT planned

600 and preformed membrane potential and cytosolic pH experiments. NM and MM

601 conceptualized, supervised and wrote the paper.

ACKNOWLEDGMENTS

603 We thank Dr. Dizza Bursztyn of the Hebrew University of Jerusalem for assisting us with

604 the statistical analysis of this work

605 This research was supported by the Israel Science Foundations, ISF (grants No. 1043/20),

606 to MM

REFERENCES

Aasamaa, K. and Sõber, A. (2012). Light sensitivity of shoot hydraulic conductance in five temperate deciduous tree species. Funct. Plant Biol. 39: 661.

Abràmoff, M.D., Magalhães, P.J., and Ram, S.J. (2004). Image processing with imageJ. Biophotonics Int. 11: 36–41.

Alvarez, J.P., Pekker, I., Goldshmidt, A., Blum, E., Amsellem, Z., and Eshed, Y. (2006). Endogenous and Synthetic MicroRNAs Stimulate Simultaneous, Efficient, and Localized Regulation of Multiple Targets in Diverse Species. Plant Cell 18: 1134–1151.

Assmann, S. (1993). Signal Transduction in Guard Cells. Annu. Rev. Cell Dev. Biol. 9: 345–375.

Assmann, S.M., Simoncini, L., and Schroeder, J.I. (1985). Blue light activates electrogenic ion pumping in guard cell protoplasts of Vicia faba. Nature 318: 285–287.

Aubrey, D.P., Boyles, J.G., Krysinsky, L.S., and Teskey, R.O. (2011). Spatial and temporal patterns of xylem sap pH derived from stems and twigs of Populus deltoides L. Environ. Exp. Bot. 71: 376–381.

Ben Baaziz, K., Lopez, D., Rabot, A., Combes, D., Gousset, A., Bouzid, S., Cochard, H., Sakr, S., and Venisse, J.-S. (2012). Light-mediated Kleaf induction and contribution of both the PIP1s and PIP2s aquaporins in five tree species: walnut (Juglans regia) case study. Tree Physiol. 32: 423–434.

Briggs, W.R. and Christie, J.M. (2002). Phototropins 1 and 2: Versatile plant blue-light receptors. Trends Plant Sci. 7: 204–210.

Briggs, W.R. and Huala, E. (1999). Blue-Light Photoreceptors in Higher Plants. Annu. Rev. Cell Dev. Biol. 15: 33–62.

Brodribb, T.J. and Holbrook, N.M. (2004). Diurnal depression of leaf hydraulic

conductance in a tropical tree species. Plant, Cell Environ. 27: 820–827.

Chiang, C., Olsen, J.E., Basler, D., Bånkestad, D., and Hoch, G. (2019). Latitude and

Weather Influences on Sun Light Quality and the Relationship to Tree Growth.

Forests 10: 610.

Clough, S.J. and Bent, A.F. (1998). Floral dip: A simplified method for Agrobacterium-

mediated transformation of Arabidopsis thaliana. Plant J. 16: 735–743.

Doi, M., Kitagawa, Y., and Shimazaki, K.-I. (2015). Stomatal Blue Light Response Is

Present in Early Vascular Plants. Plant Physiol. 169: 1205–1213.

Domec, J.C., Noormets, A., King, J.S., Sun, G., McNulty, S.G., Gavazzi, M.J., Boggs,

J.L., and Treasure, E.A. (2009). Decoupling the influence of leaf and root hydraulic

conductances on stomatal conductance and its sensitivity to vapour pressure deficit

as soil dries in a drained loblolly pine plantation. Plant, Cell Environ. 32: 980–991.

Elmore, J.M. and Coaker, G. (2011). The Role of the Plasma Membrane H+-ATPase in

Plant–Microbe Interactions. Mol. Plant 4: 416–427.

Emi, T., Kinoshita, T., and Shimazaki, K. (2001). Specific binding of vf14-3-3a isoform

to the plasma membrane H+-ATPase in response to blue light and fusicoccin in

guard cells of broad bean. Plant Physiol. 125: 1115–1125.

Fortuna, C.A., Zerbetto De Palma, G., Aliperti Car, L., Armentia, L., Vitali, V., Zeida, A.,

Estrin, D.A., and Alleva, K. (2019). Gating in plant plasma membrane aquaporins:

the involvement of leucine in the formation of a pore constriction in the closed state.

FEBS J. 286: 3473–3487.

Gosa, S.C., Lupo, Y., and Moshelion, M. (2019). Quantitative and comparative analysis

of whole-plant performance for functional physiological traits phenotyping: New

tools to support pre-breeding and plant stress physiology studies. Plant Sci. 282: 49–

59.

Grondin, A., Rodrigues, O., Verdoucq, L., Merlot, S., Leonhardt, N., and Maurel, C.

(2015). Aquaporins Contribute to ABA-Triggered Stomatal Closure through OST1-

Mediated Phosphorylation. Plant Cell 27: 1945–1954.

Grunwald, Y., Wigoda, N., Sade, N., Yaaran, A., Torne, T., Chaka Gosa, S., Moran, N.,

and Moshelion, M. (2021). Arabidopsis leaf hydraulic conductance is regulated by

xylem-sap pH, controlled, in turn, by a P-type H + -ATPase of vascular bundle

sheath cells . Plant J.

Grunwald, Y., Wigoda, N., Sade, N., Yaaran, A., Torne, T., Gosa, S.C., Moran, N., and

Moshelion, M. (2020). Arabidopsis leaf hydraulic conductance is regulated by

xylem-sap pH, controlled, in turn, by a P-type H+-ATPase of vascular bundle sheath

cells. bioRxiv: 234286.

Guyot, G., Scoffoni, C., and Sack, L. (2012). Combined impacts of irradiance and

dehydration on leaf hydraulic conductance: insights into vulnerability and stomatal

control. Plant. Cell Environ. 35: 857–871.

Halperin, O., Gebremedhin, A., Wallach, R., and Moshelion, M. (2017). High-throughput

physiological phenotyping and screening system for the characterization of plant-

environment interactions. Plant J. 89: 839–850.

Hayashi, M., Inoue, S., Ueno, Y., and Kinoshita, T. (2017). A Raf-like protein kinase

BHP mediates blue light-dependent stomatal opening. Sci. Rep. 7: 45586.

Hub, J.S., Aponte-Santamaría, C., Grubmüller, H., and De Groot, B.L. (2010). Voltage-

regulated water flux through aquaporin channels in silico. Biophys. J. 99: L97–L99.

Hubeau, M., Thorpe, M.R., Mincke, J., Bloemen, J., Bauweraerts, I., Minchin, P.E.H., De

Schepper, V., De Vos, F., Vanhove, C., Vandenberghe, S., and Steppe, K. (2019).

High-Resolution in vivo Imaging of Xylem-Transported CO2 in Leaves Based on

Real-Time 11C-Tracing. Front. For. Glob. Chang. 2: 25.

Van Ieperen, W., Savvides, A., and Fanourakis, D. (2012). Red and blue light effects

during growth on hydraulic and stomatal conductance in leaves of young cucumber

plants. Acta Hortic. 956: 223–230.

Janacek, S.H., Trenkamp, S., Palmer, B., Brown, N.J., Parsley, K., Stanley, S., Astley,

H.M., Rolfe, S.A., Paul Quick, W., Fernie, A.R., and Hibberd, J.M. (2009).

Photosynthesis in cells around veins of the C 3 plant Arabidopsis thaliana is

important for both the shikimate pathway and leaf senescence as well as contributing

to plant fitness. Plant J. 59: 329–343.

Kaldenhoff, R. and Fischer, M. (2006). Aquaporins in plants. Acta Physiol. 187: 169–176.

Karlsson, P.E. (1986). Blue light regulation of stomata in wheat seedlings. II. Action

spectrum and search for action dichroism. Physiol. Plant. 66: 207–210.

Kinoshita, T., Doi, M., Suetsugu, N., Kagawa, T., Wada, M., and Shimazaki, K. (2001).

Phot1 and Phot2 mediate blue light regulation of stomatal opening. Nature 414: 656–

660.

Kinoshita, T. and Shimazaki, K. (1999). Blue light activates the plasma membrane H+-

ATPase by phosphorylation of the C-terminus in stomatal guard cells. EMBO J. 18:

5548–5558.

Klein, T. (2014). The variability of stomatal sensitivity to leaf water potential across tree

species indicates a continuum between isohydric and anisohydric behaviours. Funct.

Ecol. 28: 1313–1320.

Kronenberg, G.H.M. and Kendrick, R.E. (1986). The physiology of action. In

Photomorphogenesis in plants (Springer Netherlands: Dordrecht), pp. 99–114.

Locke, A.M. and Ort, D.R. (2015). Diurnal depression in leaf hydraulic conductance at

ambient and elevated [CO2] reveals anisohydric water management in field-grown

soybean and possible involvement of aquaporins. Environ. Exp. Bot. 116: 39–46.

Matthews, J.S., Vialet-Chabrand, S., Lawson, T., and Evans, J. (2020). Role of blue and

red light in stomatal dynamic behaviour. J. Exp. Bot. 71: 2253–2269.

Muries, B. et al. (2019). Aquaporins and water control in drought-stressed poplar leaves:

A glimpse into the extraxylem vascular territories. Environ. Exp. Bot. 162: 25–37.

Negin, B. and Moshelion, M. (2017). The advantages of functional phenotyping in pre-

field screening for drought-tolerant crops. Funct. Plant Biol. 44: 107.

Oishi, A.C., Oren, R., Novick, K.A., Palmroth, S., and Katul, G.G. (2010). Interannual

Invariability of Forest Evapotranspiration and Its Consequence to Water Flow

Downstream. Ecosystems 13: 421–436.

Den Os, D., Staal, M., and Elzenga, J.T.M. (2007). Signal integration by ABA in the blue

light-induced acidification of leaf pavement cells in pea (Pisum sativum L. var.

Argenteum). Plant Signal. Behav. 2: 146–152.

Palmgren, M.G. (2001). Plant plasma membrane H+ATPases: Powerhouses for nutrient

uptake.

Prado, K. and Maurel, C. (2013). Regulation of leaf hydraulics: from molecular to whole

plant levels. Front. Plant Sci. 4: 1–14.

Raschke, K. (1970). Stomatal Responses to Pressure Changes and Interruptions in the

Water Supply of Detached Leaves of Zea mays L. Plant Physiol. 45: 415–423.

Roelfsema, M.R.G. and Hedrich, R. (2005). In the light of stomatal opening: new insights

into 'the Watergate.' New Phytol. 167: 665–691.

Roelfsema, M.R.G., Steinmeyer, R., Staal, M., and Hedrich, R. (2001). Single guard cell

recordings in intact plants: Light-induced hyperpolarization of the plasma membrane.

Plant J. 26: 1–13.

Sade, N. and Moshelion, M. (2014). The dynamic isohydric-anisohydric behavior of

plants upon fruit development: taking a risk for the next generation. Tree Physiol. 34:

1199–202.

Sade, N., Shatil-Cohen, A., Attia, Z., Maurel, C., Boursiac, Y., Kelly, G., Granot, D.,

Yaaran, A., Lerner, S., and Moshelion, M. (2014). The Role of Plasma Membrane

Aquaporins in Regulating the Bundle Sheath-Mesophyll Continuum and Leaf

Hydraulics. PLANT Physiol. 166: 1609–1620.

Sade, N., Shatil-Cohen, A., and Moshelion, M. (2015). Bundle-sheath aquaporins play a

role in controlling Arabidopsis leaf hydraulic conductivity. Plant Signal. Behav. 10:

e1017177.

Sakai, A.K. et al. (2001). The Population Biology of Invasive Species. Annu. Rev. Ecol.

Syst.

Schindelin, J. et al. (2012). Fiji: an open-source platform for biological-image analysis.

Nat. Methods 9: 676–682.

Scoffoni, C., Albuquerque, C., Cochard, H., Buckley, T.N., Fletcher, L.R., Caringella,

M.A., Bartlett, M., Brodersen, C.R., Jansen, S., and Mcelrone, A.J. (2018). The

Causes of Leaf Hydraulic Vulnerability and Its Influence on Gas Exchange in

Arabidopsis thaliana. 178: 1584–1601.

Sellin, A., Sack, L., Õunapuu, E., and Karusion, A. (2011). Impact of light quality on leaf

and shoot hydraulic properties: A case study in silver birch (Betula pendula). Plant,

Cell Environ. 34: 1079–1087.

Shapira, O., Khadka, S., Israeli, Y., Shani, U., and Schwartz, A. (2009). Functional

anatomy controls ion distribution in banana leaves: Significance of Na + seclusion at

the leaf margins. Plant, Cell Environ. 32: 476–485.

Shatil-Cohen, A. and Moshelion, M. (2012). Smart pipes. Plant Signal. Behav. 7: 1088–

1091.

Shatil-Cohen, A., Sibony, H., Draye, X., Chaumont, F., Moran, N., and Moshelion, M.

(2014). Measuring the osmotic water permeability coefficient (Pf) of spherical cells:

isolated plant protoplasts as an example. J. Vis. Exp.: e51652.

Shimazaki, K., Doi, M., Assmann, S.M., and Kinoshita, T. (2007). Light Regulation of

Stomatal Movement. Annu. Rev. Plant Biol. 58: 219–247.

Spanswick, R. (1981). Eelectrogenic ion pumps. Annu. Rev. Plant Physiol. 32: 67–89.

Stutz, S.S. and Hanson, D.T. (2019). Contribution and consequences of xylem-transported

CO_2 assimilation for C_3 plants. New Phytol. 223: 1230–1240.

Svennelid, F., Olsson, A., Piotrowski, M., Rosenquist, M., Ottman, C., Larsson, C.,

Oecking, C., and Sommarin, M. (1999). Phosphorylation of Thr-948 at the C

Terminus of the Plasma Membrane H-ATPase Creates a Binding Site for the

Regulatory 14-3-3 Protein.

Takemiya, A., Sugiyama, N., Fujimoto, H., Tsutsumi, T., Yamauchi, S., Hiyama, A.,

Tada, Y., Christie, J.M., and Shimazaki, K.I. (2013). Phosphorylation of BLUS1

kinase by phototropins is a primary step in stomatal opening. Nat. Commun. 4.

Talbott, L.D., Shmayevich, I.J., Chung, Y., Hammad, J.W., and Zeiger, E. (2003). Blue

Light and Phytochrome-Mediated Stomatal Opening in the npq1 and phot1 phot2

Mutants of Arabidopsis. Plant Physiol. 133: 1522–1529.

Taylor, A.R. and Assmann, S.M. (2001). Apparent absence of a redox requirement for

blue light activation of pump current in broad bean guard cells. Plant Physiol. 125:

329–338.

Torne, T., Grunwald, Y., Dalal, A., Yaaran, A., and Moran, N. (2021). Stress is basic: 1

ABA alkalinizes both the xylem sap and the cytosol by inhibiting their P-type H^+-

ATPase 4 and stimulating their V-type H^+-ATPase AUTHOR CONTRIBUTIONS.

bioRxiv.

Tournaire-Roux, C., Sutka, M., Javot, H., Gout, E., Gerbeau, P., Luu, D.T., Bligny, R.,

and Maurel, C. (2003). Cytosolic pH regulates root water transport during anoxic

stress through gating of aquaporins. Nature 425: 393–397.

Ueno, K., Kinoshita, T., Inoue, S., Emi, T., and Shimazaki, K. (2005). Biochemical

Characterization of Plasma Membrane H+-ATPase Activation in Guard Cell

Protoplasts of Arabidopsis thaliana in Response to Blue Light. Plant Cell Physiol.

46: 955–963.

Urrestarazu, M., Sanchez, A., Lorente, F.A., and Guzmán, M. (1996). A daily rhythmic

model for pH and volume from xylem sap of tomato plants. Commun. Soil Sci. Plant

Anal. 27: 1859–1874.

Vitali, M., Cochard, H., Gambino, G., Ponomarenko, A., Perrone, I., and Lovisolo, C.

(2016). VvPIP2;4N aquaporin involvement in controlling leaf hydraulic capacitance

and resistance in grapevine. Physiol. Plant. 158: 284–296.

Voicu, M.C., Cooke, J.E.K., and Zwiazek, J.J. (2009). Aquaporin gene expression and

apoplastic water flow in bur oak (Quercus macrocarpa) leaves in relation to the light

response of leaf hydraulic conductance. J. Exp. Bot. 60: 4063–4075.

Voicu, M.C., Zwiazek, J.J., and Tyree, M.T. (2008). Light response of hydraulic

conductance in bur oak (Quercus macrocarpa) leaves. Tree Physiol. 28: 1007–1015.

Wigoda, N., Moshelion, M., and Moran, N. (2014). Is the leaf bundle sheath a "smart flux

valve" for K+ nutrition? J. Plant Physiol. 171: 715–722.

Wigoda, N., Pasmanik-Chor, M., Yang, T., Yu, L., Moshelion, M., and Moran, N. (2017).

Differential gene expression and transport functionality in the bundle sheath versus

mesophyll – a potential role in leaf mineral homeostasis. J. Exp. Bot. 68: 3179–3190.

Yamauchi, S., Takemiya, A., Sakamoto, T., Kurata, T., Tsutsumi, T., Kinoshita, T., and

Shimazaki, K. (2016). Plasma membrane H+-ATPase1 (AHA1) plays a major role in

Arabidopsis thaliana for stomatal opening in response to blue light. Plant Physiol.:

pp.01581.2016.

Zeiger, E., Assmann, S.M., and Meioner, H. (1983). The Photobiology of Paphiopedilum

stomata: opening under blue lightbut not red light. Photochem. Photobiol. 38: 627–

630.

Zeiger, E. and Helper, P.K. (1977). Light and Stomatal Function: Blue Light Stimulates

Swelling of Guard Cell Protoplasts. Science (80-.). 196: 887–889.

FIGURE LEGENDS

Figure 1: The effect of blue light on K_{leaf} of PHOT receptors mutants. K_{leaf} of fully

expanded excised leaves of WT (Col-gl) and *PHOT* mutants (*phot1-5* (Col-gl), *phot2-1*

(Col-gl) and a double mutant, *phot1-5 phot2-1* (Col-gl)) determined immediately after 15

min illumination with RL (220 µmol m^{-2} s^{-1}), or R+BL (220 µmol m^{-2} s^{-1}, consisting of

90 % RL and 10 % BL; Materials and methods). The columns are mean values (±SE) of the

indicated number of biological repeats from three independent experiments. Different

letters denote significantly different values (ANOVA, *Post hoc* Tukey's test, P< 0.05) . Note

the reduced response to RL+BL in the mutant lines.

Figure 2. amiR-silencing PHOT1 or PHOT2 (SCR:mirphot plants) in the BSCs

reduces K_{leaf} under RL+BL, generated in WT (Col; Materials and methods). K_{leaf} was

determined in fully expanded leaves of WT and *SCR:mirphot1* and *SCR:mirphot2* plants

treated with RL or RL+BL (as in Fig. 1; Materials and Methods) The columns are means

(± SE) of the indicated number of biological repeats from three independent experiments.

Different letters denote significantly different values (ANOVA, Post hoc Tukey's test, P<

0.05).

Figure 3. BSCs-directed *PHOT1* complementation of the *phot1-5* mutant restores the

normal K_{leaf}. Fully expanded leaves of WT (Col-gl), *phot1-5* (Col-gl) and *phot1-5* (Col-

gl) plants complemented with *SCR:PHOT1* underwent illumination treatments as in Fig.

1. The columns are means (± SE) of the indicated number of biological repeats from

three independent experiments. Different letters denote significantly different values

(ANOVA, *Post hoc* Tukey's test, P< 0.05).

Figure 4. Xylem-fed kinase Inhibitor tyrphostin 9 abolished the blue-light-induced

K_{leaf} increase and stomatal opening. Fully expanded leaves of WT (Col-gl) were pre-

incubated (petiole deep) in AXS without or with tyrphostin 9 (10 µm) or sprayed with AXS

without or with tyrphostin 9 (10 µm), and kept overnight in dark boxes. Immediately after

dark, leaves were illuminated for 15 min with RL (215 µmolm^{-2} s^{-1}) or RL+BL (215 µmol

m^{-2} s^{-1}, consisting of 90 % RL and 10 % BL). a. K_{leaf}. b. Stomatal conductance (g$_S$). The

columns are means (± SE) of the indicated number of biological repeats, determined in

three independent experimental days. Different letters denote significantly different values

(ANOVA, Post hoc Tukey's test, P< 0.05). Note that K_{leaf} was reduced when tyrphostin 9

was fed via the petiole but not when sprayed, and the opposite was true for the g_S.

Figure 5. - Figure 5. - Blue light (BL) hyperpolarizes the BSCs and alkalinizes their cytosol

relative to red light (RL) or Dark and tyrphostin 9 depolarizes them. a. Membrane potential

(E_M) of BSC protoplasts, without or with petiole-fed tyrphostin 9 (10 µM), determined

using the dual-excitation, ratiometric fluorescent dye di-8-ANEPPS (Materials and

Methods). After at least10-20 min in the dark, the cells were immediately imaged (Dark),

or illuminated for an additional 5-10 minutes either with RL (220 µmol m^{-2} s^{-1}) or with

RL+BL (220 µmol m^{-2} s^{-1}, including 10% BL; Materials and Methods), and then imaged.

The columns are mean values of fluorescence ratio (F $_{438}$/F$_{531}$) values (±SE; n as indicated)

derived from analyses of the images captured at the indicated excitation wavelengths

(Materials and Methods) in three independent experiments. Numbers above the columns

are the mean values of BSCs E_M, based on a calibration curve of F $_{438}$/F$_{531}$ vs. E_M obtained

in separate experiments using patch clamp (Materials and Methods, Suppl. Fig. S8).

b. Cytosol pH (pH_{CYT}) of BSCs protoplasts determined using the fluorescent pH probe

SNARF1 (Materials and methods). Protoplasts were dark-treated then further treated with

10 min of either dark, RL+BL or RL only. pH_{CYT} values were derived from analyses of the

images captured (Materials and Methods) and determined from the emitted fluorescence

ratio (F640 /F585) values, based on the in-situ calibration curve (detailed in Torne et al.,

2021). The columns are means (± SE) of the indicated number of biological repeats,

determined in three independent experimental days. Different letters denote significantly

different values (ANOVA, Post hoc Tukey's test, P< 0.05).

Figure 6. The BSCs H$^+$-ATPase, AHA2, mediates the blue-light-induced K_{leaf} increase.

Fully expanded excised leaves of WT (Col0) plants, *aha2-4* mutant plants (Col0) and *aha2-*

4 plants complemented in their BSCs with *SCR:AHA2* were illuminated for 15 min

immediately after dark, as in Fig.2. The columns are means (± SE) of the indicated number

of biological repeats from three independent experiments. Different letters denote

significantly different values (ANOVA, Post hoc Tukey's test, P< 0.05). Note the marked

restoration of K_{leaf} in the *aha2* mutant plants complemented solely in their BSCs.

Figure 7. White light (WL) illumination for 30 minutes acidifies the xylem pH of WT plants and *SCR:PHOT1*-complemented mutants, but not in the *phot1-5* mutant. The leaves of WT (Col-gl) plants, *phot1-5* (Col-gl) mutants and the *SCR:PHOT1*-complemented *phot1-5* mutant (as in Fig. 3) were illuminated by the growth room WL (150-200 μmol m^{-2} sec^{-1}, Materials and Methods). The columns are means (±SE) of the indicated number of biological repeats from three independent experiments Different letters denote significantly different values (*Post hoc* Tukey's test, P< 0.05).

Figure 8: White light increases the BSCs membrane osmotic water permeability coefficient (P$_f$). a. A time course of swelling of bundle sheath protoplasts from *SCR:GFP* plants upon exposure to a hypotonic XPS^{db} solution under the same illumination conditions (white light or darkness) as during the light treatment preceding the assay. The bath solution was buffered to pH 5.7. The arrow indicates onset of bath flush. **b**. Time course of the osmotic concentration change in the bath (C$_{out}$) during the hypotonic challenge c. Mean (±SE) initial P$_f$ values of the indicated number of bundle sheath protoplasts under two light regimes from three independent experiments. Columns are means (± SE) of 21-28 biological repetitions, determined in five independent experiments. The asterisk indicates a significant difference (Student's two-tailed unpaired t-test, P < 0.05).

Figure 9. Proposed BSCs-autonomous BL signal transduction pathway. Proposed BSCs-autonomous BL signal transduction pathway. a. An artist's rendering of a leaf radial water path, from xylem to mesophyll (K$_{leaf}$, blue hollow arrow) via the bundle sheath cells, BSCs, which tightly envelop the xylem. GC, a stomata guard cell; MC, mesophyll cell. **b**. Blue-light (BL) signalling pathway (blue arrowheads) in a bundle sheath cell, from BL perception by the phototropin receptors (PHOT), through an intermediate tyrphostin-sensitive tyrosine phosphorylation (TP), to the ultimate AHA2 (orange elipse) activation resulting in proton extrusion via the pump and xylem sap acidification, presumably, at the

expense of ATP breakdown to ADP with a transient phosphorylation (P) on the pump

protein, as expected from a P-type H^+-ATPase.

SUPPLEMENTAL FIGURES AND TABLE

Figure S1: **PHOT receptors mutants do not show the effect of blue light on the g_S, E**

and Ψ_{leaf}.

Figure S2. *SCR:mir*-**silencing of the PHOT receptors abolishes the effect of RL+BL on**

the g_S, E and Ψ_{leaf}.

Figure S3. BSCs-directed *PHOT1* **complementation of the** *phot1-5* **mutant elevates**

Ψ_{leaf} but not g_S or E.

Figure S4. **Xylem-fed kinase inhibitor tyrphostin 9 does not affect BL-increased E but**

decreases BL-increased Ψ_{leaf}, while tyrphostin 9 sprayed on leaves acts oppositely.

Figure S5. **Tyrphostin 9 perfused via the petiole does not affect the physiological**

parameters under RL-only illumination.

Figure S6. BSCs-directed *AHA2* **complementation of the** *aha2-4* **mutant restores its**

BL-induced Ψ_{leaf} increase but not the BL increases of g_S or E.

Figure S7. The 'long-illumination' g_S of intact leaves of whole plants does not differ

among the different genotypes (WT, *phot* **mutants and** *SCR:PHOT1*).**

Figure S8. *In-situ* **calibration of di-8-ANEPPs fluorescence ratio (F-ratio, F_{438}/F_{531})**

vs. the BSCs membrane potential (E_M).

Figure S9. **Leaf vein density does not depend on the expression of PHOT1 or PHOT2.**

Table S1: List of primers used in this study

Figure 1: **The effect of blue light on** K_{leaf} **of PHOT receptors mutants.** K_{leaf} of fully expanded excised leaves of WT (Col-gl) and *PHOT* mutants (*phot1-5* (Col-gl), *phot2-1* (Col-gl) and a double mutant, *phot1-5 phot2-1* (Col-gl)) determined immediately after 15 min illumination with RL (220 µmol m^{-2} s^{-1}), or R+BL (220 µmol m^{-2} s^{-1}, consisting of 90 % RL and 10 % BL; Materials and methods). The columns are mean values (±SE) of the indicated number of biological repeats from three independent experiments. Different letters denote significantly different values (ANOVA, *Post hoc* Tukey's test, P< 0.05) . Note the reduced response to RL+BL in the mutant lines.

F1. Grunwald, Gosa et al., 2021

Figure 2. amiR-silencing PHOT1 or PHOT2 (SCR:mirphot plants) in the BSCs reduces K_{leaf} under RL+BL, generated in WT (Col; Materials and methods). K_{leaf} was determined in fully expanded leaves of WT and *SCR:mirphot1* and *SCR:mirphot2* plants treated with RL or RL+BL (as in Fig. 1; Materials and Methods) The columns are means (± SE) of the indicated number of biological repeats from three independent experiments. Different letters denote significantly different values (ANOVA, Post hoc Tukey's test, P< 0.05).

Figure 3. BSCs-directed *PHOT1* complementation of the *phot1-5* mutant restores the normal K$_{leaf}$. Fully expanded leaves of WT (Col-gl), *phot1-5* (Col-gl) and *phot1-5* (Col-gl) plants complemented with *SCR:PHOT1* underwent illumination treatments as in Fig. 1. The columns are means (± SE) of the indicated number of biological repeats from three independent experiments. Different letters denote significantly different values (ANOVA, *Post hoc* Tukey's test, P< 0.05).

F3. Grunwald, Gosa et al., 2021

Figure 4. **Xylem-fed kinase Inhibitor tyrphostin 9 abolished the blue-light-induced K_{leaf} increase and stomatal opening.** Fully expanded leaves of WT (Col-gl) were pre-incubated (petiole deep) in AXS without or with tyrphostin 9 (10 μM) or sprayed with AXS without or with tyrphostin 9 (10 μM), and kept overnight in dark boxes. Immediately after dark, leaves were illuminated for 15 min with RL (215 μmolm^{-2} s^{-1}) or RL+BL (215 μmol m^{-2} s^{-1}, consisting of 90 % RL and 10 % BL). **a.** K_{leaf}. **b.** Stomatal conductance (g_S). The columns are means (± SE) of the indicated number of biological repeats, determined in three independent experimental days. Different letters denote significantly different values (ANOVA, Post hoc Tukey's test, P< 0.05). Note that K_{leaf} was reduced when tyrphostin 9 was fed via the petiole but not when sprayed, and the opposite was true for the $g_{S/}$

Figure 5. - Blue light (BL) hyperpolarizes the BSCs and alkalinizes their cytosol relative to red light (RL) or Dark and tyrphostin 9 depolarizes them. a. Membrane potential (E_M) of BSC protoplasts, without or with petiole-fed tyrphostin 9 (10 µM), determined using the dual-excitation, ratiometric fluorescent dye di-8-ANEPPS (Materials and Methods). After at least10-20 min in the dark, the cells were immediately imaged (Dark), or illuminated for an additional 5-10 minutes either with RL (220 µmol m^{-2} s^{-1}) or with RL+BL (220 µmol m^{-2} s^{-1}, including 10% BL; Materials and Methods), and then imaged. The columns are mean values of fluorescence ratio (F_{438}/F_{531}) values (±SE; n as indicated) derived from analyses of the images captured at the indicated excitation wavelengths (Materials and Methods) in three independent experiments. Numbers above the columns are the mean values of BSCs E_M, based on a calibration curve of F_{438}/F_{531} vs. E_M obtained in separate experiments using patch clamp (Materials and Methods, Suppl. Fig. S8). b. Cytosol pH (pH_{CYT}) of BSCs protoplasts determined using the fluorescent pH probe SNARF1 (Materials and methods). Protoplasts were dark-treated then further treated with 10 min of either dark, RL+BL or RL only. pH_{CYT} values were derived from analyses of the images captured (Materials and Methods) and determined from the emitted fluorescence ratio (F640 /F585) values, based on the in-situ calibration curve (detailed in Torne et al., 2021). The columns are means (± SE) of the indicated number of biological repeats, determined in three independent experimental days. Different letters denote significantly different values (ANOVA, Post hoc Tukey's test, P< 0.05).

Figure 6. The BSCs H$^+$-ATPase, AHA2, mediates the blue-light-induced K$_{leaf}$ increase. Fully expanded excised leaves of WT (Col0) plants, *aha2-4* mutant plants (Col0) and *aha2-4* plants complemented in their BSCs with *SCR:AHA2* were illuminated for 15 min immediately after dark, as in Fig.2. The columns are means (± SE) of the indicated number of biological repeats from three independent experiments. Different letters denote significantly different values (ANOVA, Post hoc Tukey's test, P< 0.05). Note the marked restoration of K$_{leaf}$ in the *aha2* mutant plants complemented solely in their BSCs.

F6. Grunwald, Gosa et al., 2021

Figure 7. White light (WL) illumination for 30 minutes acidifies the xylem pH of WT plants and *SCR*:*PHOT1*-complemented mutants, but not in the *phot1-5* mutant. The leaves of WT (Col-gl) plants, *phot1-5* (Col-gl) mutants and the *SCR:PHOT1*-complemented *phot1-5* mutant (as in Fig. 3) were illuminated by the growth room WL (150-200 µmol m^{-2} sec^{-1}, Materials and Methods). The columns are means (±SE) of the indicated number of biological repeats from three independent experiments Different letters denote significantly different values (*Post hoc* Tukey's test, $P < 0.05$).

Figure 8: White light increases the BSCs membrane osmotic water permeability coefficient (P_f). **a.** A time course of swelling of bundle sheath protoplasts from *SCR:GFP* plants upon exposure to a hypotonic XPS[db] solution under the same illumination conditions (white light or darkness) as during the light treatment preceding the assay. The bath solution was buffered to pH 5.7. The arrow indicates onset of bath flush. **b.** Time course of the osmotic concentration change in the bath (C_{out}) during the hypotonic challenge **c.** Mean (±SE) initial P_f values of the indicated number of bundle sheath protoplasts under two light regimes from three independent experiments. Columns are means (± SE) of 21-28 biological repetitions, determined in five independent experiments. The asterisk indicates a significant difference (Student's two-tailed unpaired t-test, $P < 0.05$).

a

b

Figure 9. Proposed BSCs-autonomous BL signal transduction pathway.
a. An artist's rendering of a leaf radial water path, from xylem to mesophyll
(K_{leaf}, blue hollow arrow) via the bundle sheath cells, BSCs, which tightly
envelop the xylem. GC, a stomata guard cell; MC, mesophyll cell. **b.** Blue-light
(BL) signalling pathway (blue arrowheads) in a bundle sheath cell, from BL
perception by the phototropin receptors (PHOT), through an intermediate
tyrphostin-sensitive tyrosine phosphorylation (TP), to the ultimate AHA2
(orange elipse) activation resulting in proton extrusion via the pump and xylem
sap acidification, presumably, at the expense of ATP breakdown to ADP with a
transient phosphorylation (P) on the pump protein, as expected from a P-type
H^+-ATPase.

F9. Grunwald, Gosa et al., 2021