Myogenesis modelled by human pluripotent stem cells uncovers Duchenne muscular dystrophy phenotypes prior to skeletal muscle commitment

Virginie Mournetas[1*], Emmanuelle Massouridès[2], Jean-Baptiste Dupont[1], Etienne Kornobis[4], Hélène Polvèche[2], Margot Jarrige[2], Maxime Gosselin[5]; Spiros D. Garbis[6]; Dariusz C. Górecki[5]; Christian Pinset[3]

[*] Correspondence: vmournetas@istem.fr

[1]INSERM UEVE UMR861, I-STEM, AFM, 28 rue Henri Desbruères, 91100 Corbeil-Essonnes, France

[2]CECS, I-STEM, AFM, 28 rue Henri Desbruères, 91100 Corbeil-Essonnes, France

[3]CNRS, I-STEM, AFM, 28 rue Henri Desbruères, 91100 Corbeil-Essonnes, France

[4]Institut Pasteur, 25-28 Rue du Dr Roux, 75015 Paris, France

[5]Molecular Medicine, School of Pharmacy and Biomedical Sciences, University of Portsmouth, PO1 2DT, Portsmouth, UK

[6]Proteome Exploration Laboratory, Beckman Institute, California Institute of Technology, Division of Biology and Biological Engineering, 1200 E. California Blvd., MC 139-74, Pasadena, California, 91125, USA

ABSTRACT

Duchenne muscular dystrophy (DMD) causes severe disability of children and death of young men, with an incidence of approximately 1/5,000 male births. Symptoms appear in early childhood, with a diagnosis made around 4 years old, a time where the amount of muscle damage is already significant, preventing early therapeutic interventions that could be more efficient at halting disease progression. In the meantime, the precise moment at which disease phenotypes arise – even asymptomatically – is still unknown. Thus, there is a critical need to better define DMD onset as well as its first manifestations, which could help identify early disease biomarkers and novel therapeutic targets.

In this study, we have used human induced pluripotent stem cells (hiPSCs) from DMD patients to model skeletal myogenesis, and compared their differentiation dynamics to healthy control cells by a comprehensive multi-omics analysis. Transcriptome and miRnome comparisons combined with protein analyses at 7 time points demonstrate that hiPSC differentiation 1) mimics described DMD phenotypes at the differentiation endpoint; and 2) homogeneously and robustly recapitulates key developmental steps - mesoderm, somite, skeletal muscle - which offers the possibility to explore dystrophin functions and find earlier disease biomarkers.

Starting at the somite stage, mitochondrial gene dysregulations escalate during differentiation. We also describe fibrosis as an intrinsic feature of skeletal muscle cells that starts early during myogenesis. In sum, our data strongly argue for an early developmental manifestation of DMD whose onset is triggered before the entry into the skeletal muscle compartment, data leading to a necessary reconsideration of dystrophin functions during muscle development.

INTRODUCTION

Duchenne muscular dystrophy (DMD) is a rare genetic disease, but it is the most common form of myopathy affecting approximately one in 5,000 male births and very rarely female. In this recessive X-linked monogenic disorder, mutations in the DMD gene lead to the loss of a functional dystrophin protein, resulting in a progressive - yet severe - muscle wasting phenotype (1). In patients, symptoms usually appear in early childhood (2-5 years old) and worsen with age, imposing the use of wheelchair before 15 and leading to premature death by cardiac and/or respiratory failure(s) mostly around 30 years of age (2).

At the age of diagnosis, around 4 years old, muscles of DMD patients have already suffered from the pathology (3,4). Several reviews pointed out the limitations of current disease biomarkers, which fail to detect the development of DMD specifically and at an early age (5,6). Meanwhile, no treatment is available to stop this degenerative disease yet. Developing therapies aim at restoring the expression of dystrophin in muscle cells but, so far, the level stays too low to be beneficial to patients (7). The absence of both reliable biomarkers and effective therapies stress the need of better defining the first steps of DMD in Human to be able to 1) find specific markers of disease initiation in order to increase diagnosis sensitivity and, therefore, improve patient management by accelerating their access to better healthcare; and 2) develop alternative therapeutic approaches by finding targets that compensate the lack of dystrophin and complement current attempts at restoring its expression (8).

In 2007, a seminal publication reported that the gene expression profile of muscles from asymptomatic DMD children younger than 2 years old is already distinguishable from healthy muscles, suggesting that DMD molecular dysregulations appear before disease symptomatic manifestations (4). Evidence obtained in multiple animal models, such as neonatal *GRMD* dogs (9), DMD zebrafish (10) and *mdx* mouse embryos (11), as well as in human foetuses (12–14) even suggest that DMD starts before birth, during prenatal development. Our team recently identified the embryonic dystrophin isoform Dp412e expressed in early mesoderm-committed cells (15), another indication that DMD can start *in utero*. Further exploring DMD onset in human foetuses is extremely challenging for obvious ethical and practical reasons. A way to overcome these issues is to develop a human DMD model *in vitro*, recapitulating embryonic development from human pluripotent stem cells to skeletal muscle lineages.

To our knowledge, none of the existing human DMD *in vitro* models, either based on tissue-derived myoblasts (16) or on the differentiation of induced pluripotent stem cells (17–21), have been used for studying DMD during the ontogeny of the skeletal muscle lineage. Moreover, original protocols for *in vitro* myogenesis from human pluripotent stem cells (reviewed in (22)) use transgene overexpression or/and cell sorting procedures, and thereby, miss the steps preceding skeletal muscle commitment, *e.g.* paraxial mesoderm and myotome. Novel protocols have recently used transgene-free directed differentiation to recapitulate human embryonic development in a dish, giving theoretical access to the developmental steps (19,23–25).

In the present study, we compared the myogenic differentiation dynamics of healthy and DMD hiPSCs using a multi-omic approach to identify early disease manifestations *in vitro*. DMD cells showed marked transcriptome dysregulations from day 10, before the detection of skeletal muscle regulatory factors at day 17. Specifically, we identified the dysregulation of mitochondrial genes as one of the earliest detectable phenotypes. These alterations escalated over the course of muscle specification. In addition, we showed an early induction of Sonic hedgehog signalling pathway, followed by collagens as well as fribrosis-related genes suggesting the existence of an intrinsic fibrotic process solely driven by DMD muscle cells. Overall, our data highlight that human pluripotent stem cells are a suitable cell model to study muscle development in both healthy and disease conditions. In the context of DMD, they strongly argue for the existence of early disease manifestations during somite development.

 RESULTS

To establish the early/developmental impact of *DMD* gene mutations, human induced pluripotent stem cells
(hiPSCs) from three DMD patients and three healthy individuals were generated as described previously (15).
These cells, when subjected to a standardised differentiation protocol without utilisation of feeder cells, cell
sorting or gene overexpression, formed elongated and plurinucleated myotubes within 25 days (23), with an
amplification fold of 2918 ± 480 (mean ± SEM). Skeletal muscle progenitor cells after 17 days of differentiation
could be cryopreserved (Figure S1A). Whole transcriptome and miRnome profiles were compared at 7
differentiation time points (tissue-derived myoblasts and myotubes, as well as hiPSC-derived cells at days 0, 3,
10, 17 and 25) and complemented by iTRAQ proteomics and Western blot analyses (Table S1).

DMD is initiated prior to the expression of skeletal muscle markers

First, the expression profile of the *DMD* variants was studied by RT-qPCR in healthy and DMD hiPSCs during the
differentiation process described in Figure S1A. The *Dp427m* variant, which is normally observed in muscle
cells (26), appeared from day 17, in contrast with *Dp412e* – the embryonic variant of dystrophin present in
mesoderm cells (15) – which was expressed at differentiation day 3. Therefore, the expression of the *DMD*
locus is initiated in the very first steps of our differentiation protocol, well before myotube formation. The
ubiquitous variant *Dp40-71* was detected at every time points, in contrast with *Dp116* (Schwann cell variant
(27)), *Dp140* (kidney and foetal brain variant (28)) and *Dp427p1p2* (Purkinje cell variant (29)), which were all
undetected at the examined time points (Figure S1B). Interestingly, *Dp260* (retinal variant (30)) followed a
similar expression pattern than *Dp427m*. *Dp427c* (cerebral variant (31)) was also detected at days 17 and 25,
but at a very low level. As expected, DMD cells expressed lower levels of *Dp427m* and *Dp260* (Figure S1B).

A strong correlation in the transcriptomic data was observed by mRNA-seq and miRNA-seq between samples
collected at an individual time point, as opposed to samples from two distinct time points. In addition, the
correlation coefficient between samples taken at two successive time points increased as differentiation
progressed (Figure 1A). Differential expression analysis between two successive collection days (days 3/0, days
10/3, days 17/10, days 25/17) in healthy controls showed that the proportion of regulated genes decreased
from 20 % to 12 % of the whole transcriptome (2223 to 1284 mRNAs, adjusted pvalue ≤ 0.01) through the
course of differentiation. These observations demonstrate the robustness of the differentiation protocol and
are in agreement with an early specialisation and a later refinement of the transcriptome as cells quickly exit
pluripotency and become progressively restricted to the skeletal muscle lineage.

To characterise the developmental stages achieved by the cells, the expression of lineage-specific markers
(both mRNAs and miRNAs) was determined at each time point, together with gene enrichment analyses
(Figure 1B-2A, Figure S2B-C, Table S2).
Pluripotency was similarly maintained in healthy and DMD cells at day 0 (Figure 2A – Table S2), as already
shown by our group (15). At day 3, cells lost pluripotency and became paraxial mesoderm cells expressing
marker genes such as *PAX3* and *PAX7* (11) (Figure 2A, Table S2). Importantly, markers of lateral plate (*GATA4*
(32) and *NKX2-5* (33)) and intermediate mesoderm (*PAX8* (34) and *ORS1* (35)) were undetected at this stage
(Table S2). Similarly, earlier markers of primitive streak (*EOMES* (36) and *TBX6* (37)), mesendoderm (*NODAL*
(38) and *MIXL1* (39)), as well as markers of the other germ layers endoderm (*AFP* (40) and *SOX17* (41)) and
ectoderm (*PAX6* (42) and *SOX2* (43)) were either not expressed or greatly downregulated (Table S2),
suggesting cell homogeneity in the differentiation process.

At that early time point, DMD-associated gene dysregulation represented less than 3 % of the entire
transcriptome (adjusted pvalue ≤ 0.05, Figure 2B) but already contained genes important for development
(*e.g. MEIS2* (44)) and muscle formation (*e.g. ACTA1* (45)). However, mesoderm markers were not significantly
dysregulated, attesting that mesoderm commitment was mostly unimpaired (Figure 2A – Table S2). No

increase in the expression of primitive streak, mesendoderm, endoderm or ectoderm markers was detected,
suggesting no differences in the differentiation process of DMD cells at that stage (Table S2).

In contrast, a sharp increase in the proportion of dysregulated genes appeared at day 10 compared to day 0
and day 3, mostly including gene downregulations (DMD/Healthy expression ratio ≤ 0.76, adjusted pvalue ≤
0.05). This concerned almost 10 % of the transcriptome at day 10 (against 3 % at day 3) and remained stable
from 10 to 12 % (1226 mRNAs) until day 25 (Figure 2B). At day 10, healthy cells started to express genes
typically observed during somitogenesis, such as *PAX3* (46) *NR2F2* (47), *PTN* (48), *MET* (49), *H19* and *IGF2* (50)
(Table S2). More precisely, their transcriptome exhibits a mixed profile between dermomyotome (expression
of *GLI3* (51) and *GAS1* (52) but not *ZIC3* (53)) and myotome (expression of *MET* (54) and *EPHA4* (55) but not
LBX1 (56)) (Table S2). Neither markers of presomitic mesoderm cells (*MSGN1* (57) and *FGF8* (58)) and neural
plate cells (*SIX1* (59) and *FOXD3* (60)), nor markers of sclerotome (*PAX1* (61) and *PAX9* (62)) and dermatome
(*EGFL6* (63)) were expressed (Table S2). In DMD cells, no increase of presomitic mesoderm, neural plate,
sclerotome or dermatome markers was observed (Table S2). In the meantime, several somite markers were
downregulated, including *H19, IGF2*, *MET* and *SEMA6A* (64) (validated at the protein level for SEMA6A, Figure
2A-S3A – Table S2), while a slight upregulation of dermomyotome and chondrocyte markers was highlighted
and confirmed at the protein level for GLI3 (Figure S3B), together with a significant enrichment of the gene
ontology term 'neurogenesis', suggesting potential lineage bifurcations at day 10 (Figure 2A-S2C – Table S2).

The study of differentiation dynamics presented above highlights that mesoderm commitment is not impaired
by the absence of dystrophin, and shows that DMD onset takes place at the somite cell stage, before the
expression of the skeletal muscle program and especially before the expression of *Dp427m*, the muscle variant
of the *DMD* gene.

DMD hiPSC can become skeletal muscle progenitor cells, but exhibit specific muscle gene dysregulations

Healthy and DMD cells were in the skeletal muscle compartment at day 17, as evidenced by the expression of
multiple lineage-specific genes and proteins, such as transcription factors (*e.g. MYOD1* (65)), cell surface
markers (*e.g. CDH15* (66)), sarcomere genes (*e.g. TNNC2* (67)), dystrophin-associated protein complex (DAPC)
genes (*e.g. SGCA* (68)), Calcium homeostasis genes (*e.g. RYR1* (69)) and muscle-specific miRNAs (myomiR, *e.g.*
miR-1 (70)), (Figure 1B, Table S2). They both showed an embryonic/foetal phenotype characterised by *ERBB3*
expression, in contrast with tissue-derived myoblasts that expressed *NGFR* (21). Here again, alternative cell
lineages were absent or greatly downregulated, such as tenocytes (*MKX* (71) and *SCX* (72)), chondrocytes
(*SOX5* and *SOX9* (73)), osteoblasts (*SP7* (74) and *SPP1* (75)) or nephron progenitors (*SIX2* (76) and *SALL1* (77))
(Table S2).
Interestingly, DMD cells did not show dysregulated expression of skeletal muscle transcription factors (Table
S2). However, several myomiRs were found downregulated (*e.g. miR-1*, Figure 2C), together with genes related
to calcium homeostasis (*e.g. ATP2A2* (78), at both mRNA and protein level, Figure 2D-E) as well as members of
the DAPC (*e.g. SNTA1* (79)) (Table S2). Concerning cell lineages, there was no visible difference when
compared to healthy controls, except an upregulation of markers associated with chondrocytes, which was
confirmed at the protein level for GLI3 (Figure S3C), and a significant enrichment of the gene ontology term
'neurogenesis' previously seen at day 10 (Figure 2A-S2C – Table S2).
Therefore, DMD cells efficiently enter the skeletal muscle compartment at day 17, but exhibit dysregulations in
several features typically associated with dystrophic muscles, which could be a direct consequence of the early
manifestations of DMD detected at day 10.

hiPSC differentiation lead to embryonic/foetal myotubes that reproduce DMD phenotypes

As previously described (23), both healthy and DMD hiPSC-derived myotubes (day 25) were able to twitch
spontaneously in culture, and fluorescent staining of nuclei and α-actinin confirmed cell fusion and the
formation of striation patterns typical of muscle fibres *in vivo* (Figure 3A). Western blot analyses on protein

extracts from DMD cells confirmed that dystrophin was either undetectable or slightly expressed (Figure 3B),
as in the corresponding patient muscle biopsies.
We selected representative mRNAs and miRNAs and showed that both hiPSC-derived and tissue-derived
myotubes have exited the cell cycle and upregulated genes expressed in skeletal muscles (Figure S4A, Figure
4A, Table S2). This included skeletal muscle myomiRs (miR-1, miR-133 and miR-206 (80,81)), transcription
factors involved in skeletal myogenesis including those of the MRF family (*e.g. MYOD1* (65), *MYOG* (82)),
specific muscle cell surface markers (*e.g. CDH15* (66), *ITGA7* (83)) as well as genes involved in the formation of
the DAPC (*e.g. SGCA* (68), *DTNA* (84)), sarcomeres (*e.g. TNNC2* (67), *TNNT3* (85)), myofibril organisation (*e.g.*
UNC45B (86), *NACA* (87)) and the execution of excitation-contraction coupling at the neuromuscular junction
(NMJ, *e.g. MUSK* (88), *DOK7* (89)) (Figure 4A, Table S2).
Even though global analysis showed that hiPSC-derived myotubes were similar to their tissue-derived
counterparts in term of lineage commitment, they displayed an embryonic/foetal phenotype – as suggested in
progenitors at day 17. This can be illustrated by the expression of the embryonic/foetal myosin heavy/light
chains *MYH3* (90), *MYH8*(91), *MYL4* (92) and *MYL5* (93) but not the postnatal transcripts *MYH1* and *MYH2* (94),
which were detected in tissue-derived myotubes. Myotubes derived from hiPSCs had also higher levels of *IGF2*,
which is downregulated at birth (95), and expressed *DLK1*, which is known to be extinct in adult muscles (96)
(Figure S4B).
Despite the embryonic/foetal phenotype, hiPSC-derived myotubes showed evidence of terminal
differentiation and cellular maturation. First, their total level of myosin heavy chain proteins was significantly
higher than in tissue-derived myotubes, as confirmed by Western blotting (Figure 3B). RNAs and proteins
involved in DAPC formation (*e.g.* DMD, SGCA (68) and SGCG (97)), as well as in excitation-contraction coupling
(*e.g.* RYR1 (69) and CACNA1S / CAV1.1 (98)) were also present at higher levels (Figure 3B-4A). Finally, higher
expression of skeletal muscle transcription factors (*e.g. MEF2C* (99)), and of multiple genes involved in muscle
contraction (*e.g. TNNT3* (85)), NMJ formation (*e.g. RAPSN* (100)), and creatine metabolism (*e.g. CKM* (101))
indicates that hiPSC-derived cells expressed features of fully differentiated muscle cells (Figure 4A). Similar to
previous time points, day 25 cells were negative for markers of alternative muscle lineages, *i.e.* cardiac (*miR-*
208a (102), *MYL7* (103) and *RYR2* (104)) and smooth muscle cells (*MYH11* (105), *CNN1* (106) and
CHRNA3/B2/B4 (107)).
In DMD cells, there was a global trend toward downregulation of muscle transcription factors, which was only
significant for *MEF2A* and *MEF2D* in hiPSC-derived myotubes and *EYA4* and *MYOD1* in tissue-derived myotubes
(Figure S4C). In addition, myomiRs previously associated with muscle dystrophy (dystromiRs, *e.g.* miR-1 (70),
Figure 2C) were found downregulated (Table S2). Similarly, a global downregulation phenotype was observed
in both tissue- and hiPSC-derived DMD myotubes, and concerned multiple genes associated with known
disease phenotypes, such as cell surface markers (*e.g. ITGA7* (83)), DAPC organisation (*e.g. SGCA* (68)),
myofibril organisation (*e.g. UNC45B* (86)), sarcomere formation (*e.g. MYO18B* (108)), NMJ function (*e.g.*
CHRNB1 (109)) and calcium homeostasis (*e.g. ATP2A2* (78), Figure 2D)(Figure 4B-S2C).
Altogether, these data indicate that hiPSC-derived myotubes recapitulate a full skeletal muscle differentiation
program, and exhibit an embryonic/foetal phenotype. Importantly, it shows that disease phenotypes usually
observed in young adult animal models are detectable at least at the transcriptional level in human DMD
myotubes that display an otherwise embryonic/foetal profile. This validates the quality of this cell system to
model the DMD pathology.

Fibrosis, an intrinsic feature of DMD cells independent of TGF-β pathway

As presented above, the upregulation of chondrocyte markers in DMD cells, although already present at day
10, became significant from day 17 (Figure 2A – Table S2). It was accompanied by the upregulations of the
Sonic hedgehog (SHH) signalling pathway and of multiple collagens (Figure 5A, Table S2). Genes encoding the
P4H collagen synthases, were not dysregulated while *RRBP1* (that stimulates collagen synthesis (110)) together
with *PLOD1* and *PLOD2* (that stabilise collagens (111,112)) were significantly upregulated. Moreover, *SETD7*, a
gene known for activating collagenases (113), was significantly downregulated.

At the myotube stage, a fibrosis-related gene set was clearly upregulated in DMD cells, as illustrated by the
overexpression of *ANGPT1* (114), *CTGF* (115), collagens (*e.g. COL1A2* (116)), matrix metallopeptidases (*MMPs*)
and tissue inhibitors of metallopeptidase (*TIMPs*) (117) (Figure 5B). Conversely, the myomiR miR-133b that
controls *CTGF* expression (118) was repressed (Table S2). Interestingly, gene members of the transforming
growth factor (TGF)-β pathway, a well-known inducer of fibrosis (119), were not found dysregulated (Figure
5B).
Altogether, these data argue for fibrosis as an intrinsic feature of DMD skeletal muscle cells, rather than a
process solely driven by interstitial cell populations in the niche. Furthermore, this muscle-driven fibrosis
seems independent of the TGF-β pathway, and could rather depend on the SHH pathway, together with an
intrinsic upregulation of chondrocyte markers and collagens.

Mitochondria, a key organelle impacted by the absence of dystrophin prior calcium dysregulation

As previously described (120) and illustrated on Figure S5A, the energy metabolism of DMD hiPSC-derived
myotubes was dysregulated at the creatine and carbohydrate levels, up to the respiration (Figure 6A-B, Figure
S2C). The creatine transporter was not impacted while mRNAs coding for enzymes of both creatine and
creatine phosphate biosynthesis were underrepresented. Neither glucose nor glutamate transporter
expression were impaired. However, glutamine biosynthesis (followed by gluconeogenesis that feeds glycolysis
from glutamine) as well as glycogenesis (followed by glycogenolysis that feeds glycolysis from glycogen) were
all downregulated, together with glycolysis itself. In contrast, the pentose phosphate pathway, which is in
parallel to glycolysis, was upregulated, especially the oxidative part. Pyruvate decarboxylation and generation
of acetyl-CoA to feed the tricarboxylic acid (TCA) cycle was also impaired. Finally, the TCA cycle itself (Figure
6A, Figure S2C) and the mitochondrial electron transport chain were downregulated Figure 6B, Figure S2C).
This is particularly reinforced by lower levels of a member of the ATP synthase complex ATP5A1 at both mRNA
and protein levels (Figure 6C-D). Moreover, transcripts encoded by the mitochondrial DNA and mitochondrial
DNA itself were decreased in DMD hiPSC-derived myotubes at day 25 (Figure S5B-S5E).
In the presented cell model, a significant downregulation of a mRNA set coding for mitochondrial proteins was
primarily observed at day 10 with the downregulation of 11 % (10 mRNAs, DMD/Healthy expression ratio ≤
0.76, adjusted pvalue ≤ 0.05) of the mitochondrial outer membrane genes, and amplified during the
differentiation of DMD cells (Figure 7A). Therefore, defects depicted at day 25 rooted before the expression of
the skeletal muscle program at day 17. Among them, mRNA downregulation of *TSPO*, a channel-like molecule
involved in the modulation of mitochondrial transition pore (121), occurred from day 10 to day 25. This
downregulation was also observed at the protein level at day 17 (Figure 7B). Moreover, the protein import
system was affected from day 17 at both mRNA and protein levels (Figure S5C-S5F). Simultaneously, mRNAs
involved in mitochondrial genome transcription started to be downregulated, followed by genes involved in
mitochondrial DNA replication at day 25 (Figure S5D-S5G). This progressive increase of dysregulations was also
observed at the level of the entire mRNA set related to mitochondria (around 1,000 mRNAs) as illustrated by
the volcano plots as well as the gene ontology enrichments (Figure 7C, Figure S2C).
Our data highlight early impairments in genes coding for mitochondria that start at the somite stage - prior
calcium homeostasis dysregulation - and increase with the differentiation in an orderly manner. These
elements complete the mitochondrial DMD phenotype described at the myotube stage.

Altogether, our study demonstrates that DMD starts prior to the expression of well-described markers of
muscle differentiation. It shows that hiPSC-based experimental models of DMD can help identify early disease
manifestations and stratify multiple pathological features over the course of muscle development.

DISCUSSION

Since the discovery of the *DMD* gene in 1987 (1), DMD cellular phenotypes were considered under the unique scope of a "mechanical hypothesis" in which dystrophin deficiency led to membrane leakage and ultimately muscle cell rupture. However, over the last 15-20 years, studies have brought unequivocal evidence that multiple additional factors are in play, such as calcium intracellular overloads (122,123), excessive oxidative stress (124,125), metabolic switches (126,127), as well as an overall tissue context where aberrant interactions between resident cells lead to inflammation and fibro-adipogenesis (128–130). This has progressively led to a complex picture involving interdependent homeostatic perturbations and to date, the identification of prevalent pathological features driving the initiation of DMD is hardly feasible.

The skeletal myogenesis modelled here by the differentiation of hiPSCs, without gene overexpression or cell sorting, homogeneously and robustly recapitulates key developmental steps – pluripotency, mesoderm, somite and skeletal muscle – without any trace of other lineages. Therefore, it is a suitable dynamic model for studying human skeletal muscle development in both healthy and DMD cells, offering the possibility to clarify the consequences of the absence of dystrophin at each step of the differentiation process, as well as to explore dystrophin functions and find earlier and more specific disease biomarkers.

As previously observed with pluripotent stem cells (131), hiPSC-derived myotubes at day 25 displayed an embryonic/foetal gene expression profile. However, a clear distinction must be made between the nature of the expressed isoforms – embryonic / foetal / postnatal – and the degree of differentiation. For instance, hiPSC-derived myotubes expressed multiple markers of terminally differentiated muscles at levels higher than those measured in tissue-derived myotubes. With the idea of exploring human DMD phenotypes during muscle development, we argued that generating embryonic/foetal myotubes from hiPSCs would not be a limitation.

In qualitative terms, DMD hiPSC-derived myotubes showed an overall morphology similar to healthy controls, with cell fusion and clear striation patterns, suggesting that the potential impact of dystrophin during *in vitro* differentiation is subtle and does not prevent myotube formation. However, our unbiased mRNA-seq analysis highlighted striking transcriptome dysregulations with 3,578 differentially expressed genes at day 25. This includes numerous genes which can be linked to previously described DMD phenotypes such as 1) DAPC dissociation (132); 2) rupture of calcium homeostasis (122); 3) myomiR downregulation (70,133); 4) sarcomere destabilisation (134–136); 5) mitochondrial and metabolism dysregulations (126,127); 6) NMJ fragmentation (137,138) and 7) fibrosis (130,139). It is interesting to note that these phenotypes are already detected at the transcriptional level in embryonic/foetal myotubes, while they usually appear postnatally in Human and other animal models. In addition, most of them are often considered as consequences of degeneration-regeneration cycles typical of DMD muscles *in vivo* (135,140,141) which are absent in our *in vitro* model, indicating that a part of these defects are primarily due to the absence of dystrophin itself. In particular, our data suggest that fibrosis is an intrinsic feature of DMD skeletal muscle cells, and therefore, it does not absolutely require a specific tissue context or additional cell populations to be detected *in vitro*. Fibrosis is a major hallmark of DMD pathophysiology, and the regulation of this process has been largely investigated in the past (119,142). A long-debated question is the implication of the TGFβ signalling pathway (143). In our model, TGFβ signalling was inhibited up to day 17 by inhibitors in the cell culture media, and TGFβ-related genes were not upregulated at day 25, suggesting that the observed fibrosis is TGFβ- independent.

Since several studies on Human and other animal models had described dystrophic phenotypes in DMD foetuses/infants (9–14), we investigated the precise timing of disease onset in our hiPSC-derived myotubes. First, the absence of dystrophin does not modify the capacity of cells derived from adult tissue biopsies to be reprogramed using the approach developed by Takeshi and Yamanaka (144). Both healthy and DMD cells retained pluripotency and the capacity to enter the mesoderm compartment at day 3. At that time, the embryonic dystrophin Dp412e is expressed and only marginal dysregulations are observed in DMD cells, *a*

priori unrelated to cell fate choice as cells only express paraxial mesoderm markers at level similar to healthy
controls.

DMD dysregulations are greatly increased at day 10, when cells express somite markers without any trace of
transcripts coding for the long dystrophin isoform Dp427m expressed in skeletal muscles (only the ubiquitous
variant Dp71-40 is found). At that time, we noticed few significant dysregulations of cell lineage markers,
which became more prevalent at day 17 and 25. This might be an indication that to some extent, cell fate is
misguided in DMD cells, where skeletal muscle markers are underexpressed and replaced by markers of
alternative lineages, such as chondrocytes.

First visible at day 10, we identified mitochondrial dysregulation as one of the key processes happening in an
orderly manner. Interestingly, early observations prior the discovery of the *DMD* gene had hypothesised that
DMD was a mitochondrial/metabolic disease based on protein quantifications and enzyme activities (126,145).
Later, mitochondria was identified as a key organelle in DMD, responsible for metabolic perturbations but also
calcium accumulation and generation of reactive oxygen species (122–125). In this study, numerous genes
coding for proteins located in the outer mitochondrial membrane start to be downregulated from day 10 in
DMD cells, such as the benzodiazepine receptor TSPO, a member of the controversial mitochondrial
permeability transition pore (mPTP) (121). The mPTP is a multiprotein complex whose members are not all
precisely identified, and several studies suggest that it might be involved in DMD pathophysiology (146,147). A
chicken-and-egg question currently debated relates to the initiation of these homeostatic breakdowns, as
positive feedbacks exist between mitochondria, oxidative stress and calcium homeostasis dysregulations
(123,124). In our model, dysregulations of genes controlling calcium homeostasis were detected after day 10,
suggesting that mitochondrial impairment starts early and has predominant consequences in DMD, as
hypothesised by Timpari *et al*. (120). Further experiments are needed to better evaluate the impact of
mitochondrial dysregulations at the functional level.

Day 17 marks the entry in the skeletal muscle compartment with the expression of specific transcription
factors, cell surface markers, the skeletal muscle variant of dystrophin (*Dp427m*), as well as myomiRs. It also
marks the initiation of the skeletal muscle gene dysregulations observed at the myotube stage (*i.e.*
downregulation of genes involved in DAPC and calcium homeostasis). For instance, the upregulation of
fibrosis-related genes observed in DMD myotubes at day 25 is already visible at day 17, with the upregulation
of the SHH pathway as well as collagen-related genes. In this study, it is seen as an early indicator of DMD
physiopathology, confirming previous observations in DMD infants, both transcriptionally (4) and histologically
(148,149).

Moreover, several myomiRs were found downregulated at days 17 and 25 and seem to play a central part in
multiple DMD phenotypes. Besides their role in myogenesis (80,81), myomiRs can be involved in calcium
homeostasis (150), metabolism and mitochondrial functions (151,152), and fibrosis (118,153). In particular,
miR-1 and miR-206 are known to target key genes such as CACNA1C (150), CTGF (118), RRBP1 (153), several
regulators of the pentose phosphate pathway (151), and even transcripts encoded by the mitochondrial
genome (152). Even though the functional consequences of the multiple gene and myomiR dysregulations
highlighted in this study is virtually impossible to anticipate, we believe that myomiRs can be key players in
DMD biology.

Only few studies argued that DMD starts before the expression of the muscular dystrophin protein (18,154).
Our data supports this idea as disease phenotypes seem to be initiated at the somite stage where Dp427m was
not even transcribed. This could be explained by the deficit in other dystrophin isoforms expressed prior day
10, such as Dp412e at day 3 (15), but also by the decrease or loss of other RNA products expressed from the
DMD locus, such as the ubiquitous isoform Dp71-40 or long non-coding RNAs (155). The lack of knowledge
around these additional products contrasts with the extensive amount of data on the structure and function of
the main muscular isoform Dp427m whose most studied role is to stabilise muscle cell membrane during
contraction (156). Other tissue specific isoforms have been described, *e.g.* in the retina (Dp260 (30)) and in the
brain (Dp427c (31), Dp427p (29) and Dp140 (28)), some of which are also slightly expressed in skeletal muscles
under certain circumstances (157), but their role remains mostly unknown. Interestingly, in our data, the

expression of Dp260 follows the same pattern of expression as Dp427m. It has been shown that the expression
of Dp260 in *mdx*/utrnK/K mice can rescue the *mdx* phenotype (158), indicating overlapping functions between
Dp427m and Dp260. On the other hand, it is now well established that a third of DMD patients display
cognitive deficiencies – which might be correlated with mutations affecting Dp140 (159) – attesting that
dystrophin can be involved in other cell functions.

To date, the standard of care for DMD patients helps mitigate and delay some of the most severe symptoms
but remains insufficient to have a curative effect. Despite decades of work with the *mdx* mouse model, only a
few pharmacological candidate molecules have moved forward to clinical trials, with variable efficiency. As
several gene therapy trials have been recently initiated with promising preliminary data, we believe that our
human *in vitro* model system might be useful for the development of combination therapies. Recent studies
have proved that the association of two different therapeutic approaches could have a synergistic effect on
the overall treatment outcome, and can be used for instance to boost the effect of dystrophin re-expression by
antisense oligonucleotides or gene therapy (8,160,161). Here, our extensive RNA-seq data could help identify
relevant therapeutic targets for pharmacological intervention, such as CTGF – involved in fibrosis and found
upregulated in DMD myotubes – which can be inhibited by monoclonal antibodies (162), or TSPO receptor – a
receptor potentially member of the mPTP downregulated in DMD cells – targetable with benzodiazepines
(163). In addition, our model might also be used as a platform to screen pharmacological compounds in an
unbiased high-throughput manner. Indeed, skeletal muscle progenitor cells at day 17 can be robustly
amplified, cryopreserved and plated in a 384-well plate format (data not shown). Thus, they could be an
interesting tool to highlight pharmacological compounds to be used alone, or in combination with gene
therapy.

To summarise, the directed differentiation of hiPSCs without gene overexpression or cell sorting
homogeneously and robustly recapitulates key developmental steps of skeletal myogenesis and generates
embryonic/foetal myotubes without any trace of other lineages. The absence of dystrophin does not
compromise cell reprogramming, pluripotency or the entry into the mesoderm compartment. While none of
the long dystrophin isoform is expressed, a significant transcriptome dysregulation can be observed at the
somite stage that implicates mitochondria prior to defects in calcium homeostasis. Although being able to
enter the skeletal lineage compartment and become myotubes, DMD cells exhibit an imbalance in cell fate
choice as they express lower amount of key muscle proteins and retain basal expressions of other lineages,
leading to the well-characterised DMD phenotypes including muscle features and metabolism dysregulations
as well as fibrosis. Altogether, these data argue for 1) a deficit and not a delay in DMD differentiation; 2)
seeing DMD as a progressive developmental disease as well as a metabolic pathology whose onset is triggered
before the entry into the skeletal muscle compartment; and 3) fibrosis as an intrinsic feature of DMD muscle
cells. Future studies could explore the additional roles of *DMD* locus products with the impact of their loss all
along the skeletal muscle development, as well as find earlier and more specific disease biomarkers and
develop combination therapeutic strategies using high-throughput drug screening.

Materials and methods

Ethics, consent, and permissions

At the Cochin Hospital-Cochin Institute, the collection of primary cultures of myoblasts was established from
patient muscle biopsies conducted as part of medical diagnostic procedure of neuromuscular disorders. For
each patient included in this study, signed informed consent was obtained to collect and study biological
resources, and establish primary cultures of fibroblasts and myoblasts at the Hospital Cell Bank-Cochin
Assistance Publique—Hôpitaux de Paris (APHP). This collection of myoblasts was declared to legal and ethical
authorities at the Ministry of Research (number of declaration, 701, n° of the modified declaration, 701–1) via

the medical hosting institution, APHP, and to the "Commission Nationale de l'Informatique et des Libertés"
(CNIL, number of declaration, 1154515).

Cells

Human primary adult myoblasts from healthy individuals and DMD patients were provided by Celogos and
Cochin Hospital-Cochin Institute (Table S3). In Celogos laboratory, cell preparation was done according to
patent US2010/018873 A1.

Cell culture

Human tissue-derived myoblasts – Primary myoblasts were maintained in a myoblast medium: DMEM/F-12,
HEPES (31330–038, Thermo Fisher Scientific) supplemented with 10 % fetal bovine serum (FBS, Hyclone,
Logan, UT), 10 ng/mL fibroblast growth factor 2 (FGF2, 100-18B, Peprotech), and 50 nM Dexamethasone
(D4902, Sigma-Aldrich) on 0.1 % gelatin (G1393, Sigma-Aldrich) coated culture ware.

Human tissue-derived myotubes – Primary myoblasts were differentiated into myotubes. Cells were seeded at
600 cells/cm^2 on 0.1 % gelatin coated cultureware in myoblast medium containing 1 mM Acid ascorbic 2P
(A8960, Sigma-Aldrich).
Human induced pluripotent stem cells - Primary myoblasts were reprogrammed into hiPSCs following the
protocol described in (164), using the Yamanaka's factors POU5F1, SOX2 and KLF4 transduction by ecotropic or
amphotropic vectors (Table S3). HiPSCs were adapted and maintained with mTeSR™1 culture medium (05850,
Stemcell Technologies) on Corning® Matrigel® Basement Membrane Matrix, lactose dehydrogenase elevating
virus (LDEV)-Free-coated cultureware (354234, Corning Incorporated). Cells were then seeded at 20,000
cells/cm^2, passaged and thawed each time with 10 µM StemMACS™ Y27632.

Human iPSC-derived cell – Six hiPSCs (3 healthy and 3 DMD) were differentiated three times toward skeletal
muscle lineage using commercial media designed from Caron's work (23) (Skeletal Muscle Induction
medium SKM01, Myoblast Cell Culture Medium SKM02, Myotube Cell Culture Medium SKM03, AMSbio). This
protocol is a 2D directed differentiation that uses 3 consecutive defined media (SKM01 from day 0 to 10,
SKM02 from day 10 to 17 and SKM03 from day 17 to d25) and only one cell passage at day 10.

DNA and RNA experiments

RNA extraction and quality – RNA extraction was done in the six cell lines at 7 different time points: tissue-
derived myoblast and tissue-derived myotube, as well as during hiPSC differentiation at day 0, 3, 10, 17 and 25
(hiPSC-derived myotube) using the miRNeasy Mini kit (217004, QIAgen) on the QIAcube instrument. RNAs
coming from the part A of the extraction protocol was used for mRNA-seq and RT-qPCR. RNAs coming from the
part B of the extraction protocol was used for miRseq. PartA RNA was quantified on Nanodrop
spectrophotometer (ND-1000, Thermo Fisher Scientific) and purity/quality (RIN ≥ 7) was assessed on the 2200
TapeStation using the Agilent RNA ScreenTape (5067-5576 / 5067-5577 / 5067-5578, Agilent). PartB RNA was
quantified and purity/quality was assessed on the 2100 Agilent BioAnlayzer using the Agilent small RNA kit
(5067-1548, Agilent).

Reverse transcription – 500 ng of total RNA were reverse transcribed with random primers (48190–011,
Thermo Fisher Scientific), oligo(dT) (SO131, Thermo Fisher Scientific), and deoxynucleotide (dNTP, 10297–018,
Thermo Fisher Scientific) using Superscript® III reverse transcriptase (18080–044, Thermo Fisher Scientific).
Thermocycling conditions were 10 min, 25 °C; 60 min, 55 °C; and 15 min, 75 °C.

qPCR – We amplified cDNA/total DNA using primers (Thermo Fisher Scientific) listed in Table S4. They were
designed using Primer blast (http://www.ncbi.nlm.nih.gov/tools/primer-blast). The amplification efficiency of

each primer set was preliminarily determined by running a standard curve. Detection was performed using a
QuantStudio™ 12K Flex Real-Time PCR System (Thermo Fisher Scientific). Reactions were carried out in a 384-
well plate, with 10 µL containing 2.5 µL of 1/10 cDNA or 6.25 ng/uL total DNA, 0.2 µL of mixed forward and
reverse primers at 10 µM each, and 5 µL of 2X Luminaris Color HiGreen qPCR Master Mix Low Rox (K0973,
Thermo Fisher Scientific). Thermocycling conditions were 50 °C during 2 min, 95 °C during 10 min, followed by
45 cycles including 15 sec at 95 °C, 1 min at 60 °C plus a dissociation stage. All samples were measured in
triplicate. Experiments were normalised using UBC as reference gene and relative quantification was done
with the ΔΔCt method.

mRNA-seq – Libraries are prepared with TruSeq Stranded mRNA kit protocol according supplier
recommendations. Briefly, the key stages of this protocol are successively, the purification of PolyA containing
mRNA molecules using poly-T oligo attached magnetic beads from 1µg total RNA, a fragmentation using
divalent cations under elevated temperature to obtain approximately 300bp pieces, double strand cDNA
synthesis and finally Illumina adapter ligation and cDNA library amplification by PCR for sequencing.
Sequencing is then carried out on paired-end 100 b/75 b of Illumina HiSeq 4000.
An RNA-seq analysis workflow was designed using snakemake 3.5.4 (165) for read quality estimation, mapping
and differential expression analysis. Quality estimation was obtained with FastQC 0.11.5
(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Mapping to the human genome assembly
Ensembl GRCh37.87 (43,695 transcripts) was performed with STAR 2.5.0a (166). According to STAR manual and
for more sensitive novel junction discovery, the junctions detected in a first round of mapping were used in a
second mapping round. Read strandness was confirmed using RSeQC (167). Expression counts at the gene
level were calculated using FeatureCounts 1.4.6 (168). Analysis results were summarised using MultiQC 1.0
(169). Normalised counts (median ratio normalisation, MRN) and differential expression analysis was
performed with DESeq2 1.16.1 (170), considering pairwise comparisons with all developmental stages and
comparing DMD versus healthy cells within developmental stages. BiomaRt 2.30.0 (171) was used to fetch
gene annotations from Ensembl. Transcripts with |log2FoldChange| ≥ 0.4 (equivalent of DMD/healthy ratio ≤
0.76 or ≥ 1.32) and adjusted pvalue ≤ 0.05 were considered differentially expressed. Data are going to be
submitted to GEO.
miRNA-seq – 10 ng of miRNA was reverse transcribed using the Ion Total RNA-seq kit v2
(4475936, Thermofisher Scientific) following the protocol of the manufacturer for small RNA libraries. The
cDNA libraries were amplified and barcoded using Ion Total RNA-seq kit v2 and Ion Xpress RNA-seq Barcode
Adapters 1-16 Kit (Thermofisher Scientific). The amplicons were quantified using Agilent High Sensitivity DNA
kit before the samples were pooled in sets of fifteen. Emulsion PCR and Enrichment was performed on the Ion
OT2 system Instrument using the Ion PI Hi-Q OT2 200 kit (A26434, Thermofisher Scientific). Samples were
loaded on an Ion PI v3 Chip and sequenced on the Ion Proton System using Ion PI Hi-Q sequencing 200 kit
chemistry (200 bp read length; A26433, Thermofisher Scientific). Sequencing reads were trimmed with Prinseq
(172) (v0.20.4) (--trim-right 20) and filtered by average quality score (--trim-qual 20). Reads with a size less
than 15 bp have been removed and reads with a size greater than 100 bp have been trimmed with Cutadapt
(v1.16)(173). Mapping to the human genome assembly Ensembl GRCh37.87 (3111 transcripts) was performed
with STAR 2.5.3a (166)**.** Normalised counts (median ratio normalisation, MRN) and differential expression
analysis was performed with DESeq2 1.16.1 (170), considering pairwise comparisons with all developmental
stages and comparing DMD versus healthy cells within developmental stages. Transcripts with
|log2FoldChange| ≥ 0.4 (equivalent of DMD/healthy ratio ≤ 0.76 or ≥ 1.32) and pvalue ≤ 0.05 were considered
differentially expressed. The use of pvalue instead of adjusted pvalue is justified by biological meaning(174)
(i.e. well-known regulated / dysregulated miRNAs had a pvalue ≤ 0.05 but not an adjusted pvalue ≤ 0.05). Data
are going to be submitted to GEO.

High-throughput data analyses – Graphs were realised using RStudio. Viridis library (175) was used for the
colour palette easier to read with colour blindness and print well in grey scale. For unsupervised analyses,
normalised counts were standardised with scale function (center = TRUE, scale = TRUE) and plotted with
corrplot function from corrplot library (176). Spearman correlation was done with the cor function (method =
"spearman", use = "pairwise.complete.obs") on standardised data. Hierarchical clustering and heatmap were
performed with gplots library (177) heatmap.2 function on standardised data. Gene enrichment data were
retrieved from DAVID database using RDAVIDWebService library (178) on supervised list of mRNAs (mRNA-seq
data: adjusted pvalue ≤ 0.05, ratio ≤ 0.76 or ≥ 1.32, normalised counts ≥ 10 in at least one sample; enrichment
data: Benjamini value ≤ 0.05, enrichment ≥ 1.5). Only Gene Ontology terms were processed.

Protein experiments

Immunolabelling – Cells (healthy hiPSC 1/ DMD hiPSC 2, Table S3) at day 17 of culture were thawed and
seeded at 10,000 cells/cm^2 in SKM02 medium in Falcon® 96-well microplate (353219, Corning) coated with
0.1% gelatin (G1393, Sigma-Aldrich) and 2.4 µg/mL laminin (23017015, Thermofischer Scientific) in PBS 1X
(D8537, Sigma-Aldrich). After 4 days, cells were switched to DMEM/F-12, HEPES (31330038, Thermofischer
Scientific) with 2% Horse serum (H1270, Sigma-Aldrich). Before staining, after removing the culture medium,
cells were fixed 15 min at 4°C with PFA 4% (15710, Euromedex) after 7 days of culture. A first quick Phosphate
buffered saline (PBS) 1X tablets (P4417, Sigma-Aldrich) wash was done, followed by another lasting 10 min.
Then, a solution with PBS 1X, Triton™ X-100 0.25% (T8787, Sigma-Aldrich) and Bovine serum albumin 2.5%
(BSA, A9418, Sigma-Aldrich) was added and incubated 30 min at room temperature. Primary antibody was
finally added, diluted in the same buffer (α-actinin 1/500, A7811, Sigma-Aldrich), overnight at 4°C. The next
day, two quick PBS 1X washes were followed by a third incubated 10 min at room temperature. An incubation
was done 45 min at room temperature with a mix of 4′,6-Diamidine-2′-phenylindole dihydrochloride (DAPI,
1µg/mL, 10236276001, Sigma-Aldrich) and the secondary antibody Donkey anti-Mouse Alexa Fluor 555 in PBS
1X, (1/1000, A-31570, Thermofischer Scientific). Finally, two quick PBS 1X washes were followed by a third
incubated 10 min at room temperature. The stained cells were kept in PBS 1X at 4°C before imaging with a
Zeiss LSM880 Airyscan confocal and Zen software (Black edition).

Western blotting – For tissue-derived myotubes, after three rinses with cold PBS 1X (w/o Ca2+ and Mg2+,
D8537, Sigma-Aldrich), protein extracts were isolated from cultured cells by scraping (010154, Dutscher) with
an extraction protein buffer (NaCl 150 mM, Tris 50 mM, EDTA 10 mM (AM9260G, ThermoFisher Scientific),
Triton 1X, 1/100 Protease Inhibitor Cocktail (P8340, Sigma-Aldrich), PhosSTOP tablet (04906845001, Roche
Diagnostics)). For hiPSC-derived myotubes, cell pellets were rinsed once with cold PBS 1X, spun 5 min at 300 g
and resuspended in the same extraction protein buffer. Protein Extracts were centrifuged at 4°C 10 min at
16,000 g and supernatants were kept at −80 °C. Quantitation of total protein was done with Pierce BCA protein
assay kit (23225, ThermoFischer Scientific). Before gel loading, protein extracts were mixed with 9µL of loading
buffer (Urea 4M, SDS 3.8%, Glycerol 20%, Tris 75mM pH 6.8, 5% β-mercaptoethanol, 0.1mg/mL Bromophenol
blue) and completed to 28µL (for one well) with extraction protein buffer, then heated once 5 min at 95 °C.
Western blots were performed either with Criterion ™ XT Tris-Acetate Precast Gels 3–8 % (3450130, Bio-Rad,
Hercules, CA), XT Tricine running buffer (161–0790, Bio-Rad) and ran at room temperature for 1 hour and 15
min at 150 V for RYR1 (1/1000, MA3-925, ThermoFisher Scientific), MF20 (1/500, DSHB, concentrate),
Manex50 (1/30, DSHB), α-sarcoglycane (1/150, A-SARC-L-CE, Leica biosystems), γ-sarcoglycane (1/150, G-
SARC-CE, Leica biosystems), or with 4–15% Criterion™ TGX™ Precast Midi Protein Gel (5671084, Bio-Rad), 10x
Tris/Glycine/SDS Running Buffer (1610772), and ran at room temperature for 1 hour at 200 V for CaV1.1
(1/1000, MA3-920, ThermoFisher Scientific), ATP5A (1/1,000, ab14748, ABCAM), Semaphorin 6A (1/55,
AF1146, R&D systems) and GLI3 (1/200, AF3690, R&D systems). Gels were rinsed once in water and blotted
either with "high molecular weight" or "mixed molecular weight" program of TransBlot® Turbo™ transfer
system (Bio-Rad) using Trans-Blot®Turbo™ Midi Nitrocellulose Transfer Packs (170–4159, Bio-Rad). Blots were
then processed with the SNAP i.d.® 2.0 Protein Detection System following the manufacturer's protocol, with

Odyssey® Blocking Buffer (927-40003, LI-COR) for blocking and with 0,2% Tween® 20 added for antibody
dilutions (28829.296, VWR), washes were done with phosphate-buffered saline tween (PBST) buffer (PBS 1X
tablets, P4417, Sigma-Aldrich; 0.1 % Tween® 20). Every primary antibody was pooled with either α-actinin
(1/12,500, sc-17829, Santa Cruz or 1/7000, A7811, Sigma-Aldrich) or α-tubulin (1/6666, Ab7291, Abcam). For
secondary antibodies, either IRDye 800CW donkey anti-mouse and/or IRDye® 680RD donkey anti-goat were
used (1/5000-1/10000, 926-32212, 926-68074, LI-COR). After completion of SNAP i.d.® general protocol, with
the membrane still in the blot holder, two PBS 1X washes were finally done before band visualisations with
Odyssey® CLx Imaging System and quantification with Image Studio Lite software (Version 5.2). Statistical
analysis was performed using unpaired t test on GraphPad Prism software.

iTRAQ proteomics –
Samples Preparation: Cells at day 17 were collected and resuspended in 90% FBS (Hyclone), 10% DMSO
(A3672.0050, VWR), cooled down until -90°C with the CryoMed™ device (ThermoFisher Scientific), before
storage in liquid nitrogen. Cells were then thawed and washed 5 times with cold PBS and air was replaced by
Argon to thoroughly dry the pellet that was flash frozen in liquid nitrogen. 5-10 times the approximate cell
pellet volume of 0.5 M triethyl ammonium bicarbonate (TEAB) with 0.05% SDS was added to the cell pellet for
protein extraction. Cell pellet was re-suspended and triturated by passing through a 23-gauge needle and 1ml
syringe for 30 times. Samples were then sonicated on ice at amplitude of 20% for 30 x 2 sec bursts and
centrifuged at 16000g for 10 min at 4°C. Supernatant was transferred to a fresh Eppendorf tube. Protein was
quantified by nanodrop. 100-150µg of protein was aliquoted for each individual sample and 2µl TCEP (50mM
tris-2-carboxymethyl phosphine) was added for every 20µl of protein used for reducing the samples. After 1 hr
incubation at 60°C, 1µl MMTS (200mM methylmethane thiosulphonate) was added for every 20µl of protein
used for alkylating/'blocking' the samples. Finally, after a 10 min incubation at RT, samples were trypsinised by
addition of 6-7.5µl of 500ng/µl trypsin. The ration between enzyme: substrate was 1:40. Samples were
incubated overnight at 37°C in the dark. *iTRAQ labelling:* When iTRAQ reagents reached room temperature,
50µl of isopropanol/[acetonitrile] was added to each iTRAQ 8-plex reagent and was incubated at RT for 2 hrs,
in the dark. 8 µl of 5% hydroxylamine was added to neutralise the reaction. Each sample was separately
lyophilised at 45°C. Samples have been stored at -20°C or used immediately.
Offline C4 High Performance Liquid Chromatography (HPLC): All 8 samples were pooled together in 60µl of 97%
mobile phase A (99.92% % H2O, 0.08% NH4OH) and 3% mobile phase B (99.92% % Acetonitrile, 0.02% NH4OH)
by serially reconstituting each sample. Extra 40µl of mobile phase was added to sample 1, after sample has
been well vortexed, all the contents of sample 1 tube were transferred to the tube with the sample 2 (and
serially repeated until all samples were pooled). Final volume of samples needed to be 100µl. After sample was
centrifuged at 13000g for 10 min, supernatant was collected with an HPLC injection syringe. 100µl was injected
onto the sample loop. Fractions were collected in a peak dependent manner. Finally, fractions were lyophilised
at 45°C and stored at -20°C until required. The used column was a Kromasil C4 column 100Å pore size, 3.5µm
particle size, 2.1mm inner diameter and 150mm length. The gradient for C4 separation was (RT in min - %B): 0-
3; 10-3; 11-5; 16-5; 65-20; 100-30; 15-80; 120-80; 125-3.
Solid Phase Extraction Cleaning of peptides fractions: A GracePureTMT SPE C18-Aq cartridge was used for pre-
cleaning of samples (Support: Silica, % Carbon: 12.5%, With endcapping, Surface area: 518m^2/g, Particle size:
50µm, Pore size: 60Å, Water-wettable). Samples were reconstituted using in total 400µl of 1% ACN, 0.01% FA.
Cartridge was washed with 600µl of ACN. ACN was then completely flushed out of the column at dropwise
speed. This activated the ligands. Then 1% ACN, 0.01% FA (600µl) was flushed through the cartridge to
equilibrate the sorbent. 400µl of the sample was loaded in the cartridge. It was then very slowly flushed
through the cartridge and recovered into a fresh tube. This process was repeated 3 times. 2 volumes of 250µl
of 1%ACN, 0.01%FA were used to clean and de-salt the sample. It was flushed through very slowly. 2 volumes
(250µl each) were used per step (2% ACN, 10% ACN, 30% ACN, 50% ACN, 70% ACN). This cycle was repeated
twice. Each particular concentration was pooled in one tube. Samples were dried to dryness in a Speedvac at
RT overnight and stored at -20°C. Like previously, samples were pooled with 100µl of 97% mobile phase A

(99.92% % H2O, 0.08% NH$_4$OH) and 3% mobile phase B (99.92% % Acetonitrile, 0.02% NH$_4$OH) and injected

onto the sample loop. Fractions were collected in a peak dependent manner. The gradient for SPE cleaned

peptides C4 separation (RT in min - %B): 0-2; 10-2; 20-5; 25-5; 35-20; 55-35; 60-35; 70-80; 75-80; 80-3.

Online C18 High Precision Liquid Chromatography (HPLC): 30µl of loading phase (2% acetonitrile, 1.0% formic

acid) was added to each fraction-containing Eppendorf tube. Samples were vortexed and centrifuged. Blanks

(30µl mobile phase) were added into well A1 to A12. 30µl of sample 1 was pipetted into well B1, sample 2 in

well B2 and so on. An orthogonal 2D-LC-MS/MS analysis was performed with the Dionex Ultimate 3000 UHPLC

system coupled with the ultra-high-resolution nano ESI LTQ-Velos Pro Orbitrap Elite mass spectrometer

(Thermo Scientific).

Data analysis: HCD and CID tandem mass spectra were collected and submitted to Sequest search engine

implemented on the Proteome Discoverer software version 1.4 for peptide and protein identifications. All

spectra were searched against the UniProtKB SwissProt. The level of confidence for peptide identifications was

estimated using the Percolator node with decoy database searching. False discovery rate (FDR) was set to 0.05,

and validation was based on the q-Value. Protein ratios were normalised to protein median and peptides with

missing iTRAQ values were rejected from protein quantification. Phosphorylation localisation probability was

estimated with the phosphoRS node. Protein ratios were transformed to log$_2$ ratios and significant changes

were determined by one sample T-test. To reduce the impact of possible false positive identifications, more

parameters were set: 1) only proteins with more than two quantified unique peptides. 2) DMD/Healthy ratio ≥

1.32 or ≤ 0.76 and 3) only FDR corrected p value ≤ 0.05 were retained for bioinformatics analysis.

Acknowledgements

We thank the Fondation Maladies Rares (GenOmics grant), Labex Revive (Investissement d'Avenir; ANR-10-

LABX-73) and the AFM Téléthon for funding this project. The RNA-Sequencing libraries were processed and

sequenced by Integragen (Evry, France). We gratefully acknowledge support from the PSMN (Pôle Scientifique

de Modélisation Numérique) of the ENS de Lyon for the computing resources. We thank Dr Nacira Tabti, Dr

Elisabeth Le Rumeur, Dr Nathalie Deburgrave and Dr Malgorzata Rak for providing us with specific reagents

and antibodies. We thank Dr David Israeli for his feedback on the manuscript and overall discussion on our

project.

Figure 1 – Differentiation dynamics of hiPSCs (D0) into MyoT (D25) in healthy cells at the transcriptomic
levels. A) Spearman correlation matrix of transcriptomes (mRNAs, right) and miRnomes (miRNAs, left). Yellow
dots indicate a stronger correlation. **B)** Heatmap of selected differentiation markers. (D: day; hiPSC: human
induced pluripotent stem cell; MyoT: myotube).

Figure 2 – Differentiation dynamics of hiPSCs (D0) into MyoT (D25) in DMD cells. A) Dotplot of DMD/healthy
expression ratios of selected markers. Statistical differences are indicated in brackets after gene names, and
grey circles around the corresponding dots. **B)** Proportions of significantly dysregulated mRNAs (adjusted
pvalue ≤ 0.05) in DMD cells at each time points. Expression of **C)** hsa-miR-1 and **D)** *ATP2A2* mRNA during
differentiation, as well as **E)** ATP2A2 protein level at D17. (*adjusted pvalue ≤ 0.05, **adjusted pvalue ≤ 0.01,
***adjusted pvalue ≤ 0.001, ****adjusted pvalue ≤ 0.0001; D: day; hiPSC: human induced pluripotent stem
cell; MyoT: myotube).

Figure 3 – Comparison of healthy and DMD MyoT from hiPSCs and tissues at the protein level. A) hiPSC-
derived MyoT immunolabelling of α-actinin (red) and nuclei (DAPI, blue) in healthy (left) and DMD cells (right).
B) Representative Western blots and related quantifications of DMD, SGCA, SGCG, myosin heavy chains,
CACNA1S and RYR1 from protein extracts in healthy and DMD hiPSC-derived and tissue-derived MyoT (X: 0.25
µg of total protein was used in hiPSC-derived MyoT instead of 7µg in tissue-derived MyoT - *pvalue ≤ 0.05,
**pvalue ≤ 0.01, ***pvalue ≤ 0.001, ****pvalue ≤ 0.0001)). (hiPSC: human induced pluripotent stem cell;
MyoT: myotube).

Figure 4 – Manifestation of the DMD phenotype in the transcriptomes and miRnomes of myotubes derived
from hiPSCs and tissues. A) Hierarchical clustering and heatmap in healthy hiPSCs (D0), hiPSC-derived MyoT
and tissue-derived MyoT with selected skeletal muscle transcripts and miRNAs. **B)** Volcano plots of
dysregulated mRNAs/miRNAs in hiPSC-derived MyoT (left) and tissue-derived MyoT (right) – vertical grey
dashed lines represent DMD/Healthy ratio thresholds at 0.76 or 1.32 - the horizontal grey dashed line
represents the adjusted pvalue threshold at 0.05. (DAPC: dystrophin-associated protein complex; hiPSC:
human induced pluripotent stem cell; MyoT: myotube; NMJ: neuromuscular junction; TF: transcription factor).

Figure 5 – Illustration of the fibrosis phenotypes in DMD cells. Volcano plots of dysregulated mRNAs/miRNAs
related to **A)** the SHH pathway and collagen metabolism at D10/17/25; and **B)** fibrosis at D25 – vertical grey
dashed lines represent DMD/Healthy ratio thresholds at 0.76 or 1.32 - the horizontal grey dashed line
represents the adjusted pvalue threshold at 0.05. (D: day; MMP: matrix metallopeptidase; SHH: sonic
hedgehog pathway; TIMP: tissue inhibitor of metallopeptidase; TGF: transforming growth factor).

Figure 6 – Illustration of the metabolic and mitochondrial phenotypes in DMD cells. Volcano plots of
dysregulated mRNAs/miRNAs related to **A)** principal metabolic pathways; and **B)** the constitution of the five
mitochondrial respiratory complexes in DMD hiPSC-derived MyoT – vertical grey dashed lines represent
DMD/Healthy ratio thresholds at 0.76 or 1.32 - the horizontal grey dashed line represents the adjusted pvalue
threshold at 0.05. Quantification of ATP5A1 expression **C)** at the mRNA level during differentiation, and **D)** at
the protein level at D17 (iTRAQ data, left) and D25 (Western blot data, right). (*adjusted pvalue ≤ 0.05,
**adjusted pvalue ≤ 0.01, ***adjusted pvalue ≤ 0.001, ****adjusted pvalue ≤ 0.0001). (D: day; hiPSC: human
induced pluripotent stem cell, MyoT: myotube)

Figure 7 – Mitochondrial dysregulations in DMD cells during differentiation. A) Absolute (top) and relative
numbers (%, bottom) of dysregulated genes from the different mitochondrial compartments over the course
of DMD hiPSC differentiation. **B)** Expression ratios of selected mitochondrial proteins. Statistical differences
are indicated in brackets (*adjusted pvalue ≤ 0.05, **adjusted pvalue ≤ 0.01, ***adjusted pvalue ≤ 0.001,
****adjusted pvalue ≤ 0.0001). **C)** Volcano plots of mitochondria-related genes over the course of DMD hiPSC

differentiation. Statistical differences are symbolised with orange dots – vertical grey dashed lines represent
DMD/Healthy ratio thresholds at 0.76 or 1.32 - the horizontal grey dashed line represents the adjusted pvalue
threshold at 0.05 – The percentage of significantly dysregulated genes is indicated at the bottom right in grey.
(D: day).

Figure S1 – DMD variant expression over the course of hiPSC differentiation. A) Bright field microscope
pictures at the 7 differentiation points giving rise to hiPSC-derived and tissue-derived MyoT. Possible
cryopreservation time points are indicated by snowflakes. **B)** RT-qPCR relative quantification of *DMD* variants
expression during differentiation of hiPSCs (D0) into MyoT (D25) with the related cycle threshold (CT) values
(Ct: cycle threshold; D: day; hiPSC: human induced pluripotent stem cell; MyoB: myoblast; MyoT: myotube).

Figure S2 – Gene ontology enrichments over the course of healthy and DMD hiPSC differentiation A)
Proportions of significantly regulated mRNAs (adjusted pvalue ≤ 0.01) between successive differentiation time
points during the differentiation of healthy hiPSCs. Gene ontology enrichments on **B)** significantly regulated
mRNAs between successive differentiation time points in healthy cells (number of genes in brackets) and **C)**
significantly dysregulated mRNAs at each differentiation time points in DMD cells. The number of genes
involved in these significant enrichments is indicated in brackets next to each GO term. In green, GO terms
related to downregulated genes and in yellow, GO terms related to upregulated genes and in yellow (BP:
biological process; CC: cellular component; D: day; hiPSC: human induced pluripotent stem cell; MyoT:
myotube).

Figure S3 – Comparison of healthy and DMD cells at D10 and D17, protein analyses. Western blots and
quantifications of **A)** SEMA6A at D10, **B)** GLI3 at D10 and **C)** GLI3 at D17. (*pvalue ≤ 0.05, **pvalue ≤ 0.01,
***pvalue ≤ 0.001, ****pvalue ≤ 0.0001; D: day; GLI3FL: GLI3 full length; GLI3R: GLI3 repressor).

**Figure S4 – Comparison of hiPSC-derived and tissue-derived MyoT for the expression of cell cycle genes and
myogenic regulators.** Hierarchical clustering and heatmap of **A)** selected cell cycle transcripts and miRNAs, and
B) DLK1, IGF2 and selected myosin transcripts in hiPSCs (D0), hiPSC- and tissue-derived MyoT. **C)** Dotplot of
DMD/healthy expression ratio of muscle transcription factors. Significant statistical differences are shown in
brackets (*adjusted pvalue ≤ 0.05, **adjusted pvalue ≤ 0.01, ***adjusted pvalue ≤ 0.001, ****adjusted pvalue
≤ 0.0001) . (hiPSC: human induced pluripotent stem cell; MyoT: myotube).

**Figure S5 – Dysregulations of metabolic pathways and mitochondrial genes during differentiation of DMD
hiPSCs. A)** Scheme of metabolism dysregulations at day 25. Dotplots of **B)** mitochondrial transcripts, **C)**
transcripts coding mitochondrial protein import, and **D)** transcripts coding mitochondrial
transcription/replication; **E)** Mitochondrial DNA quantification by RT-qPCR at D25. Dotplots of mitochondrial
proteins expressed at D17 involved in **F)** protein import, **G)** mitochondrial transcription/replication. Statistics
are in brackets (*adjusted pvalue ≤ 0.05, **adjusted pvalue ≤ 0.01, ***adjusted pvalue ≤ 0.001, ****adjusted
pvalue ≤ 0.0001; D: day).

References

1. Hoffman EP, Brown RH, Kunkel LM. Dystrophin: the protein product of the Duchenne muscular dystrophy locus. Cell. 1987;51(6):919–28.

2. Koeks Z, Bladen CL, Salgado D, van Zwet E, Pogoryelova O, McMacken G, et al. Clinical Outcomes in Duchenne Muscular Dystrophy: A Study of 5345 Patients from the TREAT-NMD DMD Global Database. J Neuromuscul Dis [Internet]. 2017;4(4):293–306. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29125504%0Ahttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC5701764

3. Liu M, Chino N, Ishihara T. Muscle damage progression in Duchenne muscular dystrophy evaluated by a new quantitative computed tomography method. Arch Phys Med Rehab. 1993;74(5):507–14.

4. Pescatori M, Broccolini A, Minetti C, Bertini E, Bruno C, D'amico A, et al. Gene expression profiling in the early phases of {DMD:} a constant molecular signature characterizes {DMD} muscle from early postnatal life throughout disease progression. {FASEB} J. 2007;21(4):1210–26.

5. Szigyarto C, Spitali P. Biomarkers of Duchenne muscular dystrophy: current findings. Degener Neurol Neuromuscul Dis. 2018;8:1–13.

6. Moat SJ, Bradley DM, Salmon R, Clarke A, Hartley L. Newborn bloodspot screening for Duchenne muscular dystrophy: 21 years experience in Wales {(UK).}. Eur J Hum Genet. 2013;21(10):1049–53.

7. Crone M, Mah JK. Current and Emerging Therapies for Duchenne Muscular Dystrophy. Curr Treat Options Neurol. 2018;20(8):31.

8. Ngoc L-N, Malerba A, Popplewell L, Schnell F, Hanson G, Dickson G. Systemic Antisense Therapeutics for Dystrophin and Myostatin Exon Splice Modulation Improve Muscle Pathology of Adult mdx Mice. Mol Ther - Nucleic Acids. 2017;6:15–28.

9. Nguyen F, Cherel Y, Guigand L, I G-L, Wyers M. Muscle lesions associated with dystrophin deficiency in neonatal golden retriever puppies. J Comp Pathol. 2002;126(2–3):100–8.

10. Bassett DI. Dystrophin is required for the formation of stable muscle attachments in the zebrafish embryo. Development. 2003;130(23):5851–60.

11. Merrick D, Stadler LK, Larner D, Smith J. Muscular dystrophy begins early in embryonic development deriving from stem cell loss and disrupted skeletal muscle formation. Dis Model Mech. 2009;2(7–8):374–88.

12. Emery AE. Muscle histology and creatine kinase levels in the foetus in Duchenne muscular dystrophy. Nature. 1977;266(5601):472–3.

13. Toop J, Emery AE. Muscle histology in fetuses at risk for Duchenne muscular dystrophy. Clin Genet. 1974;5(3):230–3.

14. Vassilopoulos D, Emery AE. Muscle nuclear changes in fetuses at risk for Duchenne muscular dystrophy. J Med Genet. 1977;14(1):13–5.

15. Massouridès E, Polentes J, Mangeot PE, Mournetas V, Nectoux J, Deburgrave N, et al. Dp412e: A novel human embryonic dystrophin isoform induced by BMP4 in early differentiated cells. Skelet Muscle [Internet]. 2015 Dec 14 [cited 2017 Feb 27];5(1):40. Available from: http://www.skeletalmusclejournal.com/content/5/1/40

16. Nesmith AP, Wagner MA, Pasqualini FS, B OB, Pincus MJ, August PR, et al. A human in vitro model of Duchenne muscular dystrophy muscle formation and contractility. J Cell Biol. 2016;215(1):47–56.

17. Shoji E, Sakurai H, Nishino T, Nakahata T, Heike T, Awaya T, et al. Early pathogenesis of Duchenne muscular dystrophy modelled in patient-derived human induced pluripotent stem cells. Sci Rep. 2015;5:12831.

18. Choi IY, Lim HT, Estrellas K, Mula J, Cohen T V., Zhang Y, et al. Concordant but Varied Phenotypes among Duchenne Muscular Dystrophy Patient-Specific Myoblasts Derived using a Human iPSC-Based Model. Cell Rep. 2016;15(10):2301–12.

19. Chal J, Oginuma M, Al Tanoury Z, Gobert B, Sumara O, Hick A, et al. Differentiation of pluripotent stem cells to muscle fiber to model Duchenne muscular dystrophy. Nat Biotechnol. 2015;33(9):962–9.

20. Young CS, Hicks MR, Ermolova N V, Nakano H, Jan M, Younesi S, et al. A Single {CRISPR-Cas9} Deletion Strategy that Targets the Majority of {DMD} Patients Restores Dystrophin Function in {hiPSC-Derived} Muscle Cells. Cell Stem Cell. 2016;18(4):533–40.

21. Hicks MR, Hiserodt J, Paras K, Fujiwara W, Eskin A, Jan M, et al. {ERBB3} and {NGFR} mark a distinct skeletal muscle progenitor cell in human development and {hPSCs.}. Nat Cell Biol. 2018;20(1):46–57.

22. Kodaka Y, Rabu G, Asakura A. Skeletal Muscle Cell Induction from Pluripotent Stem Cells. Stem Cells Int. 2017;2017:1376151.

23. Caron L, Kher D, Lee KL, McKernan R, Dumevska B, Hidalgo A, et al. A Human Pluripotent Stem Cell Model of Facioscapulohumeral Muscular Dystrophy-Affected Skeletal Muscles. Stem Cells Transl Med. 2016;5(9):1145–61.

24. Shelton M, Metz J, Liu J, Carpenedo RL, Demers S-PP, Stanford WL, et al. Derivation and expansion of {PAX7-positive} muscle progenitors from human and mouse embryonic stem cells. Stem Cell Reports. 2014;3(3):516–29.

25. Xi H, Fujiwara W, Gonzalez K, Jan M, Liebscher S, Van Handel B, et al. {In~Vivo} Human Somitogenesis Guides Somite Development from {hPSCs.}. Cell Rep. 2017;18(6):1573–85.

26. Monaco AP, Neve RL, Chris C-F, Bertelson CJ, Kurnit DM, Kunkel LM. Isolation of candidate {cDNAs} for portions of the Duchenne muscular dystrophy gene. Nature. 1986;323(6089):646–50.

27. Byers TJ, Lidov HGW, Kunkel LM. An alternative dystrophin transcript specific to peripheral nerve. Nat Genet. 1993;4(1):ng0593-
77.

28. Lidov HG, Selig S, Kunkel LM. Dp140: a novel 140 {kDa} {CNS} transcript from the dystrophin locus. Hum Mol Genet.
1995;4(3):329–35.

29. Górecki DC, Monaco AP, Derry JM, Walker AP, Barnard EA, Barnard PJ. Expression of four alternative dystrophin transcripts in
brain regions regulated by different promoters. Hum Mol Genet. 1992;1(7):505–10.

30. D'souza VN, Man NT, Morris GE, Karges W, Pillers DAM, Ray PN. A novel dystrophin isoform is required for normal retinal
electrophysiology. Hum Mol Genet. 1995;4(5):837–42.

31. Nudel U, Zuk D, Einat P, Zeelon E, Levy Z, Neuman S, et al. Duchenne muscular dystrophy gene product is not identical in muscle
and brain. Nature. 1989;337(6202):337076a0.

32. Heikinheimo M, Scandrett JM, Wilson DB. Localization of Transcription Factor GATA-4 to Regions of the Mouse Embryo Involved
in Cardiac Development. Dev Biol. 1994;164(2):361–73.

33. Lyons I, Parsons LM, Hartley L, Li R, Andrews JE, Robb L, et al. Myogenic and morphogenetic defects in the heart tubes of murine
embryos lacking the homeo box gene Nkx2-5. Genes Dev. 1995;9(13):1654–66.

34. Pfeffer PL, Gerster T, Lun K, Brand M, Busslinger M. Characterization of three novel members of the zebrafish Pax2/5/8 family:
dependency of Pax5 and Pax8 expression on the Pax2.1 (noi) function. Development [Internet]. 1998;125(16):3063–74. Available
from: http://www.ncbi.nlm.nih.gov/pubmed/9671580

35. So P, Danielian PS. Cloning and expression analysis of a mouse gene related to Drosophila odd-skipped. Mech Dev. 1999;84(1–
2):157–60.

36. Hancock SN, Agulnik SI, Silver LM, Papaioannou VE, Lowe LA, Yamada S, et al. Mapping and expression analysis of the mouse
ortholog of Xenopus Eomesodermin. Mech Dev. 1999;81(1–2):205–8.

37. Chapman DL, Cooper-Morgan A, Harrelson Z, Papaioannou VE. Critical role for Tbx6 in mesoderm specification in the mouse
embryo. Mech Dev. 2003;120(7):837–47.

38. Lowe LA, Yamada S, Kuehn MR. Genetic dissection of nodal function in patterning the mouse embryo. Development [Internet].
2001;128(10):1831–43. Available from: http://www.ncbi.nlm.nih.gov/pubmed/11311163

39. Hart AH, Hartley L, Sourris K, Stadler ES, Li R, Stanley EG, et al. Mixl1 is required for axial mesendoderm morphogenesis and
patterning in the murine embryo. Development [Internet]. 2002;129(15):3597–608. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/12117810

40. Kwon GS, Fraser ST, Eakin GS, Mangano M, Isern J, Sahr KE, et al. Tg(Afp-GFP) expression marks primitive and definitive
endoderm lineages during mouse development. Dev Dyn. 2006;235(9):2549–58.

41. Kanai-Azuma M, Kanai Y, Gad JM, Tajima Y, Taya C, Kurohmaru M, et al. Depletion of definitive gut endoderm in Sox17-null
mutant mice. Development [Internet]. 2002;129(10):2367–79. Available from: http://www.ncbi.nlm.nih.gov/pubmed/11973269

42. Zhang X, Huang CT, Chen J, Pankratz MT, Xi J, Li J, et al. Pax6 is a human neuroectoderm cell fate determinant. Cell Stem Cell.
2010;7(1):90–100.

43. Rex M, Orme A, Uwanogho D, Tointon K, Wigmore PM, Sharpe PT, et al. Dynamic expression of chicken Sox2 and Sox3 genes in
ectoderm induced to form neural tissue. Dev Dyn. 1997;209(3):323–32.

44. Machon O, Masek J, Machonova O, Krauss S, Kozmik Z. Meis2 is essential for cranial and cardiac neural crest development. BMC
Dev Biol. 2015;15(1):40.

45. Laing NG, Dye DE, Wallgren-Pettersson C, Richard G, Monnier N, Lillis S, et al. Mutations and polymorphisms of the skeletal
muscle α-actin gene (ACTA1). Hum Mutat. 2009;30(9):1267–77.

46. Kardon G, Heanue TA, Tabin CJ. Pax3 and Dach2 positive regulation in the developing somite. Dev Dyn. 2002;224(3):350–5.

47. Pereira FA, Yuhong Q, Zhou G, Tsai MJ, Tsai SY. The orphan nuclear receptor COUP-TFII is required for angiogenesis and heart
development. Genes Dev. 1999;13(8):1037–49.

48. Mitsiadis TA, Salmivirta M, Muramatsu T, Muramatsu H, Rauvala H, Lehtonen E, et al. Expression of the heparin-binding
cytokines, midkine (MK) and HB-GAM (pleiotrophin) is associated with epithelial-mesenchymal interactions during fetal
development and organogenesis. Development [Internet]. 1995;121(1):37–51. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/7867507

49. Yang XM, Vogan K, Gros P, Park M. Expression of the met receptor tyrosine kinase in muscle progenitor cells in somites and limbs
is absent in Splotch mice. Development [Internet]. 1996;122(7):2163–71. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/8681797

50. Sasaki H, Ferguson-Smith AC, Shum AS, Barton SC, Surani MA. Temporal and spatial regulation of H19 imprinting in normal and
uniparental mouse embryos. Development [Internet]. 1995;121(12):4195–202. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/8575319

51. Borycki AG, Mendham L, Emerson CP. Control of somite patterning by Sonic hedgehog and its downstream signal response
genes. Development [Internet]. 1998;125(4):777–90. Available from: http://www.ncbi.nlm.nih.gov/pubmed/9435297

52. Lee CS, Buttitta L, Fan C-M. Evidence that the {WNT-inducible} growth arrest-specific gene 1 encodes an antagonist of sonic

hedgehog signaling in the somite. Proc Natl Acad Sci. 2001;98(20):11347–52.

53.   McMahon AR, Merzdorf CS. Expression of the zic1, zic2, zic3, and zic4 genes in early chick embryos. BMC Res Notes.
2010;3(1):167.

54.   Bladt F, Riethmacher D, Isenmann S, Aguzzi A, Birchmeier C. Essential role for the c-met receptor in the migration of myogenic
precursor cells into the limb bud. Nature. 1995;376(6543):768–71.

55.   Swartz ME, Eberhart J, Pasquale EB, Krull CE. EphA4/ephrin-A5 interactions in muscle precursor cell migration in the avian
forelimb. Development [Internet]. 2001;128(23):4669–80. Available from: http://www.ncbi.nlm.nih.gov/pubmed/11731448

56.   Schäfer K, Braun T. Early specification of limb muscle precursor cells by the homeobox gene Lbx1h. Nat Genet.
1999;23(2):ng1099_213.

57.   Jeong Kyo Yoon, Wold B. The bHLH regulator pMesogenin1 is required for maturation and segmentation of paraxial mesoderm.
Genes Dev. 2000;14(24):3204–14.

58.   Crossley PH, Minowada G, MacArthur CA, Martin GR. Roles for FGF8 in the induction, initiation, and maintenance of chick limb
development. Cell. 1996;84(1):127–36.

59.   Brugmann SA. Six1 promotes a placodal fate within the lateral neurogenic ectoderm by functioning as both a transcriptional
activator and repressor. Development. 2004;131(23):5871–81.

60.   Stewart RA, Arduini BL, Berghmans S, George RE, Kanki JP, Henion PD, et al. Zebrafish foxd3 is selectively required for neural
crest specification, migration and survival. Dev Biol. 2006;292(1):174–88.

61.   Deutsch U, Dressler GR, Gruss P. Pax 1, a member of a paired box homologous murine gene family, is expressed in segmented
structures during development. Cell. 1988;53(4):617–25.

62.   Neubüser A, Koseki H, Balling R. Characterization and developmental expression of Pax9, a paired-box-containing gene related to
Pax1. Dev Biol. 1995;170(2):701–16.

63.   Buchner G, Broccoli V, Bulfone A, Orfanelli U, Gattuso C, Ballabio A, et al. {MAEG,} an {EGF-repeat} containing gene, is a new
marker associated with dermatome specification and morphogenesis of its derivatives. Mech Dev. 2000;98(1–2):179–82.

64.   Xu X-M, Fisher DA, Zhou L, White FA, Ng S, Snider WD, et al. The Transmembrane Protein Semaphorin {6A} Repels Embryonic
Sympathetic Axons. J Neurosci. 2000;20(7):2638–48.

65.   Davis RL, Weintraub H, Lassar AB. Expression of a single transfected cDNA converts fibroblasts to myoblasts. Cell.
1987;51(6):987–1000.

66.   Donalies M, Cramer M, Ringwald M, A S-P. Expression of M-cadherin, a member of the cadherin multigene family, correlates
with differentiation of skeletal muscle cells. Proc Natl Acad Sci. 1991;88(18):8024–8.

67.   Gahlmann R, Kedes L. Cloning, structural analysis, and expression of the human fast twitch skeletal muscle troponin C gene. J
Biol Chem. 1990;265(21):12520–8.

68.   Roberds SL, Anderson RD, Ibraghimov-Beskrovnaya O, Campbell KP. Primary structure and muscle-specific expression of the 50-
kDa dystrophin-associated glycoprotein (adhalin). J Biol Chem [Internet]. 1993;268(32):23739–42. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/8226900

69.   MacKenzie AE, Korneluk RG, Zorzato F, Fujii J, Phillips M, Iles D, et al. The human ryanodine receptor gene: its mapping to
19q13.1, placement in a chromosome 19 linkage group, and exclusion as the gene causing myotonic dystrophy. Am J Hum Genet
[Internet]. 1990;46(6):1082–9. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/1971150%0Ahttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC1683814

70.   Greco S, De Simone M, Colussi C, Zaccagnini G, Fasanaro P, Pescatori M, et al. Common {micro-RNA} signature in skeletal muscle
damage and regeneration induced by Duchenne muscular dystrophy and acute ischemia. {FASEB} J. 2009;23(10):3335–46.

71.   Milet C, Duprez D. The Mkx homeoprotein promotes tenogenesis in stem cells and improves tendon repair. Ann Transl Med.
2015;3(Suppl 1):S33.

72.   Brent AE, Schweitzer R, Tabin CJ. A somitic compartment of tendon progenitors. Cell. 2003;113(2):235–48.

73.   Lefebvre V, Li P, De Crombrugghe B. A new long form of Sox5 (L-Sox5), Sox6 and Sox9 are coexpressed in chondrogenesis and
cooperatively activate the type II collagen gene. EMBO J. 1998;17(19):5718–33.

74.   Gao Y, Jheon A, Nourkeyhani H, Kobayashi H, Ganss B. Molecular cloning, structure, expression, and chromosomal localization of
the human Osterix (SP7) gene. Gene. 2004;341(1–2):101–10.

75.   NODA M, DENHARDT DT. Regulation of Osteopontin Gene Expression in Osteoblasts. Ann N Y Acad Sci. 1995;760(1):242–8.

76.   Kobayashi A, Valerius MT, Mugford JW, Carroll TJ, Self M, Oliver G, et al. Six2 Defines and Regulates a Multipotent Self-Renewing
Nephron Progenitor Population throughout Mammalian Kidney Development. Cell Stem Cell. 2008;3(2):169–81.

77.   Atala A. Re: Sall1 Maintains Nephron Progenitors and Nascent Nephrons by Acting as Both an Activator and a Repressor: Editorial
Comment. J Urol. 2015;194(2):592–3.

78.   Lytton J, DH M. Molecular cloning of {cDNAs} from human kidney coding for two alternatively spliced products of the cardiac
{Ca2+-ATPase} gene. J Biol Chem. 1988;263(29):15024–31.

79.   Yoshida M, Ozawa E. Glycoprotein complex anchoring dystrophin to sarcolemma. J Biochem. 1990;108(5):748–52.

80.   Chen JF, Mandel EM, Thomson JM, Wu Q, Callis TE, Hammond SM, et al. The role of microRNA-1 and microRNA-133 in skeletal
850   muscle proliferation and differentiation. Nat Genet. 2006;38(2):228–33.

851   81.   Hak KK, Yong SL, Sivaprasad U, Malhotra A, Dutta A. Muscle-specific microRNA miR-206 promotes muscle differentiation. J Cell
852   Biol. 2006;174(5):677–87.

853   82.   Hasty P, Bradley A, Morris JH, Edmondson DG, Venuti JM, Olson EN, et al. Muscle deficiency and neonatal death in mice with a
854   targeted mutation in the myogenin gene. Nature. 1993;364(6437):364501a0.

855   83.   Vignier N, Moghadaszadeh B, Gary F, Beckmann J, Mayer U, Guicheney P. Structure, genetic localization, and identification of the
856   cardiac and skeletal muscle transcripts of the human integrin α7 gene (ITGA7). Biochem Biophys Res Commun. 1999;260(2):357–
857   64.

858   84.   Newey SE, Howman E V., Ponting CP, Benson MA, Nawrotzki R, Loh NY, et al. Syncoilin, a Novel Member of the Intermediate
859   Filament Superfamily That Interacts with α-Dystrobrevin in Skeletal Muscle. J Biol Chem. 2001;276(9):6645–55.

860   85.   WU Q-L, JHA PK, RAYCHOWDHURY MK, DU Y, LEAVIS PC, SARKAR S. Isolation and Characterization of Human Fast Skeletal β
861   Troponin T cDNA: Comparative Sequence Analysis of Isoforms and Insight into the Evolution of Members of a Multigene Family.
862   DNA Cell Biol. 2009;13(3):217–33.

863   86.   Bernick EP, Zhang PJ, Du S. Knockdown and overexpression of Unc-45b result in defective myofibril organization in skeletal
864   muscles of zebrafish embryos. BMC Cell Biol. 2010;11(1):70.

865   87.   Li H, Randall WR, Du S-J. skNAC (skeletal Naca), a muscle-specific isoform of Naca (nascent polypeptide-associated complex
866   alpha), is required for myofibril organization. FASEB J. 2009;23(6):1988–2000.

867   88.   DeChiara TM, Bowen DC, Valenzuela DM, Simmons M V., Poueymirou WT, Thomas S, et al. The receptor tyrosine kinase MuSK is
868   required for neuromuscular junction formation in vivo. Cell. 1996;85(4):501–12.

869   89.   Okada K, Inoue A, Okada M, Murata Y, Kakuta S, Jigami T, et al. The Muscle Protein Dok-7 Is Essential for Neuromuscular
870   Synaptogenesis. Science (80-). 2006;312(5781):1802–5.

871   90.   Ilene K-M, Travis M, Blau H, Leinwand LA. Expression and {DNA} sequence analysis of a human embryonic skeletal muscle mvosin
872   heavy chain gene. Nucleic Acids Res. 1989;17(15):6167–79.

873   91.   Weiss A, Schiaffino S, Leinwand LA. Comparative sequence analysis of the complete human sarcomeric myosin heavy chain
874   family: implications for functional {diversity11Edited} by J. Karn. J Mol Biol. 1999;290(1):61–75.

875   92.   Strohman RC, J M-E, Glass CA, Matsuda R. Human fetal muscle and cultured myotubes derived from it contain a fetal-specific
876   myosin light chain. Science (80-). 1983;221(4614):955–7.

877   93.   Collins C, Hayden MR, Schappert K. The genomic organization of a novel regulatory myosin light chain gene (MYL5) that maps to
878   chromosome 4p16.3 and shows different patterns of expression between primates. Hum Mol Genet. 1992;1(9):727–33.

879   94.   Type IIx myosin heavy chain transcripts are expressed in type IIb fibers of human skeletal muscle. Am J Physiol - Cell Physiol.
880   1994;267(6 36-6):C1723-8.

881   95.   Rotwein P, Pollock KM, Watson M, Milbrandt JD. Insulin-like growth factor gene expression during rat embryonic development.
882   Endocrinology. 1987;121(6):2141–4.

883   96.   Andersen DC, Laborda J, Baladron V, Kassem M, Sheikh SP, Jensen CH. Dual role of delta-like 1 homolog (DLK1) in skeletal muscle
884   development and adult muscle regeneration. Development. 2013;140(18):3743–53.

885   97.   Noguchi S, EM M, Othmane BK, Hagiwara Y, Mizuno Y, Yoshida M, et al. Mutations in the dystrophin-associated protein gamma-
886   sarcoglycan in chromosome 13 muscular dystrophy. Sci New York N Y. 1995;270(5237):819–22.

887   98.   Campbell KP, Leung AT, Sharp AH. The biochemistry and molecular biology of the dihydropyridine-sensitive calcium channel.
888   Trends Neurosci. 1988;11(10):425–30.

889   99.   A fourth human MEF2 transcription factor, hMEF2D, is an early marker of the myogenic lineage. Development.
890   1993;118(4):1095–106.

891   100.   Gautam M, Noakes PG, Mudd J, Nichol M, Chu GC, Sanes JR, et al. Failure of postsynaptic specialization to develop at
892   neuromuscular junctions of rapsyn-deficient mice. Nature. 1995;377(6546):377232a0.

893   101.   Dawson DM, Eppenberger HM, Eppenberger ME. Multiple Molecular Forms of Creatine Kinases. Ann N Y Acad Sci.
894   1968;151(1):616–26.

895   102.   van Rooij E, Sutherland LB, Qi X, Richardson JA, Hill J, Olson EN. Control of {Stress-Dependent} Cardiac Growth and Gene
896   Expression by a {MicroRNA}. Science (80-). 2007;316(5824):575–9.

897   103.   Hailstones D, Barton P, P C-T, Sasse S, Sutherland C, Hardeman E, et al. Differential regulation of the atrial isoforms of the myosin
898   light chains during striated muscle development. J Biol Chem. 1992;267(32):23295–300.

899   104.   Marx SO, Reiken S, Hisamatsu Y, Jayaraman T, Burkhoff D, Rosemblit N, et al. PKA Phosphorylation Dissociates FKBP12.6 from
900   the Calcium Release Channel (Ryanodine Receptor). Cell. 2004;101(4):365–76.

901   105.   Kuro-o M, Nagai R, Tsuchimochi H, Katoh H, Yazaki Y, Ohkubo A, et al. Developmentally regulated expression of vascular smooth
902   muscle myosin heavy chain isoforms. J Biol Chem. 1989;264(31):18272–5.

106.   Gimona M, Herzog M, Vandekerckhove J, Small JV. Smooth muscle specific expression of calponin. FEBS Lett. 1990;274(1–2):159–62.

107.   Eglen RM, Reddy H, Watson N, Challiss RAJ. Muscarinic acetylcholine receptor subtypes in smooth muscle. Trends Pharmacol Sci. 1994;15(4):114–9.

108.   Ajima R, Akazawa H, Kodama M, Takeshita F, Otsuka A, Kohno T, et al. Deficiency of Myo18B in mice results in embryonic lethality with cardiac myofibrillar aberrations. Genes to Cells. 2008;13(10):987–99.

109.   Heidmann O, Buonanno A, Geoffroy B, Robert B, Guenet JL, Merlie JP, et al. Chromosomal localization of muscle nicotinic acetylcholine receptor genes in the mouse. Science (80-). 1986;234(4778):866–8.

110.   Ueno T, Tanaka K, Kaneko K, Taga Y, Sata T, Irie S, et al. Enhancement of procollagen biosynthesis by p180 through augmented ribosome association on the endoplasmic reticulum in response to stimulated secretion. J Biol Chem. 2010;285(39):29941–50.

111.   Hautala T, Byers MG, Eddy RL, Shows TB, Kivirikko KI, Myllyla R. Cloning of human lysyl hydroxylase: Complete cDNA-derived amino acid sequence and assignment of the gene (PLOD) to chromosome 1p36.3→p36.2. Genomics. 1992;13(1):62–9.

112.   Valtavaara M, Papponen H, Pirttilä AM, Hiltunen K, Helander H, Myllylä R. Cloning and characterization of a novel human lysyl hydroxylase isoform highly expressed in pancreas and muscle. J Biol Chem. 1997;272(11):6831–4.

113.   Martens JHA, Verlaan M, Kalkhoven E, Zantema A. Cascade of Distinct Histone Modifications during Collagenase Gene Activation. Mol Cell Biol. 2003;23(5):1808–16.

114.   Long DA, Price KL, Ioffe E, Gannon CM, Gnudi L, White KE, et al. Angiopoietin-1 therapy enhances fibrosis and inflammation following folic acid-induced acute renal injury. Kidney Int. 2008;74(3):300–9.

115.   Lipson KE, Wong C, Teng Y, Spong S. CTGF is a central mediator of tissue remodeling and fibrosis and its inhibition can reverse the process of fibrosis. Fibrogenesis Tissue Repair [Internet]. 2012;5(S1):S24. Available from: https://fibrogenesis.biomedcentral.com/articles/10.1186/1755-1536-5-S1-S24

116.   Fragiadaki M, Witherden AS, Kaneko T, Sonnylal S, Pusey CD, George B-G, et al. Interstitial fibrosis is associated with increased {COL1A2} transcription in {AA-injured} renal tubular epithelial cells in vivo. Matrix Biol. 2011;30(7–8):396–403.

117.   Hemmann S, Graf J, Roderfeld M, Roeb E. Expression of {MMPs} and {TIMPs} in liver fibrosis – a systematic review with special emphasis on anti-fibrotic strategies. J Hepatol. 2007;46(5):955–75.

118.   Duisters RF, Tijsen AJ, Schroen B, Leenders JJ, Lentink V, Van Der Made I, et al. MiR-133 and miR-30 Regulate connective tissue growth factor: Implications for a role of micrornas in myocardial matrix remodeling. Circ Res. 2009;104(2):170–8.

119.   Vidal B, Serrano AL, Tjwa M, Suelves M, Ardite E, Mori R, et al. Fibrinogen drives dystrophic muscle fibrosis via a {TGFβ/alternative} macrophage activation pathway. Gene Dev. 2008;22(13):1747–52.

120.   Timpani CA, Hayes A, Rybalka E. Revisiting the dystrophin-ATP connection: How half a century of research still implicates mitochondrial dysfunction in Duchenne Muscular Dystrophy aetiology. Med Hypotheses. 2015;85(6):1021–33.

121.   Šileikyte J, Blachly-Dyson E, Sewell R, Carpi A, Menabò R, Di Lisa F, et al. Regulation of the mitochondrial permeability transition pore by the outer membrane does not involve the peripheral benzodiazepine receptor (translocator protein of 18 kDa (TSPO)). J Biol Chem. 2014;289(20):13769–81.

122.   Emery AE, Burt D. Intracellular calcium and pathogenesis and antenatal diagnosis of Duchenne muscular dystrophy. Br Med J. 1980;280(6211):355–7.

123.   Shkryl VM, Martins AS, Ullrich ND, Nowycky MC, Niggli E, Shirokova N. Reciprocal amplification of ROS and Ca2+ signals in stressed mdx dystrophic skeletal muscle fibers. Pflugers Arch Eur J Physiol. 2009;458(5):915–28.

124.   Whitehead NP, Yeung EW, Froehner SC, Allen DG. Skeletal muscle NADPH oxidase is increased and triggers stretch-induced damage in the mdx mouse. PLoS One. 2010;5(12):e15354.

125.   Rodriguez MC, Tarnopolsky MA. Patients with dystrophinopathy show evidence of increased oxidative stress. Free Radic Biol Med. 2003;34(9):1217–20.

126.   Scholte HR, Busch HFM. Early changes of muscle mitochondria in duchenne dystrophy Partition and activity of mitochondrial enzymes in fractionated muscle of unaffected boys and adults and patients. J Neurol Sci. 1980;45(2–3):217–34.

127.   Sharma U, Atri S, Sharma MC, Sarkar C, Jagannathan NR. Skeletal muscle metabolism in Duchenne muscular dystrophy {(DMD):} an in-vitro proton {NMR} spectroscopy study. Magn Reson Imaging. 2003;21(2):145–53.

128.   Lemos DR, Babaeijandaghi F, Low M, Chang C-K, Lee ST, Fiore D, et al. Nilotinib reduces muscle fibrosis in chronic muscle injury by promoting {TNF-mediated} apoptosis of fibro/adipogenic progenitors. Nat Med. 2015;21(7):786–94.

129.   Villalta AS, Nguyen HX, Deng B, Gotoh T, Tidball JG. Shifts in macrophage phenotypes and macrophage competition for arginine metabolism affect the severity of muscle pathology in muscular dystrophy. Hum Mol Genet. 2009;18(3):482–96.

130.   Desguerre I, Mayer M, Leturcq F, Barbet J-P, Gherardi RK, Christov C. Endomysial Fibrosis in Duchenne Muscular Dystrophy: A Marker of Poor Outcome Associated With Macrophage Alternative Activation. J Neuropathol Exp Neurol. 2009;68(7):762–73.

131.   Fusako S-T, Narita A, Masuda S, Wakamatsu T, Watanabe N, Nishiyama T, et al. Premyogenic progenitors derived from human pluripotent stem cells expand in floating culture and differentiate into transplantable myogenic progenitors. Sci Rep-uk. 2018;8(1):6555.

132. Matsumura K, Tome FMS, Ionasescu V, Ervasti JM, Anderson RD, Romero NB, et al. Deficiency of dystrophin-associated proteins in Duchenne muscular dystrophy patients lacking COOH-terminal domains of dystrophin. J Clin Invest. 1993;92(2):866–71.

133. Yuasa K, Hagiwara Y, Ando M, Nakamura A, Takeda S, Hijikata T. {MicroRNA-206} is highly expressed in newly formed muscle fibers: implications regarding potential for muscle regeneration and maturation in muscular dystrophy. Cell Struct Funct. 2008;33(2):163–9.

134. Cullen MJ, Fulthorpe JJ. Stages in fibre breakdown in duchenne muscular dystrophy An electron-microscopic study. J Neurol Sci. 1975;24(2):179–200.

135. Brouilly N, Lecroisey C, Martin E, Pierson L, Mariol MC, Mounier N, et al. Ultra-structural time-course study in the C. elegans model for Duchenne muscular dystrophy highlights a crucial role for sarcomere-anchoring structures and sarcolemma integrity in the earliest steps of the muscle degeneration process. Hum Mol Genet. 2015;24(22):6428–45.

136. Consolino CM, Brooks S V. Susceptibility to sarcomere injury induced by single stretches of maximally activated muscles of mdx mice. J Appl Physiol Bethesda Md 1985. 2004;96(2):633–8.

137. Kong J, Anderson JE. Dystrophin is required for organizing large acetylcholine receptor aggregates. Brain Res. 1999;839(2):298–304.

138. Kong J, Yang L, Li Q, Cao J, Yang J, Chen F, et al. The absence of dystrophin rather than muscle degeneration causes acetylcholine receptor cluster defects in dystrophic muscle. Neuroreport. 2012;23(2):82–7.

139. Bell CD, Conen PE. Histopathological changes in Duchenne muscular dystrophy. J Neurol Sci. 1968;7(3):529–44.

140. Haddix SG, il Lee Y, Kornegay JN, Thompson WJ. Cycles of myofiber degeneration and regeneration lead to remodeling of the neuromuscular junction in two mammalian models of Duchenne muscular dystrophy. PLoS One. 2018;13(10):e0205926.

141. Luz MAM, Marques MJ, Neto SH. Impaired regeneration of dystrophin-deficient muscle fibers is caused by exhaustion of myogenic cells. Braz J Med Biol Res. 2002;35(6):691–5.

142. Zhou L, Porter JD, Cheng G, Gong B, Hatala DA, Merriam AP, et al. Temporal and spatial {mRNA} expression patterns of {TGF-beta1,} 2, 3 and {TbetaRI,} {II,} {III} in skeletal muscles of mdx mice. Neuromuscul Disord Nmd. 2005;16(1):32–8.

143. Bernasconi P, Torchiana E, Confalonieri P, Brugnoni R, Barresi R, Mora M, et al. Expression of transforming growth factor-beta 1 in dystrophic patient muscles correlates with fibrosis. Pathogenetic role of a fibrogenic cytokine. J Clin Invest. 1995;96(2):1137–44.

144. Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, et al. Induction of Pluripotent Stem Cells from Adult Human Fibroblasts by Defined Factors. Cell. 2007;131(5):861–72.

145. DREYFUS JC, SCHAPIRA G, SCHAPIRA F. Biochemical study of muscle in progressive muscular dystrophy. J Clin Invest. 1954;33(5):794–7.

146. Ascah A, Khairallah M, Daussin F, Bourcier-Lucas C, Godin R, Allen BG, et al. Stress-induced opening of the permeability transition pore in the dystrophin-deficient heart is attenuated by acute treatment with sildenafil. Am J Physiol Circ Physiol. 2011;300(1):H144–53.

147. Pauly M, Daussin F, Burelle Y, Li T, Godin R, Fauconnier J, et al. AMPK activation stimulates autophagy and ameliorates muscular dystrophy in the mdx mouse diaphragm. Am J Pathol. 2012;181(2):583–92.

148. M PC. {HISTOPATHOLOGICAL} {FEATURES} {OF} {MUSCLE} {IN} {THE} {PRECLINICAL} {STAGES} {OF} {MUSCULAR} {DYSTROPHY}. Brain. 1962;85(1):109–20.

149. Bradley WG, Hudgson P, Larson PF, Papapetropoulos TA, Jenkison M. Structural changes in the early stages of Duchenne muscular dystrophy. J Neurol Neurosurg Psychiatry. 1972;35(4):451.

150. Rau F, Freyermuth F, Fugier C, Villemin JP, Fischer MC, Jost B, et al. Misregulation of miR-1 processing is associated with heart defects in myotonic dystrophy. Nat Struct Mol Biol. 2011;18(7):840–5.

151. Singh A, Happel C, Manna SK, George A-M, Carrerero J, Kumar S, et al. Transcription factor {NRF2} regulates {miR-1} and {miR-206} to drive tumorigenesis. J Clin Invest. 2013;123(7):2921–34.

152. Zhang X, Zuo X, Yang B, Li Z, Xue Y, Zhou Y, et al. MicroRNA directly enhances mitochondrial translation during muscle differentiation. Cell. 2014;158(3):607–19.

153. Fry CS, Kirby TJ, Kosmac K, McCarthy JJ, Peterson CA. Myogenic Progenitor Cells Control Extracellular Matrix Production by Fibroblasts during Skeletal Muscle Hypertrophy. Cell Stem Cell. 2017;20(1):56–69.

154. Blau HM, Webster C, Pavlath GK. Defective myoblasts identified in Duchenne muscular dystrophy. Proc Natl Acad Sci. 1983;80(15):4856–60.

155. Bovolenta M, Erriquez D, Valli E, Brioschi S, Scotton C, Neri M, et al. The DMD Locus Harbours Multiple Long Non-Coding RNAs Which Orchestrate and Control Transcription of Muscle Dystrophin mRNA Isoforms. PLoS One. 2012;7(9).

156. Ervasti JM, Campbell KP. A role for the dystrophin-glycoprotein complex as a transmembrane linker between laminin and actin. J Cell Biol. 1993;122(4):809–23.

157. Muntoni F, Melis MA, Ganau A, Dubowitz V. Transcription of the dystrophin gene in normal tissues and in skeletal muscle of a family with X-linked dilated cardiomyopathy. Am J Hum Genet. 1995;56(1):151–7.

1013   158.   Warner LE. Expression of Dp260 in muscle tethers the actin cytoskeleton to the dystrophin-glycoprotein complex and partially
1014   prevents dystrophy. Hum Mol Genet. 2002;11(9):1095–105.

1015   159.   Doorenweerd N, Mahfouz A, Van Putten M, Kaliyaperumal R, T'Hoen PAC, Hendriksen JGM, et al. Timing and localization of
1016   human dystrophin isoform expression provide insights into the cognitive phenotype of Duchenne muscular dystrophy. Sci Rep.
1017   2017;7(1).

1018   160.   Lu-Nguyen N, Ferry A, Schnell FJ, Hanson GJ, Popplewell L, Dickson G, et al. Functional muscle recovery following dystrophin and
1019   myostatin exon splice modulation in aged mdx mice. Hum Mol Genet. 2019;

1020   161.   Peccate C, Mollard A, Le Hir M, Julien L, McClorey G, Jarmin S, et al. Antisense pre-treatment increases gene therapy efficacy in
1021   dystrophic muscles. Hum Mol Genet. 2016;25(16):3555–63.

1022   162.   Morales MG, Gutierrez J, Cabello-Verrugio C, Cabrera D, Lipson KE, Goldschmeding R, et al. Reducing CTGF/CCN2 slows down
1023   mdx muscle dystrophy and improves cell therapy. Hum Mol Genet. 2013;22(24):4938–51.

1024   163.   Gatliff J, Campanella M. {TSPO:} kaleidoscopic {18-kDa} amid biochemical pharmacology, control and targeting of mitochondria.
1025   Biochem J. 2016;473(2):107–21.

1026   164.   Massouridès E, Polentes J, Mangeot P-EP-E, Mournetas V, Nectoux J, Deburgrave N, et al. Dp412e: a novel human embryonic
1027   dystrophin isoform induced by BMP4 in early differentiated cells. Skelet Muscle [Internet]. 2015 Dec 14 [cited 2017 Feb
1028   27];5(1):40. Available from: http://www.skeletalmusclejournal.com/content/5/1/40

1029   165.   Köster J, Rahmann S. Snakemake—a scalable bioinformatics workflow engine. Bioinformatics. 2012;28(19):2520–2.

1030   166.   Dobin A, Davis C a, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics
1031   [Internet]. 2013 Jan 1 [cited 2014 Jul 13];29(1):15–21. Available from:
1032   http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3530905&tool=pmcentrez&rendertype=abstract

1033   167.   Wang L, Wang S, Li W. RSeQC: quality control of RNA-seq experiments. Bioinformatics [Internet]. 2012 Aug 15 [cited 2017 Jul
1034   6];28(16):2184–5. Available from: https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/bts356

1035   168.   Liao Y, Smyth GK, Shi W. FeatureCounts: An efficient general purpose program for assigning sequence reads to genomic features.
1036   Bioinformatics [Internet]. 2014 Apr 1 [cited 2017 Jul 6];30(7):923–30. Available from:
1037   http://www.ncbi.nlm.nih.gov/pubmed/24227677

1038   169.   Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: Summarize analysis results for multiple tools and samples in a single report.
1039   Bioinformatics [Internet]. 2016 Oct 1 [cited 2017 Jul 6];32(19):3047–8. Available from:
1040   https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/btw354

1041   170.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol
1042   [Internet]. 2014;15(12):550. Available from: http://genomebiology.biomedcentral.com/articles/10.1186/s13059-014-0550-8

1043   171.   Durinck S, Spellman PT, Birney E, Huber W. Mapping identifiers for the integration of genomic datasets with the R/ Bioconductor
1044   package biomaRt. Nat Protoc [Internet]. 2009 [cited 2019 Jun 12];4(8):1184–91. Available from:
1045   http://www.ncbi.nlm.nih.gov/pubmed/19617889

1046   172.   Schmieder R, Edwards R. Quality control and preprocessing of metagenomic datasets. Bioinformatics. 2011;27(6):863–4.

1047   173.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. Embnet J. 2011;17(1):10–2.

1048   174.   Amrhein V, Greenland S, Blake M. Scientists rise up against statistical significance. Nature. 2019;567(7748):305–7.

1049   175.   Garnier S. viridis: Default Color Maps from "matplotlib". R package version 0.5.1. 2018; Available from: https://cran.r-
1050   project.org/package=viridis

1051   176.   Simko TW and V. R package "corrplot": Visualization of a Correlation Matrix (Version 0.84). 2017; Available from:
1052   https://github.com/taiyun/corrplot

1053   177.   Gregory R. Warnes, Ben Bolker, Lodewijk Bonebakker, Robert Gentleman, Wolfgang Huber Andy Liaw, Thomas Lumley M,
1054   Maechler, Arni Magnusson, Steffen Moeller MS and BV. gplots: Various R Programming Tools for Plotting Data. R package
1055   version 3.0.1. 2016; Available from: https://cran.r-project.org/package=gplots

1056   178.   Fresno C, Fernández EA. {RDAVIDWebService:} a versatile R interface to {DAVID}. Bioinformatics. 2013;29(21):2810–1.

1057

Figure 1

Figure 5

A Metabolism

B Respiratory complexes

C *ATP5A1* mRNA expression

D ATP5A1 protein expression