1

Small molecule modulation of a redox-sensitive stress granule protein dissolves stress granules with beneficial outcomes for familial amyotrophic lateral sclerosis models

Hiroyuki Uechi[1]   (uechi@mpi-cbg.de)

Sindhuja Sridharan[2,24]   (sindhuja.sridharan@embl.de)

Jik Nijssen[1]   (nijssen@mpi-cbg.de)

Jessica Bilstein[1,25]   (jessica.bellmann@uniklinikum-dresden.de)

Juan M. Iglesias-Artola[1]   (jiglesia@mpi-cbg.de)

Satoshi Kishigami[3]   (satoshi.kishigami@exeter.ox.ac.uk)

Virginia Casablancas-Antras[3]   (virginia.casablancasantras@st-annes.ox.ac.uk)

Ina Poser[4]   (iposer@dewpointx.com)

Eduardo J. Martinez[5]   (emartinez@dewpointx.com)

Edgar Boczek[4]   (eboczek@dewpointtherapeutics.com)

Michael Wagner[4]   (mwagner@dewpointx.com)

Nadine Tomschke[1]   (tomschke@mpi-cbg.de)

António M. de Jesus Domingues[1,26]   (amjdomingues@gmail.com)

Arun Pal[6,27]   (a.pal@hzdr.de)

Thom Doeleman[6]   (T.Doeleman-3@umcutrecht.nl)

Sukhleen Kour[7,8,9]   (suk106@pitt.edu)

Eric Nathaniel Anderson[7,8,9]   (ena15@pitt.edu)

Frank Stein[10]   (frank.stein@embl.de)

Hyun O. Lee[1,28]   (hyunokate.lee@utoronto.ca)

Xiaojie Zhang[1,29]   (zhangxj2@shanghaitech.edu.cn)

Anatol W. Fritsch[1]   (fritsch@mpi-cbg.de)

Marcus Jahnel[1,11,12]   (marcus.jahnel@tu-dresden.de)

Julius Fürsch[13,14]   (juliusfuersch@gmail.com)

Anastasia C. Murthy[15]   (anastasia_murthy@alumni.brown.edu)

Simon Alberti[12]   (simon.alberti@tu-dresden.de)

Marc Bickle[1,30]   (marc.bickle@roche.com)

Nicolas L. Fawzi[16]   (nicolas_fawzi@brown.edu)

André Nadler[1]   (nadler@mpi-cbg.de)

Della C. David[17,31]   (della.david@babraham.ac.uk)

Udai B. Pandey[7,8,9]   (udai@pitt.edu)

Andreas Hermann[6,18,19,20,21]   (Andreas.Hermann@med.uni-rostock.de)

Florian Stengel[13,14]   (florian.stengel@uni-konstanz.de)

Benjamin G. Davis[3,22]   (ben.davis@chem.ox.ac.uk)

Andrew J. Baldwin[3,22]   (andrew.baldwin@chem.ox.ac.uk)

Mikhail M. Savitski[2]   (mikhail.savitski@embl.de)

Anthony A. Hyman[1]   (hyman@mpi-cbg.de)

Richard J. Wheeler[1,23]   (richard.wheeler@ndm.ox.ac.uk)

Affiliations

[1]Max Planck Institute of Molecular Cell Biology and Genetics, Pfotenhauerstraße 108, 01307

Dresden, Germany

[2]Genome Biology Unit, European Molecular Biology Laboratory, Meyerhofstraße 1, 69117

Heidelberg, Germany

[3]Department of Chemistry, University of Oxford, South Parks Road, Oxford OX1 3TA, UK

[4]Dewpoint Therapeutics GmbH, Tatzberg 47, Dresden, Germany

[5]Dewpoint Therapeutics Inc, 451 D Street, Suite 104, Boston, MA 02210, USA

[6]Department of Neurology, Technische Universität Dresden, Dresden, Germany

[7]Division of Child Neurology, Department of Pediatrics, Children's Hospital of Pittsburgh, University of

Pittsburgh Medical Center, Pittsburgh, PA, USA

[8]Department of Human Genetics, University of Pittsburgh Graduate School of Public Health,

Pittsburgh, PA, USA

[9]Department of Neurology, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA

[10]Proteomics Core Facility, European Molecular Biology Laboratory (EMBL), Heidelberg, Germany

[11]Cluster of Excellence Physics of Life, TU Dresden, 01062 Dresden, Germany

[12]Biotechnology Center (BIOTEC), CMCB, TU Dresden, Tatzberg 47/48, 01307 Dresden, Germany

[13]University of Konstanz, Department of Biology, Universitätsstrasse 10, 78457 Konstanz, Germany

[14]Konstanz Research School Chemical Biology, University of Konstanz, Universitätsstrasse 10, 78457

Konstanz, Germany

[15]Graduate Program in Molecular Biology, Cell Biology and Biochemistry, Brown University,

Providence, RI, USA

[16]Department of Molecular Biology, Cell Biology, and Biochemistry, Brown University, Providence, RI,

USA

[17]German Centre for Neurodegenerative Diseases, Otfried-Müller-Straße 23, 72076 Tübingen,

Germany

[18]Center for Regenerative Therapies TU Dresden (CRTD), Technische Universität Dresden, Dresden,

Germany

[19]Center for Transdisciplinary Neurosciences Rostock (CTNR), University Medical Center Rostock,

University of Rostock, 18147 Rostock, Germany

[20]Translational Neurodegeneration Section "Albrecht Kossel", Department of Neurology, University

Medical Center Rostock, 18147 Rostock, Germany

[21]Deutsches Zentrum für Neurodegenerative (DZNE) Rostock/Greifswald, 18147 Rostock, Germany

[22]The Rosalind Franklin Institute, Oxfordshire, OX11 0FA, UK

[23]Peter Medawar Building for Pathogen Research, Nuffield Department of Medicine, University of

Oxford, Oxford OX1 3SY, UK

Present address

[24]Present address: Blizard Institute, Barts and the London School of Medicine and Dentistry, Queen

Mary University of London, London, UK

[25]Present address: Faculty of Medicine, University Hospital Carl Gustav Carus, Technische Universität

Dresden, Dresden, Germany.

[26]Present address: Dewpoint Therapeutics GmbH, Tatzberg 47, Dresden, Germany

[27]Present address: Dresden High Magnetic Field Laboratory (HLD), Helmholtz-Zentrum Dresden-

Rossendorf (HZDR), 01328 Dresden, Germany

[28]Present address: Department of Biochemistry, University of Toronto, Toronto, ON M5G 1M1 Canada

[29]Present address: iHuman Institute, School of Life Science and Technology, ShanghaiTech University,

Shanghai, China

[30]Present address: Institute for Translational Bioengineering, pRED, Roche, Grenzachertrasse 124,

4070, Basel, Switzerland

[31]Present address: Babraham Institute, Cambridge CB22 3AT, UK

Abstract

Neurodegenerative diseases such as amyotrophic lateral sclerosis (ALS) are often associated with mutations in proteins that are associated with stress granules. Stress granules are condensates formed by liquid-liquid phase separation which, when aberrant, can lead to altered condensation behaviours and disease phenotypes. Here, we identified lipoamide, a small molecule which specifically prevents cytoplasmic condensation of stress granule proteins. Thermal proteome profiling showed that lipoamide preferentially stabilises intrinsically disordered domain-containing proteins. These include SRSF1 and SFPQ, stress granule proteins necessary for lipoamide activity. The redox state of SFPQ correlates with its condensate-dissolving behaviour, in concordance with the importance of the dithiolane ring for lipoamide activity. In animals, lipoamide ameliorates aging-associated aggregation of a stress granule reporter, improves neuronal morphology, and recovers motor defects caused by expression of ALS-associated FUS and TDP-43 mutants. In conclusion, lipoamide is a well-tolerated small molecule modulator of stress granule condensation and dissection of its molecular mechanism identified a cellular pathway for redox regulation of stress granule formation.

Introduction

Amyotrophic lateral sclerosis (ALS) is a fatal neurodegenerative disease, primarily affecting motor neurons, with poor prognosis and few options for therapy[1]. Currently three FDA approved drugs are available: riluzole, edaravone, and, recently approved, relyvrio™ (a combination of sodium phenylbutyrate and taurursodiol)[2–4]. However, none blocks disease progression, and thus investigating new therapeutic routes important to overcome ALS. Many mutations associated with familial ALS are found in RNA-binding proteins. Notably, TAR DNA-binding protein 43 (TDP-43) and fused in sarcoma (FUS), with >40 ALS-associated mutations described in each[5–7]. These RNA-binding proteins have large intrinsically disordered regions (IDRs) with low sequence complexity.

TDP-43 and FUS are examples of stress granule proteins which normally localise to the nucleus, where they have crucial functions in gene expression regulation and DNA damage responses. For example, FUS localises to paraspeckles and DNA damage foci in the nucleus[8,9]. Upon cellular stress, TDP-43 and FUS are exported to the cytoplasm where they become incorporated into stress granules, although neither are necessary for stress granule formation[10]. Stress granules are liquid-like cytoplasmic assemblies, or condensates, which are formed by liquid-liquid phase separation of both nuclear exported and constitutively cytoplasmic proteins, along with mRNA[11–13]. Stress granule formation is triggered by cellular stresses, such as oxidative stress. This is often dependent on the cytoplasmic stress granule protein G3BP1[10]. When the cellular stress is alleviated, stress granules dissolve and proteins that normally reside in the nucleus, including FUS and TDP-43 translocate back to the nucleus.

It has been proposed that ALS-linked FUS and TDP-43 mutants cause diseases in part by inducing aberrant phase transition of stress granules[12,14,15]. This reduces the dynamics of stress granule proteins, preventing them from dissolving when stress is removed, thus trapping nuclear proteins in the cytoplasm. Supportive evidence is that FUS and TDP-43 mutants often show constitutive mislocalisation to the cytoplasm, and FUS tends to aggregate more readily in the cytoplasm[16]. Both mechanisms may cause a loss-of-function phenotype in the nucleus or a gain-of-function (cytotoxic) phenotype in the cytoplasm as cytoplasmic aggregates or fibrils: these are associated with motor neuron dysfunction leading to neurodegenerative disease[17–19]. Either way, dissolving aberrant stress granules, reducing sensitivity to triggers of stress granule formation, preventing or reversing stress granule protein aggregation, and/or driving protein back to the nucleus might be an efficient way to prevent or reverse the consequences of ALS.

Small molecules have been an essential tool for relating cytology to function. For instance, the
microtubule polymerization inhibitor nocodazole have helped us to study cytokinesis and microtubule-
mediated intracellular trafficking. Inhibitors against actin/myosin II activities such as cytochalasin and
Y-27632 have illuminated pivotal roles of these cytoskeletal proteins in cell shape, division, and
migration. Indeed compounds have been identified that can disrupt stress granule condensation,
especially 1,6-hexanediol[20] and similar alcohols[21]. However, these compounds are both toxic and non-
specific, as they affect multiple condensates[21,22]. There is an unmet need for non-toxic and specific
compounds able to specifically dissolve condensates.   Here we searched for compounds that both
prevented the formation of stress granules, and induced their dissolution, and identify lipoamide,
which partitions into stress granules in cells and alleviates pathology caused by ALS-associated FUS
and TDP-43 mutants in both motor neurons *in vitro* and in fly models of ALS. Using lipoamide as a tool
compound, we identify a pathway that allows stress granules to sense the oxidative state of the cell.

Results

We performed a cell-based screen of 1,600 small molecules to identify compounds which affect stress
granule formation upon arsenate treatment, by monitoring GFP-tagged FUS (FUS-GFP) localisation in
HeLa cells (Fig. 1A–C). Many compounds altered FUS-GFP localisation in stressed cells, often reducing
the number of FUS-GFP-containing stress granules (Fig. 1B,C). Emetine, known to prevent stress
granule formation by stabilising polysomes[23], was present in the library and acted as a positive control.
Edaravone, an FDA-approved ALS therapeutic[24], had no significant effect.

The 47 strongest hits in HeLa cells were further tested *in vitro* for an effect on condensation of purified
FUS-GFP under physiological (low salt, 50 mM KCl and reducing, 1 mM DTT) conditions, with the aim
of selecting for compounds which can directly affect stress granule proteins (Fig. S1A). Seven
compounds significantly affected FUS-GFP condensates *in vitro* and fell into three compound classes
(Fig. S1B,C). Of these, surfactants have no plausibility as a systemic therapeutic, and heterotri- and
tetracyclic compounds have previously been investigated for anti-prion properties with limited
success[25]. Lipoamide was a novel hit for stress granule modulation. Lipoic acid, a related compound
featuring a carboxylic acid instead of the carboxamide, was also a good hit in HeLa cells (Fig. 1B).

Lipoamide prevents and reverses stress granule formation in cultured cells

To test whether lipoamide and lipoic acid affect stress granule formation or solely partitioning of FUS
into stress granules, we treated HeLa cells expressing five different GFP-tagged stress granule proteins
with lipoamide or lipoic acid. Pre-treatment with either compound for 1 h prior to 1 h arsenate stress
prevented cytoplasmic condensation for all proteins we tested, including G3BP1 (Fig. 1D). Addition of
lipoamide and lipoic acid to arsenate-stressed cells, in continued presence of arsenate, also led to
dissolution of pre-existing stress granules (Fig. S1D).

To assess whether lipoamide acts specifically on stress granules, we tested its effects on other
intracellular condensates such as P-bodies, Cajal bodies, and DNA damage foci and found that these
nuclear or cytoplasmic condensates were not affected (Fig. S1E). The specificity extended to stressor
types, as stress granule formation induced by oxidative stress and osmotic shock was inhibited
whereas stress granules still formed after heat treatment or inhibition of glycolysis (Fig. S1F). Therefore,
we conclude that the lipoamide activity is comparatively specific with regard to modulating the
properties of a cellular condensate.

To confirm that lipoamide enters cells and to determine its intracellular concentration, we synthesised
[15N]-lipoamide, which can be quantitatively detected by ^{15}N NMR (Fig. S2). Upon treatment of HeLa
cells with 100 µM [15N]-lipoamide, loss of [15N]-lipoamide signal from the growth medium indicated

that it accumulates in millimolar concentrations in cells. There was also a corresponding gain in [15N]-
lipoamide signal in the cell pellet (Fig. S2A–C). These indicate that lipoamide is taken into cells.

We used two strategies to ask whether lipoamide partitions into stress granules. Firstly, we used [15N]-
lipoamide with FUS condensates, as a minimal *in vitro* model of the stress granule environment. [15N]-
lipoamide from the dilute phase following FUS-GFP condensation under low salt reducing conditions
(Fig. S3A–D) partitioned into the FUS-GFP condensate phase by a factor of ten (Fig. 2A). To analyse
partitioning of lipoamide into stress granules in cells, we synthesized a lipoamide analogue derivatised
with a diazirine (for UV-induced crosslinking) and alkyne (for click chemistry) groups (Fig. 2B). This
dissolved stress granules with a slightly lower potency than lipoamide (Fig S3E), but allowed us to
crosslink this analogue to proteins in the vicinity by UV irradiation, subsequently labelling it *via* click
reaction with a fluorophore. Using the click-crosslink analogue at 30 µM (insufficient to dissolve stress
granules), we observed signal particularly in nuclei, mitochondria, and stress granules both without
and with crosslinking. Crosslinking increased colocalization with stress granules (Figs. 2C,D and S3F).
Comparison of signal intensity without and with crosslinking suggests that >50% of the analogues are
part of high affinity complexes with fixable macromolecules and/or are covalently bound to proteins
in each compartment (Fig. 2E). We suspect that this represents nonspecific binding of the reduced
dithiolane to proteins. Taken together, these data indicate that lipoamide partitions into stress
granules, along with other organelles.

SAR study identifies more potent lipoamide analogues and the dithiolane as a key feature

To determine which chemical features of lipoamide are required for activity, we synthesised a panel of
lipoamide-like compounds and tested the structure-activity relationship (SAR). As a reference, we
confirmed lipoamide potency: lipoamide pre-treatment, in both HeLa and induced pluripotent stem
cells (iPSCs), caused a dose-dependent decrease in stress granule numbers while an increase in
partition of FUS-GFP back to the nucleus with a similar dose-dependency (Fig. 3A). Titration analyses
of the series of lipoamide analogues identified 16 compounds with more than approximately five-fold
increased potency (EC_{50} < 2.5 µM, summarised in Fig. 3B–J), compared to lipoamide. Specifically,
lipoamide derivatives of 6-amino-3-substituted-4-quinazolinones and five-membered
aminoheterocyclic amides represented the most potent analogues (Fig. 3I).

The (*R*) and (*S*) isomers of lipoamide and lipoic acid had a similar EC_{50}, indicating little stereoisomer
specificity (Fig. 3B). The chemical structure of lipoamide is similar to that of the lipoyl moiety, used as
a hydrogen-accepting cofactor by two mitochondrial Krebs cycle enzymes, which is recycled to the
oxidised state by dihydrolipoamide dehydrogenase (DLD, also mitochondrial)[26]. Cells exclusively use
the (*R*)-lipoyl moiety stereoisomer. However, the comparable EC_{50} between the (*R*)- and (*S*)-isomers of
lipoamide and the absence of a lipoate ligase in eukaryotic cells[26] indicate that lipoamide does not
primarily function through these mitochondrial proteins to effect stress granule dissolution.

Mono-methylation of lipoamide on the amide improved the activity, while di-methylation reduced it
(Fig. 3C). However, other disubstituted amide analogues showed activity in the context of more
complex amide structures (Fig 3D). Indeed, many mono-substitutions of the lipoamide amide
improved the activity, with no clear trend for beneficial substitutions, *i.e.*, a relatively 'flat' SAR space
in the carboxamide group. Dissimilar substitutions could similarly increase potency to low µM (Fig. 3E),
while some similar heterocycle substitutions could have a wide range of potencies (Fig. 3F). Activity
could be retained and even increased by shortening the alkane 'backbone' (Fig. 3G). A compound
without a carboxamide or carboxylic acid moiety (*i.e.*, unlike both lipoamide and lipoic acid) was active,
with increased potency (Fig. 3H), although the most potent compounds were mono-substituted
amides (Fig. 3I).

Importantly, the dithiolane ring is necessary for activity, indicating a redox activity for stress granule
dissolution. Lipoamide derivatives are likely reduced in the cellular environment. Indeed, the reduced
dihydrolipoamide form was active (Fig. 3J). Furthermore, a labile thiol modification (two thioesters)
was active while non-labile derivatives (thiol benzylation and substitution to a tetrahydrothiophene, a
thiolane ring) were not (Fig. 3J). A six-membered disulfide ring removed activity (Fig. 3J). As redox
potential is linked to disulfide ring size, this also indicated a redox-linked mechanism. Since Edaravone
and ascorbic acid, other redox active compounds[27,28], did not reduce stress granules at micromolar
concentrations comparable to lipoamide (Fig. 1B and S1G), this suggests that lipoamide is more potent
than those compounds to control stress granule dynamics. Taken together, the SAR suggests that
lipoamide acts through a non-enzymatic route and likely through a redox-associated process.

Lipoamide weakly increases liquidity of FUS condensates *in vitro*

To test if lipoamide interacts with known stress granule proteins, we turned again to FUS, using the
classical methods of isothermal titration calorimetry (ITC) and chemical shift perturbation in
"fingerprint" ^{1}H-^{15}N 2D protein NMR spectra. We could not detect interaction of FUS-GFP with
lipoamide *in vitro* by ITC. NMR of the N terminal prion-like domain of FUS showed only extremely weak
^{1}H and ^{15}N shifts in the presence of lipoamide (Fig S4A,B). To test if lipoamide alters FUS condensate
formation, we examined the critical salt concentration and temperature of *in vitro* FUS condensates,
but found no detectable change in the presence of lipoamide (Fig. S4C).

We then tested whether lipoamide alters FUS condensate properties, first testing the effect on *in vitro*
condensate liquidity using laser optical tweezers to assay droplet fusion. This showed significantly
decreased droplet fusion times in the presence of lipoamide and thus increased liquidity (higher ratio
of surface tension to viscosity) (Fig. S4D,E). Over time FUS condensates gradually harden, visible as an
increasing viscosity and decreasing mobile fraction of FUS, and eventually forming solid fibres. This is
particularly prominent for condensates of ALS-linked mutant FUS G156E, which hardens then forms
fibres rapidly[12]. We tested whether lipoamide maintains condensate liquidity, using fluorescence
recovery after photobleaching (FRAP). Both lipoamide and lipoic acid reduced FUS G156E-GFP
condensate hardening and fibre formation (Fig. S4F–H).

Finally, we turned to mass spectrometry to analyse changes in FUS G156E self-interaction *in vitro* in
the presence of lipoamide. We used lysine-lysine (K-K) chemical crosslinking and, following tryptic
digest, mass spectrometry detection of the cross-linked peptides as evidence for inter- and intra-
molecular interactions under different conditions. This technique requires lysine residues, which the
N-terminal IDR of WT FUS lacks. Therefore, we also analysed a FUS mutant with 12 lysine substitutions
in the N-terminal domain. Lipoamide caused a change, predominantly decrease, in the intensity of
identified K-K cross-linking sites and therefore suggested reduced FUS-FUS interactions (Fig. S4I,J).
Taken together, lipoamide has a weak effect on FUS condensate properties *in vitro* by modulating FUS-
FUS interactions and does so without strong small molecule-protein binding typically detectable by ITC
or NMR.

Lipoamide stabilises proteins with arginine/tyrosine-rich low complexity domains in cells

As the effects of lipoamide on FUS *in vitro* were likely too small to explain the effect of lipoamide on
stress granule dynamics in cells, we turned to thermal proteome profiling (TPP). Here, aliquots of HeLa
cells treated with DMSO (solvent control), lipoamide, arsenate, or lipoamide and arsenate were heated
to a range of different temperatures, and the abundance of soluble proteins was measured by
quantitative mass spectrometry. Relative increase in abundance with temperature is indicative of
protein thermal stability[29,30] (Fig. 4A), summarised as z-scores (Fig. 4B and S5A,B). Increased protein

thermal stability in the presence of a small molecule often indicates interaction[31,32]. Thermal stabilities
of proteins in lipoamide vs. lipoamide and arsenate-treated cells showed a strong positive correlation,
but arsenate vs. lipoamide and arsenate-treated cells did not, indicating a dominant effect of lipoamide.
Furthermore, lipoamide treatment also broadly reversed the thermal stability changes occurring due
to arsenate treatment (Fig. S5A,B). Therefore, we focused on the analysis of the sample treated with
both lipoamide and arsenate. As a positive control, we confirmed that the thermal stability of DLD was
weakly but significantly increased, consistent with lipoamide binding to the active site (z = 0.66±0.007,
adjusted p-value: false discovery rate [FDR] = 2.6×10^{-4}). As we would predict from its mitochondrial
localization and enzymatic function, RNAi of DLD affected neither stress granule formation nor the
lipoamide activity to prevent it (Fig. S5C). Histone deacetylase 1 (HDAC1) and HDAC2 were also
stabilized (z = 3.58, FDR = 1.8×10^{-14} and z = 4.75, FDR = 7.4×10^{-5}, respectively), consistent with a
recent report[33].

Many proteins had higher TPP z-scores than DLD (Fig. 4B): Lipoamide and arsenate treatment resulted
in significantly increased thermal stability of 70 proteins, while reducing the thermal stability of 144
proteins compared to no treatment (Fig. 4B). Stabilised proteins had disproportionately long IDRs
which contained a disproportionly high proportion of arginine (R), tyrosine (Y), and phenylalanine (F)
residues, while destabilised proteins showed the opposite trend (Fig. 4B–D). R and Y-rich IDRs are
characteristic of stress granule proteins such as the FET family (FUS, TAF15, and EWSR1)[34], although
their thermal stability was not significantly increased (z = 0.84±1.3, FDR = 0.73; z = 2.08±1.5, FDR =
0.20; z = 0.05±0.82, FDR = 0.87, respectively). This is consistent with no obvious interaction between
FUS and lipoamide *in vitro*. Also, individual FET family proteins are neither necessary for stress granule
formation[10] nor lipoamide activity (Fig. S5C), further indicating that they are not primary targets of
lipoamide for stress granule dissolution.

Specific stress granule proteins are necessary for lipoamide activity in cells

To assess which of the proteins identified as interacting with lipoamide by TPP are necessary for
lipoamide activity, we performed an endoribonuclease-prepared small interfering RNA (esiRNA)[35]-
mediated gene knockdown screen of all 122 proteins with increased thermal stability (z > 2) (Table S1,
Fig S5A). We looked for the genes whose depletion reduced lipoamide activity in preventing stress
granule formation, which identified two IDR-rich proteins: splicing factor proline- and glutamine-rich
(SFPQ) and splicing factor serine/arginine-rich splicing factor 1 (SRSF1)[36,37]: Lipoamide pre-treatment
failed to prevent stress granule formation in cells with either gene depletion (Figs. 4E,F and S5D). RNAi
of SFPQ or SRSF1 also prevented dissolution of pre-existing stress granules following lipoamide
treatment (Fig. S5E). Stress granule formation was neither exacerbated in stressed cells nor induced in
non-stressed cells by these RNAis, showing that the phenotype of lipoamide pre-treated cells is not
simply due to basal increase in stress granule formation (Figs. 4E,F and S5F).

Both SFPQ and SRSF1 are stress granule proteins: like FUS and TDP-43, both localise to the nucleus in
unstressed cells and, in stressed cells, stress granules (Fig. 4G). Therefore, lipoamide activity dissolving
stress granules is dependent on at least two stress granule proteins.

Redox state controls SPFQ-mediated dissolution of stress granule protein condensates

313 The activity of lipoamide requires the redox active dithiolane (Fig. 3J), and SFPQ is notably rich in redox-
314 sensitive methionine (28 out of 707 amino acids; Fig. 5A). SRSF1 is not methionine-rich with only three
315 out of 248 amino acids. Pioneering work has previously shown that methionine oxidation modulates
316 function and material properties of phase separated yeast ataxin-2[38], and methionine oxidation in
317 SFPQ has been detected in cells[39]. These suggest that SFPQ may be a main target of lipoamide in a

redox-based mechanism of action. We used in vitro experiments using pure protein to analyse the effect of oxidation of SFPQ on condensate formation, using the oxidizing agent hydrogen peroxide (H_2O_2). SFPQ condensation was induced at low salt concentration (75 mM KCl; Fig. S6A). Oxidation of SFPQ, confirmed by modulated electrophoretic migratory aptitude in non-reducing SDS-PAGE (Fig. S6B), led to dissolution of SFPQ condensates in an H_2O_2 concentration-dependent manner (Fig. S6A), similar to the behavior of axaxin-2 condensates[38]. This suggests that oxidation alters properties of SFPQ proteins. In contrast, H_2O_2 alone did not lead to FUS condensate dissolution (Fig. S6A). Therefore, oxidation-mediated condensate dissolution is specific to a subset of proteins.

We next examined functions of SFPQ and its oxidation on FUS condensate dynamics at a physiological salt concentration (150 mM KCl), which kept SFPQ proteins in diffused state (Fig. S6C). We found that adding SFPQ prevented FUS condensation (Fig. S6C). This effect did not occur by adding only GFP proteins (Fig. S6C). However, H_2O_2 treatment restored FUS condensates even in the presence of SFPQ (Fig. S6D). These suggest that SFPQ proteins dissolve stress granule protein condensates in a redox state-dependent manner.

Based on these *in vitro* results, we hypothesized that stress granule formation would be attenuated if SPFQ is not oxidizable. To assess this possibility, we aimed to replace methionine with a non-oxidizable, non-natural analogue L-azidohomoalanine (AHA), normally used for protein labeling[40] (Fig. 5B). Cells were cultured in methionine-free medium supplemented with AHA for 2 h, resulting in methionine-to-AHA replacement in newly synthesized proteins, and then stressed with arsenate for 1 h (Fig. S6E). This resulted in attenuated stress granule formation (Fig. 5C), and normal stress granule formation was rescued by depletion of SFPQ (Fig. 5C). This suggests that SFPQ in the reduced state is responsible preventing stress granule formation. One possibility is that cellular stress leads to oxidation of SFPQ, allowing stress granule condensation. In this scenario, lipoamide reduces SFPQ and restores its stress granule dissolution activity (Fig. 5D).

Lipoamide treatment rescues nuclear localization and functions of FUS and TDP-43

FUS and TDP-43 have important nuclear functions in unstressed cells. We asked whether lipoamide treatment not only dissolves stress granules but also returns these proteins to the nucleus. Similar to FUS-GFP (Fig. 3A), lipoamide pre-treatment increased partition of TDP-43 and the ALS-associated nuclear localisation sequence mutant FUS P525L-GFP to the nucleus in stressed HeLa cells (Fig. S7A). To confirm that these effects also occur in cells prominently defective in ALS, we analysed iPSC-derived motor neurons (MNs). Lipoamide had similar effect on nuclear partitioning of wild-type TDP-43 in stressed (prolonged oxidative stress with low dose [10 μM] of arsenite) and FUS P525L-GFP in non-stressed but long-cultured conditions (Figs. 6A,B and S7B,C)[41].

We characterised the functional importance of the re-localization to the nucleus by considering FUS and TDP-43 nuclear functions. FUS forms condensates at DNA damage sites to engage in DNA damage repair, and this malfunction caused by ALS-linked mutations on FUS is implicated to underlie neuronal dysfunction in ALS[42]. Lipoamide increased recruitment of FUS-GFP (WT in iPSCs and P525L in iPSC-derived MNs) to laser-induced DNA damage foci (Fig. 6C,D). TDP-43 contributes to normal transcript splicing in the nucleus, particularly of Stathmin-2 (STMN2) transcript, and altered *STMN2* splicing leading to reduced transcript levels is a hallmark of ALS[6,43]. In iPSC-derived MNs, the prolonged oxidative stress recapitulated reduction in *STMN2* mRNA levels. This reduction in splicing was rescued by lipoamide treatment, concomitant with TDP-43 nuclear partitioning (Fig. 6B,E). Lipoamide action therefore dissolves stress granules, allows return of those ALS-linked proteins to the nucleus, and restores nuclear functions of FUS and TDP-43.

Lipoamide alleviates ALS phenotypes in familial ALS models

The ultimately lethal phenotype of ALS is thought to be caused by axon defects in motor neurons. Indeed, iPSC-derived MNs expressing FUS P525L show a motor neuron survival defect *in vitro*, with reduced neurite growth and defective axonal transport[41]. Lipoamide treatment rescued neurite growth of iPSC-derived MNs stressed with arsenite, shown by increased area covered in neurites in a non-polarised culture (Fig. 7A). We tested if this correlated with improved axonal transport, by tracking lysosome transport in iPSC-derived MN axons grown through silicone channels (Fig. 7B). As previously observed, in an unstressed condition, distal axonal transport of lysosomes was disrupted by expression of FUS P525L[41], and lipoamide recovered transport to a similar level to that in WT FUS iPSC-derived MNs (Fig. 7C,D). Motor neuron degeneration caused by an ALS-associated FUS mutant can therefore be rescued by lipoamide.

Aggregation of TDP-43 and FUS in neurons is a hallmark pathology of ALS, and aggregation of proteins is also a phenotype of aging more generally, including in *C. elegans*[44]. Feeding lipoic acid (it has higher solubility in food media than lipoamide) caused a dose-dependent reduction in the number of aggregates of transgenic PAB-1, an orthologue of the human stress granule protein PABPC1 (Fig. 7E), but not those of a non-stress granule protein RHO-1 (Fig. S8A)

In *D. melanogaster*, motor neuron-specific expression of human FUS and TDP-43 induces ALS-like phenotypes, including motor defects manifesting as a reduced ability for negative geotaxis[45,46]. Feeding either lipoamide or lipoic acid improved climbing ability in flies expressing FUS nuclear localizing signal (NLS) mutants, either FUS P525L or R521C (Figs. 7F and S8B). Similarly, lipoamide feeding alleviated climbing defects in flies expressing TDP-43, either WT or an ALS-linked mutant M337V (Fig. 7F). The severe phenotype caused by TDP-43 M337V was associated with abnormal neuromuscular junction morphology, the presence of satellite boutons, similar to previously described phenotypes of another ALS-linked mutant TDP-43 G298S[47]. Lipoamide treatment suppressed appearance of satellite boutons (Fig. S8C). Collectively, our data show that lipoamide can alleviate ALS-like phenotypes in patient-derived motor neurons and animal models caused by expression of two ALS-associated stress granule protein mutants.

Discussion

Stress granules are an example of a liquid cellular compartment formed by phase separation. Due to the strong genetic association of ALS with stress granule proteins, we sought small molecules which alter the physiological function of stress granule proteins in forming biological condensates. Our screen identified lipoamide, which partitioned into stress granules in cells and their *in vitro* model (condensates formed with the stress granule protein FUS). Lipoamide caused rapid disassembly and prevented formation of stress granules. Our SAR showed the potency of lipoamide for stress granule dissolution could be increased, particularly by mono-substituted amines and some alkane backbone modifications.

The dithiolane is always required for activity of lipoamide-like molecules. However, the degree to which other areas of lipoamide could be modified while retaining activity was large. This shows the lipoamide compound family has potential for medicinal chemistry development. The high degree to which the non-dithiolane regions could be altered would be surprising for a molecule that binds a protein at a structured binding site, although this is perhaps unsurprising given the interaction of lipoamide with intrinsically disordered proteins. Overall, lipoamide is likely therefore acting by delivering a redox-active dithiolane payload to physicochemical environments formed by its interacting

proteins, including stress granules. More active lipoamide derivatives are likely improving targeting
(cell uptake and partition into stress granules) while leaving the dithiolane payload intact.

Our work has identified a plausible candidate for the protein likely to sense the redox state of the cell.
Among all the proteins stabilised by lipoamide, our RNAi screen showed only SFPQ and SRSF1 were
necessary for the rapid (<20 minute) lipoamide effect on stress granule dynamics. Both are stress
granule proteins. Uniquely, SFPQ is very methionine-rich, which likely confers high redox sensitivity not
found in most arginine/tyrosine-rich IDR-containing proteins. SFPQ activity was indeed methionine-
and redox-dependent.   SFPQ appears the primary target for early lipoamide activity, which overrides
the ability of SFPQ to act as an oxidation sensor, promoting stress granule dissolution only when
reduced. While we saw some effect of lipoamide on FUS condensate liquidity *in vitro*, we suspect that
this is a similar effect of secondary importance on a less redox-sensitive stress granule protein or a
minor effect from strong partitioning of lipoamide into this condensate. Overall, our identification of
lipoamide allowed us to dissect a mechanism where dissolution of stress granule condensates involves
direct redox sensitivity of a key stress granule protein. Lipoamide therefore represents productive
small molecule intervention in an emerging paradigm: redox-sensitivity of proteins able to phase
separate as a homeostatic mechanism[38,48,49].

Oxidative stress is a common theme in ALS pathogenesis mechanisms[50–52]. We showed that lipoamide
can recover pathology in motor neurons and animals expressing ALS-associated stress granule mutants
(FUS and TDP43) with no explicit oxidative stress. This leaves the link between the redox-associated
cellular effects of lipoamide on stress granules and neuron/animal model outcomes ambiguous, but
consistent with modulated stress granule formation in response to stochastic oxidative stresses.
Previous works described that HDAC1 and 2 are responsible for long-term changes in histone
acetylation in motor neurons[53] and that they are inhibited by lipoamide[33]. However, we did not find
them necessary for short-term lipoamide activity on stress granules, instead detecting the two stress
granule proteins both necessary for lipoamide activity. However, this does not preclude long-term
nuclear effects. FUS NLS mutations are strongly associated with ALS[41,54–56] and dissolution of stress
granules by lipoamide leads to return of FUS and TDP-43 to the nucleus. Indeed, we saw lipoamide
does not prevent nuclear FUS condensation and rescues nuclear TDP-43 functions. Ultimately, this
relates to central questions about ALS pathogenesis. Does persistent stress granules or stress granule
protein aggregation lead to harmful (gain-of-function) effects, or is stress granule formation
fundamentally beneficial over a short time scale but, over a long time scale, leads to defects from
nuclear loss-of-function by sequestration of proteins in the stress granules? – overall our results are
consistent with the latter.

Although lipoamide does not have the characteristics of a typical therapeutic, it is notable that lipoic
acid is used to treat diabetic neuropathy and, in humans, a 600 mg daily dose gives plasma
concentrations of 8 to 30 µM[57,58]. This is comparable to the concentrations used in our cell-based
assays and we saw beneficial effects in patient-derived motor neurons and *D. melanogaster* models of
ALS. Therefore lipoamide, in addition to allowing our discovery of direct redox sensation by SFPQ for
stress granule dissolution, has some plausibility as the basis of a therapeutic with medicinal chemistry
potential for further improvements. However it is important to point out that we have not shown a
direct relationship between stress granule dissolution and phenotype rescue on our disease models.
Future work will be required to understand the relationship between stress granule formation and
disease in animal models. However the identification of lipoamide provides a powerful tool for such
investigation.

Methods

Cells and cell lines

Kyoto HeLa cells were maintained in high glucose Dulbecco's modified Eagle's medium (DMEM, Thermo Fisher Scientific) supplemented with 10% FBS and 1% penicillin-streptomycin at 37°C with 5% CO_2. Stable Kyoto HeLa BAC cell lines expressing proteins with a C-terminal fluorescent protein were generated using BAC recombineering[59]. This gives near-endogenous expression levels of the fusion protein[12,60]. In these lines, GFP is part of a modified localisation and affinity purification (LAP) tag[61], providing a short linker. The stable expression was kept under G418 selection (400 µg/ml; Gibco). The following BAC lines were used: FUS (MCB_005340) (used also for the compound screen [see Small molecule screen]), COIL (MCB_0002582), DCP1A (MCB_0003876), EWSR1 (MCB_0008863), PABPC1 (MCB_0006901), TIAL1 (MCB_0008967), and TRP53BP1 (MCB_0003740). HeLa FUS P525L-GFP cells were generated similarly to the iPS cell lines as described previously[41].

Human iPS cells were grown in either TeSR E8 or mTeSR1 medium (Stem Cell Technologies) at 37°C with 5% CO_2 (ref.[62]). iPS cells lines, derived from two different donors, expressing FUS with a C-terminal GFP fluorescent marker were used. All were generated using CRISPR/Cas9 assisted tagging and mutagenesis and have been previously described[41]. KOLF iPS cell lines expressing wild-type FUS-GFP or FUS P525L-GFP were previously generated from the KOLF-C1 clonal iPS cell line produced as part of the human induced pluripotent stem cell initiative (HipSci)[63]. KOLF-C1 cells were derived from a healthy male donor. In these lines, GFP is part of a modified localisation and affinity purification (LAP) tag[61], giving an identical fusion protein sequence to the Kyoto HeLa BAC cell line. AH-ALS1-F58 iPS cell expressing FUS P525L with a C-terminal GFP fluorescent marker were previously generated from a clonal iPS cell line from a female ALS patient expressing FUS P521C. The P525L mutation and GFP tag were introduced and the P521C mutation corrected by simultaneous tagging and mutagenesis[41,64,65].

MNs for FUS P525L dynamics were induced and maintained as described previously[66]. MNs for the prolonged arsenite stress assay were derived from commercially available WTC-11 iPS cells (Coriell Institute GM25256) and differentiated as described previously[67]. MNs for axonal lysosome mobility assays were generated from AH-ALS1-F58 iPS cells expressing FUS P525L. In short, the iPS cells were differentiated into neuronal progenitor cells and matured to spinal MNs in Matrigel-coated plates as previously described[41,62]. The coating and assembly of silicone microfluidic chambers (MFC; RD900, Xona Microfluidics) to prepare for the subsequent seeding of MNs was performed as described previously[41,68,69]. MNs were eventually seeded into one side of an MFC for maturation to obtain a fully compartmentalized culture with proximally clustered somata and their dendrites being physically separated from their distal axons, as only the latter type of neurite was able to grow from the proximal seeding site through a microgroove barrier of 900 µm-long microchannels to the distal site (Fig. 7B). All subsequent imaging in MFCs was performed at day 21 of axonal growth and MN maturation (day 0 = day of seeding into MFCs).

All procedures using human cell samples were in accordance with the Helsinki convention and approved by the Ethical Committee of the Technische Universität Dresden (EK45022009, EK393122012).

Recombinant protein purification

Recombinant proteins were purified using baculovirus/insect cell expression system, as previously described[12]. Briefly, 6×His-MBP-FUS-GFP and 6×His-MBP-FUS-SNAP were purified from Sf9 cell lysate by Ni-NTA (QIAGEN) affinity purification. The 6×His-MBP tag was cleaved by 3C protease, concentrated by dialysis, and further purified by size exclusion chromatography. 6xHis-MBP-SFPQ-GFP was purified

from Sf9 cell lysate by affinity purification using Ni-NTA and amylose resin (New England Biolabs). The
6×His-MBP tag and, if necessary, the GFP tag were cleaved by 3C protease and TEV protease,
respectively, and the target proteins were concentrated by dialysis and further purified by cation
exchange chromatography. The composition of the storage buffer for the purified proteins was 1 M or
500 mM KCl, 50 mM Tris-HCl pH 7.4, 5% glycerol and 1 mM DTT, and FUS concentration was adjusted
to 30 µM in storage buffer prior to use.

Small molecule screen

For the small molecule screen we used the PHARMAKON 1600 library of small molecules, prepared as
10 mM solutions in DMSO. The Kyoto HeLa BAC cell line stably expressing FUS-GFP was seeded at 4000
cells per well in 384 well plates 24 h before the assay. The cells were pre-treated with 10 µM compound
for 1 h and then stressed with 1 mM potassium arsenate (A6631, Sigma Aldrich). After 1 h, cells were
fixed in 4% formaldehyde and stained with 1 µg/ml Hoechst 33342 and CellMask blue (1:10,000;
H32720, Thermo Fisher Scientific) before imaged on a CellVoyager CV7000 automated spinning disc
confocal microscope (Yokogawa) with a 40× NA 1.1 air objective to assess FUS-GFP localisation.

FUS-GFP signal was analysed using CellProfiler[70], and the data were processed with KNIME. Cytoplasm
and nuclei were distinguished with weak (CellMask blue) and strong (Hoechst 33342) blue fluorescent
signals, respectively. Particle number and sum area, granularity (at 9, 10, and 11 px in the cytoplasm
or 1, 5, 6, 7, 8, and 9 px in the nucleus) scale, texture (at 10 px scale), and integrated signal intensity of
FUS-GFP in the nucleus and cytoplasm were measured. Z scores ($z = (x - \mu)/\sigma$ where x is the observed
value, μ the control mean and σ the control standard deviation) relative to the DMSO treated control
wells on each plate were calculated for these parameters and combined into the Mahalanobis distance.
Compounds of interest were selected on the criteria of: treatment returned the cells to the unstressed
state (ie. reducing stress granule number, increasing nuclear signal), a clear monotonic dose
dependent response, and by manual prioritisation by known mechanism (e.g. emetine, cardiac
glycosides) or implausibility as a cell-compatible compound (e.g. surfactants, used as topical
antiseptics).

The follow-up *in vitro* assay of compounds on FUS-GFP condensates was assessed in a 384 well plate
format. The compound volume (in DMSO) necessary for 1, 3, 10, 30 or 100 µM final concentration
were added by acoustic dispensing (Labcyte Echo 550) to 96 well plate wells containing FUS-GFP in 3 µl
of 50 mM Tris-HCl pH 7.4, 1 mM DTT, and 170 mM KCl. Final DMSO concentration was 0.01 to 1%.
Using a Freedom Evo 200 liquid handling workstation (TECAN) the FUS-GFP/compound mixture was
diluted in 7 µl 50 mM Tris-HCl pH 7.4 to reach the final composition of 50 mM Tris-HCl pH 7.4, 1 mM
DTT, 50 mM KCl, the indicated concentrations of each compound and DMSO, and 0.7 µM FUS-GFP.
Compound/FUS-GFP and assay buffer were mixed by a standardised pipetting procedure, split to four
wells in clear bottom 384 well plates, and then immediately imaged using a CellVoyager CV7000
automated spinning disc confocal microscope (as above). Condensates in suspension for six fields of
view were imaged as a maximum intensity projection of 6 focal planes at 2 µm steps per sample.
Condensate number and FUS-GFP partition into condensates were analysed with a fixed intensity
threshold using Fiji. Number of condensates and partition were weakly time dependent due to
condensate sedimentation, so normalised assuming a linear change over time by reference to DMSO
controls at the start and end of each plate row.

Compound characterisation on cells

Compound effects were assessed under a variety of conditions in HeLa cells, iPS cells, or iPSC-derived
motor neurons. Unless otherwise indicated, cells were pre-treated for 1 h using 10 µM compounds

from 10 mM stock in DMSO (or an equal volume of DMSO control) then stressed for 1 h with 1 mM
potassium arsenate still in the presence of the compounds. Live cells were imaged by widefield
epifluorescence using an inverted Olympus IX71 microscope with a 100× NA 1.4 Plan Apo oil immersion
objective (Olympus) and a CoolSNAP HQ CCD camera (Photometrics), using a DeltaVision climate
control unit (37°C, 5% CO_2) (Applied Precision).

Various cellular stresses were achieved by replacing 1 h 1 mM potassium arsenate treatment with
other conditions: 0.4 M sorbitol (S1876, Sigma Aldrich) from a 4 M stock in water for 1 h (osmotic
stress); 42°C in normal growth medium for 30 min (heat stress); 100 mM 6-deoxyglucose (D9761,
Sigma Aldrich) from a 1 M stock in H_2O in glucose free DMEM (11966025, Thermo Fisher Scientific)
supplemented with 10% FCS for 1 h (glycolysis stress). L-ascorbic acid (A4544, Sigma Aldrich) was used
from 1 M stock in H_2O. Sodium arsenite (S7400, Sigma Aldrich) was used from 10 mM stock in H_2O.

Compound dose responses

Dose dependent effect of lipoamide on HeLa and iPS cells expressing FUS-GFP were assessed with pre-
treatment of lipoamide for 1 h followed by 1 h treatment with 1 mM potassium arsenate similar to the
ex vivo HeLa cell screen, except serial compound dilutions in medium were prepared manually from
80 µM to ~0.4 nM at 1.189× dilution steps. Small dilution steps rather than concentration replicates
were selected as it provides greater statistical power from a set number of samples[71]. Final DMSO
concentration was 0.08% in all samples, and each plate included at least 12 control wells with 0.08%
DMSO. Cytoplasmic FUS-GFP condensate number and nuclear/cytoplasm partition of FUS-GFP were
analysed using custom macros in Fiji. Nuclei were identified by intensity thresholding of DNA images
labelled with Hoechst following a 5 px Gaussian blur. Cytoplasmic FUS-GFP condensates were identified
by intensity thresholding of the FUS-GFP images following a 10 px weight 0.9 unsharp filter masked by
the thresholded nuclei. The ratio of the number of cytoplasmic FUS-GFP condensates to that of nuclei
was taken as cytoplasmic FUS-GFP condensates per cell per field of view, and p, the ratio of partition
of FUS-GFP to the nucleus and the cytoplasm, was derived from $a = v_n/v_t$, the ratio of nuclear to total
green signal per field of view, where $p = a/(1 - a)$. These data were log transformed and fitted to a
Rodbard sigmoidal curve[72] to determine EC_{50}. Six fields of view were captured and analysed per
condition.

The series of lipoamide analogues including lipoamide and lipoic acid were newly synthesized by Wuxi
AppTec and provide through Dewpoint Therapeutics. To assess those dose response effect, the HeLa
BAC cells of FUS-GFP were seeded in 384-well plates (4000 cell per well) 24 h prior treatment, pre-
treated with the compounds in a half-log dilution series (from 30 µM to 3 nM: seven concentrations)
using an Echo 650, and followed by a 1 h treatment with 1.5 mM potassium arsenate before fixation
with 4% formaldehyde for 15 min, permeabilization with 0.1% Triton X-100 for 10 min, and counter-
staining with Hoechst and cell mask blue as described above. Imaging was performed using an Opera
Phenix (PerkinElmer), 20×, 9 FOV, binning 2, and using Harmony 4.9 software to determine cytoplasmic
FUS-GFP condensates number as well as cytoplasmic and nuclear FUS-GFP intensities to calculate
nuclear to cytoplasmic ratio of FUS-GFP intensities. EC_{50} was calculated either using CDD Vault curve
fitting or Harmony 4.9 software.

***In vitro* protein condensation, solidification, and oxidation assays**

For the condensation assay at different KCl concentrations, FUS-GFP proteins in storage buffer was
diluted with 20 mM HEPES pH 7.25 containing DMSO and lipoamide to give 20 µl of indicated
concentrations of the protein and KCl, 0.3 mM DTT, and 300 µM lipoamide (0.3% DMSO), and placed
on a 384-well plate (781096, Greiner). Condensates were imaged on a Nikon TiE inverted microscope

with a Nikon Apo 60× NA 1.2 water immersion objective using a Yokogawa CSU-X1 spinning disk head
and an Andor iXon EM+ DU-897 EMCCD camera

The assay to determine dilute phase concentrations at different temperature was performed with a
newly established technique, which will be reported in detail elsewhere. In brief, the technique is
based on mass and volume conservation and defined reaction volumes. We can use this method to
determine accurate values for both dilute and condensed branch protein concentrations. Here, FUS-
GFP phase separation was induced for a protein concentration titration in water-in-oil emulsions in a
buffer containing 25 mM Tris-HCl pH 7.4, 150 mM KCl, 1 mM DTT, and the indicated concentrations of
lipoamide (or DMSO as control) and imaged with a CSU-W1 (Yokogawa) spinning disk confocal system
at an IX83 microscope with a 40x UPlanSApo 0.95 NA air objective, controlled via CellSens (Olympus).
The dilute phase protein concentration was derived from a linear fit to the volume fractions of FUS-
GFP condensed phase versus the total concentrations of FUS-GFP. Temperature was controlled using a
custom-made stage[73].

For solidification assays, FUS-GFP in storage buffer was diluted in 50 mM Tris-HCl pH 7.4, 1 mM DTT to
give 10 µM protein, 50 mM Tris-HCl pH 7.4, 1 mM DTT, 50 mM KCl in a 20 µl volume in non-binding
clear bottom 384 well plates (781906, Greiner). Compounds, or an equal volume of DMSO, were then
added for a final compound concentration of 30 µM and 0.3% DMSO. 'Aging' to cause fibre formation
was induced by horizontal shaking at 800 rpm at room temperature, as previously described[12]. Fibre
and condensate formation were analysed by widefield epifluorescence using a DeltaVision Elite
microscope (GE Healthcare Life Sciences) with a Plan ApoN 60× NA 1.4 oil immersion objective
(Olympus) and an sCMOS camera (PCO). Fluorescence recovery after photobleaching (FRAP) of FUS-
GFP condensates and fibres was performed on a Nikon TiE inverted microscope with a Nikon Apo 100×
NA 1.49 oil immersion objective using a Yokogawa CSU-X1 spinning disc head and an Andor iXon EM+
DU-897 EMCCD camera. 10×10 px regions were bleached for 50 ns with a 6 mW 405 nm laser using an
Andor FRAPPA beam delivery unit then imaged for 5 min at 5 Hz. Recovery curves were derived using
scripts in Fiji.

Oxidation of SFPQ was detected by change in mobility in SDS-PAGE without reducing agents. 10 µM of
untagged SFPQ in buffer (20 mM HEPES pH 7.25 and 150 mM KCl) was incubated with H_2O_2 at RT for
30 min before subjecting to SDS-PAGE. For condensation assays of individual proteins with H_2O_2,
condensates of SFPQ-GFP and FUS-GFP were induced in buffer (20 mM HEPES pH 7.25 and 75 mM KCl).
The assays for dissolution and revival of FUS-SNAP condensates were performed in buffer (20 mM
HEPES pH 7.25 and 150 mM KCl). FUS-SNAP was labelled with SNAP-Surface Alexa Fluor 546 (New
England Biolabs), and protein mixtures were oxidized with H_2O_2 at RT for 1 h before image acquisition.
Proteins were imaged similarly to FUS-GFP condensates above.

Controlled droplet fusion using optical tweezers

Liquidity of FUS protein condensates was assessed by controlled fusion experiments using dual-trap
optical tweezers, as detailed previously[12,34]. In short, for each independent fusion event, two FUS
protein droplets in the presence of 300 µM lipoamide or equivalent amount of DMSO (0.3%) as the
control were trapped in each optical trap and brought into contact to initiate droplet coalescence.
Fusion relaxation times were accurately recorded as changes to the laser signal as condensate material
flows into the space between the two optical traps during coalescence. The laser signal was recorded
at 1 kHz, smoothed at 100 Hz and used to extract the characteristic relaxation time. After fusion was
complete – as indicated by a stable laser signal– the fused droplet was discarded, and two new droplets
were captured for quantifying an independent fusion event.

 Ex vivo DNA cut assays

UV micro-irradiation was performed as previously described[12,41]. Briefly, iPSCs expressing wild-type
FUS-GFP were stressed by addition of 1 mM arsenate for 1 h, then treated with lipoamide or an equal
volume of DMSO for 1 h. A single point in the nucleus was subject to 3 UV pulses as described for FRAP,
but at 10% laser power. GFP fluorescence was imaged at 1 Hz, and intensity of response was analysed
on Fiji. iPSC-MNs expressing FUS P525L-GFP were pre-treated with 20 µM Lipoamide for 24 h before
laser irradiation. The UV laser cutter setup utilized a 355-nm UV-A laser with a pulse length of <350 ps.
A Zeiss alpha Plan-Fluar 100× 1.45 oil immersion objective was used, and 12 laser shots in 0.5 µm-steps
were administered over a 12 µm linear cut.

NMR for FUS-lipoamide interaction

635 Untagged FUS low complexity domain (residues 1 to 163) was expressed, purified, and analysed using
636 ^{1}H-^{15}N heteronuclear single quantum coherence NMR and sample conditions previously described[74] in
637 the presence of 500 µM lipoamide or equivalent DMSO solvent control (1%).

NMR for lipoamide concentrations

Synthesis/validation of [15N]-labelled lipoamide

640 [15N]-racemic (±) and (*R*)-(+)-lipoamide were synthesized from racemic and (*R*)-(+)-lipoic acid,
641 respectively, by activating the carboxylic acid using N-hydroxysuccinimide and (1-ethyl-3-(3-
642 dimethylaminopropyl) carbodiimide hydrochloride. The NHS derivative was reacted with ^{15}NH$_4$Cl to
643 incorporate the [15N]-labelling. Full details of the synthesis and subsequent biophysical validation are
644 included in Supplementary Information.

NMR based detection and quantification of [15N]-labelled lipoamide

^{1}H detected ^{15}N edited ^{1}H sensitivity enhanced HSQC NMR ((^{15}N)^{1}H) spectra were acquired on a 14.1 T
Varian Inova spectrometer equipped with a 5 mm z-axis gradient triple resonance room temperature
probe. The free induction decay was recorded for an acquisition time of 0.0624 s and a sweep width
of 8 kHz recorded over 1000 points and a recovery delay of 1 s. Typically, 10000 transients were
collected giving a total experiment time of 3 h 1 min. The J coupling between the amide protons and
the ^{15}N in H$_2$O samples was determined to be 88 Hz, and so the transfer times of 1/4 J in the INEPT
portions of the pulse sequence were set to 2.6 ms. With these settings, ^{15}N ammonia or ammonium
ions would not be detectable. Chemical modification of [15N]-lipoamide (including covalent
attachment to an apoenzyme) would give a substantial change in the (^{15}N)^{1}H NMR spectrum. Similarly,
dissolution of lipoamide in a phospholipid membrane would give substantial peak broadening in the
cell samples. We observed neither, consistent with freely diffusing lipoamide.

Optimisation of the NMR measurement conditions of [15N]-labelled lipoamide

Solvent, pH and temperature sensitivity of the primary amide proton chemical shifts were determined
using dummy samples assembled from the appropriate solvent and added compounds.

Integrated NMR signal intensity is proportional to concentration if provided conditions (temperature
and pH) are identical[75]. Chemical exchange[76], expected as the amide protons in lipoamide should be
labile in water, must also be accounted for. To select appropriate conditions, we determined
temperature (Figure S2F) and pH (Figure S2G) sensitivity of the amide proton signal of 1 mM [15N]-
lipoamide in cell medium. Both amide protons showed chemical exchange under high temperature,
high pH conditions, with the trans-amide proton affected weakly (Figure S2F and S2G). We then
assessed degradation of the trans-amide proton over 10 h (Figure S2H). At 37°C, but not 10°C, the

signal intensity decayed slowly, suggesting slow hydrolysis to form ammonia. We concluded that at
10°C and below pH 8.6 the integrated signal from the trans-amide proton resonance is a good measure
of [15N]-lipoamide concentration.

Quantification of [15N]-lipoamide cellular uptake

Sample preparation workflow

HeLa cells expressing FUS-GFP were grown in 6 well plates to 10^6 cells/well in DMEM supplemented
with 10% FCS. To simultaneously stress and treat cells, the medium was replaced with 0.6 ml medium
supplemented with potassium arsenate and 100 µM [15N]-racemic (±) or (R)-(+)-lipoamide for 1 h at
37°C. High concentrations of compound were used to maximise the signal. The medium was then
removed and retained (medium sample), the cells washed with ~2 ml PBS, then the cells removed by
trypsinisation: addition of 0.3 ml TrypLE Express (12604013, Thermo Fisher Scientific) and incubation
at 37°C for 5 min, then addition of 0.3 ml medium to quench the trypsin. The resuspended cells were
retained (cell sample). All samples were frozen at -80°C. Wells were prepared for all combinations of
no compound (1% DMSO control), [15N]-(±)-lipoamide or [15N]-(R)-(+)-lipoamide, with or without
potassium arsenate and with or without cells.

Calculation of cell volume and uptake

The concentrations of [15N]-lipoamide inside (C_{cell}) and outside (C_{out}) the cells were calculated from
measurements of signal intensity S of the trans-amide proton of lipoamide acquired in the absence
($-cells$, sample i, Figure S2A) and presence ($+cells$, sample ii, Figure S2A) of HeLa cells, using the
following equations (full derivation included in Supplementary Information):

687
$$C_{out} = (1 - U)\frac{c_{add}V_{add}}{V_{out}}$$

688
$$C_{cell} = U\frac{c_{add}V_{add}}{V_1 N_{cell}}$$

689 where $N_{cell} = 10^6$, $c_{add} = 100$ µM and V_{add} (added volume) = 600 µl, $V_1 = 4.19$ x 10^{-15} m^3
690 (approximating HeLa cells as spheres of radius 10^{-5} m) and U represents measured fractional uptake as
691 given   by:

692
$$U = 1 - \frac{S_{+cells}}{S_{-cells}}$$

***In vitro* partitioning of [15N]-(±)-lipoamide in FUS-GFP condensate phase**

Sample preparation workflow

Phase separation of FUS-GFP, at room temperature, was achieved by diluting 12.5 µl of protein stock
at 170 µM concentration (in salty HEPES buffer – 50 mM HEPES, 500 mM KCl and 5% glycerol, at pH
7.25, DTT 1 mM) with 247 µl of salt-free buffer containing [15N]-(±)-lipoamide (50 mM HEPES, 5% D$_2$O,
105 µM [15N]-(±)-lipoamide, 1.05% DMSO, at pH 8), resulting into samples of 260 µl with 8 µM FUS-
GFP, 100 µM [15N]-(±)-lipoamide and 25 mM KCl.

The sample was centrifuged for 10 min at 4000 × *g* and room temperature and the supernatant kept
for NMR analysis. The remaining supernatant was carefully pipetted out, without disturbing the
condensate pellet, and discarded. Perpendicular view photographs of the pellet were taken. Finally,

the condensate pellet was resuspended in 260 μl of buffer with the same buffer conditions as the
phase separated sample (50 mM HEPES, 25 mM KCl, pH ~7.4).

Resuspended condensate or supernatant were loaded in D₂O-matched 5 mm Shigemi tubes and
analysed by (^{15}N)^{1}H NMR. To achieve adequate signal to noise, the resuspended condensate was
scanned for 20 h, while the supernatant was scanned for 4 h. The signal factor (intensity ratio between
the dilute and resuspended condensate samples) was adjusted accounting for differences in sample
volume and number of scans.

Calculation of condensate phase volume

The volume of condensate was calculated from perpendicular photographs (see Fig. S3C), from the
pellet radius (a) and inner radius of the semi-spherical bottom of the microcentrifuge tube (r):

714
$$V = \frac{\pi}{3}r^3(2 + \cos\theta)(1 - \cos\theta)^2$$

715
$$\theta = \sin^{-1}\frac{a}{r}$$

716 θ = subtended angle, as showed in Fig. S3C.

717 Calculation of partition coefficient

718 The ratio between the concentration of lipoamide in the condensate and dilute phases (partition
719 coefficient - PC) was calculated using:

720
$$PC = \frac{V_{Cond} + V_{Added}}{SF * V_{Cond}}$$

721 where V_{Cond} is the volume of condensate phase, V_{Added} is the volume added to resuspend the
722 condensate phase (260 μl) and SF is the signal intensity ratio between the dilute phase and the
723 resuspended condensate measured by NMR.

724 The concentrations of lipoamide in the condensate ([L]$_{Cond}$) and dilute ([L]$_{Dil}$) phases were calculated
725 as:

726
$$[L]_{Cond} = \frac{[L]_{Tot} * V_{Tot}}{V_{Cond} + \frac{V_{Tot} - V_{Cond}}{PC}}$$

727
$$[L]_{Dil} = \frac{[L]_{Cond}}{PC}$$

728 Where [L]$_{Tot}$ is the total concentration of [15N]-($\pm$)-lipoamide and V$_{Tot}$ is the total volume of the phase
729 separated sample. The fraction (%) of [15N]-($\pm$)-lipoamide signal in the condensate phase (see Fig. S3D)
730 was calculated as:

731
$$Lipoamide\ condensate\ signal\ fraction\ (\%) = \frac{1}{1 + SF} * 100$$

The full derivation of these expressions can be found in the Supplementary Information.

Crosslinking coupled to Mass Spectrometry (XL-MS)

FUS condensates were processed and analyzed essentially as described previously[77]. In short,
reconstituted droplets of lysine-rich FUS K12 or FUS G156E were generated by low salt (80 mM KCl)

and crosslinked by addition of H12/D12 DSS (Creative Molecules) in the presence or absence of
lipoamide for 30 min at 37°C, shaking at 600 rpm. Protein samples were quenched by addition of
ammonium bicarbonate to a final concentration of 50 mM and directly evaporated to drynes. The
dried protein samples were denatured in 8 M Urea, reduced by addition of 2.5 mM TCEP at 37°C for
30 min and subsequently alkylated using 5 mM Iodoacetamide at RT for 30 min in the dark. Samples
were digested by addition of 2% (w/w) trypsin (Promega) over night at 37°C after adding 50 mM
ammonium hydrogen carbonate to a final concentration of 1 M urea. Digested peptides were
separated from the solution and retained by a C18 solid phase extraction system (SepPak Vac 1cc tC18
(50 mg cartridges, Waters) and eluted in 50% ACN, 0.1% FA. Dried peptides were reconstituted in 30%
ACN, 0.1% TFA and then separated by size exclusion chromatography on a Superdex 30 increase
3.2/300 (GE Life Science) to enrich for crosslinked peptides. Peptides were subsequently separated on
a PepMap C18 2 µM, 50 µM x 150 mm (Thermo Fisher Scientific) using a gradient of 5 to 35% ACN for
45 min. MS measurement was performed on an Orbitrap Fusion Tribrid mass spectrometer (Thermo
Fisher Scientific) in data dependent acquisition mode with a cycle time of 3 s. The full scan was done
in the Orbitrap with a resolution of 120000, a scan range of 400-1500 m/z, AGC Target 2.0e5 and
injection time of 50 ms. Monoisotopic precursor selection and dynamic exclusion was used for
precursor selection. Only precursor charge states of 3-8 were selected for fragmentation by collision-
induced dissociation (CID) using 35% activation energy. MS2 was carried out in the ion trap in normal
scan range mode, AGC target 1.0e4 and injection time of 35 ms. Data were searched using *xQuest* in
ion-tag mode. Carbamidomethylation (+57.021 Da) was used as a static modification for cysteine.
Crosslinks were quantified relative to the condition containing no lipoamide.

Transfection

Transfection for gene depletion was performed with Lipofectamine2000 (Thermo Fisher Scientific) and
esiRNA oligos targeting human genes (Eupheria Biotech), as listed in Table S1. esiRNA targeting Renilla
luciferase was used for a negative control. The medium was replaced 5 h after transfection, and the
cells were cultured for 3 days before analysis.

Immunocytochemistry of cultured cells

HeLa cells were fixed with 4% paraformaldehyde (PFA) in PBS at room temperature for 15 min, then
washed with PBS containing 30 mM glycine. After permeabilization with 0.1% Triton X-100 in PBS at
4°C and the following wash with glycine-containing PBS, the cells were blocked with 0.2% fish skin
gelatin (Sigma) in PBS (blocking buffer) at room temperature for 20 min, incubated with primary
antibodies in blocking buffer overnight at 4°C, washed with blocking buffer, and incubated with
secondary antibodies and DAPI in blocking buffer at room temperature for 1 h. After washed with
blocking buffer, the samples were stored in PBS until imaging. For detecting endogenous SFPQ, cells
were fixed with cold methanol on ice for 10 min, and blocked with blocking buffer at room temperature
for 20 min before treated with primary antibodies. Samples were imaged on Zeiss LSM 700 or 880
confocal microscopes with a 40× NA 1.2 water objective (Zeiss). Segmentation of nuclei, the cytoplasm,
stress granules, and mitochondria, and measurement of fluorescence intensities at each segment were
performed using CellProfiler. The data were then processed using KNIME to calculate number of stress
granules per cell, nuclear-to-cytoplasmic intensity ratio of stress granule proteins, intensity ratio of the
click-crosslink lipoamide analogue (i) at stress granules, mitochondria, and nuclei over the cytoplasm
(excluding stress granules and mitochondria) or (ii) without crosslinking over with crosslinking, and
percent of cells that have more than 2 stress granules.

iPSC-MNs were fixed for 15 min at room temperature in 4% PFA in PBS. Permeabilization and blocking
was performed simultaneously using 0.1% Triton X-100, 1% BSA and 10% FBS in PBS at room

temperature for 45 min. Subsequently, the primary antibodies were applied overnight at 4 °C in 0.1%
BSA in PBS. The cells were washed with 0.1% BSA in PBS and incubated with secondary antibodies for
1 h at room temperature. Finally, the cells were washed three times with 0.1% BSA in PBS-T (0.005%
Tween-20), including Hoechst or DAPI in the second washing step. Neurofilament H (NF-H) was used
for a marker of MNs. Samples were imaged on either a CellVoyager CV7000 automated spinning disc
confocal microscope (Yokogawa) with a 40× NA 1.3 water objective or a Zeiss LSM880 laser scanning
confocal microscope.

The following primary antibodies were used: rabbit anti-G3BP1 (PA5-29455, Thermo Fisher Scientific);
mouse anti-Tom20 (F-10, Santa Cruz); mouse anti-SPFQ (C23, MBL); mouse anti-SRSF1 (103,
Invitrogen); rabbit anti-TDP-43 (80002-1-RR, Proteintech); mouse anti-Neurofilament H (SMI-32,
Millipore); mouse anti-β3 Tubulin (T5076, Sigma-Aldrich). The secondary antibodies used are as
follows: Alexa Fluor 488-conjugated anti-mouse, Alexa Fluor 594-conjugated anti-rabbit, anti-mouse,
and Alexa Fluor 647-conjugated anti-rabbit and anti-mouse (Thermo Fisher Scientific).

UV cross linking and click reaction

HeLa cells were treated with 3 mM arsenate for 1 h, followed by 30 µM of the click-crosslink lipoamide
analog for 30 min in the presence of arsenate. Then the cells were irradiated with a 305 nm light-
emitting diode for 10 sec for cross linking just before fixation with 4% PFA in PBS at room temperature.
The fixed cells were subjected to immunostaining as described above. After staining, the cells were
subjected to click reaction with 2 µM AF594-Picolyl-Azide (CLK-1296-1, Jena Bioscience) in buffer
containing 100 mM HEPES pH 7.25, 5 mM L-ascorbic acid, 0.5 mM THPTA, and 0.1 mM $CuSO_4$ at 37°C
for 40 min. Cells were then washed three times with 0.1% Triton X-100 in PBS to remove free dye.
Imaging was performed on CSU-W1 (Yokogawa) spinning disk confocal system on an IX83 microscope
(Olympus) with a 100× UPlanSApo 1.4 NA oil objective (Olympus).

Treatment with L-azidohomoalanine (AHA)

Wilde-type HeLa cells were firstly washed with and cultured in methionine (Met)-free medium (Gibco
21013-24) supplemented with 10% FBS for 1 h. Then the medium was replaced with complete medium
or Met-free medium supplemented with 1 mM of Met (Sigma M9625) or AHA (Invitrogen C10102) for
2 h before the cells were stressed with 1 mM arsenate for 1 h. After fixation with 4% PFA in PBS, the
cells were subjected to immunostaining to label G3BP1.

Time-lapse imaging

Time-lapse imaging was performed at 37°C with 5% CO_2. iPSC-derived MNs were treated with 20 µM
lipoamide or 0.02% DMSO (control) for 1 h and then treated with 20 µM arsenite just before image
acquisition. Maximum projection images were generated, and number of FUS-GFP foci was quantified
by Fiji.

Axonal transport assays

AH-ALS1-F58 iPS MNs expressing P525L FUS were treated with 2 µM compound or an equal volume of
DMSO for 3 days. Longer incubation was selected to ensure penetration and action of compounds
along the length of the axon channel. 2 µM was selected as the highest concentration where there
were no toxic effects on this iPS line (assessed qualitatively). Analysis of axonal transport of lysosomes
were performed as previously described[41]. Briefly, lysosomes were labelled by addition of 50 nM
lysotracker red (Thermo Fisher Scirntific) and imaged using a Leica DMI6000 inverted microscope with
a 100× NA 1.46 oil immersion objective and an Andor iXON 897 EMCCD camera in an incubator
chamber (37°C, 5% CO2) at 3 Hz for 120 s at either the proximal or distal end of the silicone channels

harbouring the axons. Kymographs were generated on Fiji. Particle tracking was used to identify
proportion of particles moving faster than 0.2 µm/s for five videomicrographs. Each video includes a
variable population of non-moving background particles, therefore, for each biological replicate, data
were normalised to the mean proportion of moving lysosomes (>0.2 µm/s) at either MFC site (proximal
and distal) in the DMSO (solvent control)-treated FUS P525L samples in Fig. 7D.

Protein aggregation in *C. elegans*

The effect of lipoic acid on stress granule protein aggregation *in vivo* was analysed using a *C. elegans*
model for stress granule formation and aggregation. As previously described[44,78], fluorescent-tagged
PAB-1 forms abundant stress granules and large solid aggregates during aging or upon chronic stress.
RHO-1 also aggregates during aging, but is not an RNA binding or stress granule protein. Two lines
were used: Fluorescently tagged PAB-1 (DCD214: N2; *uqls24[pmyo- 2::tagrfp::pab1gene]*) and RHO-1
(DCD13: N2; *uqls9[pmyo-2::rho-1::tagrfp+ptph-1::gfp]*). Each were analysed as below, except DCD13
were maintained at 20°C.

The animals were exposed to lipoic acid in liquid culture in a 96 well plate starting from larval stage L4
in a total volume of 50 µl S-Complete per well (100 mM NaCl, 50 mM Potassium phosphate pH 6,
10 mM potassium citrate, 3 mM $MgSO_4$, 3 mM $CaCl_2$, 5 µg/mL cholesterol, 50 µM
ethylenediaminetetraacetic acid (EDTA), 25 µM $FeSO_4$, 10 µM $MnCl_2$, 10 µM $ZnSO_4$, 1 µM $CuSO_4$)
supplemented with heat-killed OP50 and 50 µg/ml carbenicillin. Per experiment, a minimum of nine
wells each with 13 animals were treated with R-(+) or S-(−)-lipoic acid or an equivalent volume of
DMSO.

48 h after switching the L4s from 20°C to 25°C (day 2 of adulthood) extensive aggregation of
fluorescently tagged PAB-1 and RHO-1 occurs in the pharyngeal muscles. After immobilization with
2 mM levamisole aggregation was scored using a fluorescent stereo microscope (Leica M165 FC, Plan
Apo 2.0× objective). For PAB-1, aggregates occurred primarily in the terminal bulb of the pharynx, and
aggregation was scored by the number of aggregates (>10 per animal). For RHO-1, aggregates were
scored in the isthmus of the pharynx and aggregation was scored as high (>50% of the isthmus),
medium (<50%) or low (no aggregation). High-magnification images were acquired with a Leica SP8
confocal microscope with a HC Plan Apo CS2 63× NA 1.40 oil objective using a Leica HyD hybrid detector.
tagRFP::PAB-1 was detected using 555 nm as excitation and an emission range from 565-650 nm.
Representative confocal images are displayed as maximum z stack projection.

***D. melanogaster* ALS models**

All fly stocks were maintained on standard cornmeal at 25°C in light/dark controlled incubator. *w[1118]*,
UAS-eGFP, *D42-GAL4*, and *OK6-Gal4* flies were obtained from Bloomington Drosophila Stock Center.
UAS-FUS WT, *UAS-FUS P525L*, and *UAS-FUS R521C* flies were previously described[45,79]. *UAS-TDP-43 WT*
and *UAS-TDP-43 M337V* flies were provided by J. Paul Taylor[80].

Tissue-specific expression of the human genes was performed with the Gal4/UAS-system[81]. Climbing
assays were performed as previously described[79]. Briefly, flies expressing eGFP, human FUS, or TDP-43
were grown in the presence or absence of Lipoic Acid (430 µM, ethanol as the vehicle control) or
Lipoamide (430 µM, DMSO as the vehicle control), then anesthetised, placed into vials and allowed to
acclimatise for 15 min in new vials. Feeding these compounds did not show obvious lethality or toxicity
at this concentration. For each fly genotype, the vial was knocked three times on the base on a bench
and counted the flies climbing up the vial walls. The percentage of flies that climbed 4 cm in 30 s was
recorded. TDP-43-expressing flies were raised at 18°C to suppress lethality.

For immunohistochemistry of neuromuscular junctions, parent flies were crossed on food supplemented with DMSO or lipoamide, and the offspring were raised on the same food. Wandering third instar larvae were dissected and subjected to immunostaining as described previously[82]. Briefly, the dissected larvae were fixed with 4% PFA in PBS at room temperature for 20 min, then washed with PBS. After removing unnecessary tissues, the samples were blocked with 0.2% fish skin gelatin (Sigma) and 0.1% Triton X-100 in PBS (blocking buffer) at room temperature for 1 h, incubated with anti-HRP-Cy3 (1:200, Jackson Immunoresearch) in blocking buffer overnight at 4°C, washed with 0.2% Triton X-100 in PBS (PBT), and incubated with Alexa Fluor 488 Phalloidin (1:5000, Thermo Fisher Scientific) at room temperature for 2 h to visualize muscles. The samples were then washed with PBT and mounted with 70% glycerol in PBS. Synaptic boutons of muscle 4 in abdominal segments 2, 3, and 4 (A2–A4) were imaged using Zeiss LSM 700 or 880 confocal microscopes with a 40× NA 1.2 water objective (Zeiss). Numbers of synaptic boutons and satellite boutons were counted manually.

Quantitative RT-PCR (qPCR)

qPCR was performed with primers targeting GAPDH (control) and full-length STMN2, as described previously[83].

Thermal proteome profiling (TPP)

Thermal proteome profiling was performed as described previously[31]. In brief, two 150 mm dishes of HeLa cells (~6 million cells per dish) were treated with 0.1% (v/v) DMSO (control) or 100 μM lipoamide for 1 h. At the end of incubation one lipoamide- and one DMSO-treated dishes of HeLa cell were stressed with 1 mM arsenate for 1 h. The second set of cells served as the control (treatment with water, vehicle in which arsenate was dissolved) for only lipoamide treatment and only DMSO treatment. All incubations were performed at 37°C with 5% CO_2. Following the incubation, the cells were washed with PBS and trypsinized. The cells were collected by centrifugation at $300 \times g$ for 3 min. The cell pellet was re-suspended in PBS containing the appropriate treatment concentrations of the compounds (lipoamide, DMSO, and arsenate) at cell density of 4×10^6 cells/ml. This cell suspension was split into 10×100 μl aliquots on a PCR plate, spun at $1000 \times g$ for 3min, and finally 80 μl of supernatant (PBS) was subsequently removed. The cell aliquots were then heated to ten to different temperatures (37.0, 40.4, 44.0, 46.9, 49.8, 52.9, 55.5, 58.6, 62.0, and 66.3°C) for 3 min in a thermocycler (SureCycler 8800, Agilent) and left at room temperature for 3 min. Subsequently, the cells were lysed with 30 μl of lysis buffer (PBS containing protease inhibitors, 1.12% NP-40, 2.1 mM $MgCl_2$, and phosphatase inhibitors), and the PCR plates containing the cell lysate was centrifuged at $1000 \times g$ for 5 min to remove cell debris. Next, the heat-induced protein aggregates were removed from the cleared supernatant by passing it through a 0.45 μm 96-well filter plate (Millipore) at $500 \times g$ for 5 min. Equal volumes of the flow-through and 2× sample buffer (180 mM Tris-HCl pH 6.8, 4% SDS, 20% glycerol, and 0.1 g bromophenol blue) were mixed and stored in −20°C until used for mass spectrometry sample preparation. Protein digestion, peptide labelling and mass spectrometry-based proteomics were performed as previously described[32].

Data analysis of TPP

Abundance and thermal stability scores for every protein was calculated as described previously[30,84]. Briefly, the ratio of the normalized tandem mass tag (TMT) reporter ion intensity in each treatment (only lipoamide, only arsenate, lipoamide and arsenate) and the control (only DMSO) was calculated for each temperature. The abundance score for each protein was calculated as the average log_2 fold change (FC) at the two lowest temperatures:

$$Abundance\ score = \frac{\log_2 FC_{37.0°C} + \log_2 FC_{40.4°C}}{2}$$

The thermal stability score for each protein was computed by subtracting the abundance score from the log$_2$ fold changes of all temperatures, and then summing the resulting fold changes (requiring that there were at least ten data points to calculate this score):

$$Thermal\ stability\ score = \sum_T (\log_2 FC_T - Abundanace\ score)$$

where *T* is the ten temperatures. Both the abundance and the thermal stability scores were transformed into a z-distribution by subtracting the mean and dividing by the standard deviation. The significance of the abundance and thermal stability scores was further assesses using *limma*[85] (the two scores were weighted for the number of temperatures in which a protein was identified), followed by FDR estimation using the *fdrtool* package[86]. Proteins with |z-score| > 1.5 and with FDR < 0.05 were considered to be significant changes for the IDR analyses. Thermal stability scores indicated in the result section are those in cells treated with lipoamide and arsenate.

Bioinformatics

Positions of amino acid sequences with disordered tendency were visualized using IUPred3 (https://iupred.elte.hu/). Length of IDRs in each protein was estimated using d2p2 database (https://d2p2.pro/)[87]. The IDR is defined as a region which is regarded as being disordered in more than 75% of all predictors in the database as well as with more than ten successive disordered amino acids. Then proportion of IDR(s) to the whole protein amino acid length was calculated. Enrichment of individual amino acids in IDRs were calculated using Composition Profiler (http://www.cprofiler.org/)[88]. IDRs of all the proteins detected in TPP were used as a background. Positive and negative scores indicate enrichment and depletion of each amino acid compared to the background, respectively.

Statistical analysis

Statistical analyses were performed using the R statistical software or GraphPad Prism. The statistical details (the *p*-value, the number of samples, and the statistical test used) are specified in the figure panels or legends. A *p*-value below 0.05 was considered statistically significant.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE[89] partner repository with the dataset identifiers PXD039670 (for the crosslinking assay) and PXD039501 (for the TPP assay).

Acknowledgements

We thank J.P. Taylor for providing fly strains; the following services and facilities of the MPI-CBG for their support: fly team, LMF, PEPC, TDS, Scientific computing facility, and OSCF; J. Robertson for his advice in the characterization of [15N]-labelled lipoamide; C. Möbius for helping on plate-based cell imaging; C. Höge for valuable comments on the manuscript; members in the Hyman lab and in the MPI-CBG for valuable discussion.

Funding

Hiroyuki Uechi      Uehara Memorial Foundation; JSPS Overseas Research Fellowships; The Osamu Hayaishi Memorial Scholarship for Study Abroad

948	Marcus Jahnel	The Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2068 – 390729961
951	Simon Alberti	European Research Council (grant agreement number 725836)
952	Nicolas L. Fawzi	National Institute of General Medical Sciences (NIGMS) of the National Institutes of Health (R01GM147677) and the National Science Foundation (1845734)
955	Della C. David	Deutsches Zentrum für Neurodegenerative Erkrankungen (DZNE)
956	Andreas Hermann	NOMIS foundation, an unrestricted grant by a family of a deceased ALS patient, the Stiftung zur Förderung der Hochschulmedizin in Dresden and the Hermann und Lilly Schilling-Stiftung für medizinische Forschung im Stifterverband.
960	Florian Stengel	The German Research Foundation (STE 2517/1-1 and STE 2517/5-1)
961	Richard J. Wheeler	Wellcome Trust Sir Henry Wellcome (211075/Z/18/Z) and Sir Henry Dale (103261/Z/13/Z) Fellowships
963	Satoshi Kishigami	Ishizaka Memorial Foundation; Ezoe Memorial Foundation; Shigeta Foundation
965	BGD and AJB	Next Generation Chemistry at the Rosalind Franklin Institute is funded by the EPSRC EP/V011359/1

Conflict of interest statement

Anthony Hyman is the Scientific Founder of Dewpoint Therapeutics; Anthony Hyman and Simon Alberti are Dewpoint Therapeutics shareholders; Richard Wheeler is a Scientific Advisor for Dewpoint Therapeutics; António M. de Jesus Domingues is an employee of Dewpoint Therapeutics, but his contribution was prior to his employment. Anthony Hyman, Mark Bickle and Richard Wheeler filed a patent related to this work (US20200150107A1 and synchronized worldwide applications). Dewpoint Therapeutics contributed intellectually to this work in the structure-activity relationship analysis of lipoamide analogs. All other experimental work either predates foundation of Dewpoint Therapeutics or was carried out independently.

References

1. Robberecht, W. & Philips, T. The changing scene of amyotrophic lateral sclerosis. *Nature Reviews Neuroscience* **14**, 248–264 (2013).

2. Miller, R. G., Mitchell, J. D. & Moore, D. H. Riluzole for amyotrophic lateral sclerosis (ALS)/motor neuron disease (MND). *Cochrane Database Syst. Rev.* (2012). doi:10.1002/14651858.CD001447.PUB3/INFORMATION/EN

3. Abe, K. *et al.* Safety and efficacy of edaravone in well defined patients with amyotrophic lateral sclerosis: a randomised, double-blind, placebo-controlled trial. *Lancet Neurol.* **16**, 505–

512 (2017).

4.   Paganoni, S. *et al.* Trial of Sodium Phenylbutyrate–Taurursodiol for Amyotrophic Lateral
Sclerosis. *N. Engl. J. Med.* **383**, 919–930 (2020).

5.   Deng, H., Gao, K. & Jankovic, J. The role of FUS gene variants in neurodegenerative diseases.
Nature Reviews Neurology **10**, 337–348 (2014).

6.   Lee, E. B., Lee, V. M. Y. & Trojanowski, J. Q. Gains or losses: Molecular mechanisms of TDP43-
mediated neurodegeneration. *Nature Reviews Neuroscience* **13**, 38–50 (2012).

7.   Taylor, J. P., Brown, R. H. & Cleveland, D. W. Decoding ALS: From genes to mechanism.
Nature **539**, 197–206 (2016).

8.   Yamazaki, T. *et al.* Functional Domains of NEAT1 Architectural lncRNA Induce Paraspeckle
Assembly through Phase Separation. *Mol. Cell* **70**, 1038-1053.e7 (2018).

9.   Pessina, F. *et al.* DNA Damage Triggers a New Phase in Neurodegeneration. *Trends in Genetics*
37, 337–354 (2021).

10. Yang, P. *et al.* G3BP1 Is a Tunable Switch that Triggers Phase Separation to Assemble Stress
Granules. *Cell* **181**, 325-345.e28 (2020).

11. Protter, D. S. W. & Parker, R. Principles and Properties of Stress Granules. *Trends Cell Biol.* **26**,
668–679 (2016).

12. Patel, A. *et al.* A Liquid-to-Solid Phase Transition of the ALS Protein FUS Accelerated by
Disease Mutation. *Cell* **162**, 1066–1077 (2015).

13. Molliex, A. *et al.* Phase Separation by Low Complexity Domains Promotes Stress Granule
Assembly and Drives Pathological Fibrillization. *Cell* **163**, 123–133 (2015).

14. Conicella, A. E., Zerze, G. H., Mittal, J. & Fawzi, N. L. ALS Mutations Disrupt Phase Separation
Mediated by α-Helical Structure in the TDP-43 Low-Complexity C-Terminal Domain. *Structure*
24, 1537–1549 (2016).

15. Li, Y. R., King, O. D., Shorter, J. & Gitler, A. D. Stress granules as crucibles of ALS pathogenesis.
J. Cell Biol. **201**, 361–372 (2013).

16. Neumann, M. *et al.* BRAIN A JOURNAL OF NEUROLOGY FET proteins TAF15 and EWS are
selective markers that distinguish FTLD with FUS pathology from amyotrophic lateral sclerosis
with FUS mutations. (2022). doi:10.1093/brain/awr201

17. Sreedharan, J. *et al.* TDP-43 mutations in familial and sporadic amyotrophic lateral sclerosis.
Science **319**, 1668–1672 (2008).

18. Kwiatkowski, T. J. *et al.* Mutations in the FUS/TLS gene on chromosome 16 cause familial
amyotrophic lateral sclerosis. *Science (80-.).* **323**, 1205–1208 (2009).

19. Arseni, D. *et al.* Structure of pathological TDP-43 filaments from ALS with FTLD. *Nature* **601**,
139–143 (2022).

20. Kroschwald, S. *et al.* Promiscuous interactions and protein disaggregases determine the
material state of stress-inducible RNP granules. *Elife* **4**, (2015).

21. Kroschwald, S., Maharana, S. & Simon, A. Hexanediol: a chemical probe to investigate the
material properties of membrane-less compartments. *Matters* (2017).
doi:10.19185/matters.201702000010

22.   Perdikari, T. M., Murthy, A. C. & Fawzi, N. L. Molecular insights into the effect of alkanediols
1026   on FUS liquid-liquid phase separation. *bioRxiv* 2022.05.05.490812 (2022).
1027   doi:10.1101/2022.05.05.490812

1028   23.   Kedersha, N. *et al.* Dynamic Shuttling of TIA-1 Accompanies the Recruitment of mRNA to
1029   Mammalian Stress Granules 7. *J. Cell Biol.* **151**, 1257–1268 (2000).

1030   24.   Petrov, D., Mansfield, C., Moussy, A. & Hermine, O. ALS clinical trials review: 20 years of
1031   failure. Are we any closer to registering a new treatment? *Front. Aging Neurosci.* **9**, 68 (2017).

1032   25.   Krance, S. H. *et al.* Cellular models for discovering prion disease therapeutics: Progress and
1033   challenges. *J. Neurochem.* **153**, 150–172 (2020).

1034   26.   Solmonson, A. & DeBerardinis, R. J. Lipoic acid metabolism and mitochondrial redox
1035   regulation. *J. Biol. Chem.* **293**, 7522–7530 (2018).

1036   27.   Arumugam, S. *et al.* Beneficial effects of edaravone, a novel antioxidant, in rats with dilated
1037   cardiomyopathy. *J. Cell. Mol. Med.* **16**, 2176–2185 (2012).

1038   28.   Yoshida, H. *et al.* Neuroprotective Effects of Edaravone: a Novel Free Radical Scavenger in
1039   Cerebrovascular Injury. *CNS Drug Rev.* **12**, 9–20 (2006).

1040   29.   Becher, I. *et al.* Pervasive Protein Thermal Stability Variation during the Cell Cycle. *Cell* **173**,
1041   1495-1507.e18 (2018).

1042   30.   Mateus, A. *et al.* The functional proteome landscape of Escherichia coli. *Nature* **588**, 473–478
1043   (2020).

1044   31.   Savitski, M. M. *et al.* Tracking cancer drugs in living cells by thermal profiling of the proteome.
1045   *Science (80-.).* **346**, (2014).

1046   32.   Sridharan, S. *et al.* Proteome-wide solubility and thermal stability profiling reveals distinct
1047   regulatory roles for ATP. *Nat. Commun.* **10**, 1–13 (2019).

1048   33.   Lechner, S. *et al.* Chemoproteomic target deconvolution reveals Histone Deacetylases as
1049   targets of (R)-lipoic acid. *Nat. Commun. 2023 141* **14**, 1–10 (2023).

1050   34.   Wang, J. *et al.* A Molecular Grammar Governing the Driving Forces for Phase Separation of
1051   Prion-like RNA Binding Proteins. *Cell* **174**, 688-699.e16 (2018).

1052   35.   Kittler, R. *et al.* Genome-wide resources of endoribonuclease-prepared short interfering RNAs
1053   for specific loss-of-function studies. *Nat. Methods 2007 44* **4**, 337–344 (2007).

1054   36.   Lim, Y. W., James, D., Huang, J. & Lee, M. The emerging role of the rna-binding protein sfpq in
1055   neuronal function and neurodegeneration. *International Journal of Molecular Sciences* **21**, 1–
1056   16 (2020).

1057   37.   Das, S. & Krainer, A. R. Emerging functions of SRSF1, splicing factor and oncoprotein, in RNA
1058   metabolism and cancer. *Molecular Cancer Research* **12**, 1195–1204 (2014).

1059   38.   Kato, M. *et al.* Redox State Controls Phase Separation of the Yeast Ataxin-2 Protein via
1060   Reversible Oxidation of Its Methionine-Rich Low-Complexity Domain. *Cell* **177**, 711-721.e8
1061   (2019).

1062   39.   Ghesquière, B. *et al.* Redox proteomics of protein-bound methionine oxidation. *Mol. Cell.
1063   Proteomics* **10**, (2011).

1064   40.   Van Kasteren, S. I. *et al.* Expanding the diversity of chemical protein modification allows post-
1065   translational mimicry. *Nat. 2007 4467139* **446**, 1105–1109 (2007).

41. Naumann, M. *et al.* Impaired DNA damage response signaling by FUS-NLS mutations leads to
neurodegeneration and FUS aggregate formation. *Nat. Commun.* **9**, (2018).

42. Wang, H. *et al.* Mutant FUS causes DNA ligation defects to inhibit oxidative damage repair in
Amyotrophic Lateral Sclerosis. *Nat. Commun. 2018 91* **9**, 1–18 (2018).

43. Klim, J. R. *et al.* ALS-implicated protein TDP-43 sustains levels of STMN2, a mediator of motor
neuron growth and repair. *Nat. Neurosci.* **22**, 167–179 (2019).

44. Lechler, M. C. *et al.* Reduced Insulin/IGF-1 Signaling Restores the Dynamic Properties of Key
Stress Granule Proteins during Aging. *Cell Rep.* **18**, 454–467 (2017).

45. Lanson, N. A. *et al.* A Drosophila model of FUS-related neurodegeneration reveals genetic
interaction between FUS and TDP-43. *Hum. Mol. Genet.* **20**, 2510–2523 (2011).

46. Uechi, H. *et al.* Ubiquitin-Binding Protein CG5445 Suppresses Aggregation and Cytotoxicity of
Amyotrophic Lateral Sclerosis-Linked TDP-43 in Drosophila. *Mol. Cell. Biol.* **38**, (2018).

47. Coyne, A. N. *et al.* Fragile X protein mitigates TDP-43 toxicity by remodeling RNA granules and
restoring translation. *Hum. Mol. Genet.* **24**, 6886–6898 (2015).

48. Saito, Y. & Kimura, W. Roles of Phase Separation for Cellular Redox Maintenance. *Front.*
Genet. **12**, 691946 (2021).

49. Huang, X. *et al.* ROS regulated reversible protein phase separation synchronizes plant
flowering. *Nat. Chem. Biol. 2021 175* **17**, 549–557 (2021).

50. Mead, R. J., Shan, N., Reiser, H. J., Marshall, F. & Shaw, P. J. Amyotrophic lateral sclerosis: a
neurodegenerative disorder poised for successful therapeutic translation. *Nat. Rev. Drug*
Discov. 2022 223 **22**, 185–212 (2022).

51. Cunha-Oliveira, T. *et al.* Oxidative Stress in Amyotrophic Lateral Sclerosis: Pathophysiology
and Opportunities for Pharmacological Intervention. *Oxid. Med. Cell. Longev.* **2020**, (2020).

52. Motataianu, A., Serban, G., Barcutean, L. & Balasa, R. Oxidative Stress in Amyotrophic Lateral
Sclerosis: Synergy of Genetic and Environmental Factors. *Int. J. Mol. Sci. 2022, Vol. 23, Page*
9339 **23**, 9339 (2022).

53. Kuta, R. *et al.* Depending on the stress, histone deacetylase inhibitors act as heat shock
protein co-inducers in motor neurons and potentiate arimoclomol, exerting neuroprotection
through multiple mechanisms in ALS models. *Cell Stress Chaperones* **25**, 173–191 (2020).

54. Naumann, M. *et al.* Phenotypes and malignancy risk of different FUS mutations in genetic
amyotrophic lateral sclerosis. *Ann. Clin. Transl. Neurol.* **6**, 2384–2394 (2019).

55. Dormann, D. & Haass, C. TDP-43 and FUS: A nuclear affair. *Trends in Neurosciences* **34**, 339–
348 (2011).

56. Dormann, D. *et al.* Arginine methylation next to the PY-NLS modulates Transportin binding
and nuclear import of FUS. *EMBO J.* **31**, 4258–4275 (2012).

57. Amenta, F., Traini, E., Tomassoni, D. & Mignini, F. Pharmacokinetics of Different Formulations
of Tioctic (Alpha-Lipoic) Acid in Healthy Volunteers.
http://dx.doi.org/10.1080/10641960802563568 **30**, 767–775 (2009).

58. Teichert, J., Hermann, R., Ruus, P. & Preiss, R. Plasma Kinetics, Metabolism, and Urinary
Excretion of Alpha-Lipoic Acid following Oral Administration in Healthy Volunteers. *J. Clin.*
Pharmacol. **43**, 1257–1267 (2003).

59. Poser, I. *et al.* BAC TransgeneOmics: A high-throughput method for exploration of protein
function in mammals. *Nat. Methods* **5**, 409–415 (2008).

60. Hein, M. Y. *et al.* A Human Interactome in Three Quantitative Dimensions Organized by
Stoichiometries and Abundances. *Cell* **163**, 712–723 (2015).

61. Cheeseman, I. M. & Desai, A. A combined approach for the localization and tandem affinity
purification of protein complexes from metazoans. *Sci. STKE* **2005**, (2005).

62. Reinhardt, P. *et al.* Genetic correction of a lrrk2 mutation in human iPSCs links parkinsonian
neurodegeneration to ERK-dependent changes in gene expression. *Cell Stem Cell* **12**, 354–367
(2013).

63. Kilpinen, H. *et al.* Common genetic variation drives molecular heterogeneity in human iPSCs.
Nature **546**, 370–375 (2017).

64. Japtok, J. *et al.* Stepwise acquirement of hallmark neuropathology in FUS-ALS iPSC models
depends on mutation type and neuronal aging. *Neurobiol. Dis.* **82**, 420–429 (2015).

65. Higelin, J. *et al.* FUS mislocalization and vulnerability to DNA damage in ALS patients derived
hiPSCs and aging motoneurons. *Front. Cell. Neurosci.* **10**, 290 (2016).

66. Bellmann, J. *et al.* A customizable microfluidic platform for medium-throughput modeling of
neuromuscular circuits. *Biomaterials* **225**, 119537 (2019).

67. Nijssen, J., Aguila, J., Hoogstraaten, R., Kee, N. & Hedlund, E. Axon-Seq Decodes the Motor
Axon Transcriptome and Its Modulation in Response to ALS. *Stem Cell Reports* **11**, 1565–1578
(2018).

68. Pal, A. *et al.* High content organelle trafficking enables disease state profiling as powerful tool
for disease modelling. *Sci. Data 2018 51* **5**, 1–15 (2018).

69. Kreiter, N. *et al.* Age-dependent neurodegeneration and organelle transport deficiencies in
mutant TDP43 patient-derived neurons are independent of TDP43 aggregation. *Neurobiol.
Dis.* **115**, 167–181 (2018).

70. McQuin, C. *et al.* CellProfiler 3.0: Next-generation image processing for biology. *PLoS Biol.* **16**,
e2005970 (2018).

71. Dushek, O. *et al.* Antigen potency and maximal efficacy reveal a mechanism of efficient T cell
activation. *Sci. Signal.* **4**, (2011).

72. Oerter, K. E., Munson, P. J., McBride, W. O. & Rodbard, D. Computerized estimation of size of
nucleic acid fragments using the four-parameter logistic model. *Anal. Biochem.* **189**, 235–243
(1990).

73. Mittasch, M. *et al.* Non-invasive perturbations of intracellular flow reveal physical principles
of cell organization. *Nat. Cell Biol. 2018 203* **20**, 344–351 (2018).

74. Burke, K. A., Janke, A. M., Rhine, C. L. & Fawzi, N. L. Residue-by-Residue View of In Vitro FUS
Granules that Bind the C-Terminal Domain of RNA Polymerase II. *Mol. Cell* **60**, 231–241
(2015).

75. Wider, G. & Dreier, L. Measuring protein concentrations by NMR spectroscopy. *J. Am. Chem.
Soc.* **128**, 2571–2576 (2006).

76. Baldwin, A. J. & Kay, L. E. NMR spectroscopy brings invisible protein states into focus. *Nature
Chemical Biology* **5**, 808–814 (2009).

77.   Boczek, E. E. *et al.* HspB8 prevents aberrant phase transitions of FUS by chaperoning its
folded RNA binding domain. *Elife* **10**, (2021).

78.   Lechler, M. C. & David, D. C. More stressed out with age? Check your RNA granule
aggregation. *Prion* **11**, 313–322 (2017).

79.   Anderson, E. N. *et al.* Traumatic injury induces stress granule formation and enhances motor
dysfunctions in ALS/FTD models. *Hum. Mol. Genet.* **27**, 1366–1381 (2018).

80.   Ritson, G. P. *et al.* TDP-43 mediates degeneration in a novel Drosophila model of disease
caused by mutations in VCP/p97. *J. Neurosci.* **30**, 7729–7739 (2010).

81.   Brand, A. H. & Perrimon, N. Targeted gene expression as a means of altering cell fates and
generating dominant phenotypes. *Development* **118**, 401–415 (1993).

82.   Smith, R. & Taylor, J. P. Dissection and imaging of active zones in the drosophila
neuromuscular junction. *J. Vis. Exp.* e2676 (2011). doi:10.3791/2676

83.   Melamed, Z. *et al.* Premature polyadenylation-mediated loss of stathmin-2 is a hallmark of
TDP-43-dependent neurodegeneration. *Nat. Neurosci.* **22**, 180–190 (2019).

84.   Selkrig, J. *et al.* SARS-CoV-2 infection remodels the host protein thermal stability landscape.
Mol. Syst. Biol. **17**, e10188 (2021).

85.   Ritchie, M. E. *et al.* Limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).

86.   Strimmer, K. fdrtool: A versatile R package for estimating local and tail area-based false
discovery rates. *Bioinformatics* **24**, 1461–1462 (2008).

87.   Oates, M. E. *et al.* D2P2: Database of disordered protein predictions. *Nucleic Acids Res.* **41**,
D508–D516 (2013).

88.   Vacic, V., Uversky, V. N., Dunker, A. K. & Lonardi, S. Composition Profiler: A tool for discovery
and visualization of amino acid composition differences. *BMC Bioinformatics* **8**, 1–7 (2007).

89.   Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass spectrometry-
based proteomics evidences. *Nucleic Acids Res.* **50**, D543–D552 (2022).

1174

A HeLa FUS^WT GFP cells

HeLa cells FUS-GFP

Add compounds → 1600 compounds 10 μM, 1 h

Add arsenate → 1 mM, 1 h

?

Imaging ↓

Measurement of FUS localization

B HeLa FUS^WT GFP cells

Mahalanobis distance (FUS localization)

Mahalanobis cut-off

Lipoamide
Emetine

Lipoic acid

Edaravone

Compounds (ranked)

Condensates remained → Less condensates

C HeLa FUS^WT GFP cells

Arsenate 1 mM

Unstressed | DMSO | Emetine

50 μm

D HeLa cells (stress granule markers)

FUS | G3BP1 | PABPC1 | TIAL1 | EWSR1

Unstressed DMSO

Arsenate 1 mM

DMSO

Lipoamide 10 μM

Lipoic acid 10 μM

20 μm

Figures

Fig. 1. Lipoamide reduces cytoplasmic condensation of stress granule proteins

A, Workflow for screening small molecules for effects on FUS-GFP localisation in HeLa cells *ex vivo*. **B**, Ranked Mahalanobis distances for all 1600 compounds screened (mean from six fields of view) where high values indicate more compound effect. Several automated measures of FUS localisation were combined into a single Mahalanobis distance score; the largest contributors were cytoplasmic FUS condensate number and area (see the method section). A cut-off of 130 was used to select 47 compounds for further analysis. **C**, The sub-cellular localisation of FUS-GFP in unstressed HeLa cells, stressed cells with compound solvent (DMSO) negative control, and with the positive control emetine. Stress causes nuclear export of FUS and formation of stress granules (cytoplasmic liquid FUS-containing condensates). Insets, magnified images in the boxed areas. **D**, Representative images of HeLa cells expressing GFP-tagged stress granule markers (G3BP1, PABPC1, TIAL1, or EWSR1) from 3 independent experiments. The cells were pre-treated with 10 µM lipoamide or lipoic acid (with DMSO solvent control) for 1 h followed by 1 mM arsenate for 1 h, or DMSO without arsenate.

A

Dilute phase
Condensate phase
Lipoamide

dilute

condensate

10.3 ± 1.7×

0.0   0.5   1.0   1.5
[^{15}N lipoamide] (mM)

B

C

DMSO +UV   Analog -UV   Analog +UV

Analog / G3BP1

20 μm

D

Partition coefficient
(each compartment / cytoplasm)

p = 0.0013   p = 0.74   p = 0.91

-UV +UV   -UV +UV   -UV +UV

Stress granule   Mitochondria   Nuclei

E

Signal ratio (-UV/+UV)

Cytoplasm
Stress granules
Mitochondria
Nuclei

Fig. 2. Lipoamide partitions into compartments of stress granule proteins

A, Top, schema of lipoamide partitioning into FUS condensates *in vitro*. Bottom, mean ± s.e.m. of concentration of racemic [15N]-lipoamide in the condensate and the surrounding dilute phase of FUS-GFP *in vitro*, quantified using ^{15}N(^{1}H) NMR from 4 independent experiments. **B**, Chemical structure of the click-crosslink lipoamide analog, with the lipoamide backbone (orange) and the groups for UV cross linking and click reaction (green). **C**, Representative images of HeLa cells treated with 3 mM arsenate for 1 h followed by 30 μM of the analogue or the control DMSO in the presence of arsenate for additional 30 min before either irradiated with UV for cross-linking (+UV) or not (-UV), fixed, immunostained, and subjected to the click reaction with the fluorophore (for all conditions). Stress granules were labelled with G3BP1. Insets, stress granules in the boxed areas (analogue and G3BP1 boxed in green and magenta, respectively). **D**, Boxplot of the partition coefficient of the analogue into stress granules, mitochondria, or nuclei relative to the cytoplasm (excluding stress granules and mitochondria) based on signal intensity of the fluorophore. Boxplot shows median (bold bar), 25th and 75th percentiles, and outliers (open dots); whiskers extend to the most extreme values. *n* = 344 (-UV) and 345 (+UV) cells from 3 experiments. *p* values by unpaired *t*-test. **E**, Mean ± s.d. of signal intensity ratio (-UV against +UV) of the fluorophore at indicated subcellular compartments.

A

carboxamide — backbone — chiral centre — dithiolane — disulfide

HeLa FUS GFP

iPSC FUS GFP

Stress granules per cell / Nucleus/cytoplasm partition vs Lipoamide (µM)

for D, E, F & I

B stereoisomer

stereoisomer	Lipoamide EC$_{50}$ (µM)	Lipoic acid EC$_{50}$ (µM)
Racemic-(±)	13.9	17.0
R-(+)	12.1	~30
S-(−)	14.0	23.1

C amide methylation   EC$_{50}$ (µM)

	EC$_{50}$ (µM)
R-(+)-lipoamide	12.1
Monomethylated	3.48
Dimethylated	>30

D secondary amides   R = / EC$_{50}$ (µM)

EC$_{50}$ (µM)
9.36
13.3
19.5

E   R = / EC$_{50}$ (µM)

EC$_{50}$ (µM)
2.12
2.23
2.28
2.43
2.59
2.94
2.94

F   R = / EC$_{50}$ (µM)

EC$_{50}$ (µM)
7.72
28.4
5.12
2.39

G backbone length   EC$_{50}$ (µM)

EC$_{50}$ (µM)
13.9
3.48
3.58

H carboxamide replacement   EC$_{50}$ (µM)

EC$_{50}$ (µM)
1.07

I most potent   R = / EC$_{50}$ (µM)

EC$_{50}$ (µM)
1.20
1.30
1.30
1.41
1.44
1.63
1.72
1.85
1.93
2.01

J dithiolane modifications   EC$_{50}$ (µM)

EC$_{50}$ (µM)
19.8
14.9
>30
>30
>30

 Fig. 3. Structure-activity relationship shows lipoamide activity is dependent on the dithiol but is non-enzymatic

 A, Top, schema of the chemical structure of lipoamide (racemic) with highlighting its features. Bottom, lipoamide dose
 response using HeLa and iPS cells, showing FUS-GFP condensate (stress granule) number (solid circles, left axis) and
 nuclear/cytoplasmic signal ratio (open circles, right axis) with 1 h pre-treatment with lipoamide followed by 1 h
 arsenate stress under continued lipoamide treatment. **B-J,** Chemical structures and EC_{50} (μM) of lipoamide and its
 derivatives, using HeLa cells and the treatment scheme in A. EC_{50} was calculated from dose response curves (details in
 the method section), and each concentration of each compound was tested in duplicated wells (n = 1750–2650 cells
 per well) with 2 independent experiments. **B**, Enantiomers of lipoamide and lipoic acid. **C**, Comparison of mono- and
 di-methylated lipoamide. **D–F**, Additional carboxamide analogs of lipoamide. **G**, Modifications of the linker length
 between the carboxamide and the dithiolane ring of lipoamide. **H**, Substitution of the carboxamide of lipoamide. **I**,
 Carboxamide analogs of 6-amino-3-substituted-4-quinazolinones and five-membered aminoheterocyclic amides. **J**,
 Modifications of the dithiolane ring of lipoamide, lipoic acid or similar compounds.

Fig. 4. Lipoamide interacts with disordered proteins in cells

A, Schema of TPP to see the effect of lipoamide treatment on protein thermal stability. In our TPP, HeLa cells treated with 100 μM lipoamide and/or 1 mM arsenate were heated to ten different temperatures. Heating causes protein denaturing and precipitation, and lipoamide could prevent precipitation of the target proteins. The soluble protein amount was quantified by mass spectrometry and normalized with the result of non-stressed 0.1% DMSO-treated samples (control). **B**, Volcano plot of z-scores (mean from 3 individual experiments) and FDRs of protein thermal stability in HeLa cells treated with lipoamide and arsenate. A larger z-score indicates more thermal stabilization. Black broken and solid lines indicate cutoffs of z-score (±1.5) and FDR (<0.05), respectively, used to classify stabilized (green) and destabilized (blue) proteins in F and G. The positions of DLD, SFPQ, SRSF1, and several stress granule proteins are indicated. **C**, Violin and box plots showing proportions of IDRs in each protein from cells treated with both lipoamide and arsenate. The proteins were categorized into stabilised (z > 1.5, FDR < 0.05; 70 proteins), destabilised (z < −1.5, FDR < 0.05; 144 proteins), and unaffected (the others detected; 5811 proteins). Boxplots show median (bold bar), 25th and 75th percentiles, and outliers (closed dots); whiskers extend to the most extreme values. *p* values by a Wilcoxon signed-rank test followed by Holm's test. **D**, Enrichment (> 0) or depletion (< 0) of each amino acid in IDRs of the stabilized (green) and destabilized (blue) proteins in cells treated with both lipoamide and arsenate, in comparison to IDRs of all the detected proteins as background. *p* values by unpaired *t*-test followed by Bonferroni's test. **E**, Representative images of HeLa cells from >3 independent experiments, depleted of SFPQ or SRSF1, treated with 10 μM lipoamide or 0.1% DMSO for 1 h followed by 1 mM arsenate for 1 h in the presence of lipoamide. Stress granules were labelled with G3BP1. **F**, Mean ± s.d. of percentage of stressed HeLa cells with ≥3 G3BP1-positive stress granules (SGs). *n* = 292–615 cells from 3 independent experiments. Dots indicate means of each experiment. *p* values by Tukey's test. **G**, Representative images of HeLa cells from >3 independent experiments, treated with 1 mM arsenate for 1 h and subjected to immunostaning with indicated antibodies. Outer and inner broken lines indicate edges of the cytoplasm and nucleus of one cell each condition, respectively. Note that the SFPQ and SRSF1 signals were diminished by individual RNAis.

A

Met

Cys

(707 a.a.)

B

L-Methionine
(Met)

L-Azidohomoalanine
(AHA)

C

Arsenate 1 mM

RNAi

Control

SFPQ

Complete | -Met +Met | -Met +AHA

20 µm

$p = 9 \times 10^{-7}$

$p < 2 \times 10^{-6}$ $p > 0.1$

% cells with ≥3 SGs

RNAi   Control   *SFPQ*

D

Reduced state

SFPQ

Lipoamide

Stress granules

SFPQ

Oxidized state

Fig. 5. SFPQ redox state may mediate the lipoamide activity

A, Schema of distributions of methionine (Met; 28 residues) and cysteine (Cys; 2) residues in human SFPQ. PLD, prion-
like domain; RRM, RNA recognition motif; NOPS, NonA/paraspeckle domain; NLS, nuclear localizing signal. **B**, Chemical
structures of Met and its non-natural analogue azidohomoalanine (AHA). **C**, Left, representative images of HeLa cells
subjected to indicated RNAis, cultured in complete medium (light grey) or Met-free medium supplemented with 1 mM
of Met (dark grey) or AHA (green) for 2 h followed by 1 mM arsenate for 1 h (experimental schema in Fig. S6E). Stress
granules (SGs) were labelled with G3BP1. Right, mean ± s.d. of percentage of stressed HeLa cells with ≥3 G3BP1-
positive SGs. n = 325–407 cells from 3 independent experiments. p values by Tukey's test. **D**, Schema of SFPQ as a
redox sensor to modulate stress granule condensation.

Fig. 6. Lipoamide improves nuclear localization of FUS and TDP-43

A, Representative images of iPSC-derived MNs from 3 independent experiments, treated with 0.1% DMSO or 10 μM lipoamide for 1 day followed by 10 μM arsenite for 5 days in the presence or absence of lipoamide, and labelled with TDP-43. Broken line, outline of nuclei. **B**, mean ± s.e.m of nuclear TDP-43 levels normalized to those of unstressed DMSO-treated MNs (control). *n* = 417–1741 cells from 3 independent experiments. *p*-values, Tukey's test. **C**, (Left) images showing recruitment of FUS-GFP to sites of UV laser-induced DNA damage (yellow arrow) in nuclei (outlined with broken lines) of iPS cells at indicated times after laser irradiation. Cells were subjected after 1 h treatment with lipoamide followed by 1 h arsenate stress. (Right) mean ± s.d. of relative FUS-GFP signal intensity in response to DNA damage. *n* = 5 (DMSO) and 7 (lipoamide) cells. **D**, (Left) images of nuclei (outlined with broken lines) of iPSC-derived MNs expressing FUS P525L-GFP from 3 independent experiments, cultured for 21 days and then treated with 0.02 % DMSO or 20 μM lipoamide for 24 h, at indicated times after laser irradiation. Yellow lines indicate laser-irradiated sites. (Right) mean ± s.e.m. of relative intensity of FUS-GFP at DNA damage sites after ablation. *n* = 14 (DMSO) and 18 (lipoamide) cells from 3 independent experiments. **E**, Mean ± s.d. of relative STMN2 full length mRNA levels normalized to those of GAPDH from 2 independent experiments. *P*-values by Tukey's test. In B and E, MNs were treated as in A.

Fig. 7. Lipoamide improves cellular fitness in ALS disease models of iPSC-derived motor neurons and animals

A, (Left) representative images of iPSC-derived MNs treated as in Fig.5A. (Right) mean ± s.e.m of percentage of neurite (tubulin-positive) area. 18 image fields from 3 independent experiments. *P*-values, Tukey's test. **B**, Schematic of neuron culture, showing the channels through which the axons grow from the soma on the right. **C**, Kymographs of lysosome movement in the distal portion of FUS P525L MN axons 3 days after treatment with compound solvent (DMSO) or 2 μM lipoamide, visualized with lysotracker. **D**, mean ± s.e.m. of relative proportion of lysotracker-labelled lysosomes moving with an average speed greater than 0.2 μm/s following 3 days treatment with 2 μM lipoamide or equivalent DMSO concentration solvent control for iPSC-derived MNs expressing either P525L or WT FUS, normalized to mean of proportion moving (proximal and distal) in the DMSO-treated P525L FUS MNs. 6 (P525L) or 3 (wild-type) biological replicates, analyzing 5 axon bundles per replicate. *p* values by Tukey's test. **E**, (Left) Representative images of the pharynx of worms expressing fluorescently tagged PAB-1 with or without lipoic acid treatment (2 mM). Broken lines, the edge of pharynges. (Right) Mean ± s.e.m. of incidence of each protein aggregation in the pharyngeal muscles. Incidence of PAB-1 aggregation was scored from the proportion of animals with >10 aggregates. ***$p < 0.0001$ by Fisher's exact test. *n* > 100 for each sample. **F**, Mean ± s.e.m. of proportion of flies that climbed, fed with 0.1% DMSO (solvent control) or 430 μM lipoamide. Human WT or ALS-linked mutants of FUS (left) or TDP-43 (right) were expressed in motor neurons. *p* values by unpaired *t*-test. *n* = 30–40 (FUS) and 130–202 (TDP-43) flies from 3 or 4 independent experiments.

A *in vitro* proteins

FUS-GFP

Add compounds → 47 *in vivo* hits 1 to 100 μM

Reduce salt → ↓ KCl to 50 mM

Robotic mixing & plate transfer

Imaging

Droplet number, signal partition

B

Droplet number Z score at 100 μM

Z score cut-off

In/out signal partition Z score at 100 μM

Z score cut-off

Compounds (ranked)

● Lipoamide   ● Surfactants
● Heterotri-/tetra-cyclic   ● Other compounds

C

DMSO | Cetylpyridinium chloride | Lipoamide | Mitoxantrone

20 μm

D HeLa cells

% cells with ≥3 SGs

Time (min)

● DMSO
● Lipoamide
● Lipoic acid
(both 10 μM)

E HeLa cells (other condensates)

P body — DCP1A | Nuclear compartment — COIL | TRP53BP1

DMSO

Lipoamide 10 μM

20 μm

F HeLa FUS WT-GFP cells (under different stresses)

Solvent control | Arsenate 1 h 1 mM | Sorbitol 1 h 0.4 M | Heat stress 30 m 42°C | 6-deoxy-glucose 1 h 100 mM

DMSO

Lipoamide 10 μM

Lipoic acid 10 μM

20 μm

G HeLa cells

Arsenate 1 mM

DMSO

Lipoamide 10 μM

Ascorbic acid
10 μM | 100 μM | 1 mM

% cells with ≥3 SGs

DMSO | Lipoamide | 10 μM 100 μM 1 mM Ascorbic acid

20 μm

Supplemental figures

Fig. S1. *In vitro* **follow-up screening and lipoamide characterization in HeLa cells**

A, Workflow for screening small molecules for effects on FUS condensation of purified FUS-GFP *in vitro*. **B,** Ranked Z scores of change in condensate droplet number and signal partition into FUS-GFP droplets (formed under low salt conditions) where larger positive or negative values mean more compound effect. Scores were calculated at the maximum concentration at which the compound solvent (DMSO) negative control had no significant effect; 100 µM. Lipoamide, surfactant and heterotri-/tetracyclic compounds are indicated by data point colour. **C,** Appearance of the droplets with compound solvent (DMSO) negative control or examples of compound classes: cetylpyridinium chloride (surfactant), lipoamide or mitoxantrone (heterotricyclic). Note the larger drops with cetylpyridinium chloride and lipoamide and the fewer smaller drops with mitoxantrone. **D,** Mean ± s.d. of percentage of HeLa cells with ≥ 3 G3BP1-positive stress granules (SGs). Cells were treated with 1 mM arsenate for 1 h to induce SGs, followed by 10 µM lipoamide, lipoic acid, or 0.1% DMSO (control) in the presence of arsenate for indicated minutes. *n* = 52–248 cells from 3 independent experiments. **E,** Images of HeLa cells expressing GFP-tagged markers of other membrane-less organelle compartments subjected to 1 h treatment with 10 µM lipoamide (or DMSO control). Where unclear, the position of nuclei is indicated with a broken outline. Lipoamide does not disrupt P bodies (DCP1A), Cajal bodies (COIL), or DNA damage foci (TRP53BP1). **F,** Images of HeLa cells expressing FUS-GFP subjected to different stresses – arsenate, sorbitol (osmotic), heat, or 6-deoxyglucose (glycolysis) – with concurrent treatment with 10 µM lipoamide or lipoic acid. **G,** Representative images of HeLa cells treated with 1 mM arsenate for 1 h, followed by 0.1% DMSO (control), 10 µM lipoamide, or indicated concentrations of ascorbic acid for 15 min. SGs were labelled with G3BP1.

 Fig. S2. Tracking cellular uptake of [15N]-Lipoamide using NMR

A, Methodology for quantitation of [15N]-lipoamide uptake by HeLa cells, using the trans-amide proton to measure
[15N]-lipoamide concentration (see F- H). Medium with 100 μM [15N]-lipoamide was incubated for 1 h in the absence
or presence of HeLa cells. Following removal of medium, the cells were washed with medium (without arsenate) and
detached using EDTA-trypsin. Solution or cell pellet/in-cell NMR was used to determine [15N]-lipoamide concentration.
Example spectra for cells stressed with 3 mM arsenate and incubated with R-(+)-lipoamide are shown with the same y
axis scale. **B,** Cellular uptake was determined by subtracting signal from medium incubated with cells (red) from signal
from medium without cells (blue). This was carried out for all four combinations of stressed (3 mM arsenate) or
unstressed cells with [15N]-(R)-(+) or (±)-lipoamide. For stressed cells treated with [15N]-(R)-(+)-lipoamide the high
signal intensity from the washed cell sample (green) is consistent with the large uptake from the medium calculated
from the with (red) and without cell (blue) signal intensity. **C,** Quantitation of B showing percentage uptake and
calculated intracellular concentration, assuming that lipoamide is uniformly distributed within cells (see Supplemental
Methods). Uncertainty in measurement was approximately 30% and there was no significant difference in uptake
between conditions. All measurements indicated substantial uptake of lipoamide and cellular concentrations >1 mM.
D, Overview of synthesis of [15N]-lipoamide, highlighting the trans amide proton (14). **E,** [1]H NMR spectrum of [15N]-
lipoamide in CDCl$_3$. Peaks can be unambiguously assigned to individual proton environments. F-H, Controls determining
reliability of quantitation of [15N]-lipoamide using the amide protons in [15]N edited [1]H NMR experiments. **F,**
Dependency of the cis (13) and trans (14) amide proton signal on temperature, at a constant pH of 8.3. Both resonances
decreased with increasing temperature, indicating local molecular dynamics and/or interactions with H$_2$O on ms to μs
timescale reduce the signal. Trans amide proton resonance approaches a plateau towards 10°C. **G,** Dependency of the
cis and trans amide proton signal on pH, at a constant temperature of 10°C. Together, indicating at 10°C and below
pH 8.6 integrated signal intensity of the trans-amide proton of lipoamide in [15]N edited [1]H NMR experiments is a reliable
proxy for concentration. **H,** Signal intensity of the trans-amide proton of lipoamide, when dissolved in growth medium,
decreased over time at 37°C but not at 10°C. At 10°C signal intensity is stable for >10 h experiments.

A

centrifugation

condensate phase
dilute phase

calculate volume

supernatant (dilute phase)

resuspended pellet (diluted condensate phase)

$(^{15}N)^{1}H$ NMR

B

Supernatant
Resuspended Condensate * Signal Factor

NMR intensity (au)

$_1H\ \omega$ (ppm)

8   7.8   7.6   7.4   7.2   7   6.8   6.6

C

Supernatant

Condensate Pellet

r = Tube inner radius
h = Condensate pellet height
a = Condensate pellet radius
θ = angle subtended ($\sin^{-1}(a/r)$)

D

i)

Percentage (%)

^{15}N lipoamide condensate signal fraction

Condensate volume fraction

ii)

Partition Coefficient

E

DMSO   Lipoamide 10 µM   Analog

5   10   20   µM

G3BP1

20 µm

F

Merge   Analog   G3BP1   Tom-20   DNA

20 µm

Fig. S3. Experimental set up for portioning assays of lipoamide and its analogue

A-D, Methodology for determination of partition of [15N]-lipoamide into FUS condensates *in vitro.* (A) Schematic
showing the sample preparation process. (B) Example of ^{15}N edited ^{1}H NMR signal around the ^{15}N cis and trans amide
protons for a dilute phase and condensate phase. Condensate phase spectrum is shown scaled by an experimentally
determined signal factor, used in calculation of the signal fraction. (C) Measurements for calculation of condensate
pellet volume from macro photographs of the sample within a microcentrifuge tube. (D) Mean ± s.d. of measured ^{15}N
edited ^{1}H NMR signal fraction and condensate volume fraction (from 4 independent experiments) and calculated
partition coefficient. Alternative presentation of the data in Fig. 2A. **E**, Representative images of HeLa cells pre-treated
with indicated concentrations of lipoamide or the click-crosslink lipoamide analog in Fig. 2B for 1 h followed by 1 mM
of arsenate for additional 1 h in the presence of compounds. SGs were labelled with G3BP1. **F**, The images of HeLa cells
treated with the analog and subjected to arsenate treatment and UV cross-linking from Fig. 2C, with a channel of Tom-
20 as a mitochondrial marker.

Fig. S4. Lipoamide weakly increases liquidity of FUS condensates *in vitro*

A, NMR chemical shift deviations per residue for the FUS N-terminal PLD (residues 1 to 163) with 500 µM lipoamide compared to the drug solvent control (1% DMSO). Light grey bars indicate tyrosine residues and residues neighbouring a tyrosine. **B,** Average ^{1}H and ^{15}N shifts across residues zero, one, two, three or more than three residues from a tyrosine in the presence of lipoamide. **C**, Top, mean ± s.d. of fractions of FUS proteins condensed at indicated salt (KCl) concentrations in the presence of 300 µM lipoamide or the DMSO control (0.3% v/v). n = 16 image fields. Bottom, dilute phase concentrations (equivalent to saturation concentrations) of FUS-GFP at 150 mM KCl at different temperatures and lipoamide concentrations (errors are s.d.) **D**, Schematic illustrating the quantitation of condensate droplet liquidity using optical tweezers. Two droplets are brought into contact and begin to fuse: the time taken to relax to a single spherical droplet (once adjusted for the geometric mean radius as the characteristic droplet size) is a measure of the viscosity to surface tension ratio of the droplet – a proxy of liquidity. **E**, Droplet size-corrected relaxation times for droplet fusions with either 300 µM lipoamide (n = 93 independent fusion event) or equivalent DMSO solvent control (0.3%, n = 60). Box represents the 25th, 50th and 75th percentiles, whiskers represent 5th and 95th percentiles. p value by unpaired t-test. Lipoamide reduces fusion time, indicating lower viscosity and/or greater surface tension. **F– H**, Effect of 30 µM lipoamide or lipoic acid on FUS G156E-GFP condensates 'aging', relative to an equivalent DMSO solvent control (0.3%). Condensates were formed under 50 mM of KCl while shaking. (F) Representative images after 1 and 3 h aging, showing fibre formation in the DMSO sample in contrast to the lipoamide or lipoic acid samples. (G) Representative fluorescence recovery after photobleaching (FRAP) time series of FUS condensates and fibres at corresponding time points. (H) Mean ± s.d. of relative intensity of FUS-GFP FRAP in (G). Aged (3 h) condensates treated with lipoamide or lipoic acid maintain large FUS-GFP mobile fraction. Both compounds delay fibre formation. **I**, Changes in intramolecular crosslinking due to lipoamide of FUS in *in vitro* low salt (80 mM KCl) condensates using the lysine-rich FUS K12 or FUS G156E. Significantly changed crosslinking sites with a change in intensity of more than two-fold and FDR ≤ 0.1; 3 independent experiments) are shown coloured in green (increased) or red (decreased). Other crosslinking sites are shown in grey. **J,** Dose-dependent effect of lipoamide on FUS K12, plotting absolute change in crosslink intensity relative to no lipoamide. Crosslinking sites with false discovery rate (FDR) > 0.1 are shown in blue, those with FDR ≤ 0.1 in orange (2 independent experiments). Two-fold change is indicated with a dashed red line.

Fig. S5. SFPQ and SRSF1 are cellular targets of lipoamide not necessary for stress granule formation

A, Z-scores of protein thermal stability in HeLa cells treated with only lipoamide and both lipoamide and arsenate. Proteins categorized as stabilized and destabilized in Fig. 4C,D are depicted in green and blue, respectively. The positions of FUS, TAF15, DLD, SFPQ, and SRSF1 are indicated. Black solid and broken lines indicate cutoffs of z-scores used for the IDR analysis in Fig. 4C (±1.5) and the targeted RNAi screen (+2), respectively. **B**, Z-scores of protein thermal stability in HeLa cells treated with only arsenate and both lipoamide and arsenate. Black lines indicate |z-score| = 1.5. In most proteins with Increased or decreased thermal stability by only arsenate treatment (|z-score [arsenate]| > 1.5), the shifts were prevented by lipoamide pre-treatment (|z-score [arsenate + lipoamide]| < 1.5; masked in orange). Proteins categorized in stabilized and destabilized in Fig. 4C,D are depicted in the same colours; note that shifts in their thermal stability was not primarily due to treatment with arsenate. **C**, Mean ± s.d. of percentage of HeLa cells with ≥ 3 G3BP1-positive stress granules. Cells depleted of indicated genes were treated with 10 µM lipoamide or 0.1% DMSO for 1 h followed by 1 mM arsenate for 1 h in the presence of lipoamide before stained with G3BP1. *n* = 324–393 cells from 3 independent experiments. *p* values by Tukey's test. **D**, Domain compositions and distributions of IDRs of human SFPQ (left) and SRSF1 (right). PLD, prion-like domain; RRM, RNA recognition motif; NOPS, NonA/paraspeckle domain; CC, coiled-coil domain; NLS, nuclear localizing signal; G-rich, glycine-enriched domain; SR-rich, serine/arginine-enriched domain. **E**, Mean ± s.d. of percentage of cells with ≥3 stress granules. HeLa cells depleted of indicated genes were treated with 3 mM arsenate for 1 h, and then with 100 µM lipoamide or the control DMSO in the presence of arsenate for indicated minutes. *n* = 213–467 cells from 3 independent experiments. **F**, Representative images of HeLa cells treated and labeled as in Fig. 4E but without arsenate.

A

H₂O₂ (%) 0   0.01   0.03   0.1

SFPQ-GFP

FUS-GFP

10 µm

B

H₂O₂ (%) 0   0.01 0.03 0.1 0.3   kDa

SFPQ

—80

—65

C

SFPQ-mGFP added (µM)

0   10   30   50

FUS-SNAP

SFPQ-GFP

10 µm

Fraction FUS-SNAP condensed

1
0.8
0.6
0.4
0.2
0

○ GFP
● SFPQ-GFP

0 10 20 30 40 50
(SFPQ-)GFP (µM)

D

H₂O₂ (%) 0   0.01   0.03   0.1

GFP

SFPQ-GFP

Added (40 µM)

10 µm

E

Met   Each medium   Arsenate (1 mM)

                                              Fixation

1 h   2 h   1 h

Complete   [SFPQ][Met]   [SFPQ][Met]=O

-Met +Met   [SFPQ][Met]   [SFPQ][Met]=O

-Met +AHA   [SFPQ][AHA]   [SFPQ][AHA]

 Fig. S6. SFPQ dissolves FUS condensates *in vitro*

A, Representative images of SFPQ-GFP (top, 10 µM) and FUS-GFP (bottom, 7 µM) protein condensates at a low salt condition (75 mM KCl) and in the presence of H_2O_2 from >3 independent experiments. **B**, SDS-PAGE (non-reduced condition) of 10 µM of the purified and untagged SFPQ proteins in diluted state oxidized with the indicated percentages of H_2O_2 for 30 min. **C**, Left, representative images of co-incubation of indicated concentrations of SFPQ-GFP and 6 µM of FUS-SNAP at a physiological salt concentration (150 mM KCl) from >3 independent experiments. SFPQ proteins do not form condensates at 150 mM KCl while they suppress condensation of FUS proteins in dosage-dependent manner. Right, mean ± s.d. of FUS condensate fraction in the presence of GFP (control) or SFPQ-GFP. *n* = 16 image fields. **D**, Representative images of FUS-SNAP condensates (4 µM) co-incubated with 40 µM of GFP or SFPQ-GFP at a physiological salt concentration (150 mM KCl) in the presence of indicated percentages of H_2O_2 from 3 independent experiments. **E**, Schema of the time course used in Fig. 5C. Cells were firstly cultured in methionine (Met)-free medium and then in each medium (complete medium or Met-free medium supplemented with Met or AHA) before arsenate treatment.

A **HeLa cells**

Nucleus/cytoplasm partition

$p = 0.029$ — FUS^{WT}

$p = 0.046$ — FUS^{P525L}

$p = 0.039$ — TDP-43

■ DMSO
■ Lipoamide 10 µM

B **iPSC-derived FUS^{P525L}-GFP MNs**

Arsenite 20 µM

DMSO

Lipoamide 20 µM

0   20   30   (min)

10 µm

FUS P525L foci per cell

DMSO
Lipoamide

$p = 0.014$

Time (min)

C **iPSC-derived FUS^{P525L}-GFP MNs**

DMSO

Lipoamide 20 µM

10 µm

$p = 0.0086$

Nuclear FUS^{P525L} intensity (a.u.; normalized)

■ DMSO
■ Lipoamide

 Fig. S7. Lipoamide rescues nuclear localisation and functioning of ALS-linked proteins

A, Mean ± s.d. of nuclear-cytoplasmic intensity ratio of FUS and TDP-43 in HeLa cells pre-treated with 10 µM lipoamide
(0.1% DMSO as the control) for 1 h followed by 1 mM arsenate for 1 h in the presence of lipoamide. n = 290–603 cells
from 3 independent experiments. p value by unpaired t-test. **B**, (Left) time-lapse images of iPSC-derived MNs
expressing FUS P525L-GFP cultured for 14 days. Cells were treated with 0.02% DMSO or 20 µM lipoamide for 1 h
followed by 20 µM arsenite for indicated minutes in the presence of lipoamide. Broken lines indicate outline of some
nuclei. Arrowheads indicate some cytoplasmic FUS P525L foci. (Right) mean ± s.e.m. of number of FUS P525L foci per
MN after arsenite treatment. n = 16 (DMSO) and 18 (lipoamide) cells from 3 independent experiments. p value by
unpaired t-test. **C**, (Left) representative images of iPSC-derived MNs expressing FUS P525L-GFP cultured for 5 days and
then 30 days in the presence of 0.02% DMSO or 20 µM lipoamide. Broken lines indicate outline of some nuclei. (Right)
mean ± s.e.m. of nuclear intensity of FUS P525L-GFP, normalized to that in the control (DMSO). n = 64–198 cells from
5 independent experiments. p value by one-sample t-test.

A *C. elegans* **pharynx**

RHO-1

Proportion showing aggregation

100%
50%
0%

DMSO

R-(+)-Lipoic acid 1.5 mM

B *D. melanogaster* **locomotion**

Proportion climbing 4 cm in 30 s

100%
50%
0%

■ Ethanol
■ Lipoic acid 430 μM

$p = 0.18$

$p = 0.044$

$p = 0.0065$

eGFP   WT FUS   FUS P525L   FUS R521C

C *D. melanogaster* **neuromuscular junctions**

HRP (neuronal membrane)

DMSO

Lipoamide 430 μM

10 μm

$p = 1.1 \times 10^{-4}$ $p = 8.1 \times 10^{-4}$

$p = 0.34$

% Satelite boutons

■ DMSO
■ Lipoamide 430 μM

Control   TDP-43 M337V

Fig. S8. Extended analysis of *C. elegans* and *D. melanogaster* animal models of ALS

A, Mean ± s.e.m. of incidence of each protein aggregation in the pharyngeal muscles. Incidence of RHO-1 was scored on a low, medium, high scale (see methods). **B,** Mean ± s.e.m. of percentage of flies that climbed, with lipoic acid feeding in place of lipoamide in Fig. 6F. *p* values by unpaired *t*-test. **C,** (Left) Representative images of synaptic boutons of TDP-43 M337V-expressing flies, immunostained with an antibody against horseradish peroxidase (HRP), which labels the neuronal membrane. Arrowheads indicate appearance of satellite boutons. (Right) mean ± s.d. of percentage of satellite boutons (number of satellite boutons/number of total boutons) per fly. The control flies fed with 0.1% DMSO (grey; *n* = 9) and TDP-43 M337V-expressing flies fed with 0.1% DMSO (*n* = 19) or that containing 430 µM lipoamide (orange; *n* = 19) were examined. *p* value by Tukey's test.