Potential Neurocognitive Biomarkers for Post Traumatic Stress Disorder (PTSD) Severity in Recent Trauma Survivors

Short Running Title: Potential Neurocognitive Biomarkers for PTSD Severity

Ziv Ben-Zion, M.Sc.[1,2], Yoav Zeevi, M.Sc.[2,3,], Nimrod Jackob Keynan, Ph.D.[1,4], Roee Admon, Ph.D.[5], Tal Kozlovski, M.Sc.[3,6], Haggai Sharon, M.D., Ph.D.[1,7,8,9], Pinchas Halpern, M.D., Ph.D.[7,10], Israel Liberzon, M.D., Ph.D.[11], Arieh Y. Shalev, M.D., Ph.D.[12], Yoav Benjamini, Ph.D.[2,3*] & Talma Hendler[1,2,4,7*],M.D., Ph.D.

*equal contribution to senior authorship for the last two authors

[1]Sagol Brain Institute Tel-Aviv, Wohl Institute for Advanced Imaging, Tel Aviv Sourasky Medical Center, Tel-Aviv, Israel

[2]Sagol School of Neuroscience, Tel-Aviv University, Tel-Aviv, Israel

[3]Department of Statistics and Operations Research, Tel-Aviv University, Israel;

[4]School of Psychological Sciences, Faculty of Social Sciences, Tel-Aviv University, Tel-Aviv, Israel

[5]Department of Psychology, University of Haifa, Haifa, Israel

[6]Laboratory of Early Markers of Neurodegeneration (LEMON), Department of Neurology, Tel-Aviv Sourasky Medical Center, Tel-Aviv, Israel

[7]Sackler Faculty of Medicine, Tel-Aviv University, Tel-Aviv, Israel

[8]Institute of Pain Medicine, Department of Anesthesiology and Critical Care Medicine, Tel Aviv Sourasky Medical Center, Tel-Aviv, Israel

[9]Pain Management & Neuromodulation Centre, Guy's & St Thomas' NHS Foundation Trust, London, UK

[10]Department of Emergency Medicine, Tel Aviv Sourasky Medical Center, Tel-Aviv, Israel

[11]Department of Psychiatry, Texas A&M Health Science Center, TX, USA

[12]Department of Psychiatry, NYU Langone Medical Center, New York, NY, USA

Abstract

Contemporary symptom-based diagnosis of Post-traumatic Stress Disorder (PTSD) largely overlooks related neurobehavioral findings and rely entirely on subjective interpersonal reporting. Previous studies associating objective biomarkers with PTSD have mostly used the disorder's symptom-based diagnosis as main outcome measure, overlooking the actual clustering and richness of phenotypical features associated with PTSD. Here, we aimed to computationally derive potential neurocognitive biomarkers that could efficiently differentiate PTSD subtypes, based on an observational cohort study of recent trauma survivors. A three-staged semi-unsupervised method ("3C") was used to *categorize* trauma survivors based on current PTSD diagnostics, derive *clusters* of PTSD based on features related to symptom load, and to *classify* participants' cluster membership using objective features. A total of 256 features were extracted from psychometrics, cognitive, structural and functional neuroimaging data, obtained from 101 adult civilians (age=34.80±11.95, 51 females) evaluated within a month of trauma exposure. Multi-domain features that best differentiated cluster membership were indicated by using importance analysis, classification trees, and ANOVA. Results revealed that entorhinal and rostral anterior cingulate cortices volumes (structural

domain), in-task amygdala's functional connectivity with the insula and thalamus (functional domain), executive function and cognitive flexibility (cognitive domain) best differentiated between two clusters related to PTSD severity. Cross-validation established the results' robustness and consistency within this sample. Multi-domain biomarkers revealed by the 3C analytics offer objective classifiers of post-traumatic morbidity shortly following trauma. They also map onto previously documented neurobehavioral PTSD features, supporting the future use of standardized and objective measurements to more precisely identify psychopathology subgroups shortly after trauma.

<u>Introduction</u>

Post-traumatic stress disorder (PTSD) symptoms are commonly observed shortly

after exposure to trauma, with the extent of their initial intensity associated with a high

risk for poor-recovery[1–3]. Categorization is currently based on symptoms reported via

structured interviews, such as the Clinician-Administered PTSD Scale (CAPS). While

such scales show good reliability, construct and predictive validity, they also have

notable limitations. The CAPS does not capture various symptoms often co-expressed

with PTSD, such as depression and general anxiety[4]. It is also only weakly linked with

cognitive measures and other objective measures of putative biological mechanisms[5–7].

By solely accounting for reported- or observed symptoms, the CAPS entirely rely on

subjective interpersonal reporting. These limitations may be responsible for its long term

diagnoses instability [8] as well as its suboptimal use as a guide for individualized clinical

management[9].

Cognitive functioning is one of many dimensions often overlooked in the clinical

context of post-traumatic psychopathology. To date, reported associations between

cognitive deficits and PTSD included working memory, information processing speed,

verbal learning, short-term and declarative memory ,attention and executive functioning,

response inhibition and attentional switching (see studies reported in recent meta-

analyses[10,11]). Adaptive cognitive functioning has been linked with resilience and a

reduced likelihood of post-traumatic stress symptoms development and maintenance[11,12].

The investigation of *brain structure and function* are also crucial for understanding the

underlying clinical manifestations of PTSD. Indeed, accumulating evidence from

neuroimaging points to structural and functional brain abnormalities among PTSD

patients[5,13–16], including lower hippocampal volume[17–19] and altered activity and

connectivity of the amygdala, insula, ACC, mPFC and dlPFC; structures known to be

involved in threat detection, executive function, emotion regulation, and contextual

processing[5,13,20–24]. Some structural and functional neuroimaging studies also point to

early predisposing factors in the development of PTSD[13,21,25–27]. Despite promising

findings from cognitive and neuroimaging studies, these objective measures have yet to

be integrated in the routine assessment and management of post-traumatic

psychopathology. One obstacle in the clinical translation of these findings is an unclear

understanding of how these indicators cluster together into PTSD subtype to provide a

better guide for potential interventions. Moreover, most studies associating objective

biomarkers with PTSD have used the disorder's symptom-based formal diagnostic

criteria as main outcome measure, thereby overlooking the actual clustering and richness

of phenotypical features associated with post-traumatic psychopathology within a dataset,

and imposing an extraneous construct (PTSD diagnosis) as an outcome of interest.

The above-mentioned limitations could benefit from advanced data-driven

computational and statistical methods that can evaluate wider arrays of potential

biomarkers and disorder indicators, quantifying their relationship with various clinical

manifestations. Machine learning methods are particularly well-suited to confront such

computational challenges, as they are able to account for the complex interrelation of

many relevant factors[28]. Indeed, the last decade has shown an exponential increase in the

use of machine learning for the study of posttraumatic stress, including both supervised

and unsupervised approaches. Supervised analytic approaches are limited by the accuracy

of the prior knowledge they rely on[29]. While unsupervised approaches make few to no

assumptions, this limits the subpopulations revealed, as they are not tied to any specific

questions of interest[29].

A possible solution is the use of hybrid analytic methods that combine both

supervised and unsupervised approaches, which may lead to more precise diagnosis by

identifying novel combinations of potential biomarkers for specific disorders[29]. The

present study propose to evaluate objective biomarkers' ability to identify data-driven

clusters, which hopefully represent post-traumatic psychopathology phenotypes. To do

so, we applied a recently developed three-staged hybrid analytic methodology, termed 3C

(Categorize, Cluster, and Classify); a semi-unsupervised method that combines theory-

and data-driven approaches[30]. Analysis included both clinical wide-used knowledge

assumptions and state-of-the-art data-driven methods apply on objective measures

obtained from neuroimaging and cognitive testing of recent trauma survivors. We

assumed that such a hybrid analytic approach would unveil a unique set of potential

mechanism-related neurocognitive biomarkers for PTSD, closely tied to pre-existing

diagnostic methods. The term "subtypes" accounts for different demographics, clinical

sub-scales or symptom severity.

Materials and Methods

The 3C methodology was used on a multi-domain data set composed of clinical

interviews and questionnaires, computerized cognitive testing and neuroimaging of brain

structure and function. Data was obtained from recent trauma survivors discharged from

a general hospital's emergency department (ED) after experiencing a traumatic event.

This data is a part of a larger on-going project that examines PTSD development in

trauma survivors (for full study protocol please see Ben-Zion et al. (2019)[31]). Here we

present multi-domain dataset obtained from participants within one-month of the

traumatic incident.

Participants. 101 recent trauma survivors (age=34.80±11.95, range 18-65, 51 females)

admitted to ED following a traumatic experience. The most common trauma types were

motor-vehicle accidents (n=79, 78%), and other traumatic events included assaults, terror

attacks, hostilities, drowning, mass casualty incidents, robbery and electrocution. Out of

101 participants, n=58 individuals met a full PTSD symptom diagnosis during the face-

to-face clinical interview ("PTSD" group), and n=43 did not meet this criterion even

though they still suffered from a variety of PTSD symptoms ("No PTSD" group).

Clinical Instruments. PTSD symptom severity was quantified using the Clinician-

Administered PTSD Scale for DSM-IV (CAPS-4)[32]. Four additional self-report clinical

questionnaires were administered: PTSD Checklist for DSM-IV (PCL-4)[33] evaluating

post-traumatic symptoms; Beck Depression Inventory (BDI)[34] assessing current

depressive symptoms; Anxiety Inventory (BAI)[35] measuring concurrent anxiety

symptoms; and Participants' Clinical Global Impression Scale (CGI-P)[36] evaluating

patients' subjective impression, on a scale between 1 ('normal feeling') to 7 ('the worst

feeling there is'). For more details on these clinical instruments, please see[31].

Cognitive Functioning. WebNeuro[37], an internet-based comprehensive cognitive

assessment battery previously validated against traditional cognitive tests, was used to

assess cognitive functioning. To standardize testing conditions, all tests were conducted

at our laboratory in Hebrew. Performance on the different tasks was calculated using an

automated software program that derived standardized Z-scores for each participant on

each of the following eleven cognitive domains: motor coordination, processing speed,

sustained attention, controlled attention, cognitive flexibility, response inhibition,

working memory, recall memory, executive function, emotion identification, and

emotional bias. For detailed description of the cognitive tasks, please see[12,31].

Imaging Data Acquisition. Structural and functional scans were performed in a 3.0

Tesla Siemens MRI system (MAGNETOM Prisma, Germany), using a twenty-channel

head coil, located in our lab at Tel-Aviv Sourasky Medical Center. To allow high-

resolution whole-brain structural images, a T1-weighted magnetization prepared rapid

gradient echo (MPRAGE) (TR/TE=2400/2.29ms, flip angle = 8°, voxel size $0.7 \times 0.7 \times$

0.7 mm, FOV = 224×224 mm) was acquired. Functional whole-brain scan was

performed in an interleaved order, using a T2*-weighted gradient echoplanar imaging

pulse sequence (TR/TE = 2000/28 ms, flip angle = 90°, voxel size $2.2 \times 2.2 \times 2.2$ mm,

FOV =220×220 mm, slice thickness = 3mm, 36 slices per volume).

Structural Imaging Data Analysis. Cortical reconstruction and volumetric segmentation

were performed with the FreeSurfer image analysis suite[38] (version 1.379.2.73), which is

documented   and   freely   available   for   download   online

(http://surfer.nmr.mgh.harvard.edu/). Briefly, this processing included motion correction

and the averaging[39] of multiple volumetric T1 weighted images, removal of non-brain

tissue using a hybrid watershed/surface deformation procedure[40], automated Talairach

transformation, segmentation of the subcortical white matter and deep gray matter

volumetric   structures   (including   hippocampus,   amygdala,   caudate,   putamen,

ventricles)[41,42], intensity normalization, tessellation of the gray matter-white matter

boundary, automated topology correction, and surface deformation following intensity

gradients to optimally place the gray/white and gray/cerebrospinal fluid borders at the

location where the greatest shift in intensity defines the transition to the other tissue class.

The automatic subcortical segmentation of brain volume is based upon the existence of

an atlas ("aseg") containing probabilistic information on the location of structures[43]. The

maps are created using spatial intensity gradients across tissue classes and are therefore

not simply reliant on absolute signal intensity. The maps produced are not restricted to

the voxel resolution of the original data thus are capable of detecting submillimeter

differences between groups.

Functional Imaging Data Analysis. Preprocessing and statistical analysis of the

functional images were performed in a voxel-based approach using Statistical Parametric

Mapping (SPM)[44] version 12. In short, this process included: Slice time correction, using

one slice before the last as the reference slice. Head motions correction by six-parameter

rigid body spatial transformations, using three translations and three rotation parameters,

with the first image serving as a volume reference. A 4^{th} degree interpolation was applied

to detect and correct head motions. Functional maps were automatically co-registered to

corresponding structural maps using an objective function of normalized mutual

information (NMI). The complete dataset was transformed into MNI space and spatially

smoothed with an isotropic 6mm full-width at half-maximum (FWHM) Gaussian kernel.

During this scan, participants performed the **Emotional Faces Matching Task**[45]**,**

which was used to evaluate their emotional reactivity. In this task, subjects were

instructed to select the face/shape (located at the bottom right or bottom left of the

screen) that matches the target face/shape (located at the top of the screen), as accurately

and as quickly as possible. The tasks included four blocks of shapes (that were used as a

baseline) and four blocks of emotional faces (angry, fearful, surprised and neutral faces).

The order of the blocks of emotional faces was counterbalanced between subjects using

four different versions for this task. Both whole-brain activations and functional

connectivity of the amygdala (seed region) were calculated for the following contrasts:

angry faces (vs. shapes), fearful faces (vs. shapes), surprised faces (vs. shapes), and

neutral faces (vs. shapes). For a full list of the functional brain measures derived from

this analysis, please refer to 'brain function variables' in Supplementary Table 1.

Procedure. A member of the research team identified potentially trauma-exposed

patients using the ED medical records. Within 10–14 days after potential trauma

exposure, and after being discharged from the hospital, these individuals were contacted

for an initial telephone screening, which was conducted by MA-level clinicians that were

trained in the specific assessment tools. After obtaining verbal consent, the PCL-5 was

administered to assess the risk of PTSD development. Those who met PTSD symptom

criteria (except the "one-month duration" criteria) and did not meet any of the exclusion

criteria (see under Participants section), received verbal information about the study.

Participants were subsequently invited to participate in two meetings, both within 30 days

from ER admission: (1) In-person clinical assessment including administration of the

CAPS, self-report questionnaires (BDI, BAI, PCL, CGI) and WebNeuro cognitive

battery; (2) Structural and functional MRI scan. Each meeting lasted approximately 3

hours (total of 6 hours for both), for which participants received financial remuneration,

in accordance with the ethics committee regulations and approval.

Statistical Approach. The 3C method[30] assumes that existing medical knowledge of a

given disorder is critical but not sufficient for an accurate diagnosis. It suggests to build

upon the current diagnostics and expands upon it with unsupervised data-driven methods.

Thus, using well-used clinical measures to divide patients into homogeneous clusters

based on common characteristics. Clustering allows exploration of objective measures

related to the specific disorder-subtype. A recent study using the 3C[46] discovered new

sub-phenotypic groups and their specific biomarkers in a large Alzheimer's disease

dataset ("ADNI"[47]). The 3C finds homogeneous groups via unsupervised clustering using

well-used clinical measures, relevant to the disease diagnosis, it then further characterizes

those groups by exploring potential biomarkers. The 3C procedure is composed of three

stages; Categorize, Cluster and Classify**.** During Categorization, multi-domain variables

are sorted into three categories; 1) *Assigned diagnosis*; as currently applied in the field. 2)

Clinical measurement; variables that describe the patient's condition and the expression

of the disease (symptoms and signs). 3) *Potential biomarkers*; variables that could

improve existing diagnostic procedures, but not currently in use in the diagnostic

procedure. Clustering includes two steps: 1) A supervised selection of the most relevant

clinical variables through some statistical testing, to be used as predictors to the *assigned*

diagnosis. This feature selection was done based on a permissive threshold of Benjamini-

Hochberg[48] FDR-adjusted of p=0.2 (From now on, the use of "FDR" in this manuscript

will specifically refer to the BH procedure). This was done to eliminate clinical variables

that are not relevant to the disease as defined by *the assigned diagnosis* category. 2)

Unsupervised clustering utilizing the selected *clinical measurements* was performed,

using k-medoids with Manhattan distance metrics. This allowed us to discover data-

driven homogeneous clusters that are related to existing diagnostics (i.e., dividing

participants into subtypes based on common–used variables). Nevertheless, these clusters

are not limited to formal symptom-based PTSD diagnosis (as indicated by CAPS), but

rather capture the actual richness of phenotypical features associated with post-traumatic

psychopathology within a dataset. Prior to clustering we determined the optimal number

of clusters based on two metrics: Gap statistics[49] and Silhouette[50]. Lastly, <u>Classification</u>

includes characterization of the clusters based on the objective *potential biomarkers,* via

three distinct approaches: importance analysis (mean decrease GINI[51]); classification

trees; and a marginal one-way analysis of variance (ANOVA) between clusters for each

potential bio-marker.

Algorithms, Codes, and Software. Algorithms and codes were performed using R

software version 3.4.4. Data imputation was performed using the 5-nn method in order to

deal with missing data (less than 1% of the full dataset). Variables were monotonic

transformed to gain symmetry when needed, using a semi-automated Shiny App[52].

Importance, measured as the marginal loss of classification accuracy for each variable by

randomly permuting it on the out-of-bag validation set, was calculated using the

{randomForest} R package[51]. R package {cluster}[53] was used for clustering, and R

package {rpart} was used for classification and regression trees (CART).

<u>**Results**</u>

During <u>Categorization</u>, features were divided into three distinct categories

required by the 3C method: *Assigned Diagnosis* was based on the CAPS-4 total scores,

Clinical Measurements included the total scores of the four self-report questionnaires

(PCL, BDI, BAI, and CGI), and *Potential Biomarkers* included 11 standardized total

scores obtained from computerized cognitive testing, 192 features from structural

imaging (including volumes and thickness of subcortical and cortical areas), 48 features

extracted from fMRI during the emotional faces matching task (including whole-brain

activations and functional connectivity of left and right amygdala during the task). For a

full list of neurocognitive features, see Supplementary Table 1.

214   For <u>Clustering</u>, we used the *Clinical Measurements* that were found highly

correlated with PTSD symptom severity as indicated by CAPS-4 total scores: PCL, BDI,

BAI, and CGI (based on the 0.2 FDR threshold). Participants were divided into an

optimal number of two clusters, based on both Gap statistics[49] and Silhouette[54] methods,

which presented the best separation on all four clinical measurements (PCL, BDI, BAI,

and CGI). Participants within the first cluster showed an average higher severity on all

four clinical measurements, compared to those within the second cluster (see Fig. 1). To

test the association between these clusters (representing high and low "disease load") and

PTSD diagnosis (PTSD or not according to CAPS-4), a two-sample test for equality of

PTSD proportions between the two clusters was conducted. Results revealed a significant

link between the proposed clusters in PTSD dichotomous diagnosis *($F_{1,99}=35.47$,*

$p<0.001$, $\chi^2_{1}=20.911$, $p<0.001$, CI of the difference of proportions = (0.28,0.63)) (see

Table 1). Furthermore, a one-way ANOVA was performed on the CAPS-4 total scores

(continuous PTSD symptom severity measure) with clusters as the group variable.

Results indicated that individuals in cluster 2 had significantly higher CAPS-4 total

scores *(M=61.91±18.16)* compared to those in cluster 1 *(M=37.45±23.12)(p<0.001)* (see

Supplementary Figure 1). It is important to note that these clusters do not represent a

simple division into high and low CAPS severity scores, but rather a data-driven division

into severity subtypes, based on self-reported symptoms of depression (BDI), anxiety

(BAI), post-trauma (PCL) and the patients' subjective global impression (CGI). As

mentioned, these subtypes (clusters) were found to correlate with PTSD clinical diagnosis

and severity, but were not identical to it, and rather represent clusters of "disease load".

Accordingly, cluster 1 will now be referred to as the "Low-Symptomatic" Cluster

(LoClus, low PTSD severity), and cluster 2 as the "High-Symptomatic" Cluster (HiClus,

high PTSD severity).

During the Classification stage, clusters were based on objective variables that

could serve as potential biomarkers, using a mean decrease GINI index (i.e. importance

index) for each biomarker. The most significant potential neurocognitive biomarkers

associated with the resulted clusters included left EC volume *(Importance=0.884),*

cognitive flexibility *(Importance=0.487),* rostral anterior cingulate cortex (rACC) volume

(Importance=0.429), and average amygdala functional connectivity with the thalamus

while watching angry faces vs. shapes (Importance=0.419). The top ten potential

neurocognitive biomarkers for the clustering are presented in Figure 2 and

Supplementary Table 2.

To further characterize patients within each cluster according to the identified

biomarkers, we built classification trees (see Fig. 3). Results showed that left hemisphere

EC volume had the greatest influence on clustering - 70 out of 101 participants had left

EC volume equal to or greater than 1449mm^3, out of which there was almost an equal

distribution between the two clusters (56%/44%). The other 31 participants had a left EC

volume smaller than 1449mm^3, out of which the vast majority (84%) belonged to HiClus;

indicating that subjects with lower EC volume were more likely to belong to the HiClus.

Further down on the left branch of the tree, HiClus subjects had larger left caudal middle

frontal gyri volume. Down the right branch of the tree, high executive functioning was

more associated with LoClus, and vice versa. Further down, low supramarginal gyrus

cortical thickness together with high paracentral volume were related to LoClus, while

low executive functions with low functional connectivity between the amygdala and the

left insula while watching fearful faces vs. shapes was strongly related to HiClus.

Finally, a marginal one-way ANOVA was conducted on the potential biomarkers

with HiClus/LoClus as the dependent variable. Benjamini-Hochberg adjustment to

control the FDR at level 0.05, yielded no significant difference for any potential

neurocognitive biomarker between HiClus/LoClus, FDR-corrected (partly due to

hundreds of p-values that were adjusted).

To assess robustness and consistency within this sample, the procedure detailed

above was repeated with cross-validation. The 3C method was performed on a different

percentage of the total subjects (P=20%, 30%, …, 90% of all participants). For each

percentage of subjects (P), n=1000 iterations were performed. In each iteration, the 3C

methodology was executed on a randomly selected subset of individuals, resulting in

different clusters and classification tress. The remaining individuals (which were not

included in this iteration), were classified to the new clusters based on the created

decision trees. For each P, the mean percentage (and SD) of individuals correctly

classified (according to the results of the 3C methodology based on all subjects, see Table

2). For example, when the 3C algorithm was based on only P=20% of the subjects, 83%

of the remaining individuals were correctly classified on average.

Discussion

This work illustrates the efficacy of a novel 3-staged (3C) data analytic approach

in classifying recent trauma survivors to different "disease load" using a set of potential

neurocognitive biomarkers. The 3C 'hybrid' data- and theory-driven analytic approach

combined current diagnostic-based *categorization*, symptom severity-based *clustering*

and the data-driven retrieval of *classifying* using *potential biomarkers*. Unlike supervised

machine learning models that 'flatten' information and dimensions into one data matrix,

the 3C method uniquely and in a stepwise manner combines current PTSD diagnostics

with two layers of data-driven exploration: clinical symptom severity and concurrently-

recorded objective measurements of cognitive functioning, brain structure and function. It

does not impose an extraneous construct (PTSD diagnosis) as an outcome of interest, but

rather focus on the actual clustering and richness of phenotypical features associated with

post-traumatic psychopathology, thereby may able to create symptom clusters that more

reliably represent phenotypical subclasses of theoretical and clinical interest. The

biomarkers revealed by the 3C approach are in line with previously documented neural

and cognitive correlates of PTSD. Thus, they provide a template for the early objective

mechanism-based categorization of trauma survivor psychopathology.

Potential neurocognitive biomarkers for PTSD

The potential neurocognitive biomarkers for PTSD severity included features

obtained from both neuroimaging and cognitive testing. These are not yet proven PTSD

biomarkers, as they did not underwent analytical and clinical validations[55], but rather

serve as potential biomarkers from neurocognitive domains. The structural brain features

that had the most influence on PTSD classification was entorhinal cortex (EC) volume

(both according to importance analysis and classification trees, see Fig. 2 and 3); lower

EC volume was associated with higher post-traumatic stress symptom severity. The EC

plays an important role in memory, a key feature of post-traumatic psychopathology, as

uncontrolled recall of the traumatic event determines symptom severity[56–58]. Another

structural feature of importance to classification was the volume of the rostral anterior

cingulate cortex (rACC); lower rACC volume was associated with higher PTSD severity.

Indeed, rACC volume has previously been associated with PTSD[21,59,60], and shown to

predict cognitive behavioral treatment response for individuals suffering from this

disorde[61]; suggesting its potential as a guide for early mechanism-based intervention.

To note, our classifier did not identify several structural abnormalities found in

previous cross-sectional PTSD studies[5,13,21,25]. This includes the most replicated finding

of small hippocampus volume[62], but also abnormal amygdala volume, insular cortex,

medial and dorsal prefrontal cortices (mPFC and dlPFC respectively)[16,19,20,63–65]. This

could stem from our classifier identifying early stage severity-related (or "disease load")

biomarkers, rather than DSM-based categories of PTSD diagnosis. Our approach

identified structural abnormalities within one-month after trauma as related to symptom

severity; since major changes in gray matter volume within this time frame are less likely

to occur[66], these abnormalities might be early predisposing risk factors for chronic PTSD

development. Future studies on populations prone to trauma exposure with longitudinal

measurement could shed more light on the causal inference of our findings[67–69].

Functional neuroimaging found amygdala functional connectivity with both the

insula and the thalamus to be particularly important for classification (both according to

importance analysis and classification trees, see Fig. 2 and 3). Aberrant connectivity of

the amygdala with other structures is consistent with previous studies, and abnormal

amygdala activation had been hypothesized to contribute to PTSD pathophysiology[70–72].

Moreover, thalamic dysfunction has been found in patients with PTSD, suggesting its

role in the disorder's psychopathology[73,74]. Therefore, although neuroimaging studies

have implicated several functional brain abnormalities in the pathophysiology of PTSD,

our computational analysis showed that some of these abnormalities are involved in the

PTSD severity subtype at the early aftermath of the trauma.

The most significant cognitive factors were cognitive flexibility (according to

importance analysis, see Fig. 2) and executive function (according to classification trees,

see Fig. 3). Indeed, meta-analyses regarding the role of cognitive functions in PTSD

consistently show an impaired ability in executive functioning, including cognitive

flexibility among PTSD patients[75,76]. Cognitive flexibility; the ability to switch between

two different tasks or strategies[75] is of particular interest as it has recently been found to

correlate with PTSD trajectories in two independent longitudinal cohorts[12]. Specifically,

lower scores of cognitive flexibility correlated with increased PTSD symptoms one

month after trauma and predicted persistent symptoms one year later. Moreover,

disturbances in cognitive flexibility was ameliorated by a cognitive intervention and

associated with better treatment outcomes[12], implying its role as an early recovery related

process. It is indeed possible that intact cognitive flexibility enables the individual to

better differentiate between threat-related and neutral situations, thus assisting in the

extinction of fear-motivated learning, a core-element in PTSD recovery[77].

Methodological considerations

Our 3C approach revealed two PTSD subtypes (classes) in recent trauma

survivors, correlated with high and low clinical severity, according to total CAPS scores,

and across all CAPS subscales (re-experiencing, avoidance, negative alterations in

cognitions and mood, and alternations in arousal and reactivity). Our analysis did not find

classes representing different clinical subtypes (e.g. dissociative subtype, greater

avoidance, etc.). This could be because 3C clustering is based on a given set of clinical

measures, and classification is thus based on predetermined measures of potential

biomarkers. A larger dataset (i.e., more measures and more participants), could allow for

the identification of unique clinical classes of different symptomatic subtypes (possibly

more than two). Indeed, in an effort to account for the heterogeneity in PTSD expression,

several studies attempted to characterize different subtypes of the disorder. For example,

an externalizing subtype characterized by low constraint and high negative emotionality,

compared to an internalizing cluster with high negative emotionality and low positive

emotionality[78–80]. Another example is a dissociative subtype for patients with PTSD and

de-personalization and/or de-realization symptoms, introduced by the fifth edition of the

DSM[81–83]. Note that there is no definitive ad-hoc threshold for selection of clinical

measurements, different measures strongly efffect the clusters created and the derived

classification of potential biomarkers. Lastly, adding longitudinal measures from

different time-points following trauma may reveal classes corresponding to PTSD clinical

trajectories. This may be cruical for the identification of individuals at risk for developing

PTSD, as well as providing appropriate early-stage treatment.

Conclusion

Our study implemented an innovative computational approach that unveiled novel

variables correlated with the morbidity classification of recent trauma survivors,

associated with the mechanisms underlying post-traumatic stress symptoms. The method

utilized current DSM-based PTSD diagnostic categories and other clinical severity

measures of depression and anxiety, as well as a data-driven classification of

neurocognitive potential biomarkers. Our results point to an alternative approach for

identifying objective variables linked to PTSD severity subtypes (high and low), based on

testing within a single session closely after exposure to trauma. If successful, this

objective computational classification of potential neurocognitive biomarkers may further

guide mechanism-driven interventions for PTSD (e.g. cognitive remediation or

neuromodulation treatments). If performed on a larger data set and with more clinical

measures and potential biomarkeres, this computational approach may further refine post-

traumatic diagnostic subtypes, playing an important role in the treatment management of

recent trauma survivors.

Acknowledgments: This work was supported by award number R01-MH-103287 from the National Institute of Mental Health (NIMH) given to AS (PI), IL and TH (co-Investigators, subcontractors), and had undergone critical review by the NIMH Adult Psychopathology and Disorders of Aging study section. This work was also supported by funding to the Human Brain Project from the European Union Seventh Framework Program (FP7/2007-2013) under grant agreement no. 604102 (HBP). Sagol School of Neuroscience at Tel-Aviv University and the HBP supported ZB fellowships, and Sagol Brain Institute at Tel-Aviv Sourasky Medical Center supported NK and HS fellowships.

The authors would like to thank the research team at Tel-Aviv Sourasky Medical Center - including Nili Green, Mor Halevi, Sheli Luvton, Yael Shavit, Olga Nevenchannaya, Iris Rashap, Efrat Routledge and Ophir Leshets - for their major contribution in carrying out this research, including subjects' recruitment and screening, and performing clinical, cognitive and neural assessments. We also extend our gratitude to all the participants of this study, which completed all the assessments at three different time-points after experiencing a traumatic event.

Conflict of Interest: The authors declare that they have no financial disclosures and no conflict of interests.

Trial Registration: Neurobehavioral Moderators of Post-traumatic Disease Trajectories. ClinicalTrials.gov registration number: NCT03756545.

https://clinicaltrials.gov/ct2/show/NCT03756545

<u>References</u>

1.   Galatzer-Levy IR, Karstoft KI, Statnikov A, Shalev AY. Quantitative forecasting

of PTSD from early trauma responses: A Machine Learning application. *J*

Psychiatr Res. 2014;59:68-76. doi:10.1016/j.jpsychires.2014.08.017

2.   Kessler RC, Rose S, Koenen KC, et al. How well can post-traumatic stress

disorder be predicted from pre-trauma risk factors? An exploratory study in the

WHO World Mental Health Surveys. *World Psychiatry*. 2014;13(3):265-274.

doi:10.1002/wps.20150

3.   Perkonigg A, Pfister H, Stein MB, et al. Longitudinal course of posttraumatic

stress disorder and posttraumatic stress disorder symptoms in a community sample

of adolescents and young adults. *Am J Psychiatry*. 2005;162(7):1320-1327.

doi:10.1176/appi.ajp.162.7.1320

4.   Keane TM, Kaloupek DG. Comorbid Psychiatric Disorders in PTSD. *Ann N Y*

Acad Sci. 1997;821(1 Psychobiology):24-34. doi:10.1111/j.1749-

6632.1997.tb48266.x

5.   Shalev A, Liberzon I, Marmar C. Post-Traumatic Stress Disorder. *N Engl J Med*.

2017;376(25):2459-2469. doi:10.1056/NEJMra1612499

6.   Hyman SE. Can neuroscience be integrated into the DSM-V? *Nat Rev Neurosci*.

2007;8(9):725-732. doi:10.1038/nrn2218

7.   Galatzer-Levy IR, Bryant RA. 636,120 Ways to Have Posttraumatic Stress

Disorder. *Perspect Psychol Sci*. 2013;8(6):651-662.

doi:10.1177/1745691613504115

8.   Bryant RA, O'Donnell ML, Creamer M, McFarlane AC, Silove D. A multisite

405  analysis of the fluctuating course of posttraumatic stress disorder. *JAMA*

406  *Psychiatry*. 2013;70(8):839-846. doi:10.1001/jamapsychiatry.2013.1137

407 9. Bisson JI, Roberts NP, Kitchiner NJ, Kenardy J. Systematic review and meta-

408  analysis of multiple-session early interventions following traumatic events. *Am J*

409  *Psychiatry*. 2009;166(3):293-301. doi:10.1176/appi.ajp.2008.08040590

410 10. Schuitevoerder S, Rosen JW, Twamley EW, et al. A meta-analysis of cognitive

411  functioning in older adults with PTSD. *J Anxiety Disord*. 2013;27(6):550-558.

412  doi:10.1016/j.janxdis.2013.01.001

413 11. Scott JC, Matt GE, Wrocklage KM, et al. A quantitative meta-analysis of

414  neurocognitive functioning in posttraumatic stress disorder. *Psychol Bull*.

415  2015;141(1):105-140. doi:10.1037/a0038039

416 12. Ben-Zion Z, Fine NB, Keynan NJ, et al. Cognitive Flexibility Predicts PTSD

417  Symptoms: Observational and Interventional Studies. *Front Psychiatry*.

418  2018;9:477. doi:10.3389/fpsyt.2018.00477

419 13. Pitman RK, Rasmusson AM, Koenen KC, et al. Biological studies of post-

420  traumatic stress disorder. *Nat Rev Neurosci*. 2012;13(11):769-787.

421  doi:10.1038/nrn3339

422 14. Hayes JP, Hayes SM, Mikedis AM, et al. Quantitative meta-analysis of neural

423  activity in posttraumatic stress disorder. *Biol Mood Anxiety Disord*. 2012;2(1):9.

424  doi:10.1186/2045-5380-2-9

425 15. Akiki TJ, Averill CL, Abdallah CG. A Network-Based Neurobiological Model of

426  PTSD: Evidence From Structural and Functional Neuroimaging Studies. *Curr*

427  *Psychiatry Rep*. 2017;19(11):81. doi:10.1007/s11920-017-0840-4

428   16.   Bremner JD, Elzinga B, Schmahl C, Vermetten E. Structural and functional

429   plasticity of the human brain in posttraumatic stress disorder. *Prog Brain Res*.

430   2007;167:171-186. doi:10.1016/S0079-6123(07)67012-5

431   17.   Wang Z, Neylan TC, Mueller SG, et al. Magnetic resonance imaging of

432   hippocampal subfields in posttraumatic stress disorder. *Arch Gen Psychiatry*.

433   2010;67(3):296-303.

434   18.   Smith ME. Bilateral hippocampal volume reduction in adults with post-traumatic

435   stress disorder: A meta-analysis of structural MRI studies. *Hippocampus*.

436   2005;15(6):798-807. doi:10.1002/hipo.20102

437   19.   Karl A, Schaefer M, Malta LS, et al. A meta-analysis of structural brain

438   abnormalities in PTSD. *Neurosci Biobehav Rev*. 2006;30(7):1004-1031.

439   doi:10.1016/j.amp.2006.02.006

440   20.   Shin LM, Liberzon I. The Neurocircuitry of Fear, Stress and Anxiety Disorders.

441   *Neuropsychopharmacology*. 2010;35(1):169-191. doi:10.1038/npp.2009.83

442   21.   Bolsinger J, Seifritz E, Kleim B, Manoliu A. Neuroimaging Correlates of

443   Resilience to Traumatic Events—A Comprehensive Review. *Front Psychiatry*.

444   2018;9:693. doi:10.3389/fpsyt.2018.00693

445   22.   Elzinga BM, Bremner JD. Are the neural substrates of memory the final common

446   pathway in posttraumatic stress disorder (PTSD)? *J Affect Disord*. 2002;70(1):1-

447   17. http://www.ncbi.nlm.nih.gov/pubmed/12113915.

448   23.   Rauch SL, Shin LM, Phelps EA. Neurocircuitry Models of Posttraumatic Stress

449   Disorder and Extinction: Human Neuroimaging Research-Past, Present, and

450   Future. *Biol Psychiatry*. 2006;60(4):376-382. doi:10.1016/j.biopsych.2006.06.004

24.   Liberzon I, Abelson JL. Context Processing and the Neurobiology of Post-

Traumatic Stress Disorder. *Neuron*. 2016;92(1):14-30.

doi:10.1016/j.neuron.2016.09.039

25.   Admon R, Milad MR, Hendler T. A causal model of post-traumatic stress disorder:

disentangling predisposed from acquired neural abnormalities. *Trends Cogn Sci*.

2013;17(7):337-347. doi:10.1016/J.TICS.2013.05.005

26.   Koenen KC, Moffit TE, Poulton R, Martin J, Caspi avshalom. Early childhood

factors associated with the development of post-traumatic stress disorder: results

from a longitudinal birth cohort. *Psychol Med*. 2007;37(2):181-192.

doi:10.1017/s0033291706009019

27.   Hendler T, Admon R. Predisposing Risk Factors for PTSD: Brain Biomarkers. In:

Comprehensive Guide to Post-Traumatic Stress Disorder. Cham: Springer

International Publishing; 2015:1-12. doi:10.1007/978-3-319-08613-2_64-1

28.   Schultebraucks K, Galatzer□Levy IR. Machine Learning for Prediction of

Posttraumatic Stress and Resilience Following Trauma: An Overview of Basic

Concepts and Recent Advances. *J Trauma Stress*. 2019:1-11.

doi:10.1002/jts.22384

29.   Feczko E, Miranda-Dominguez O, Marr M, Graham AM, Nigg JT, Fair DA. The

Heterogeneity Problem: Approaches to Identify Psychiatric Subtypes. *Trends

Cogn Sci*. 2019:1-18. doi:10.1016/j.tics.2019.03.009

30.   Galili T, Mitelpunkt A, Shachar N, Marcus-Kalish M, Benjamini Y. Categorize,

Cluster, And classify: A 3-c strategy for scientific discovery in the medical

informatics platform of the human brain project. *Lect Notes Comput Sci (including*

Subser Lect Notes Artif Intell Lect Notes Bioinformatics). 2014;8777:73-86.

31.   Ben-Zion Z, Fine NB, Keynan NJ, et al. Neurobehavioral Moderators of Post-

Traumatic Stress Disorder Trajectories: Prospective fMRI Study of Recent Trauma

Survivors. *Eur J Psychotraumatol*. 2019;10(1).

doi:10.1080/20008198.2019.1683941

32.   Weathers FW, Keane TM, Davidson JRT. Clinician-administered PTSD scale: A

review of the first ten years of research. *Depress Anxiety*. 2001;13(3):132-156.

doi:10.1002/da.1029

33.   Blevins CA, Weathers FW, Davis MT, Witte TK, Domino JL. The Posttraumatic

Stress Disorder Checklist for DSM-5 (PCL-5): Development. *J Trauma Stress*.

2015;28:489-498. doi:10.1002/jts.22059

34.   Beck, A. T., Steer, R. A., & Brown GK. Beck depression inventory-II. *San

Antonio*. 1996;78 (2)(2):490-498.

35.   Beck A, Epstein N, Brown G, Steer RA. An inventory for measuring clinical

anxiety: Psychometric properties. *J Consult Clin Psychol*. 1988;56(6):893.

36.   Busner J, Targum SD. The clinical global impressions scale: applying a research

tool in clinical practice. *Psychiatry (Edgmont)*. 2007;4(7):28-37.

37.   Silverstein S, Berten S, Olson P, et al. Development and validation of a World-

Wide-Web-based neurocognitive assessment battery: WebNeuro. *Behav Res

Methods*. 2007;39(4):940-949. doi:10.3758/BF03192989

38.   Fischl B. FreeSurfer. *Neuroimage*. 2012;62(2):774-781.

39.   Reuter M, Rosas H, - BF. Highly accurate inverse consistent registration: a robust

approach. *Neuroimage*. 2010.

497   https://www.sciencedirect.com/science/article/pii/S1053811910009717. Accessed

498   November 12, 2019.

499   40.   Ségonne F, Dale AM, Busa E, et al. A hybrid approach to the skull stripping

500   problem in MRI. *Neuroimage*. 2004;22(3):1060-1075.

501   doi:10.1016/j.neuroimage.2004.03.032

502   41.   Fischl B, Salat DH, Van Der Kouwe AJW, et al. Sequence-independent

503   segmentation of magnetic resonance images. *Neuroimage*. 2004;23(SUPPL. 1).

504   doi:10.1016/j.neuroimage.2004.07.016

505   42.   Fischl B, Salat DH, Busa E, et al. Whole brain segmentation: Automated labeling

506   of neuroanatomical structures in the human brain. *Neuron*. 2002;33(3):341-355.

507   doi:10.1016/S0896-6273(02)00569-X

508   43.   Fischl B, Salat DH, Busa E, et al. Whole Brain Segmentation□: Neurotechnique

509   Automated Labeling of Neuroanatomical Structures in the Human Brain. *Neuron*.

510   2002;33:341-355.

511   44.   Friston KJ, Ashburner JT, Kiebel SJ, Nichols TE, Penny WD. *Statistical*

512   *Parametric Mapping: The Analysis of Functional Brain Images*. Vol 8.; 2007.

513   doi:10.1016/B978-0-12-372560-8.50052-8

514   45.   Hariri AR, Bookheimer SY, Mazziotta JC. Modulating emotional responses:

515   effects of a neocortical network on the limbic system. *Neuroreport*. 2000;11(1):43-

516   48. doi:10.1097/00001756-200001170-00009

517   46.   Mitelpunkt A, Galili T, Shachar N, Marcus-Kalish M, Benjamini Y. Categorize,

518   Cluster & Classify - The 3C Strategy Applied to Alzheimer's Disease as a

519   Case Study. *Proc Int Conf Heal Informatics*. 2015:566-573.

doi:10.5220/0005275705660573

47.   Jack CR, Bernstein MA, Fox NC, et al. The Alzheimer's Disease Neuroimaging

Initiative (ADNI): MRI methods. *J Magn Reson Imaging*. 2008;27(4):685-691.

doi:10.1002/jmri.21049

48.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *J R Stat Soc Ser B*. 1995;57(1):289-300.

doi:10.1111/j.2517-6161.1995.tb02031.x

49.   Tibshirani R, Walther G, Hastie T. Estimating the number of clusters in a data set

via the gap statistic. *J R Stat Soc Ser B Stat Methodol*. 2001;63(2):411-423.

doi:10.1111/1467-9868.00293

50.   Kodinariya TM, Makwana PR. Review on determining number of Cluster in K-

Means Clustering. *Int J Adv Res Comput Sci Manag Stud*. 2013;1(6).

www.ijarcsms.com.

51.   Archer KJ, Kimes R V. Empirical characterization of random forest variable

importance measures. *Comput Stat Data Anal*. 2008;52(4):2249-2260.

doi:10.1016/j.csda.2007.08.015

52.   Shachar N, Mitelpunkt A, Kozlovski T, et al. The Importance of Nonlinear

Transformations Use in Medical Data Analysis. *JMIR Med informatics*.

2018;6(2):e27. doi:10.2196/medinform.7992

53.   Maechler M, Rousseeuw P, Struyf A, Hubert M, Hornik K. Cluster Analysis

Basics and Extensions. R package version 1.14.4. *Cran*. 2013. http://cran.r-

project.org/web/packages/cluster/index.html.

54.   Rousseeuw PJ. *Silhouettes: A Graphical Aid to the Interpretation and Validation*

of Cluster Analysis. Vol 20.; 1987.

55.   Zhang L, Li H, Benedek D, Li X, Ursano R. A strategy for the development of

545   biomarker tests for PTSD. *Med Hypotheses*. 2009;73(3):404-409.

546   doi:10.1016/j.mehy.2009.02.038

547   56.   Rubin DC, Berntsen D, Bohni MK. A Memory-Based Model of Posttraumatic

548   Stress Disorder: Evaluating Basic Assumptions Underlying the PTSD Diagnosis.

549   *Psychol Rev*. 2008;115(4):985-1011. doi:10.1037/a0013397

550   57.   Schuff N, Zhang Y, Zhan W, et al. Patterns of altered cortical perfusion and

551   diminished subcortical integrity in posttraumatic stress disorder: An MRI study.

552   *Neuroimage*. 2011;54(SUPPL. 1):S62-S68. doi:10.1016/j.neuroimage.2010.05.024

553   58.   Tsao A, Sugar J, Lu L, et al. Integrating time from experience in the lateral

554   entorhinal cortex. *Nature*. 2018;561(7721):57-62. doi:10.1038/s41586-018-0459-6

555   59.   Kitayama N, Quinn S, Bremner JD. Smaller volume of anterior cingulate cortex in

556   abuse-related posttraumatic stress disorder. *J Affect Disord*. 2006;90(2-3):171-174.

557   doi:10.1016/j.jad.2005.11.006

558   60.   Woodward SH, Kaloupek DG, Streeter CC, Martinez C, Schaer M, Eliez S.

559   Decreased anterior cingulate volume in combat-related PTSD. *Biol Psychiatry*.

560   2006;59(7):582-587. doi:10.1016/j.biopsych.2005.07.033

561   61.   Bryant RA, Felmingham K, Whitford TJ, et al. Rostral anterior cingulate volume

562   predicts treatment response to cognitive-behavioural therapy for posttraumatic

563   stress disorder. *J Psychiatry Neurosci*. 2008;33(2):142-146.

564   http://www.ncbi.nlm.nih.gov/pubmed/18330460.

62.   Woon FL, Sood S, Hedges DW. Hippocampal volume deficits associated with

exposure to psychological trauma and posttraumatic stress disorder in adults: A

meta-analysis. *Prog Neuro-Psychopharmacology Biol Psychiatry*.

2010;34(7):1181-1188. doi:10.1016/j.pnpbp.2010.06.016

63. Ferrari MCF, Busatto GF, McGuire PK, Crippa JAS. Structural magnetic

ressonance imaging in anxiety disorders: An update of research findings. *Rev Bras*

Psiquiatr. 2008;30(3):251-264. doi:10.1109/ICCIS.2013.425

64. Gilbertson MW, Shenton ME, Ciszewski A, et al. Smaller hippocampal volume

predicts pathologic vulnerability to psychological trauma. *Nat Neurosci*.

2002;5(11):1242-1247. doi:10.1038/nn958

65. Dedovic K, D'Aguiar C, Pruessner JC. What stress does to your brain: A review of

neuroimaging studies. *Can J Psychiatry*. 2009;54(1):6-15.

doi:10.1177/070674370905400104

66. Maclaren J, Han Z, Vos SB, Fischbein N, Bammer R. Reliability of brain volume

measurements: A test-retest dataset. *Sci Data*. 2014;1:140037.

doi:10.1038/sdata.2014.37

67. Admon R, Leykin D, Lubin G, et al. Stress-induced reduction in hippocampal

volume and connectivity with the ventromedial prefrontal cortex are related to

maladaptive responses to stressful military service. *Hum Brain Mapp*.

2013;34(11):2808-2816. doi:10.1002/hbm.22100

68. Bonne O, Brandes D, Gilboa A, et al. Longitudinal MRI Study of Hippocampal

Volume in Trauma Survivors With PTSD. *Am J Psychiatry*. 2001;158(8):1248-

1251. doi:10.1176/appi.ajp.158.8.1248

69. De Bellis MD, Hall J, Boring AM, Frustaci K, Moritz G. A pilot longitudinal study

of hippocampal volumes in pediatric maltreatment-related posttraumatic stress

disorder. *Biol Psychiatry*. 2001;50(4):305-309. doi:10.1016/S0006-

3223(01)01105-2

70.　RK S, AP K, RC W, et al. Neural dysregulation in posttraumatic stress disorder:

evidence for disrupted equilibrium between salience and default mode brain

networks. *Psychosom Med*. 2012;74(9):904-911 8p.

doi:10.1097/PSY.0b013e318273bf33

71.　Rabinak CA, Angstadt M, Welsh RC, et al. Altered amygdala resting-state

functional connectivity in post-traumatic stress disorder. *Front Psychiatry*.

2011;2(NOV):62. doi:10.3389/fpsyt.2011.00062

72.　Nicholson AA, Sapru I, Densmore M, et al. Unique insula subregion resting-state

functional connectivity with amygdala complexes in posttraumatic stress disorder

and its dissociative subtype. *Psychiatry Res - Neuroimaging*. 2016;250:61-72.

doi:10.1016/j.pscychresns.2016.02.002

73.　Lanius RA, Williamson PC, Densmore M, et al. Neural correlates of traumatic

memories in posttraumatic stress disorder: A functional MRI investigation. *Am J*

Psychiatry. 2001;158(11):1920-1922. doi:10.1176/appi.ajp.158.11.1920

74.　Yin Y, Jin C, Hu X, et al. Altered resting-state functional connectivity of thalamus

in earthquake-induced posttraumatic stress disorder: A functional magnetic

resonance imaging study. *Brain Res*. 2011;1411:98-107.

doi:10.1016/j.brainres.2011.07.016

75.　Aupperle RL, Melrose AJ, Stein MB, Paulus MP. Executive function and PTSD:

Disengaging from trauma. *Neuropharmacology*. 2012;62(2):686-694.

doi:10.1016/j.neuropharm.2011.02.008

76.   Polak AR, Witteveen AB, Reitsma JB, Olff M. The role of executive function in

posttraumatic stress disorder: A systematic review. *J Affect Disord*.

2012;141(1):11-21. doi:10.1016/j.jad.2012.01.001

77.   Scott WA. Cognitive Complexity and Cognitive Flexibility. *Sociometry*.

1962;25(4):405. doi:10.2307/2785779

78.   Miller MW, Resick PA. Internalizing and Externalizing Subtypes in Female

Sexual Assault Survivors: Implications for the Understanding of Complex PTSD.

Behav Ther. 2007;38(1):58-71. doi:10.1016/J.BETH.2006.04.003

79.   Miller MW, Greif JL, Smith AA. Multidimensional Personality Questionnaire

profiles of veterans with traumatic combat exposure: Externalizing and

internalizing subtypes. *Psychol Assess*. 2003;15(2):205-215. doi:10.1037/1040-

3590.15.2.205

80.   Miller MW, Kaloupek DG, Dillon AL, Keane TM. Externalizing and internalizing

subtypes of combat-related PTSD: A replication and extension using the PSY-5

scales. *J Abnorm Psychol*. 2004;113(4):636-645. doi:10.1037/0021-

843X.113.4.636

81.   Lanius RA, Brand B, Vermetten E, Frewen PA, Spiegel D. The dissociative

subtype of posttraumatic stress disorder: rationale, clinical and neurobiological

evidence, and implications. *Depress Anxiety*. 2012;29(8):701-708.

doi:10.1002/da.21889

82.   Lanius RA, Vermetten E, Loewenstein RJ, et al. Emotion modulation in PTSD:

Clinical and neurobiological evidence for a dissociative subtype. *Am J Psychiatry*.

2010;167(6):640-647. doi:10.1176/appi.ajp.2009.09081168

83. Swart S, Wildschut M, Draijer N, Langeland W, Smit JH. Dissociative Subtype of

Posttraumatic Stress Disorder or PTSD With Comorbid Dissociative Disorders:

Comparative Evaluation of Clinical Profiles. *Psychol Trauma Theory, Res Pract*

Policy. 2019. doi:10.1037/tra0000474

Tables Titles and Legends

Table 1. Confusion matrix of PTSD diagnosis clusters assignments. Table rows represent individuals' current clinical DSM-based PTSD diagnosis (PTSD / No PTSD), while the columns represent the two proposed clusters (LoClus=Low-Symptomatic Cluster / HiClus=High-Symptomatic Cluster).

Proposed Clusters / Clinical Diagnosis	LoClus	HiClus	Total
No PTSD	30	13	**43**
PTSD	14	44	**58**
Total	**44**	**57**	**101**

Table 2. Cross-Validation Results. The table present the mean percentage and standard deviation (SD) of subjects which were correctly classified (according to the results of the 3C methodology based on all subjects). For each P, n=1,000 iterations were performed.

P	20%	30%	40%	50%	60%	70%	80%	90%
Mean	83%	86%	86%	88%	89%	90%	90%	95%
SD	10%	7%	10%	9%	7%	7%	6%	7%

Fi`gure Titles and Legends

Figure 1. Parallel Coordinates Plot of Clinical Measurements. The Y-axis depicts the

four variables on which the clusters were built, all of them with an FDR-adjusted p-value

smaller than 0.2, while the x-axis depicts their percentiles. The figure presents, for each

cluster, its median value (bold line) and their 0.025 and 0.975 percentiles (i.e., 95% CI,

the scattered "cloud" around it) of 400 bootstrap samplings' medians. Participants

belonging to the first cluster (turquoise) were, on average, higher on all four variables,

compared to those belonging to the second cluster (red). "T1_CGI_Total" = Total Score

of Clinical Global Impression Scale Questionnaire; "T1_PCL_Total" = Total Score of

PTSD Checklist Questionnaire; "T1_BAI_Total" = Total Score of Beck Anxiety

Inventory Questionnaire; "T1_BDI_Total"= Total Score of Beck Depression Inventory

Questionnaire.

Figure 2. Parallel Coordinates Plot of Potential Neurocognitive Biomarkers. The Y-axis depicts the top ten most important potential neurocognitive biomarkers in classifying to the two clusters, together with their mean decrease GINI measure (i.e. importance index). Biomarkers from cognitive domains are colored in yellow, biomarkers from structural brain measurements are colored in dark blue, and biomarkers from functional brain measurements are colored in green. Average CAPS-4 total score is presented for both "Low-Symptomatic" Cluster (cluster 1, turquoise) and "High-Symptomatic" Cluster (cluster 1, red). The medians of 400 Bootstrap samplings were drawn, and their median and 0.025 and 0.975 quantiles are plotted per cluster. "rh_rostralanteriorcingulate_volume" = right rostral anterior cingulate cortex volume; "lh_entorhinal_volume" = left entorhinal cortex volume; "rh_transversetemporal_volume" = right transverse temporal gyrus volume;

"log.X3rd.Ventricle" = third ventricle volume (after log transformation to create symmetrical distribution); "T1_Flexibility_Norm" = cognitive flexibility score; "lh_parsorbitalis_volume" = left pars orbitalis volume; "log.X4th.Ventricle" = fourth ventricle volume (after log transformation to create symmetrical distribution); "neutral.Visual.avg" = average visual cortex activity in response to neutral faces; "rh_paracentral_thickness" = para central lobule cortical thickness; "PPI_angry.amy.caudate.thalamus.avg" = average amygdala functional connectivity with the thalamus while watching angry faces vs. shapes; "Total CAPS" = Total scores of CAPS-4.

Figure 3. Classification Tree based on the Two Clusters. The classification tree

depicts variables important for the division of the participants to the two clusters, starting

from the most important one at the top of the tree (left entorhinal cortex volume). Each

block in tree is labeled either HC or LC, indicating whether most of the subjects in that

block belong to the HiClus or the LoClus (colored bluer or greener, respectively).

Furthermore, each block has the number of subjects belonging to the dominant cluster

(either HC or LC), and the total number of subjects in that specific block. Inspecting the

top block for example, 70 out of 101 participants had left EC volume equal to or greater

than 1449mm^3, out of which 39 belonged to the LoClus. The other 31 participants had a

left EC volume smaller than 1449mm^3, out of which the most (n=26) belonged to the

HiClus. For full description please see eResults. "lh_entorhinal_volume" = left entorhinal

cortex volume; "T1_Executive_Function_Norm" = executive function score;

"lh_caudalmiddlefrontal_volume" = left caudal middle frontal gyrus volume;

"lh_supramarginal_ thickness" = left supramarginal gyrus cortical thickness;

"PPI_fear.amy.L.insula.avg" = average amygdala functional connectivity with the left

insula while watching fearful faces vs. shapes; "log.rh_precentral_volume" = right

precentral gyrus volume (after log transformation to create symmetrical distribution).