Impact of second-generation antipsychotics on white matter microstructure in

adolescent-onset psychosis.

Claudia Barth*[a], PhD; Vera Lonning [a], MD; Tiril Pedersen Gurholt [a,b], PhD; Ole A.

Andreassen [a,c], Professor; Anne M. Myhre [c,d], Associate Professor, MD; Ingrid Agartz [a,b,e],

Professor.

[a] Norwegian Centre for Mental Disorders Research, KG Jebsen Centre for Psychosis

Research, Institute of Clinical Medicine, University of Oslo, Oslo, Norway

[b] Department of Psychiatric Research, Diakonhjemmet Hospital, Oslo, Norway

[c] Norwegian Centre for Mental Disorders Research, KG Jebsen Centre for Psychosis

Research, Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway

[d] Child & Adolescent Mental Health Research Unit, Oslo University Hospital, Oslo, Norway

[e] Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden

*Corresponding author: Claudia Barth, PhD, E-Mail: claudia.barth@medisin.uio.no, phone:

+47 22 02 99 67, fax: +47 22 02 99 01, postal address: Institute of Clinical Medicine, Section

Vinderen, University of Oslo, P.O. Box 85, Vinderen, N-0319 Oslo, Norway.

Abstract

White matter abnormalities are well-established in adult patients with psychosis. Yet less is known about changes in early onset psychosis (EOP) during adolescence, especially whether antipsychotic medication might impact white matter microstructure in this sensitive phase. Here, we utilized Magnetic Resonance Imaging (MRI) in unmedicated and medicated adolescent EOP patients in comparison to healthy controls to examine the impact of antipsychotic medication status on indices of white matter microstructure. Twenty-two EOP patients (11 unmedicated) and 33 healthy controls, aged between 12-18 years, underwent 3T diffusion-weighted MRI. Using Tract-based Spatial Statistics, we calculate case-control differences in scalar diffusion measures, e.g. fractional anisotropy (FA), and investigated their association with antipsychotic medication. We replicated previous results from studies in EOP patients showing significantly decreased mean FA including the left genu of the corpus callosum, the left anterior corona radiata and the right superior longitudinal fasciculus in patients relative to healthy controls. Mean FA in the left anterior corona radiata was significantly associated with antipsychotic medication status, showing higher FA values in medicated compared to unmedicated EOP patients. Increased regional FA values might be a first hint towards an early effect of antipsychotic medication on white matter microstructure in adolescent EOP patients.

Introduction

A variety of hypotheses has been proposed to explain the etiology of psychotic disorders, including aberrant dopamine neurotransmission [1], altered neurodevelopmental trajectories [2], and active neuroinflammation [3]. Such theories are not mutually exclusive and are more likely complementary. Patients with early onset psychosis (EOP), with defined age of onset before 18 years, provide an unprecedented opportunity to specifically investigate the perspective of aberrant neurodevelopment.

The application of diffusion weighted imaging (DWI) can relate white matter organization to disease. DWI maps the Brownian movement of water molecules in the brain in vivo, and as axon membranes and myelin provide natural barriers for water diffusion, DWI can be used to infer local tissue properties [4]. The most commonly used scalar measure is fractional anisotropy (FA), which characterizes the degree of diffusion directionality. For an in-depth evaluation of FA, the relative contribution of axial diffusion (AD) along the primary axis, and radial diffusion (RD) perpendicular to it, can be informative. RD has been associated with changes in myelin [5], and a disruption of myelin sheaths may be reflected in an increased RD. Conversely, AD has been linked to axonal integrity and axonal damage may be characterized by decreased AD [4].

Using DWI, a number of studies show widespread FA reductions in many different brain regions with low spatial overlap such as corpus callosum, cingulum, superior longitudinal fasciculus, inferior longitudinal fasciculus and fronto-occipital fasciculus in EOP patients compared to healthy controls [6-10]. Scalar DWI measures beyond FA are rarely analyzed in EOP populations. However, Lagopoulos and colleagues report on increased RD values, indicative of potential demyelinating processes underlying the observed white matter abnormalities [10].

The low degree of regional specificity of whiter matter changes seems to be attributed to a number of factors including differences in image acquisition, different analysis approaches (ROI vs voxel-wise), small sample sizes, low prevalence of EOP (estimated prevalence of 17.6 in 10,000 at age of 18 years, [11]) and differing sample characteristics such as age of onset. Further, antipsychotic medication status might also affect the pattern of white matter microstructure in EOP.

Studies investigating white matter microstructure in EOP mainly focus on case-control differences, either reporting antipsychotic effects as secondary findings or using antipsychotic medication status as a covariate of no interest. So far, studies in EOP patients do not indicate an impact of either current [12-14] or cumulative antipsychotic exposure [6,13,15] on scalar DWI measures. The absence of an antipsychotic medication effect could reflect small sample sizes

and young patients with shorter medication histories. The apparent limitations of the
adolescent study population also hold potential advantages: EOP patients are less affected by
chronic exposure to antipsychotic medication in comparison to their adult counterparts, which
allows for dissecting medication-mediated from disease-related effects on brain structure.
Furthermore, according to the World Health Organization guideline for pharmacological
interventions in adolescents with psychotic disorders (2015), antipsychotic medication use is
significantly less recommended in comparison to adult patients with psychosis [16]. This partly
translates to a clinical practice of a higher reluctance in starting antipsychotic treatment early
in the course of psychosis in children and adolescents, leading to a higher percentage of
antipsychotic-naïve EOP patients relative to adult first episode patients with psychosis. Thus,
EOP patients represent an ideal population to investigate the impact of antipsychotic
medication on white matter structure early in disease progression.

Here, we use a thoroughly clinically characterized adolescent EOP sample to (1) investigate
white matter microstructure in comparison to healthy controls, and (2) explore the association
between second-generation antipsychotic medication and white matter microstructure in
medicated compared to currently unmedicated/antipsychotic-naïve EOP patients. We utilize
DWI and, by using Tract-Based Spatial Statistics (TBSS), we calculate FA and its scalar sub-
measures, RD and AD, and investigate their association with antipsychotic medication and
other clinical measures (e.g. Positive and Negative Syndrome Scale, etc.). Based on the
existing literature, we hypothesized that EOP patients show widespread reduced FA attended
by increased RD and unchanged AD compared to healthy controls, mainly in the corpus
callosum and superior/inferior longitudinal fasciculus. As there are no established effects of
antipsychotic medication on white matter structure in EOP patients, our post hoc analysis is
exploratory by nature.

Results

Demographic and clinical data

As presented in Table 1, EOP patients did not differ significantly from controls in general demographic variables such as age, handedness and IQ. However, there was a significant case-control difference in clinical measures such as CGAS and MFQ, reflecting the clinical diagnosis. EOP patients showed higher impairment of general functioning evaluated with CGAS and exhibited significantly more depressive symptoms using MFQ, relative to their healthy counterparts. Furthermore, patients report significantly more cannabis use than controls.

Within the patient group (Table 2), patients on antipsychotic medication were significantly older than unmedicated patients (t = -4.186, p = 0.0006). In line with antipsychotic medication status, duration of untreated psychosis (DUP) was significantly longer in the unmedicated EOP subgroup in comparison to the medicated group. Further demographic and clinical variables did not differ significantly between the patient subgroups.

TBSS analyses

Voxel-wise statistical analysis of case-control differences revealed decreases in mean FA including the left genu of the corpus callosum, the left anterior corona radiata (ACR), and the right superior longitudinal fasciculus (SLF) in EOP patients compared to healthy controls (see Figure 1 and Table 3). There was no increase in mean FA for the opposing contrast.

Applying the TBSS pipeline to diffusion-derived data other than FA, namely RD and AD, did not yield significant case-control differences for RD, but decreases in mean AD overlapping with FA findings in EOP patients in comparison to healthy controls. In detail, mean AD shows significantly decreases in the left ACR (see supplementary Table S2), in addition to decreases in the right posterior limb of the internal capsule (PLIC) and right superior fronto-occipital fasciculus (SFOF).

Extracted mean values of all scalar diffusion measures for all significant clusters stratified by group are displayed in supplementary Figure S1 for descriptive purposes only. In the interest of transparency, TBSS case-control differences in mean FA and mean AD (patients < controls) at a cluster-forming threshold of p ≤ 0.05 are also presented (see supplementary Figure S2).

Linear regression analyses

To evaluate the potential influence of duration of illness and antipsychotic treatment on regional mean FA and mean AD values within significant TBSS clusters, in patients, linear regression

analyses were performed. While duration of illness was not significantly associated with mean FA in any of the clusters, we found a significant negative association with mean AD in the left ACR (t = -2.364, p = 0.029). For latter association, however, the regression equation was not significant (p = 0.085) and the overall explanatory power of the model was low (R^2 = 0.147), rendering this finding most likely spurious (see supplementary Table S1).

Exposure to antipsychotic medication was significantly associated with mean FA values in the left ACR (t = 2.991, p = 0.008), showing higher mean FA in medicated relative to the unmedicated patients (Figure 2). There was no association between antipsychotic medication and mean FA or mean AD in the other significant TBSS clusters (see supplementary Table S1). Cohen's d effect size values suggest a high (d = 1.48) and a low (d = -0.08) standardized difference in mean FA of the left ACR in unmedicated and medicated EOP patients relative to healthy controls, respectively (Figure 3). Based on visual inspection, higher mean FA in medicated patients seems driven by an increase in AD and a decrease in RD (Figure 2). However, mean AD and RD did not differ significantly between medicated and unmedicated patients (Welch Two Sample t-test, AD: t = 0.183, p = 0.857; RD: t = 1.887, p = 0.079).

Association with clinical measures

We found no association between current antipsychotic medication evaluated as chlorpromazine equivalent at scan day (CPZ, Spearman ρ = 0.13, n = 11, p = 0.695) or cumulative CPZ (Pearson ρ = -0.15, n = 11, p = 0.665) with regional mean FA values in the left ACR.

In addition, we also found no significant correlations, which survived correction for multiple comparisons (FA cluster: Bonferroni, α = 0.003; AD cluster: Bonferroni, α = 0.004), between neither extracted mean FA nor mean AD values of all significant TBSS clusters and clinical measures such as PANSS (neither positive nor negative), CGAS and MFQ scores.

Discussion

Our case-control results replicate prominent brain regions with known white matter abnormalities implicated in EOP, namely corpus callosum [6], right SLF [14,17,18] and the left ACR [9,10]. Intriguingly, lower FA in these regions seems to occur early in the disease process [10,18,19]. For instance, Lagopoulos and colleagues found a decrease in FA in the left ACR in both patients with established psychiatric disorder and patients exhibiting sub-syndromal symptoms, aged 14-30 years. Based on these findings, the authors proposed that abnormalities in the left ACR are a putative precursor to the development of a psychiatric condition [10].

However, the ACR is a highly heterogeneous structure with three long-range association fiber tracts traversing through it [10]: anterior thalamic radiation (ATR), inferior fronto-occipital fasciculus (IFOF) and uncinate fasciculus (UF). All three association fibers form connections to the frontal lobe and have been implicated in the pathophysiology of psychiatric disorders [10,20-22]. In the current study, the left ACR peak voxel shows a 16% probability of IFOF involvement based on the JHU White-Matter Tractography Atlas. The IFOF connects the occipital and temporal lobes with the orbitofrontal cortex as part of the ventral visual and language stream. In particular, the left IFOF seems to subserve language semantics [23]. Already in 1996, Aloia and colleagues proposed that the disruption of semantic networks have potential implications for the origin of "thought disorder" in schizophrenia [24]. Adding to this hypothesis, patients with 22q11.2 deletion syndrome, who are genetically at high risk for developing schizophrenia, showed lower FA values in left IFOF [25]. Furthermore, DeRosse and colleagues found that lower FA proximal to the SLF and corticospinal tract bilaterally, and left IFOF and left inferior longitudinal fasciculus (ILF), were associated with higher levels of psychotic-like experiences in otherwise healthy volunteers [26]. In early-onset schizophrenia (EOS) patients, lower FA in the left IFOF and the left ILF predicted worse neurocognitive performance [9]. The authors also detected a shared decrease in FA in the left IFOF among patients with clinical high risk for schizophrenia and patients with established EOS, in comparison to healthy controls. Together, these findings suggest that white matter abnormalities in the left ACR, putatively in the left IFOF, may represent a potential candidate for understanding the etiology of psychosis.

This assumption seems further supported by effects of antipsychotic medication on diffusion metrics in the left ACR. We found that FA values in the left ACR were significantly predicted by antipsychotic medication status, with higher FA values in medicated relative to unmedicated EOP patients. No such association was found with the other brain regions showing significantly decreased FA values. Besides the high Cohen's d effect size estimate (Figure 2), we found no significant association of regional FA with either current or cumulative antipsychotic exposure.

This lack of significant associations is, however, in line with previous studies in EOP patients
[6,12-15] and likely due to the fairly short medication history in EOP patients compared to their
adult-onset counterparts or limited sample size. Hence, the presence of antipsychotic
medication rather than the actual dose might induce the observed changes in white matter
microstructure.

In medicated patients, increased FA values seem to be driven by an increase in AD and a
decrease in RD, relative to their unmedicated counterparts (Figure 2). Thus, FA might be
enhanced by antipsychotic medication as a result of both facilitated parallel diffusivity (AD,
potentially mediated by an increase in axon numbers, and restricted perpendicular diffusivity
(RD), indicative of changes in myelin.

Converging evidence from multiple studies suggests oligodendroglial dysfunction, with
subsequent abnormalities in myelin maintenance and repair, to underpin white matter
abnormalities observed in psychotic patients [27]. In the framework of schizophrenia, it has been
proposed that myelin dysfunction, especially in frontal regions, contributes to psychotic
symptoms [13,27]. Based on findings from cell culture studies using aripiprazole [28] and rodent
work using quetiapine [29,30], second-generation antipsychotic medication may promote
oligodendrocyte recovery and myelin repair leading to reduced white matter abnormalities and,
subsequently, reduced psychotic symptoms. A recent study in patients with schizophrenia also
reports on promyelinating effects of antipsychotics [31]. Tishler and colleagues found an increase
in intracortical myelin predominantly mediated by risperidone and other second-generation
antipsychotics in adult patients with schizophrenia compared to healthy controls within the first
year of treatment. In the current study, given that medicated EOP patients received either
aripiprazole, quetiapine or risperidone, one might speculate that early medication with second-
generation antipsychotics might affect white matter microstructure by remediating
oligodendroglial dysfunction, leading to an increase in FA detected by DWI.

Even though FA is highly sensitive to microstructural changes in general, it lacks
neurobiological specificity to the exact type of change [4]. For instance, a decrease in FA can
reflect alternations in fiber organization, including packing density and fiber crossing, and
myelin loss or myelin remodeling [32]. We found a widespread decrease of AD on a whole brain
level, indicative of axonal damage, but no changes were found in RD, relative to healthy
controls. This finding is not in line with previous work from Lagopoulos and colleagues, who
found a decrease in FA in the left ACR associated with increases in RD and no changes in AD
[10].

Although there are likely several reasons for these conflicting findings, neuroinflammatory
processes might pose particular difficulties in interpreting DWI signals in psychotic populations

[33]. For instance, in an animal model of cuprizone-induced demyelination of corpus callosum,
regions with extensive axonal edema and prominent cellular inflammation showed no change
in RD, while AD values were diminished at the beginning of demyelination [34]. Given the
neuroinflammation hypothesis of schizophrenia [3], it seems likely that the disease progression
encompasses a dynamic evolution of inflammation, axonal injury, and myelin degeneration. In
the current study, one might speculate that EOP patients are on the verge of undergoing
demyelination processes, reflected by widespread decreases in AD. However, the timing of
neuroinflammation in psychotic disorders relative to tissue injury is unclear, leading to a
heightened risk of misinterpreting changes in DWI measures. According to a recent review of
Winklewski and colleagues, in cases of neuroinflammation linked to tissue damage, DWI
seems to underestimate the extent of demyelination (undervalued RD), and overestimate the
extent of axonal injury (overvalued AD) [35]. This pattern seems replicated in our study, with
significant changes in AD and no changes in RD. As the consistency of DWI metrics seems
affected by brain edema and inflammatory response, future studies can benefit from using
tools such as free water imaging which provide the opportunity to separate the contribution of
extracellular water from the diffusion of water molecules inside the fiber tracts, leading to a
higher specificity in detecting structural changes [36].

Deviations in scalar DWI measures in the current study relative to previous studies could also
be due to ongoing white matter maturation processes in our adolescent EOP sample. In
healthy individuals, age-related increases in FA during childhood, adolescence and early
adulthood have been consistently reported [37-40]. This increase in FA seems primarily driven by
a reduction in RD, while AD remains fairly stable or decreases slightly [41,42]. Findings for AD
changes during the transition to adulthood are less consistent [37-40]. Thus, the AD difference
found in the current study could also be attributed to developmental processes, which may
fade as adolescents mature into adulthood.

However, it should be noted that the neurodevelopmental trajectories of white matter structure
relative to disease progression in EOP patients are unclear. So far, three different studies
yielded inconclusive results, postulating either diverging [13], converging [43], or parallel [44]
trajectories relative to healthy controls. In the current study, we did not find any predictive value
of duration of illness for regional FA. This is in line with previous findings from Kumra and
colleagues, who speculated that the decrease in FA in EOP patients compared to healthy
controls reflects developmental abnormalities rather than secondary effects of the disease
progression [13]. In addition, Epstein and Kumra found lower FA in the inferior longitudinal
fasciculus, IFOF and corticospinal tract, but no significant group differences in longitudinal
changes in FA [44]. Thus, the observed changes in the current study might persist but do not

affect the overall white matter maturation trajectories. However, the cross-sectional nature of

the current study precludes the assessment of developmental effects over time.

The results of the current study should be considered in the context of several limitations.

Unmedicated EOP patients were significantly younger than those receiving medication.

Although the analysis was corrected for age, we cannot exclude that the age of the patients

contributes to the observed antipsychotic-medication related changes in white matter structure.

It is possible that time-of-measurement effects, with older patients having higher FA values

than younger patients due to more advanced white matter maturation, could confound our

results. However, while we acknowledge this possibility, we consider this unlikely to be the

driving mechanism for the following reason: we would expect differences in FA values in other

regions showing a similar maturation trajectory, such as the SLF [41], if our results were mainly

driven by age differences. This was not the case, as we found no significant difference in mean

FA values of the SLF between medicated and unmedicated patients (t = -0.56934, p-value =

0.576). We further stress that we did not find any associations between regional FA variation

and clinical measures such a PANSS scores, which is likely due to the relatively small sample

size.

In summary, the present study is the first to link antipsychotic medication status to altered

regional FA in the left ACR in patients with EOP. Understanding the significance of white matter

abnormalities in the left ACR in adolescents with EOP and the putatively remediating effect of

antipsychotic medication, may help to phenotype the disease and to develop new

pharmacological regimes to subsequently improve functional outcome. Assuming that

antipsychotic medication reverses the hypothesized myelin dysfunction in psychosis, early

interventions with antipsychotic medication, already in individuals at risk of developing

psychosis, could provide the opportunity to normalize white matter maturation. Although

exciting, further work is needed to draw firm conclusions about the beneficial effects of

antipsychotic medication early in the disease process. Building on our first results, longitudinal

studies with larger samples sizes using high resolution DWI in combination with clinical, genetic

and neurocognitive measures are warranted to delineate heritability, affected brain regions,

antipsychotic medication effects, and directions of FA changes over time.

Methods

Participants

The study sample was drawn from the ongoing longitudinal Youth-Thematic-Organized-Psychosis (Youth-TOP) research study, which is a subdivision of the TOP research group/NORMENT and KG Jebsen center of psychosis research in Oslo, Norway. EOP patients, aged between 12-18 years, were recruited from in- and outpatient clinics in the Oslo region. Healthy controls were randomly selected from the Norwegian National Registry in the same catchment area. All participants and their respective parents/guardians provided written informed consent. The study was approved by the Regional Committee for Medical Research Ethics (REK-Sør) and the Norwegian Data Inspectorate and was conducted in accordance with the Declaration of Helsinki.

For study inclusion, participants were required to have an intelligence quotient (IQ) > 70, a good command of the Norwegian language, no previous moderate to severe head injuries, no diagnosis of substance-induced psychotic disorder, and no organic brain disease. IQ was measured by the Wechsler Abbreviated Scale of Intelligence [45]. Diagnosis was established according to the Diagnostic and Statistical Manual of Mental Disorder- IV criteria using the Norwegian version of the Kiddie-Schedule for Affective Disorders and Schizophrenia for School Aged Children (6-18 years): Present and Lifetime Version (K-SADS-PL[46]). The clinical characterization was conducted by trained psychologists or psychiatrists.

A total of 67 participants (27 patients/40 controls) satisfied the above-mentioned criteria and underwent MRI examination. All MRI scans were visually inspected by a trained neuroradiologist to rule out any pathological changes. Out of the initial sample, seven control participants and five patients were excluded due to (i) clinical/radiological reasons (five patients/ three controls), or (ii) strong motion artefacts in the diffusion imaging data (four controls), resulting in a final sample of 55 participants (22 patients/ 33 controls) being entered in the statistical analysis.

Sample demographics and clinical characteristics separated by antipsychotic medication status of EOP patients are reported in Table 1 and Table 2, respectively.

Clinical measures

Presence and severity of psychopathological symptoms of EOP patients were assessed using the Positive and Negative Syndrome Scale (PANSS[47]). Children Global Assessment Scale (CGAS[48]) and Mood and Feelings Questionnaire (MFQ, long version [49]) were evaluated in all participants to measure general functioning level and to screen for depressive symptoms,

respectively. Recreational drug use was assessed within the structured K-SADS interview and
scored with 0 or 1 for absent or present. For EOP patients, current and lifetime cumulative use
of medication was recorded and converted into chlorpromazine equivalents (CPZ), using
formulas published elsewhere [50]. While 11 EOP patients were off any antipsychotic medication
at scan, yielding a lack of current CPZ values, 3 patients had received pharmacological
treatment prior to inclusion, resulting in a low cumulative CPZ dosage for this subgroup (see
Table 2).

MRI data acquisition

MR images were acquired on a 3-Tesla General Electric Signa HDxt scanner equipped with
an 8-channel head coil at the Oslo University Hospital, Norway. The diffusion imaging data
was acquired using a 2D spin-echo whole-brain echo-planar imaging sequence with the
following parameters: slice thickness = 2.5 mm, repetition time = 15 s, echo time = 85 ms, flip
angle = 90°, acquisition matrix = 96 x 96, in-plane resolution = 1.875 x 1.875 mm. A total of 32
volumes with different gradient directions (b = 1000 s/mm^2), including two b0-volumes with
reversed phase-encode (blip up/down), were acquired.

Diffusion data analysis

Diffusion data were analyzed with FSL version 5.0.9 using the FMRIB's software library
(https://fsl.fmrib.ox.ac.uk/fsl/fslwiki). Before creating voxel wise maps of diffusion parameters,
the following steps of the standard processing pipeline were used: (i) *topup* to correct for
susceptibility-induced distortions [51,52], (ii) *eddy* current correction to correct for gradient-coil
distortions and head motion [53], (iii) removal of non-brain tissue using the Brain Extraction Tool
(*bet*) [54], and (iv) local fitting of the diffusion tensor at each voxel using *dtifit* (FMRIB's Diffusion
Toolbox (FDT) [55]). *Dtifit* yielded in voxel wise participant-specific maps of FA, mean diffusion
(MD), and axial diffusivity (AD, derived from eigenvector $\lambda 1$). Based on the outputted
eigenvectors $\lambda 2$ and $\lambda 3$, radial diffusivity (RD) was computed (($(\lambda 2+\lambda 3)/2$)). Next, voxel wise
statistical analysis of the FA data was carried out using TBSS [56]. First, all FA images were
nonlinearly aligned to the most representative FA image out of all images and transformed into
1x1x1 mm^3 MNI152 standard space by means of affine registration. Secondly, TBSS projects
all participant's FA data onto a mean FA tract skeleton (threshold FA > 0.25), before applying
voxel wise cross-participant statistics. After TBSS for FA was completed, results were used to
generate skeletonized RD and AD data for additional voxel-wise group comparisons using the
TBSS non-FA pipeline.

Statistical analyses

For contrasting case-control differences, we run voxel-wise statistics, co-varied for age and
gender, using a nonparametric permutation-based approach (*Randomise*, implemented in FSL,
5000 permutations). All variables were demeaned. The statistical threshold was set at $p \leq 0.01$,
after family-wise error correction for multiple comparisons using threshold-free cluster
enhancement. We chose a highly conservative threshold for FA to minimize type I errors and
to better account for the exploratory nature of the study concerning the impact of antipsychotic
medication status. For RD and AD, the same statistical model was used.

Regions identified with TBSS (FA) and TBSS non-FA (RD or AD were subsequently used as
masks to extract mean FA, RD and AD values for plotting and further analysis. We refrained
from using MD values, as a measure of overall diffusivity within a voxel, in further analysis due
to its lack of specificity [4]. As scalar diffusion measures largely vary in their value ranges,
extracted mean values were z-standardized for plotting purposes using the following formula:
z = (participant's value – group mean) / standard deviation.

A linear regression model was performed to examine whether patients' mean values of
significant TBSS and TBSS non-FA clusters were associated with duration of illness and
antipsychotic medication status as categorical variable (coded as yes (1)/no (0)). The effect
size was reported as Cohen's d [57].

If there is an association between patients' regional mean values and antipsychotic medication,
follow-up correlation analysis with current and cumulative CPZ were performed using
Spearman's rank correlation rho for non-normal data.

Further analysis of regional mean values and its association with clinical measures (PANSS,
CGAS, MFQ) were performed using Pearson's product moment correlation coefficient.

Statistical tests were conducted in R, version 3.5.2 (www.r-project.org).

References

1. Jauhar, S. *et al.* A Test of the Transdiagnostic Dopamine Hypothesis of Psychosis Using Positron Emission Tomographic Imaging in Bipolar Affective Disorder and Schizophrenia. *JAMA Psychiatry* **74**, 1206-1213, doi:10.1001/jamapsychiatry.2017.2943 (2017).

2. Rapoport, J. L., Giedd, J. N. & Gogtay, N. Neurodevelopmental model of schizophrenia: update 2012. *Mol Psychiatry* **17**, 1228-1238, doi:10.1038/mp.2012.23 (2012).

3. Najjar, S. & Pearlman, D. M. Neuroinflammation and white matter pathology in schizophrenia: systematic review. *Schizophr Res* **161**, 102-112, doi:10.1016/j.schres.2014.04.041 (2015).

4. Alexander, A. L., Lee, J. E., Lazar, M. & Field, A. S. Diffusion tensor imaging of the brain. *Neurotherapeutics* **4**, 316-329, doi:10.1016/j.nurt.2007.05.011 (2007).

5. Song, S. K. *et al.* Dysmyelination revealed through MRI as increased radial (but unchanged axial) diffusion of water. *Neuroimage* **17**, 1429-1436 (2002).

6. Douaud, G. *et al.* Anatomically related grey and white matter abnormalities in adolescent-onset schizophrenia. *Brain* **130**, 2375-2386, doi:10.1093/brain/awm184 (2007).

7. James, A. *et al.* Greater white and grey matter changes associated with early cannabis use in adolescent-onset schizophrenia (AOS). *Schizophr Res* **128**, 91-97, doi:10.1016/j.schres.2011.02.014 (2011).

8. Ashtari, M. *et al.* Disruption of white matter integrity in the inferior longitudinal fasciculus in adolescents with schizophrenia as revealed by fiber tractography. *Arch Gen Psychiatry* **64**, 1270-1280, doi:10.1001/archpsyc.64.11.1270 (2007).

9. Epstein, K. A. *et al.* White matter abnormalities and cognitive impairment in early-onset schizophrenia-spectrum disorders. *J Am Acad Child Adolesc Psychiatry* **53**, 362-372 e361-362, doi:10.1016/j.jaac.2013.12.007 (2014).

10. Lagopoulos, J. *et al.* Microstructural white matter changes are correlated with the stage of psychiatric illness. *Transl Psychiatry* **3**, e248, doi:10.1038/tp.2013.25 (2013).

11. Gillberg, C., Wahlstrom, J., Forsman, A., Hellgren, L. & Gillberg, I. C. Teenage psychoses--epidemiology, classification and reduced optimality in the pre-, peri- and neonatal periods. *J Child Psychol Psychiatry* **27**, 87-98 (1986).

12. Moran, M. E. *et al.* Comparing fractional anisotropy in patients with childhood-onset schizophrenia, their healthy siblings, and normal volunteers through DTI. *Schizophr Bull* **41**, 66-73, doi:10.1093/schbul/sbu123 (2015).

13. Kumra, S. *et al.* White matter abnormalities in early-onset schizophrenia: a voxel-based diffusion tensor imaging study. *J Am Acad Child Adolesc Psychiatry* **44**, 934-941, doi:10.1097/01.chi.0000170553.15798.94 (2005).

14. Kyriakopoulos, M., Vyas, N. S., Barker, G. J., Chitnis, X. A. & Frangou, S. A diffusion tensor imaging study of white matter in early-onset schizophrenia. *Biol Psychiatry* **63**, 519-523, doi:10.1016/j.biopsych.2007.05.021 (2008).

15. Kyriakopoulos, M. *et al.* Effect of age at onset of schizophrenia on white matter abnormalities. *Br J Psychiatry* **195**, 346-353, doi:10.1192/bjp.bp.108.055376 (2009).

16. Organization, W. H. Pharmacological interventions in adolescents with psychotic disorders. (2015).

17. Schwehm, A. *et al.* Age and Sex Effects on White Matter Tracts in Psychosis from Adolescence through Middle Adulthood. *Neuropsychopharmacology* **41**, 2473-2480, doi:10.1038/npp.2016.47 (2016).

18. Karlsgodt, K. H. *et al.* Diffusion tensor imaging of the superior longitudinal fasciculus and working memory in recent-onset schizophrenia. *Biol Psychiatry* **63**, 512-518, doi:10.1016/j.biopsych.2007.06.017 (2008).

19. DeRosse, P. *et al.* White Matter Abnormalities Associated With Subsyndromal Psychotic-Like Symptoms Predict Later Social Competence in Children and Adolescents. *Schizophr Bull* **43**, 152-159, doi:10.1093/schbul/sbw062 (2017).

20 Kawashima, T. *et al.* Uncinate fasciculus abnormalities in recent onset schizophrenia
and affective psychosis: a diffusion tensor imaging study. *Schizophr Res* **110**, 119-
126, doi:10.1016/j.schres.2009.01.014 (2009).
21 Sprooten, E. *et al.* The relationship of anterior thalamic radiation integrity to psychosis
risk associated neuregulin-1 variants. *Mol Psychiatry* **14**, 237-238, 233,
doi:10.1038/mp.2008.136 (2009).
22 Joo, S. W. *et al.* Altered white matter connectivity in patients with schizophrenia: An
investigation using public neuroimaging data from SchizConnect. *PLoS One* **13**,
e0205369, doi:10.1371/journal.pone.0205369 (2018).
23 Almairac, F., Herbet, G., Moritz-Gasser, S., de Champfleur, N. M. & Duffau, H. The
left inferior fronto-occipital fasciculus subserves language semantics: a multilevel
lesion study. *Brain Struct Funct* **220**, 1983-1995, doi:10.1007/s00429-014-0773-1
(2015).
24 Aloia, M. S., Gourovitch, M. L., Weinberger, D. R. & Goldberg, T. E. An investigation
of semantic space in patients with schizophrenia. *J Int Neuropsychol Soc* **2**, 267-273
(1996).
25 Kikinis, Z. *et al.* Genetic contributions to changes of fiber tracts of ventral visual
stream in 22q11.2 deletion syndrome. *Brain Imaging Behav* **7**, 316-325,
doi:10.1007/s11682-013-9232-5 (2013).
26 DeRosse, P. *et al.* Adding insult to injury: childhood and adolescent risk factors for
psychosis predict lower fractional anisotropy in the superior longitudinal fasciculus in
healthy adults. *Psychiatry Res* **224**, 296-302, doi:10.1016/j.pscychresns.2014.09.001
(2014).
27 Davis, K. L. *et al.* White matter changes in schizophrenia: evidence for myelin-related
dysfunction. *Arch Gen Psychiatry* **60**, 443-456, doi:10.1001/archpsyc.60.5.443
(2003).
28 Seki, Y. *et al.* Pretreatment of aripiprazole and minocycline, but not haloperidol,
suppresses oligodendrocyte damage from interferon-gamma-stimulated microglia in
co-culture model. *Schizophr Res* **151**, 20-28, doi:10.1016/j.schres.2013.09.011
(2013).
29 Bi, X. *et al.* Quetiapine prevents oligodendrocyte and myelin loss and promotes
maturation of oligodendrocyte progenitors in the hippocampus of global cerebral
ischemia mice. *J Neurochem* **123**, 14-20, doi:10.1111/j.1471-4159.2012.07883.x
(2012).
30 Zhang, Y. *et al.* Quetiapine enhances oligodendrocyte regeneration and myelin repair
after cuprizone-induced demyelination. *Schizophr Res* **138**, 8-17,
doi:10.1016/j.schres.2012.04.006 (2012).
31 Tishler, T. A. *et al.* Abnormal Trajectory of Intracortical Myelination in Schizophrenia
Implicates White Matter in Disease Pathophysiology and the Therapeutic Mechanism
of Action of Antipsychotics. *Biol Psychiatry Cogn Neurosci Neuroimaging* **3**, 454-462,
doi:10.1016/j.bpsc.2017.03.007 (2018).
32 Zatorre, R. J., Fields, R. D. & Johansen-Berg, H. Plasticity in gray and white:
neuroimaging changes in brain structure during learning. *Nat Neurosci* **15**, 528-536,
doi:10.1038/nn.3045 (2012).
33 Lodygensky, G. A. *et al.* In vivo MRI analysis of an inflammatory injury in the
developing brain. *Brain Behav Immun* **24**, 759-767, doi:10.1016/j.bbi.2009.11.005
(2010).
34 Xie, M. *et al.* Rostrocaudal analysis of corpus callosum demyelination and axon
damage across disease stages refines diffusion tensor imaging correlations with
pathological features. *J Neuropathol Exp Neurol* **69**, 704-716,
doi:10.1097/NEN.0b013e3181e3de90 (2010).
35 Winklewski, P. J. *et al.* Understanding the Physiopathology Behind Axial and Radial
Diffusivity Changes-What Do We Know? *Front Neurol* **9**, 92,
doi:10.3389/fneur.2018.00092 (2018).

515	36	Pasternak, O., Sochen, N., Gur, Y., Intrator, N. & Assaf, Y. Free water elimination and
516		mapping from diffusion MRI. *Magn Reson Med* **62**, 717-730, doi:10.1002/mrm.22055
517		(2009).
518	37	Lebel, C. & Beaulieu, C. Longitudinal development of human brain wiring continues
519		from childhood into adulthood. *J Neurosci* **31**, 10937-10947,
520		doi:10.1523/JNEUROSCI.5302-10.2011 (2011).
521	38	Bava, S. *et al.* Longitudinal characterization of white matter maturation during
522		adolescence. *Brain Res* **1327**, 38-46, doi:10.1016/j.brainres.2010.02.066 (2010).
523	39	Brouwer, R. M. *et al.* White matter development in early puberty: a longitudinal
524		volumetric and diffusion tensor imaging twin study. *PLoS One* **7**, e32316,
525		doi:10.1371/journal.pone.0032316 (2012).
526	40	Giorgio, A. *et al.* Longitudinal changes in grey and white matter during adolescence.
527		*Neuroimage* **49**, 94-103, doi:10.1016/j.neuroimage.2009.08.003 (2010).
528	41	Lebel, C., Walker, L., Leemans, A., Phillips, L. & Beaulieu, C. Microstructural
529		maturation of the human brain from childhood to adulthood. *Neuroimage* **40**, 1044-
530		1055, doi:10.1016/j.neuroimage.2007.12.053 (2008).
531	42	Giorgio, A. *et al.* Changes in white matter microstructure during adolescence.
532		*Neuroimage* **39**, 52-61, doi:10.1016/j.neuroimage.2007.07.043 (2008).
533	43	Douaud, G. *et al.* Schizophrenia delays and alters maturation of the brain in
534		adolescence. *Brain* **132**, 2437-2448, doi:10.1093/brain/awp126 (2009).
535	44	Epstein, K. A. & Kumra, S. White matter fractional anisotropy over two time points in
536		early onset schizophrenia and adolescent cannabis use disorder: A naturalistic
537		diffusion tensor imaging study. *Psychiatry Res* **232**, 34-41,
538		doi:10.1016/j.pscychresns.2014.10.010 (2015).
539	45	Wechsler, D. Wechsler Abbreviated Scale of Intelligence. *The Psychological*
540		*Corporation: Harcourt Brace & Company. New York, NY* (1999).
541	46	Kaufman, J. *et al.* Schedule for Affective Disorders and Schizophrenia for School-Age
542		Children-Present and Lifetime Version (K-SADS-PL): initial reliability and validity data.
543		*J Am Acad Child Adolesc Psychiatry* **36**, 980-988, doi:10.1097/00004583-199707000-
544		00021 (1997).
545	47	Kay, S. R., Fiszbein, A. & Opler, L. A. The positive and negative syndrome scale
546		(PANSS) for schizophrenia. *Schizophr Bull* **13**, 261-276 (1987).
547	48	Shaffer, D. *et al.* A children's global assessment scale (CGAS). *Arch Gen Psychiatry*
548		**40**, 1228-1231 (1983).
549	49	Angold, A. *et al.* Development of a short questionnaire for use in epidemiological
550		studies of depression in children and adolescents. *Int J Method Psych* **5**, 237-249
551		(1995).
552	50	Andreasen, N. C., Pressler, M., Nopoulos, P., Miller, D. & Ho, B. C. Antipsychotic
553		dose equivalents and dose-years: a standardized method for comparing exposure to
554		different drugs. *Biol Psychiatry* **67**, 255-262, doi:10.1016/j.biopsych.2009.08.040
555		(2010).
556	51	Andersson, J. L., Skare, S. & Ashburner, J. How to correct susceptibility distortions in
557		spin-echo echo-planar images: application to diffusion tensor imaging. *Neuroimage*
558		**20**, 870-888, doi:10.1016/S1053-8119(03)00336-7 (2003).
559	52	Smith, S. M. *et al.* Advances in functional and structural MR image analysis and
560		implementation as FSL. *Neuroimage* **23 Suppl 1**, S208-219,
561		doi:10.1016/j.neuroimage.2004.07.051 (2004).
562	53	Andersson, J. L. R. & Sotiropoulos, S. N. An integrated approach to correction for off-
563		resonance effects and subject movement in diffusion MR imaging. *Neuroimage* **125**,
564		1063-1078, doi:10.1016/j.neuroimage.2015.10.019 (2016).
565	54	Smith, S. M. Fast robust automated brain extraction. *Hum Brain Mapp* **17**, 143-155,
566		doi:10.1002/hbm.10062 (2002).
567	55	Behrens, T. E. *et al.* Characterization and propagation of uncertainty in diffusion-
568		weighted MR imaging. *Magn Reson Med* **50**, 1077-1088, doi:10.1002/mrm.10609
569		(2003).

56   Smith, S. M. *et al.* Tract-based spatial statistics: voxelwise analysis of multi-subject
diffusion data. *Neuroimage* **31**, 1487-1505, doi:10.1016/j.neuroimage.2006.02.024
(2006).
57   Nakagawa, S. & Cuthill, I. C. Effect size, confidence interval and statistical
significance: a practical guide for biologists. *Biol Rev Camb Philos Soc* **82**, 591-605,
doi:10.1111/j.1469-185X.2007.00027.x (2007).

Acknowledgements

We thank the study participants and the Youth-TOP clinicians involved in recruitment and assessment at the Norwegian Centre for Mental Disorders (NORMENT) and the Diakonhjemmet Hospital, Oslo, Norway (Runar Elle Smelror, Kirsten Wedervang-Resell, Cecilie Haggag Johannessen, Tarje Tinderholt, Tove Matzen Drachmann). Further, we like to thank Kristine Engen and Brian Frank O'Donnell for proofreading. This work was supported by the Research Council of Norway, grant numbers 223273, 213700, and 250358; the South-Eastern Norway Regional Health Authority, grant numbers 2016-118 and 2017-097; and KG Jebsen Centre for Psychosis Research.

Author Contributions

CB undertook the processing of the imaging data, the statistical analysis, the literature search, interpreted the results and wrote the first draft of the manuscript. TPG helped with the processing of the imaging data. VL was significantly involved in the participant inclusion and data acquisition for Youth-TOP and calculated current and cumulative chlorpromazine equivalents. IA designed the ongoing longitudinal Youth-TOP research study the data is drawn from. IA, AMM and OAA obtained funding and contributed to the data acquisition. All authors contributed to the critical revision of the manuscript and approved the final draft for submission.

Additional Information

Competing interests

The authors declare no competing interests.

Figures

Figure 1| Lower fractional anisotropy (FA) in early onset psychosis (EOP) patients in
comparison to healthy controls. Displayed are significant FWE-corrected TBSS results (red-
yellow, p ≤ 0.01), contrasting EOP patients against healthy controls, overlaid on the study-specific mean
FA skeleton in green and the mean FA image. Results shown underwent threshold-free cluster
enhancement and are corrected for age and sex. CC = corpus callosum, ACR = anterior corona radiata,
SLF = superior longitudinal fasciculus, R = right, L = left.

Figure 2| Extracted scalar diffusion values of the left anterior corona radiata (ACR)
cluster stratified by antipsychotic use, in comparison to the healthy controls (HC). Data
is z-standardized and presented as boxplots for the different scalar diffusion measures overlaid with raw
data points. HC are depicted in white, EOP patients on antipsychotic medication in dark grey and EOP
patients off antipsychotic medication in light grey. EOP = Early onset psychosis, AP = Antipsychotic use
(on = yes, off = no), AD = axial diffusion, FA = fractional anisotropy, RD = radial diffusion. Significant
differences in scalar measures between patient subgroups, based on linear regression models, are
indicated with a star.

Figure 3| Effect size of antipsychotic medication effect on fractional anisotropy (FA) in left anterior corona radiata (ACR). Effect size is presented as Cohen's d with 95% confidence interval (CI) to show the standardized difference of antipsychotic medication status (AP) of early onset psychosis (EOP) patients relative to healthy controls (HC, N = 28). EOP patients on AP (N = 8) are depicted in dark grey and EOP patients off AP (N = 11) are depicted in light grey. Results are controlled for age, sex and BMI.

Tables

Table 1| General sample characteristics.

	EOP patients overall N = 22	Healthy controls N = 33	Statistics group-level
Sex (m/f)	7/15	13/20	$X^2 = 0.33$, p = 0.775
Age at MRI (y)	16.69 ± 1.13	16.08 ± 1.43	t = 1.76, p = 0.085
Range	14.53 – 18.25	12.67 – 18.15	
Handedness (r/l)	18/2	30/2	FET, p = 0.634
Missing N (%)	2 (9.1)	1 (3)	
Parental Education (y)			
Mother	15.05 ± 1.94	16.13 ± 2.09	*t = - 1.94, p = 0.058*
Range	11 – 19	12 – 22	
Missing N (%)	0	2 (6.1)	
Father	14.67 ± 2.87	15.57 ± 2.39	t = - 1.18, p = 0.245
Range	10 – 23	11 – 20	
Missing N (%)	1 (4.5)	3 (9.1)	
IQ	102.74 ± 11.82	101.81 ± 11.32	t = 0.27, p = 0.785
Range	83 – 132	70 – 116	
Missing N (%)	3 (13.63)	1 (3)	
CGAS	44.95 ± 8.57	89.15 ± 6.56	MWU test, **p < 0.001**
Range	32 – 59	75 – 98	
MFQ	29.05 ± 12.79	6.19 ± 6.5	MWU test, **p < 0.001**
Range	5 – 52	0 – 31	
Missing N (%)	1 (4.5)	1 (3)	
BMI (kg/m^2)	21.65 ± 5.6	20.81 ± 2.6	MWU test, p = 0.704
Range	15.4 ± 35.4	16.7 ± 26.0	
Missing N (%)	3 (13.63)	5 (15.15)	
Cannabis (yes/no)	7/15	1/32	FET, **p = 0.005**

All values in mean ± standard deviation, N = Number of participants, m = male, f = female, y = years, r = right, l = left, IQ = Intelligence Quotient, BMI = Body Mass Index, CGAS = Children's Global Assessment Scale, MFQ = Mood and Feelings Questionnaire, MWU = Mann-Whitney-U, FET = Fisher's Exact Test

Table 2| Patient clinical characteristics stratified by antipsychotic medication status

		EOP patients	*EOP patients*	*Statistics*
		Off AP at scan	*On AP at scan*	*patient-level*
		N = 11	*N = 11*	
Sex *(m/f)*		5/6	2/9	FET, p = 0.361
Age Scan *(y)*		15.94 ± 1.49	17.44 ± 0.65	***t = - 4.19, p = 0.0006***
	Range	14.53 – 17.25	16.53 – 18.25	
BMI *(kg/m²)*		20.53 ± 4.32	23.2 ± 7.02	MWU test, p = 0.492
	Range	15.4 – 28.7	16.3 – 35.4	
	Missing N (%)	0	3 (27.3)	
CGAS		46 ± 8	43.91 ± 9.38	t = 0.56, p = 0.580
	Range	34 – 59	32 – 58	
MFQ		30.82 ± 12.6	27.1 ± 13.39	t = 0.65, p = 0.521
	Range	5 – 52	8 – 49	
	Missing N (%)	0	1 (9.1)	
PANSS				
positive		19.09 ± 3.83	17.18 ± 3.84	t = 1.17, p = 0.257
	Range	12 – 25	13 – 26	
negative		22.09 ± 7.13	18.27 ± 6.96	t = 1.27, p = 0.218
	Range	9 – 32	7 – 32	
general		38.73 ± 8.44	36.27 ± 7.89	t = 0.70, p = 0.489
	Range	28 – 54	24 – 52	
Age of Onset *(y)*		14.39 ± 1.92	14.83 ± 2.07	t = - 0.51, p = 0.614
	Range	10 – 16	12 – 17.6	
DUP *(w)*		67.36 ± 68.33	24.73 ± 35.58	MWU test, ***p = 0.003***
	Range	14 – 227	3 – 125	
DUI *(y)*		1.54 ± 1.44	2.61 ± 2.17	MWU test, p = 0.401
	Range	0.47 – 4.53	0.33 – 5.97	
Diagnosis				
SCZ		7	7	
SCA		1	0	
NOS		3	4	
Antipsychotics				
Aripiprazole			5	
Risperidone			3	
Quetiapine			3	
CPZ				
current			272.3 ± 140.83	
	Range		151.52 – 559.44	

					MNI coordinates in mm			
				cumulative	0.08 ± 0.28	21.6 ± 19.13		
				(AP-naïve, N = 9)				
				Range	0 – 0.92	1.69 – 59.28		
				Cannabis use (yes/no)	3/8	4/7		FET, p = 1

Table 3| White matter cluster of reduced fractional anisotropy in early onset psychosis patients relative to healthy controls.

				MNI coordinates in mm				
Cluster	**Region**[1]	**Side**	**Voxels**	**X**	**Y**	**Z**		**t-values**
4	Genu of corpus callosum	L	150	-14	35	6		3.35
3	Anterior corona radiata[2]	L	46	-26	13	14		3.69
2	Superior longitudinal fasciculus	R	12	29	-37	35		4.81
1	Anterior corona radiata (16% Inferior fronto-occipital fasciculus) [3]	L	9	-26	24	16		4.05

[1] Johns Hopkins University International Consortium for Brain Mapping (JHU ICBM)-DTI-81 white matter atlas and JHU white matter tractography atlas (in brackets) were utilized to label significant clusters with specific tract names

[2] 6% uncinate fasciculus/ 5% inferior fronto-occipital fasciculus according to JHU White-Matter Tractography Atlas

[3] 11% anterior thalamic radiation, 8% uncinate fasciculus according to JHU White-Matter Tractography Atlas