1

Structural Basis and Mechanism of TrkA Activation by NGF through Ligand-Induced Rotation of Transmembrane Domain Dimers

María L. Franco[1,4], Kirill D. Nadezhdin[2,3,4], Sergey A. Goncharuk[2,3], Konstantin S Mineev[2,3], Alexander S. Arseniev[2,3]* and Marçal Vilar[1]*

[1]Molecular Basis of Neurodegeneration Unit. Institute of Biomedicine of València (IBV-CSIC). C/ Jaume Roig 11, 46010 València, Spain.

[2]Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry of the Russian Academy of Sciences, Moscow 117997, Russian Federation.

[3]Moscow Institute of Physics and Technology (State University), Institutskiy Pereulok 9, Dolgoprudny, Moscow Region 141700, Russian Federation.

[4]Co-First author.

*co-corresponding authors: aars@nmr.ru , mvilar@ibv.csic.es

Running Title: Mechanism of Activation of TrkA by NGF

Abstract

Trk receptors are essential for the nervous system development. The molecular mechanism of TrkA activation by its ligand NGF is still unsolved. Recent data indicates that at endogenous levels most of TrkA is in an equilibrium monomer-dimer and the binding of NGF induces an increase of the dimer and oligomer forms of the receptor. An unsolved issue is the role of the transmembrane domain (TMD) in the dimerization of TrkA and the structural details of the TMD in the active dimer receptor. We found that TrkA-TMD can form dimers, identified the structural determinants of the dimer interface in the active receptor and validated this interface using site-directed mutagenesis together with functional and cell differentiation studies. As around 20% of TrkA is in an inactive dimeric form in the absence of ligand how NGF binding is able to active this pre-formed dimer is unknown. Using *in vivo* crosslinking we identified a reordering of the extracellular juxtamembrane (JTM) region after ligand binding. This conformational change could be mimicked by replacement of some residues in the JTM with cysteine that form ligand-independent active dimers and reveal a preferred dimer interface. In addition to that, insertion of leucine residues into the TMD helix induces a ligand-independent TrkA activation suggesting that a rotation of the TMD dimers could be behind TrkA activation by NGF. Altogether our data indicate that the transmembrane and juxtamembrane regions of the receptor play a key role in the dimerization and conformational activation of TrkA by NGF.

Introduction

Nerve growth factor (NGF) is a member of the mammalian neurotrophin (NT) protein family implicated in the maintenance and survival of the peripheral and central nervous systems (1-3). NGF is a dimer that interact with two distinct receptors, TrkA a cognate member of the Trk receptor tyrosine kinase family and the p75 neurotrophin receptor, which belongs to the tumor necrosis factor receptor (TNFR) superfamily of death receptors (4-6). TrkA signalling is essential for sensory and sympathetic neuron survival during development (7). Genetic mutations in the NTRK1 gene cause Congenital Insensitivity to Pain with Anhidrosis (CIPA), a rare disease (8), and somatic mutations and chromosomal rearrangements generate aberrant protein fusions with constitutive kinase activation causing several types of cancer (9, 10).

Despite all these important roles, the molecular mechanisms of TrkA activation have been poorly studied in comparison with those of other receptor tyrosine kinase (RTK) family members (11, 12). The first three extracellular domains of TrkA consist of a leucine-rich region, LRR (Trk-d1) that is flanked by two cysteine-rich domains (Trk-d2 and Trk-d3). The fourth and fifth domains (Trk-d4 and Trk-d5) are immunoglobulin (Ig)-like domains, and these are followed by a 30-residue-long linker that connects the extracellular portion of the receptor to the single transmembrane domain and a juxtamembrane intracellular region that is connected to the kinase domain. TrkA is activated by nerve growth factor (NGF) a member of the neurotrophin family (3). The NGF binding domain is located in the Trk-d5(Ig2) domain (13) although other domains also participate in activation by NTs through an unknown mechanism (14, 15).

Two models for TrkA activation are postulated; a ligand-induced dimerization of TrkA monomers and a ligand-induced conformational activation of pre-formed inactive dimers. The first model, which is based on the crystal structure of NGF with the ligand-binding

domain of TrkA (13, 16), assumed that the dimerization of TrkA is solely ligand-mediated

and that receptor-receptor interactions are not present in the absence of its ligand. In the

second model TrkA exists as a pre-formed inactive dimer suggesting receptor-receptor

contacts in the absence of NGF (17, 18). The most recent data using single-particle tracking

(19) and FRET studies (20) suggests that TrkA is, at endogenous levels, predominantly

monomer (80%) and NGF binding induces an increase and a stabilization of the TrkA

dimers and the formation of oligomers, together with a conformational change leading to

kinase activation. This mechanism of activation has been called the "transition model" (20)

and postulates a dynamic transition from a monomer to an inactive dimer to a ligand-bound

active dimer, suggesting that Trk receptors are activated through a combination of the two

mentioned models.

Whatever the model, it is clear that dimerization of TrkA is required for its activation.

Deletion constructs suggested that dimerization of TrkA in the absence of NGF is mediated

by the transmembrane (TM) and by the intracellular domains (ICD) (20). In the case of the

ICD this is supported by the crystal structure of the kinase domain of TrkA that showed the

presence of dimers in the crystallographic unit (21, 22). However the structural determinants

of the TMD dimerization are not know and in this regard, it is important to understand the

conformation of the TrkA-TMD dimer and identify the active dimer interface that may

represent the functional state of the full-length receptor. In addition biochemical data

supporting a conformational activation of TrkA pre-formed dimers are lacking.

In the present work we investigated the role of TrkA-TMDs in its activation, using several

complementary structural and molecular biological approaches. Here we present the

structure of the dimeric TMD in the receptor active state and show that the coupling of

ligand-binding to rotation of the TMD may underlie the TrkA activation by NGF.

Results

Structural basis of TrkA transmembrane domain dimerization.

It has been shown that the isolated TMDs of all human RTKs form dimers in bacterial

membranes (23). In addition functional studies indicate that TMDs play an important role

as a modulator of RTK homodimerization and kinase activation (reviewed in (23-25)).

Switching between two dimerization modes of the transmembrane helix has recently been

described as part of the activation mechanism of the EGF, VEGF and FGF receptors (26-

29). However, to date the role of TrkA-TMD dimerization in TrkA receptor activation has

not been studied in detail. Here we used two different approaches to characterize the

dimerization of the TrkA-TMD. First, to study TrkA-TMD dimerization in a biological

membrane context we used ToxRED, a genetic bacterial system, to assess TrkA

transmembrane dimerization using ToxCAT as a reporter system (30, 31). The TrkA-TMD

self-associated in biological membranes in this system (Figure 1A). To determine the TrkA

amino acid residues implicated in this dimerization we assayed the effect in this assay of

individual mutation of each of the small residues (Ala, Gly or Ser) in the TMD sequence to

a bulky residue such as isoleucine. This approach to disruption of dimerization was similar

to that used in classical studies of glycophorin-A, in which the mutation G83I impaired its

TMD dimerization (31). In our study, less TMD dimerization was observed in the S419I

and G423I mutants compared to wt (Figure 1B), implicating the sequence $S_{419}xxxG_{423}$ as

part of the TrkA-TMD dimerization motif in bacterial membranes.

To obtain a structural insight into TrkA-TMD dimerization we solved the structure of

human TrkA-TMD dimers in lipid micelles using NMR. For this study, the human TrkA-

TMD was produced in a cell-free system (see Experimental Procedures) as previously

described (32). When the peptide is solubilized in dodecylphosphocholine (DPC) micelles

at a lipid-to-protein molar ratio (LPR) of 50:1 the TrkA-TMD is in equilibrium between

monomeric, dimeric and other oligomeric states. The ratio of these states varies as the LPR

value is altered (Figure S1A). We then titrated TrkA-TMD in DPC micelles using a

standard technique in our laboratory (33) to measured the standard free energy of

dimerization (ΔG_0) (Figure S1B). The ΔG_0 value obtained (-1.9±0.2 kcal/mol) suggested

that the TrkA-TMD dimer is quite stable compared to the TMD dimers of other RTKs.

Thus, although its dimerization energy is weaker than that of the VEGFR2 dimer (ΔG_0 = -

2.5 kcal/mol in DPC) (34), it is stronger than that of FGFR3 (ΔG_0 = -1.4 kcal/mol in

DPC/SDS 9:1 mixture) and ErbB4 (ΔG_0 = -1.4 kcal/mol in DMPC/DHPC 1:4 bicelles)

(35). The ^{15}N-HSQC spectrum of ^{15}N labeled-TrkA-TMD (Figure S2) contained the

expected number of cross-peaks, and the good quality of the spectra allowed solving of the

structure of the dimer in DPC micelles (Figures 1C, Figure S3, Figure S4, Figure S5 and

Table S1). The α-helical region of the TrkA-TM dimer starts at G417, ends at N440, and is

~38 Å in length (Figure S3). The crossing angle of the TrkA-TM helices is 40º, and the

minimal distance between two monomers is 8.8 Å (Figure 1C and Table S1). The

hydrophobicity plot and contact surface area of the dimer is shown in Figure S5. The

dimerization interface lies along the sequence motif $L_{424}xxF_{427}A_{428}xxF_{431}$ (Figure 1C and

Figure S5) that is conserved in the TrkA-TMD of several species and also in TrkC but not

in TrkB (Figure 1D).

These analyses of TrkA-TMD dimerization suggested the existence of two possible

dimerization motifs; $S_{419}xxxG_{423}$ and $L_{424}xxF_{427}A_{428}xxF_{431}$, which are located on opposite

faces of the TMD helix (Figure 1C). The latter motif was found in analysis of the

dimerization of TMDs in micelles using a direct approach and in the context of the isolated

domain, while the former motif was observed in analysis of the dimerization of TMDs in

real membranes, using an indirect approach and in the context of a chimeric protein. Thus,

the biological relevance of each motif can be questioned, as the presence of large

extracellular and intracellular globular domains or the lipid environment of the plasma

membrane may favour or hinder a specific interaction interface (36). We therefore used

different functional assays to verify the found dimer interfaces in the context of the full-

length receptor.

Functional identification of the dimer interface upon NGF stimulation.

The state of full-length TrkA was followed by assay of three different aspects of its

activity: dimerization of the receptor, phosphorylation of intracellular tyrosine residues and

neurite differentiation of PC12nnr5 cells. To investigate the dimerization of TrkA, we

individually mutated most of the N-terminal residues of the rat TrkA-TMD to cysteine

(Figure 2A), expressed these constructs in Hela cells, and then measured the amount of

cross-linked species. To facilitate cross-linking via these cysteine residues in the

transmembrane domains we used oxidation with molecular iodine (I_2) as previously

described (37). Such oxidation allows the formation of a disulfide bond between two close

cysteine residues inside the lipid bilayer. Plasma membrane fractions from cells expressing

different single-cysteine mutants were incubated in the absence or presence of NGF,

together with molecular I_2 and were then analyzed by non-reducing SDS-PAGE and

western immunoblotting. As shown in Figures 2B and 2C the mutants F416C, G417C and

A421C formed covalent dimers in the absence of NGF. Upon NGF stimulation the amount

of F416C, G417C and A421 dimers increased and a dimer was newly seen in the V418C

mutant (Figures 2B and 2C). The residues G417, F416 and A421 are very close to each

other in the NMR structure (Figures 2D and 2E), supporting the NMR-derived

conformation as the active state of the TrkA dimer.

To further study the significance of the found interfaces, we mutated the small residues Ala,

Gly and Ser within this region to the bulky Ile residue and we then assayed TrkA activation

upon NGF stimulation (Figures 2F and 2G). The rationale behind this approach was that the

mutation of a small residue to a bulky one on the relevant interface would prevent the

formation of the active dimeric state by inducing steric clashes, and would, therefore,

reduce TrkA activation. To perform this assay, we transfected Hela cells that do not express

endogenous TrkA with these mutants, stimulated these cells with non-saturating

concentrations (10 ng/mL) of NGF, and then assayed TrkA activation by analysis of TrkA

autophosphorylation using western blotting. Upon transfection, two TrkA electrophoretic

bands are present in the TrkA immunoblots of Hela cells; a lower band (approx.110 kDa) of

intracellular immature TrkA that has not completed Golgi-mediated processing of high-

mannose N-glycans (38) and an upper band (approx.140 kDa) with mature sugars that is

expressed in the plasma membrane. Exposure to NGF substantially increased the

phosphorylation of the upper TrkA band as assessed by blotting with a phospho-specific

antibody against the phospho-tyrosine residues of the activation loop, Tyr674 and Tyr675

(P-Y674/5). This autophosphorylation was quantified to follow TrkA activation.

Constitutive (t=0, no NGF added) and ligand-dependent phosphorylation of plasma

membrane-localized TrkA after 5 and 15 minutes were measured. Since overexpression of

TrkA induces ligand-independent autophosphorylation, we first transfected the Hela cells

with increasing concentrations of TrkA to determine a TrkA level that could still be

detected but that displayed no autophosphorylation in the upper band in the absence of

NGF (Figure S6). It is noteworthy that all mutants are expressed at the plasma membrane as

evidenced by both immunofluorescence localization in the absence of Triton X-100 using

an antibody against an epitope in the TrkA N-terminus (Figure S7) and by flow cytometry

(Figure 2H). Out of the seven single-point mutants tested only the A428I substitution

demonstrated a pronounced inhibitory effect on receptor autophosphorylation. A428 is the

only small-chain residue that is found deep in the dimerization interface of the TrkA TMD

conformation determined using NMR, which further supports the relevance of the obtained

NMR structure. The inhibitory effect of A428 substitution on receptor activity was further

enhanced when all three Ala residues that are at least somehow involved in the TMD

dimerization in the NMR-based structure, A421, A425 and A428, were simultaneously

substituted (TrkA-3A/3I).

Lastly, we studied the effect of the same mutations on the NGF-induced differentiation of

transfected PC12nnr5 cells (Figure 2H). Again, the A428I mutant displayed substantial

inhibition of this TrkA activity. Unexpectedly, although the mutation S419I had no effect

on TrkA activation by NGF in these two assays, the mutation G423I did have an impact on

cell differentiation (Figure 2H). This result suggested that G423 is located at a helix

interface that is important in the downstream activation and signalling of TrkA leading to

cell differentiation (See Discussion).

The combined results of the functional assays agree regarding the importance of the NMR-

derived TMD structure for TrkA activation. The critical role of the residue A428 in TrkA

activation by NGF together with its location deep in the dimer interface indicates that the

NMR-based structure corresponds to the mode of TMD interaction in the NGF-induced

active state of the receptor

A specific conformation of extracellular juxtamembrane residues is necessary for

TrkA activation.

Stimulation of Hela cells transfected with TrkA-wt with NGF induces the formation of

TrkA dimers that are crosslinked with BS3 (Figure 3). While it is known that TrkA dimers

are formed in the absence of NGF, no cross-linking was observed without the ligand, which

suggests that ligand binding is accompanied by changes in the conformation of the

extracellular part of the protein. As BS3 reacts only with free amines (the side-chains of

Lys residues or a free N-terminus) we searched for possible sites in TrkA that might have

caused the observed crosslinking. According to the crystal structure of the TrkA/NGF

complex (15) (Figure 3A), there are no lysine residues in the TrkA-ECD that are located in

a position where crosslinking of the side chains of Lys residues of two monomers could

occur. Since BS3 does not cross the plasma membrane, and since we used the full-length

TrkA receptor in our assays, we wondered if the observed BS3 cross-linking was mediated

via crosslinking of K410 and K411 in the extracellular juxtamembrane region (eJTM) of

TrkA (Figure 3A) since this region is not observed in the crystal structure (16). To verify

this hypothesis, we mutated both K410 and K411 to Arg and repeated the initial experiment

using HEK293 cells transfected with this TrkA-KK/RR construct (Figure 3B). No BS3-

induced TrkA crosslinking was observed in the TrkA-KK/RR-transfected cells suggesting

that NGF binding brings this region of the eJTM into close proximity.

We considered that, if NGF indeed induces contacts between these eJTM regions, then we

should be able to mimic this activity of NGF by forcing the dimerization of eJTMs in the

absence of NGF. For this purpose, we individually mutated most of the residues in the

eJTM of TrkA to cysteine and subsequently analyzed the dimerization of these transfected

single point mutants (Figure 4A). After transfection of Hela cells, disulfide dimers were

spontaneously formed in all constructs but the amount of dimers differed between the

various mutants (Figure 4B). The amount of dimer is significantly higher in the positions

D412C and K411C. As a functional assay we then transfected these mutants into HeLa

cells, which do not express endogenous TrkA, and quantified the phosphorylation of the

tyrosines from the kinase activation loop (Y674/675) in the absence and presence of NGF

(Figure 4C). This analysis showed the presence of active dimers (D406C, K410C and

K411C) which are activated constitutively in the absence of NGF, and dimers that are not

active in the absence of NGF (V408C, D412C, E413C and T414C). In general there is a

poor correlation (Figure 4D) between the amount of dimer formation and constitutive

activation, suggesting that dimerization by itself is not enough for TrkA activation.

However the mutants with higher constitutive activation (D406C, K410C and K411C)

showed a good correlation between dimer formation and activation (Figure 4D, green dots,

r^2=0.92). We then transfected some of the active mutants in PC12nnr5 cells. In the absence

of NGF the R405C, K410C and K411C mutants induced the formation of neurites in

PC12nnr5 cells (Figures 4E and 4F) supporting the constitutive activation of these mutants

and suggested that disulfide bond formation through this interface mimics the binding of

NGF. If we assume that the TMD α-helix continues in the juxtamembrane region the

residues R405, K410 and K411 are in one face of the helix (in green in the Figure 4G). By

contrast the residues whose mutation to cysteine induces inactive dimers are located in

another face (in red in the Figure 4G). All the mutants are correctly expressed at the plasma

membrane as found by flow cytometry (Figure 4H).

In the presence of NGF these mutants showed no further activation by NGF (Figure 4C)

suggesting they are fully active and the dimer interface adopted by the cysteine dimers is

similar or identical to the one obtained with NGF binding. By contrast the mutants that

form inactive dimers respond to NGF in a similar manner or even higher than the wt

(Figure 4C), indicating that covalent dimer formation do not preclude the activation by the

ligand. How can NGF activate a covalent dimer if the cysteine are located downstream of

the ligand binding domain? The most plausible explanation is that NGF can induce the

dimerization of two pre-formed inactive dimers as shown in the Figure 4H.

In summary, our results supports the notion of the insufficiency of TrkA dimerization alone

for receptor activation and support the existence of an preferred active dimer interface that

continues through the TMD.

An Induced rotation activates TrkA in the absence of NGF

It is known that overexpression of TrkA induce a ligand-independent activation of TrkA.

However the levels of autophosphorylation are lower than in the presence of NGF. The

western blotting shown in Figure 5A shows the autophosphorylation of the intracellular

tyrosines of TrkA (Tyr-674/5, Tyr-490 and Tyr-751) and downstream signaling (P-ERK)

due to TrkA overexpression (absence of NGF) and in the presence of NGF. The blots

indicate that, although the overexpression of TrkA induces a ligand-independent activation

(probably by facilitating receptor dimerization), the presence of NGF is required for a

higher and complete activation of the receptor. The complete sequence of the events caused

by the ligand is also of great importance for receptor activation. As we have shown above,

ligand binding brings eJTM regions into proximity at certain specific positions. In turn,

such proximity can either result in a mutual rotation of the receptor TMDs or change the

spacing between the N-termini of the TMD helices in the dimer. We found two dimer

interfaces in TrkA-TMD that reside on opposite sides of the α-helix suggesting the

possibility of a rotation from one interface to the other upon NGF binding. If this model

were correct, mutual rotation of two motifs on the surface of the TM helix should activate

TrkA in the absence of NGF or increase its basal activation by overexpression. To test this

mechanism we introduced a different number of leucine residues into the TrkA-TMD and

analyzed the resulting TrkA activation in the absence of NGF (Figure 5B). The insertion of

each Leu should rotate the intracellular region around an angle of approximately 100º.

Thus, Leu insertion allows evaluation of whether a change in the rotation angle of the

intracellular domain plays any role in TrkA activation (Figure 5C). We transfected the

constructs TrkA-ins1L, TrkA-ins2L, TrkA-ins3L and TrkA-ins4L that included 1 to 4

inserted leucines respectively. The insertion of one Leu, TrkA-ins1Leu, significantly

increased both the constitutive activation of TrkA in transfected Hela cells compared to that

of transfected TrkA-wt (Figure 5D and 5E) and the differentiation of PC12nnr5 cells

compared to wt-transfected cells (Figure 5F) in a ligand-independent manner. Equal levels

of all constructs were expressed at the plasma membrane as determined using flow

cytometry (Figure S8). These data indicate that a rotation of the TMD is likely to underlie

TrkA activation.

Discussion

TrkA belongs to a subfamily of RTKs that includes the other family members TrkB and

TrkC. These RTKs are essential for the formation of the nervous system and mediate a

variety of cellular responses in normal biological processes and in pathological states (39).

An understanding of their mechanism of action is necessary to facilitate the design of new

pharmacological agents targeted to the processes in which they play a role.

Both TrkA and TrkB are in a pre-formed dimeric state in the absence of ligand and are

inactive in this state (18, 40). Using *in vivo* crosslinking and fluorescence complementation

assays in PC12 cells that express endogenous TrkA, Maruyama et al. found that TrkA, as

well as TrkB, form inactive dimers in the absence of NGF (18, 40). Marchetti et al., using

single-tracking of TrkA receptors, recently found that around 20% of plasma membrane-

expressed TrkA receptors are in a pre-dimer state in the absence of NGF (19). Additionally,

the existence of pre-formed dimers of full-length TrkA expressed in the plasma membrane

of Xenopus oocytes was shown by freeze-fracture electron microscopy (17). A recent study

has provided a further evidence for the existence of pre-formed dimers in the Trk family

(20). The laboratory of Dr. Hristova found that TrkA, as well TrkB and TrkC, exist in an

equilibrium monomer-dimer state in the absence of ligand depending on the expression

levels (20). That study also showed that the TMD and the intracellular domains of TrkA

contribute significantly to its unliganded dimerization. In the same work the authors

suggested that the equilibrium constant of TrkA dimerization indicated that endogenous

levels, for instance in PC12 cells, the most predominant form of TrkA is a monomer (80%)

in equilibrium with a 20% of pre-formed inactive dimers, supporting the data from

Marchetti et al. (19). Single-particle tracking showed that TrkA monomer form dimers and

oligomers upon NGF stimulation. Altogether TrkA activation follows a *transition* from a

monomer to a pre-former inactive dimer and to a full active ligand-bound dimer and

oligomers, in what has been called the "*transition model of RTK activation*" (20).

In this regard, in the present work we posed two major questions: (1) what are the

conformations of TrkA TMDs in the active/inactive states? and (2) how is the coupling

between the ligand binding and conformational switch accomplished? To answer the first

question we employed structural characterization using NMR spectroscopy together with

mutagenesis studies, disulfide cross-linking and functional assays. We described the high-

resolution NMR structure of the TrkA transmembrane domain dimer, which is the first such

description of a neurotrophin receptor of the Trk family. The obtained spatial structure was

verified using functional assays and cross-linking, which confirmed both the relevance of

this structure for TrkA activity and assignation of the found dimer conformation to the

receptor active state. This result, combined with the crystal structure of the extracellular

domain of the TrkA complex with NGF (16, 41), the crystal structure of the TrkA inactive

kinase domain (21, 22) and the recently reported structures of the entire TrkA intracellular

region (42, 43) provides an almost complete picture of the full length Trk receptor family,

lacking only the structure of the small JTM regions.

Additionally, we found another possible TMD dimerization interface using a genetic assay,

and functional tests indicated that this interface was also relevant for TrkA activation.

Although destruction of this interface by replacement of a small amino acid residue with a

bulky residue inhibited the dimerization in the bacterial membranes, it does not affect the

activation of TrkA by NGF, suggesting that this dimer interface is not populated in the full-

length TrkA receptor at least in the ligand-bound state. The structural data indicates that

this interface is an opposite side as the active dimer interface. Interestingly in an ideal α-

helix connected to the TMD the residues from the eJTM that form inactive dimers when

mutated to cysteine are located in the same helix side than the SxxxG motif. This suggests

that rotation of the eJTM will also rotate the TMD. Based on this it is possible that the

SxxxG interface forms part of an inactive pre-dimer state, although we cannot detect the

formation of dimers when the SxxxG motif is mutated to cysteine. One possibility is that, in

the inactive state, the distance between the SxxxG motifs are longer than the required for

the disulfide bond formation. This could be due to the presence of steric clashes from the

TrkA ICD or ECD, that are absent in the ToxRed constructs. However the finding that the

mutation of the G423I reduces significantly the differentiation of PC12 cells with NGF but

not the activation of the kinase domain (as shown by the phosphorylation of the Tyr-

674/675) is intriguing. PC12 cells differentiation needs of a sustained MAPK activation to

differentiate and it is plausible that a higher activation of TrkA is reached by the lateral

association of liganded dimers; i.e oligomerization of TrkA dimers. Actually the formation

of TrkA oligomers by NGF has been already described by single-particle tracking in

neuroblastoma cells (19). And oligomerization of ligand-bound RTKs seems a general

mechanism to regulate differential intracellular signalling (44). The important role of G423

in cell differentiation and its location in an opposite side of the α-helix to the active dimer

interface, suggests that G423 may participate in a protein-protein interface corresponding to

a supra-dimer (oligomer) form of the TrkA receptor that is populated upon NGF binding

(19). In this context further investigation on the TrkA oligomer formation is warranted.

To answer the second question posed above, we investigated the role of the eJTM regions

of the TrkA receptor upon NGF binding in receptor activation and dimerization. We

showed that full-length TrkA receptors can be activated by specific single-point mutations

in the eJTM, in which the position of the mutation relative to the TMD was more important

for activation than the dimerization propensity of the mutant.

Bearing all of these findings in mind, we propose a mechanism of receptor activation that is

outlined in Figure 6. In the model of the Figure 6, NGF binding to the Ig2 will induce a

rotation of the eJTM to facilitate the dimerization of the TMD via a preferred interface. This

model is supported by our data that showed that cysteine mutants in the eJTM in some

specific positions can activate TrkA without ligand and NGF binding brings the N-termini of

the α-helices into close proximity. The insertion of Leu residues downstream of the TMD

dimerization motif activated TrkA in the absence of ligand. This suggests that rotation of the

downstream domain may be behind the activation of the kinase domain. The rotational

mechanism of RTK activation has been proposed by other authors (45, 46). In this model the

ligand would induce a rotation of the TMD dimer interface that will re-orient the kinase

domains in order to facilitate the trans-phosphorylation. Although our results support this

model of activation other alternative possibilities may exist. For instance the insertion of

extra-residues increases the TMD length and could induce a piston-like mechanism of

activation by a pulling-down force. However an increase in the length by two, three or four

residues should also activate the kinase and this was not the case, as only the Ins1L mutant

showed activation. Also as the insertion of the residues are located into the TMD it may alter

the dimer interface of the TMD dimer leading to the formation of another dimer interface

compatible with a higher activation of the kinase domains. Although we cannot discard this

possibility, in the constructs we made the Leu residues are inserted upstream of the active

dimer interface in order to not alter the dimer interface found by NMR studies. In any case

the results obtained from the insertion mutagenesis suggest a conformational activation of

TrkA activation. Future structural and functional studies of the TrkA intracellular domain

should address how the kinase domain is activated if they are connected to the TMD by a

flexible intracellular juxtamembrane region.

In summary our data supports a conformational activation of TrkA by NGF and we have

provided functional and structural evidences of the roles played by the TMD in TrkA

dimerization and activation by NGF.

References

1.　Chao, M. V., Neurotrophins and their receptors: a convergence point for many signalling
pathways. *Nat. Rev. Neurosci.* **2003,** *4* (4), 299-309.
2.　Bothwell, M., NGF, BDNF, NT3, and NT4. *Handb. Exp. Pharmacol.* **2014,** *220*, 3-15.
3.　Ceni, C.; Unsain, N.; Zeinieh, M. P.; Barker, P. A., Neurotrophins in the regulation of cellular
survival and death. *Handb. Exp. Pharmacol.* **2014,** *220*, 193-221.
4.　Friedman, W. J.; Greene, L. A., Neurotrophin signaling via Trks and p75. *Exp. Cell Res.* **1999,**
253 (1), 131-42.
5.　Bothwell, M., Recent advances in understanding neurotrophin signaling. *F1000Res* **2016,** *5*.
6.　Vilar, M., Structural Characterization of the p75 Neurotrophin Receptor: A Stranger in the
TNFR Superfamily. *Vitam Horm* **2017,** *104*, 57-87.
7.　Smeyne, R. J.; Klein, R.; Schnapp, A.; Long, L. K.; Bryant, S.; Lewin, A.; Lira, S. A.;
Barbacid, M., Severe sensory and sympathetic neuropathies in mice carrying a disrupted Trk/NGF
receptor gene. *Nature* **1994,** *368* (6468), 246-9.
8.　Indo, Y.; Tsuruta, M.; Hayashida, Y.; Karim, M. A.; Ohta, K.; Kawano, T.; Mitsubuchi, H.;
Tonoki, H.; Awaya, Y.; Matsuda, I., Mutations in the TRKA/NGF receptor gene in patients with
congenital insensitivity to pain with anhidrosis. *Nat. Genet.* **1996,** *13* (4), 485-8.
9.　Martin-Zanca, D.; Hughes, S. H.; Barbacid, M., A human oncogene formed by the fusion of
truncated tropomyosin and protein tyrosine kinase sequences. *Nature* **1986,** *319* (6056), 743-8.
10.　Cocco, E.; Scaltriti, M.; Drilon, A., NTRK fusion-positive cancers and TRK inhibitor therapy.
Nat. Rev. Clin. Oncol. **2018,** *15* (12), 731-747.

11.   Endres, N. F.; Barros, T.; Cantor, A. J.; Kuriyan, J., Emerging concepts in the regulation of the
EGF receptor and other receptor tyrosine kinases. *Trends Biochem. Sci.* **2014,** *39* (10), 437-46.
12.   Lemmon, M. A.; Schlessinger, J., Cell signaling by receptor tyrosine kinases. *Cell* **2010,** *141*
(7), 1117-34.
13.   Wiesmann, C.; Ultsch, M. H.; Bass, S. H.; de Vos, A. M., Crystal structure of nerve growth
factor in complex with the ligand-binding domain of the TrkA receptor. *Nature* **1999,** *401* (6749), 184-
8.
14.   Arevalo, J. C.; Conde, B.; Hempstead, B. I.; Chao, M. V.; Martin-Zanca, D.; Perez, P., A novel
mutation within the extracellular domain of TrkA causes constitutive receptor activation. *Oncogene*
2001, *20* (10), 1229-34.
15.   Zaccaro, M. C.; Ivanisevic, L.; Perez, P.; Meakin, S. O.; Saragovi, H. U., p75 Co-receptors
regulate ligand-dependent and ligand-independent Trk receptor activation, in part by altering Trk
docking subdomains. *J. Biol. Chem.* **2001,** *276* (33), 31023-9.
16.   Wehrman, T.; He, X.; Raab, B.; Dukipatti, A.; Blau, H.; Garcia, K. C., Structural and
mechanistic insights into nerve growth factor interactions with the TrkA and p75 receptors. *Neuron*
2007, *53* (1), 25-38.
17.   Mischel, P. S.; Umbach, J. A.; Eskandari, S.; Smith, S. G.; Gundersen, C. B.; Zampighi, G. A.,
Nerve growth factor signals via preexisting TrkA receptor oligomers. *Biophys. J.* **2002,** *83* (2), 968-76.
18.   Shen, J.; Maruyama, I. N., Nerve growth factor receptor TrkA exists as a preformed, yet
inactive, dimer in living cells. *FEBS Lett.* **2011,** *585* (2), 295-9.
19.   Marchetti, L.; Callegari, A.; Luin, S.; Signore, G.; Viegi, A.; Beltram, F.; Cattaneo, A., Ligand
signature in the membrane dynamics of single TrkA receptor molecules. *J. Cell Sci.* **2013,** *126* (Pt 19),
4445-56.
20.   Ahmed, F.; Hristova, K., Dimerization of the Trk receptors in the plasma membrane: effects of
their cognate ligands. *Biochem. J.* **2018,** *475* (22), 3669-3685.
21.   Artim, S. C.; Mendrola, J. M.; Lemmon, M. A., Assessing the range of kinase autoinhibition
mechanisms in the insulin receptor family. *Biochem. J.* **2012,** *448* (2), 213-20.
22.   Bertrand, T.; Kothe, M.; Liu, J.; Dupuy, A.; Rak, A.; Berne, P. F.; Davis, S.; Gladysheva, T.;
Valtre, C.; Crenne, J. Y.; Mathieu, M., The crystal structures of TrkA and TrkB suggest key regions for
achieving selective inhibition. *J. Mol. Biol.* **2012,** *423* (3), 439-53.
23.   Bocharov, E. V.; Bragin, P. E.; Pavlov, K. V.; Bocharova, O. V.; Mineev, K. S.; Polyansky, A.
A.; Volynsky, P. E.; Efremov, R. G.; Arseniev, A. S., The Conformation of the Epidermal Growth
Factor Receptor Transmembrane Domain Dimer Dynamically Adapts to the Local Membrane
Environment. *Biochemistry* **2017,** *56* (12), 1697-1705.
24.   Li, E.; Hristova, K., Role of receptor tyrosine kinase transmembrane domains in cell signaling
and human pathologies. *Biochemistry* **2006,** *45* (20), 6241-51.
25.   Li, E.; Hristova, K., Receptor tyrosine kinase transmembrane domains: Function, dimer
structure and dimerization energetics. *Cell adhesion migration* **2010,** *4* (2), 249-54.
26.   Jura, N.; Endres, N. F.; Engel, K.; Deindl, S.; Das, R.; Lamers, M. H.; Wemmer, D. E.; Zhang,
X.; Kuriyan, J., Mechanism for activation of the EGF receptor catalytic domain by the juxtamembrane
segment. *Cell* **2009,** *137* (7), 1293-307.
27.   King, C.; Hristova, K., Direct measurements of VEGF:VEGFR2 binding affinities reveal the
coupling between ligand binding and receptor dimerization. *J. Biol. Chem.* **2019**.
28.   Sarabipour, S.; Ballmer-Hofer, K.; Hristova, K., VEGFR-2 conformational switch in response
to ligand binding. *eLife* **2016,** *5*, e13876.
29.   Sarabipour, S.; Hristova, K., Mechanism of FGF receptor dimerization and activation. *Nature*
communications **2016,** *7*, 10262.

30.   Berger, B. W.; Kulp, D. W.; Span, L. M.; DeGrado, J. L.; Billings, P. C.; Senes, A.; Bennett, J.
S.; DeGrado, W. F., Consensus motif for integrin transmembrane helix association. *Proc. Natl. Acad.*
Sci. U. S. A. **2010,** *107* (2), 703-8.
31.   Russ, W. P.; Engelman, D. M., TOXCAT: a measure of transmembrane helix association in a
biological membrane. *Proc. Natl. Acad. Sci. U. S. A.* **1999,** *96* (3), 863-8.
32.   Nadezhdin, K. D.; Garcia-Carpio, I.; Goncharuk, S. A.; Mineev, K. S.; Arseniev, A. S.; Vilar,
M., Structural Basis of p75 Transmembrane Domain Dimerization. *J. Biol. Chem.* **2016,** *291* (23),
12346-57.
33.   Mineev, K. S.; Lesovoy, D. M.; Usmanova, D. R.; Goncharuk, S. A.; Shulepko, M. A.;
Lyukmanova, E. N.; Kirpichnikov, M. P.; Bocharov, E. V.; Arseniev, A. S., NMR-based approach to
measure the free energy of transmembrane helix-helix interactions. *Biochim. Biophys. Acta* **2014,** *1838*
(1 Pt B), 164-72.
34.   Mineev, K. S.; Goncharuk, S. A.; Arseniev, A. S., Toll-like receptor 3 transmembrane domain
is able to perform various homotypic interactions: an NMR structural study. *FEBS Lett.* **2014,** *588* (21),
3802-7.
35.   Bocharov, E. V.; Mineev, K. S.; Goncharuk, M. V.; Arseniev, A. S., Structural and
thermodynamic insight into the process of "weak" dimerization of the ErbB4 transmembrane domain
by solution NMR. *Biochim Biophys Acta* **2012,** *1818* (9), 2158-70.
36.   Cymer, F.; Veerappan, A.; Schneider, D., Transmembrane helix-helix interactions are
modulated by the sequence context and by lipid bilayer properties. *Biochim. Biophys. Acta* **2012,** *1818*
(4), 963-73.
37.   Hughson, A. G.; Lee, G. F.; Hazelbauer, G. L., Analysis of protein structure in intact cells:
crosslinking in vivo between introduced cysteines in the transmembrane domain of a bacterial
chemoreceptor. *Protein Sci.* **1997,** *6* (2), 315-22.
38.   Schecterson, L. C.; Hudson, M. P.; Ko, M.; Philippidou, P.; Akmentin, W.; Wiley, J.;
Rosenblum, E.; Chao, M. V.; Halegoua, S.; Bothwell, M., Trk activation in the secretory pathway
promotes Golgi fragmentation. *Mol. Cell. Neurosci.* **2010,** *43* (4), 403-13.
39.   Deinhardt, K.; Chao, M. V., Trk receptors. *Handb. Exp. Pharmacol.* **2014,** *220,* 103-19.
40.   Shen, J.; Maruyama, I. N., Brain-derived neurotrophic factor receptor TrkB exists as a
preformed dimer in living cells. *J Mol Signal* **2012,** *7* (1), 2.
41.   Ultsch, M. H.; Wiesmann, C.; Simmons, L. C.; Henrich, J.; Yang, M.; Reilly, D.; Bass, S. H.;
de Vos, A. M., Crystal structures of the neurotrophin-binding domain of TrkA, TrkB and TrkC. *J. Mol.*
Biol. **1999,** *290* (1), 149-59.
42.   Furuya, N.; Momose, T.; Katsuno, K.; Fushimi, N.; Muranaka, H.; Handa, C.; Ozawa, T.;
Kinoshita, T., The juxtamembrane region of TrkA kinase is critical for inhibitor selectivity. *Bioorg.*
Med. Chem. Lett. **2017,** *27* (5), 1233-1236.
43.   Su, H. P.; Rickert, K.; Burlein, C.; Narayan, K.; Bukhtiyarova, M.; Hurzy, D. M.; Stump, C. A.;
Zhang, X.; Reid, J.; Krasowska-Zoladek, A.; Tummala, S.; Shipman, J. M.; Kornienko, M.; Lemaire, P.
A.; Krosky, D.; Heller, A.; Achab, A.; Chamberlin, C.; Saradjian, P.; Sauvagnat, B.; Yang, X.; Ziebell,
M. R.; Nickbarg, E.; Sanders, J. M.; Bilodeau, M. T.; Carroll, S. S.; Lumb, K. J.; Soisson, S. M.;
Henze, D. A.; Cooke, A. J., Structural characterization of nonactive site, TrkA-selective kinase
inhibitors. *Proc. Natl. Acad. Sci. U. S. A.* **2017,** *114* (3), E297-E306.
44.   Needham, S. R.; Roberts, S. K.; Arkhipov, A.; Mysore, V. P.; Tynan, C. J.; Zanetti-Domingues,
L. C.; Kim, E. T.; Losasso, V.; Korovesis, D.; Hirsch, M.; Rolfe, D. J.; Clarke, D. T.; Winn, M. D.;
Lajevardipour, A.; Clayton, A. H.; Pike, L. J.; Perani, M.; Parker, P. J.; Shan, Y.; Shaw, D. E.; Martin-
Fernandez, M. L., EGFR oligomerization organizes kinase-active dimers into competent signalling
platforms. *Nature communications* **2016,** *7,* 13307.

45.   Bell, C. A.; Tynan, J. A.; Hart, K. C.; Meyer, A. N.; Robertson, S. C.; Donoghue, D. J.,
Rotational coupling of the transmembrane and kinase domains of the Neu receptor tyrosine kinase. *Mol*
Biol Cell **2000,** *11* (10), 3589-99.
46.   Maruyama, I. N., Activation of transmembrane cell-surface receptors via a common
mechanism? The "rotation model". *Bioessays* **2015,** *37* (9), 959-67.
47.   Aoki, M.; Matsuda, T.; Tomo, Y.; Miyata, Y.; Inoue, M.; Kigawa, T.; Yokoyama, S.,
Automated system for high-throughput protein production using the dialysis cell-free method. *Protein*
Expr. Purif. **2009,** *68* (2), 128-36.
48.   Kai, L.; Roos, C.; Haberstock, S.; Proverbio, D.; Ma, Y.; Junge, F.; Karbyshev, M.; Dotsch, V.;
Bernhard, F., Systems for the cell-free synthesis of proteins. *Methods Mol. Biol.* **2012,** *800*, 201-25.
49.   Schwarz, D.; Junge, F.; Durst, F.; Frolich, N.; Schneider, B.; Reckel, S.; Sobhanifar, S.; Dotsch,
V.; Bernhard, F., Preparative scale expression of membrane proteins in Escherichia coli-based
continuous exchange cell-free systems. *Nat. Protoc.* **2007,** *2* (11), 2945-57.
50.   Goncharuk, S. A.; Goncharuk, M. V.; Mayzel, M. L.; Lesovoy, D. M.; Chupin, V. V.;
Bocharov, E. V.; Arseniev, A. S.; Kirpichnikov, M. P., Bacterial Synthesis and Purification of Normal
and Mutant Forms of Human FGFR3 Transmembrane Segment. *Acta naturae* **2011,** *3* (3), 77-84.
51.   Keller, R. L. J., The Computer Aided Resonance Assignment Tutorial. CANTINA Verlag:
Goldau, Switzerland, 2004.
52.   Cavanagh, J., Fairbrother, W.J., Palmer, A.G. and Skelton, N.J., *Protein NMR spectroscopy:*
principles and practice. 2nd ed.; Academic Press,: San Diego, CA, USA, 2006.
53.   Güntert, P., Automated NMR protein structure calculation. *Prog. NMR Spectrosc* **2003,** *43*, 105.
54.   Shen, Y.; Bax, A., Protein backbone and sidechain torsion angles predicted from NMR
chemical shifts using artificial neural networks. *J. Biomol. NMR* **2013,** *56* (3), 227-41.
55.   Koradi, R.; Billeter, M.; Wuthrich, K., MOLMOL: a program for display and analysis of
macromolecular structures. *J. Mol. Graph.* **1996,** *14* (1), 51-5, 29-32.
56.   Polyansky, A. A.; Volynsky, P. E.; Efremov, R. G., Multistate organization of transmembrane
helical protein dimers governed by the host membrane. *J. Am. Chem. Soc.* **2012,** *134* (35), 14390-400.
549
550

Material and Methods

DNA constructs

A plasmid encoding rat TrkA with an N-terminal HA tag was kindly provided by Dr. Y

Barde. All TrkA mutants and constructs were derived from this plasmid. Mutagenesis was

done using the Site-directed mutagenesis kit (Agilent) according to the manufacturer's

protocol. The oligonucleotide sequences of all of the constructs are available upon request.

All DNA constructs were sequenced using local facilities.

Cell culture and transfection

Hela cells, which do not express endogenous TrkA, were cultured in DMEM medium (Fisher) supplemented with 10% FBS (Fisher) at 37 °C in a humidified atmosphere with 5% CO_2. PC12 and PC12nnr5 cells were cultured in DMEM with 10% FBS and 5% horse serum (Fisher). Transfection of Hela cells was performed using polyethylenimine (PEI; Sigma) at a concentration of 1-2 µg/µl. The use of PEI as the transfection reagent for Hela cells resulted in suboptimal transfection (10-15% of cells transfected) and in the expression of only a small amount of TrkA in the cells. In contrast, when the same PEI/DNA ratio was used for transfection of Hek293 cells TrkA was expressed in higher amounts and ligand-independent activation of TrkA was seen. A concentration of 500-1000 ng of DNA per 10 cm cell plate was used for the TrkA activation experiments. Twenty four hours after transfection the cells were lifted and re-plated into 12-well plates at a density of 100,000 cells per well. By using this procedure the percentage of cells transfected was identical in all the wells. Forty eight hours after transfection the cells were starved in serum free medium for 2 h and were then stimulated with NGF (Alomone) at the indicated concentrations and time intervals. Cells were lysed with TNE buffer (Tris-HCl pH 7.5, 150 mM NaCl, 1 mM EDTA) supplemented with 1% triton X-100 (Sigma), protease inhibitors (Roche), 1 mM PMSF (Sigma), 1 mM sodium orthovanadate (Sigma), and 1 mM sodium fluoride (Sigma). In the experiments involving the TrkA cysteine mutants, 10 mM iodoacetamide (Sigma) was added to the lysis buffer. Lysates were kept on ice for 10 minutes and centrifuged at 12,000 g for 15 minutes in a tabletop centrifuge. The protein level of the lysates was quantified using a Bradford kit (Pierce) and lysates were analyzed by SDS-PAGE

Western blot analysis

Cellular debris was removed by centrifugation at <u>12,000 g </u>for 15 minutes and the protein

level of cell lysates was quantified using the Bradford assay (Pierce). Proteins were resolved

in SDS-PAGE gels and transferred to nitrocellulose membranes that were incubated

overnight at 4 ºC with one of the following antibodies: mouse monoclonal anti-HA (1:2000,

Sigma); rabbit polyclonal MBP-probe (1:1000, Santa Cruz); rabbit anti-phosphoTyr674/5

(1:1000, Cell Signaling); rabbit anti-TrkA (1:1000, Millipore). Following incubation with the

appropriate secondary antibody, membranes were imaged and bands quantified using

enhanced chemiluminescence and autoradiography.

ToxRED assays

Bacterial colonies were inoculated into LB medium (with 50 µg/ml ampicillin). Fresh LB

cultures (with 50 µg/ml ampicillin) were inoculated from fresh plates and were grown at 37

°C until they reached an O.D. of approximately 600≈0.6. They were then harvested by

centrifugation. Cells were subsequently lysed in lysis buffer (TNE, see recipe above) under

gentle conditions to avoid mCherry denaturation. After transfer to 1.5-mL Eppendorf tubes,

the mixture was incubated at room temperature with gentle agitation for 30 min. Samples

were then centrifuged for 10 min at $12000 \times g$ to remove cell debris and to clarify the

supernatants for analysis. Aliquots (150 µl) of clarified supernatant were transferred to black,

opticlear 96-well plates and mCherry emission spectra were collected using a plate reader

(Tecan, Maennedorf, CH) with an excitation wavelength of 587 nm and emission

wavelengths of 610–650 nm. Subsequently, aliquots were transferred from black to clear 96-

well plates and the absorbance was measured from 450 to 750 nm. Measurements of

mCherry and of construct expression were performed from at least 10 different colonies and

were normalized for the relative expression level of each construct using western blotting

with the anti-MBP antibody. For western blots, samples were mixed with equal volumes of

2× SDS-PAGE sample buffer, heated to 95 °C for 10 min, separated on 10% (w/v)

polyacrylamide mini-gels, and were blotted onto nitrocellulose membranes that were probed

with anti-MBP antibody. To analyze disulfide bond formation bacterial colonies were lysed

using TNE plus 10 mM iodoacetamide and were analyzed using non-reducing SDS-PAGE.

TrkA-TMD constructs for cell-free expression

The gene encoding the transmembrane domain of the human TrkA-TM

(MK$_{410}$KDETPFGVSVAVGLAVFACLFLSTLLLVLNK$\underline{A}$GRRNK$_{447}$) was amplified by PCR

from six chemically synthesized oligonucleotide templates (Evrogen, Russia) whose

sequences partially overlapped along its sequence. The PCR products were cloned into a

pGEMEX-1 vector by three-component ligation using the NdeI, AatII and BamHI restriction

sites.

Cell-free gene expression.

A bacterial S30 cell-free extract was prepared from a 10 L culture of the E. coli

Rosetta(DE3)pLysS strain according to a previously described protocol. The S30 cell extract

was stored in 500 μl aliquots at -80 °C. The continuous exchange mode of preparation using

a 12.5 kDa membrane was used in this study. Preparative-scale reactions (2-3 ml of reaction

mixture) were carried out in 50 mL tubes. Optimal reaction conditions such as Mg2+ and K+

concentrations, the ratio of the reaction mixture (RM) to the feeding mixture (FM) or the

DNA concentration were established using homemade reactors based on the Mini-CECF-

Reactor previously described (47, 48). The final reaction mixture was: a standard FM:RM

ratio of 8:1 and a cell-free reaction mixture containing 100 mM HEPES ∕0.83 mM EDTA

with KOH added to achieve a pH of 8.0, 0.1 mg/mL Folinic acid, 20 mM acetyl phosphate,

1.2 mM ATP, 0.8 mM each of G/C/UTP, 2 mM 1,4-dithiothreitol, 0.05% sodium azide, 2%

PEG-8000, 20 mM magnesium acetate, 270 mM potassium acetate, 60 mM creatine

phosphate, 1 mM each of 20 amino acids or 0.25% of a 20 amino-acid mix (CIL, USA), 1

tablet / 50 mL of complete protease inhibitor (Roche, Switzerland), 0.5 mg/mL E.coli tRNA

(Roche, Switzerland), 0.25 mg/mL creatine kinase from rabbit muscle (Roche, Switzerland),

0.05 mg/mL T7 RNA polymerase prepared using a previously described protocol (49), 0.1

U/μL Ribolock (Fermentas), 0.02 μg/μL plasmid DNA, and 30% S30 cell-free extract. All

reagents were provided by Sigma unless otherwise specified. Plasmid DNA was purified

using a Promega MaxiPrep kit. Reactions were conducted overnight at 34 °C and in an

Innova 44R shaker (New Brunswick) set at 150 rpm.

Protein purification.

The cell-free reaction mixture was diluted three-times with buffer A (50 mM Tris pH 8.0 and

200 mM NaCl). After 10 minutes of incubation the mixture was centrifuged for 10 min at

18000 g at room temperature. The precipitate was washed consecutively with buffer A

containing 30 μg/ml RNAse A (Fermentas) and buffer B (50 mM Tris pH 8.0 and 100 mM

NaCl). The target protein was solubilized with 200 μl buffer B containing 1% lauryl

sarcosine. After each step the protein was centrifuged for 10 min at <u>18.000</u> g at room

temperature and aliquots of the supernatant were analyzed using 12,5% Tricine SDS-PAGE

(53). The clarified protein solution was applied onto a 10/300 Tricorn column prepacked

with Superdex 200 (GE Healthcare) and pre-equilibrated with buffer B containing 0.2%

lauryl sarcosine. Protein-containing fractions were combined and precipitated using the

TCA/acetone procedure (50).

Preparation of NMR samples in a membrane mimetic media

The so-called "isotopic-heterodimer" (1:1 mixture of unlabeled and 15N/13C-labeled peptides)

samples were prepared corresponding to the TrkA-TMD construct in order to solve its

structure. The powder containing the peptides of both samples was first dissolved in a 1:1

trifluoroethanol–H_2O mixture with the addition of deuterated DPC (d_{38}, 98%, CIL) and

phosphate buffer, and was then kept for several minutes in an ultrasound bath and

lyophilized. Subsequently, the dried samples were dissolved in 350 µl of a 9:1 $H_2O:D_2O$

mixture. To attain a uniform micelle size and uniform distribution of the peptide throughout

the micelles, the samples were sonicated in an ultrasound bath for several minutes until the

solution was completely transparent. The TrkA-TMD concentration in the isotopic-

heterodimer sample was 1.9 mM, and other conditions were: LPR 50:1, pH 5.9, and 20 mM

phosphate buffer. Samples were placed in Shigemi NMR tubes with a glass plunger.

Selective-residue labeling was implemented to avoid peaks overlapping while processing the

NMR spectra.

NMR spectroscopy and spatial structure calculation

NMR spectra were acquired at 45 °C using 600 and 800 MHz AVANCE III spectrometers

(Bruker BioSpin, Germany) equipped with pulsed-field gradient triple-resonance cryoprobes.

1H, 13C, and 15N resonances of TrkA-TMD were assigned with CARA software (51) using

two- and three-dimensional heteronuclear experiments (52): $^1H/^{15}N$-HSQC, $^1H/^{15}N$-TROSY,

$^1H/^{13}C$-HSQC, $^1H/^{15}N$-HNHA, $^1H/^{13}C/^{15}N$-HNCA, $^1H/^{13}C/^{15}N$-HN(CO)CA, $^1H/^{13}C/^{15}N$-HNCO,

3D-HccH-TOCSY, ^{13}C- and ^{15}N-edited NOESY-HSQC (recorded on 600 and 800 MHz

spectrometers, respectively). Dimeric spatial structures were calculated with the CYANA

program (53) based on torsion angle restraints estimated from the chemical shift values

obtained with the standard protocol of the TALOS-N program (54) and with intra- and inter-

monomeric NOE distance restraints derived through analysis of the three-dimensional ^{15}N-

and [13]C-edited NOESY and [15]N,[13]C-F1-filtered/F3-edited-NOESY spectra (52) acquired for

"isotopic-heterodimer" samples. MOLMOL software was used to calculate the contact areas

between the dimer subunits and to visualize the structures (55). Hydrophobic properties of

the α-helices in the TrkA-TMD dimers were calculated using the molecular hydrophobicity

potential (MHP) approach implemented in the PREDDIMER program (56).

Isolation of Membrane fractions

TrkA mutants were overexpressed in Hela cells by transfection using PEI (1mg/ml). 48

hours after transfection cells were resuspended in 1ml ice-cold homogenization buffer (250

mM sucrose, 1mM EDTA, 10 mM tris buffer pH 7,1 plus protease inhibitors) and broken by

sonication in two time intervals (30,30 s) with 50W and frequency at 30 MHz on ice. The

broken cell homogenate was centrifugated for 10 minutes at 500g at 4ºC to remove whole

cells and nuclei. To collect the membrane fraction the cleared supernatant was centrifuged at

100,000*g* at 4ºC for 1h in a Beckman Optima MAX ultracentrifuge with a TLA110 rotor

using polycarbonate thick wall centrifuge tubes (Beckman coulter ref. 362305). The

supernatant that contains soluble proteins was removed and the pellet-containing membranes

was resuspended in 1ml ice-cold homogenization buffer by sonication and re-centrifuged at

100,000*g*. The final pellet contains the membrane fraction used to the iodine oxidation

protocol.

Iodine oxidation for crosslinking of TrkA mutants

As cysteine residues are buried deep in the phospholipid bilayer, membranes were isolated

and I2 was used as the oxidation agent following as in (Schwem & Fillingame, 2006).

Membrane fractions were diluted in the homogenization buffer (250 mM sucrose, 1mM

EDTA, 10 mM Tris buffer pH 7,1 plus protease inhibitors). Protein content was quantified

using a Bardford kit (Pierce) and equal amounts of isolated membranes fractions were

incubated for 10 min with or without NGF (10 ng/mL). A solution of 2.5 mM I2 in absolute

ethanol was freshly prepared immediately before the crosslinking of cysteine residues and

was added to the incubated membrane fractions with NGF (250 mM final Iodine

concentration) for 30 seconds at room temperature. The reaction was stop adding 1/10

volume of a freshly made solution of sodium thiosulfate (60 mM final concentration). Non-

reducing SDS-PAGE sample buffer was added and the samples were boiled for 5 minutes

before analyzed by non-reducing SDS-PAGE.

Differentiation of PC12nnr5 cells.

Transfection in PC12nnr5 cells was performed using Lipofectamine 2000 as per

manufacturer instructions. The mutant or the wild-type TrkA-transfected cells and mock-

transfected cells as well as non-transfected cells were treated under the same conditions in a

six-well tissue culture plate. The cells were washed three times with serum-free medium and

incubated for 48 h in a medium containing 1% fetal bovine serum and 50 ng/mL of NGF

(Alomone). At time 0, 24h and 48h, cells were washed with cold PBS and fixed with 4%

paraformaldehyde during 15 minutes at room temperature. Cells were imaged using a Leica

SP8 spectral confocal microscope. The percentage of cells with a neurite twice longer as the

cell body was counted as differentiated.

Accession codes

Atomic coordinates and experimental restraints were deposited in the Protein Data Bank

under the accession codes PDB ID: 2n90 for TrkA-TM-wt.

Acknowledgments

We thank Dr. Carlos Ibañez for critical reading of the manuscript, Dr. W. DeGrado for

providing ToxRed plasmids, and Dr. Yves Barde for the TrkA plasmid. This study was

supported by the Spanish Ministry of Economy and Competitiveness (MINECO; project

BFU2013-42746-P and SAF2017-84096-R) and by the Generalitat Valenciana Prometeo

Grant 2018/055 to MV). Studies of TrkA-TM homodimerization were supported by the

Russian Science Foundation (grant No# 19-74-30014 to A.S.A).

Author Contributions

M.L.F. constructed the TrkA mutants and performed the TrkA activation, differentiation,

functional and crosslinking experiments. K.D.N. designed and performed NMR

experiments and analyzed NMR data. S.A.G. developed the expression protocols and

synthesized samples for NMR analysis. K.S.M. analyzed NMR data and contributed to the

preparation of the manuscript. A.S.A. and M.V. designed and supervised the entire project

and wrote the manuscript.

 Figure Legends

Figure 1. Analysis of the structure of TrkA transmembrane domain dimers in the

ToxRED system and in DPC micelles.

A) ToxRed system analysis and quantification of the percentage of dimerization of the rat

TrkA-TMD normalized to glycophorin A (GpA) wt (100%). GpA-wt* refers to an inactive

ToxR that cannot induce the RedFP.

B) Percentage of dimerization of rat TrkA-wt and TrkA-TMD mutants normalized to

dimerization of GpA wt (100%) in the ToxRed system. (In A and B results represent the

average of at least 10 colonies analyzed per experiment. Plotted is the result of at least

seven independent experiments). Error bars represent the standard error calculated from the

seven independent measurements. Statistics were performed using two-way ANOVA

analysis and Dunnet's multiple comparison test using GraphPad software. P values of

conditions significantly different from wt are shown on top of the error bars (****

P<0.0001).

C) Schematic representation of the NMR spatial structure of the human TrkA-TMD dimer

in lipid micelles, from different angles. The surface of one TMD helix is represented with

the $S_{419}xxxG_{423}$ motif in blue and the NMR dimer interface in red. The TMD of the other

monomer within the dimer is represented as ribbons, and side chains are indicated as sticks.

The location of the NMR dimer interface LxxFAxxF on the NMR structure is colored red

and other mutated residues are colored orange. The PDB accession code is 2n90.

D) Alignment of the amino acid sequences of the transmembrane helical domain of TrkA

from different species (h, homo sapiens, r, ratus norvegicus, m, mus musculus; c Gallus

gallus; z Danio rerio; x Xenopus laevis). The residues participating in the NMR dimer

interface are highlighted in red, and residues participating in dimerization based on the

ToxRED analysis are highlighted in blue.

Figure 2. Functional identification of the two dimer interfaces in the TrkA-TMD

A) Amino acid sequence of the transmembrane domain of rat TrkA showing the location of the single cysteine residue substitutions.

B) Western blot of Hela cells transfected with the indicated TrkA-TMD constructs and analyzed using non-reducing SDS PAGE, showing the formation of covalent cysteine dimers (d) arising from monomers (m) crosslinked using molecular iodine (I_2) in the presence or absence of NGF.

C) Quantification of the data in B) derived from at least three independent experiments. Bars represent standard error of the mean. Statistics were performed using two-way ANOVA analysis and Dunnet's multiple comparison test using GraphPad software. (**** $P<0.0001$, *** $P=0.0004$).

D) Structure of the TrkA-TMD dimer showing the location of the indicated residues in red.

E) Cβ-Cβ distance between the indicated residues in the TrkA-TMD.

F) Hela cells transfected with the indicated TrkA constructs were stimulated with NGF (10 ng/mL) for 0, 5 or 15 minutes. The immunoblots show the activation (autophosphorylation) of TrkA as detected using an antibody specific for TrkA P-Tyr674/675. The levels of total TrkA are shown below each blot. Molecular weight markers are shown at left.

G) Quantification of the data in F) derived from at least three independent experiments. Bars represent standard error of the mean. Statistics were performed using two-way ANOVA analysis and Dunnet's multiple comparison test using GraphPad software. P values of conditions significantly different from wild type are shown on top of the error bars.

H) Cytometric analysis of the expression of the TrkA mutants at the plasma membrane (orange bars) and quantification of the percentage of PC12nnr5 differentiated cells (colored

bars) transfected with the indicated TrkA constructs together with GFP. EV, empty vector.

The percentage of the total GFP-positive transfected cells with a neurite twice as long as

the length of the cell body was quantified. Bars represent the standard error of at least three

independent experiments. Statistical analysis was performed using ordinary one-way

ANOVA, using Bonferroni's multicomparison test compared to wt. The P values of

significant differences are shown. ns, not significant.

Figure 3. K410 and K411 of eJTM are crosslinked with BS3 upon NGF binding.

A) Location of the Lys residues in the crystal structure of the rat TrkA-ECD/NGF complex

(PDB: 2IFG) (Wehrman et al., 2007). The Lys residues are shown as green circles are

located in the extracellular juxtamembrane region of the TrkA-ECD (in gray).

B) Western immunoblots of lysates of Hek293 cells transfected with the indicated TrkA

constructs (see Figure 3A) and incubated with or without NGF in the presence the cross

linker BS3. Molecular weights are indicated at left. Actin was assayed as a loading control.

Figure 4. A preferred dimer interface in the TrkA juxtamembrane region.

A) Amino acid sequence of the rat TrkA cysteine mutant constructs that are mutated in the

region of the eJTM closest to the TMD. B) Quantification of the ratio of dimer:monomer of

the TrkA mutants as determined using non-reducing SDS-PAGE. C) Quantification of the

activation of TrkA (with and without NGF) by quantifying the signal from the

phosphorylation of the Tyr 674/675 signal using western blotting of the cysteine mutants in

the JTM region. D) Scatter plot of the dimerization of TrkA cysteine mutants respect to its

activation in the absence of NGF. A regression fit using the active (green) and inactive

(red) cysteine mutant dimers is shown with the indicated r^2. E) PC12nnr5 cell

differentiation assay of TrkA-wt and TrkA-K410C in the presence and absence of NGF. E)

Quantification of the differentiation of PC12nnr5 cells transfected with the indicated TrkA

constructs and incubated in the absence (blue bars) or presence (green bars) of NGF. G)

Cytometric analysis of the expression of the TrkA mutants at the plasma membrane of Hela

cells. H) Model of the eJTM into an ideal α-helix showing the spacial location of the

indicted residues. Error bars represent the standard error of the mean. Statistics was done

using two-way ANOVA analysis and Dunnet's multiple comparison test using GraphPad

software. The P values of significant differences are shown (**** $P<0.0001$).

Figure 5. Rotation of the TrkA-TMD as a mechanism of TrkA activation by NGF.

A) Effect of NGF on the activation of overexpressed TrkA. Hela cells transfected with the

indicated constructs were stimulated with NGF (10 ng/mL) for 0, 5, 15 or 30 minutes. The

immunoblots show the activation of TrkA as detected using an antibody specific for the

autophosphorylated P-Tyr674/675, P-Tyr490 and P-Tyr751 residues and for the

downstream activated P-ERK. The levels of total TrkA and total ERK are shown below

each blot. Actin was blotted as a loading control. A representative blot of at least three

independent experiments is shown.

B) Amino acid sequences of the rat TrkA Leu insertion mutants indicating the location of

the inserted Leu residues.

C) Schematic drawing of the TrkA-TMD-ICD showing how the different numbers of

inserted Leu residues (blue) in the TMD induce a different rotation of the intracellular

domain (green).

D) Hela cells were transfected with the indicated TrkA-TMD insertion mutants. The

immunoblots show the activation of TrkA as detected using an antibody specific for the

TrkA autophosphorylation P-Tyr674/675 residues. The levels of total TrkA are shown

below. Actin was assayed as a loading control.

E) Quantification of the data in D) from at least three independent experiments. Bars

represent standard error of the mean. Statistical analysis was performed using one-way

ANOVA with Dunnet's multicomparison test and TrkA-wt as a control. The P value of the

significant difference is shown.

F) PC12nnr5 cell differentiation of TrkA-TMD insertion mutant-transfected cells incubated

in the absence of NGF. The percentage of the total GFP positive-transfected cells with a

neurite twice as long as the length of the cell body was quantified. Bars represent the

standard error of at least four independent experiments. Statistical analysis was performed

using one-way ANOVA with Dunnet's multicomparison test and TrkA-wt as a control. The

P values of significant differences are shown.

Figure 6. Model of TrkA activation by NGF.

TrkA exists in an equilibrium between a monomer and an inactive pre-dimer, which is

stabilized by interactions between Ig2 domains and between the inactive kinase domains.

Upon NGF binding a rotation of the Ig2 domain induces the formation of a crossed TMD

dimer that is stabilized by the protein interface determined in the NMR and functional

studies. The rotation of the TMD is transmitted to the intracellular region in order to

activate TrkA kinase autophosphorylation.

Figure 1- Table supplement 1

NMR distance & dihedral restraints	
Total unambiguous NOE restraints	360
intra-residue	136
inter-residue	228
sequential ($|i-j|=1$)	138
medium-range ($1<|i-j|\leq4$)	76
long-range ($|i-j|>4$)	0
inter-monomeric NOE	16
Hydrogen bond restraints (upper/lower)	144/144
Total torsion angle restraints	118
backbone φ	52
backbone ψ	51
side-chain χ^1	15

Structure calculation statistics	
CYANA target function (Å^2)	0.64±0.21
Average pairwise r.m.s.d. (Å)	
TM α-helical region	
backbone atoms	0.18±0.07
all heavy atoms	0.36±0.07

Ramachandran analysis [a]	
% residues in most favoured regions	90.6
% residues in additional allowed regions	9.4
% residues in generously allowed regions	0.0
% residues in disallowed regions	0.0

Helix-helix packing	
Contact surface area per dimer subunit (Å^2)	370±40
Angle θ between the helix axes (deg.)	40±5
Distance d between the helix axes (Å)	
TM region	8.8±0.5

Table note:
[a] Ramachandran statistics was determined using CYANA
[b] Residues from unfolded and flexible regions

Figure 1

A

% diemrization respect to GpA wt

GpA-wt   GpA-wt*   GpA-G83I   TrkA-TMD

**** (on GpA-wt*)
**** (on GpA-G83I)

B

% dimerization

P = 0.0209
P = 0.0002

wt   G417I   S419I   A421I   G423I   A425I   A428I   A429I

*** (on S419I)
* (on G423I)

C

S419
G423
G417
L424
A428
F427
F431
F431
40º
A421
A425
S433

D

TM helix

hTrkA	TPFGV**S**VAV**G****L**AV**FA**CL**F**LSTLLLVL
rTrkA	TPFGV**S**VAV**G****L**AVS**A**AL**F**LSALLLVL
mTrkA	TPFGV**S**VAV**G****L**AVS**A**AL**F**LSALLLVL
cTrkA	HTFGV**S**VAVA**L**AV**FA**SL**F**LSVMLIAL
xTrkA	NTFGV**S**VAV**G****L**AV**FA**CL**F**LLLMLIAL
zTrkA	RVFGV**S**VAV**G****L**AV**FA**CT**F**LLIMVVVI
hTrkC	DTFGV**S**IAV**G****L**AA**FA**CVLLVVLFVMI
hTrkB	EHLSVYAVVV**I**ASVVGFCLLVMLFLL

A

α-helix

```
wt      PFGVSVAVGLAVSAALFLSALLLVL
F416C   PCGVSVAVGLAVSAALFLSALLLVL
G417C   PFCVSVAVGLAVSAALFLSALLLVL
V418C   PFGCSVAVGLAVSAALFLSALLLVL
S419C   PFGVCVAVGLAVSAALFLSALLLVL
V420C   PFGVSCAVGLAVSAALFLSALLLVL
A421C   PFGVSVCVGLAVSAALFLSALLLVL
V422C   PFGVSVACGLAVSAALFLSALLLVL
G423C   PFGVSVAVCLAVSAALFLSALLLVL
L424C   PFGVSVAVGCAVSAALFLSALLLVL
```

Figure 3

A

K100

K174

K133

K240
K241

K197

K260
K261
K292

NGF

TrkA

absent in the
X-ray Structure

K410
K411

membrane

B

TrkA-wt   TrkA-KK/RR

NGF   -   +   -   +
BS3   +   +   +   +

250-

dimer

140-

100-

monomer

50-

actin

A

wt   SRDPVEKKDETPF
S404C CRDPVEKKDETPF
R405C SCDPVEKKDETPF
D406C SRCPVEKKDETPF
V408C SRDPCEKKDETPF
E409C SRDPVCKKDETPF
K410C SRDPVECKDETPF
K411C SRDPVEKCDETPF
D412C SRDPVEKKCETPF
E413C SRDPVEKKDCTPF
T414C SRDPVEKKDECPF

Figure 5

A

EV   TrkA   TrkA + NGF
0 5 15 30 0 5 15 30 0 5 15 30

P-Tyr 674
P-Tyr490
P-Tyr751
TrkA
P-ERK
ERK
actin

B

wt   TPFGV**S**VAVG**L**AVFACLFLSTLLLVL
ins1L TPFGVSVAVGL**L**AVFACLFLSTLLLVL
ins2L TPFGVSVAVGL**LL**AVFACLFLSTLLLVL
ins3L TPFGVSVAVGL**LLL**AVFACLFLSTLLLVL
ins4L TPFGVSVAVGL**LLLL**AVFACLFLSTLLLVL

C

wt   ins1L   ins2L   ins3L   ins4L
           100°   200°   300°   400°

D

wt ins1L ins2L ins3L ins4L

P-Tyr674/675
total TrkA
Actin

E

P=0.039

P-Tyr/total TrkA

wt InsL Ins2L Ins3L Ins4L

F

% Cells with a neurite

****

wt InsL Ins2L Ins3L Ins4L

Figure 6

Figure S1. Dimerization of the TrkA-TMD in micelles.

(A) Percentage of the monomeric (green), dimeric (blue) and oligomeric (red) forms of the human TrkA-TMD at different LPRs. Population values were calculated from the cross-peak intensities in the ^{1}H/^{15}N-TROSY spectra.

(B) Apparent free energy of dimerization of the TrkA-TMD (ΔG_{app}) = $RT\ln([M]2/[D])$ plotted as a function of $RT\ln(X)$, where $X = ([Det]-N_m[M]-N_d[D])/N_e$. [M], [D] and [Det] represent monomer, dimer and DPC concentration, respectively. N_e represents the number of DPC molecules per empty micelle, and N_m and N_d represent the number of DPC molecules per micelle for monomers and dimers, respectively.

Figure S2. The ^{1}H/^{15}N-HSQC spectrum of the human TrkA-TMD-wt.

The human TrkA-TMD was solubilized in an aqueous suspension of DPC micelles at an LPR of 50:1, and at 45 °C and pH 5.9. ^{1}H–^{15}N backbone and side-chain resonance assignments are shown.

Figure S3. Summary of the NMR data for the TrkA-TMD embedded into DPC micelles.

A) NOE connectivities observed in the ^{1}H/^{15}N-NOESY-HSQC spectrum for the TrkA-TM peptide.

B) Secondary ^{13}Cα chemical shifts of the TrkA-TMD are given by the difference between the actual chemical shift and a typical random-coil chemical shift for a specific residue. Pronounced positive or negative $\Delta\delta(^{13}C\alpha)_{h-c}$ values indicate a helical structure or an extended conformation of a protein.

Figure S4. Assessment of the quality of the NMR-derived TrkA-TMD dimer structure.

Structural alignment of the 10 lowest energy NMR structures of TrkA-TMD dimers.

Figure S5. **Hydrophobicity map of the TrkA-TMD dimer.**

The black oval delineates an helix-packing interface as calculated with the PREDIMER software. The residues within this black oval comprise the TrkA-TMD dimerization interface. The panels to the right show the percentage of inter-monomer contact surface area.

Figure S6. **TrkA-concentration-dependence of ligand-independent activation of TrkA**. Increasing amounts of TrkA were transfected into Hela cells using PEI. After 48 h, the cells were stimulated or not with NGF (10 ng/mL). Cell lysates were analyzed by SDS-PAGE immnoblotting using antibodies specific for TrkA phospho-Tyr674/675 and total TrkA.

Figure S7. Membrane localization of TrkA mutants in Hela cells. Confocal fluorescence images of Hela cells transiently transfected with the indicated TrkA constructs. The cells were fixed and analyzed using an antibody against HA located in the N-terminus of TrkA, which was detected with a secondary antibody conjugated with the fluorophore alexa-555 (Red). Blue, nuclear DAPI staining. The cells shown are representative of 10 transfected cells analyzed for each condition**.**

Figure S8. Membrane localization of TrkA mutants in Hela cells by cytometric analysis. Percentage of TrkA constructs expressed at the plasma membrane relative to TrkA-wt (100%) obtained in Hela cells transfected with the indicated constructs and

analyzed by flow cytometry using an HA primary antibody and a secondary antibody

labeled with a fluorephore for alexa-488. EV is empty vector.

Figure S1

A

B

Figure S3

A

B

Figure S4

Figure S5

Figure S6

Phospho-Tyr 674/675

Total TrkA

Figure S7

Figure S8

EV

TrkA wt

InsL

Ins2L

Ins3L

Ins4L