1 **Myelin-reactive B cells exacerbate the severity of CD4[+] T cell-driven CNS**

2 **autoimmunity in an IL-23-dependent manner.**

Mohamed Reda Fazazi[1], Prenitha Mercy Ignatius Arokia Doss[1], Resel Pereira[2],

Neva Fudge[3,4], Aryan Regmi[2,5], Charles Joly-Beauparlant[1], Irshad Akbar[1], Asmita

Pradeep Yeola[1], Benoit Mailhot[1], Joanie Baillargeon[1], Philippe Grenier[6], Nicolas

Bertrand[6,7], Steve Lacroix[1,8], Arnaud Droit[6, 8], Craig S. Moore[3,4], Olga L. Rojas[2,5],

Manu Rangachari[1,8,9].

[1] axe Neurosciences, Centre de recherche du CHU de Québec – Université

Laval, Pavillon CHUL, 2705 boul Laurier, Quebec City, QC, CANADA G1V 4G2; [2]

Krembil Research Institute, University Health Network, Toronto ON CANADA

M5T 0S8; [3] Division of BioMedical Sciences and [4] Department of Neurology,

Faculty of Medicine, Memorial University of Newfoundland, St. John's NL,

CANADA, A1B 3V6; [5] Department of Immunology, University of Toronto, Toronto

ON, CANADA, M5S 1A1; [6] axe Endocrinologie et nephrologie, Centre de

recherche du CHU de Québec – Université Laval, Pavillon CHUL, 2705 boul

Laurier, Quebec City, QC, CANADA G1V 4G2; [7] Faculty of Pharmacy, Laval

University, 1050 ave de la Médecine, Quebec City, CANADA; [8] Department of

Molecular Medicine, Faculty of Medicine, Laval University, 1050 ave de la

Médecine, Quebec City, CANADA; [9] Corresponding author.

Abstract

Multiple sclerosis (MS) is an autoimmune disorder of the central nervous system

(CNS) that has traditionally been considered a T cell-mediated disease.

However, accumulating evidence points to a crucial role for B cells in disease

processes. IgH[MOG] mice possess a transgenic immunoglobulin heavy chain

derived from a monoclonal antibody specific to myelin oligodendrocyte

glycoprotein (MOG), a key target for autoimmune responses. Here, using the

experimental autoimmune encephalomyelitis (EAE) model of MS, we investigated

the susceptibility of IgH[MOG] mice to CD4$^+$ T cell-driven disease that was induced

by active immunization with MOG$_{[35-55]}$ autoantigen. We found that immunized

IgH[MOG] mice rapidly developed severe EAE, characterized by the aggregation of

T and B cells in the CNS meninges. We observed an increased presence of

class-switched and inflammatory cytokine-positive B cells in the IgH[MOG] CNS, as

well as a greater frequency of IL-17- and GM-CSF-producing CD4$^+$ T cells.

Production of the Th17 maintenance factor IL-23 was increased from IgH[MOG]

CNS-infiltrating B cells, and *in vivo* blockade of IL-23p19 strongly attenuated

disease severity in IgH[MOG] mice. Strikingly, we observed an increased frequency

of PD-1$^+$CXCR5$^-$ T peripheral helper (Tph)-like cells in the CNS parenchyma and

dura mater of IgH[MOG] mice. Both Tph accumulation in the CNS, as well as

meningeal inflammation, were again sharply reduced upon IL-23p19 blockade *in*

vivo. Notably, CNS-infiltrating B and Tph cells from IgH[MOG] mice showed an

upregulation of genes related to neurodegeneration and oxidative

phosphorylation, and IL-23 blockade reduced ROS production from these cells *in*

vivo. Altogether, these data show that MOG-specific B cells contribute to severe

CD4$^+$ T cell-driven EAE by promoting CNS accumulation of Th17 and Tph cells,

as well as tertiary lymphoid organs in the CNS meninges, in an IL-23 dependent

manner.

Keywords: experimental autoimmune encephalomyelitis, multiple sclerosis, B

cell, CD4$^+$ T cell, myelin oligodendrocyte glycoprotein (MOG), IgH[MOG], 1C6,

non-obese diabetic (NOD), Th17, IL-23, T-peripheral helper cells (Tph cells),

tertiary lymphoid organ.

Introduction

Multiple sclerosis (MS) is a chronic autoimmune disease in which the adaptive immune system launches an attack against central nervous system (CNS) proteins, such as myelin. MS affects more than 2 million people worldwide (1). Approximately 80% of patients present an initially relapsing-remitting (RR) disease course for which there are now more than 10 disease-modifying therapies available. However, 30-60% of these RR patients will eventually transition to a chronically worsening secondary progressive (SP) phase, for which treatment options are limited (2). Pathophysiological mechanisms in progressive MS are thus of intense current interest (3).

T cells, and CD4$^+$ T cells in particular, have historically been the most intensively studied players in the immune pathogenesis of MS. However, it has become increasingly clear that B cells additionally play important roles in MS. Clonally expanded B cells are present in the cerebrospinal fluid (CSF) and MS plaques (4-6), and the presence of meningeal follicles adjacent to cortical lesions is associated with disease progression (7, 8). Further, antibodies against myelin oligodendrocyte glycoprotein (MOG), a key component of myelin, were found in active MS lesions (9). Crucially, the B cell-targeting anti-CD20 drugs rituximab(10), ocrelizumab (11) and ofatumumab (12) improve RRMS, and ocrelizumab is the only FDA approved drug for primary progressive MS (13). Curiously, though, antibody-secreting plasmablasts do not express CD20 (14). This suggests that the pathogenic role of B cells in MS might lie less with the

generation of autoantibodies, and more with their capacity to interact with other

immune cell types, such as T cells.

Experimental autoimmune encephalomyelitis (EAE) is an animal disease

that recapitulates many of the immune aspects of MS pathogenesis. Use of this

model has helped us to understand the role of T cells, and CD4$^+$ T cells in

particular, in the initiation and maintenance of autoreactive inflammation in the

CNS (15). However, studies using the transgenic IgH$^{[MOG]}$ mouse strain have

indicated that B cells may also play a crucial role in EAE pathology (16-18).

These mice (also known as "Th") express a knocked-in IgH chain derived from a

MOG-specific antibody; thus, around 30% of their B cells are therefore specific

for MOG protein (16). IgH$^{[MOG]}$ animals develop severe EAE when immunized

with either whole MOG protein (16) or with its extracellular domain (MOG$_{[1-125]}$)

(19), indicating an important role for MOG-reactive B cells in neuroimmune

processes. Further, they develop spontaneous EAE when crossed to myelin

peptide-specific T cell receptor transgenic lines on the B6 (17) and SJL/J (18)

backgrounds. Indeed, while "T cell help" is required for the full activation of B

cells in the majority of cases, there is increasing evidence that B cells can

reciprocally promote effector T cell responses, notably those of the Th17 lineage

(20-24). However, the potential mechanisms by which MOG-reactive B cells

facilitate T cell-driven pathogenicity, such as in class II-restricted peptide

immunization models of EAE, remain incompletely understood.

Here, we studied the co-operative role of B cells and T cells in CNS

autoimmunity using IgH$^{[MOG]}$ mice on the non-obese diabetic (NOD) genetic

background on which EAE develops with a relapsing/chronic disease pattern

(25).When immunized with the T cell-obligate peptide MOG[35-55], IgH[MOG] mice

developed a rapid, severe form of EAE characterized by meningeal clusters of

CD4$^+$ T cells, B cells and CD11c$^+$ dendritic cells (DCs) with alterations to the

underlying extracellular matrix (ECM) reminiscent of tertiary lymphoid organs

(TLOs). Th17 responses were specifically upregulated in the inflamed IgH[MOG]

CNS and, surprisingly, CNS-infiltrating B cells produced the Th17 maintenance

factor IL-23. Further, PD1$^+$CXCR5$^-$ "T peripheral helper-like cells", a recently

described effector T cell subset that associates with B cells in the inflamed

synovium in rheumatoid arthritis (RA) (26), were presented at a high frequency in

the IgH[MOG] CNS parenchyma and dura mater. Transcriptomics analysis of CNS-

infiltrating B and Tph cells revealed that genes related to neurodegeneration and

oxidative phosphorylation were upregulated in both cell types in IgH[MOG] mice,

suggesting that these processes contribute to the heightened pathology observed

in these animals. Notably, IL-23 blockade abrogated severe disease in IgH[MOG]

mice, as well as both CNS Tph accumulation and ROS production from both B

cells and Tph cells. Together, our results demonstrate that MOG-specific B cells

play a crucial role in augmenting CD4$^+$ T cell-driven EAE in an IL-23-dependent

manner.

Results

IgH[MOG] mice develop severe EAE upon active immunization with MOG[35-55]

When immunized with MOG$_{[35-55]}$, wildtype WT NOD mice display a disease

course characterized by relapse remitting disease in the early phase that

transitions to a chronic worsening phase in some animals (25). NOD-EAE has

thus been considered a possible model of SPMS (27, 28), though others argue

against this interpretation (29). We confirmed that MOG$_{[35-55]}$-immunized WT

NOD mice develop disease characterized by relatively mild symptoms over the

first ~100 days (Supplemental Figure 1A). We next compared the development of

EAE between NOD-background WT and IgH$^{[MOG]}$ mice upon MOG$_{[35-55]}$

immunization. Both male and female IgH$^{[MOG]}$ mice developed extremely severe

disease within 25 days (Figure 1A), with a substantial frequency of these mice

attaining ethical endpoints (10/14 IgH$^{[MOG]}$ males vs. 0/14 WT males, p=0.0002;

12/16 IgH$^{[MOG]}$ females vs. 0/16 WT females, p<0.0001). Histopathological

analyses revealed increased lymphocyte infiltration and demyelination in the

spinal cords of IgH$^{[MOG]}$ mice relative to controls (Figure 1B).

We next examined peripheral T cell responses prior to disease onset. No

differences in IL-17 or IFNγ were detected by either *ex vivo* flow cytometry

(Figure 1C) or by ELISA after MOG$_{[35-55]}$ peptide recall (Supplemental Figure 1B).

Next, to identify a possible role for IgH$^{[MOG]}$ B cells in directly driving T cell

responses, we pulsed IgH$^{[MOG]}$ or WT B cells with MOG$_{[35-55]}$ and co-cocultured

them with MOG$_{[35-55]}$-specific 1C6 transgenic T cells (30-32). However, no

differences were observed in T cell proliferation (Figure 1D), indicating that

IgH$^{[MOG]}$ B cells were not intrinsically better at presenting antigen to T cells.

Furthermore, while the cervical LNs are a potential site of T cell reactivation in

MS (33), we observed only limited production of IL-17 and IFNγ in this

compartment. While a large proportion of CD4[+] T cells in cervical LNs were

positive for TNFα, no differences were detected between WT and IgH[MOG]

(Supplemental Figure 1C).

Antigen-specific antibody (Ab) secretion is the primary function of B cells.

Further, oligoclonal immunoglobulin (Ig) banding in cerebrospinal fluid (CSF) is

an important diagnostic marker for MS (34). However, no differences in MOG-

specific circulating IgG were found in IgH[MOG] serum relative to controls (Figure

1E), suggesting that severe disease in transgenic animals was not accompanied

by an increase in MOG-specific autoantibodies. Together, these data showed

that the presence of myelin-reactive B cells can exacerbate CNS autoimmunity

and tissue damage when EAE is induced in a CD4[+] T cell dependent manner.

However, increased disease severity did not appear to be accompanied by

markers of peripheral T and B cell activation.

Accumulation of meningeal TLOs in IgH[MOG] mice

Target organ-infiltrating immune cells are essential to the pathogenesis of

autoimmune disease. We therefore enumerated the frequency and absolute

number of CNS-infiltrating immune cells – CD19[+] B cells, CD4[+] T cells, CD8[+] T

cells, CD11c[+] dendritic cells (DCs), CD11b[+]CD11c[-] macrophages and Ly6G[+]

neutrophils – in immunized WT and IgH[MOG] mice. The relative frequencies

(Figure 2A) and absolute numbers (Supplemental Figure 2) of CD4[+] T cells,

CD8[+] T cells, DCs, macrophages and neutrophils were significantly increased in

the CNS of IgH[MOG] mice relative to WT; however, B cell frequency and absolute

numbers were unchanged.

Meningeal TLOs have been documented in progressive MS and their presence

correlates with poor outcomes (7, 35). Their presence in EAE is dependent on

murine genetic background and the manner of disease induction; while TLOs

arise upon active immunization of SJL/J mice with PLP$_{[139-151]}$ (36), they do not

appear in MOG$_{[35-55]}$-immunized C57BL/6J mice, rather arising only upon

adoptive transfer of Th17 cells (37). Whether or not they are a feature of NOD-

EAE is as-yet unknown. As B cells are crucial to TLO formation (37), we

assessed their presence in WT versus IgH[MOG] mice. Strikingly, we observed

clustering of CD4[+] T cells, B cells and CD11c[+] DCs of varying magnitude in the

CNS of IgH[MOG] relative to WT mice (Figure 2B).

TLOs are defined as tissue immune cell aggregates organized within a

stroma cell network in inflamed non-lymphoid tissues(38). We found an increase

of elaborate stroma network near the meninges in IgH[MOG] mice as demonstrated

by the organized expansion of fibronectin, PDGFR$\alpha\beta$a/b IF staining underneath

the B cell clustering (Figure 2C). Together, these data demonstrate that TLOs

accumulate in the meninges of MOG$_{[35-55]}$-immunized IgH[MOG] mice.

The IgH[MOG] CNS is characterized by inflammatory B cells and Th17 cells

As we did not see an increase in frequency or absolute number of B cells

in the CNS of immunized IgH[MOG] mice (Figure 2A), we next assessed whether

there might be differences in the proportion of functional B cell subsets between

the groups. Indeed, we observed a stark increase in the frequency of class-

switched (CS) IgM$^-$IgD$^-$ B cells in the CNS (Figure 3A) of IgH[MOG] mice, which

was accompanied by a decrease in the frequency of immature and mature B

cells. In line with our finding that serum MOG-specific IgG was comparable

between WT and IgH[MOG], we observed no differences in B cell subpopulation

frequency in the spleen (Supplemental Figure 3A). Further, we observed no

difference in MHC class II expression between CNS-infiltrating WT and IgH[MOG]

B cells (Supplemental Fig 3B). On the other hand, an increase in GL-7$^+$Fas$^+$

germinal center (GC) B cells was seen in the IgH[MOG] CNS (Supplemental Figure

3C).

B cells can generate canonical inflammatory cytokines such as IL-6 (ref

(39)), TNFα (40) and GM-CSF (41), with their production of such cytokines being

linked to worsened outcomes in MS. We observed that a significantly greater

proportion of IgH[MOG] CNS B cells produced these cytokines as compared to WT

counterparts, with the differences in IL-6 and TNFα being most apparent (Fig

3B).

Next, as we had observed an elevated frequency of CD4$^+$ T cells in the

CNS of sick IgH[MOG] mice, we next examined the capacity of these cells to

produce inflammatory Th1 and Th17 cytokines by flow cytometry, due to the well-

established role of these CD4$^+$ effector T cell subsets in EAE (42). We observed

no differences between IgH[MOG] and WT CD4+ T cells in their production of IFNγ

and TNFα in the CNS. By contrast we saw a strong upregulation of IL-17

production from CNS-infiltrating CD4+ T cells from IgH[MOG] at disease endpoint

(Figure 3C). Notably, production of GM-CSF, a key pathogenic cytokine

implicated in Th17-driven tissue inflammation, was also augmented in IgH[MOG]

CD4+ T cells.

B cells have been reported as promoting Th17 differentiation (21); further,

both the congenital absence (22) and therapeutic depletion (23, 24) of B cells

correlates with diminished circulating Th17 cell frequency. Thus, to test whether

IgH[MOG] B cells facilitate Th17 responses *in vivo*, we adapted our previously

described Th17 adoptive transfer protocol (32) by seeding NOD.*Scid* mice with

immunologically naïve WT or IgH[MOG] B cells, and subsequently injecting $MOG_{[35\text{-}55]}$-specific 1C6 transgenic (30) Th17 cells. Mice that had been infused with

IgH[MOG] B cells developed substantially more severe Th17-driven EAE than

those reconstituted with WT B cells (Figure 3D). This was characterized by an

increased frequency of IL-17+ and GM-CSF+ CNS-infiltrating Th17 cells in

IgH[MOG] B cell-reconstituted hosts (Figure 3E, Supplemental Figure 3D). Taken

together, these data show that severe EAE in IgH[MOG] mice is characterized by

the presences of inflammatory B cells and Th17 cells, and that IgH[MOG] B cells

can promote Th17-mediated disease processes.

Augmented IL-23 production by IgH[MOG] mice leads to exacerbated disease

These findings led us to investigate whether IgH[MOG] B cells may be

characterized by increased production of the Th17 stabilization factor IL-23,

which is essential to Th17 pathogenicity *in vivo* (43, 44) and which can be

secreted by human B cells (45). The frequency of B cells co-expressing both

chains of IL-23 (p40$^+$p19$^+$) was strongly upregulated in the IgH[MOG] CNS, despite

showing no differences in co-expression of the chains of the related cytokine, and

Th1 differentiation factor, IL-12 (p40$^+$p35$^+$) (Supplemental Figure 4A). Further, no

differences in the co-expression of p19 and p40 were observed between WT and

IgH[MOG] DCs, astrocytes (46) or microglia (Supplemental Figure 4B). Use of an

antibody that detected IL-23 heterodimer confirmed that CNS IgH[MOG] B cells

show higher expression of this cytokine (Figure 4A). As we had observed an

increase in B cell-rich aggregates in the IgH[MOG] meninges, we assessed if these

B cells were also a source of IL-23. Indeed, immunofluorescence staining

showed co-localization of IL-23 with B cells in the IgH[MOG] meninges (Figure 4B).

The robust upregulation of IL-23 by B cells in the IgH[MOG] CNS led us to

ask whether it might be essential for the exacerbated disease seen in these

animals. We therefore actively immunized IgH[MOG] mice and administered either

anti-IL-23p19 blocking antibody (47) or isotype control. *In vivo* blockade

abrogated severe EAE (Figure 4C), characterized by reduced lymphocyte

infiltration and demyelination in the spinal cord (Supplemental Figure 4D). As

Th17 responses are crucial to TLO formation (37), we next examined the effects

of IL-23 blockade on the presence of TLOs in the meninges. Administration of

anti-IL23p19 diminished the number of lymphocyte clusters in the cerebellar

meninges of IgH[MOG] mice (Figure 4D). Concomitantly, we observed a decrease

in the intricate ECM network underneath, suggesting a decrease in meningeal

TLOs in IgH[MOG] mice after administration of anti-IL23p19 (Supplemental Figure

4C).

Turning to the impact of IL-23 in cellular inflammation in the IgH[MOG] CNS,

we found that the frequency of class-switched B cells was reduced in immunized

animals treated with anti-p19 (Figure 4E). Further, CNS-infiltrating CD4$^+$ T cells

from anti-p19-treated IgH[MOG] mice showed decreased IL-17 and GM-CSF

expression (Figure 4F); however, no differences in IFNγ production between T

cells from anti-19 or isotype-treated mice were observed.

The cerebrospinal fluid is a useful proxy for pathologic processes within

the CNS parenchyma (48) and is routinely sampled in MS diagnosis. We

therefore asked whether expression of IL-23p19 transcript in the CSF of people

with MS (pwMS) were altered in relation to the presence of B cells. We examined

the CSF of 11 pwMS for expression of *Il23a* (IL-23p19 gene name) and

correlated this to the frequency of CD19$^+$ B cells in the CSF of the same

individuals. A robust correlation was identified (Spearman r = 0.7107) between

IL23a and the proportion of CD45$^+$ leukocytes that were CD19$^+$ (Fig. 4G). In sum,

these data show that excessive production of IL-23 by IgH[MOG] B cells may

underpin the severe pathology observed in these animal and that IL-23 transcript

correlates with the presence of B cells in CNS-associated tissues from pwMS.

T peripheral helper (Tph)-like cells accumulate in the IgH[MOG] CNS

T follicular helper (Tfh) cells play a critical role in germinal center formation and B cell immune activation, and were recently shown to collaborate with B cells in driving Th17-dependent EAE (49). We therefore examined the presence of PD-1^+CXCR5$^+$ Tfh cells in splenic and CNS tissues from WT and IgH$^{[MOG]}$ mice. Intriguingly, while Tfh were relatively rare in all cases, we instead observed an increase in PD-1$^+$CXCR5$^-$ CD4$^+$ T cells in the IgH$^{[MOG]}$ CNS (Figure 5A). These cells were reminiscent of the recently discovered "T peripheral helper" (Tph) cell subset (26, 50). In contrast to the CNS, we observed no differences in the frequency of PD-1$^+$CXCR5$^-$ CD4$^+$ T cells in the spleen (Supplemental Figure 5A).

Originally discovered in the context of rheumatoid arthritis (RA), Tph are recruited to the inflamed synovium, where they associate with B cells. They are distinct from Tfh (26), and indeed we found that PD-1$^+$CXCR5$^-$ cells from both WT and IgH$^{[MOG]}$ mice were negative for Bcl6 (Supplemental Figure 5B), the Tfh master transcription factor (51). While present in human RA (26, 50) and lupus (52), evidence for the accumulation of Tph in murine autoimmune disease is, thus far, limited. Interestingly the frequency of PD-1$^+$CXCR5$^-$ CD4$^+$ T cells was higher at the peak of EAE in IgH$^{[MOG]}$ mice as opposed to at onset or recovery (Figure 5B).

In addition to being PD-1$^+$CXCR5$^-$Bcl6$^-$, Tph are characterized as highly positive for ICOS, CXCL13 and IL-21 (26). We therefore compared expression of these markers between CNS-infiltrating PD-1^{neg} and PD-1$^+$ CD4$^+$ T cells of both WT and IgH$^{[MOG]}$ mice. Expression of all three markers was upregulated in the

PD-1$^+$ subpopulation in both WT and IgH$^{[MOG]}$ (Figure 5C), showing that CNS-

infiltrating PD1$^+$CXCR5$^-$ cells express known Tph markers in the context of NOD-

EAE. Interestingly, expression of IL-21 was significantly higher in IgH$^{[MOG]}$ Tph-

like cells relative to WT (Supplemental Figure 5C).

As we had found that the frequency of CNS-infiltrating IL-17$^+$ CD4$^+$ T cells

is increased in immunized IgH$^{[MOG]}$ mice (Figure 3C), we next asked whether

production of IL-17 differed between the WT and IgH$^{[MOG]}$ Tph populations.

Indeed, CNS-infiltrating IgH$^{[MOG]}$ CD4$^+$PD-1$^+$ T cells showed strikingly higher

positivity for IL-17 than their WT counterparts (Figure 5D). By contrast, we saw

no differences in IL-17 between WT and IgH$^{[MOG]}$ CD4$^+$PD-1$^-$ T cells, nor were

strain-specific differences in IFNγ seen in either the CD4$^+$PD-1$^+$ or CD4$^+$PD-1$^+$

subpopulations. Thus, Tph cells are the primary source of IL-17 from the CD4+ T

cell compartment in the IgH[MOG] CNS.

Altogether, our data indicated that CD4$^+$PD-1$^+$CXCR5$^-$ Tph cells not only

accumulate in the CNS of IgH$^{[MOG]}$ mice, but also show important phenotypic

differences with WT Tph cells. To define these alterations globally, we performed

mRNAseq analysis to compare the transcriptome of WT and IgH$^{[MOG]}$ Tph cells

taken from the CNS of immunized mice. We uncovered 2565 differentially

expressed genes ($p_{adj} < 0.05$ & $L_2FC > |1.5|$) of which 2154 were upregulated and

411 downregulated in IgH$^{[MOG]}$ versus WT (Supplemental Figure 5D). In

analyzing known Tph marker expression, we found that *Il21* was upregulated in

IgH$^{[MOG]}$ Tph cells relative to WT (Supplemental Figure 5E), in line with what was

observed at the protein level (Supplemental Figure 5C). Further, *Cx3cr1* and

Cxcl13 were also augmented in IgH[MOG] Tph cells. By contrast, *Ccr2* and *Icos*

were downregulated in IgH[MOG] Tph cells as compared to WT (Supplemental

Figure 5E). Notably, and again mirroring our observations at the protein level

(Figure 5D), expression of *Il17a*, as well as of *Il17f*, were increased in IgH[MOG]

Tph cells (Supplemental Figure 5F). As Tph cells were described to interact with

B cells in inflammatory tissue, and having seen an increase of B cell aggregates

in the meningeal TLOs, we then assessed by flow cytometry the presence of Tph

cells in the dura mater. A robust upregulation of Tph cells was observed in the

dura mater of IgH[MOG] mice compared to WT mice (Figure 5E).

Finally, we examined whether *in vivo* depletion of IL-23 could impact Tph

frequency. Indeed, the frequency of Tph was reduced in the CNS, but not spleen,

of immunized IgH[MOG] treated with anti-p19 (Figure 5F). Altogether, our data

show that CD4$^+$PD1$^+$CXCR5$^-$ Tph cells accumulate in the CNS of immunized

IgH[MOG] mice in an IL-23-dependent manner, where they are an important source

of IL-17. Further, in addition to differences in their frequency, qualitative

differences also exist between IgH[MOG] and WT CNS-infiltrating Tph cells.

Reactive oxygen species (ROS) generation is enhanced from IgH[MOG] B cells

and Tph cells

We next sought to identify molecular features, characteristic of IgH[MOG] B cells,

that might underpin the exacerbated disease seen in these mice. Pathway

analysis of the IgH[MOG] versus WT CNS-infiltrating B cell transcriptome revealed,

unsurprisingly, a number of terms linked to leukocyte function and immunity.

Strikingly, however, we also observed an over-representation of terms linked to

neurodegeneration and oxidative stress (Figure 6A). Notably, similar terms were

found to be upregulated when we compared the transcriptome of IgH[MOG] CNS

Tph cells to WT (Figure 6A); indeed, neurodegeneration and oxidative stress

gene sets were enriched in both IgH[MOG] B cells and Tph cells in the CNS (Figure

6B).

These findings suggested to us that an oxidative stress program might

underly the EAE phenotype of IgH[MOG] mice. We therefore examined whether

CNS-infiltrating B cells and Tph cells could produce reactive oxygen species

(ROS). Notably, IgH[MOG] B cells (Figure 6C) and Tph cells (Figure 6D) had higher

ROS production than their WT counterparts, as assessed CM-H2DCFDA, a

fluorescent oxidative stress indicator. Then, we examined whether *in vivo*

depletion of IL-23 could impact ROS production from IgH[MOG] B and Tph cells.

Interestingly, IL-23 blockade was decreased CM-H2DCFDA positivity from CNS-

infiltrating B cells (Figure 6E) and Tph cells (Figure 6F). In sum, these data show

that CNS-infiltrating IgH[MOG] B cells and Tph cells mount oxidative stress

responses that are IL-23 dependent.

Discussion

The role of CD4$^+$ T cells in the autoimmune pathogenesis of MS is well-

established: genome-wide association studies (GWAS) have revealed that

polymorphisms in the human leukocyte antigen class II region are strongly

associated with MS susceptibility (53), and many of the current treatments of MS

are believed to target T cell responses (54-56). However, the success of the B

cell-depleting drug ocrelizumab in both RRMS and PPMS has intensified recent

interest in the contribution of B cells to disease processes (57).

IgH$^{[MOG]}$ mice were initially described on the C57BL/6 (B6) and SJL/J

genetic backgrounds. On the B6 background, IgH$^{[MOG]}$ mice showed an increased

incidence of EAE, relative to WT, when immunized with whole MOG protein(16).

Notably, IgH$^{[MOG]}$ SJL/J mice developed EAE of greater severity than controls

when immunized with the myelin-derived epitope proteolipid protein (PLP)$_{[139-}$

$_{154]}$(16), which induces a relapsing/remitting disease pattern in SJL/J background

mice(58). This suggested that the presence of MOG-reactive B cells could

contribute to EAE pathology that was driven by a class II-restricted peptide. It

was later shown that when IgH$^{[MOG]}$ mice were crossed to the 2D2 MOG$_{[35-55]}$

TcR-transgenic strain on the C57BL/6 background, the resulting double-

transgenics spontaneously developed Devic's-like disease while single

transgenic mice did not (17). Similarly, IgH$^{[MOG]}$ x TCR^{1640} (ref (18)) and IgH$^{[MOG]}$

x 1C6 (ref (30)) double-transgenic mice, on the SJL/J and NOD backgrounds

respectively, develop spontaneous EAE at a high rate of incidence(30). These

findings indicated that the collaboration of both myelin-specific B and T cells in

the same animal could induce CNS autoimmunity; however, it was difficult to

determine whether B or T cell-driven responses were initially responsible for

disease induction in this model, and thus the nature of a putative collaboration

between B cells and T cells in disease processes remained incompletely defined.

In our study, we actively immunized IgH[MOG] mice on NOD background

with the class II-restricted peptide MOG$_{[35-55]}$. This permitted us to study the

contribution of MOG-reactive B cells in a model of EAE that is initiated by CD4$^+$ T

cells. While WT NOD mice developed a gradual, chronic MOG$_{[35-55]}$-driven

disease course, with advanced symptoms appearing as late as >100 days post

immunization, we found that immunized IgH[MOG] NOD mice develop severe

disease within a matter of weeks. Thus, by using a myelin-derived, class II-

restricted, immunogen, we show that B cells augment EAE even when CD4$^+$ T

cells initiate disease. Interestingly, no differences in circulating anti-MOG IgG

were identified between immunized WT and IgH[MOG], suggesting that worsened

disease in IgH[MOG] mice is not the result of increased production of

encephalitogenic autoantibody.

The role of CD4$^+$ T cells in licensing peripheral B cell responses is well-

established. Th17 cells, in particular, can foster B cell responses by inducing

class switching to inflammatory IgG1 and IgG2a (59) and by recruiting B cells to

TLO structures (60). Here, we provide evidence that the converse is also true,

and that B cells may themselves shape Th17 responses. CNS-infiltrating CD4$^+$ T

cells in IgH[MOG] mice showed significantly higher expression of IL-17 and GM-

CSF, but not of IFNγ, indicating that Th17 responses are elevated in these mice.

Further, CNS-infiltrating IgH[MOG] B cells showed increased expression of IL-23,

an innate immune-associated cytokine that stabilizes the Th17 lineage and is

required for the pathogenicity of Th17 cells (43, 44), and *in vivo* blockade of IL-23

sharply reduced EAE in IgH[MOG] mice.

Overall, our findings show that B cells can play an important supportive

role in facilitating pathogenic Th17 function and are consistent with a model by

which bidirectional B:Th17 interactions are essential in CNS autoimmunity.

Indeed, there is emerging evidence that B cells can promote Th17 responses.

Both peritoneal $CD5^+B220^{lo}$ (ref (20)) and LPS-stimulated splenic

$CD80^+CD86^+CD44^+$ (ref (21)) B cells can elicit Th17 differentiation. Further,

production of IL-17 is defective from T cells taken from agammaglobulinemia

patients that lack B cells(22) and therapeutic depletion of B cells with rituximab

can reduce Th17 function in the context of MS (23) as well as in rheumatoid

arthritis (24). The mechanisms underpinning bidirectional B:T cell interactions

remain to be fully elucidated.

Human B cells can secrete IL-23, and repression of this capacity is posited

as a mechanism of action for interferon-beta in MS (45). It was previously

observed that peripheral B cells from naïve IgH[MOG] mice possess the capacity to

secrete IL-23 (61); both upon pulsing with rMOG, and most strikingly upon co-

culture with $MOG_{[35-55]}$-specific 2D2 T cells in the presence of either rMOG or

$MOG_{[35-55]}$. B cell production of IL-23 in response to LPS is repressed by the

inhibitory receptor Tim-1 (ref (62)). We now show that immunization with $MOG_{[35-}$

$_{55]}$ elicits IL-23 expression from CNS-infiltrating B cells *in vivo*, and that IL-23 is

essential to both EAE and TLO formation from these animals.

Using mass cytometry, Brenner and colleagues found that $PD1^+CXCR5^-$

$CD4^+$ T cells accumulate in the inflamed synovium of RA patients. These Tph

cells secrete B cell attractant factors such as CXCL13 and IL-21 and associate

with B cells in the inflamed synovium, both in defined lymphoid aggregates as

well as more diffusely. However, in contrast to Tfh cells, Tph are not defined by

expression of the transcription factor Bcl6 (26). While these cells are also

observed in patients with lupus (52) and ANCA-associated vasculitis (63), their

potential roles in murine models of autoimmunity remain obscure. We have found

an enrichment of $PD1^+CXCR5^-$ $CD4^+$ T cells in the CNS, but not immune

periphery, of immunized $IgH^{[MOG]}$ mice. These cells are $Bcl6^{neg}$ and show

increased expression of the Tph markers ICOS, CXCL13 and IL-21, relative to

$PD-1^-$ counterparts. Intriguingly, we find that the presence of $PD-1^+CXCR5^-$ cells

is highest in mice of peak EAE severity; Tph cells were previously shown to

correlate with the severity of rheumatoid arthritis (26) and lupus (52). Here, we

also show that Tph cells can be found in the dura mater of immunized mice and

their frequency was higher in $IgH^{[MOG]}$ mice. Further, the accumulation of Tph

cells in the CNS is dependent on IL-23, revealing a heretofore unknown function

of this cytokine.

Oxidative stress has been described as an important factor for chronic

inflammation and different studies have identified a role for oxidative stress in MS

(64, 65). We now show that Tph cells as well as B cells have a general oxidative

stress programming that is IL-23 dependent.

In conclusion, we provide evidence that in the presence of a B cell

repertoire that is skewed towards MOG, NOD-background mice develop

unusually rapid and severe CD4$^+$ T cell-mediated EAE. Disease is dependent on

IL-23, which is produced by B cells and is required for the augmented Th17

responses seen in these mice. Furthermore, these mice show accumulation of

TLOs in their meninges that are again dependent on the presence of IL-23.

Finally, Tph cells aggregate in the CNS and the dura mater of these mice.

Together, our findings support a critical role for myelin-reactive B cells in

bolstering T cell-driven CNS autoimmunity in an IL-23-dependent manner.

Methods

Ethics

All mouse experiments and breedings were approved by the Animal Protection

Committee of Université Laval (protocols 2021-820 and 2021-830, to M.R).

Protocols and experiments involving human participants were approved by the

Newfoundland Health Research Ethics Board (to C.S.M).

Animals

IgH[MOG] mice on the NOD background (30), and 1C6 mice (30-32), were

obtained from Dr. Vijay Kuchroo (Brigham & Women's Hospital, Boston), and

were maintained at the animal facility of the Centre de recherche du CHU de

Québec-Université Laval. NOD/ShiLtJ and NOD.*Scid* mice were purchased from

Jackson Laboratories.

EAE induction and scoring

WT NOD and IgH[MOG] mice were immunized subcutaneously with 200µg MOG$_{[35-55]}$ (Feldan), emulsified in incomplete Freund's adjuvant (BD Difco) that was supplemented with 500 µg *M. tuberculosis* extract (BD Difco). On day 0 and day 2 post-immunization, mice received 200 ng pertussis toxin (List Biological Laboratories) intraperitoneally. Mice were monitored daily for signs of EAE, which were assessed using a semi-quantitative 0-5 scale: 0; no disease, 0.5; ruffled fur, 1; limp tail, 1.5; mild impairment in gait, 2; severe impairment in gait, 2.5; partial hind limb paralysis, 3; hind limb paralysis, 4; forelimb paralysis, 5; moribund (66). Pre-onset analyses were conducted a minimum of 5 days post-immunization but before the onset of symptoms. For comparison of disease burden, area under curve was calculated for individual disease curves and was divided by the number of days in the protocol (AUC d^{-1}). For p19 blockade, mice were administered anti-p19 (clone G23-8) or istotype (rIgG1, clone HRPM; both BioXcell), 1 mg on d^{-1} and d^{+6} (47).

Histopathology

Mice were euthanized and intracardially perfused with PBS. The spinal cord was extracted and fixed in 10% formalin and 30% sucrose prior to being embedded in paraffin. Five-to-seven-micron coronal sections of the spinal cords were collected on superfrost microscope slides (using microtome, Leica Biosystems HistoCore Autocut) and dried overnight on a drying bench at 37°C. Slides were stored at

room temperature in a slide storage box until staining was performed. Sections

for Hematoxylin and Eosin (H&E) were deparaffinated with xylene, while toluene

was used for Luxol Fast Blue (LFB) sections. Histology was performed as

previously described (67), using standard H&E to visualize lymphocyte infiltration

and LFB to stain myelin in order to visualize areas of demyelination. Briefly,

stained slides were scanned at 20x using a brightfield microscope scanner

(Aperio AT2 DX System), and representative RGB images of the cervical spine

were acquired. Quantification of the staining was performed on Image J

(software, v 1.53K) in a blinded manner. Images were split into separate single

colour channels using the Colour Deconvolution plugin. Thresholding was

performed on the single colour channels for hematoxylin and LFB, and an area

fraction measurement was performed on the white matter of the spinal cord.

Staining is expressed as the percent of total white matter that is stained.

Immunofluorescence

Immunofluorescence (IF) analysis was performed on flash-frozen brain tissue

embedded in optimum cutting temperature medium (OCT). The tissue was cryo-

sectioned (7-10um) and mounted onto positively charged slides. IF staining were

performed with antibodies against surface immune antigens such as CD4, B220,

CD45 and CD11c and in some cases against ECM antigens: Fibronectin,

PDGFRα/β simultaneously. Slides were imaged using a widefield inverted

microscope (AxioObserver 7) and analysis was done using Image J software.

ELISA

Splenocytes and draining lymph nodes were isolated from NOD and IgH[MOG] mice 5d post-immunization. Cells were cultured at a concentration of 10×10^6 cells mL^{-1} in T cell media and stimulated, or not, with 10 mg mL^{-1} MOG$_{[35-55]}$. Supernatants were collected at d5 of the culture for the analysis of T cell cytokine production using a combination of commercially available antibodies (Biolegend). Briefly, the capture antibodies used were: purified anti-mouse IFNg (clone R4-6A2) and purified anti-mouse IL-17 (clone TC11-18H10.1). The detection antibodies used were: biotin anti-mouse IFNg; clone XMG1.2, biotin anti-mouse IL-17; clone TC11-8H4. Following the incubation with the avidin-horseradish peroxidase (Biolegend) and TMB substrate (Mendel Scientific), colorimetric readings were performed using a SpectraMax i3 Microplate Reader.

Measurement of serum immunoglobulin

Blood was collected from Igh[MOG] mice and WT mice at endpoint. Serum was collected by centrifuging blood samples at 2000g for 10min. Total anti-MOG IgG was quantified by using SensoLyte Anti-Mouse MOG(1-125) IgG Quantitative ELISA Kit (Anaspec).

Isolation of CNS-infiltrating mononuclear cells

Mice were euthanized and perfused intracardially with PBS. Brain and spinal cord were dissected from the skull and vertebral column respectively and were

prepared as previously described (66). Briefly, CNS tissues were digested with

liberase (Roche) and DNAse I (Sigma) and cells were enriched using a 35%

Percoll (GE Healthcare) gradient.

Isolation of dura mater cells

Mice were injected intravenously with fluorochrome-labeled anti-CD45 antibody

15 minutes prior to euthanasia. Mice were then perfused intracardially with PBS

and skull was gently removed mechanically to peel the dura from the edges of

the skull cap. Samples were then incubated for 30 minutes at 37°C in digestion

buffer (1mg/mL collagenase D (Sigma) and 50ug/mL of DNAse 1 (Sigma) in

RPMI). Samples were mechanically dissociated through a 100um cell strainer

and flowthrough was centrifuged at 300g for 5 minutes at 4°C. Finally, the

supernatant was discarded, and cell pellet was resuspended for flow cytometry

analysis.

Flow cytometry

Single cell suspensions were obtained from spleens, lymph nodes and CNS of

EAE mice. For detection of surface antigens, cells were stained with Fixable

Viability Dye (eBioscience) and incubated with Fc Block (Biolegend) prior to

staining with antibodies against surface antigens (CD45, CD4, CD8, CD19,

CD11b, CD11c, B220, Ly6G, IgD, IgM, FAS, GL7, PD-1, CXCR5, ICOS, I-Ag^7,

GFAP; details in following section). For detection of intracellular cytokines, cells

were first stimulated with 50ng ml^{-1} PMA (Sigma), 1µM ionomycin (Sigma) and

1μL mL^{-1} GolgiStop (BD) for 4 hours at 37°C, prior to being labeled with viability

indicator, Fc Block and relevant surface antigens as above. They were then fixed

and permeabilized (Fixation Buffer and Intracellular Staining Perm Wash Buffer,

both Biolegend) and stained for intracellular markers (IFN-γ, IL-17A, TNF-α, IL-6,

GM-CSF, IL-12p40, IL-12p35, IL-23p19, IL-23; Bcl6, IL-21, CXCL13; details in

following section). CM-H2DCFDA, a general oxidative stress indicator, was used

to detect total ROS production. Cells were incubated with 1 μM CM-H2DCFDA

(Thermo Fisher Scientific, C6827) at 37°C for 30 minutes, followed by surface

proteins staining as described above. Samples were analyzed on a FACS Aria

(BD) and data were analyzed using FlowJo software (Treestar).

Flow cytometry antibodies

The following monoclonal antibodies against mouse antigens were used: CD45,

clone A20 (Biolegend); CD11b, clone M1/70 (eBioscience); CD11c, clone N418

(Biolegend); Ly6G, clone 1A8 (BD Biosciences); CD4, clone RM4-5

(eBioscience); CD8, clone 53-6.7 (Biolegend); CD19, clone 1D3 (eBioscience);

B220, clone RA3-6B2 (eBioscience); IgD, clone 11-26c (eBioscience); IgM, clone

11/41 (eBioscience); FAS, clone 15A7 (eBioscience); GL7, clone GL-7

(eBioscience); PD-1, clone J43 (eBioscience), CXCR5, clone SPRCL5

(eBioscience); ICOS, clone C398.4A (eBioscience); GFAP, clone 1B4 (BD

Biosciences); I-Ag^7, clone 39-10-8 (Biolegend); IFN-γ, clone XMG1.2

(eBioscience); TNF-α, clone MP6-XT22 (eBioscience); IL-17a, clone TC11-

18H10.1 (Biolegend); GM-CSF, clone MP1-22E9 (eBioscience); IL-6, clone MP5-

20F3 (eBioscience); IL-21, clone mhalx21 (eBioscience); IL-23 heterodimer,

clone IC18871P (R&D Systems); IL-23p19, clone fc23pg (eBioscience); IL-

12p35, clone 4D10p35 (eBioscience); IL-12p40, clone C17.8 (eBioscience);

BCL6, clone IG191E/A8 (eBioscience); CXCL13, clone D58CX13 (eBioscience);

CM-H2DCFDA, clone C6827 (Thermo Fisher Scientific).

Immunofluorescence antibodies

The following monoclonal antibodies against mouse antigens were used: CD4

(RM4-5), B220 (RA3-6B2), CD11c (N418), CD45.2 (104), PDGFRα (APA5), and

PDGFRβ (APB5), all of which are from eBioscience; and fibronectin (GW20021F,

Sigma); smooth muscle actin (1A4, Sigma); IL-23 heterodimer (IC18871P, R&D

Systems).

Antigen presentation assay

Single cell suspensions were obtained from the spleens of unimmunized WT and

IgH[MOG] mice. Cells were labeled with CD43 (Ly-48) Microbeads (Miltenyi), and

$CD43^+$ leukocytes (all leukocytes except resting B cells) were depleted on a

magnetic MACS column (Miltyeni). Unlabelled $CD43^-$ cells were collected and

were subsequently stained with anti-mouse CD19. $CD19^+$ B cells were purified

using high-speed cell sorting. In parallel, $CD4^+$ T cells were purified from the

spleens of 1C6 mice using mouse CD4 MicroBeads (Miltenyi) and labeled with

CellTrace Violet (CTV; Thermo Fisher Scientific). CTV-labeled 1C6 $CD4^+$ T cells

were cultured with B cells at a ratio of 1 CD4: 1 B, with 0, 1 or 10µg ml^{-1} MOG$_{[35-}$

$_{55]}$ for 72 hours. CTV dilution was assessed by flow cytometry.

Tfh culture

The protocol of Gao et al was adapted (68). Briefly, 5×10^5 splenocytes from NOD

mice were stimulated 24h with LPS (1 µg mL^{-1}). 1×10^4 MOG$_{[35-55]}$-specific 1C6

CD4$^+$CD62L^{hi} T cells were then plated on top of the splenocytes with 1µg mL^{-1}

MOG$_{[35-55]}$, plus 50 ng ml^{-1} IL-21 and 100 ng mL^{-1} of IL-6 (Tfh) or no additional

cytokines (Th0) for 72h.

Analysis of CSF from MS-affected individuals

5mL of CSF was collected from consenting adult participants with MS at The

Health Sciences Centre neurology clinic in St. John's, NL between July 2021 and

May 2022. For CSF immune phenotyping, 2.5mL of CSF was centrifuged for 10

minutes at 300g and the CSF was removed. The CSF cell pellet was

resuspended in 100uL of flow buffer (1% bovine albumin serum, 2mM EDTA,

2mM sodium azide in PBS). The CSF cell suspension was added to a

DURAclone IM Phenotyping BASIC tube (Beckman Coulter), mixed and

incubated at 4°C for 30 minutes. The cells were washed with 4mL flow buffer,

centrifuged at 300g for 5 minutes, decanted and resuspended in 100µL 2%

paraformaldehyde. Data was acquired from the whole sample using a Cytoflex

flow cytometer (Beckman Coulter) and analysed using FlowJo software. The

CD19$^+$ cell count was expressed as a percentage of the total number of CD45$^+$

cells.

For measurement of CSF cell *Il23a* expression, 2.5 mL of human CSF was

centrifuged at 300g for 10 minutes and the CSF supernatant was carefully

removed. The CSF cell pellet was resuspended in 500ul of Trizol and stored at -

80°C. Total RNA was extracted using chloroform extraction followed by an

RNeasy Micro kit (Qiagen). 200ng of RNA was used to synthesize cDNA with the

M-MLV Reverse Transcriptase kit (Invitrogen). *Il23a* transcript expression was

quantified using *Il23a* (FAM labelled) and GAPDH (VIC labelled) Taqman

probe/primer assays (Invitrogen), Fast Advanced Master mix (Invitrogen) and a

ViiA7 Real-time PCR system (Applied Biosystems).

mRNA-seq

At experimental endpoints, CNS tissues from WT and IgH[MOG] mice were

necropsied and mononuclear cells were isolated. Live CD19$^+$ B cells and

CD4$^+$PD1$^+$CXCR5$^-$ Tph cells were purified by high-speed sorting (FACSAria II)

and were directly processed using NEBNext Single Cell/Low Input RNA Library

Prep Kit (New England Biolabs), according to the manufacturer's instructions, for

mRNA sequencing library preparation. Total mRNA was enriched using poly(A)+

primers and template-switching oligo synthesize double-stranded cDNA with

reverse transcriptase. A PCR amplification of 21 cycles was performed to

increase cDNA yields followed by a purification step with AxyPrep Mag PCR

Clean-up kit (Axygen, Big Flats, NY, USA). Enzymatic fragmentation of cDNA

and end-repair steps were followed by ligation of NEBNext adaptors and PCR

enrichment step of 11 cycles to incorporate specific indexed adapters for the

multiplexing. The quality of final amplified libraries was examined with a DNA

screentape D1000 on a TapeStation 2200 and the quantification was done on the

QuBit 3.0 fluorometer (ThermoFisher Scientific, Canada). Subsequently, mRNA-

seq libraries with unique dual indexes were pooled together in equimolar ratio

and sequenced for paired-end 100 pb sequencing on NovaSeq 6000 at the Next-

Generation Sequencing Platform, Genomics Center, CHU de Québec-Université

Laval Research Center. The average insert size for the libraries was 253 bp. The

mean coverage/sample was 24M paired-end reads.

Bioinformatics analyses

Paired-end mRNAseq fastq datasets were trimmed with trimmomatic v0.39 using

the following command:

java -jar trimmomatic-0.39.jar PE -phred 33 sample_R1.fastq sample_R2.fastq

trimmed/paired_sample_R1.fastq   trimmed/unpaired_sample_R1.fastq

trimmed/paired_sample_R2.fastq   trimmed/unpaired_sample_R2.fastq

ILLUMINACLIP:TruSeq3-PE.fa:2:30:10   LEADING:30   TRAILING:30

SLIDINGWINDOW:4:15 MINLEN:30

Trimmed fastq files were mapped on the *M. musculus* genome (Ensemble;

Mus_musculus.GRCm39.cds.all.fa.idx) and quantified using kallisto 0.44.0.

Differential expression between WT and IgH[MOG] conditions cells was performed

using DESeq2 v1.38.3. For identifying relevant pathways, we used Gene

Ontology enrichment analysis. We focused on biological processes on all

significantly differentially expressed genes using Benjamini-Hochberg method to

adjust for the false discovery rate. We also used the simplify function from

ClusterProfiler package to remove redundant terms. Finally, GSEA from the

Broad Institute of UCSD was used to calculate the enrichment score of different

gene sets.

703
Statistical analyses

Comparisons between two groups were made by *t*-test. Fisher's exact test was

used to test for differences in the frequency of mice attaining ethical endpoints.

For calculation of Tph frequency at different disease stages, Tukey's multiple

comparisons test was used. Two-tailed analyses were used in all instances. All

error bars represent s.e.m. All statistical analyses were conducted using Prism

software (GraphPad).

Abbreviations

Ab, antibody; CNS, central nervous system; CSF, cerebrospinal fluid; CTV,

CellTrace Violet; DC, Dendritic cell; DEG, differentially expressed gene; EAE,

experimental autoimmune encephalomyelitis; GC, Germinal center; GM-CSF,

granulocyte and macrophage colony stimulating factor; H&E, hematoxylin &

eosin; IFN, interferon; Ig, immunoglobulin; IL, interleukin; LFB, Luxol fast blue;

MHC, major histocompatibility complex; MOG, myelin oligodendrocyte

glycoprotein; MS, multiple sclerosis; NOD, nonobese diabetic; PLP, proteolipid

protein; preonset, PO; RR, relapsing/remitting; SP, secondary progressive; Tfh, T

follicular helper; TLO, Tertiary lymphoid organ; TNF, tumor necrosis factor; Tph,

T peripheral helper; WT, wildtype.

Declarations

Competing interests

M.R. held a research contract with Remedy Pharmaceuticals (2019-2020), with

funds paid to the CHU de Québec. This contract is unrelated to the work in this

manuscript.

Funding

The work was supported by a Biomedical Discovery Research Grant from the MS

Society of Canada (#3781) and a CIHR Project Grant. MRF and PG are

supported by Ph.D. studentships from the Fonds recherche du Québec – Santé

(FRQS). PMIAD was supported by a doctoral studentship from the Multiple

Sclerosis Society of Canada. NB is Junior-2 scholar, and MR is a Senior scholar,

of the FRQS.

Authors' contributions

MRF directed the project conducted experiments and helped write the manuscript. PMIAD, NF, IA, APY and JB conducted experiments. RP, AR, BM, SL and OR conducted histological analyses. PG and NB assisted with serum detection of immunoglobulins. CJ-B and AD assisted with bioinformatics analyses. CSM contributed to writing of the manuscript. MR supervised the project and wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank Vijay Kuchroo for providing us with NOD-background IgH[MOG] and 1C6 mice respectively. We thank Françoise Morin for critical discussions; Vincent Desrosiers, Alexandre Brunet, Stéphanie Fiola for technical assistance with flow cytometry; and Kim Larose, Andrée Brisson, Mathieu Vallière-St-Amant, Cindy Ouellet and the veterinary service of Université Laval for technical assistance and collaboration regarding animal care.

Data availability statement

All datasets will be made upon reasonable request to the corresponding author.

Figure legends

Figure 1. Active immunization with MOG$_{[35-55]}$ induces severe EAE in

IgH$^{[MOG]}$ mice in the absence of increased peripheral inflammation. A. *Left,*

Disease curves and linear regression of males (WT, n=10, IgH$^{[MOG]}$, n=10) and

females (WT, n=10, IgH$^{[MOG]}$, n=10) immunized with MOG$_{[35-55]}$ and monitored for

development of EAE. ****, p<0.0001. Representative of 2 experiments in each

sex. *Right,* AUC analysis of disease curves for all mice studied (30 WT, 30

IgH$^{[MOG]}$). **B.** WT and IgH$^{[MOG]}$ spinal cords were sectioned and stained with

hematoxylin & eosin (H&E; immune infiltration) and Luxol fast blue (LFB; myelin).

Slides were scanned at 10X magnification for quantification. WT, n=4, IgH$^{[MOG]}$,

n=4. *, p<0.05, *t*-test. **C.** WT (n=3) and IgH$^{[MOG]}$ (n=3) mice were immunized with

MOG$_{[35-55]}$. LN cells were isolated at disease onset and IFNγ and IL-17

expression were determined by flow cytometry. Gated on live CD4$^+$CD44$^+$

events. **D.** WT or IgH$^{[MOG]}$ B cells were co-cultured with CellTrace Violet-labeled

MOG$_{[35-55]}$-specific 1C6 T cells that were pulsed, or not, with the indicated

concentrations of MOG$_{[35-55]}$. Gate frequencies indicate the percentage of cells

undergoing at least one cell division. Representative of triplicate cultures. **E.** Sera

were collected from MOG$_{[35-55]}$-immunized WT (n=5) and IgH$^{[MOG]}$ (n=5) mice,

and concentration of MOG-specific IgG was assessed by ELISA. n.s., not

significant; *, p<0.05; ****, p<0.0001; linear regression analysis (**A**), *t*-test (**A-C,**

E).

Figure 2. Increased lymphocytic aggregation in the meninges of immunized

IgH$^{[MOG]}$ mice. A. Mononuclear cells were isolated from the CNS of immunized

WT (n=6) and IgH[MOG] (n=12) mice. Frequency of CD19$^+$ B cells, CD4$^+$ and CD8$^+$

T cells, Ly6G$^+$ neutrophils, CD11b$^+$CD11c$^-$ macrophages and CD11c$^+$ dendritic

cells were enumerated by flow cytometry. **B.** Brain and spinal cord meningeal

sections from MOG$_{[35-55]}$-immunized WT (n=7) or IgH[MOG] (n=7) mice were

assessed by immunofluorescence (IF) for expression of B220 (red), CD4 (blue),

and CD11c (green), and lymphocyte aggregates from the cerebellar meningeal

compartment were enumerated. **C.** Assessment of the ECM near meninges in

the cerebellum was determined by IF. Expansion on the meningeal ECM is

shown by fibronectin (green) and PDGFRα/β (blue) which overlap with B220 B

cell clusters. Dotted lines outline the meningeal membrane and (*) denotes

vascular endothelium.

Figure 3. Inflammatory B cells facilitate Th17 responses in IgH[MOG] mice. A.

The frequency of immature (IgM^{mid}IgD^{mid}), mature (IgM^{hi}IgD^{hi}) and class-switched

(CS; IgM$^-$IgD$^-$) B cells were assessed from WT (n=5) and IgH[MOG] (n=5) CNS.

Gated on live B220$^+$ events. **B.** Expression of the indicated cytokines was

assessed from WT (n=5) and IgH[MOG] (n=5) CNS B cells. Gated on live CD19$^+$

events. **C.** WT (n=5) and IgH[MOG] (n=5) were immunized with MOG$_{[35-55]}$ and

CNS-infiltrating CD4$^+$ T cells were isolated at endpoints. Expression of the

indicated cytokines was assessed by intracellular flow cytometry. Gated on live

CD4$^+$ events. **D**. NOD.*Scid* mice were passively infused with 2×10^6 WT or

IgH[MOG] B cells, and received 2×10^6 1C6 Th17 cells 7 days later. Mice were

monitored for signs of EAE. *Left*, disease curve from one of 2 transfer

experiments; *right*, AUC comparison of mice from both transfers (n=7, both

groups of recipient mice). **E**. CNS-infiltrating CD4$^+$ T cells were isolated from

adoptive transfer recipients in (**D**) and the indicated cytokines were assessed by

flow cytometry. Gated on live CD4$^+$ events. n.s., not significant; *, p<0.05; **,

p<0.01; ***, p<0.001; ****, p<0.0001; *t*-test (**A-E**); linear regression (**D**).

Figure 4. CNS IgH$^{[MOG]}$ B cells upregulate IL-23. A. NOD (n=6) and IgH$^{[MOG]}$

(n=6) mice were immunized with MOG$_{[35-55]}$ and B cells were isolated from CNS

tissues at endpoints. Expression of IL-23 heterodimer was assessed by

intracellular flow cytometry. **B.** Brain and spinal cord meningeal sections from

MOG$_{[35-55]}$-immunized WT (n=3) or IgH$^{[MOG]}$ (n=3) mice were assessed by

immunofluorescence (IF) for expression of IL-23 (aqua), B220 (red), CD4 (blue).

C. IgH$^{[MOG]}$ mice were immunized with MOG$_{[35-55]}$ and were treated on day (-1)

and day 6 with 1 mg anti-p19 (n=5) or rIgG1 isotype (n=5). Disease curves

representative of 3 experiments. AUC calculations include all mice studied; anti-

p19 (n=14) or rIgG1 isotype (n=14). **D.** Cerebellar meningeal sections from

MOG$_{[35-55]}$-immunized IgH$^{[MOG]}$ mice, treated with isotype (n=6) or anti-p19 (n=6),

were assessed for lymphocyte clusters. **E.** CNS-infiltrating mature, immature and

class-switched B cells were assessed by flow cytometry from anti-p19 (n=4) or

rIgG1 isotype (n=4)-treated immunized IgH$^{[MOG]}$ mice. Gated on live B220+

events. **F.** CD4$^+$ T cells were isolated from p19 (n=5) or rIgG1 isotype (n=5)-

treated immunized IgH$^{[MOG]}$ mice and the indicated cytokines were assessd by

flow cytometry. **G.** *Il23a* was assessed from CSF immune cells of MS-affected

individuals (10 RRMS, 1 PPMS; 9 females, 2 males) by qPCR and was

correlated to the frequency of CD19$^+$ B cells within the CD45$^+$ CSF leukocyte

population of each patient. n.s., not significant; *, p<0.05; **, p<0.01; ***,

p<0.001, *** *, p<0.0001; *t*-test (**A-F**); linear regression (**G**).

Figure 5. Tph cells infiltrate the IgH$^{[MOG]}$ CNS. A. WT (n=6) and IgH$^{[MOG]}$ (n=6)

mice were immunized with MOG$_{[35-55]}$ and T cells were isolated from CNS at

endpoints. Expression of PD-1 and CXCR5 was assessed by flow cytometry. **B.**

WT or IgH$^{[MOG]}$ mice were immunized and sacrificed at onset (n=4 each), peak

(n=3 each), recovery (n=3 each). CNS-infiltrating CD4$^+$ T cells were isolated from

these groups, as well as from mice that did not develop disease (n=4 each) and

the frequency of PD-1$^+$CXCR5$^-$ cells was assessed. **C.** Expression of IL-21,

CXCL13 and ICOS was assessed from CNS-infiltrating CD4$^+$ T cells taken from

immunized WT (n=6) and IgH$^{[MOG]}$ (n=6) mice, and expression between PD-1$^-$

and PD-1$^+$ CD4$^+$ subpopulations was compared. **D.** Expression of IFNγ and IL-17

was assessed from PD-1$^-$ and PD-1$^+$ CD4$^+$ T cells from immunized WT or

IgH$^{[MOG]}$ mice. **E.** WT (n=5) and IgH$^{[MOG]}$ (n=5) mice were immunized with

MOG$_{[35-55]}$ and T cells were isolated from the dura mater at endpoints.

Expression of PD-1 and CXCR5 was assessed by flow cytometry. **F.** T cells were

isolated from spleen and CNS at endpoints from anti-p19 (n=5) or isotype-treated

(n=5) mice. Expression of PD-1 and CXCR5 was assessed by flow cytometry. All

flow cytometry data gated on live CD4$^+$ events. n.s., not significant; *, p<0.05; **,

p<0.01; ***, p<0.001, ****, p<0.0001; unpaired *t*-test (**A, F**), Tukey's multiple

comparisons test (**B, D**), paired *t*-test (**C**).

Figure 6. CNS-infiltrating IgH[MOG] B cells and Tph cells show increased

production of reactive oxygen species. A. GO term analysis of the

transcriptome of CNS-infiltrating B cells and Tph cells from immunized IgH$^{[MOG]}$

(n=3) vs WT (n=3) at endpoint. **B.** Normalized enrichment scores (NES) for

neurodegeneration and oxidative stress pathways from B cell and Tph cell

transcriptomes. **C-F.** Uptake of ROS indicator CM-H2DCFDA from B cells (**C, E**)

or Tph cells (**D, F**) from immunized IgH$^{[MOG]}$ vs WT mice (**C, D**; n=5 each) or from

immunized and anti-p19 vs isotype-treated IgH$^{[MOG]}$ mice (E, F; n=5 each).

Supplemental Figure 1. A. WT NOD mice (n=5) were immunized with MOG$_{[35-55]}$

and were monitored for signs of EAE for 120d. **B.** Splenic and LN cells were

isolated from immunized WT (n=3) and IgH$^{[MOG]}$ (n=3) cells at disease onset, and

were stimulated with 10 μg mL^{-1} MOG$_{[35-55]}$. Supernatants were collected at d5

and IFNγ and IL-17 were assessed by ELISA. **C.** WT (n=8) and IgH$^{[MOG]}$ (n=8)

mice were immunized with MOG$_{[35-55]}$, and CD4^{+} T cells were isolated from

cervical LNs at endpoints and assessed for expression of the indicated cytokines.

Supplemental Figure 2. Mononuclear cells were isolated from the CNS of

immunized WT (n=6) and IgH$^{[MOG]}$ (n=12) mice. Frequency and absolute

numbers of CD19^{+} B cells, CD4^{+} and CD8^{+} T cells, Ly6G^{+} neutrophils,

$CD11b^+CD11c^-$ macrophages and $CD11c^+$ dendritic cells were enumerated by

flow cytometry.

Supplemental Figure 3. A. Splenic immature, mature and CS B cells were

assessed from the same mice as in **Figure 3A**. **B.** MHC class II (I-Ag^7)

expression was assessed from CNS B cells at endpoint from $MOG_{[35-55]}$-

immunized NOD (n=5) or IgH$^{[MOG]}$ (n=5) mice. **C.** Germinal center B cells (GL-

7$^+$Fas$^+$) were assessed from the spleens and CNS of $MOG_{[35-55]}$-immunized NOD

(n=5) or IgH$^{[MOG]}$ (n=5) mice. **D.** Representative plots for the data presented in

Figure 3E. *, $p<0.05$; ***, $p<0.001$, n.s., not significant.

Supplemental Figure 4. A. WT (n=10) and IgH$^{[MOG]}$ (n=10) mice were

immunized with $MOG_{[35-55]}$ and B cells were isolated from CNS tissues at

endpoints. Expression of IL-12p35, IL-12p40 and IL-23p19 were assessed by

intracellular flow cytometry. **B.** Dendritic cells (CD11c$^+$), astrocytes (GFAP$^+$),

resting microglia (CD45^{dim}CD11b^{dim}) and activated microglia (CD45^{hi}CD11b^{hi})

were assessed for expression of p19 and p40 from the CNS of $MOG_{[35-55]}$-

immunized WT (n=5) or IgH$^{[MOG]}$ (n=5) mice. DCs, microglia; n=5 each group;

astrocytes, n=10 each group. **C.** ECM deposition underneath meningeal

lymphoid aggregates was assessed from the samples in **Figure 4D**. **D**.

Inflammatory foci (H&E) and demyelination (LFB) were quantified from cervical

spine of immunized IgH$^{[MOG]}$ mice treated with anti-p19 (n=3) or rIgG1 isotype

(n=3).

Supplemental Figure 5. A. Splenic CD4$^+$ T cells were assessed for expression

of PD-1 and CXCR5 from the same mice as in **Figure 5A**. **B.** Expression of Bcl6

was assessed from WT or IgH$^{[MOG]}$ PD-1$^+$CXCR5$^-$ CNS-infiltrating T cells versus

in vitro generated 1C6 Tfh or Th0. **C.** Expression of IL-21 was compared between

WT (n=5) and IgH$^{[MOG]}$ (n=5) CNS-infiltrating CD4$^+$ T cells. All flow cytometry data

gated on live CD4$^+$ events. n.s., not significant; ***, $p<0.001$; *t*-test. **D.** Volcano

plot of mRNA-seq data depicting DEG (1.5x-fold increase, $p_{adj}<0.05$) between

WT (n=3) and IgH$^{[MOG]}$ (n=3) CNS-infiltrating PD1$^+$CXCR5$^-$CD4$^+$ T cells. The top

10 DEG are labeled on the plot. **E.** Normalized counts for Tph markers *Il21*,

Cx3cr1, *Cxcl13*, *Ccr2*, *Icos* between WT and IgH$^{[MOG]}$ Tph from mRNAseq. **F.**

Normalized counts for Il17a, Il17f. p-values from mRNA-seq data are p_{adj}.

References

915
1. Kingwell E et al. Incidence and prevalence of multiple sclerosis in Europe: a
systematic review. [Internet]. *BMC Neurol* 2013;13(1):128.

2. Larochelle C, Uphaus T, Prat A, Zipp F. Secondary Progression in Multiple
Sclerosis: Neuronal Exhaustion or Distinct Pathology? *Trends in Neurosciences*
2016;39(5):325–339.

3. Thompson A. Overview of primary progressive multiple sclerosis (PPMS):
similarities and differences from other forms of MS, diagnostic criteria, pros and
cons of progressive diagnosis. [Internet]. *Mult. Scler.* 2004;10 Suppl 1:S2–7.

4. Baranzini SE et al. B cell repertoire diversity and clonal expansion in multiple
sclerosis brain lesions. [Internet]. *J. Immunol.* 1999;163(9):5133–5144.

5. Owens GP et al. Restricted use of VH4 germline segments in an acute multiple
sclerosis brain. [Internet]. *Ann. Neurol.* 1998;43(2):236–243.

6. Owens GP et al. Single-cell repertoire analysis demonstrates that clonal
expansion is a prominent feature of the B cell response in multiple sclerosis
cerebrospinal fluid. [Internet]. *J. Immunol.* 2003;171(5):2725–2733.

7. Magliozzi R et al. Meningeal B-cell follicles in secondary progressive multiple
sclerosis associate with early onset of disease and severe cortical pathology.
Brain 2007;130(Pt 4):1089–1104.

8. Uccelli A, Aloisi F, Pistoia V. Unveiling the enigma of the CNS as a B-cell
fostering environment. [Internet]. *Trends Immunol.* 2005;26(5):254–259.

9. Genain CP, Cannella B, Hauser SL, Raine CS. Identification of autoantibodies
associated with myelin damage in multiple sclerosis. [Internet]1999;5(2):170–
175.

10. Hauser SL et al. B-cell depletion with rituximab in relapsing-remitting multiple
sclerosis. *N. Engl. J. Med.* 2008;358(7):676–688.

11. Kappos L et al. Ocrelizumab in relapsing-remitting multiple sclerosis: a phase
2, randomised, placebo-controlled, multicentre trial. [Internet]. *Lancet*
2011;378(9805):1779–1787.

12. Hauser SL et al. Ofatumumab versus Teriflunomide in Multiple Sclerosis. *N.*
Engl. J. Med. 2020;383(6):546–557.

13. Montalban X et al. Ocrelizumab versus Placebo in Primary Progressive
Multiple Sclerosis. *N. Engl. J. Med.* [published online ahead of print: December
21, 2016]; doi:10.1056/NEJMoa1606468

14. Lehmann-Horn K, Kronsbein HC, Weber MS. Targeting B cells in the
treatment of multiple sclerosis: recent advances and remaining challenges.
[Internet]2013;6(3):161–173.

15. Rangachari M, Kuchroo VK. Using EAE to better understand principles of
immune function and autoimmune pathology. *J. Autoimmun.* 2013;45:31–39.

16. Litzenburger T et al. B lymphocytes producing demyelinating autoantibodies:
development and function in gene-targeted transgenic mice. *J. Exp. Med.*
1998;188(1):169–180.

17. Krishnamoorthy G, Lassmann H, Wekerle H, Holz A. Spontaneous
opticospinal encephalomyelitis in a double-transgenic mouse model of
autoimmune T cell/B cell cooperation. [Internet]2006;116(9):2385–2392.

18. Pöllinger B et al. Spontaneous relapsing-remitting EAE in the SJL/J mouse:
MOG-reactive transgenic T cells recruit endogenous MOG-specific B cells. *J.*
Exp. Med. 2009;206(6):1303–1316.

19. Dang AK, Jain RW, Craig HC, Kerfoot SM. B cell recognition of myelin
oligodendrocyte glycoprotein autoantigen depends on immunization with protein
rather than short peptide, while B cell invasion of the CNS in autoimmunity does
not. *J. Neuroimmunol.* 2015;278:73–84.

20. Zhong X et al. Reciprocal generation of Th1/Th17 and T(reg) cells by B1 and
B2 B cells. *Eur. J. Immunol.* 2007;37(9):2400–2404.

21. Wang Y, Rothstein TL. Induction of Th17 cell differentiation by B-1 cells.
Front Immunol 2012;3:281.

22. Barbosa RR et al. Primary B-cell deficiencies reveal a link between human IL-
17-producing CD4 T-cell homeostasis and B-cell differentiation. *PLoS ONE*
2011;6(8):e22848.

23. Bar-Or A et al. Abnormal B-cell cytokine responses a trigger of T-cell-
mediated disease in MS? *Ann. Neurol.* 2010;67(4):452–461.

24. van de Veerdonk FL et al. The anti-CD20 antibody rituximab reduces the
Th17 cell response. *Arthritis Rheum* 2011;63(6):1507–1516.

25. Encinas JA et al. QTL influencing autoimmune diabetes and
encephalomyelitis map to a 0.15-cM region containing Il2. *Nat. Genet.*
1999;21(2):158–160.

26. Rao DA et al. Pathologically expanded peripheral T helper cell subset drives
B cells in rheumatoid arthritis. *Nature* 2017;542(7639):110–114.

27. Mayo L et al. IL-10-dependent Tr1 cells attenuate astrocyte activation and
ameliorate chronic central nervous system inflammation. *Brain* 2016;139(Pt
7):1939–1957.

28. Farez MF et al. Toll-like receptor 2 and poly(ADP-ribose) polymerase 1
promote central nervous system neuroinflammation in progressive EAE.
[Internet]2009;10(9):958–964.

29. Baker D et al. Autoimmune encephalomyelitis in NOD mice is not initially a
progressive multiple sclerosis model. *Ann Clin Transl Neurol* 2019;6(8):1362–
1372.

30. Anderson AC et al. A transgenic model of central nervous system
autoimmunity mediated by CD4+ and CD8+ T and B cells. *J. Immunol.*
2012;188(5):2084–2092.

31. Yeola AP et al. Endogenous T Cell Receptor Rearrangement Represses
Aggressive Central Nervous System Autoimmunity in a TcR-Transgenic Model
on the Non-Obese Diabetic Background. *Front Immunol* 2019;10:3115.

32. Doss PMIA et al. Male sex chromosomal complement exacerbates the
pathogenicity of Th17 cells in a chronic model of central nervous system
autoimmunity. *Cell Rep* 2021;34(10):108833.

33. Legroux L, Arbour N. Multiple Sclerosis and T Lymphocytes: An Entangled
Story. *J Neuroimmune Pharmacol* 2015;10(4):528–546.

34. Obermeier B et al. Matching of oligoclonal immunoglobulin transcriptomes
and proteomes of cerebrospinal fluid in multiple sclerosis.
[Internet]2008;14(6):688–693.

35. Howell OW et al. Meningeal inflammation is widespread and linked to cortical
pathology in multiple sclerosis. [Internet]. *Brain* 2011;134(Pt 9):2755–2771.

36. deLuca LES et al. Substrain differences reveal novel disease-modifying gene
candidates that alter the clinical course of a rodent model of multiple sclerosis. *J.*
Immunol. 2010;184(6):3174–3185.

37. Pikor NB et al. Integration of Th17- and Lymphotoxin-Derived Signals Initiates
Meningeal-Resident Stromal Cell Remodeling to Propagate Neuroinflammation.
Immunity 2015;43(6):1160–1173.

38. Pikor NB, Prat A, Bar-Or A, Gommerman JL. Meningeal Tertiary Lymphoid
Tissues and Multiple Sclerosis: A Gathering Place for Diverse Types of Immune
Cells during CNS Autoimmunity. *Front Immunol* 2015;6:657.

39. Barr TA et al. B cell depletion therapy ameliorates autoimmune disease
through ablation of IL-6-producing B cells. *J. Exp. Med.* 2012;209(5):1001–1010.

40. Duddy M et al. Distinct effector cytokine profiles of memory and naive human
B cell subsets and implication in multiple sclerosis. *J. Immunol.*
2007;178(10):6092–6099.

41. Li R et al. Proinflammatory GM-CSF-producing B cells in multiple sclerosis
and B cell depletion therapy. *Sci Transl Med* 2015;7(310):310ra166.

42. Domingues HS, Mues M, Lassmann H, Wekerle H, Krishnamoorthy G.
Functional and pathogenic differences of Th1 and Th17 cells in experimental
autoimmune encephalomyelitis. *PLoS ONE* 2010;5(11):e15531.

43. McGeachy MJ et al. The interleukin 23 receptor is essential for the terminal
differentiation of interleukin 17-producing effector T helper cells in vivo. *Nat.*
Immunol. 2009;10(3):314–324.

44. Haines CJ et al. Autoimmune memory T helper 17 cell function and
expansion are dependent on interleukin-23. *Cell Rep* 2013;3(5):1378–1388.

45. Ramgolam VS et al. B cells as a therapeutic target for IFN-β in relapsing-
remitting multiple sclerosis. *J. Immunol.* 2011;186(7):4518–4526.

46. Nitsch L et al. Astrocyte-specific expression of interleukin 23 leads to an
aggravated phenotype and enhanced inflammatory response with B cell
accumulation in the EAE model. *J Neuroinflammation* 2021;18(1):101–16.

47. Chen Y et al. Anti-IL-23 therapy inhibits multiple inflammatory pathways and
ameliorates autoimmune encephalomyelitis. *J. Clin. Invest.* 2006;116(5):1317–
1326.

48. Schröder JB et al. Immune Cell Activation in the Cerebrospinal Fluid of
Patients With Parkinson's Disease. *Front Neurol* 2018;9:1081.

49. Quinn JL, Kumar G, Agasing A, Ko RM, Axtell RC. Role of TFH Cells in
Promoting T Helper 17-Induced Neuroinflammation. *Front Immunol* 2018;9:382.

50. Zhang F et al. Defining inflammatory cell states in rheumatoid arthritis joint
synovial tissues by integrating single-cell transcriptomics and mass cytometry.
Nat. Immunol. 2019;20(7):928–942.

51. Johnston RJ et al. Bcl6 and Blimp-1 are reciprocal and antagonistic
regulators of T follicular helper cell differentiation. *Science* 2009;325(5943):1006–
1010.

52. Bocharnikov AV et al. PD-1hiCXCR5- T peripheral helper cells promote B cell
responses in lupus via MAF and IL-21. *JCI Insight* 2019;4(20).
doi:10.1172/jci.insight.130062

53. Field J et al. A polymorphism in the HLA-DPB1 gene is associated with
susceptibility to multiple sclerosis. [Internet]. *PLoS ONE* 2010;5(10):e13454.

54. Noronha A, Toscas A, Jensen MA. Interferon beta decreases T cell activation
and interferon gamma production in multiple sclerosis. *J. Neuroimmunol.*
1993;46(1-2):145–153.

55. Duda PW, Schmied MC, Cook SL, Krieger JI, Hafler DA. Glatiramer acetate
(Copaxone) induces degenerate, Th2-polarized immune responses in patients
with multiple sclerosis. [Internet]. *J. Clin. Invest.* 2000;105(7):967–976.

56. Stüve O et al. Altered CD4+/CD8+ T-cell ratios in cerebrospinal fluid of
natalizumab-treated patients with multiple sclerosis. [Internet]. *Arch. Neurol.*
2006;63(10):1383–1387.

57. Myhr K-M, Torkildsen Ø, Lossius A, Bø L, Holmøy T. B cell depletion in the
treatment of multiple sclerosis. *Expert Opin Biol Ther* 2019;19(3):261–271.

58. Vanderlugt CL, Miller SD. Epitope spreading in immune-mediated diseases:
implications for immunotherapy. [Internet]2002;2(2):85–95.

59. Mitsdoerffer M et al. Proinflammatory T helper type 17 cells are effective B-
cell helpers. [Internet]. *Proc. Natl. Acad. Sci. U.S.A.* 2010;107(32):14292–14297.

60. Hartlehnert M et al. Bcl6 controls meningeal Th17-B cell interaction in murine
neuroinflammation. *Proc. Natl. Acad. Sci. U.S.A.* 2021;118(36).
doi:10.1073/pnas.2023174118

61. Lee Y et al. IL-21R signaling is critical for induction of spontaneous
experimental autoimmune encephalomyelitis. *J. Clin. Invest.* 2015;125(11):4011–
4020.

62. Xiao S et al. Checkpoint Receptor TIGIT Expressed on Tim-1+ B Cells
Regulates Tissue Inflammation. *Cell Rep* 2020;32(2):107892.

63. Liu Z et al. Increased circulating PD-1hiCXCR5- peripheral T helper cells are
associated with disease activity of ANCA-associated vasculitis. *Clin. Exp.*
Immunol. 2022;207(3):287–296.

64. Bartova R et al. Changes in levels of oxidative stress markers and some
neuronal enzyme activities in cerebrospinal fluid of multiple sclerosis patients.
Neuro Endocrinol Lett 2016;37(2):102–106.

65. Irani DN. Cerebrospinal fluid protein carbonylation identifies oxidative
damage in autoimmune demyelination. *Ann Clin Transl Neurol* 2017;4(2):145–
150.

66. Boivin N, Baillargeon J, Doss PMIA, Roy A-P, Rangachari M. Interferon-β
suppresses murine Th1 cell function in the absence of antigen-presenting cells.
PLoS ONE 2015;10(4):e0124802.

67. Rojas OL et al. Recirculating Intestinal IgA-Producing Cells Regulate
Neuroinflammation via IL-10. *Cell* 2019;176(3):610–624.e18.

68. Gao X, Wang H, Chen Z, Zhou P, Yu D. An optimized method to differentiate
mouse follicular helper T cells in vitro. *Cell. Mol. Immunol.* 2020;17(7):779–781.

1094

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.