Measuring Robustness of Brain Networks in Autism Spectrum Disorder with Ricci Curvature

Anish K. Simhal[1, *, +]**, Kimberly L.H. Carpenter**[2, +]**, Saad Nadeem**[3, +]**, Joanne Kurtzberg**[4]**, Allen Song**[5]**, Allen Tannenbaum**[6]**, Guillermo Sapiro**[1, 7]**, and Geraldine Dawson**[2, 4, 8]

[1]Department of Electrical and Computer Engineering, Duke University
[2]Duke Center for Autism and Brain Development, Department of Psychiatry and Behavioral Sciences, Duke University School of Medicine
[3]Department of Medical Physics, Memorial Sloan Kettering Cancer Center
[4]Marcus Center for Cellular Cures, Duke University Medical Center
[5]Brain Imaging and Analysis Center, Duke University, Durham, North Carolina
[6]Departments of Computer Science, Applied Mathematics and Statistics, Stony Brook University
[7]Departments of Biomedical Engineering, Computer Sciences, and Math, Duke University
[8]Duke Institute for Brain Sciences, Duke University
[*]aksimhal@gmail.com, corresponding author
[+]These authors contributed equally to this work

ABSTRACT

Ricci curvature is a method for measuring the robustness of networks. In this work, we use Ricci curvature to measure robustness of brain networks affected by autism spectrum disorder (ASD). Subjects with ASD are given a stem cell infusion and are imaged with diffusion MRI before and after the infusion. By using Ricci curvature to measure changes in robustness, we quantify both local and global changes in the brain networks correlated with the infusion. Our results find changes in regions associated with ASD that were not detected via traditional brain network analysis.

Autism spectrum disorder (ASD) is a heterogeneous neurodevelopmental disorder characterized by social communication deficits and the presence of restricted interests and repetitive behaviors [1]. It is well established that individuals with ASD have different patterns of white matter development in their brains as compared to individuals without ASD, and this differential development of white matter has been linked to autism symptom severity [2, 3]. As a potential therapy, stem cells from umbilical cord blood may alleviate neuroinflammation in the brain that has been linked in ASD [4, 5]. The resulting decrease in inflammation may promote reconfiguration of connectivity patterns among brain regions by promoting white matter development [6]. In a previous study, improvements in social functioning and communication abilities following treatment were correlated with increased connectivity in white matter networks that underlie those social and communicative functions [6, 7]. While examining the difference in volume and strength of white matter connectivity between two brain regions from diffusion tensor imaging (DTI) is a useful measure, it does not account for all the connections into the brain region or pairs of brain regions or consider the robustness of such brain networks and connections. To fully take advantage of the network imaged by DTI, we need a measure that reflects a specific brain region's relationship with every other region in the brain and that quantifies the robustness of such broad network connections.

One solution to measure the robustness of brain regions is Ricci curvature. Broadly speaking, curvature is a 'measure by which a geometrical object deviates from being flat' [8]. Although there are multiple notions of graph curvature [9], this work focuses on the Ricci curvature because it directly measures the robustness of a network and because of its natural physical interpretation and computational efficiency. The link between curvature and robustness is as follow: curvature correlates positively with entropy; entropy correlates positively with robustness [8]. Robustness measures the extent to which a network can withstand perturbations. For brain networks, robustness measures the extent a region in a brain or a connection between two regions in the brain can withstand damage or be affected by a disease or a treatment.

The formal connection of curvature to robustness arises from several sources, including systems and control theory. Feedback tends to make a given system less sensitive, i.e., more robust, to parameter variations and external disturbances [10]. For a weighted graph derived from DTI, feedback is represented by the number of invariant triangles at a given node. Therefore, the

greater the number of triangles [11], the higher the curvature value.

The curvature between two brain regions is computed by using a distance derived from the theory of optimal transport, and gives a novel measure of connectivity based on both local and global network geometry [12]. Thus, the curvature between two brain regions not only considers the strength of connection between those two brain regions, but also the rest of the brain. This measurement takes into account the context of a brain region pair and serves as a useful lens through which to analyze the robustness of the brain.

To measure the efficacy of cord blood infusions for improving social functions of children with ASD, 19 participants were imaged via DTI and participated in a series of behavioral exams before and after the treatment. The brain regions for each subject were delineated and defined as nodes of a network, while edges described structural connectivity between them. Ricci curvature was calculated for the edges as well as for the nodes by taking the weighted average of the curvature for all the neighboring edges.

The results highlight regions which have been previously indicated in ASD, but were not evident when constrained to the differences in white matter connectivity between pairs of individual brain regions. Furthermore, these differences were not found when using traditional brain network analysis techniques, as listed in [13]. First, using Ricci curvature, we see a relationship between clinical improvement and altered robustness in white matter pathways that are implicated in the social and communication abilities that improved following treatment. For example, we demonstrate a relationship between clinical improvement and increased curvature (robustness) in a white matter pathway connecting the left pars triangularis to the left temporal pole, which has been implicated in language abilities (Figure 1). Additionally, we found a path between the right paracentral gyrus (i.e. somatosensory cortex) and the amygdala, which was not identified using conventional DTI analyses [7] as the path from the paracentral gyrus to the amygdala first passes through the sensory thalamus [14]. In addition to identifying novel white matter pathways, the curvature analysis also identified two regions for which the robustness of their connections with the rest of brain increased in relation to clinical outcomes: the left caudal middle frontal (i.e., dorsolateral prefrontal cortex, dlPFC) and the left insula. The finding that changes in insular robustness correlated with clinical outcomes is particularly exciting given a wealth of data supporting the insula as a key structural and functional brain hub across health and disease [15]. With respect to the dlPFC, previous research has implicated the dlPFC as a putative important region in the pathophysiology of ASD [16]. These results support scalar curvature (robustness) as a novel method for the identification of patterns of white matter connectivity that are not identified by common measures of DTI, but yet are related to treatment outcomes in ASD.

These results must be considered in light of some limitations. Because this was an open label trial, it is not possible to determine whether the clinical and curvature changes were a result of normal trajectories of improvement and development or whether they were a consequence of the treatment itself. However, the current data provides targets for exploring brain-related changes in future placebo-controlled double-blind trials, which are currently taking place.

Despite significant progress in understanding the underlying neurobiology of ASD, there are still few reliable and objective measures of change in social and communication function ASD and their relationship with underlying brain structures. We show that curvature identifies changes in robustness in brain regions that may be attributed to the changes in brain connectivity caused by the cord blood infusion. Thus, this study lays the foundation for a new approach to assess both the robustness of a specific brain region and brain region pairs.

Methods

Study Design and Sample

The current study is a secondary data analysis of DTI data collected as part of a phase 1 open-label trial of a single intravenous infusion of autologous umbilical cord blood in 25 children with ASD who were between 24-72 months of age at baseline. The methods of this trial and the accompanying DTI analyses have been described in detail elsewhere [6, 7, 17, 18]. Children with a confirmed diagnosis of ASD and a banked autologous umbilical cord blood unit of adequate size and quality participated in the trial. Nineteen participants provided high quality, artifact-free data for the DTI at both baseline and 6-month visits (17 males and 2 females). All caregivers/legal guardians of participants gave written, informed consent, and the study protocol was approved by the Duke University Health System Institutional Review Board. Methods were carried out in accordance with institutional, state, and federal guidelines and regulation. All methods and the trial were approved by the Duke Hospital Institutional Review Board and conducted under IND #15949. The ClinicalTrials.gov Identifier is NCT02176317.

Clinical Measures

Social abilities were measured with the Vineland Adaptive Behavior Scales-II Socialization Subscale (VABS-SS) [19]. The change in the VABS-SS (6 month-baseline) was used to measure change in social behavior. The Expressive One Word Picture Vocabulary Test 4 (EOW) is a clinician-administered assessment which measures an individual's ability to match a spoken word with an image of an object, action, or concept [20]. The change in the raw score (6 month-baseline) was used to measure change in expressive language. Finally, overall clinical severity and improvement was measured with the Clinical Global Impression Severity (CGI-S) and Improvement (CGI-I) scales.

Magnetic Resonance Imaging Acquisition and Analysis

MRI scanning was conducted on a 3.0 T GE MR750 whole-body 60cm bore MRI scanner (GE Healthcare, Waukesha, WI). Participants were sedated to reduce motion artifacts in the MRI. Diffusion weighted images were acquired using a 25-direction gradient encoding scheme at $b = 1000s/mm^2$ with three non-diffusion-weighted images, an average (std) echo time (TE) of $85ms$ ($2ms$), and a repetition time (TR) of $12,000ms$. An isotropic resolution of $2mm^3$ was achieved using a 96×96 acquisition matrix in a field of view (FOV) of $192 \times 192mm^2$ at a $2mm$ slice thickness. T1-weighted images were obtained with an inversion-prepared 3D fast spoiled-gradient-recalled (FSPGR) pulse sequence with a TE of $2.7ms$, an inversion time (TI) of $450ms$, a TR of $7.2ms$, and a flip angle of $12°$, at a $1mm^3$ isotropic resolution.

Connectome Analysis Pipeline

The full connectome analysis pipeline is described in detail elsewhere [7]. Briefly, each participant's T1 image and the first non-diffusion weighted image (b0) of the DTI acquisition were skull-stripped using the FSL brain extraction tool [21, 22]. The T1 image was registered to the b0 image with an affine registration created using FSL FLIRT [23, 24]. Region of interest (ROI) parcellation was performed by warping the dilated UNC Pediatric Brain atlas (available publicly at http://www.nitrc.org/projects/unc_brain_atlas/) into each participant's T1 in diffusion image space via the Advanced Normalization Tools (ANTs) toolkit [25, 26]. A total of 83 regions were defined for each participant, 41 gray matter regions in each hemisphere, and a single region encompassing the brainstem. FMRIB's Automated Segmentation Tool (FAST) was used to calculate whole brain white matter volume for each participant at both baseline and 6 month visits [27]. Following this, a standardized pipeline for deterministic tractography based on the Connectome Mapper (CMP) was used to analyze participant data at both baseline and 6 month visits (http://www.cmtk.org) [28, 29]. The parcellated gray matter ROIs included in this analysis are defined as nodes. Edges are defined as the volume of voxels containing valid streamlines that originate and terminate within a pair of nodes. For each participant, edge volumes were calculated and normalized by whole-brain white matter volume at both baseline and 6-month visits.

Curvature Analysis

In this section, we outline how we compute the Ricci curvature on discrete metric measure spaces including weighted graphs. The motivation of the Olliver-Ricci [12] definition of curvature on a weighted graph is based on the following characterization of Ricci curvature from Riemannian geometry [30]. For X a Riemannian manifold, consider two very close points $x, y \in X$ and two corresponding small geodesic balls. Positive curvature is reflected in the fact that the distance between two balls is less than the distance between their centers. Similar considerations apply to negative and zero curvature.

For this work, the brain network is represented as an undirected and positively weighted graph, $G = (V, E)$, where V is the set of n vertices (nodes) in the network and E is the set of all edges (links) connecting them with weights $\{w\}$. Consider the graph metric $d : V \times V \to \mathbb{R}^+$ on the set of vertices V where $d(x, y)$ is the number of edges in the shortest path connecting x and y. (d may be any intrinsic metric defined on V.) We let denote $w_{xy} > 0$ denoting the weight of the edge between node x and y. (If there is no edge, then $w_{xy} = 0$.) For any two distinct points $x, y \in V$, the Ollivier-Ricci (OR) curvature is defined as

$$k(x, y) := 1 - \frac{W_1(\mu_x, \mu_y)}{d(x, y)},$$

where W_1 denotes the Earth Mover's Distance (Wasserstein 1-metric). We define the weighted degree at node x, d_x as

$$d_x := \sum_z w_{xz}, \text{ the sum is taken over all nodes } z \text{ adjacent to } x,$$

and we define the probability measure at x, μ_x as

$$\mu_x(y) := \frac{w_{xy}}{d_x}, \ y \in V.$$

The *scalar curvature* at a given node *x* (the contraction of Ricci curvature) is defined as

$$\hat{S}_{OR}(x) := \sum_y \mu_x(y) k(x,y),$$

where the sum is taken over all neighbors of *x*. Given the discussed, positive correlation of robustness and curvature, in our work, we propose to use curvature as a proxy for robustness. Various advantages of using Ricci curvature in this framework are described in more detail in [8].

Acknowledgements

This work would not have been possible without the participation of the participants and their families. We thank the Marcus Foundation (NCT02176317), PerkinElmer, Inc., and the Stylli Translational Neuroscience Award for their financial support for this study, the children who participated and their families, and the following staff members: Jennifer Baker, Colleen McLaughlin, Rebecca Durham, Barbara Waters-Pick, Todd Calnan, Crystal Chiang, Kendyl Cole, Michelle Perry, Mallory Harris, Jennifer Newman, Katherine S. Davlantis, Elizabeth Paisley, Charlotte Stoute, and Elizabeth Sturdivant. This work was supported by the following: National Science Foundation (NSF), United States Office of Naval Research (ONR), United States Army Research Office (ARO), United States Air Force Office of Research (AFOSR), National Institute of Aging (NIA), Breast Cancer Research Foundation (BCRF), and the National Geospatial-Intelligence Agency (NGA). Gifts to G.S. from Google, Microsoft, and Amazon are also acknowledged.

Author Contributions

AKS, KC, SN, AT, GS, and GD designed the study and wrote the paper. AS is in charge of the MRI data collection. JK leads the blood cord effort. JK and GD designed the blood cord intervention in ASD. AKS and SN implemented the algorithms. AKS, KC, and GD analyzed the results.

Author Declarations

G.D. is on the Scientific Advisory Board and receives research funding from Janssen Research and Development, LLC, is a consultant for Roche Pharmaceuticals, LabCorps, Inc., Apple, Gerson Lehrman Group, Guidepoint, and Akili, Inc., received research funding from PerkinElmer and Janssen, and receives royalties from Guilford Press, Springer and University of Oxford Press. J.K. is Director of the Carolinas Cord Blood Bank and Medical Director of Cord:Use Cord Blood Bank. G.S. consults for Apple Inc., and Volvo Cars, and is on the Board of Directors of SIS.

References

1. Association, A. P. *Diagnostic and statistical manual of mental disorders (5th ed.)* (American Psychiatric Association, Washington, DC, 2013), 5th edn.

2. Solso, S. *et al.* Diffusion tensor imaging provides evidence of possible axonal overconnectivity in frontal lobes in autism spectrum sisorder toddlers. *Biol. Psychiatry* **79**, 676–684, DOI: 10.1016/j.biopsych.2015.06.029 (2016).

3. Wolff, J. J. *et al.* Neural circuitry at age 6 months associated with later repetitive behavior and sensory responsiveness in autism. *Mol. Autism* **8**, 8, DOI: 10.1186/s13229-017-0126-z (2017).

4. Suzuki, K. *et al.* Microglial activation in young adults with autism spectrum disorder. *Arch. Gen. Psychiatry* **70**, 49–58, DOI: 10.1001/jamapsychiatry.2013.272 (2013).

5. McAllister, A. K. Immune contributions to cause and effect in autism spectrum disorder. *Biol. Psychiatry* **81**, 380–382, DOI: 10.1016/j.biopsych.2016.12.024 (2017).

6. Dawson, G. *et al.* Autologous cord blood infusions are safe and feasible in young children with autism spectrum disorder: Results of a single-center phase I open-label trial. *Stem Cells Transl. Medicine* **6**, 1332–1339, DOI: 10.1002/sctm.16-0474 (2017).

7. Carpenter, K. L. H. *et al.* White matter tract changes associated with clinical improvement in an open-label trial assessing autologous umbilical cord blood for treatment of young children with autism. *Stem Cells Transl. Medicine* 1–10, DOI: 10.1002/sctm.18-0251 (2019).

8. Sandhu, R. S. *et al.* Graph curvature for differentiating cancer networks. *Sci. Reports* **5**, 12323 (2015).

9. Pouryahya, M., Mathews, J. & Tannenbaum, A. Comparing three notions of discrete ricci curvature on biological networks. *arXiv preprint arXiv:1712.02943* (2017).

10. Doyle, J. C., Francis, B. A. & Tannenbaum, A. R. *Feedback control theory* (Courier Corporation, 2013).

11. Bauer, F., Jost, J. & Liu, S. Ollivier-Ricci curvature and the spectrum of the normalized graph Laplace operator. *Math. Res. Lett.* **19**, 1185–1205 (2012).

12. Ollivier, Y. Ricci curvature of Markov chains on metric spaces. *J. Funct. Analysis* **256**, 810–864 (2009).

13. Rubinov, M. & Sporns, O. Complex network measures of brain connectivity: Uses and interpretations. *NeuroImage* **52**, 1059–1069, DOI: 10.1016/j.neuroimage.2009.10.003 (2010).

14. Abivardi, A. & Bach, D. R. Deconstructing white matter connectivity of human amygdala nuclei with thalamus and cortex subdivisions in vivo. *Hum. brain mapping* **38**, 3927–3940 (2017).

15. van den Heuvel, M. P. & Sporns, O. Network hubs in the human brain. *Trends Cogn. Sci.* **17**, 683–696, DOI: 10.1016/j.tics.2013.09.012 (2013).

16. Qian, L. *et al.* Alterations in hub organization in the white matter structural network in toddlers with autism spectrum disorder: A 2-year follow-up study. *Autism Res.* **11**, 1218–1228, DOI: 10.1002/aur.1983 (2018).

17. Murias, M. *et al.* Validation of eye-tracking measures of social attention as a potential biomarker for autism clinical trials. *Autism Res.* **11**, 166–174 (2018).

18. Murias, M. *et al.* Electrophysiological biomarkers predict clinical improvement in an open-label trial assessing efficacy of autologous umbilical cord blood for treatment of autism. *Stem Cells Transl. Medicine* DOI: 10.1002/sctm.18-0090 (2018).

19. Sparrow, S., Cicchetti, D. & Balla, D. *Vineland adaptive behavior scales (Vineland II): caregiver/caregiver rating form* (NCS Pearson Inc., Minneapolis, MN, 2005), 2 edn.

20. Martin, N. A. & Brownell, R. *Expressive one-word picture vocabulary test (EOWPVT-4)* (Academic Therapy Publications Inc., Novato, CA, 2011), 4 edn.

21. Smith, S. M. Fast robust automated brain extraction. *Hum. Brain Mapp.* **17**, 143–155, DOI: 10.1002/hbm.10062 (2002). NIHMS150003.

22. Jenkinson, M., Pechaud, M. & Smith, S. BET2: MR-based estimation of brain, skull and scalp surfaces. In *Eleventh annual meeting of the organization for human brain mapping*, vol. 17, 167 (Toronto., 2005).

23. Jenkinson, M. & Smith, S. A global optimisation method for robust affine registration of brain images. *Med. Image Analysis* **5**, 143–156, DOI: 10.1016/S1361-8415(01)00036-6 (2001).

24. Jenkinson, M., Bannister, P., Brady, M. & Smith, S. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage* **17**, 825–841, DOI: 10.1006/nimg.2002.1132 (2002).

25. Avants, B. B., Tustison, N. & Song, G. Advanced normalization tools (ANTS). *Insight j* **2**, 1–35 (2009).

26. Avants, B. B. *et al.* A reproducible evaluation of ANTs similarity metric performance in brain image registration. *Neuroimage* **54**, 2033–2044, DOI: 10.1016/j.neuroimage.2010.09.025 (2011).

27. Zhang, Y., Brady, M. & Smith, S. Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Transactions on Med. Imaging* **20**, 45–57, DOI: 10.1109/42.906424 (2001).

28. Gerhard, S. *et al.* The connectome viewer toolkit: an open source framework to manage, analyze, and visualize connectomes. *Front. Neuroinformatics* **5**, 1–15, DOI: 10.3389/fninf.2011.00003 (2011).

29. Englander, Z. A. *et al.* Brain structural connectivity increases concurrent with functional improvement: evidence from diffusion tensor MRI in children with cerebral palsy during therapy. *Neuroimage Clin* **7**, 315–324, DOI: 10.1016/j.nicl.2015.01.002 (2015).

30. Do Carmo, M. P. *Riemannian Geometry* (Birkhauser, 2015).

31. Xia, M., Wang, J. & He, Y. BrainNet Viewer: a network visualization tool for human brain connectomics. *PloS one* **8**, e68910 (2013).

	Region Pairs	VABS	EOW	CGI
a	R. Paracentral - R. Amygdala	-0.53	-0.50	0.55
b	R. Superior Temporal - R. Accubens Area	-0.77	-0.67	0.51
c	R. Caudate - R. Amygdala	-0.55	-0.54	0.58
d	L. Pars Orbitalis - L. Entorhinal	0.56	0.60	-0.49
e	L. Pars Triangularis - L. Temporal Pole	0.66	0.50	-0.46
f	L. Paracentral - L. Inferior Temporal	-0.52	-0.60	0.59
g	L. Pericalcarine - L. Lingual	-0.60	-0.49	0.57
h	L. Temporal Pole - L. Pallidum	0.46	0.51	-0.57

Figure 1. Overview of the study and results. Top) Table showing the brain region pairs and their associated behavioral scores. Bottom) Axial projection of pairs where the change in curvature correlated significantly with behavioral exams. Cross hemisphere brain connections were removed for this analysis. The brain graphics were visualized with the BrainNet Viewer (http://www.nitrc.org/projects/bnv/) [31].

Supplemental Curvature Information

Introduction

Geometric techniques can be used to study the properties of complex networks, including as a method to measure a network's robustness to certain perturbations [1, 2, 3, 4, 5]. Curvature, for this task, is based on the theory of optimal mass transport [6, 7]. First, note that the space of probability densities on a given Riemannian manifold also inherits a natural Riemannian structure [8, 9, 10] for which changes in entropy and Ricci curvature are positively correlated [11, 12]. In conjunction with the Fluctuation Theorem [13], we can conclude that an increase in Ricci curvature is positively correlated with an increase in network robustness, herein expressed as $\Delta \text{Ric} \times \Delta R \geq 0$. This is the key observation that we will exploit below. This observation leads to a notion of curvature on the space of probability distributions on rather general metric measure spaces.

There have been several notions of graph curvature proposed, including those based on the Bochner-Weitzenböck decomposition of the Laplace operator [14] (Forman-Ricci curvature), and those based on Bakry-Émery theory [5, 15]. In this work, we use the formulation of Ollivier [16] because of its direct link to the geometry of optimal mass transport, and therefore its link to entropy and network robustness. There are several other properties of the Ollivier-Ricci curvature that make it attractive for studying the robustness of networks. There is a direct correlation of changes in the rate function from large deviations theory and Ollivier-Ricci curvature [1, 16]. Second, the Ollivier-Ricci curvature is a natural measure of feedback connectivity. This is because the number of triangles in a network (redundant pathways) can be characterized by an explicit lower bound based on Ollivier-Ricci curvature [17]. Similar to control theory, redundancy in feedback is the main characteristic of a robust system architecture. Finally, positive Olliver-Ricci curvature governs the rate of convergence to an invariant distribution on a weighted graph which is modeled as a Markov chain. For these reasons, we chose to use the Ollivier-Ricci curvature for our analysis.

Ricci curvature and entropy

We briefly sketch the connection of curvature and entropy and curvature on a Riemannian manifold. Accordingly, let X denote a complete connected Riemannian manifold equipped with metric d. Ricci curvature provides a way of measuring the degree to which the geometry determined by a given Riemannian metric differs from that of ordinary Euclidean space [18]. The Ricci curvature tensor is obtained as the trace of the sectional curvature [18], and a lower bound on Ricci curvature provides an estimate of the tendency of geodesics to converge or diverge. Through the work of Lott-Sturm-Villani [11, 12], optimal transport connects the bounds on Ricci curvature with entropy, as sketched out below.

First, let $\mathscr{P}(X)$ denote the space of probability densities with finite second moments on X. One can show that $\mathscr{P}(X)$ also has a Riemannian structure [8, 10] via the Wasserstein metric W_2 from optimal mass transport theory (OMT). Next, for a probability measure $\mu \in \mathscr{P}(X)$, the *Boltzmann entropy* is defined as

$$H(\mu) = -\int_X \mu \log \mu \, \mathrm{dvol}.$$

Lott, Sturm, and Villani [11, 12] found a very interesting connection between Ricci curvature and the Boltzmann entropy via OMT. Namely, the Ricci curvature $\text{Ric} \geq k$ if and only if the entropy functional is displacement k-concave along the 2-Wasserstein geodesics, that is, for $\mu_0, \mu_1 \in \mathscr{P}(X)$ we have

$$H(\mu_t) \geq (1-t)H(\mu_0) + tH(\mu_1) + k\frac{t(1-t)}{2}W_2(\mu_0, \mu_1)^2 \quad \forall t \in [0,1], \tag{1}$$

where $(\mu_t)_{0 \leq t \leq 1}$ is a 2-Wasserstein geodesic between μ_0 and μ_1. This inequality indicates the positive correlation between changes in entropy and changes in curvature, which we denote as

$$\Delta H \times \Delta \text{Ric} \geq 0.$$

Entropy and robustness

As previously mentioned, we are interested in the connection between robustness and the network entropy. Informally, functional robustness is defined as the ability of a system to adapt to random perturbations in its environment. This has been exploited in several works [1, 2, 3, 4] in order to employ changes in curvature as a proxy for robustness. This has been extensively explored in the aforementioned papers, and so we will only briefly describe the necessary results.

The starting point is the paper [13], in which the authors characterize robustness via the fluctuation decay rate after random perturbations. Denote by $p_\varepsilon(t)$, the probability that the deviation of the sample mean is more than ε at time t from the original value. Following large deviations theory [19], the *fluctuation decay rate* is defined as

$$R := \lim_{t \to \infty, \varepsilon \to 0} \left(-\frac{1}{t} \log p_\varepsilon(t) \right).$$

The larger the value of R, the faster the systems converges to a stationary state. The Fluctuation Theorem [13] states that the changes in entropy and robustness are positively correlated:

$$\Delta H \times \Delta R \geq 0. \tag{2}$$

Combining (1) and (2) give a relationship which indicates the correlation of changes in curvature and robustness:

$$\Delta R \times \Delta \text{Ric} \geq 0. \tag{3}$$

Thus changes in curvature serve as a measure of changes in system robustness. Positive changes indicate increase in robustness, while negative changes indicate the increase in system fragility.

Wasserstein distance on discrete spaces

We will not need a general definition of the Wasserstein distance in defining the curvature, and so we will suffice with a definition that applies to a weighted connected undirected graph $G = (V, E)$ with n nodes (V) and m edges (E). See [6, 7, 20] and the references therein for the details.

Given two probability densities $\rho^0, \rho^1 \in \mathbb{R}^n$ on the graph, the Kantorovich formulation of the optimal transport problem [6, 7] seeks a joint distribution $\rho \in \mathbb{R}^{n \times n}$ with marginals ρ^0 and ρ^1 minimizing the total cost $\sum c_{ij} \rho_{ij}$:

$$W_C(\rho^0, \rho^1) = \min_\rho \left\{ \sum_{i,j=1}^n c_{ij} \rho_{ij} \mid \sum_k \rho_{ik} = \rho_i^0, \sum_k \rho_{kj} = \rho_j^1, \forall i, j \right\}. \tag{4}$$

Here, c_{ij} is the cost of moving unit mass from node i to node j. If c_{ij} is defined via some intrinsic distance on G (e.g., the hop metric), the minimum of (4) defines a metric W_1 (the Earth Mover's Distance) on the space probability densities on G.

An alternative formulation may be formulated via the fluxes $u \in \mathbb{R}^m$ on the edges. Letting $D \in \mathbb{R}^{n \times m}$ denote the oriented incidence matrix of $\mathscr{G}$, we have

$$W_1(\rho^0, \rho^1) = \min_u \left\{ \sum_{i=1}^m |u_i| \mid \rho^0 - \rho^1 - Du = 0 \right\}.$$

The incidence matrix $D = [d_{ik}] \in \mathbb{R}^{n \times m}$ is defined by associating an orientation to each edge $e_k = (i,j) = (j,i)$ of the graph: one of the nodes i, j is defined to be the head and the other the tail, and then we set $d_{ik} = +1(-1)$ if i is the head (tail) of e_k and 0 otherwise. Compared to the Kantorovich formulation which has n^2 variables, the above formulation has only m variables. It may greatly reduce the computational load when the graph G is sparse, i.e., $m << n^2$ and is often the case in real data.

Olliver-Ricci curvature

In this section, we define the Ricci curvature on discrete networks. Specifically, we assume that our network is presented by an undirected and positively weighted graph, $G = (V, E)$, where V is the set of n vertices (nodes) in the network and E is the set of all edges connecting them, with $w_{xy} > 0$ denoting the weight of the edge between node x and y. (If there is no edge connecting the two nodes, then $w_{xy} = 0$.) We should note that in our treatment above we connected curvature and entropy via W_2, and now we will be using W_1. This technical point is discussed and justified in detail in [3].

Let $d_x := \sum_z w_{xz}$, where the sum is taken over all nodes z in a neighborhood of x. At each node, we define the probability measure

$$\mu_x(y) := \frac{w_{xy}}{d_x}.$$

Let $d : V \times V \to \mathbb{R}^+$ denote an intrinsic metric on G, e.g., the hop distance. Then for any two distinct points $x, y \in V$, the *Ollivier-Ricci (OR) curvature* [16] is defined as

$$k(x, y) := 1 - \frac{W_1(\mu_x, \mu_y)}{d(x, y)}.$$

Using this edge based notion of curvature, we can define the *scalar curvature* at a given node by

$$\hat{S}_{OR}(x) := \sum_y \mu_x(y) k(x, y), \tag{5}$$

where the sum is taken over all neighbors of x.

Given our above arguments, the positive correlation of robustness and curvature, we use curvature as a proxy for robustness in our study of autism. Curvature can be defined either *nodally* (scalar) or relative to *edges* (Ricci). Various advantages of using Ricci curvature in this framework are described in more detail in [1].

References

1. Sandhu, R. S. *et al.* Graph curvature for differentiating cancer networks. *Sci. Reports* **5**, 12323 (2015).

2. Sandhu, R. S., Georgiou, T. T. & Tannenbaum, A. R. Ricci curvature: An economic indicator for market fragility and systemic risk. *Sci. advances* **2**, e1501495 (2016).

3. Farooq, H., Chen, Y., Georgiou, T. T., Tannenbaum, A. R. & Lenglet, C. Network curvature as a hallmark of brain structural connectivity. *bioRxiv* 162875 (2017).

4. Pouryahya, M. *et al.* Bakry-Émery Ricci curvature on weighted graphs with applications to biological networks. In *Int. Symp. on Math. Theory of Net. and Sys*, vol. 22 (2016).

5. Bakry, D. & Emery, M. Diffusions hypercontractives. *Lect. Notes Math. Seminaire de Probab. XIX 1983/84* 177–206, DOI: 10.1007/bfb0075847 (1985).

6. Rachev, S. T. & Rüschendorf, L. *Mass Transportation Problems: Volume I: Theory*, vol. 1 (Springer Science & Business Media, 1998).

7. Villani, C. *Topics in Optimal Transportation*. 58 (American Mathematical Soc., 2003).

8. Benamou, J.-D. & Brenier, Y. A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem. *Numer. Math.* **84**, 375–393 (2000).

9. Jordan, R., Kinderlehrer, D. & Otto, F. The variational formulation of the Fokker–Planck equation. *SIAM journal on mathematical analysis* **29**, 1–17 (1998).

10. Otto, F. The geometry of dissipative evolution equations: the porous medium equation. *Commun. Partial. Differ. Equations* (2001).

11. Lott, J. & Villani, C. Ricci curvature for metric-measure spaces via optimal transport. *Annals Math.* 903–991 (2009).

12. Sturm, K.-T. & Others. On the geometry of metric measure spaces. *Acta mathematica* **196**, 65–131 (2006).

13. Demetrius, L. & Manke, T. Robustness and network evolution—an entropic principle. *Phys. A: Stat. Mech. its Appl.* **346**, 682–696, DOI: 10.1016/j.physa.2004.07.011 (2005).

14. Forman, R. Bochners method for cell complexes and combinatorial ricci curvature. *Discret. Comput. Geom.* **29**, 323–374, DOI: 10.1007/s00454-002-0743-x (2003).

15. Klartag, B., Kozma, G., Ralli, P. & Tetali, P. Discrete curvature and abelian groups. *Can. J. Math.* **68**, 655–674, DOI: 10.4153/cjm-2015-046-8 (2016).

16. Ollivier, Y. Ricci curvature of metric spaces. *Comptes Rendus Math.* **345**, 643–646, DOI: 10.1016/j.crma.2007.10.041 (2007).

17. Bauer, F., Jost, J. & Liu, S. Ollivier-Ricci curvature and the spectrum of the normalized graph Laplace operator. *Math. Res. Lett.* **19**, 1185–1205 (2012).

18. Do Carmo, M. P. *Riemannian Geometry* (Birkhauser, 2015).

19. Varadhan, S. R. S. *Large Deviations and Applications* (Society for Industriel and Applied Mathematics, 1984).

20. Cedric, V. *Optimal Transport: Old and New* (Springer, 2009).