Culex saltanensis and *Culex interfor* (Diptera: Culicidae) are susceptible and competent to transmit St. Louis encephalitis virus (Flavivirus: Flaviviridae) in central Argentina

Mauricio D. Beranek[1,2], Agustín I. Quaglia[2], Giovana C. Peralta[3], Fernando S. Flores [2,4], Marina Stein[1], Luis A. Diaz [2,5], Walter R. Almirón[5] & Marta S. Contigiani[2]

1. Área Entomología, Instituto de Medicina Regional, CONICET, Universidad Nacional del Nordeste, Resistencia, Chaco, Argentina.
2. Laboratorio de Arbovirus y Arenavirus, Instituto de Virología "Dr. *J. M. Vanella*", CONICET, Facultad de Medicina, Universidad Nacional de Córdoba, Córdoba, Argentina.
3. Instituto de Diversidad y Ecología Animal, CONICET, Universidad Nacional de Córdoba, Córdoba, Argentina.
4. Instituto Nacional de Tecnología Agropecuaria, CONICET, Rafaela, Santa Fe, Argentina.
5. Instituto de Investigaciones Biológicas y Tecnológicas, CONICET, Universidad Nacional de Córdoba, Córdoba, Argentina.

ABSTRACT

Infectious diseases caused by mosquito-borne viruses constitute health and economic problems worldwide. St. Louis encephalitis virus (SLEV) is endemic and autochthonous in the American continent. *Culex pipiens quinquefasciatus* is the primary urban vector of SLEV; however, *Culex interfor* and *Culex saltanensis* have also been found naturally infected with the virus, suggesting their potential role as vectors.

OBJECTIVE The aim of this study was to determine the vector competence of *Cx. interfor* and *Cx. saltanensis* for SLEV from central Argentina in comparison to *Cx. p. quinquefasciatus*.

METHODS Adult female mosquitoes of the three *Culex* species were orally infected by feeding on viremic chicks that had been inoculated with SLEV. Then, abdomens, legs and saliva blood-fed mosquitoes were analyzed by viral plaque assay and the presence of cytopathic effect on the cell culture monolayer.

RESULTS Mosquitoes were permissive to orally acquired infections, to virus dissemination, and transmission of SLEV in the saliva. *Cx. saltanensis* and *Cx. interfor* are potential vectors of SLEV.

CONCLUSIONS Our results demonstrate that in Argentina both *Cx. saltanensis* and *Cx. interfor* are susceptible to SLEV and competent for its transmission. Moreover they are abundant during SLEV epidemic period in urban area, positive for this virus in nature, and found to feed on natural hosts.

Keywords: Arbovirus, Argentina, Culicidae, *Flavivirus*, Infectious diseases, Vector competence.

Introduction

Infectious diseases caused by vector-borne pathogens constitute health and economic problems worldwide [1]. Mosquitoes are an important group of arthropod vectors. Due to the hematophagous habit of females, many mosquito species are vectors of infectious agents, including viruses (arthropod-borne viruses; 'arboviruses') [2]. Arbovirus is maintained by biologic transmission among vectors and hosts. Sometimes this biological transmission is specific and includes few vector and host species such as Chikungunya (CHIKV), dengue (DENV), urban yellow fever (YFV) and Zika viruses (ZIKV). However, most of the arboviruses are generalist and they use many vectors and hosts species such as St. Louis encephalitis virus (SLEV) and West Nile virus (WNV) [3]. The emergence and reemergence of diseases caused by arbovirus are a global phenomenon, in particular, those caused by CHIKV, DENV, WNV and ZIKV [1].

SLEV is an endemic neurothropic flavivirus in temperate and subtropical areas of the New World. This virus is maintained between multiple avian hosts and *Culex* vectors, although incidental infections are possible in humans and other mammals, which are typically dead-end hosts [4]. In 2002, SLEV reemerged in the central area of Argentina and southern Brazil causing neurological diseases in humans. In 2005, the first outbreak occurred in Córdoba City with 47 confirmed cases and nine fatalities. After the 2005 outbreak, additional SLEV outbreaks in Argentina occurred in Parana (2006), Buenos Aires (2010), and San Juan (2011) [5]. Factors that promoted this emergence in Argentina include the introduction of a more virulent SLEV strain into a highly susceptible avian hosts community along with possible land use changes (urbanization, agriculture) [5]. Phylogenetic analyses indicate that the emerging SLEV in US (2015) is related to the epidemic strains isolated during a human encephalitis outbreak in Argentina (2005), suggesting introduction from South America [5].

SLEV, as similar to other arboviruses transmitted by mosquitoes, is dependent upon a complex interaction between the virus and vector [2, 6]. The virus in the ingested bloodmeal has to infect and replicate in the epithelial cells of the midgut (midgut infection barrier-MIB). The virus must then successfully escape from the midgut (midgut escape barrier-MEB) and infect the salivary glands (gland infection barrier-SGIB), followed by release of the virus into the salivary ducts for transmission orally to vertebrates. Salivary gland infection and escape barriers (salivary gland escape barrier-SGEB) determine if the virus can replicate and shed into the mosquito's saliva for final transmission to the vertebrate host during [2, 7]. Mosquito vector capacity [susceptibility, extrinsic incubation period (EIP), transmission proportions], longevity, and host bloodmeal preferences are included among intrinsic factors. Each of these factors is affected by extrinsic factors such as larval density and nutrition, temperature, rainfall, avian host availability and avian host immunity [2, 7].

In Argentina, the Eared Dove (*Zenaida auriculata*) and Picui Ground-Dove (*Columbina picui*) are natural hosts of SLEV, while *Culex pipiens quinquefasciatus* is an efficient vector for the virus [3, 8]. For example, SLEV was detected and isolated in *Cx. p. quinquefasciatus* mosquitoes during an outbreak of encephalitis in humans in Córdoba City [9]. Infected *Cx. p. quinquefasciatus* have been detected during periods without reports of clinical disease symptoms in Santa Fe province between 1978-1983 [10] and Córdoba between 2001-2004 [11]. In addition viral isolations from field collected mosquitoes, and the population abundance throughout the Córdoba City [9], laboratory studies have confirmed horizontal transmission [8, 12], and the infrequent vertical transmission of SLEV in *Cx. p. quinquefasciatus* [13]. The presence of naturally SLEV-infected *Culex interfor* and *Culex saltanensis*, dominant species in urban-vegetated sub-assemblage that frequently feed on competent hosts, suggest they could participate as vectors in the transmission network of SLEV [3, 8, 9, 14]. Based on these evidences we argue that *Cx. saltanensis* and *Cx. interfor* could be playing a role as vectors of SLEV in central Argentina; yet vector competence experiments have not been conducted on these species. Therefore, we evaluated the vector competence of *Cx. interfor* and *Cx. saltanensis* against SLEV from central Argentina compared to the primary urban vector, *Cx. p. quinquefasciatus*.

Methods

Capture and maintenance of mosquito colonies

Egg rafts of *Cx. p. quinquefasciatus*, *Cx. interfor* and *Cx. saltanensis* were collected during 2015 at the Bajo Grande sewage treatment plant (31°24'13"S 64°06'08"W) located in east of Córdoba City. The site is surrounded by aquatic vegetation, reservoirs, low income human settlements and crop lands (vegetables and fruits). Authorization for mosquito field collections was obtained from the Municipality of Córdoba.

Egg rafts were collected with a dipper (350 ml) and transferred to polypropylene bottles with a fine brush and transported to the Instituto de Virología "Dr. *J. M. Vanella*" Facultad de Medicina, Universidad Nacional de Córdoba (FM-UNC). Mosquitoes were maintained 27°C, 70% humidity, and 12:12 h light: dark (L: D) photoperiodic. Adult F1 females from the field collected rafts were used in the vector competence assay (Table 1). Egg rafts were placed in a plastic container with distilled water and hatched larvae were fed with suspension of liver powder (0.25mg/0.5ml distilled water) every 48 h. After identification, larvae were grouped by species, pupae of the same species were placed inside screened cages (26 x 22 cm) until emergence, and adults provided a 10% sugar solution by soaked cotton balls. Identification of adult females and 4th instars were based on morphological keys [15]. Adult males were also collected to confirm the taxonomic identification based on the genitalia morphology [16, 17]. The small number of adult females used in this study, was caused by low feeding success and high mortality, which limited the sample size (Table 1).

Table 1: The egg collections and number females of Cx. p. quinquefasciatus, Cx. interfor and Cx. saltanensis in Córdoba City.

Species	N° of egg rafts per species*	Total N° of females fed on chicks†	N° of engorged females‡	N° of SLEV positive females§
Culex p. quinquefasciatus	12	64	39	13
Culex interfor	15	32	25	14
Culex saltanensis	9	18	12	8

*Number of egg rafts collected for each mosquito species, assuming that each raft was from a different female. Number of eggs not determined. Male/female ratio of hatched individuals was ~50%.

† Number of females provided blood meals on viremic chicks.

‡Number of engorged females after feeding on viremic chicks.

§Number of females positives for SLEV.

Viral stock

Adult female mosquitoes were orally infected by feeding on viremic chicks that had been inoculated with SLEV CbaAr-4005 [9]. Viral stock was obtained from collections stored at the Instituto de Virología "Dr. J. M. Vanella" (FM-UNC). Viral stock was prepared from brain of an infected Swiss albino suckling mice homogenized in 10% P/V solution in Eagle's minimal essential medium (MEM) (Gibco, Ireland), supplemented with 10% fetal bovine serum (FBS) (Natocor, Argentina) and 1% of gentamycin (Klonal, Argentina).

Viral titration was carried out by viral plaque assay in VERO cell monolayer (African green monkey kidney, Cercophitecus aethiops). We inoculated 0.1 ml of the samples onto VERO cell monolayer on a 12-well plate and incubated the plates for 60 min at 37°C with 5% CO_2 and a humid atmosphere to favor the adsorption of the virus to the cell. After incubation 0.5 ml of MEM 2x was mixed with 1% methylcellulose and added to plate followed by incubation at 37°C for 7 days, until the formation of lytic plaques. The plates were then fixed with 10% formalin solution for 2 h and stained with crystal violet for the observation and plaque forming units (PFU) counts. Viral concentrations were then expressed as $\log_{10}$ PFU per milliliter (PFU/ml).

Vector competence assay

Twenty-four hour-old chicks (Gallus gallus) were inoculated intraperitoneally with 0.1 ml of a viral suspension containing approximately 400 PFU of SLEV. In accordance with viremia kinetic [8], 48 h post-inoculation starved female mosquitoes were fed on chicks (Table 1). Prior to and following blood feeding by mosquitoes, 0.1 ml of blood was taken from the chick jugular vein to determine the viremia titer (potential viral load ingested by mosquitoes). The chick blood was diluted in 0.45 ml of MEM supplemented with 10% FBS and 1% gentamycin, followed by centrifugation at 4°C for 20 min at 2,300 g before supernatant stored at -80°C. Titers were determined by plaque assay and expressed as the mean viremia load before and after feeding (log10 PFU/ml). Mosquitoes were anesthetized by cold and fully engorged females were maintained in a screened cage at 27oC, H° 70%, 12:12 h L: D photoperiod and provided a 10% sugar solution. After 14 days EIP, the females were aspirated and anesthetized for 2 min at 4oC, placed on a refrigerated plate where the legs and wings were gently removed. Saliva samples were recovered after live females were placed on a flat surface with adhesive tape and the proboscis inserted for 30 min into a capillary tube with 0.001 ml glycerin (Todo Droga, Argentina) [18]. The abdomen was then removed, and saliva, legs and abdomen samples individually stored at-80oC with 1 ml of MEM supplemented with 10% FBS and 1% gentamycin. Legs and abdomen were individually homogenized by agitation with glass beads for 4 min and centrifuged at 11,200 g for 30 min. Subsequently, 0.1 ml of the sample was inoculated with two cellular and viral controls and infective virus particles detected by plaque assay as described above.

SLEV-negative saliva samples from females with positive abdomen samples were re-analyzed to reduce false negatives due to dilution. Thus, 0.2 ml of each sample was inoculated in two cell monolayer plates ('A' and 'B' plates) using the same controls as mentioned above. 'A' plates followed the same culture protocol described above. For 'B' plates, MEM, FBS (2%) and gentamycin (1%) were added, and after 96 h the cytopathic effect (cell detachment, rounding and nonconfluent monolayer) was determined under an inverted microscope. To amplify viral particles not detected in the first passage, monolayer from the 'B' plates were harvested on the fourth day post inoculation, by collecting 0.75 ml from each of the duplicates wells followed by centrifugation at 9,300 g for 30 min (passage 1). Finally, another plate was inoculated (duplicate, same 'B' plates protocol) with 0.2 ml of the samples without cytopathic effect in the 'B' plates. Therefore, PFU/ml was recorded in 'A' plates and the cytopathic effect presence in 'B' plates.

Ethical Guidelines

The protocol used was approved by Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) and Instituto de Virología "Dr. J. M. Vanella" (FM-UNC). The chicks were maintained in cages with a substrate of wood shavings and with a 12:12 h L: D photoperiod, 40-80% humidity and 20-26°C temperature. Food and water were available ad libitum (balanced concentrate feed, Gepsa Feeds, Argentina). The experimental procedures used were to minimize or eliminate pain and distress. Awareness were taken to avoid chicks suffering, as anesthetic protocols were not used. Euthanasia was carried out with neck dislocation by a trained laboratory technician.

Statistics analysis

Abdomens, legs, and saliva were considered positive when they showed at least one PFU or the presence of cytopathic effect on the cell culture monolayer. Infection rates were defined as the numbers of positive mosquito abdomens of the total number of blood feed mosquitoes. Dissemination rates were calculated as the number of mosquitoes with positive legs out of the number of mosquitoes fed analyzed. Transmission rates were determined by the number of mosquitoes with positive saliva of the total number of mosquitoes that blood fed. Confidence intervals (0.95%) for infection, dissemination and transmission rates and graphical presentations were made in the R [19]. Differences in rates for each species Culex spp. were compared by a Fisher exact test, considering statistically significant $\alpha < 0.05$ [20].

Results

The numbers of eggs collected *Cx. p. quinquefasciatus*, *Cx. interfor*, *Cx. saltanensis* and females that fed and were positive for SLEV are shown in Table 1. Mosquitoes were permissive to orally acquire infections, to virus dissemination, and transmission of SLEV in saliva (Figure 1). There was a narrow range of viremia during blood feeding (*Cx. p. quinquefasciatus* = 2.9 $\log_{10}$ PFU/ml, *Cx. interfor* = 3.5 $\log_{10}$ PFU/ml and *Cx. saltanensis* = 3.2 $\log_{10}$ PFU/ml).

Figure 1: Vector competence for SLEV of *Cx. p. quinquefasciatus*, *Cx. interfor* and *Cx. saltanensis*. Infection rate (number of positive mosquito abdomens/total number of mosquitoes fed) in light gray, dissemination rate (number of mosquitoes with positive legs/total number of mosquitoes fed) in gray and transmission rate (number of mosquitoes with positive saliva/total number of mosquitoes fed) in dark gray; with their 0.95% CIs.

Females *Cx. p. quinquefasciatus* were equally susceptible to infection and transmission of SLEV, because were not statistically significant difference between infection (13/39, 33%), dissemination (7/13, 54%) and transmission rates (18%, 7/39) (Table 2).

Table 2: Vector competence of *Culex p. quinquefasciatus*, *Cx. interfor* and *Cx. saltanensis* for SLEV measured as infection, dissemination and transmission.

Species Vector competence	*Culex p. quinquefasciatus*			*Culex interfor*			*Culex saltanensis*		
	N*	Rates† (0.95%CIs)	Viral load‡ (0.95%CIs)	N*	Rates† (0.95%CIs)	Viral load‡ (0.95%CIs)	N*	Rates† (0.95%CIs)	Viral load‡ (0.95%CIs)
Infection	13/39	33 (19.1-50.2)	2.4 (1.3-2.9)	14/25	56 (34.9-75.6)	4.8 (2.4-5.3)	8/12	67 (34.9-90.1)	4.8 (3.8-5.1)
Dissemination	7/39	18 (7.5-33.5)	2.9	10/25	40 (21.1-61.3)	4.8 (1-5.4)	8/12	67 (34.9-90.1)	4.6 (1.6-5.1)
Transmission	7/39	18 (7.5-33.5)	No data	5/25	20 (6.8-40.7)	1.8 (1.1-2.3)	2/12	17 (2.1-48.4)	1.8

*Number of mosquitoes positive for SLEV/total number of mosquitoes assayed

†Infection rate (number of positive mosquito abdomens/total number of mosquitoes fed); dissemination rate (number of mosquitoes with positive legs/total number of mosquitoes fed) and transmission rate (number of mosquitoes with positive saliva/total number of mosquitoes fed).

‡ Average SLEV titers ($\log_{10}$ PFU/ml) in abdomen, legs and saliva of each mosquito species.

Viral loads were evaluated for abdomens 2.4 $\log_{10}$ PFU/ml and legs 2.9 $\log_{10}$ PFU/ml. Because lysis plaques in *Cx. p. quinquefasciatus* were not confluent the SLEV load in saliva was not able to measure (Table 2). For *Cx. interfor*, statistically significant differences were observed across infection (14/25, 56%) and transmission rates (20%, 5/25) (Fisher's exact test, p = 0.0186) (Table 3).

Table 3: Infection, dissemination and transmission rates of *Cx. p. quinquefasciatus*, *Cx. interfor* and *Cx. saltanensis* in Córdoba City.

Culex p. quinquefasciatus	Infection rate[†]	Dissemination rate[†]	Transmission rate[†]
Infection rate		0.1942	0.1942
Dissemination rate	0.1942		1
Transmission rate	0.1942	1	
Culex interfor			
Infection rate		0.3961	**0.0186***
Dissemination rate	0.2165		0.3961
Transmission rate	**0.0186***	0.2165	
Culex saltanensis			
Infection rate		1	**0.0361***
Dissemination rate	1		**0.0361***
Transmission rate	**0.0361***	**0.0361***	

Fisher exact test used to compare infection, dissemination and transmission rates for each species *Culex* spp.

†Infection rate (number of positive mosquito abdomens/total number of mosquitoes fed); dissemination rate (number of mosquitoes with positive legs/total number of mosquitoes fed) and transmission rate (number of mosquitoes with positive saliva/total number of mosquitoes fed).

*Significance was tested at a level of α = 0.05.

Viral loads were evaluated for abdomens 4.8 $\log_{10}$ PFU/ml, legs 4.8 $\log_{10}$ PFU/ml and saliva 1.8 $\log_{10}$ PFU/ml (Table 2). For *Cx. saltanensis,* statistically significant differences were observed between infection (8/12, 67%) and transmission rates (17%, 2/12), and this rate with respect to the dissemination rate (8/12, 67%) (Fisher's exact test, p = 0.0361) (Table 3). Viral loads were evaluated for

abdomens 4.8 $\log_{10}$ PFU/ml, legs 4.6 $\log_{10}$ PFU/ml and saliva 1.8 $\log_{10}$ PFU/ml (Table 2).

Discussion

Culex p. quinquefasciatus, *Cx. interfor* and *Cx. saltanensis* were competent vectors for SLEV based on 1) acquired infections, 2) disseminated virus, and 3) transmission of SLEV in the saliva after feeding on a viremic chick. SLEV was able to cross the midgut barriers showing disseminated infection (positive legs) and was able to cross salivary gland barriers in positive saliva samples. However, the number of tested mosquitoes was low, thus our results are not conclusive.

To be considered a vector, a mosquito species must fulfill several biological and ecological characteristics [21, 22]. The coevolution between pathogen and arthropods determine the vector competence, and thus the ability to acquire, maintain and eventually transmit it [21]. Variation in vector competence has been documented with all of the major disease agents they transmit (i.e. malaria and filarial parasites, and arboviruses) [6, 23, 24]. Although the life cycle of each pathogen is distinct, they all face the common events of being ingested, exposed to the midgut environment, and traversing hemocoel to reach their tissue site of development and/or suitable site for transmission back to a new vertebrate host. Each of these migratory steps presents potential barriers that might be manipulated to interfere with normal pathogen migration and/or development [22]. Along with these barriers, some other factors like digestive enzymes, midgut microbiota, and innate immune responses might be responsible for vector's refractoriness and ineffective horizontal transmission [2]. Understanding the vector competence is crucial for assessing the risks of arbovirus transmission and maintenance in nature.

There is considerable specificity in the vector-arbovirus relationship, and some of this specificity comes from the ability of a particular arbovirus to overcome tissue barriers in the vector to establish a persistent infection. Factors that strongly affect vector competence of a mosquito for a particular arbovirus include MIB, MEB, SGIB, and SGEB [6]. In our findings, among infected mosquitoes, dissemination was achieved in 100% of those individuals tested (8/8) for *Cx. saltanensis*, while are 71% (10/14) of *Cx. interfor* and 54% (7/13) of *Cx. p. quinquefasciatus* demonstrated disseminated infections. These results suggest the possible existence of a midgut barrier to SLEV in *Cx. p. quinquefasciatus*. Kramer et al. were the first to demonstrate that the inability of infected *Cx. tarsalis* mosquitoes to transmit western equine encephalitis virus (WEEV) was associated with a MEB [25]. Also following studies detected existence the midgut barrier for Rift Valley Fever virus in *Cx. pipiens* [26], SLEV and WNV in *Cx. p. quinquefasciatus* respectively [27, 28].

Virus transmission is a critical component of laboratory studies of vector competence and is essential to understanding the epidemiology of arboviruses [18]. Reisen et al. quantified the viral particles of SLEV expectorated in the saliva of *Cx. tarsalis* (1.1-2.2 $\log_{10}$ PFU) [29] and we obtained similar data for *Cx. interfor* (range = 1.1-2.3 $\log_{10}$ PFU) and *Cx. saltanensis* (1.8 $\log_{10}$ PFU). Moreover, all *Cx. p. quinquefasciatus* females with disseminated infections demonstrated SLEV in their saliva (7/7); this rate was only 50% (5/10) in *Cx. interfor* and 25% (2/8) in *Cx. saltanensis*, indicating a potential salivary gland barrier in both *Cx. interfor* and *Cx. saltanensis*. Similar results were obtained for Japanese encephalitis virus in *Cx. p. molesus* [30], WEEV in *Cx. tarsalis* [25] and Venezuelan equine encephalitis virus in *Psorophora cingulata* and *Coquillettidia venezuelensis* [31]. Further work are needed should evaluate and explore the relationship between midgut and salivary gland barriers.

Our results corroborate the findings reported by Diaz et al. on the susceptibility of *Cx. p. quinquefasciatus* for SLEV infection [8]. These authors observed an infection rate of 70% using the same viral strain CbaAr-4005 and feeding viremia level of 5.2 $\log_{10}$ PFU/ml. In our study only one third of the *Cx. p. quinquefasciatus* females became infected. This difference could be related to the lower viremia level that the mosquitoes were exposed to in this assay (2.9 $\log_{10}$ PFU/ml). Mitchell et al. obtained a transmission rate of (90.5%, 19/21) with strain 78V-6507 for *Cx. p. quinquefasciatus* from Santa Fe Province [12]. In our study, the transmission rate was (18%, 7/39). Dissemination and transmission rate formulation in Mitchell and herein are different. Even through dissemination rates are higher with a higher infectious dose (Mitchell, 4.1-4.8 $\log_{10}$ PFU/ml vs our study, 2.9 $\log_{10}$ PFU/ml). The effect of viremia could be affected by the small number of individuals used in this study. Viremia, virus dose, extrinsic incubation temperature, mosquito age, and colony are all important factors influencing the vector competence of *Cx. p. quinquefasciatus* [27, 32]. Future studies to determine the Minimum Infection Threshold and the Extrinsic Incubation Periods are needed.

The maintenance of SLEV in nature is complex and requires the coexistence in time and space of mosquito vectors and avian hosts. The detection of naturally infected mosquitoes does not represent, by itself, a reliable proof of their role as a competent vector. In the case of *Cx. p. quinquefasciatus*, *Cx. interfor* and *Cx. saltanensis*, there is evidence supporting their intervention as vectors in SLEV transmission. *Culex p. quinquefasciatus* is considered the primary vector, because it was abundant and *pools* assayed were positive for SLEV [9-11, 14]. *Culex saltanensis* and *Cx. interfor* participate in the maintenance of SLEV and could assist in the spillover of SLEV to humans. In 2004, prior to the outbreak of encephalitis in Córdoba City, SLEV infected *Cx. interfor* were detected [11]. In 2010, there were small outbreaks of SLEV in provinces, e.g., Buenos Aires, Córdoba and San Juan [33]. Not long thereafter, SLEV infected *Cx. saltanensis* were detected for first time in Córdoba City [14]. *Culex interfor* and *Cx. saltanensis* are mainly ornithophiles, and bloodmeal from Columbiformes and Passeriformes have been also detected, although the pattern of host preference and its drivers have not been established yet [14, 34, 35]. In Argentina, *Z. auriculata* and *C. picui* are amplifier hosts of SLEV and have been recorded in engorged *Cx. saltanensis*, *Cx. interfor* and *Cx. p. quinquefasciatus* sustaining that SLEV maintenance could relied on multiple vectors [3, 36]. However, the transmission load in SLEV episystem could be unequal between the three *Culex*, despite they showed similar transmission rate experimentally (ranged 17-20%). For instance, among other traits, lifespan difference among mosquito species is expected to impact vector capacity as longest lifespan increase the odds of extrinsic incubation completeness and delivering infectious bites [7, 21]. Here, *Cx. p. quinquefasciatus* was less able to survive after a viremic bloodmeal than *Cx. saltanensis* and *Cx. interfor* suggesting that the role of the last species has been neglected. In addition, it has been proposed that *Cx. interfor* could transmit SLEV from birds to mammals and thus fulfill a role of "bridge vector" [14, 34, 37] as *Cx. interfor* was recorded in human baited barley traps [37] and *Cx. interfor* and *Cx. saltanensis* switch between bird feeding profile in spring-summer to bird-mammals in autumn in a rural environment [38]. The local populations of *Culex* spp. increase in abundance with peaks in summer, with are temporal distribution of *Culex* spp. coinciding with the activity peaks of SLEV in human infection [3, 39]. Adult mosquitoes belonging to the species *Cx. saltanensis* and *Cx. interfor* have been found in increasing numbers in Córdoba City, with a higher abundance in urban and periurban areas where vegetation is more robust. This differs from *Cx. p. quinquefasciatus*, which is predominant throughout a vast range of city-type environments [39]. Our results support the hypothesis that SLEV is transmitted by multiple sympatric *Culex* spp., and that both *Cx. saltanensis* and *Cx. interfor* can be considered potential vectors of SLEV. In the United States, this has been observed as well; however, different mosquito species serve as the primary vector transmitting SLEV in different geographical areas. *Culex quinquefasciatus* and *Cx. nigripalpus* are vectors for the virus in Florida, *Cx. tarsalis* in the western and *Cx. pipiens* in the northern United States [4].

SLEV is a multi-host and multi-vector flavivirus in the process of an ongoing reemergence in Argentina. Further studies are required to understand the spatial compartmentalization of these mosquito species in the transmission network of SLEV by performing vector capacity studies. By having insights into its ecoepidemiology, we will have a better understanding of which factors are causing this reemergence and how biological and environmental factors interact with and affect its activity. In addition, knowledge of the potential mosquito species vectors of SLEV will provide information to be used by different public agencies related to human health for the control of vector mosquito populations and improve efficiency in SLEV prevention programs.

Data accessibility

Data are available online: link or DOI of the webpage hosting the data.

Supplementary material

End of the Manuscript

Acknowledgements

We thank Kristin Sloyer for sharing suggestions and improving the English of the manuscript.

Version 6 of this preprint has been peer-reviewed and recommended by Peer Community In Entomology (https://doi.org/10.24072/pci.entomol.100002).

Conflict of interest disclosure

The authors of this preprint declare that they have no financial conflict of interest with the content of this article.

Adrian Diaz is one of the PCI Entomology recommenders.

References

[1] Gould E, Pettersson J, Higgse S *et al*. Emerging arboviruses: Why today? One Health 4 (2017), 1-13. DOI: 10.1016/j.onehlt.2017.06.001.

[2] Agarwal A, Parida M, Dash PK. Impact of transmission cycles and vector competence on global expansion and emergence of arboviruses. Rev Med Virol (2017), e1941. DOI: 10.1002/rmv.1941.

[3] Diaz LA, Flores FS, Quaglia A *et al*. Intertwined arbovirus transmission activity: Reassessing the transmission cycle paradigm. Front Physiol 3 (2013), 493. DOI: 10.3389/fphys.2012.00493.

[4] Reisen WK. Epidemiology of St. Louis encephalitis virus. Adv Virus Res 61 (2003), 139-184. DOI: 10.1016/s0065-3527(03)61004-3.

[5] Diaz A, Coffey LL, Burket-Cadena N *et al*. Reemergence of St. Louis encephalitis virus in the Americas. Emerg Infect Dis 24 (2018), 2150-2157. DOI: 10.3201/eid2412.180372.

[6] Franz AWE, Kantor AM, Passarelli AL *et al*. Tissue barriers to arbovirus infection in mosquitoes. Viruses 7 (2015), 3741-3767. DOI: 10.3390/v7072795.

[7] Kramer LD & Ebel GD. Dynamics of *Flavivirus* infection in mosquitoes. Adv Virus Res 60 (2003), 187-232. DOI: 10.1016/s0065-3527(03)60006-0.

[8] Diaz LA, Flores FS, Beranek M *et al*. Transmission of endemic St Louis encephalitis virus strains by local *Culex quinquefasciatus* populations in Córdoba, Argentina. Trans R Soc Trop Med Hyg 107 (2013), 332-334. DOI: 10.1093/trstmh/trt023.

[9] Diaz LA, Ré V, Almirón W *et al*. Genotype III Saint Louis encephalitis virus outbreak, Argentina, 2005. Emerg Infect Dis 128 (2006), 1752-1754. DOI: 10.3201/eid1211.060486.

[10] Sabattini MS. Historical, epidemiological and ecological aspects of arboviruses in Argentina: Flaviviridae, Bunyaviridae and Rhabdoviridae. An overview of Arbovirology in Brazil and neighbouring countries (ed. by Travassos da Rosa APA, Vasconcelos PFC & Travassos da Rosa JFS). Brasil: Instituto Evandro Chagas (1998), 113-134.

[11] Diaz LA, Albrieu Llinás G, Vázquez A *et al*. Silent circulation of St Louis encephalitis virus prior to an encephalitis outbreak in Córdoba, Argentina (2005). PLoS Negl Trop Dis 6 (2012), 1489.

[12] Mitchell CJ, Monath TP & Sabattini MS. Transmission of St. Louis encephalitis virus from Argentina by mosquitoes of the *Culex pipiens* (Diptera: Culicidae) Complex. J Med Entomol 17 (1980), 282-285. DOI: 10.1093/jmedent/17.3.282.

[13] Flores FS, Diaz LA, Batallán GP *et al*. Vertical transmission of St. Louis encephalitis virus in *Culex quinquefasciatus* (Diptera: Culicidae) in Córdoba, Argentina. Vector Borne Zoonotic Dis 10 (2010), 999-1002. DOI: 10.1089/vbz.2009.0136.

[14] Batallán GP. Binomía de *Culex interfor* Dyar en Córdoba, Argentina. [theses]. Córdoba: Universidad Nacional de Córdoba (2013), 197 p.

[15] Darsie RF. Mosquitoes of Argentina. Part I. Keys for identification of adult females and fourth stages larvae in English and Spanish (Diptera: Culicidae). Mosquito Systematics 1985: 17: 153-253.

[16] Harbach RE, Jakov WL & Peyton EL. Recognition of *Culex bidens* Dyar and *Culex interfor* Dyar (Diptera: Culicidae) as separate species. Mosquito Systematics 18 (1986), 139-144.

[17] Laurito M, Visintin AM & Almirón WR. *Culex saltanensis* morphological redescription of the inmature and adult stage. J Am Mosq Control Assoc 24 (2008), 203-210.

[18] Anderson SL, Richards SL & Smartt CT. A Simple method for determining arbovirus transmission in mosquitoes. J Am Mosq Control Assoc 26 (2010), 108-111. DOI: 10.2987/09-5935.1.

[19] R Core Team. R: A language and environment for statistical computing. [cited 2015]. https://www.R-project.org.

[20] GraphPad Prism [www.graphpad.com/quickcalcs/contingency1.cfm].

[21] Gubler DJ. Vector-borne diseases. Rev sci tech 28 (2009), 583-588. DOI: 10.20506/rst.28.2.1904.

[22] Beerntsen BT, James AA & Christensen BM. Genetics of Mosquito Vector Competence. Microbiology and Molecular Biology Reviews 64 (2000) 115-137. DOI: 10.1128/mmbr.64.1.115-137.2000.

[23] Cohuet A, Harris C, Vincent R *et al*. Evolutionary forces on *Anopheles*: what makes a malaria vector? Trends Parasitol 26 (2010), 130-136. DOI: 10.1016/j.pt.2009.12.001.

[24] Ahid SMM, da Silva Vasconcelos PS & Lourenço-de-Oliveira R. Vector Competence of *Culex quinquefasciatus* Say from Different Regions of Brazil to *Dirofilaria immitis*. Mem Inst Oswaldo Cruz 95 (2000), 769-775.

[25] Kramer LD, Hardy JL, Presser SB *et al*. Dissemination barriers for western equine encephalomyelitis virus in *Culex tarsalis* infected after ingestion of low viral doses. Am J Trop Med Hyg 30 (1981), 190-197. DOI:10.4269/ajtmh.1981.30.190.

[26] Romoser WS, Faran ME, Bailey CL *et al*. An immunocytochemical study of the distribution of Rift Valley fever virus in the mosquito *Culex pipiens*. Am J Trop Med Hyg 46 (1992), 489–501. DOI: 10.4269/ajtmh.1992.46.489.

[27] Richards SL, Lord CC, Pesko K *et al*. Environmental and Biological Factors Influencing *Culex pipiens quinquefasciatus* Say (Diptera: Culicidae) Vector Competence for Saint Louis Encephalitis Virus. Am J Trop Med Hyg 81 (2009), 264–272.

[28] Richards SL, Anderson SL, Lord CC *et al*. Relationships between infection, dissemination, and transmission of West Nile virus RNA in *Culex pipiens quinquefasciatus* (Diptera: Culicidae). J Med Entomol 49 (2012), 132-142. DOI: 10.1603/me10280.

[29] Reisen WK, Chiles RE, Kramer LD *et al*. Method of infection does not alter response of chicks and house finches to western equine encephalomyelitis and St. Louis encephalitis viruses. *J Med Entomol* 37 (2000), 250-258.

[30] Turell MJ, Mores CN, Dohm DJ *et al.* Laboratory transmission of Japanese encephalitis and West Nile viruses by molestus form of *Culex pipiens* (Diptera: Culicidae) collected in Uzbekistan in 2004. J Med Entomol 43 (2006) 296-300, DOI: 10.1603/0022-2585(2006)043[0296:ltojea]2.0.co;2.

[31] Turell MJ, Jones JW, Sardelis MR *et al*. Vector competence of Peruvian mosquitoes (Diptera: Culicidae) for epizootic and enzootic strains of Venezuelan equine encephalomyelitis virus. J Med Entomol 37 (2000), 835-839. DOI: 10.1603/0022-2585-37.6.835.

[32] Richards SL, Mores CN, Lord CC *et al*. Impact of Extrinsic Incubation Temperature and Virus Exposure on Vector Competence of *Culex pipiens quinquefasciatus* Say (Diptera: Culicidae) for West Nile Virus. Vector Borne Zoonotic Dis 7 (2007), 629–636. DOI: 10.1089/vbz.2007.0101.

[33] Vergara Cid C, Spinsanti L, Rivarola M *et al*. Detección de infecciones humanas por *Flavivirus* en la ciudad de Córdoba durante el año 2010. Rev Argent Microbiol 43 (2011), 37.

[34] Beranek MD. Mosquitos del género *Culex* (Diptera: Culicidae) como vectores del virus Saint Louis encephalitis (Flavivirus) en ambientes urbanos de la ciudad de Córdoba. [theses]. Córdoba: Universidad Nacional de Córdoba (2019), 188 p.

[35] Berrón CI. Preferencia de hospedadores aviares en especies de mosquitos Género *Culex* asociadas a la transmisión de Flavivirus (Flaviviridae) en el arco sur de la Laguna Mar Chiquita. [theses]. Córdoba: Universidad Nacional de Córdoba (2014), 155 p.

[36] Diaz LA, Flores FS, Quaglia IA et al. Evaluation of Argentinean birds species as amplifying hosts for St. Louis encephalitis virus (Flavivirus, Flaviviridae). Am J Trop Med Hyg 99 (2018), 216-221. DOI:10.4269/ajtmh.17-0856.

[37] Stein M, Zalazar L, Willener JA *et al*. Culicidae (Diptera) selection of humans, chickens and rabbits in three different environments in the province of Chaco,

Argentina. Mem Inst Oswaldo Cruz 108 (2013), 563-571. DOI: 10.1590/s0074-02762013000500005.

[38] Quaglia AIE. Comunidades de Mosquitos Vectores y Hospedadores Aviares y su Asociación en el Mantenimiento del virus St. Louis encephalitis (Flavivirus). [theses]. Córdoba: Universidad Nacional de Córdoba (2017), 217 p.

[39] Batallán GP, Estallo EL, Flores FS *et al*. St. Louis encephalitis virus mosquito vectors dynamics in three different environments in relation to remotely sensed environmental conditions. Acta Trop 146 (2015), 53-59. DOI: 10.1016/j.actatropica.2015.03.009.

Supplementary material

Table: Female mosquitoes *Cx. interfor*, *Cx. saltanensis* and *Cx. p. quinquefasciatus* used in vector competence assay with the SLEV

Specie	ID	Viremia	Results	Rate	N	PFU/ml	log₁₀PFU/ml
Cx.interfor	1.1	3.5	0	Infection	25	0	
Cx.interfor	1.1	3.5	0	Dissemination	25	0	
Cx.interfor	1.1	3.5	0	Transmission	25	0	
Cx.interfor	1.2	3.5	0	Infection	25	0	
Cx.interfor	1.2	3.5	0	Dissemination	25	0	
Cx.interfor	1.2	3.5	0	Transmission	25	0	
Cx.interfor	1.3	3.5	0	Infection	25	0	
Cx.interfor	1.3	3.5	0	Dissemination	25	0	
Cx.interfor	1.3	3.5	0	Transmission	25	0	
Cx.interfor	1.4	3.5	0	Infection	25	0	
Cx.interfor	1.4	3.5	0	Dissemination	25	0	
Cx.interfor	1.4	3.5	0	Transmission	25	0	
Cx.interfor	1.5	3.5	0	Infection	25	0	
Cx.interfor	1.5	3.5	0	Dissemination	25	0	
Cx.interfor	1.5	3.5	0	Transmission	25	0	
Cx.interfor	1.6	3.5	1	Infection	25	145000	5,161368002
Cx.interfor	1.6	3.5	1	Dissemination	25	82000	4,913813852
Cx.interfor	1.6	3.5	1	Transmission	25	12,5	1,096910013
Cx.interfor	1.7	3.5	1	Infection	25	750	2,875061263
Cx.interfor	1.7	3.5	0	Dissemination	25	0	
Cx.interfor	1.7	3.5	0	Transmission	25	0	
Cx.interfor	1.8	3.5	1	Infection	25	1320	3,120573931
Cx.interfor	1.8	3.5	0	Dissemination	25	0	
Cx.interfor	1.8	3.5	0	Transmission	25	0	
Cx.interfor	1.9	3.5	1	Infection	25	280	2,447158031
Cx.interfor	1.9	3.5	0	Dissemination	25	0	
Cx.interfor	1.9	3.5	0	Transmission	25	0	
Cx.interfor	1.10	3.5	0	Infection	25	0	
Cx.interfor	1.10	3.5	0	Dissemination	25	0	
Cx.interfor	1.10	3.5	0	Transmission	25	0	
Cx.interfor	1.11	3.5	0	Infection	25	0	
Cx.interfor	1.11	3.5	0	Dissemination	25	0	
Cx.interfor	1.11	3.5	0	Transmission	25	0	
Cx.interfor	1.12	3.5	1	Infection	25	140000	5,146128036
Cx.interfor	1.12	3.5	1	Dissemination	25	230000	5,361727836

Cx.interfor	1.12	3.5	0	Transmission	25	0	
Cx.interfor	1.13	3.5	0	Infection	25	0	
Cx.interfor	1.13	3.5	0	Dissemination	25	0	
Cx.interfor	1.13	3.5	0	Transmission	25	0	
Cx.interfor	1.14	3.5	0	Infection	25	0	
Cx.interfor	1.14	3.5	0	Dissemination	25	0	
Cx.interfor	1.14	3.5	0	Transmission	25	0	
Cx.interfor	1.15	3.5	1	Infection	25	182000	5,260071388
Cx.interfor	1.15	3.5	1	Dissemination	25	23000	4,361727836
Cx.interfor	1.15	3.5	1	Transmission	25	212,5	2,327358934
Cx.interfor	1.16	3.5	0	Infection	25	0	
Cx.interfor	1.16	3.5	0	Dissemination	25	0	
Cx.interfor	1.16	3.5	0	Transmission	25	0	
Cx.interfor	1.17	3.5	1	Infection	25	116000	5,064457989
Cx.interfor	1.17	3.5	1	Dissemination	25	20000	4,301029996
Cx.interfor	1.17	3.5	1	Transmission	25	20	1,301029996
Cx.interfor	1.18	3.5	1	Infection	25	127000	5,103803721
Cx.interfor	1.18	3.5	1	Dissemination	25	13000	4,113943352
Cx.interfor	1.18	3.5	1	Transmission	25	20	1,301029996
Cx.interfor	1.19	3.5	1	Infection	25	960	2,982271233
Cx.interfor	1.19	3.5	1	Dissemination	25	10	1
Cx.interfor	1.19	3.5	0	Transmission	25	0	
Cx.interfor	1.20	3.5	1	Infection	25	75000	4,875061263
Cx.interfor	1.20	3.5	1	Dissemination	25	60000	4,77815125
Cx.interfor	1.20	3.5	0	Transmission	25	0	
Cx.interfor	1.21	3.5	1	Infection	25	930	2,968482949
Cx.interfor	1.21	3.5	1	Dissemination	25	150	2,176091259
Cx.interfor	1.21	3.5	1	Transmission	25	NA	
Cx.interfor	1.22	3.5	1	Infection	25	20000	4,301029996
Cx.interfor	1.22	3.5	0	Dissemination	25	0	
Cx.interfor	1.22	3.5	0	Transmission	25	0	
Cx.interfor	1.23	3.5	0	Infection	25	0	
Cx.interfor	1.23	3.5	0	Dissemination	25	0	
Cx.interfor	1.23	3.5	0	Transmission	25	0	
Cx.interfor	1.24	3.5	1	Infection	25	180000	5,255272505
Cx.interfor	1.24	3.5	1	Dissemination	25	80000	4,903089987
Cx.interfor	1.24	3.5	0	Transmission	25	0	
Cx.interfor	1.25	3.5	1	Infection	25	1050	3,021189299
Cx.interfor	1.25	3.5	1	Dissemination	25	69000	4,838849091
Cx.interfor	1.25	3.5	0	Transmission	25	0	

Cx.saltanensis	1.1	3.2	1	Infection	12	43000	4,633468456
Cx.saltanensis	1.1	3.2	1	Dissemination	12	40	1,602059991
Cx.saltanensis	1.1	3.2	1	Transmission	12	NA	
Cx.saltanensis	1.2	3.2	1	Infection	12	88000	4,944482672
Cx.saltanensis	1.2	3.2	1	Dissemination	12	70000	4,84509804
Cx.saltanensis	1.2	3.2	0	Transmission	12	0	
Cx.saltanensis	1.3	3.2	1	Infection	12	74000	4,86923172
Cx.saltanensis	1.3	3.2	1	Dissemination	12	132000	5,120573931
Cx.saltanensis	1.3	3.2	1	Transmission	12	65	1,812913357
Cx.saltanensis	1.4	3.2	0	Infection	12	0	
Cx.saltanensis	1.4	3.2	0	Dissemination	12	0	
Cx.saltanensis	1.4	3.2	0	Transmission	12	0	
Cx.saltanensis	1.5	3.2	0	Infection	12	0	
Cx.saltanensis	1.5	3.2	0	Dissemination	12	0	
Cx.saltanensis	1.5	3.2	0	Transmission	12	0	
Cx.saltanensis	1.6	3.2	1	Infection	12	120000	5,079181246
Cx.saltanensis	1.6	3.2	1	Dissemination	12	45000	4,653212514
Cx.saltanensis	1.6	3.2	0	Transmission	12	0	
Cx.saltanensis	1.7	3.2	0	Infection	12	0	
Cx.saltanensis	1.7	3.2	0	Dissemination	12	0	
Cx.saltanensis	1.7	3.2	0	Transmission	12	0	
Cx.saltanensis	1.8	3.2	1	Infection	12	73000	4,86332286
Cx.saltanensis	1.8	3.2	1	Dissemination	12	1040	3,017033339
Cx.saltanensis	1.8	3.2	0	Transmission	12	0	
Cx.saltanensis	1.9	3.2	1	Infection	12	7000	3,84509804
Cx.saltanensis	1.9	3.2	1	Dissemination	12	1650	3,217483944
Cx.saltanensis	1.9	3.2	0	Transmission	12	0	
Cx.saltanensis	1.10	3.2	1	Infection	12	79000	4,897627091
Cx.saltanensis	1.10	3.2	1	Dissemination	12	1260	3,100370545
Cx.saltanensis	1.10	3.2	0	Transmission	12	0	
Cx.saltanensis	1.11	3.2	0	Infection	12	0	
Cx.saltanensis	1.11	3.2	0	Dissemination	12	0	
Cx.saltanensis	1.11	3.2	0	Transmission	12	0	
Cx.saltanensis	1.12	3.2	1	Infection	12	76000	4,880813592
Cx.saltanensis	1.12	3.2	1	Dissemination	12	37000	4,568201724
Cx.saltanensis	1.12	3.2	0	Transmission	12	0	
Cx.p.quinquefasciatus	1.1	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.1	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.1	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.2	2.9	0	Infection	39	0	

Cx.p.quinquefasciatus	1.2	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.2	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.3	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.3	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.3	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.4	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.4	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.4	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.5	2.9	1	Infection	39	490	2,69019608
Cx.p.quinquefasciatus	1.5	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.5	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.6	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.6	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.6	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.7	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.7	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.7	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.8	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.8	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.8	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.9	2.9	1	Infection	39	NA	
Cx.p.quinquefasciatus	1.9	2.9	1	Dissemination	39	NA	
Cx.p.quinquefasciatus	1.9	2.9	1	Transmission	39	NA	
Cx.p.quinquefasciatus	1.10	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.10	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.10	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.11	2.9	1	Infection	39	370	2,568201724
Cx.p.quinquefasciatus	1.11	2.9	1	Dissemination	39	NA	
Cx.p.quinquefasciatus	1.11	2.9	1	Transmission	39	NA	
Cx.p.quinquefasciatus	1.12	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.12	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.12	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.13	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.13	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.13	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.14	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.14	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.14	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.15	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.15	2.9	0	Dissemination	39	0	

Cx.p.quinquefasciatus	1.15	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.16	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.16	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.16	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.17	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.17	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.17	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.18	2.9	1	Infection	39	170	2,230448921
Cx.p.quinquefasciatus	1.18	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.18	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.19	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.19	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.19	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.20	2.9	1	Infection	39	20	1,301029996
Cx.p.quinquefasciatus	1.20	2.9	1	Dissemination	39	NA	
Cx.p.quinquefasciatus	1.20	2.9	1	Transmission	39	NA	
Cx.p.quinquefasciatus	1.21	2.9	0	Infection	39		
Cx.p.quinquefasciatus	1.21	2.9	0	Dissemination	39		
Cx.p.quinquefasciatus	1.21	2.9	0	Transmission	39		
Cx.p.quinquefasciatus	1.22	2.9	1	Infection	39	100	2
Cx.p.quinquefasciatus	1.22	2.9	1	Dissemination	39	NA	
Cx.p.quinquefasciatus	1.22	2.9	1	Transmission	39	NA	
Cx.p.quinquefasciatus	1.23	2.9	1	Infection	39	NA	
Cx.p.quinquefasciatus	1.23	2.9	1	Dissemination	39	NA	
Cx.p.quinquefasciatus	1.23	2.9	1	Transmission	39	NA	
Cx.p.quinquefasciatus	1.24	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.24	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.24	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.25	2.9	1	Infection	39	830	2,919078092
Cx.p.quinquefasciatus	1.25	2.9	1	Dissemination	39	NA	
Cx.p.quinquefasciatus	1.25	2.9	1	Transmission	39	NA	
Cx.p.quinquefasciatus	1.26	2.9	1	Infection	39	40	1,602059991
Cx.p.quinquefasciatus	1.26	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.26	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.27	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.27	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.27	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.28	2.9	1	Infection	39	60	1,77815125
Cx.p.quinquefasciatus	1.28	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.28	2.9	0	Transmission	39	0	

Cx.p.quinquefasciatus	1.29	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.29	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.29	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.30	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.30	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.30	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.31	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.31	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.31	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.32	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.32	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.32	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.33	2.9	1	Infection	39	NA	
Cx.p.quinquefasciatus	1.33	2.9	1	Dissemination	39	720	2,857332496
Cx.p.quinquefasciatus	1.33	2.9	1	Transmission	39	NA	
Cx.p.quinquefasciatus	1.34	2.9	1	Infection	39	NA	
Cx.p.quinquefasciatus	1.34	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.34	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.35	2.9	1	Infection	39	140	2,146128036
Cx.p.quinquefasciatus	1.35	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.35	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.36	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.36	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.36	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.37	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.37	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.37	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.38	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.38	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.38	2.9	0	Transmission	39	0	
Cx.p.quinquefasciatus	1.39	2.9	0	Infection	39	0	
Cx.p.quinquefasciatus	1.39	2.9	0	Dissemination	39	0	
Cx.p.quinquefasciatus	1.39	2.9	0	Transmission	39	0	

ID: Individuals identification. Viremia ($\log_{10}$ PFU/ml). Results: 0 negative, 1 positive. Infection rate (number of positive mosquito abdomens/total number of mosquitoes fed), dissemination rate (number of mosquitoes with positive legs/total number of mosquitoes fed) of mosquitoes fed) and transmission rate (number of mosquitoes with positive saliva/total number). N: Total number of mosquitoes assayed. SLEV titers in abdomen, legs and saliva of each mosquito species in PFU/ml and $\log_{10}$ PFU/ml. NA: No data.